



**HAL**  
open science

## Assessing the ecological status of an estuarine ecosystem: linking biodiversity and food-web indicators

Camille Le Guen, Samuele Tecchio, Jean-Claude Dauvin, Gwenola de Roton, Jérémy Lobry, Mario Lepage, Jocelyne Morin, Géraldine Lassalle, Aurore Raoux, Nathalie Niquil

### ► To cite this version:

Camille Le Guen, Samuele Tecchio, Jean-Claude Dauvin, Gwenola de Roton, Jérémy Lobry, et al.. Assessing the ecological status of an estuarine ecosystem: linking biodiversity and food-web indicators. *Estuarine, Coastal and Shelf Science*, 2019, 228, pp.106339. 10.1016/j.ecss.2019.106339 . hal-02346068

**HAL Id: hal-02346068**

**<https://hal.science/hal-02346068>**

Submitted on 16 Dec 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

7  
8 3 **Assessing the ecological status of an estuarine ecosystem:**

9  
10 4 **linking biodiversity and food-web indicators**  
11  
12 5

13  
14 6 Camille Le Guen<sup>1,2</sup>, Samuele Tecchio<sup>1,3</sup>, Jean-Claude Dauvin<sup>4</sup>, Gwenola De Roton<sup>5</sup>, Jérémy  
15  
16 7 Lobry<sup>6</sup>, Mario Lepage<sup>6</sup>, Jocelyne Morin<sup>7</sup>, Géraldine Lassalle<sup>6</sup>, Aurore Raoux<sup>1,4</sup>, Nathalie  
17  
18 8 Niquil<sup>1\*</sup>  
19  
20 9

21  
22 10 <sup>1</sup> Unité Biologie des Organismes et Ecosystèmes Aquatiques (BOREA), MNHN, CNRS, IRD,  
23  
24 11 Sorbonne Université, Université de Caen Normandie, CS 14032, 14000 CAEN, France  
25

26  
27 12 <sup>2</sup> Pelagic Ecology Research Group, University of St-Andrews, KY16 9TS, St-Andrews, UK.  
28

29 13 <sup>3</sup> Sinay - Maritime Data Solution, 117 cours Caffarelli, 14000 Caen  
30

31 14 <sup>4</sup> Normandie Univ, UNICAEN, UNIROUEN, Laboratoire Morphodynamique Continentale et  
32  
33 15 Côtière M2C, UMR 6143 M2C, 24 rue des Tilleuls, 14000 Caen, France  
34

35 16 <sup>5</sup> Agence Française pour la Biodiversité, Antenne Manche – Mer du Nord – DDTM-DIRM  
36  
37 17 BP 34, 4 rue du Colonel Fabien – 76 083 LE HAVRE Cedex, France  
38

39 18 <sup>6</sup> IRSTEA, UR EABX (Aquatic Ecosystems and Global Changes Unit),  
40  
41 19 50 avenue de Verdun, 33612 Cestas CEDEX, France  
42

43  
44 20 <sup>7</sup> IFREMER, Laboratoire Ressources Halieutiques,  
45  
46 21 Avenue du Général de Gaulle, 14520 Port-en-Bessin, France  
47

48 22  
49  
50 23 \* *Corresponding author:* [cam.leguen@gmail.com](mailto:cam.leguen@gmail.com)  
51

52 24 Pelagic Ecology Research Group, University of St-Andrews, KY16 9TS, St-Andrews, UK  
53  
54 25  
55  
56  
57  
58  
59

60  
61  
62 **Highlights**  
63  
64

27

28 The Seine estuary ecosystem was described before and after the Port2000 extension.

29 A toolbox of various indices was used, describing complementary characteristics.

30 Port2000 effects were complex to separate from other human-induced and natural effects.

31 Ecological Network Analysis was useful in defining functional indicators.


32 The need to continue long-term monitoring and to use multiple tools is emphasized.

33

34

35 **Graphical abstract**  
36

37



38

39

40

41

42

43

44

119  
120  
121 **Abstract**  
122

123 43  
124  
125 44 During the last decades, the highly-anthropized Seine estuary has been impacted by  
126 45 modification of its habitats (building of a major extension of Le Havre harbour, i.e. Port2000)  
127  
128 46 and a significant natural decrease in freshwater discharge. A Before/After analysis, using a  
131  
132 47 toolbox of indicators, was applied to characterize the effects of both events on the estuarine  
133  
134 48 ecosystem status. We selected from existing tool boxes several indicators derived from food  
135  
136 49 web modelling or community composition data, such as biodiversity indicators, a guild-based  
137  
138 50 index (*i.e.* Estuarine and Lagoon Fish Index ELFI) and ecological network analysis (ENA)  
139  
140 51 indices. ENA and biodiversity indicators were applied on six spatial boxes describing the Seine  
142  
143 52 estuary and its outlet. Results showed an increase in taxonomic and functional richness over  
144  
145 53 time, mainly due to marinisation, and significant changes in food-web properties in relation to  
146  
147 54 Port2000. ENA indices appeared as a promising method in ecological status assessment,  
148  
149 55 especially for estuaries considered as inherently disturbed.  
150

151 56  
152  
153 57 **Keywords:** ecology, biodiversity, food web, ecosystem functioning, ecosystem health  
154  
155 58 indicators  
156

157 59  
158  
159 60 **1. Introduction**  
160  
161 61

162  
163  
164 62 Areas at the land-sea interface are often strongly impacted by human activities (e.g. Borja *et*  
165  
166 63 *al.*, 2012). Because estuaries are of great economic importance, they are commonly subject to  
167  
168 64 physical modifications such as port development and channelization to enhance harbour and  
169  
170 65 industrial activities (Marmin *et al.*, 2014). In addition, pollution and other anthropogenic  
171  
172 66 disturbances may cause modifications in aquatic communities (*i.e.* group of species or taxa  
173  
174  
175  
176  
177

178  
179  
180 67 present in a given area that can be characterised by the distribution of individuals or biomasses  
181  
182 68 among those species (Borja *et al.*, 2012)), that could in turn influence the whole ecosystem  
183  
184 69 functioning. A stressed ecosystem is here defined as an ecosystem undergoing changes in its  
185  
186 70 structure (e.g. productivity of its species) or functioning (e.g. food-web dynamics) following  
187  
188 71 physical, chemical or biological constraints or perturbations, either of a natural or an  
189  
190 72 anthropogenic origin. A stressed ecosystem is therefore expected to show modifications in both  
191  
192 73 its structure (e.g. species assemblages) and functioning (e.g. food-web properties), compared to  
193  
194 74 a defined baseline. Estuarine ecosystems are constituted of a mosaic of habitats linked to natural  
195  
196 75 (e.g. tide, freshwater input) and anthropogenic factors (McLusky and Elliott, 2004; Elliott and  
197  
198 76 Whitfield, 2011). However, it is difficult to distinguish between natural and human-induced  
199  
200 77 disturbances in estuaries due to their naturally high stress level and inherent variability: a  
201  
202 78 concept known as estuarine quality paradox (Dauvin, 2007; Elliott and Quintino, 2007; Dauvin  
203  
204 79 and Ruellet, 2009).

207  
208 80 Multiple tools exist to derive various indicators describing and reflecting the structure and  
209  
210 81 functioning of ecosystems (Rombouts *et al.*, 2003; de la Vega *et al.*, 2018). Biodiversity and  
211  
212 82 guild-based indices are the most commonly used and were largely developed for application to  
213  
214 83 coastal and transition water masses in Europe under the EU Water Framework Directive (WFD)  
215  
216 84 (EU, 2000; Hering *et al.*, 2010). Biodiversity indices aim to describe communities'  
217  
218 85 characteristics and enable comparisons at different spatial scales (e.g. species richness, *alpha*,  
219  
220 86 *beta* and *gamma*, diversities). Guild-based indicators are generally developed to assess the  
221  
222 87 response of ecosystems to specific or more general disturbances (*i.e.* the comparison between  
223  
224 88 the Ecology Quality Status of an estuary or a part of an estuary to a reference ecosystem with  
225  
226 89 or without moderate disturbances) (Perez-Dominguez *et al.*, 2012; Lepage *et al.*, 2016).  
227  
228 90 Delpech *et al.* (2010) suggested using a fish-based indicator called Estuarine and Lagoon Fish  
229  
230 91 Index (ELFI) to assess the ecological status of French estuaries. ELFI was based on pressure-

237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295

92 impact models and the purpose was to select relevant metrics that were known to be sensitive  
93 to anthropogenic pressures (Tableau *et al.*, 2013; Pasquaud *et al.*, 2013). However, this  
94 ecological quality indicator is mainly based on structural or taxonomic elements and do not  
95 sufficiently reflect neither the functioning of ecosystems (de Jonge *et al.*, 2006; Rombouts *et*  
96 *al.*, 2013) nor the biodiversity patterns between and within ecosystems (e.g. Vasconcelos *et al.*,  
97 2013).

98 Ecological Network Analysis (ENA) corresponds to a collection of mathematical indices  
99 describing the structure and functioning of a trophic network and characterise the overall  
100 structural properties of food webs (Ulanowicz, 1997; Niquil *et al.*, 2014). Mass-balanced food-  
101 web models such as Linear Inverse Modelling (Vézina and Platt, 1988; Niquil *et al.*, 2011;  
102 Tecchio *et al.*, 2016) and Ecopath with Ecosim (Polovina, 1984) compute values of all carbon  
103 flows in a given food web and then calculate ENA indices to assess the properties emerging  
104 from the food-web architecture.

105 A common feature of both biodiversity indices and holistic ENA indicators is that they can  
106 be related to concepts of stress, stability, resistance and resilience (Fath *et al.*, 2019; Safi *et al.*,  
107 2019). The stability of an ecosystem corresponds to its ability to maintain a comparable  
108 functioning in the presence of perturbations that drive it away from its original state  
109 (Vasconcellos *et al.*, 1997; Lobry *et al.*, 2008; Saint-Béat *et al.*, 2015). Grimm & Wissel (1997)  
110 defined resistance as the property of populations, communities or ecosystems to remain  
111 “essentially unchanged”. Resilience is the ability or the time taken by a system to recover from  
112 a change due to perturbations and is linked to stability (Pimm, 1984; Mathevet and Bousquet,  
113 2014). Although ENA indices have great potential to inform management decisions in estuaries  
114 and large tidal areas (de la Vega *et al.*, 2018), their implementation in ecosystem health  
115 assessment is not sufficiently established yet.

296  
297  
298 116 The objective of the present study was to investigate the contribution of ENA indices to the  
299  
300 117 assessment of the Seine estuarine ecosystem quality by confronting them to some more  
301  
302 118 commonly used indicators (i.e. biodiversity and guild-based indices). The Seine estuary is a  
303  
304 119 good example of a highly anthropogenic transitional area (Dauvin and Desroy, 2005; Dauvin  
305  
306 120 *et al.*, 2006). For more than 150 years, the Seine estuary has undergone many transformations,  
307  
308 121 such as shores linearization and calibration of the navigation channel from Le Havre (i.e.  
309  
310 122 entrance of the estuary) to port of Rouen (120 km upstream). Port2000 is an extension of the  
311  
312 123 existing commercial port of Le Havre, located in the Northern Channel of the Seine estuary,  
313  
314 124 that was constructed between 2002 and 2005 and that modified the morphological and  
315  
316 125 hydrological characteristics of the estuary (Dauvin *et al.*, 2006; Tecchio *et al.*, 2016). This  
317  
318 126 extension aimed to optimise the access of long-distance trade ships and to expand the available  
319  
320 127 area for those vessels in the harbour. Moreover, the navigation channel is also continuously  
321  
322 128 dredged to ensure maritime access of ships up to the port of Rouen (Marmin *et al.*, 2014). A  
323  
324 129 recent study using Ecopath models has shown that before the construction of Port2000, the  
325  
326 130 northern and navigation channels of the Seine estuary already presented signs of disturbance  
327  
328 131 (i.e. an effect, either biotic or abiotic, inducing a perturbation such as stress in an ecosystem;  
329  
330 132 relative to a specified reference state, (Rykiel JR, 1985)) due to maritime traffic, dredging, and  
331  
332 133 building works leading to a lack of connectivity between estuarine areas (Tecchio *et al.*, 2015).  
333  
334 134 Later, Tecchio *et al.* (2016) used a statistically more-advanced modelling framework (LIM-  
335  
336 135 MCMC) to estimate flows in the Seine estuary food web and compute ENA indices describing  
337  
338 136 its functional properties before and after Port2000. They showed that the northern channel  
339  
340 137 evolved towards a shorter food web and increased its resistance property. This was considered  
341  
342 138 as a potential impact of the Port2000 construction. Results from ENA indices also evidenced  
343  
344 139 that the food web in the navigation channel was constantly changing, probably due to  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354

355  
356  
357 140 continuous dredging activities, and that the southern channel could be seen as the least disturbed  
358  
359 141 area for both periods (Tecchio *et al.*, 2016).

360  
361 142 In addition to these anthropogenic disturbances, the Seine Estuary is also subject to an  
362  
363 143 increase of salinity in the lower part of the estuary, called marinisation. Marinisation is due to  
364  
365 144 a decrease in freshwater input coming from the river, associated with intrusion of marine water  
366  
367 145 in the lower part of the estuary. Marinisation is a common evolution in estuaries receiving water  
368  
369 146 from rivers and influenced by human activities (e.g. irrigation, dams) (David *et al.*, 2007;  
370  
371 147 Goberville *et al.*, 2010; Chaalali *et al.*, 2013).

372  
373 148 We used here an original combination of methods to assess the ecosystem health of the Seine  
374  
375 149 estuary before and after the construction of Port2000. These analyses were performed on six  
376  
377 150 spatial boxes previously defined to describe the estuary (Tecchio *et al.*, 2016).

380  
381 151 We first addressed the methodological question “Do the different indicators used in this  
382  
383 152 study (species-based, guild-based and ecosystem-based indices) bring similar, complementary  
384  
385 153 or different information on the ecological status of this estuarine ecosystem?” We then  
386  
387 154 addressed two ecological questions using this set of indicators: (i) How did physical distinct  
388  
389 155 areas of the estuary (spatial boxes) react to perturbations? (ii) Can we distinguish the effects  
390  
391 156 due to Port2000 from global change effects in the context of the estuary’s natural marinisation?

392  
393 157

## 394 395 158 **2. Materials and Methods**


396  
397 159

### 400 160 *2.1. Study site and periods*

401  
402 161

403  
404 162 Six spatial boxes, based on habitat characteristics and sediment types of the Seine estuary  
405  
406 163 and the adjacent eastern part of the Seine Bay, previously defined in Tecchio *et al.* (2015) were  
407  
408 164 considered: (a) three estuarine boxes: the navigation channel (CH), the northern channel (FN),  
409

414  
 415  
 416 165 and the southern channel (FS); (b) an intermediate box located at the mouth of the estuary (E4),  
 417  
 418 166 and (c) two marine spatial boxes (E2 and E14) (Fig. 1). Hydro-sedimentary and salinity  
 419  
 420 167 conditions were considered homogenous within each spatial box (Tecchio *et al.*, 2015). For  
 421  
 422 168 each zone, two sets of data were gathered corresponding to two different periods: 1996–2002,  
 423  
 424 169 representing the situation before the Port2000 construction, and 2005–2012, corresponding to  
 425  
 426 170 the recent situation, after the Port 2000 construction.



454 171  
 455  
 456 172 **Fig. 1.** Map showing the geographical position of the Seine estuary in France. The spatial boxes  
 457  
 458 173 are composed of the navigation channel (CH), the northern channel (FN), and the southern  
 459  
 460 174 channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4),  
 461  
 462 175 and two marine boxes (E2 and E14). Source: GIP Seine-Aval and SHOM.

473  
474  
475 178 2.2. Indicators based on taxonomic diversity  
476

477 179 2.2.1. Data  
478

479 180 For the subsequent indices' calculations, both fish data (from beam trawl) and  
480  
481 macrozoobenthos data (from 0.1 m<sup>2</sup> Smith McIntyre or Van Veen grabs) were collected  
482 181  
483 following identical sampling protocols between both periods, as results were standardised to  
484 182  
485 take into account differences in sampling effort. This allows quantitative and qualitative  
486 183  
487 comparisons to be made.  
488 184  
489

490 185

491  
492 186 2.2.2. Indicators  
493

494 187 The taxonomic richness is defined as the total number of taxa sampled at a given scale. It  
495  
496 was computed by pooling all sampled taxa by spatial box and period (Before Port2000 = 1996-  
497 188  
498 2002 vs After Port2000 = 2005-2012). To limit bias due to sampling design, taxonomic richness  
499 189  
500 was standardized with relation to the sampling effort by dividing it by the log-transformed total  
501 190  
502 number of samples realised for a given space-time unit (Nicolas *et al.*, 2010).  
503 191  
504

505 192 Alpha ( $\alpha$ ), beta ( $\beta$ ) and gamma ( $\gamma$ ) diversity indices give information on changes in  
506  
507 193 community composition and the processes driving them, between spatial units or periods  
508  
509 194 (Barros *et al.*, 2014).  $\alpha$ -diversity describes the taxonomic diversity on a local scale (calculated  
510  
511 for each spatial box),  $\gamma$ -diversity corresponds to the taxonomic diversity of the entire region of  
512 195  
513 interest (all six spatial boxes considered in the study) and  $\beta$ -diversity represents the variation in  
514 196  
515 taxonomic composition among the spatial boxes of the studied area (Barros *et al.*, 2014). The  
516 197  
517 mean  $\alpha$ -diversity ( $\bar{\alpha}$ ) of a region or area comprising several spatial boxes was calculated as the  
518 198  
519 average of  $\alpha$ -diversity in each spatial box (Eq. 1).  
520 199  
521

522 200  $\bar{\alpha} = \sum_{i=1}^N \alpha^i / N$  (1); with  $\alpha_i$  the taxonomic richness in the spatial box  $i$ , and  $N$  the number of  
523  
524  
525 201 spatial boxes.  
526  
527  
528  
529  
530  
531

532  
533  
534 202  $\beta$ -diversity is the ratio between  $\gamma$  and  $\bar{\alpha}$  (Eq. 2) (Anderson *et al.*, 2011) and represents the  
535  
536 203 number of times by which the taxonomic richness in the whole region considered (*i.e.* the entire  
537  
538 204 estuary) is greater than the average taxonomic richness in the different spatial boxes. In other  
539  
540 205 words, it corresponds to the ratio between regional and local species diversity and is related to  
541  
542 206 compositional heterogeneity.

543  
544  
545 207 
$$\beta = \gamma / \bar{\alpha} \quad (2)$$
  
546

547 208 with  $\gamma$ -diversity = taxonomic diversity of the entire region of interest and  $\bar{\alpha}$  = mean  $\alpha$ -diversity.

549 209 Biodiversity analyses were performed using the software package PRIMER-E v.6 (Clarke  
550  
551 210 & Gorley, 2006). A two-dimensional non-metric Multi-Dimensional Scaling (nMDS)  
552  
553 211 ordination was implemented for the six spatial boxes for each of the two periods, based on  
554  
555 212 square-root-transformed biomasses to reduce the asymmetry in species distributions, and a  
556  
557 213 Bray-Curtis similarity index computed on a species composition matrix. The nMDS plot gives  
558  
559 214 a visual representation of the proximity in species composition between samples: the closer  
560  
561 215 they are on the plot, the more similar they are in their species composition (Clarke and Warwick,  
562  
563 216 2001). The ‘stress’ index calculated by the routine in PRIMER reflects how well the ordination  
564  
565 217 is capable of preserving the rank of similarities in a bi-dimensional space, providing information  
566  
567 218 on the quality of the representation.

568  
569  
570 219

### 571 572 220 *2.3. Multimetric fish-based indicator ELFI*

#### 573 574 221 *2.3.1. Data*

575  
576 222 Fish data ‘After Port2000’ were standardized data, collected from surveys following a  
577  
578 223 normalized protocol especially designed for fish monitoring in the French estuaries for the  
579  
580 224 WFD (Appendix A). Fish data used for multimetric fish-based indicator ELFI (Delpech *et al.*,  
581  
582 225 2010) calculation ‘Before Port2000’ were collected from surveys designed for various other  
583  
584 226 monitoring purposes but were obtained using similar sampling gear (*i.e.* beam trawls with  
585  
586  
587  
588  
589  
590

591  
592  
593 227 similar net features). For standardization and comparison, we accounted for sampling effort  
594  
595 228 (trawled surface area and time spent trawling) and considered that the spatial design was  
596  
597 229 relevant regarding WFD fish sampling protocol (> 6 samples by haline zone, defined by salinity  
598  
599 230 level) following the Venice estuarine system: oligohaline (0-5); mesohaline (5-18), polyhaline  
600  
601 231 (18-30) and euhaline (<30) zones (Venice System, 1958; Elliott & McLusky, 2002). This  
602  
603 232 allowed for comparison of ELFI values between both periods.  
604  
605  
606 233

### 608 234 2.3.2. Indicator calculation

609  
610 235 ELFI has already been used to assess the ecological status of French estuaries (Tableau *et*  
611  
612 236 *al.*, 2013; Teichert *et al.*, 2018) (ecological status categories described in Appendix B). The  
613  
614 237 ELFI indicator is composed of seven metrics: density of (1) benthic fish (DB), (2) diadromous  
615  
616 238 migrant species (DDIA), (3) marine juveniles migrants (DMJ), (4) freshwater species (DFW),  
617  
618 239 (5) resident species (DER), (6) total density of fish (DT) and (7) standardised taxonomic  
619  
620 240 richness (RT/lnS) (Pasquaud *et al.*, 2013). Indeed, taxonomic richness (RT) was divided by the  
621  
622 241 log-transformed total number of samples in each spatial box (lnS) to limit the bias due to  
623  
624 242 sampling design, defining RT/lnS as the standardised taxonomic richness. These metrics are  
625  
626 243 proxies that reflect the physical quality of the habitat (DB, DER), the chemical quality of water  
627  
628 244 and sediment (DB, DMJ, DFW), but also the habitat connectivity (i.e. the degree to which  
629  
630 245 separate patches of habitat are connected; DDIA) (Hall *et al.*, 2011) and the general degradation  
631  
632 246 level, which reflects a general dysfunction in ecosystem health due to accumulation of negative  
633  
634 247 impacts (DT, RT/lnS). Contrary to the diversity indices calculated at the spatial box level, ELFI  
635  
636 248 represents the estuary level due to the high number of samples required to calculate this index.  
637  
638 249 ELFI values were computed using R (R Development Core Team, 2015). This fish index was  
639  
640 250 inter-calibrated at the European level with seven other fish indices in use in the North East  
641  
642 251 Atlantic region of Europe (Lepage *et al.*, 2016) in order to ensure that assessments provided by  
643  
644  
645  
646  
647  
648  
649

650  
651  
652 252 these indices are consistent. Data were provided by ‘*Agence de l’Eau Seine Normandie*’ for the  
653  
654 253 ‘After Port2000’ period whereas values were computed from Ifremer and CSLN (*Cellule de*  
655  
656 254 *Suivi du Littoral Normand*) fish surveys for the ‘Before Port2000’ period.  
657  
658  
659 255

## 660 256 *2.4. Indicators based on food-web models: Ecological Network Analysis (ENA)*

661  
662  
663 257

### 664 258 *2.4.1. Data sources and model construction*

667 259 We used the LIM model from Tecchio *et al.* (2016) for the three estuarine spatial boxes (FN,  
668  
669 260 FS and CH; Fig. 1) for both periods, using 13 functional groups. A summary of data sources is  
670  
671 261 given in Appendices C and D. Biomasses are expressed in gC.m<sup>-2</sup>. We constructed three  
672  
673 262 additional LIM models representing the spatial boxes E2, E4 and E14 based on the same  
674  
675 263 functional groups (Appendix C) and the same types of input data, including production, food  
676  
677 264 conversion efficiency (i.e. ratio of the biomass produced by one compartment compared to its  
678  
679 265 food intake (Christensen & Walters, 2004)), respiration, and excretion ratios. Data sources and  
680  
681 266 references used for the construction of the LIM-MCMC models are presented in detail in  
682  
683 267 Tecchio *et al.* (2015, 2016). Briefly, a system of linear equations was set up, linking species  
684  
685 268 metabolic ratios (such as production/biomass, respiration/biomass, excretion/consumption  
686  
687 269 ratios), species biomass, and diet compositions. In these equations, the unknowns were the  
688  
689 270 flows connecting the compartments. The flow constraints were established using data from  
690  
691 271 literature or other modelling works (Appendix C). Known equalities such as those including  
692  
693 272 the biomass stocks were included to constrain the model into a finite multi-dimensional space.  
694  
695 273 Metabolic parameters as well as dietary constraints (i.e. proportion of prey in predator diet)  
696  
697 274 were implemented as "inequalities": minimum and maximum values were included in the linear  
698  
699 275 equations to partially constrain the related flows. This approach produces solutions that are  
700  
701 276 robust to the effect of parameterisation uncertainty (Hines *et al.*, 2015; Hines *et al.*, 2018).  
702  
703  
704  
705  
706  
707  
708

709  
710  
711 277 Then, this space was sampled by using a Markov-Chain Monte Carlo routine to obtain 200,000  
712  
713 278 solutions of the set of flow values. Ecological Network Analysis was applied to these 200,000  
714  
715 279 solutions, using the R software packages *NetIndices* (Soetaert *et al.*, 2015) and *enaR* (Borrett  
716  
717 & Lau, 2014), along with scripts developed for the present study [see Tecchio *et al.* (2016) for  
718 280  
719 additional methodological details].  
720 281

721  
722 282 To assess statistical differences in ENA indices between both periods, the non-parametric  
723  
724 283 effect size statistic introduced by Cliff (1993) was applied [see Tecchio *et al.* (2016) for  
725  
726 284 methodological details] as we were dealing with a situation of large amount of data (each ENA  
727  
728 285 index was calculated 200,000 times before Port2000 and the same number after Port2000)  
729  
730 286 which renders the estimation of classical parametric test statistics unfeasible (Mulholland &  
731  
732 287 Jones, 1968).  
733  
734 288

#### 735 288 736 737 289 *2.4.2. Indices*

738  
739 290 We calculated six relevant ENA indices: the Total System Throughput (T..), the System  
740  
741 291 Omnivory Index (SOI), the Finn's Cycling Index (FCI), the Ascendency (A), the relative  
742  
743 292 Redundancy (R/DC) and the Detritivory/Herbivory ratio (D/H). This set of indices was used by  
744  
745 293 Tecchio *et al.* (2016) to characterize the inner part of the Seine estuary and most of them have  
746  
747 294 been proposed as management tools in the OSPAR Convention context (Safi *et al.*, 2019). The  
748  
749 295 Total system throughput (T..) corresponds to the sum of all flows occurring in the system, and  
750  
751 296 acts as an indicator of system size and activity (Rutledge *et al.*, 1976; Aoki, 1988; Latham,  
752  
753 297 2006). The Finn's Cycling Index (FCI) represents the fraction of the flows in the system that is  
754  
755 298 generated by recycling and was calculated as the proportion of T.. generated by cycling (Finn,  
756  
757 299 1976; Finn, 1980). Although other FCI definitions exist, this particular one was chosen to  
758  
759 300 remain consistent with previous work and another common modelling approach, *Ecopath with*  
760  
761 301 *Ecosim*. The Ascendency was calculated as the product of the Total system throughput (T..)

768  
769  
770 302 and the Average Mutual Information (AMI), where AMI expresses the degree of organisation  
771  
772 303 of exchanges between the different functional groups (increasing AMI indicates higher dietary  
773  
774 304 specialisation; Hirata & Ulanowicz, 1984). High ascendancy values mean that the system is  
775  
776 305 more actively channelling flows along more specific pathways, whereas low values have been  
777  
778 306 linked to system immaturity (Ortiz & Wolff, 2002). Relative redundancy (R/DC) was calculated  
780  
781 307 as the ratio between the internal flows overhead and the total development capacity of the  
782  
783 308 ecosystem (Ulanowicz, 1986; Ulanowicz, 2001) and measures the amount of parallel trophic  
784  
785 309 pathways connecting the trophic compartments. R/DC acts as an indicator of inefficiency of the  
786  
787 310 network (Hirata & Ulanowicz, 1984; Bondavalli *et al.*, 2000) but it is also a way to increase  
788  
789 311 ecosystem resilience because parallel pathways can replace each other (Ulanowicz, 1997). The  
790  
791 312 System Omnivory Index (SOI) was calculated as the mean of the omnivory indices of each  
792  
793 313 consumer compartment, weighted by the logarithm of their consumption (Christensen &  
794  
795 314 Walters, 1993; Libralato, 2008). Omnivory is defined as the variability of prey trophic levels  
796  
797 315 where high SOI values correspond to a food web with a web-like structure (i.e. with several  
798  
799 316 pathways between compartments) whereas low SOI values reflect a chain-like structure with  
800  
801 317 fewer pathways (Dimitrios *et al.*, 2018). Therefore, SOI is an indicator of the overall dietary  
802  
803 318 adaptation of the consumers, and an increase in SOI generally corresponds to a stabilizing  
804  
805 319 response to an external disturbance (Fagan, 1997; Libralato, 2008). The Detritivory/Herbivory  
806  
807 320 ratio (D/H) was calculated as the ratio between the sum of all predation flows on the detritus  
808  
809 321 compartment (detritivory, flows from detritus to consumers) and the sum of all predation flows  
810  
811 322 on primary producers (herbivory, flows from phytoplankton and microphytobenthos to  
812  
813 323 consumers). An increase in D/H would indicate that the ecosystem shifts towards a more  
814  
815 324 detritus-based food web, depending less on plant material for trophic interactions (Ulanowicz,  
816  
817 325 1992; Luong *et al.*, 2014; Niquil *et al.*, 2014).  
818  
819  
820  
821 326

827  
828  
829 **3. Results**  
830

831 328

833 329 Results concern “species-based” (species diversity and biomasses), “community-based”  
834 330 (fish index ELFI) and “ecosystem-based” (Ecological Network Analysis) indicators, which  
835  
836 331 provide information at different levels.  
837


838  
839 332

841 333 *3.1. Species-based indices related to diversity and biomasses*

843 334 *3.1.1. Biomass trend*

845  
846 335 In the estuary boxes CH and FN, fish and macrozoobenthos biomasses increased between  
847  
848 336 periods (Fig. 2, Table 1). For the southern channel (FS), while macrozoobenthos biomass  
849  
850 337 increased, fish biomass remained stable, and it appeared that this was the case for all fish groups  
851  
852 338 (Fig. 2). For instance, the biomass of piscivorous fish in FS averaged 0.057 g C.m<sup>-2</sup> in the first  
853  
854 339 period and approached 0.055 g C.m<sup>-2</sup> in the second one. In CH, while total fish biomass  
855  
856 340 increased, the biomass of planktivorous fish decreased (Fig. 2). This was mainly related to two  
857  
858 341 pelagic species: *Clupea harengus* and *Sprattus sprattus*. In the same way, for FN, while the  
859  
860 342 total fish biomass increased, the biomass of piscivorous and planktivorous fishes slightly  
861  
862 343 decreased (Fig. 2). This was quantitatively compensated by a strong increase in benthos feeders’  
863  
864 344 biomass. The decrease in planktivorous fish biomass in FN was due to four species *Clupea*  
865  
866 345 *harengus*, *Gasterosteus aculeatus*, *Osmerus eperlanus* and *Sprattus sprattus*. The overall  
867  
868 346 biomass (including fish, macrozoobenthos and the suprabenthos) increased for the three  
869  
870 347 estuarine spatial boxes. Among the three estuarine boxes, the ratio of fish biomass on benthic  
871  
872 348 invertebrate biomass (BFish/BMacrozoobenthos) was the highest in CH (0.09 in the first period  
873  
874 349 and 0.03 in the second one). Nonetheless, this ratio was divided by three in CH congruently to  
875  
876 350 the sharp increase in benthic macrofauna biomass between periods that was multiplied by 7  
877  
878 351 (Fig. 2, Table 1).  
879  
880  
881  
882  
883  
884  
885

886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944



352

353 **Fig. 2.** Barplots representing fish biomass, macrozoobenthos biomass, the sum of fish and  
 354 macrozoobenthos biomasses, and the ratio  $BFish/BMacrozoobenthos$ , for all spatial boxes  
 355 before the construction of Port2000 (first bars) and after (second bars). The spatial boxes are  
 356 composed of the navigation channel (CH), the northern channel (FN), and the southern channel  
 357 (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two  
 358 marine boxes (E2 and E14).

359

360


**Table 1.** Summary of comparisons between periods (Before and After Port2000 construction) for biomass values, biodiversity indices, and the ELFI index. See text for details on the indices. < indicates that the value of the index is lower Before than After Port2000; > indicates that the value of the index is greater Before than After Port2000.

		Spatial boxes					
		CH	FN	FS	E4	E14	E2
Biomass	Fish	<	<	<	>	<	>
	Benthos	<	<	<	<	<	<
TR	Fish	<	<	<	<	>	>
	Benthos	<	<	<	<	<	<
$\alpha$	Fish	<	<	<	<	>	>
	Benthos	<	<	<	<	<	<
Diversity	$\square^*$	Fish			<		
		Benthos			<		
	$\beta^*$	Fish			<		
		Benthos			>		
	$\gamma^*$	Fish			<		
		Benthos			<		
ELFI*				<			

\*  $\square$ ,  $\beta$  and  $\gamma$  diversity indices are computed for the whole estuary

### 3.1.2. Taxonomic richness and species composition

The taxonomic richness increased between periods, from 16 to 47 fish taxa and from 5 to 29 macrozoobenthos taxa in the navigation channel (CH) (Fig. 3). The same trend was observed in the Northern and Southern channels (FN and FS; Fig. 3). The estuary box CH showed particularly low values of macrozoobenthos taxonomic richness compared to the other spatial boxes (Fig. 3). The intermediate box E4 was the richest spatial box in terms of fish (with 51 taxa during the first period and 59 taxa during the second one) and benthic macrofauna (with 177 taxa during the first period and 307 during the second one). No strong changes were observed for the fish taxonomic richness in the marine spatial boxes E14 and E2. All taxa that were gained between periods, both for fish and macrozoobenthos, were exclusively of marine origin. Results on standardised taxonomic richness are given in Figure 3.




**Fig. 3.** Barplots representing taxonomic richness divided by the log-transformed total number of samples realised within the six spatial boxes before (first bars) and after (second bars) the construction of Port2000 for fish and macrozoobenthos. The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14).

### 3.1.3. Diversity indices

The  $\alpha$ ,  $\beta$  and  $\gamma$  taxonomic diversity analyses between periods showed that the average number of fish species  $\alpha$  increased from 32.8 to 41 species, and from 65.7 to 177.2 for macrozoobenthos.  $\gamma$ -diversity increased from 54 to 71 taxa for fish and from 195 to 353 taxa for macrozoobenthos, and  $\beta$ -diversity increased from 1.65 to 1.73 for fish and decreased from 2.97 to 1.99 for macrozoobenthos (Table 1).

The MDS ordination based on fish composition data showed a remarkable spatial segregation according to the salinity gradient between the estuarine and the marine spatial boxes (Fig. 4 top). The same pattern was observed with the MDS ordination based on macrozoobenthos species (Fig. 4 bottom). The ‘stress’ value of 0.05 was very low, underlying the good quality of the representation.

1063  
 1064  
 1065  
 1066  
 1067  
 1068  
 1069  
 1070  
 1071  
 1072  
 1073  
 1074  
 1075  
 1076  
 1077  
 1078  
 1079  
 1080  
 1081  
 1082  
 1083  
 1084  
 1085  
 1086  
 1087  
 1088  
 1089  
 1090  
 1091  
 1092  
 1093  
 1094  
 1095  
 1096  
 1097  
 1098  
 1099  
 1100  
 1101  
 1102  
 1103  
 1104  
 1105  
 1106  
 1107  
 1108  
 1109  
 1110  
 1111  
 1112  
 1113  
 1114  
 1115  
 1116  
 1117  
 1118  
 1119  
 1120  
 1121



399

**Fig. 4.** MDS ordination based on Bray-Curtis similarity matrices from biomass data (square-root transformed) for fish (on the upper part) and for macrozoobenthos (on the lower part), for all spatial boxes during the first (first bars), before Port2000, and the recent (second bars) periods.

404


405 CH appeared as the spatial box that has experienced the greatest change in fish composition  
 406 between periods, with less than 20% similarity in the community composition before and after  
 407 Port2000. In addition, the benthic macrofauna composition in this estuary box demonstrated  
 408 few similarities with the other spatial boxes, and this for both periods. For fish in the ‘Before’  
 409 period, marine boxes E2, E4 and E14 formed a group well separated from the inner estuary,

1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180

410 where FN and FS were grouped and CH isolated from these two other estuarine boxes. Similar  
411 results were obtained for macrozoobenthos. But in the ‘After’ period, the marine box E4 was  
412 grouped, for fish, with not only the estuary boxes FN and FS, but also CH (Fig. 4). This  
413 similarity in biodiversity composition was only limited to fish compartment; the benthos  
414 remained structured in a very similar way, in both periods.

415  
416 *3.2. A community-based index for fish: ELFI*

417 During the period before Port2000, the Seine estuary showed lower values of ELFI (from  
418 0.1 to 0.14) compared to the period after Port2000, the highest value recorded being 0.68 in  
419 2006 (Fig. 5). The metric DDIA, indicator of habitat connectivity related to diadromous species,  
420 showed strong changes in the recent period (after Port2000), with a value of 0.88 in 2006, 0.70  
421 in 2010 and 0.71 in 2012 (missing value in 2011), meaning that this metric gradually decreased  
422 after the Port2000 construction.



423  
424 **Fig. 5.** ELFI values for the whole estuary for 10 years, between 1995 and 2012, with the class  
425 boundaries of the Water Framework Directive.  
426

1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239

427 3.3. *Ecosystem-based indicators: food webs and ENA*

428 ENA indices showed that all indices in the two marine zones (E2 and E14) remained similar  
429 between the two periods (Table 2). R/DC and SOI decreased between before and after “Port  
430 2000” in the estuary box (FS). All indices, except SOI, increased between the two periods in  
431 the intermediate box E4 and in the estuary boxes CH and FN.

432 Total System Throughput (T..), representing the size and activity of the system, significantly  
433 increased in the estuary boxes CH and FN as well as in the intermediate box E4 and remained  
434 stable in all the other spatial boxes (Table 2); Cliff’s delta; Fig. 6).

435 The SOI, reflecting the omnivory in the system, significantly decreased in the estuary box  
436 FS between the two periods. (Fig. 6). The D/H ratio significantly increased in the estuary boxes  
437 CH, FN and in the intermediate box E4, while remaining stable in the marine boxes E14, E2  
438 and in FS (Table 2, Fig. 6).

439 The FCI reflecting the recycling in the system, significantly increased between the two  
440 periods in the estuary boxes CH, FN and in the intermediate box E4 (Table 2, Fig. 6).

441 The ascendancy (A) and relative redundancy (R/DC) indices significantly increased in CH  
442 between the two periods. The ascendancy also significantly increased in the intermediary box  
443 E4 between the two periods (Table 2, Fig. 6).

444  
445

1240  
1241  
1242  
1243  
446  
1244  
447  
1245  
448  
1246  
449  
1247  
1248  
450  
1249  
451  
1250  
1251


**Table 2.** ENA means and standard deviations, derived from LIM-MCMC models, for all spatial boxes before (1996-2002) and after (2005-2012) Port2000 construction. < indicates that the value of the index is lower Before than After Port2000; > indicates that the value of the index is greater Before than After Port2000. Only statistically significant results based on Cliff's delta are listed. ns: non-significant.

ENA	Period	CH	FN	FS	E4	E14	E2
T..	Before	1935 ± 337	2244 ± 251	2600 ± 240	2424 ± 347	2475 ± 296	2447 ± 364
T..	After	2332 ± 361	2463 ± 273	2507 ± 227	2687 ± 352	2494 ± 294	2564 ± 328
<b>Evolution T..</b>		<	<	ns	<	ns	ns
FCI	Before	0.087 ± 0.03	0.124 ± 0.03	0.163 ± 0.03	0.128 ± 0.03	0.135 ± 0.02	0.149 ± 0.04
FCI	After	0.119 ± 0.03	0.175 ± 0.04	0.178 ± 0.04	0.153 ± 0.03	0.131 ± 0.02	0.145 ± 0.03
<b>Evolution FCI</b>		<	<	Ns	<	ns	ns
Ascendency	Before	3542 ± 675	4066 ± 465	4648 ± 461	4382 ± 692	4480 ± 589	4408 ± 721
Ascendency	After	4285 ± 742	4453 ± 534	4532 ± 450	4942 ± 737	4590 ± 588	4649 ± 667
<b>Evolution Ascendency</b>		<	<	Ns	<	ns	ns
R/DC (%)	Before	32.6 ± 2	40.6 ± 2	43.7 ± 2	40.6 ± 2	40.9 ± 2	41.5 ± 2
R/DC (%)	After	39.2 ± 2	41.8 ± 1	41.7 ± 2	42.4 ± 2	41.6 ± 2	42.6 ± 2
<b>Evolution R/DC(%)</b>		<	<	>	<	ns	ns
SOI	Before	0.145 ± 0.02	0.133 ± 0.02	0.136 ± 0.01	0.146 ± 0.01	0.143 ± 0.01	0.130 ± 0.01
SOI	After	0.146 ± 0.01	0.125 ± 0.01	0.119 ± 0.01	0.148 ± 0.01	0.142 ± 0.01	0.137 ± 0.01
<b>Evolution SOI</b>		ns	ns	>	ns	ns	ns
D/H	Before	0.317 ± 0.12	0.589 ± 0.12	0.796 ± 0.13	0.600 ± 0.11	0.637 ± 0.11	0.692 ± 0.16
D/H	After	0.553 ± 0.12	0.776 ± 0.16	0.758 ± 0.15	0.760 ± 0.15	0.658 ± 0.11	0.718 ± 0.12
<b>Evolution D/H</b>		<	<	ns	<	ns	ns

1295  
1296  
1297  
1298

1299  
 1300  
 1301  
 1302  
 1303  
 1304  
 1305  
 1306  
 1307  
 1308  
 1309  
 1310  
 1311  
 1312  
 1313  
 1314  
 1315  
 1316  
 1317  
 1318  
 1319  
 1320  
 1321  
 1322  
 1323  
 1324  
 1325  
 1326  
 1327  
 1328  
 1329  
 1330  
 1331  
 1332  
 1333  
 1334  
 1335  
 1336  
 1337  
 1338  
 1339  
 1340  
 1341  
 1342  
 1343  
 1344  
 1345  
 1346  
 1347  
 1348  
 1349  
 1350  
 1351  
 1352  
 1353  
 1354  
 1355  
 1356  
 1357

452



453

454

**Fig. 6.** Ecological network analysis (ENA) indices for all spatial boxes for the period before (first bars) and after (second bars) Port2000. The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14). Means and standard deviations are represented. Asterisks represent significant changes assessed from Cliff's delta statistics (level 0.05; between before and after construction).

455

456

457

458

459

460

461

462

**4. Discussion**

1341

1342

1343

1344

*4.1. Limitations*

1345

Fish species were sampled using beam trawls, which target the first 50cm above the seafloor. The fish assemblage was therefore biased, the pelagic species likely being underestimated. In addition, the edges of each habitat were not sampled. Therefore, a few fish species might not have been sampled. However, the sampling gears and methods were similar between periods

466

467

468

1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416

469 and among spatial boxes, allowing comparison between time periods and spatial boxes. ELFI  
470 values, for both periods, need to be considered with some caution, given the differences in  
471 sampling protocol between periods that may slightly affect the comparability.

472  
473 *4.2. Signs of estuarine marinisation*

474 The biomasses of marine species increased between both periods in the Seine estuary. For  
475 example, *Pomatoschistus minutus* biomass increased in two estuarine boxes (FN, FS), and in  
476 the intermediate box (E4), and *P. microps* was newly observed in these areas after Port2000.  
477 These two fish species, especially *P. minutus* are mostly marine species (Leitão *et al.*, 2006),  
478 suggesting a tendency of marinisation. Previously termed by David *et al.* (2007) and later used  
479 in several papers (Pasquaud *et al.*, 2012; Chevillot *et al.*, 2016, 2017), « marinisation »  
480 describes the process whereby marine waters flow far upstream into the basin along with a  
481 higher abundance of marine species in the estuarine transitional zone. The Seine estuary, as  
482 many north-eastern Atlantic Ocean estuarine ecosystems (Goberville *et al.*, 2010), is going  
483 through a process of ‘marinisation’. The time period 2005-2012 was characterised by a  
484 significantly lower discharge of the Seine river (Dauvin and Pezy, 2013), resulting in an  
485 increase of salinity up to 5 in the estuarine boxes (Bacq *et al.*, 2013). The salinity in the marine  
486 boxes remained stable between periods. The arrival of marine species finding newly suitable  
487 conditions in most spatial boxes might explain the increase in taxonomic richness. This  
488 marinisation tendency was supported by the MDS ordination showing a clear spatial (along the  
489 salinity gradient) and temporal (before and after Port2000) segregation, indicating that  
490 marinisation modified the ecological structure of communities. This is also supported by the  
491 higher ELFI values in the Seine estuary after Port2000 which is likely a consequence of the  
492 marinisation enhancing the taxonomic richness and biomass of fish in the estuarine spatial  
493 boxes. The same trend has already been observed in the Gironde estuary where changes in fish

1417  
1418  
1419 494 assemblages and an increase of the ELFI index were attributed to an increase in temperature  
1420  
1421 495 and salinity indicating marinisation (Pasquaud *et al.* 2012; Chevillot *et al.*, 2016, 2017).  
1422

1423 496

### 1426 497 4.3. Signs of anthropogenic impacts

#### 1428 498 4.3.1. The global estuary

1430 499 In the Seine estuary, the ELFI indicator increases with ecosystem quality (Le Pape *et al.*,  
1431  
1432 500 2015). The ecological status of the Seine estuary therefore improved from a ‘bad’ status before  
1433  
1434 501 Port2000 to a ‘poor’ status after Port2000. Diadromous species abundance was associated to  
1435  
1436 502 habitat connectivity, as opposed to habitat fragmentation (barriers to migratory fish) (Pasquaud  
1437  
1438 503 *et al.*, 2013). After Port2000, the highest DDIA value recorded was 0.88 in 2006 and then the  
1439  
1440 504 general pattern of this metric for that period was characterized by an overall decrease of DDIA,  
1441  
1442 505 with values dropping down to 0.71 in 2012, likely due to a lower connectivity for migrant  
1443  
1444 506 species such as eel *Anguilla anguilla* or grey mullet *Liza ramada*, associated with the extension  
1445  
1446 507 of the breakwaters between FN and CH during the Port2000 construction. As an example, smelt  
1447  
1448 508 *O. eperlanus* is a diadromous species which was formerly abundant in the upper part of the  
1449  
1450 509 northern channel for reproduction (Morin *et al.*, 2015). Extending the breakwaters between FN  
1451  
1452 510 and CH during the Port2000 construction might have impacted the connectivity between these  
1453  
1454 511 two zones, and *O. eperlanus* has been negatively affected by this rupture of connectivity.  
1455  
1456 512 Moreover, it seemed that the differences in fish assemblages between the spatial boxes widened  
1457  
1458 513 ( $\beta$ -diversity increased from 1.65 to 1.73), which might be related to availability of new habitats  
1459  
1460 514 for marine species inside the estuary. Concerning benthic species, the diversity trend between  
1461  
1462 515 spatial boxes showed the reverse trend, showing signs of standardisation ( $\beta$ -diversity decreased  
1463  
1464 516 from 2.97 to 1.99).  
1465  
1466 517

#### 1470 518 4.3.2. Estuarine disturbed boxes

1476  
1477  
1478 519 Comparatively to the rest of the estuary, strong effects of perturbations were expected in the  
1479  
1480 520 navigation channel (CH) because of the continuous dredging and the high hydrodynamics. The  
1481  
1482 521 estuarine box FN was the spatial box the most directly affected by the construction of Port2000  
1483  
1484 522 and by the mitigation measures such as the artificial formation of meanders, which might have  
1485  
1486 523 modified the hydrodynamics (Cuvilliez *et al.*, 2015). In addition, an increase in water turbidity  
1487  
1488 524 was noted in this box (S. Lesourd, personal communication).

1491 525 The increase in detritivory/herbivory ratio (D/H) observed in both CH and FN between the  
1492  
1493 526 two periods suggests an ecosystem more stressed after the Port2000 construction (Niquil *et al.*,  
1494  
1495 527 2014), probably reacting to perturbations by enhancing opportunistic behaviours such as  
1496  
1497 528 detritivory. The higher carbon recycling (FCI) observed after Port2000 in these two estuarine  
1498  
1499 529 spatial boxes can also be a sign of increasing stress (Vasconcellos *et al.*, 1997). However,  
1500  
1501 530 increasing FCI and D/H values could also be interpreted by a sign of increasing maturity  
1502  
1503 531 (Odum, 1969; Christensen, 1995; Allesina and Ulanowicz, 2004; Vasconcelos *et al.*, 2007) but  
1504  
1505 532 other ENA indices support the statement of increasing stress. Indeed, the relative redundancy  
1506  
1507 533 (R/DC) significantly increased across periods in CH and FN, highlighting the fact that the  
1508  
1509 534 system was enduring a higher level of stress (Hirata & Ulanowicz, 1984; Bondavalli *et al.*,  
1510  
1511 535 2000) but was also more able to resist to disturbances as the energy transfers could be  
1512  
1513 536 maintained through the trophic network via other pathways (Ulanowicz, 1997; Rybarczyk &  
1514  
1515 537 Elkaïm, 2003).

#### 1518 538 1519 1520 1521 539 *4.3.3. Estuarine reference box*

1522  
1523 540 The FS, which was initially considered as the least varying spatial box (Tecchio *et al.*, 2015)  
1524  
1525 541 and thus a reference, showed a reduction in redundancy (R/DC), omnivory (SOI) and detritivory  
1526  
1527 542 (D/H), while maintaining a stable system activity (T..) and carbon recycling (FCI). The trophic  
1528  
1529 543 specialisation was evidenced by the congruent reduction in omnivory (leading to a more chain-

1535  
1536  
1537 544 like structure of the food web) and redundancy (reduction of parallel pathways), which can be  
1538  
1539 545 a sign of maturity. Results for FS therefore suggested that the reduced disturbance conditions  
1540  
1541 546 here favoured an expected ecological succession.  
1542  
1543  
1544 547

#### 1546 548 *4.3.4. Intermediate box*

1548 549 E4 is the transitional area between the inner estuary and the marine environment. Results  
1549  
1550 550 from ENA indices show an increase in system activity (T..), redundancy (R/DC), carbon cycling  
1551  
1552 551 (FCI), omnivory (SOI) and detritivory (D/H). Therefore, as for CH and FN, following the same  
1553  
1554 552 interpretation framework, we could assume that the system in E4 was increasing its stabilizing  
1555  
1556 553 mechanisms when facing stronger disturbances.  
1557  
1558 554

#### 1560 555 *4.3.5. Marine boxes*

1563 556 Results from ENA indices show no significant differences between the two periods for both  
1564  
1565 557 spatial boxes E14 and E2. Therefore, we can assume that the construction of Port2000 has no  
1566  
1567 558 negative effect on the food web dynamics of these two marine boxes. However, the spatial box  
1568  
1569 559 E2 contains the dumping site of Octeville for dredged sediments from the Le Havre harbour  
1570  
1571 560 (Marmin *et al.*, 2014). For these two spatial boxes, we noticed that the fish taxonomic richness  
1572  
1573 561 decreased whereas the benthic macrofauna richness increased. The relative importance of  
1574  
1575 562 benthic communities had intensified in E2, as shown by the decrease in the  
1576  
1577 563 BFish/BMacrozoobenthos ratio between both periods, contrary to what was noticed in E14.  
1578  
1579 564 This might be explained by a potential increasing turbidity in E2 due to the dumping of  
1580  
1581 565 sediments that could affect fish more than the benthic communities. However, the effect of the  
1582  
1583 566 Port2000 construction on these two marine boxes cannot be evidenced.  
1584  
1585  
1586 567  
1587  
1588  
1589  
1590  
1591  
1592  
1593

1594  
1595  
1596 568 *4.4. Relative effects of marinisation and port construction*  
1597

1598 569 *4.4.1. The global estuary*  
1599

1600 570 The higher taxonomic diversity and ELFI index values observed after Port2000 in a context  
1601  
1602 571 of marinisation suggest a decrease of ecosystem stress after Port2000, as taxonomic richness is  
1603  
1604 572 considered a key factor in ecosystem stability in response to anthropogenic pressures (e.g.  
1605  
1606 573 McCann, 2000; Chapin *et al.*, 2000). However, investigations in the different spatial boxes were  
1607  
1608 574 necessary to detect potential effects of the Port2000 construction.  
1609  
1610

1611 575

1612  
1613 576 *4.4.1. Estuarine disturbed boxes*  
1614

1615 577 The food web in CH is continuously disturbed due to the dredging activity, greatly limiting  
1616  
1617 578 the colonisation of the seafloor by benthic organisms, as seen in other locations (e.g. in the  
1618  
1619 579 Venice lagoon: Pranovi *et al.*, 2003). In CH, the community was characterized by an extremely  
1620  
1621 580 low benthos richness and biomass compared to the other spatial boxes. Nonetheless,  
1622  
1623 581 communities subjected to regular disturbances are also more adapted, less sensitive, and recover  
1624  
1625 582 more quickly than undisturbed communities (Dernie *et al.*, 2003; Marmin *et al.*, 2014). Results  
1626  
1627 583 from ENA indices in the two boxes CH and FN suggest an ecosystem more stressed after the  
1628  
1629 584 Port2000 construction. Species-based indices were very useful to detect the effects of dredging  
1630  
1631 585 activities which directly affect the settlement of benthic communities. This effect was also  
1632  
1633 586 reflected in the ENA indices.  
1634  
1635

1636 587

1637  
1638  
1639 588 *4.4.2. Estuarine reference box*  
1640

1641 589 In FS, both fish and macrozoobenthos taxonomic richness increased. However, the ratio  
1642  
1643 590 B<sub>Fish</sub>/B<sub>Macrozoobenthos</sub> remained stable. The MDS analysis showed that the fish  
1644  
1645 591 composition in this spatial box did not deeply change between periods. ENA indices suggest a  
1646  
1647 592 mature system for FS for both periods. Therefore, this estuarine spatial box could be confirmed  
1648  
1649

1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711

593 as the least stressed area of the Seine estuary, as previously assessed by Tecchio *et al.* (2016),  
594 and that species-based indicators, ELFI and ENA indices bring similar information for this  
595 spatial box.

596  
597 *4.4.3. Intermediate box*

598 We assumed that E4 should exhibit the highest taxonomic richness and biomasses given its  
599 intermediate position between marine and estuarine waters (Dauvin & Desroy, 2005). It had the  
600 highest fish and benthic macrofauna taxonomic richness of all spatial boxes. Similar results  
601 were found by Martino and Able (2003), who showed that in the Mullica river-Great Bay  
602 (USA), fish species richness increased from the marine stations to the ones at the mouth of the  
603 estuary, and then strongly decreased towards the estuarine sampling stations. The MDS analysis  
604 put forward that E4 could be seen as an intermediate spatial box for fish, but as a marine box  
605 for the macrobenthos community. Indeed, this confirmed that E4 itself can be considered as an  
606 ‘ecocline’, i.e. a boundary of progressive changes (spatial and ecological) between two different  
607 systems, the ocean and the estuary (Attrill and Rundle, 2002; Basset *et al.*, 2013). Finally,  
608 results from ENA indices demonstrated an increase in system activity, redundancy, cycling,  
609 omnivory and detritivory between the two periods. Therefore, as for CH and FN, ENA indices  
610 enabled us to identify that the system was increasing its stabilizing mechanisms, probably to  
611 face stronger disturbances.

612  
613 *4.4.4. Marine boxes*

614 Finally, in the bay, E14 and E2 marine boxes appeared similar in terms of fish and  
615 macrozoobenthos taxonomic composition between the two periods. Results from ENA indices  
616 show no significant differences between the two periods for both spatial boxes. Indices based

1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770

617 on diversity and biomasses showed the importance of benthic communities has intensified in  
618 E2 but the origin of this change remains unknown.

619

620 *4.4.5. Summary*

621 The estuarine marinisation effects are potentially stronger than the effects of the Port2000  
622 construction. Indeed, on the one hand, the marinisation of the Seine estuary mostly changed the  
623 species assemblages, therefore affecting the ecosystem structure (i.e. biological and physical  
624 architecture of an ecosystem), due to the arrival of numerous marine species. On the other hand,  
625 the anthropogenic pressures affected habitat connectivity and food-web dynamics (i.e. transfer  
626 of energy from one part of the food web to another, Lindeman, 1942), assessed with ENA  
627 indicators such as system activity, omnivory, recycling or detritivory. The main information  
628 brought by the combination of indices was summarized in Table 3.

629

630

631

1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829

632  
633

**Table 3.** Main information about the functioning and the status of the six spatial boxes.

Method	Main results	Conclusions
Biotic indices based on diversity and biomasses => <b>give information about the structure of communities</b>	Increase of the taxonomic richness between periods for the estuarine spatial boxes (CH, FN, FS) and the biomass of fish and macrozoobenthos increased	<b>Arrival of marine species</b> such as <i>P. minutus</i> in relation to the marination of the estuary (lower river discharge in the second period)
	CH: low values of macrozoobenthos richness	Low benthos biomass due to dredging and high hydrodynamics in this part of the estuary
	E4: the richest area for both periods	This corresponds to the rich <i>Abra alba-Lagis koreni</i> macrobenthic community and the feeding zone for fish (mainly flatfish).
	The MDS ordination showed a spatial segregation between CH and the other zones, and a temporal change between periods, both for fish and macrozoobenthos	<b>Evolution in salinity plays a determinant role in the spatial gradient of the fauna; both periods were clearly distinguished</b>
Biotic indices based on the classification of species in trophic groups => <b>give information about the functioning of the food web</b>	FN: higher carbon recycling and detritivory (increase in detritus input) + higher redundancy (more species)	<b>Stress increased in the estuary mainly for CH and FN zones. The other zones were less disturbed.</b>
	CH: increase in system activity, carbon cycling, ascendancy, redundancy and detritivory	
Biotic index based on the classification of species in ecological groups => <b>give information about the health status of ecosystems</b>	FS: still in 'high ecological status'	Reference box
	Increase in ELFI values between periods	The ecological status was supposed to have improved but it is likely due to the arrival of marine species only
	Decrease of the metric DDIA (diadromous species) in the second period, indicator of a lower habitat connectivity	<b>Lower connectivity due to the extension of the breakwaters</b> between FN and CH during the Port2000 construction

634

635

636

637

638

639

ELFI was sensitive to the arrival of marine species and therefore was highly relevant in a context of marination of the estuary. ENA indices appeared as good 'surveillance indicators', that Shephard *et al.* (2015) proposed as indicators that would be a good complement to operational indicators, for example in the context of the Marine Strategy Framework Directive application. ENA indices are also useful to take ecosystem-based management measures for

1830  
1831  
1832 640 such ecosystems (de la Vega *et al.*, 2018; Safi *et al.*, 2019). In addition, their association with  
1833  
1834 641 biodiversity indicators made the analysis more robust as suggested by Rombouts *et al.* (2013).  
1835  
1836 642 For instance, the arrival of new species and the higher biomasses observed in some spatial boxes  
1837  
1838  
1839 643 between periods can also explain increases in some ENA indicators such as system activity,  
1840  
1841 644 omnivory or redundancy but this cannot be evidenced and requires further investigations.  
1842

## 1843 645

### 1844 646 **5. Conclusions**

1847 647

1848

1849 648 The effects of the Port2000 construction were not as evident as expected due to the presence  
1850  
1851 649 of other natural (e.g. marinisation) and anthropogenic (e.g. continuous harbour activities)  
1852  
1853 650 perturbations, which makes the interpretation of ecological indicators difficult. ELFI was  
1854  
1855 651 particularly relevant in a context of marinisation, as this index is sensitive to the arrival of new  
1856  
1857 652 marine species. But the different metrics used to calculate ELFI are also indicative of  
1858  
1859 653 anthropogenic impacts (e.g. DDIA and habitat connectivity affected by the construction of  
1860  
1861 654 Port2000), making ELFI a very powerful tool to assess the health status of this estuary. The use  
1862  
1863 655 of ENA indices alone was not sufficient to describe changes in ecosystem functioning between  
1864  
1865 656 the two periods, distinguishing the effects of the Port2000 construction and the process of  
1866  
1867 657 marinisation, but they appeared as an interesting complementary set of tools for characterising  
1868  
1869 658 the evolution of food-web dynamics. Indeed, ENA indices allowed us to observe changes that  
1870  
1871 659 occurred in specific ecosystem processes and to identify how an ecosystem becomes more  
1872  
1873 660 stable (e.g. by increasing omnivory) under increasing stress. We showed that detecting changes  
1874  
1875 661 related to anthropogenic disturbances alone in the Seine requires using ecological indicators at  
1876  
1877 662 different levels (species, communities and ecosystem) and at multiple scales of observations.  
1878  
1879 663 This work emphasizes the need to use several metrics to assess ecosystem health status and to  
1880  
1881 664 continue long-term monitoring of such complex and reactive ecosystems.  
1882  
1883  
1884  
1885  
1886  
1887  
1888

1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947

665 **Acknowledgements**

666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682

The authors would like to acknowledge Valérie Girardin and Justine Lequesne (UMR CNRS LMNO) for their help in the statistical analysis, Anik Brind'Amour (IFREMER) for her help on fish data integration and statistics, Armonie Tous Rius for creating the AntropoSeine database, Alexandrine Baffreau (UMR CNRS M2C) for her help in checking this database, Barbara Leroy (Agence de l'Eau Seine-Normandie), Philippe Riou (Ifremer Port-en-Bessin), Nicolas Bacq, Stéphanie Moussard and Guillaume Daan (GIP-Seine Aval) for providing advice, maps and data from their Seine estuary sampling programs. We also thank Camille de la Vega and three anonymous referees for their helpful comments on the manuscript. This study was funded by the ANTROPOSEINE project (Seine Aval Public Interest Group, GIPSA, Rouen, France). The authors declare no competing financial interests. Nathalie Niquil is also funded by the EU DG Environment project EcApRHA ('Addressing gaps in biodiversity indicator development for the OSPAR Region from data to ecosystem assessment: Applying an ecosystem approach to (sub) regional habitat assessments'; Grant No. 11.0661/2015/712630/SUB/ENVC.2 OSPAR).

1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006

683 **References**

- 684 Allesina S, Ulanowicz RE (2004) Cycling in ecological networks: Finn's index revisited.  
685 *Computational Biology and Chemistry*, **28**, 227–233.
- 686 Anderson MJ, Crist TO, Chase JM et al. (2011) Navigating the multiple meanings of  $\beta$   
687 diversity: A roadmap for the practicing ecologist. *Ecology Letters*, **14**, 19–28.
- 688 Aoki I (1988) Entropy laws in ecological networks at steady state. *Ecological Modelling*, **42**,  
689 289–303.
- 690 Attrill M., Rundle S. (2002) Ecotone or Ecocline: Ecological Boundaries in Estuaries.  
691 *Estuarine, Coastal and Shelf Science*, **55**, 929–936.
- 692 Bacq N, Lemoine J-P, Moussard S, Le Hir P, Lobry J (2013) *Projet MESSCENE:*  
693 *modélisation des fonctions écologiques de l'estuaire de la Seine et Scénarios*  
694 *d'évolution*. Rapport final du GIP Seine-Aval pour la Fondation de France, 69 pp.
- 695 Barros F, Blanchet H, Hammerstrom K, Sauriau P, Oliver J (2014) A framework for  
696 investigating general patterns of benthic  $\beta$ -diversity along estuaries. *Estuarine, Coastal*  
697 *and Shelf Science*, **149**, 223–231.
- 698 Basset A, Barbone E, Elliott M et al. (2013) A unifying approach to understanding  
699 transitional waters: Fundamental properties emerging from ecotone ecosystems.  
700 *Estuarine, Coastal and Shelf Science*, **132**, 5–16.
- 701 Bondavalli C, Ulanowicz R, Bodini A (2000) Insights into the processing of carbon in the  
702 South Florida Cypress Wetlands: a whole-ecosystem approach using network analysis.  
703 *Journal of Biogeography*, **27**, 697–710.
- 704 Borja Á, Basset A, Bricker S et al. (2012) Classifying ecological quality and integrity of  
705 estuaries. In: *Treatise on Estuarine and Coastal Science* (ed McLusky EW& D), pp.  
706 125–162. Waltham: Academic Press.
- 707 Borrett SR, Lau MK (2014) enaR: An r package for Ecosystem Network Analysis. *Methods in*

2007  
2008  
2009 708 *Ecology and Evolution*, **5**, 1206–1213.  
2010  
2011 709 Chaalali A, Chevillot X, Beaugrand G et al. (2013) Changes in the distribution of copepods in  
2012  
2013 710 the Gironde estuary: A warming and marinisation consequence? *Estuarine, Coastal and*  
2014  
2015 711 *Shelf Science*, **134**, 150–161.  
2016  
2017  
2018 712 Chapin F, Zavaleta E, Eviner V et al. (2000) Consequences of changing biodiversity. *Nature*,  
2019  
2020 713 **405**, 234–242.  
2021  
2022 714 Chardy P (1987) Modèle de simulation du système benthique des sédiments grossiers du golfe  
2023  
2024 715 normand breton (Manche). *Oceanologica acta*, **10**, 421–434.  
2025  
2026 716 Chardy P, Dauvin J-C (1992) Carbon flows in a subtidal fine sand community from the  
2027  
2028 717 western English 703 Channel: a simulation analysis. *Marine Ecology Progress Series*,  
2029  
2030 718 **81**, 147–181.  
2031  
2032  
2033 719 Chevillot X, Pierre M, Rigaud A, Drouineau H, Chaalali A, Sautour B, Lobry J (2016) Abrupt  
2034  
2035 720 shifts in the Gironde fish community: an indicator of ecological changes in an estuarine  
2036  
2037 721 ecosystem. *Marine Ecology Progress Series*, **549**, 137–151.  
2038  
2039 722 Chevillot X, Drouineau H, Lambert P, Carassou L, Sautour B, Lobry J (2017) Toward a  
2040  
2041 723 phenological mismatch in estuarine pelagic food web? *PLoS ONE*, **12**, e0173752.  
2042  
2043 724 Christensen V (1995) Ecosystem maturity—towards quantification. *Ecological Modelling*, **77**,  
2044  
2045 725 3–32.  
2046  
2047 726 Christensen V, Walters CJ (1993) Trophic models of aquatic ecosystems. In: *ICLARM*  
2048  
2049 727 *Conference Proceedings No. 26*, p. 390.  
2050  
2051  
2052 728 Christensen V, Walters CJ (2004) Ecopath with Ecosim: Methods, capabilities and  
2053  
2054 729 limitations. *Ecological Modelling*, **172**, 109–139.  
2055  
2056 730 Clarke K, Gorley R (2006) Primer v6: user manual/tutorial. PRIMER-E, Plymouth.  
2057  
2058 731 Clarke KR, Warwick RM (2001) *Change in marine communities. An approach to statistical*  
2059  
2060 732 *analysis and interpretation.*, 2nd edn. Plymouth Marine Laboratory, UK, 1-172 pp.  
2061  
2062  
2063  
2064  
2065

2066  
2067  
2068 733 Cliff N (1993) Dominance statistics: Ordinal analyses to answer ordinal questions.  
2069  
2070 734 *Psychological Bulletin*, **114**, 494–509.  
2071  
2072 735 Cuvilliez A, Lafite R, Deloffre J, Lemoine M, Langlois E, Sakho I (2015) River flow control  
2073  
2074 on intertidal mudflat sedimentation in the mouth of a macrotidal estuary.  
2075 736  
2076 *Geomorphology*, **239**, 174–181.  
2077 737  
2078 738 Dauvin J (2007) Paradox of estuarine quality: benthic indicators and indices, consensus or  
2079  
2080 debate for the future. *Marine Pollution Bulletin*, **55**, 271–281.  
2081 739  
2082 740 Dauvin JC, Desroy N (2005) The food web in the lower part of the Seine estuary: A synthesis  
2083  
2084 of existing knowledge. *Hydrobiologia*, **540**, 13–27.  
2085 741  
2086 742 Dauvin J, Pezy J (2013) Long-term changes of the Seine estuary suprabenthos (1996-2012).  
2087  
2088 *Journal of Experimental Marine Biology and Ecology*, **448**, 93–103.  
2089 743  
2090 744 Dauvin JC, Ruellet T (2009) The estuarine quality paradox: Is it possible to define an  
2091  
2092 ecological quality status for specific modified and naturally stressed estuarine  
2093  
2094 ecosystems? *Marine Pollution Bulletin*, **59**, 38–47.  
2095 746  
2096 747 Dauvin JC, Desroy N, Janson AL, Vallet C, Duhamel S (2006) Recent changes in estuarine  
2098  
2099 benthic and suprabenthic communities resulting from the development of harbour  
2100 748  
2101 infrastructure. *Marine Pollution Bulletin*, **53**, 80–90.  
2102 749  
2103 750 David V, Sautour B, Chardy P (2007) Successful colonization of the calanoid copepod  
2104  
2105 *Acartia tonsa* in the oligo-mesohaline area of the Gironde estuary (SW France) - Natural  
2106 751  
2107 or anthropogenic forcing? *Estuarine, Coastal and Shelf Science*, **71**, 429–442.  
2108 752  
2109 753 de Jonge VN, Elliott M, Brauer V (2006) Marine monitoring: Its shortcomings and mismatch  
2110  
2111 with the EU water framework directive’s objectives. *Marine Pollution Bulletin*, **53**, 5–  
2112 754  
2113 19.  
2114 755  
2115 756 de la Vega C, Schückel U, Horn S, Kröncke I, Asmus R, Asmus H (2018) How to include  
2117  
2118 ecological network analysis results in management? A case study of three tidal basins of  
2119 757  
2120  
2121  
2122  
2123  
2124

2125  
2126  
2127 758 the Wadden Sea, south-eastern North Sea. *Ocean & Coastal Management*, **163**, 401–  
2128  
2129 759 416.  
2130  
2131 760 Delpech C, Courrat A, Pasquaud S et al. (2010) Development of a fish-based index to assess  
2132  
2133 the ecological quality of transitional waters: The case of French estuaries. *Marine*  
2134 761  
2135 *Pollution Bulletin*, **60**, 908–918.  
2136 762  
2137  
2138 763 Dernie KM, Kaiser MJ, Warwick RM (2003) Recovery rates of benthic communities  
2139  
2140 764 following physical disturbance. *Journal of Animal Ecology*, **72**, 1043–1056.  
2141  
2142 765 Dimitrios K, Konstantinos T, Athanassios M (2018) Assessing ecological and fisheries  
2143  
2144 766 implications of the EU landing obligation in Eastern Mediterranean. *Journal of Sea*  
2145  
2146 767 *Research*, **141**, 99–111.  
2147  
2148  
2149 768 Elliott M, McLusky DS (2002) The Need for Definitions in Understanding Estuaries.  
2150  
2151 769 *Estuarine, Coastal and Shelf Science*, **55**, 815–827.  
2152  
2153 770 Elliott M, Quintino V (2007) The Estuarine Quality Paradox, Environmental Homeostasis and  
2154  
2155 771 the difficulty of detecting anthropogenic stress in naturally stressed areas. *Marine*  
2156  
2157 772 *Pollution Bulletin*, **54**, 640–645.  
2158  
2159 773 Elliott M, Whitfield A (2011) Challenging paradigms in estuarine ecology and management.  
2160  
2161 774 *Estuarine, Coastal and Shelf Science*, **94**, 306–314.  
2162  
2163 775 EU (2000) Parliament and Council Directive 2000/60/EC of 23rd October 2000. Establishing  
2164  
2165 776 a Framework for Community Action in the field of Water Policy. Official Journal PE-  
2166  
2167 777 CONS 3639/1/00 REV 1, Brussels.  
2168  
2169  
2170 778 Fagan W (1997) Omnivory as a stabilizing feature of natural communities. *American*  
2171  
2172 779 *Naturalist*, **150**, 554–567.  
2173  
2174 780 Fath BD, Asmus H, Asmus R et al. (2019) Ecological network analysis metrics: The need for  
2175  
2176 781 an entire ecosystem approach in management and policy. *Ocean & Coastal*  
2177  
2178 782 *Management*, **174**, 1–14.  
2179  
2180  
2181  
2182  
2183

2184  
2185  
2186 783 Finn J (1976) Measures of ecosystem structure and function derived from analysis of flows.  
2187  
2188 784 *Journal of Theoretical Biology*, **56**, 363–380.  
2189  
2190 785 Finn J (1980) Flow analysis of models of the Hubbard Brook ecosystem. *Ecology*, **61**, 562–  
2191  
2192 571.  
2193 786  
2194  
2195 787 Goberville E, Beaugrand G, Sautour B, Tréguer P (2010) Climate-driven changes in coastal  
2196  
2197 788 marine systems of western Europe. *Marine Ecology Progress Series*, **408**, 129–147.  
2198  
2199 789 Grimm V, Wissel C (1997) Babel, or the ecological stability discussions: an inventory and  
2200  
2201 790 analysis of terminology and a guide for avoiding confusion. *Oecologia*, **109**, 323–334.  
2202  
2203 791 Le Guellec C, Bodin P (1992) Meiobenthos of the Bay of Saint-Brieuc (North Brittany,  
2204  
2205 792 France). I: quantitative distribution in subtidal and intertidal zones. *Oceanologica acta*,  
2206  
2207 793 **15**, 661–671.  
2208  
2209 794 Hall CJ, Jordaan A, Frisk MG (2011) The historical influence of dams on diadromous fish  
2210  
2211 795 habitat with a focus on river herring and hydrologic longitudinal connectivity. *Landscape*  
2212  
2213 796 *Ecology*, **26**, 95–107.  
2214  
2215 797 Hering D, Borja A, Carstensen J et al. (2010) The European Water Framework Directive at  
2216  
2217 798 the age of 10: A critical review of the achievements with recommendations for the  
2218  
2219 799 future. *Science of the Total Environment*, **408**, 4007–4019.  
2220  
2221 800 Hines DE, Lisa JA, Song B, Tobias CR, Borrett SR (2015) Estimating the effects of seawater  
2222  
2223 801 intrusion on an estuarine nitrogen cycle by comparative network analysis. *Marine*  
2224  
2225 802 *Ecology Progress Series*, **524**, 137–154.  
2226  
2227 803 Hines DE, Ray S, Borrett SR (2018) Uncertainty analyses for Ecological Network Analysis  
2228  
2229 804 enable stronger inferences. *Environmental Modelling and Software*, **101**, 117–127.  
2230  
2231 805 Hirata H, Ulanowicz R (1984) Information theoretical analysis of ecological networks.  
2232  
2233 806 *International Journal of Systems Science*, **15**, 261–270.  
2234  
2235 807 Latham LG (2006) Network flow analysis algorithms. *Ecological Modelling*, **192**, 586–600.  
2236  
2237  
2238  
2239  
2240  
2241  
2242

2243  
2244  
2245 808 Le Pape O, Lepage M, Féra P (2015) La démarche de développement d'indicateurs basés sur  
2246  
2247 809 l'ichtyofaune pour évaluer la qualité écologique des eaux de transition françaises dans le  
2248  
2249 810 cadre de la DCE : une marche forcée pour des résultats positifs. *Norois*, **235**, 37–49.  
2250  
2251 811 Leitão R, Martinho F, Neto JM, Cabral H, Marques JC, Pardal M a. (2006) Feeding ecology,  
2252  
2253 812 population structure and distribution of *Pomatoschistus microps* (Krøyer, 1838) and  
2254  
2255 813 *Pomatoschistus minutus* (Pallas, 1770) in a temperate estuary, Portugal. *Estuarine,*  
2256  
2257 814 *Coastal and Shelf Science*, **66**, 231–239.  
2258  
2259 815 Lepage M, Harrison T, Breine J et al. (2016) An approach to intercalibrate ecological  
2260  
2261 816 classification tools using fish in transitional water of the North East Atlantic. *Ecological*  
2262  
2263 817 *Indicators*, **67**, 318–327.  
2264  
2265 818 Libralato S (2008) System Omnivory Index. *Ecological Indicators*, **4**, 3472–3477.  
2266  
2267 819 Lindeman RL (1942) The Trophic-Dynamic Aspect of Ecology. *Ecology*, **23**, 399–417.  
2268  
2269 820 Lobry J, David V, Pasquaud S, Lepage M, Sautour B, Rochard E (2008) Diversity and  
2270  
2271 821 stability of an estuarine trophic network. *Marine Ecology Progress Series*, **358**, 13–25.  
2272  
2273 822 Luong A, De Laender F, Olsen Y, Vadstein O, Dewulf J, Janssen C (2014) Inferring time-  
2274  
2275 823 variable effects of nutrient enrichment on marine ecosystems using inverse modelling  
2276  
2277 824 and ecological network analysis. *Science of the Total Environment*, **493**, 708–718.  
2278  
2279 825 Marmin S, Dauvin J-C, Lesueur P (2014) Collaborative approach for the management of  
2280  
2281 826 harbour-dredged sediment in the Bay of Seine (France). *Ocean & Coastal Management*,  
2282  
2283 827 **102**, 328–339.  
2284  
2285 828 Martino EJ, Able KW (2003) Fish assemblages across the marine to low salinity transition  
2286  
2287 829 zone of a temperate estuary. *Estuarine, Coastal and Shelf Science*, **56**, 969–987.  
2288  
2289 830 Mathevet R, Bousquet F (2014) *Résilience & environnement - Penser les changements socio-*  
2290  
2291 831 *écologiques*. Buchet Chastel, Paris, 170 pp., 170 pp.  
2292  
2293 832 McCann KS (2000) The diversity-stability debate. *Nature*, **405**, 228–233.  
2294  
2295  
2296  
2297  
2298  
2299  
2300  
2301

2302  
2303  
2304 833 McLusky DS, Elliott M (2004) *The estuarine ecosystem: ecology, threats, and management.*,  
2305  
2306 834 Third edn. Oxford University Press, 214p pp.  
2307  
2308 835 Morin J, Brind'Amour A, Fisson C et al. (2015) *Peuplements halieutiques en estuaire de*  
2309  
2310 836 *Seine et baie de Seine orientale - Distribution et évolution des abondances des*  
2311  
2312 837 *principales espèces capturées au chalut à perche lors des campagnes scientifiques*  
2313  
2314 838 *réalisées par l'Ifremer entre 1995 et 2010. Rapport d'étude, 70 pp.*  
2315  
2316  
2317 839 Mulholland H, Jones C (1968) *Fundamentals of Statistics*. Butterworth-Heinemann, London,  
2318  
2319 840 301 pp.  
2320  
2321 841 Napoleon C, Raimbault V, Fiant L, Riou P, Lefebvre S, Lampert L, Claquin P (2012)  
2322  
2323 842 Spatiotemporal dynamics of physicochemical and photosynthetic parameters in the  
2324  
2325 843 central English Channel. *Journal of Sea Research*, **69**, 43–52.  
2326  
2327  
2328 844 Nicolas D, Lobry J, Lepage M et al. (2010) Fish under influence: A macroecological analysis  
2329  
2330 845 of relations between fish species richness and environmental gradients among European  
2331  
2332 846 tidal estuaries. *Estuarine, Coastal and Shelf Science*, **86**, 137–147.  
2333  
2334 847 Niquil N, Saint-Béat B, Johnson G a et al. (2011) *Inverse Modeling in Modern Ecology and*  
2335  
2336 848 *Application to Coastal Ecosystems*, Vol. 9. Elsevier Inc., 115-133 pp.  
2337  
2338 849 Niquil N, Baeta a, Marques J, Chaalali a, Lobry J, Patricio J (2014) Reaction of an estuarine  
2339  
2340 850 food web to disturbance: Lindeman's perspective. *Marine Ecology Progress Series*, **512**,  
2341  
2342 851 141–154.  
2343  
2344  
2345 852 Odum EP (1969) The Strategy of Ecosystem Development. *Science*, **164**, 262–270.  
2346  
2347 853 Ortiz M, Wolff M (2002) Trophic models of four benthic communities in Tongoy Bay  
2348  
2349 854 (Chile): comparative analysis and preliminary assessment of management strategies.  
2350  
2351 855 *Journal of Experimental Marine Biology and Ecology*, **268**, 205–235.  
2352  
2353 856 Pasquaud S, Béguer M, Larsen MH, Chaalali A, Cabral H, Lobry J (2012) Increase of marine  
2354  
2355 857 juvenile fish abundances in the middle Gironde estuary related to warmer and more  
2356  
2357  
2358  
2359  
2360

2361  
2362  
2363 858 saline waters, due to global changes. *Estuarine, Coastal and Shelf Science*, **104–105**, 46–  
2364  
2365 859 53.  
2366  
2367 860 Pasquaud S, Courrat A, Fonseca VF et al. (2013) Strength and time lag of relationships  
2368  
2369 between human pressures and fish-based metrics used to assess ecological quality of  
2370 861  
2371 estuarine systems. *Estuarine, Coastal and Shelf Science*, **134**, 119–127.  
2372 862  
2373 863 Perez-Dominguez R, MacI S, Courrat A et al. (2012) Current developments on fish-based  
2374  
2375 indices to assess ecological-quality status of estuaries and lagoons. *Ecological*  
2376 864  
2377 *Indicators*, **23**, 34–45.  
2378 865  
2379 866 Pimm SL (1984) The complexity and stability of ecosystems. *Nature*, **307**, 321–326.  
2380  
2381 867 Polovina J (1984) Model of a coral reef ecosystem. The Ecopath model and its application to  
2382  
2383 French Frigate Shoals. *Coral Reefs*, **3**, 1–11.  
2384 868  
2385 869 Pranovi F, Libralato S, Raicevich S, Granzotto a., Pastres R, Giovanardi O (2003)  
2386  
2387 Mechanical clam dredging in Venice lagoon: Ecosystem effects evaluated with a trophic  
2388  
2389 870 mass-balance model. *Marine Biology*, **143**, 393–403.  
2390  
2391 871  
2392 872 R Development Core Team (2015) R: A Language and Environment for Statistical  
2393  
2394 Computing.  
2395 873  
2396 874 Ratsimbazafy RA (1998) *Spectres de taille des peuplements benthiques de sables grossiers de*  
2397  
2398 *la Manche. Thèse de doctorat*. Paris, Muséum National d’Histoire Naturelle, 523 pp.  
2399 875  
2400 876 Rombouts I, Beaugrand G, Artigas L et al. (2013) Evaluating marine ecosystem health : Case  
2401  
2402 studies of indicators using direct observations and modelling methods. *Ecological*  
2403  
2404 877 *Indicators*, **24**, 353–365.  
2405 878  
2406 879 Rutledge R, Basorre B, Mulholland R (1976) Ecological stability: an information theory  
2407  
2408 880 viewpoint. *Journal of Theoretical Biology*, **57**, 355.  
2409  
2410 881 Rybarczyk H, Elkaïm B (2003) An analysis of the trophic network of a macrotidal estuary:  
2411  
2412 The Seine Estuary (Eastern Channel, Normandy, France). *Estuarine, Coastal and Shelf*  
2413  
2414 882  
2415  
2416  
2417  
2418  
2419

2420  
2421  
2422 883 *Science*, **58**, 775–791.  
2423  
2424 884 Rykiel JR EJ (1985) Towards a definition of ecological disturbance. *Australian Journal of*  
2425  
2426 *Ecology*, **10**, 361–365.  
2427 885  
2428  
2429 886 Safi G, Giebels D, Larissa Arroyo N et al. (2019) Vitamine ENA : A framework for the  
2430  
2431 887 development of ecosystem-based indicators for decision makers. *Ocean and Coastal*  
2432  
2433 888 *Management*, **174**, 116–130.  
2434  
2435 889 Saint-Béat B, Baird D, Asmus H et al. (2015) Trophic networks: How do theories link  
2436  
2437 890 ecosystem structure and functioning to stability properties? A review. *Ecological*  
2438  
2439 891 *Indicators*, **52**, 458–471.  
2440  
2441 892 Shephard S, Greenstreet S, Piet G, Rindorf A, Dickey-Collas M (2015) Surveillance  
2442  
2443 893 indicators and their use in implementation of the Marine Strategy Framework Directive.  
2444  
2445 894 *ICES Journal of Marine Science*, **72**, 2269–2277.  
2446  
2447 895 Soetaert K, Kones JK, van Oevelen D (2015) Package ‘NetIndices’. Estimating network  
2448  
2449 896 indices, including trophic structure of foodwebs in R.  
2450  
2451 897 Spilmont N, Migné A, Lefebvre A, Artigas L, Rauch M, Davoult D (2005) Potential  
2452  
2453 898 contribution of sporadic pelagic deposits on the benthic metabolism of an exposed sandy  
2454  
2455 899 beach during emersion (Wimereux, eastern English Channel, France). *Journal of Sea*  
2456  
2457 900 *Research*, **53**, 161–167.  
2458  
2459 901 Tableau A, Drouineau H, Delpech C et al. (2013) A fish-based index of estuarine ecological  
2460  
2461 902 quality incorporating information from both scientific fish survey and experts  
2462  
2463 903 knowledge. *Ecological Indicators*, **32**, 147–156.  
2464  
2465 904 Tecchio S, Rius AT, Dauvin J-C et al. (2015) The mosaic of habitats of the Seine estuary:  
2466  
2467 905 Insights from food-web modelling and network analysis. *Ecological Modelling*, **312**, 91–  
2468  
2469 906 101.  
2470  
2471 907 Tecchio S, Chaalali A, Raoux A et al. (2016) Evaluating ecosystem-level anthropogenic  
2472  
2473  
2474  
2475  
2476  
2477  
2478

2479  
2480  
2481 908 impacts in a stressed transitional environment: The case of the Seine estuary. *Ecological*  
2482  
2483 909 *Indicators*, **612**, 833–845.  
2484  
2485 910 Teichert N, Lepage M, Lobry J (2018) Beyond classic ecological assessment: The use of  
2486  
2487 functional indices to indicate fish assemblages sensitivity to human disturbance in  
2488 911 estuaries. *Science of the Total Environment*, **639**, 465–475.  
2489  
2490 912  
2491  
2492 913 Ulanowicz RE (1986) *Growth and development: Ecosystems phenomenology*. Springer-  
2493  
2494 914 Verlag, New York, USA.  
2495  
2496 915 Ulanowicz R (1992) Ecosystem health and trophic flow networks. In: *Ecosystem Health: New*  
2497  
2498 916 *Goals for Environmental Management* (eds Costanza R, Norton B, Haskell B). Island  
2499  
2500 917 Press, Washigton, DC & Covelo, CA, USA.  
2501  
2502 918 Ulanowicz RE (1997) *Ecology, the Ascendent Perspective*. Columbia University Press, New  
2503  
2504 919 York, 201 pp.  
2505  
2506 920 Ulanowicz RE (2001) Information theory in ecology. *Journal of Computational Chemistry*,  
2507  
2508 921 **25**, 393–399.  
2509  
2510 922 Vasconcellos M, Mackinson S, Sloman K, Pauly D (1997) The stability of trophic mass-  
2511  
2512 923 balanced models of marine ecosystems: a comparative analysis. *Ecological modelling*,  
2513  
2514 924 **100**, 125–134.  
2515  
2516 925 Vasconcelos RP, Reis-Santos P, Fonseca V et al. (2007) Assessing anthropogenic pressures  
2517  
2518 926 on estuarine fish nurseries along the Portuguese coast: A multi-metric index and  
2519  
2520 927 conceptual approach. *Science of the Total Environment*, **374**, 199–215.  
2521  
2522 928 Vasconcelos RP, Le Pape O, Costa MJ, Cabral HN (2013) Predicting estuarine use patterns of  
2523  
2524 929 juvenile fish with Generalized Linear Models. *Estuarine, Coastal and Shelf Science*, **120**,  
2525  
2526 930 64–74.  
2527  
2528 931 Venice System (1958) Symposium on the Classification of Brackish Waters, Venice, April 8-  
2529  
2530 932 14, 1958. *Archives Oceanography and Limnology*, **11**, 1–248.  
2531  
2532  
2533  
2534  
2535  
2536  
2537

2538  
2539  
2540 933 Vézina AF, Platt T (1988) Food web dynamics in the ocean. I. Best-estimates of flow  
2541  
2542 934 networks using inverse methods. *Marine Ecology Progress Series*, **42**, 269–287.  
2543  
2544 935  
2545  
2546 936  
2547  
2548  
2549  
2550  
2551  
2552  
2553  
2554  
2555  
2556  
2557  
2558  
2559  
2560  
2561  
2562  
2563  
2564  
2565  
2566  
2567  
2568  
2569  
2570  
2571  
2572  
2573  
2574  
2575  
2576  
2577  
2578  
2579  
2580  
2581  
2582  
2583  
2584  
2585  
2586  
2587  
2588  
2589  
2590  
2591  
2592  
2593  
2594  
2595  
2596

2597  
2598  
2599  
2600  
2601  
2602  
2603  
2604  
2605  
2606  
2607  
2608  
2609  
2610  
2611  
2612  
2613  
2614  
2615  
2616  
2617  
2618  
2619  
2620  
2621  
2622  
2623  
2624  
2625  
2626  
2627  
2628  
2629  
2630  
2631  
2632  
2633  
2634  
2635  
2636  
2637  
2638  
2639  
2640  
2641  
2642  
2643  
2644  
2645  
2646  
2647  
2648  
2649  
2650  
2651  
2652  
2653  
2654  
2655

937 **Figure legends**

938

939 **Fig. 1.** Map showing the geographical position of the Seine estuary in France. The spatial boxes  
940 are composed of the navigation channel (CH), the northern channel (FN), and the southern  
941 channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4),  
942 and two marine boxes (E2 and E14). Source: GIP Seine-Aval and SHOM.

943

944 **Fig. 2.** Barplots representing fish biomass, macrozoobenthos biomass, the sum of fish and  
945 macrozoobenthos biomasses, and the ratio B<sub>Fish</sub>/B<sub>Macrozoobenthos</sub>, for all spatial boxes  
946 before the construction of Port2000 (first bars) and after (second bars). The spatial boxes are  
947 composed of the navigation channel (CH), the northern channel (FN), and the southern channel  
948 (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two  
949 marine boxes (E2 and E14).

950

951 **Fig. 3.** Barplots representing taxonomic richness divided by the log-transformed total number  
952 of samples realised within the six spatial boxes before (first bars) and after (second bars) the  
953 construction of Port2000 for fish and macrozoobenthos. The spatial boxes are composed of the  
954 navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary  
955 boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2  
956 and E14).

957

958 **Fig. 4.** MDS ordination based on Bray-Curtis similarity matrices from biomass data (square-  
959 root transformed) for fish (on the upper part) and for macrozoobenthos (on the lower part), for  
960 all spatial boxes during the first (first bars), before Port2000, and the recent (second bars)  
961 periods.

962

963 **Fig. 5.** ELFI values for the whole estuary for 10 years, between 1995 and 2012, with the class  
964 boundaries of the Water Framework Directive.

965

966 **Fig. 6.** Ecological network analysis (ENA) indices for all spatial boxes for the period before  
967 (first bars) and after (second bars) Port2000. The spatial boxes are composed of the navigation  
968 channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an  
969 intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14).  
970 Means and standard deviations are represented. Asterisks represent significant changes assessed  
971 from Cliff's delta statistics (level 0.05; between before and after construction).

2656  
2657  
2658 **Appendices**  
2659

2660 **Appendix A.** Data sources used in the calculation of ELFI index  
2661  
2662  
2663  
2664  
2665  
2666

<b>Indicator</b>	<b>Before Port2000</b>	<b>After Port2000</b>	<b>Boundary classes</b>
	Biomasses from		
	Ifremer and CSLN.	Index values from	
ELFI	Use of an R script	Agence de l'Eau Seine-	Delpech et al., 2010
index	provided by Mario	Normandie	
	Lepage (Irstea, UR		
	EABX)		

2681  
2682  
2683  
2684 Fish data ‘After Port2000’ are standardized data from surveys designed for ELFI calculation  
2685 following normalized protocol. Fish data used for ELFI calculation ‘Before Port2000’ were  
2686 obtained using similar sampling gear (beam trawls with similar net features), accounting for  
2687  
2688  
2689  
2690 sampling effort (trawled surface area) and considering a relevant spatial design regarding WFD  
2691  
2692 fish sampling protocol (> 6 tows by haline zone)  
2693  
2694  
2695  
2696  
2697  
2698  
2699  
2700  
2701  
2702  
2703  
2704  
2705  
2706  
2707  
2708  
2709  
2710  
2711  
2712

**Appendix B.** Range values between the five ‘Ecological Quality Status’ levels considered for the ELFI indicator (Delpech et al., 2010). Five levels of EQS classified as ‘high’, ‘good’, ‘moderate’, ‘poor’, and ‘bad’ were used. ‘High status’ means very low human pressure. ‘Good status’ means a ‘slight’ deviation from this condition, ‘moderate status’ means moderate deviation, ‘poor status’ means high deviation and ‘bad status’ means very high human pressure.

Range values for	
EQS level	ELFI
High	$0.90 \leq \text{ELFI} \leq 1.00$
Good	$0.68 \leq \text{ELFI} < 0.90$
Moderate	$0.45 \leq \text{ELFI} < 0.68$
Poor	$0.23 \leq \text{ELFI} < 0.45$
Bad	$0.00 \leq \text{ELFI} < 0.23$

2774  
2775  
2776  
2777  
2778  
2779  
2780  
2781  
2782  
2783  
2784  
2785  
2786  
2787  
2788  
2789  
2790  
2791  
2792  
2793  
2794  
2795  
2796  
2797  
2798  
2799  
2800  
2801  
2802  
2803  
2804  
2805  
2806  
2807  
2808  
2809  
2810  
2811  
2812  
2813  
2814

**Appendix C. Data sources used in the calculation of the different indicators**

<b>Group</b>	<b>Compartment</b>	<b>Biomass data</b>	<b>Biodiversity indicators</b>	<b>ENA indices</b>	<b>ELFI</b>
1	Seabirds	-		X	
2	Fish, piscivores	Ifremer, CSLN, Le Havre port authority sampling	X	X	X
3	Fish, planktivores	programs (Liteau, Port2000, Grand Canal,	X	X	X
4	Fish, benthos feeders	COLMATAGE 2008-2009-2010), and GIP Seine-Aval	X	X	X
5	Benthic predators		X	X	
6	Benthic filter feeders	PECTOW, BENTHOSEINE, and COLMATAGE projects, Le Havre port authority, and GIP Seine-Aval	X	X	
7	Benthic deposit feeders		X	X	
8	Suprabenthos	National Natural History Museum Paris, Wimereux marine station, Caen University, Le Havre port authority, and GIP Seine-Aval sampling programs		X	
9	Zooplankton	From Rybarczyk and Elkaïm (2003)		X	
10	Meiofauna	Le Guellec and Bodin (1992); Ratsimbazafy, (1998); Spilmont <i>et al.</i> (2005)		X	
11	Bacteria	Chardy (1987); Chardy and Dauvin (1992)		X	
12	Microphytobenthos	Spilmont <i>et al.</i> (2005), Napoleon <i>et al.</i> (2012)		X	
13	Phytoplankton	SURVAL database ( <a href="http://envlit.ifremer.fr/resultats/surval">http://envlit.ifremer.fr/resultats/surval</a> )		X	
14	Detritus	Seine-Aval sampling program		X	

**Appendix D.** Biomasses entered in LIM-MCMC models for all trophic compartments and for both periods, before (1996-2002) and after (2005-2012) Port2000 construction. The values of biomass are mean over the years. Data sources are presented in Appendix A. More information can be found in Tecchio et al. 2016. For the spatial boxes, see Fig.1.

	Trophic compartments	Period	Biomass (gC m <sup>-2</sup> y <sup>-1</sup> )					
			CH	FN	FS	E4	E14	E2
1	<b>OIS</b> Seabirds	1996-2002						
		2005-2012						
2	<b>FPI</b> Fish, piscivorous	1996-2002	0.019	0.060	0.057	0.143	0.103	0.264
		2005-2012	0.044	0.048	0.055	0.061	0.543	0.004
3	<b>FPV</b> Fish, planktivorous	1996-2002	0.040	0.037	0.025	0.043	0.026	0.016
		2005-2012	0.012	0.027	0.028	0.047	0.013	0.004
4	<b>FBF</b> Fish, benthos feeders	1996-2002	0.003	0.005	0.230	0.560	0.244	0.554
		2005-2012	0.088	0.327	0.252	0.307	0.387	0.358
5	<b>IPR</b> Invertebrates, predators	1996-2002	0.163	17.15	9.830	1.000	2.590	0.981
		2005-2012	1.118	26.88	41.65	1.109	14.69	2.817
6	<b>IFF</b> Invertebrates, filter feeders	1996-2002	0.442	1.897	22.32	3.08	3.81	12.71
		2005-2012	1.963	4.340	5.262	9.425	3.123	10.82
7	<b>IDF</b> Invertebrates, deposit feeders	1996-2002	0.052	1.587	3.140	6.134	6.134	4.255
		2005-2012	1.420	1.466	0.865	7.200	3.705	5.923
8	<b>SUP</b> Suprabenthos	1996-2002	2.208	2.030	2.010	0.101	0.101	0.101
		2005-2012	0.542	0.960	0.930	0.020	0.020	0.020
9	<b>ZOO</b> Zooplankton	1996-2002						
		2005-2012						
10	<b>NEM</b> Meiofauna	1996-2002	0.427	0.119	0.119	0.377	0.402	0.167
		2005-2012	0.427	0.119	0.119	0.377	0.402	0.167
11	<b>BAC</b> Bacteria	1996-2002						
		2005-2012						
12	<b>MPB</b> Microphytobenthos	1996-2002	0.050	1.725	1.720	0.300	0.350	0.380
		2005-2012	0.050	1.725	1.720	0.300	0.350	0.380
13	<b>PHY</b> Phytoplankton	1996-2002						
		2005-2012						