



## Vocal tract modelling in fallow deer: are male groans nasalized?

David Reby, M.T. Wyman, Roland Frey, B.D. Charlton, Jean-Pierre Dalmont, Joël Gilbert

### ► To cite this version:

David Reby, M.T. Wyman, Roland Frey, B.D. Charlton, Jean-Pierre Dalmont, et al.. Vocal tract modelling in fallow deer: are male groans nasalized?. *Journal of Experimental Biology*, 2018, 221, 10.1242/jeb.179416 . hal-02348220

**HAL Id: hal-02348220**

**<https://hal.science/hal-02348220>**

Submitted on 5 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Vocal tract modelling in fallow deer: are male groans nasalized?**

**Reby D.<sup>\*</sup>, Wyman M.T.<sup>\*‡</sup>, Frey, R.<sup>#</sup>, Charlton B.D.<sup>+</sup>, Dalmont J.P.<sup>\$</sup> and Gilbert J.<sup>\$</sup>**

<sup>\*</sup>: School of Psychology, University of Sussex, United Kingdom

<sup>‡</sup>: Department of Evolutionary Biology and Environmental Studies, University of Zurich, Switzerland

<sup>#</sup>: Leibniz Institute for Zoo and Wildlife Research (IZW), Berlin, Germany

<sup>+</sup>: Recovery Ecology, San Diego Zoo's Institute for Conservation Research, Escondido, California, United States

<sup>\$</sup>: Laboratoire d'Acoustique de l'Université du Maine – UMR CNRS, le Mans, France

correspondence: reby@sussex.ac.uk

## **ABSTRACT**

Males of several species of deer have a descended and mobile larynx, resulting in an unusually long vocal tract, which can be further extended by lowering the larynx during call production. Formant frequencies are lowered as the vocal tract is extended, as predicted when approximating the vocal tract as a uniform quarter wavelength resonator. However, formant frequencies in polygynous deer follow uneven distribution patterns, indicating that the vocal tract configuration may in fact be rather complex. We CT-scanned the head and neck region of two adult male fallow deer specimens with artificially extended vocal tracts and measured the cross-sectional areas of the supra-laryngeal vocal tract along the oral and nasal tracts. The CT data was then used to predict the resonances produced by three possible configurations, including the oral vocal tract only, the nasal vocal tract only, or combining both. We found that the area functions from the combined oral and nasal vocal tracts produced resonances more closely matching the formant pattern and scaling observed in fallow deer groans than those predicted by the area functions of the oral vocal tract only or of the nasal vocal tract only. This indicates that the nasal and oral vocal tracts are both simultaneously involved in the production of a nonhuman mammal vocalisation, and suggests that the potential for nasalization in putative oral loud-calls should be carefully considered.

## 38 INTRODUCTION

39

40 A key objective of mammal vocal communication research is to determine  
41 whether the acoustic structure of vocal signals encodes functionally relevant  
42 information. To achieve this aim in a given species it is important to  
43 understand how vocal signals are produced because any potential acoustic  
44 variation is primarily constrained by the biomechanical properties and  
45 dimensions of the caller's vocal anatomy (Fitch and Hauser, 2002). While the  
46 causal links between vocal production and acoustic variation are well  
47 established in human speech (Titze, 1989; Titze, 1994), the biomechanical  
48 and physiological sources of acoustic diversity in nonhuman animal signals  
49 remain poorly understood. Nonetheless, the generalization of the source-filter  
50 theory of human voice production (Fant, 1960) to non-human mammal vocal  
51 signals has significantly advanced our understanding of the acoustic structure  
52 of mammalian calls in light of their production mechanisms (reviewed by  
53 Taylor et al., 2016). According to this theory, voiced vocalizations are the  
54 result of a two-stage production process. First, the source signal is generated  
55 in the larynx by vocal fold vibration. The rate at which the glottis opens and  
56 closes determines the fundamental frequency (F0) of the vocalisation (the  
57 main perceptual correlate of the pitch). The source signal is subsequently  
58 filtered by the supra-laryngeal vocal tract, whose resonance frequencies  
59 shape the spectral envelope of the radiated vocalization, creating broad  
60 bands of energy called 'formants' (Fitch, 2000a). Because F0 and formants  
61 are produced independently they are subject to separate biomechanical  
62 constraints.

63

64 Recent anatomical investigations of mammal supra-laryngeal vocal tracts  
65 have revealed an extensive diversity of vocal tract morphology, with e.g.,  
66 elongated noses (Frey et al., 2007b), air sacs (Frey et al., 2007a) and  
67 descended larynges (Frey and Gebler, 2003), suggesting that vocal tract  
68 resonances are under strong selection pressures. Yet, to conclusively  
69 determine how such anatomical specialisations affect vocal production  
70 requires three-dimensional cineradiography to visualise call-synchronous  
71 internal dynamic changes in vocal tract shape and document the position of

oscillating structures during call production (Fitch, 2000b). This approach is however logistically difficult - if not impossible - to perform on large wild animals. An alternative method is to obtain precise vocal tract geometries from cadavers and use this data to generate vocal tract geometries that provide predictions of formant values for multiple vocal tract configurations. Several studies of nonhuman mammals have used this approach to predict the resonance characteristics of air spaces in the upper respiratory tract, showing good concordance with actual formant patterns in species-specific vocalisations (Adam et al., 2013; Carterette et al., 1979, 1984; Gamba et al., 2012; Gamba and Giacoma, 2006b; Koda et al., 2012; Riede et al., 2005). Studies investigating the evolutionary origins of speech have also used vocal tract models to predict the potential articulatory abilities of human ancestors (Boë et al., 2002; Lieberman et al., 1972) as well as non-human primates (Boë et al., 2002; Boë et al., 2017; Fitch et al., 2016). However, attempts at predicting vocal tract resonances from anatomical data remain scant and largely focused on primate species (Gamba et al., 2012; Gamba and Giacoma, 2006a; Koda et al., 2012; Riede et al., 2005), essentially due to the lack of data on vocal tract geometries available for nonhuman terrestrial mammals.

During the autumn breeding season, male fallow deer (*Dama dama*) produce high rates of sexually selected groan vocalisations (Briefer et al., 2010; McElligott and Hayden, 1999). Groans are characterized by a very low F0 and unevenly spaced and modulated formants that obey stereotyped distribution patterns, indicating that they are produced by a consistent, but complex vocal tract shape (Reby et al., 1998; Vannoni and McElligott, 2007). Fallow bucks have a descended and mobile larynx that is retracted towards the sternum during groan production (Fitch and Reby, 2001; McElligott et al., 2006), thereby extending the vocal tract. The effect of vocal tract extension on formant frequencies has been extensively documented in fallow deer (McElligott et al., 2006) and the closely related red deer (Fitch and Reby, 2001; Frey et al., 2012). As the animal extends its vocal tract, formants are lowered until they reach a minimal plateau corresponding to maximal extension, and formant frequency spacing is inversely correlated with the



length of the vocal tract during extension (Fitch and Reby, 2001; McElligott et al., 2006).

Previous attempts at relating formant frequency spacing to vocal tract length have typically modelled the vocal tract as a simple tube of uniform diameter that is closed at the glottis and opened at the mouth (Charlton et al., 2011; Reby and McComb, 2003; Vannoni and McElligott, 2007; Fitch, 1997). Under these assumptions, the length of the vocal tract can be predicted from the formant frequency spacing (and vice-versa) using the equation:  $eVTL = c/2 \cdot DF$ , where  $eVTL$  = estimated vocal tract length,  $c$  = the speed of sound in the vocal tract of 350 m/s, and  $DF$  is the overall formant frequency spacing measured in the vocalization (Reby and McComb, 2003). Measurements of anatomical oral vocal tract length in adult male fallow deer (taken as the distance from the larynx to the tip of the snout) based on calibrated photographs have produced fully extended vocal tract lengths ranging between 46 cm and 54 cm (McElligott et al., 2006). Yet, the minimum formant spacing ( $DF$ ) measured in male fallow deer groans varies between 326 Hz to 281 Hz (Vannoni and McElligott, 2007), which, when the vocal tract is modelled as a simple cylindrical tube, corresponds to vocal tract lengths between 54 cm and 62 cm. This overestimation of the vocal tract length from the acoustic data indicates that the animal's vocal tract produces more/lower formants than expected from its anatomical length. Combined with the observation that formants are stereotypically unevenly spaced in groans (McElligott et al., 2006), this suggests the vocal tract configuration during groan production is more complex than previously assumed.

Here we investigate the hypothesis that male fallow deer simultaneously use the oral and nasal vocal tracts as resonating systems during call production, allowing this species to produce a larger number of formants and a more complex formant pattern than predicted by a simple cylindrical tube model. To this end, we performed computed tomography (CT) scans of head-and-neck specimens of male fallow deer to achieve a more detailed description of the complex anatomical structure of the male fallow deer supra-laryngeal vocal tract. The specific aims of this investigation are threefold: 1) to describe the 3-

dimensional geometry of the supra-laryngeal vocal tract in male fallow deer, including the oral and nasal vocal tracts, while the larynx is maximally retracted; 2) to predict the resonance patterns produced by the fully extended vocal tract configuration, simulating the effect of the involvement of the oral vocal tract only, of the nasal vocal tract only, or of the combined oral and nasal vocal tracts on formant patterns; and 3) to compare resonance patterns predicted by each of these configurations with formant patterns observed in actual vocalisations.

## **MATERIALS AND METHODS**

A glossary of the anatomical terms used in this article is provided in the appendix and a schematic representation of the mammalian vocal anatomy is provided in the Electronic Supplementary Material.

### **Specimen collection**

Our measures of vocal tract area functions are based on CT scans from two adult male fallow deer. Male 1 was a 7-year-old buck that died from injuries sustained during a fight with rival males during the breeding season on Oct 20, 2011 in Home Park, London, UK. Male 2 was an 11-year-old buck culled by park staff during annual population control management practices on February 7, 2011 in Richmond Royal Park, London, UK. Both specimens had similar skeletal dimensions: lower mandible length (male 1: 21.0 cm, male 2: 20.3 cm) and lower hind leg length from the calcaneal tuber to the end of the metatarsus (male 1: 31 cm, male 2: 32.5 cm). Head-and-neck specimens of both individuals were obtained by separation between the 2nd and 3rd ribs. Specimens were chilled with ice within 2 hours of death and were frozen at -20° C within 5 hours of death.

### **Specimen preparation and CT-Scanning**

Specimens were thawed, and water was flushed through the oral and nasal vocal tracts to remove debris and fluids and left upright to drain for about 30 minutes before scanning. Specimens were scanned with and without artificially extending the vocal tract. Vocal tract extension was achieved by

maximally pulling the sternothyroid muscle and the trachea towards the sternum and fastening them to the sternum using a string. This recreated a naturalistic configuration of the fully extended vocal tract in male fallow deer.

Specimens were positioned as much as possible in configurations typical of vocalising (see figure 1), although it was not possible to stretch the neck as much as desired (figure 2). The oral cavity was kept open using a block of Styrofoam. This affected the position of the tongue which was pushed backward in an unnatural position in one of the specimens. The detrimental effect on the cross-sectional area measures was moderated by extrapolating the typical position of the tongue from other specimens and anatomical examinations.

#### **Estimation of vocal tract area function**

We measured the supra-laryngeal vocal tract area functions (the cross-sectional area at 1 cm steps along the length of the vocal tract, DeBoer and Fitch 2009) using the 3D curved Multi-Planar Representation viewer in Osirix (version 6.0, 64bits for Mac, [www.osirix-viewer.com](http://www.osirix-viewer.com)) and following the three-step method described in Kim et al. (2009). First, the oral and nasal vocal tract dorso-ventral midlines were drawn manually using the “3D curved path” tool on a midsagittal section (figure 3). Second, for each vocal tract, cross-sectional areas orthogonal to the midline were produced at 1 cm intervals from the glottis to the lips or nostrils (figure 3). The vocal tract area was then measured in each cross-sectional slice using the closed polygon selection tool to delineate the VT area. Osirix automatically returned the area of the delineated zone (in cm<sup>2</sup>). Each slice/measure was saved as an image file.

#### **Prediction of vocal tract resonances**

The position and amplitude of vocal tract resonances were predicted using the transfer matrix method (Chaigne and Kergomard, 2016), where the tube geometry is approximated as a series of cylindrical elements with variable

cross sections and 1 cm in length. The transfer matrix model is commonly used for modelling acoustic propagation in tube lattices. The method is appropriate when one dimension of the tube is substantially larger than the others, as in most mammalian vocal tracts: below a cut-off frequency  $F_{co}$ , ( $F_{co} \approx \frac{c}{2D}$ , where  $c$  is the speed of sound and  $D$  the largest transverse dimension) it can be assumed that only plane waves propagate. In this study  $F_{co}$  is 4000 Hz as  $D$  remains below 4 cm in the fallow deer vocal tract. Assuming linear propagation, the internal acoustic field is perfectly defined by scalar quantities in the frequency domain: the complex amplitudes of the acoustic pressure  $P$  and that of the acoustic volume velocity  $U$ . For a given cylindrical tube portion of length  $L$  and cross sectional area  $A_o$ , a 2x2 matrix of complex elements relates vectors  $\{P, U\}$  on both sides (input, output) of the cylinder:

$$\begin{pmatrix} P_{in} \\ U_{in} \end{pmatrix} = \begin{pmatrix} \cos kL & i Z_c \sin kL \\ i Z_c^{-1} \sin kL & \cos kL \end{pmatrix} \begin{pmatrix} P_{out} \\ U_{out} \end{pmatrix},$$

where  $Z_c = \frac{\rho c}{A_o}$  is the characteristic impedance (where  $\rho$  and  $c$  are respectively the air density and the speed of sound at body temperature, and  $k = \frac{\omega}{c} = \frac{2\pi F}{c}$  is the wave number).

For complex cylindrical geometry, the tube is broken down into smaller elementary cylindrical tubes. If the tube is described by  $N$  elementary cylindrical tubes, acoustic pressure and volume flow at the input and output of the tube are related by the matrix equal to the product of the  $N$  matrices of the elementary cells:

$$\begin{pmatrix} P_{in} \\ U_{in} \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} P_{out} \\ U_{out} \end{pmatrix}$$

It is thus possible to calculate the input impedance  $Z_{in}$ :

$$Z_{in} = \frac{P_{in}}{U_{in}} = \frac{A P_{out} + B U_{out}}{C P_{out} + D U_{out}} = \frac{A Z_{out} + B}{C Z_{out} + D},$$

where  $Z_{out} = \frac{P_{out}}{U_{out}}$  is the terminal impedance of the tube. If the tube is open (at the mouth and nostrils for example),  $Z_{out}$  is the radiation impedance.

235 If several complex tubes are connected (for example at the branching point of  
236 the oral and nasal tracts), there is continuity of the acoustic pressure and  
237 conservation of the acoustic volume flow. The particular frequencies where  
238 the input impedance magnitude reaches a local maximum, correspond to the  
239 vocal tract resonances visible as formants in the spectral acoustic structure of  
240 the produced vocalisation.

241 The input parameters are the area functions of the oral and nasal vocal tracts  
242 and the branching point of the tracts. The air temperature inside the fallow  
243 deer vocal tract, which affects the absolute frequency of predicted resonances  
244 (formant frequencies are proportional to the speed of sound in the air and thus  
245 to the square root of the absolute temperature) but not their relative frequency  
246 distribution, was set to 38°C. The model assumed wall damping of rigid walls  
247 and radiation impedance of an open end, un-flanged tube (Chaigne and  
248 Kergomard, 2016).

249

250 Resonances were predicted for each male, based on the area functions of  
251 three possible configurations:

252 (1) the oral vocal tract only (common laryngopharyngeal tract, oropharynx and  
253 oral cavity),

254 (2) the nasal vocal tract only (common laryngopharyngeal tract, nasopharynx  
255 and nasal cavities),

256 (3) both the oral and nasal vocal tracts (common laryngopharyngeal tract,  
257 oropharynx, oral cavity, nasopharynx, nasal cavities).

258

259 The total cross-sectional areas of both (left and right) nasal cavities were  
260 approximated by doubling the cross-sectional areas measured for the left  
261 nasal cavity connecting to the left nostril. To approximate the potential effect  
262 of the partial opening of the sides of the mouth, we only included half of the  
263 portion of the oral cavity that is laterally open. The cross-sectional areas for  
264 this portion of the oral cavity were estimated by manually linking the upper lip  
265 to the lower lip.

266

267

268 **RESULTS**

269 **Vocal tract anatomy**

270 The potential complete vocal tract is composed of five distinct and connected  
271 sections, one of which is paired: 1) the common laryngopharyngeal tract,  
272 shared by the oral and nasal vocal tract, between the glottis and the intra-  
273 pharyngeal ostium, 2) the oropharynx, from the intra-pharyngeal ostium to the  
274 Isthmus faucium (oropharyngeal tract), 3) the oral cavity, from the Isthmus  
275 faucium to the mouth opening (oral tract), 4) the nasopharynx, from the intra-  
276 pharyngeal ostium to the choanae (nasopharyngeal tract), 5) the paired nasal  
277 cavities, from the choanae to the nostrils (nasal tract). The oral vocal tract  
278 comprises parts 1,2,3 and the nasal vocal tract parts 1,4,5 (figure 2). The  
279 choanae and the Isthmus faucium are located approximately at the same  
280 distance from the glottis.

281  
282 In the resting configuration (figures 1A, 2A and 4A, B), the pharynx, soft  
283 palate and thyrohyoid ligament are relaxed, and the larynx resides at the level  
284 of cervical vertebrae 2 and 3, however farther rostrally in male 2 than in male  
285 1. Accordingly, the rostral end of the trachea is located 41 cm from the lips in  
286 male 1 and 34 cm in male 2. The resting oral vocal tract lengths (glottis to lips)  
287 are 36 cm in male 1 and 30 cm in male 2, and the resting nasal vocal tract  
288 lengths (glottis to nostrils) are 40 cm in male 1 and 35 cm in male 2 (figures  
289 4A, B and 5). The flexible region between the rostral edge of the thyroid  
290 cartilage and the base of the epiglottis is relaxed and short (4 cm in both male  
291 1 and male 2). Correspondingly, the overall length of the larynx (from cricoid  
292 arch to epiglottal tip) is 13 cm in male 1 and 10 cm in male 2. The hyoid  
293 apparatus is folded and the distance between the basihyoid and the epiglottis  
294 is small (4.5 cm in male 1 and 2.5 cm in male 2). The rostral edge of the intra-  
295 pharyngeal ostium (the caudal tip of the palatine velum) is located about 30  
296 cm from the lips in male 1 and 25 cm in male 2. The epiglottis is in contact  
297 with the intra-pharyngeal ostium or overlaps its rostral edge, so that the  
298 laryngeal entrance comes to lie in continuation of the nasopharynx (so-called  
299 'intranarial position'). The mouth is mostly closed.

300

301 In the extended phonatory configuration (figures 1B, 2B, 3A, 3D and 4C, D),  
302 the pharynx, soft palate and thyrohyoid ligament are maximally extended, and

the larynx resides at the level of cervical vertebrae 4, 5 and 6 in male 1 and 3, 4 and 5 in male 2. Accordingly, the rostral end of the trachea is located approximately 54 cm from the lips in male 1 and 44 cm in male 2. The extended oral vocal tract lengths (glottis to lips) are 48 cm in male 1 and 40 cm in male 2, and the extended nasal vocal tract lengths (glottis to nostrils) are 50 cm in male 1 and 43 cm in male 2 (figures 4C, D and 5). The lengths of the common laryngopharyngeal tract (glottis to intra-pharyngeal ostium) are 12 cm and 9 cm for males 1 and 2, respectively (figures 4C, D and 5). The flexible region between the rostral edge of the thyroid cartilage and the base of the epiglottis is maximally tensed and considerably elongated (7.5 cm in male 1 and 5 cm in male 2). Correspondingly, the overall length of the larynx (from cricoid arch to epiglottal tip) is 15 cm in male 1 and 13 cm in male 2. The hyoid apparatus is maximally unfolded and the thyrohyoid rotated caudally. The distance between the basihyoid and the epiglottis has considerably enlarged, being now 12 cm in male 1 and 7.5 cm in male 2. The rostral edge of the intra-pharyngeal ostium (the caudal tip of the palatine velum) is located approximately 37 cm from the lips in male 1 and 31 cm in male 2. The epiglottis is retracted from the intra-pharyngeal ostium so that the laryngeal entrance is in continuation of both the oropharynx and the nasopharynx. From the intra-pharyngeal ostium onward, the pharyngeal cavity splits into two tubes (the nasopharynx connected to the nasal cavities and the oropharynx connected to the oral cavity) completely separated by the soft palate (velum). The mouth is opened for vocalizing.

326

327

### 328 **Cross-sectional areas**

329 The cross sectional areas measured along each of the male specimen's vocal  
330 tracts are shown in figure 5. The area functions from the glottis (right) towards  
331 the lips and nostrils (left) are highly comparable between the two specimens.  
332 Longitudinally, the choanae and the Isthmus faucium are located about 25-30  
333 cm (male 1) and 20-25 cm (male 2) rostral to the glottis, respectively (figure  
334 5).

335 The decrease of the cross-sectional area from the glottis towards the intra-  
336 pharyngeal ostium is a consequence of the large larynx of male fallow deer.

337 Its considerable dorsoventral height causes a relatively large intra-laryngeal  
338 cross-sectional area at the glottis (figure 3D cross-section 1). From the intra-  
339 pharyngeal ostium the cross-sectional area of the nasopharynx increases  
340 towards the choanae and, similarly, the cross-sectional area of the  
341 oropharynx increases towards the Isthmus faucium. Choanae and Isthmus  
342 faucium mark the rostral end of the pharynx, i.e. the transition from the  
343 nasopharynx to the nasal cavities and from the oropharynx to the oral cavity,  
344 respectively. The gradual, mostly uniform increase in cross-sectional area,  
345 from the intra-pharyngeal ostium towards the rostral end of the pharynx,  
346 reflects the basic funnel-shape of the pharynx, narrowest at its connection to  
347 the larynx and widest at its connection to the skull and the oral cavity. From  
348 the choanae to the nostrils there is an overall decrease of cross-sectional  
349 area as a consequence of the narrowing of the nasal cavities towards the  
350 muzzle. The particular decrease of cross-sectional area at around 42 cm in  
351 male 1 and at around 37 cm in male 2 comes from the ventral nasal conchae,  
352 which narrow the nasal cavities by their extensive, scrolled osseous lamellae  
353 (figure 3D cross-section 3). From the Isthmus faucium to the lips the cross-  
354 sectional area initially increases and then decreases. In between is a  
355 particular decrease at around 39 cm in male 1 and around 33 cm in male 2.  
356 The initial increase represents the caudal end of the oral cavity between the  
357 Isthmus faucium and the root of the tongue and the final decrease the rostral  
358 end of the oral cavity between the lingual fossa and the lips. The intermediate  
359 decrease in cross-sectional area results from the lingual torus, an elevation of  
360 the tongue in ruminants, which considerably narrows the middle oral cavity.  
361 When the mouth is opened and the lower jaw depressed for groan emission  
362 the cross-sectional areas of the (then funnel-shaped) oral cavity will increase  
363 accordingly in direction towards the lips.

364

365

### 366 **Predicted Vocal tract resonances**

367 The vocal tract resonances predicted from the vocal tract area functions of  
368 male 1 and male 2 and corresponding to the three possible configurations are  
369 presented in figure 6. F1 predicted by the combined oral and nasal vocal  
370 tracts is in intermediate position between the F1 predicted using the oral vocal



tract only, or the nasal vocal tract only. F2 and F3 predicted by the combined oral and nasal vocal tracts corresponds to the F2 predicted by the nasal vocal tract only, and to the F2 predicted by the oral vocal tract only. F4 and F5 predicted by the combined oral and nasal vocal tracts corresponds to the F3 predicted by the nasal vocal tract only, and to the F3 predicted by the oral vocal tract only (figure 6, table 1).

The centre frequencies of each predicted formant are reported in table 1. The models using both oral and nasal vocal tracts predict much lower formants overall (average formant spacing of 255 Hz) than models using the oral vocal tract only (average formant spacing of 446 Hz) or models using the nasal vocal tract only (average formant spacing of 358 Hz).

**Table 1: Predicted centre frequencies for formants F1 to F5 (Hz) and estimated formant spacing (DF) for the different vocal tract configurations for males 1 and 2.**

configuration	male	F1	F2	F3	F4	F5	DF
oral vocal tract only	1	249	494	1044	1352	1938	410
	2	305	581	1238	1629	2238	482
nasal vocal tract only	1	175	359	771	1166	1534	328
	2	194	505	981	1426	1712	388
combined oral and nasal vocal tracts	1	223	365	495	776	1060	227
	2	255	503	582	992	1302	283

**Comparison with acoustic observations**

Figure 7 plots the average centre frequencies of the first five formants observed in groans from 16 adult fallow deer males (reported in Vannoni and McElligott, 2007, see table 2) against the resonances predicted by our three

vocal tract models. The resonances predicted by a vocal tract including both the oral and nasal vocal tracts are a better fit to the observed formants than those predicted by using the oral vocal tract or the nasal vocal tract only: the slope of the regression line is closer to 1 (indicating a better fit of the scaling of the resonances), and  $R^2$  is also higher (indicating a better fit of the pattern of the resonances). Examination of the regression slopes in figure 7 shows that while model 3 (combined oral and nasal vocal tracts) underestimates the formant frequency spacing by 9%, model 1 (oral vocal tract only) overestimates DF by 37% and model 2 (nasal vocal tract only) overestimates DF by 23%. Separate correlations for male 1 and male 2 are given in the Electronic Supplementary Material.

403

**Table 2: Average centre frequencies for formants F1 to F5 (Hz) and average estimated formant spacing (DF) at maximal vocal tract extension from 16 adult males (Vannoni & McElligott 2007).**

Formant:	F1	F2	F3	F4	F5	DF
Average centre frequency (Hz) $\pm$ SE	208.5 $\pm$ 2.0	414.3 $\pm$ 1.9	575.2 $\pm$ 3.3	1060.2 $\pm$ 2.9	1265.9 $\pm$ 3.2	300.6 $\pm$ 0.8
Minimum centre frequency (Hz)	152	329	457	966	1170	281
Maximum centre frequency (Hz)	263	496	677	1149	1371	326

407

408

## 409 DISCUSSION

410

In this study, the artificially extended vocal apparatuses of two adult male fallow deer were CT-scanned, and the cross-sectional areas of the complete supra-laryngeal vocal tract (common laryngopharyngeal tract, oropharynx,

oral cavity, nasopharynx and nasal cavities) of both specimens were measured along the oral and nasal vocal tracts. We then used this data to model resonance patterns produced by these supra-laryngeal cavities including the oral vocal tract only, the nasal vocal tract only or the combined oral and nasal vocal tracts. We found that the configuration combining the oral and nasal vocal tract geometries produced a resonance pattern which more closely matches the formants observed in fallow deer groans, both in term of formant frequency pattern and in term of formant frequency scaling.

422

The formants observed in the groans of fallow deer (Briefer et al., 2010; McElligott and Hayden, 1999) and more generally in the sexually-selected calls of male polygynous deer with extensible vocal tracts (Fitch and Reby, 2001; Passilongo et al., 2013; Reby and McComb, 2003; Reby et al., 2016) obey stereotyped, uneven formant patterns incompatible with a vocal tract consisting of a simple linear tube. More specifically, in both fallow deer groans (McElligott et al., 2006) and red deer roars (Reby and McComb, 2003) the second and third formants are close to one another and the fourth formant is higher, leaving a gap between the third and fourth formant. Our simulations combining both the oral and nasal vocal tracts predict this pattern. Comparison of the models suggests that formants two and four of fallow deer groans are affiliated to the nasal vocal tract while formants three and five are affiliated to the oral vocal tract.

436

In terms of frequency scaling, our predictions resolve the aforementioned mismatch between apparent vocal tract lengths estimated from formant frequencies measured in fallow deer groans and actual anatomical vocal tract length derived from photogrammetric and anatomical data. Presumably this is because the inclusion of formants affiliated to the nasal vocal tract led to an overestimation of apparent vocal tract length in previous studies modelling the fallow deer vocal tract as a single uniform tube. Indeed, previous investigations of apparent vocal tract length (i.e. vocal tract length estimated from formant frequencies) in polygynous deer with descended and mobile larynges (Corsican deer: Kidjo et al., 2008; fallow deer: McElligott et al., 2006; Mesola deer: Passilongo et al., 2013; red deer: Reby and McComb, 2003)

448 have modelled the vocal tract as a linear tube with constant cross-section  
449 closed at one end (glottis) and open at the other (mouth), and excluded the  
450 involvement of the nasal vocal tract for loud calls produced with an open  
451 mouth. While these succeeded at characterizing inter-individual differences in  
452 formant frequency spacing (Kidjo et al., 2008; Vannoni and McElligott, 2007),  
453 apparent vocal tract length and thus body size (Reby and McComb, 2003;  
454 Vannoni and McElligott, 2008), they probably yielded over-estimations of the  
455 anatomical vocal tract length.

456

457 The inclusion of nasal resonances in future models should allow for better  
458 estimations of apparent vocal tract length from recorded mating calls in these  
459 species, thereby potentially enhancing the reliability of bioacoustics tools  
460 aimed at assessing body size from vocalisations for research, conservation or  
461 wildlife management purposes.

462

463 Taken together our observations strongly suggest that the nasal cavity and  
464 oral cavity are both simultaneously involved in the vocal production of fallow  
465 deer groans. This involvement of the nasal vocal tract due to the non-closure  
466 of the intra-pharyngeal ostium during vocal tract extension maybe widespread  
467 in species with a permanently descended larynx and extensible vocal tract  
468 (such as other polygynous deer and e.g. goitred gazelles), but also occur in  
469 species where callers lower their larynx temporarily for the production of oral  
470 (rather than nasal) calls. We suggest that the potential for nasalization of  
471 putative oral loud calls should be carefully examined across terrestrial  
472 mammals.

473

474 The role of nasal cavities in acoustic output has been investigated in humans  
475 using anatomical scans, area functions, vocal tract modeling, and/or acoustic  
476 analysis (Dang et al., 1994; Feng and Castelli, 1996; Hattori and Fujimura,  
477 1958; Pruthi et al., 2007; Story, 2005). Compared to modulation of the oral  
478 vocal tract, nasalization plays a relatively minor role in human speech  
479 variation and is often left out of vocal models. However, models that include  
480 coupling between the nasal and oral cavity can result in transfer functions that  
481 more closely match recorded acoustic output (Dang et al., 1994; Feng and

482 Castelli, 1996). The effects of nasalization are strongest in the lower  
483 frequencies (Feng and Castelli, 1996; Pruthi et al., 2007) and include the  
484 addition of low frequency formant peaks as observed here in fallow deer  
485 groans. Nasal coupling has also been suggested as a likely mechanism for  
486 the addition of low spectral peaks in Diana monkey alarm calls (Riede and  
487 Zuberbuhler, 2003).

488

489 There are obvious limitations to this investigation. Our dead specimens were  
490 scanned in artificial positions constrained by the dimensions of the CT-  
491 scanner, and thus only approximate the natural postures of live animals  
492 during vocalising. Moreover, the vocal tracts were artificially extended. The  
493 geometries are thus approximations of the vocal tract of live animals during  
494 vocalizing, and do not account for internal adjustments such as, e.g., the  
495 possible contribution of palatopharyngeal muscles. Future investigations could  
496 involve performing several scans involving different combinations of varying  
497 head/neck angle, extent of the laryngeal descent, or mouth opening; or  
498 perform simulations of these parameters (Gamba et al., 2012; Gamba and  
499 Giacomini, 2006b). Using a larger sample of specimens would also allow the  
500 assessment of inter-individual variation, including the effect of age or size.

501

502 Formants frequencies are known to provide cues to the caller's body size in  
503 fallow deer groans and red deer roars, due to a close correlation between  
504 formant frequency spacing and body size (McElligott et al., 2006; Reby and  
505 McComb, 2003), and are used by male and female receivers to assess rivals  
506 and potential mates during the breeding season (Charlton et al., 2008a;  
507 Charlton et al., 2007; Charlton et al., 2008b; Pitcher et al., 2015; Reby et al.,  
508 2005). The descended larynx and extensible vocal tract of fallow and red deer  
509 males (and some other species) are therefore considered to be adaptations  
510 that allow callers to maximise the acoustic impression of their body size  
511 conveyed to receivers (the "size exaggeration hypothesis" Fitch and Reby,  
512 2001; Ohala, 1984). Our investigations show that the involvement of the nasal  
513 vocal tract adds additional formants to the lower part of the spectrum, which  
514 increases formant density (decreasing formant spacing), and may make the  
515 caller sound larger when compared to oral or nasal only calls. Similar

functional explanations have been suggested for the evolution of air sacs (de Boer, 2009; Harris et al., 2006), which also increase formant density by adding resonances.

In conclusion, we contend that, while expensive and technically challenging, using 3-dimensional CT scanning to predict vocal tract resonances can greatly assist the interpretation of formant patterns in mammal vocalizations. We suggest that similar approaches could be generalized to the study of vocal tract resonances in other terrestrial mammals.

## **Appendix: Glossary of Anatomical Terms used in this paper (based on Constantinescu & Schaller 2012 and own descriptions)**

**Basihyoid:** unpaired, most ventral, transverse component of hyoid apparatus intercalated between the paired suspension from the skull and the paired arms to the larynx.

**Choanae:** the 'internal nares', i.e. the two openings at the caudal end of the nasal cavity, where the paired nasal meatuses lead into the nasopharynx.

**Cricoid cartilage:** the most caudal, ring-shaped cartilage of the larynx that is attached to the trachea; it consists of a dorsal plate and a ventral arch.

**Epiglottis:** the most rostral cartilage of the larynx; during quiet breathing it has a so-called intranarial position, i.e. it protrudes dorsally, through the intra-pharyngeal ostium, into the nasopharynx; during an open-mouth call the larynx is withdrawn from the intra-pharyngeal ostium so that it is now positioned in the oropharynx.

**Glottis:** forms the vocal source of the larynx; it consists of the two vocal folds, ventral parts of the arytenoid cartilages and the vocal cleft in between the vocal folds; regarding the laryngeal cavity it is positioned between the laryngeal vestibule rostrally, and the infraglottic cavity caudally.

**Hyoid apparatus:** a framework of small rod-like bones connecting dorsally to the base of the skull, rostrally to the tongue, and caudally to the larynx; it consists of 3 components: the arms of the paired dorsal part flank the pharynx on both sides and suspend the entire hyoid apparatus from the skull base, it

550 consists of several parts on both sides termed (dorsal to ventral):  
551 tympanohyoid, stylohyoid, epihyoid, ceratohyoid; the arms of the paired  
552 caudal part connect to the larynx (in fallow deer via the two thyrohyoid  
553 ligaments); it consists of one element per side termed thyrohyoid; the  
554 unpaired, transverse ventral part connects the two paired structures thereby  
555 forming a larger U-shaped fork for suspension from the skull dorsally and a  
556 smaller U-shaped fork for connection to the larynx caudally.

557 **Intra-pharyngeal ostium:** opening in the soft palate, creating a passage  
558 between nasopharynx and oropharynx; it is bordered by the palatopharyngeal  
559 muscle that can constrict the intra-pharyngeal ostium.

560 **Isthmus faucium:** narrow short passage between the mouth cavity and the  
561 oropharynx, bounded by the soft palate dorsally, the tongue ventrally, and the  
562 palatoglossal arch (a symmetrical, dorsoventral mucosa fold between soft  
563 palate and tongue) laterally.

564 **Laryngeal entrance:** rostral entrance to the larynx, bounded by the epiglottis,  
565 the aryepiglottic folds and the corniculate processes of the arytenoid  
566 cartilages.

567 **Nasopharynx:** nasal part of the pharynx, dorsal to the soft palate; it extends  
568 from the choanae to the intrapharyngeal ostium.

569 **Oropharynx:** oral part of the pharynx, ventral to the soft palate; it extends  
570 from the isthmus faucium to the base of the epiglottis.

571 **Pharynx:** musculomembraneous cross way of the respiratory and digestive  
572 tracts between the oral and nasal cavities rostrally, and the oesophagus and  
573 larynx caudally; for the sake of simplicity, and in contrast to textbooks, the  
574 pharynx is here not subdivided in 3 parts (nasal, oral and laryngeal) but only  
575 in 2 parts (nasal and oral).

576 **Soft palate:** the palatine velum or soft palate, is a soft tissue structure that is  
577 laterally fused to the pharyngeal walls; it completely separates naso- and  
578 oropharynx, except at the intra-pharyngeal ostium, which represents the only  
579 communication between Naso- and Oropharynx; as the pharynx, it extends  
580 from the choanae to the larynx.

581 **Thyrohyoid:** paired caudal element of the hyoid apparatus that, together with  
582 the basihyoid, forms the smaller U-shaped fork for establishing the connection  
583 between the hyoid apparatus and the larynx.

584 **Thyrohyoid ligament:** replaces the usual thyrohyoid articulation of most  
585 mammals by a ligamentous connection between the caudal tip of the  
586 thyrohyoid bone (of the hyoid apparatus) and the rostral horn of the thyroid  
587 cartilage.

588 **Thyroid cartilage:** The most superficial and largest cartilage of the larynx,  
589 unpaired, its two lateral laminae are ventrally fused and enclose most of the  
590 laryngeal cavity in between them; its rostral horn connects the larynx to the  
591 thyrohyoid of the hyoid apparatus, its caudal horn articulates with the cricoid  
592 cartilage in the cricothyroid articulation.

593 **Trachea:** Windpipe, connecting the lungs to the larynx, extends from its  
594 bifurcation into the main bronchi caudally to the cricoid cartilage of the larynx  
595 rostrally.

596

## 597 **Acknowledgements**

598 We thank John Bartram, John Comfort, and Paul Douglas at the London  
599 Royal Parks for their help obtaining the specimens. We are also very grateful  
600 to Jan Bush and the amazing staff at CISC for facilitating the production of the  
601 CT-Scans.

602

## 603 **Competing interests**

604 The authors declare no competing or financial interests.

605

## 606 **Author contributions**

607 D.R. supervised the research. M.T.W., R.F. and D.R. performed the  
608 anatomical and imaging investigations. J.G., J-PD. and D.R. performed the  
609 acoustical modelling. D.R., M.T.W., R.F., B.D.C., J-PD. and J.G. wrote the  
610 manuscript.

611

## 612 **Funding**

613 J.G. was supported by a Projet International de Coopération Scientifique grant  
614 reference 6188 from Centre National de la Recherche Scientifique. DR was  
615 supported by an invited professorship from Le Mans Université. MW was



616 supported by a grant from US National Science Foundation International  
617 Research Fellowship (grant number 0908569) and an award from the  
618 Systematics Research Fund.

619

#### 620 **Data availability**

621 Supporting data and scripts for predicting vocal tract resonances are available  
622 on request from the corresponding author.

623

#### 624 **Figure Legends**

625

626 **Figure 1. Fallow deer buck in a pre-groaning posture (A) and in a**  
627 **groaning posture (B) during the annual rut.** In (A) the neck is not extended,  
628 the larynx is in its resting position and the mouth is closed. In (B) the neck is  
629 extended, the larynx retracted, and the mouth is opened (no lip rounding).  
630 Single frames from a video by D. Belton ©.

631

632 **Figure 2. CT-based 3-D reconstructions of male 1.** (A): larynx and hyoid  
633 apparatus in pre-groan (resting) position. (B): larynx and hyoid apparatus in a  
634 groaning position at full artificial extension of the vocal tract. The larynx  
635 position is shifted caudally by approximately 12 cm. ochre: common  
636 laryngopharyngeal tract; green: oropharynx and oral cavity; blue: nasopharynx  
637 and nasal cavities.

638

639 **Figure 3. Multi-Planar Representation of a CT scan of male 1 illustrating**  
640 **the methodology for measuring the vocal tract area function at maximal**  
641 **artificial extension.** (A): manually drawn dorso-ventral and transverse  
642 midline through the oral vocal tract; (B): sampling transverse sections along  
643 the virtually stretched, un-curved oral vocal tract in 1 cm steps; (C): three  
644 examples of the obtained cross-sections highlighting the respective contours  
645 of the oral vocal tract in green. (D): Sagittal section of the oral vocal tract (left,  
646 midline in red), the nasal vocal tract (right, midline in green) and  
647 representative cross-sections at the level of the glottis (1), of the naso- and  
648 oro-pharynx separated by the soft palate (2) and of the nasal cavities (3).

649

650 **Figure 4. Midsagittal slices of CT scans of males 1 and 2 with the larynx**  
651 **and vocal tract in a pre-groaning (resting) position (A and B) and in a**  
652 **groaning position with the larynx maximally retracted and the vocal tract**  
653 **fully extended (C and D).** C1 – C7, T1 = cervical vertebrae 1-7, first thoracic  
654 vertebra; Cart. aryt. = arytenoid cartilage; Cart. cric. = cricoid cartilage; Cart.  
655 thyr. = thyroid cartilage; Cav. oris = oral cavity; Choan. = choanae (caudal  
656 nasal apertures, internal nares); Epigl. = epiglottis; Glott. = glottis; Ling. =  
657 tongue; Nasophar. = nasal part of pharynx; Nostr. = nostrils (rostral nasal  
658 apertures, external nares); Orophar. = oral part of pharynx; Ost. intrphar. =  
659 intra-pharyngeal ostium (IPO); Palat. dur. = hard palate; Palat. mol. = soft  
660 palate ('velum'); Trach. = trachea; o = rostral tip of sternal manubrium;  
661 asterisk: position of the basihyoid. Compared to the resting state, the  
662 elongation of the oral and nasal vocal tract is 33% in male 1 (A, C) and 25% in  
663 male 2 (B, D). Scale: 10 cm

664  
665 **Figure 5.** Estimation of oral and nasal vocal tract cross-sectional areas in the  
666 maximally extended state, proceeding from right to left, in male 1 (A) and  
667 male 2 (B).

668  
669 **Figure 6. Predicted resonances for the three possible vocal tract**  
670 **configurations of groan production in male 1 (A) and male 2 (B).** Red  
671 lines: combined oral and nasal vocal tracts; green lines: nasal vocal tract only;  
672 blue lines: oral vocal tract only. The resonance pattern produced by the  
673 combined oral and nasal vocal tracts is more similar to the observed formants  
674 in fallow deer groans than the resonance patterns of either oral or nasal vocal  
675 tract alone.

676  
677 **Figure 7. Correlations between resonances observed in male fallow deer**  
678 **groans (y axis) and resonances predicted from vocal tract geometries of**  
679 **scanned specimens (x axis).** Predicted resonances are the average centre  
680 frequencies of the first 5 peaks predicted by the cross-sectional areas of the  
681 vocal tract of male 1 and 2 including the oral vocal tract only (blue line), the  
682 nasal vocal tract only (green line), or both the oral and nasal vocal tracts (red  
683 line). The regression slopes inform the fit of the scaling (formant spacing) of

the observed resonances to the predicted resonances. The values of  $R^2$  provide the fit of the pattern of the observed resonances to the predicted resonances.

## References

- Adam, O., Cazau, D., Gandilhon, N., Fabre, B., Laitman, J. T. and Reidenberg, J. S. (2013). New acoustic model for humpback whale sound production. *Applied Acoustics* **74**, 1182-1190.
- Boë, L.-J., Heim, J., Honda, K. and Maeda, S. (2002). The potential Neandertal vowel space was as large as that of modern humans. *Journal of phonetics* **30**, 465-484.
- Boë, L.-J., Berthommier, F., Legou, T., Captier, G., Kemp, C., Sawallis, T. R., Beckers, Y., Rey, A. and Fagot, J. (2017). Evidence of a vocalic proto-system in the baboon (*Papio papio*) suggests pre-hominin speech precursors. *PLoS ONE* **12**, e0169321.
- Briefer, E., Vannoni, E. and McElligott, A. G. (2010). Quality prevails over identity in the sexually selected vocalisations of an ageing mammal. *BMC Biology* **8**, 1-15.
- Carterette, E. C., Shipley, C., and Buchwald, J. (1979). Linear prediction theory of vocalization in cat and kitten, in *Frontiers in Speech Communication Research*, edited by B. Lindblom, and S. Ohman (Academic Press, New York), pp. 245-257.
- Carterette, E. C., Shipley, C., and Buchwald, J. S. (1984). On synthesizing animal speech: The case of the cat," in *Electronic speech synthesis : techniques, technology, and applications*, edited by G. Bristow (McGraw-Hill, New York), pp. 292-302.
- Chaigne, A. and Kergomard, J. (2016). *Acoustics of musical instruments* New York: Springer.
- Charlton, B., Ellis, W., McKinnon, A., Cowin, G., Brumm, J., Nilsson, K. and Fitch, W. (2011). Cues to body size in the formant spacing of male koala (*Phascolarctos cinereus*) bellows: honesty in an exaggerated trait. *Journal of Experimental Biology* **214**, 3414-3422.
- Charlton, B., McComb, K. and Reby, D. (2008a). Free-ranging red deer hinds show greater attentiveness to roars with formant frequencies typical of young males. *Ethology* **114**, 1023-1031.
- Charlton, B., Reby, D. and McComb, K. (2007). Female red deer prefer the roars of larger males. *Biology Letters* **3**, 382-385.
- Charlton, B., Reby, D. and McComb, K. (2008b). Effect of combined source (F0) and filter (formant) variation on red deer hind responses to male roars. *Journal of the Acoustical Society of America* **123**, 2936-2943.
- Constantinescu, G.M., Schaller, O. (2012). *Illustrated Veterinary Anatomical Nomenclature*, 3<sup>rd</sup> revised edition, Enke Verlag, Stuttgart, 620 pp.
- Dang, J. W., Honda, K. and Suzuki, H. (1994). Morphological and acoustical analysis of the nasal and the paranasal cavities. *Journal of the Acoustical Society of America* **96**, 2088-2100.

730 **de Boer, B.** (2009). Acoustic analysis of primate air sacs and their effect  
731 on vocalization. *The Journal of the Acoustical Society of America* **126**, 3329–3343.

732 **Fant, G.** (1960). Acoustic Theory of Speech Production. The Hague:  
733 Mouton.

734 **Feng, G. and Castelli, E.** (1996). Some acoustic features of nasal and  
735 nasalized vowels: A target for vowel nasalization. *Journal of the Acoustical Society*  
736 *of America* **99**, 3694-3706.

737 **Fitch, W. T.** (1997). Vocal tract length and formant frequency dispersion  
738 correlate with body size in rhesus macaques. *Journal of the Acoustical Society of*  
739 *America* **102**, 1213-1222.

740 **Fitch, W. T.** (2000a). The evolution of speech: a comparative review.  
741 *Trends in Cognitive Sciences* **4**, 258-267.

742 **Fitch, W. T.** (2000b). The phonetic potential of nonhuman vocal tracts:  
743 Comparative cineradiographic observations of vocalizing animals. *Phonetica* **57**,  
744 205-218.

745 **Fitch, W. T., de Boer, B., Mathur, N. and Ghazanfar, A. A.** (2016).  
746 Monkey vocal tracts are speech-ready. *Science Advances* **2**, e1600723-e1600723.

747 **Fitch, W. T. and Hauser, M. D.** (2002). Unpacking "Honesty": Generating  
748 and extracting information from acoustic signals. In *Animal Communication*, eds.  
749 A. Megala-Simmons and A. Popper), pp. 65-137. Berlin: Springer-Verlag.

750 **Fitch, W. T. and Reby, D.** (2001). The descended larynx is not uniquely  
751 human. *Proceedings of the Royal Society of London, Series B: Biological Sciences*  
752 **268**, 1669-1675.

753 **Frey, R. and Gebler, A.** (2003). The highly specialized vocal tract of the  
754 male Mongolian gazelle (*Procapra gutturosa* Pallas, 1777 - *Mammalia, Bovidae*).  
755 *Journal of Anatomy* **203**, 451-471.

756 **Frey, R., Gebler, A., Fritsch, G., Nygren, K. and Weissengruber, G.**  
757 (2007a). Nordic rattle: the hoarse vocalization and the inflatable laryngeal air sac  
758 of reindeer (*Rangifer tarandus*). *Journal of Anatomy* **210**, 131-159.

759 **Frey, R., Volodin, I. and Volodina, E.** (2007b). A nose that roars:  
760 anatomical specializations and behavioural features of rutting male saiga. *Journal*  
761 *of Anatomy* **211**, 717-736.

762 **Frey, R., Volodin, I., Volodina, E., Carranza, J. and Torres-Porras, J.**  
763 (2012). Vocal anatomy, tongue protrusion behaviour and the acoustics of rutting  
764 roars in free-ranging Iberian red deer stags (*Cervus elaphus hispanicus*). *Journal*  
765 *of Anatomy* **220**, 271-292.

766 **Gamba, M., Friard, O. and Giacoma, C.** (2012). Vocal tract morphology  
767 determines species-specific features in vocal signals of lemurs (*Eulemur*).  
768 *International Journal of Primatology* **33**, 1453-1466.

769 **Gamba, M. and Giacoma, C.** (2006a). Vocal tract configurations and  
770 formant pattern variation in ruffed lemurs vocal production. In *Proceedings of*  
771 *the ICPV 2006, The 5th International Conference on Voice Physiology and*  
772 *Biomechanics, Variations across Cultures and Species.*, pp. 38-41. Tokyo, Japan.

773 **Gamba, M. and Giacoma, C.** (2006b). Vocal tract modeling in a prosimian  
774 primate: the black and white ruffed lemur. *Acta Acustica United With Acustica* **92**,  
775 749-755.

776 **Harris, T. R., Fitch, W. T., Goldstein, L. M. and Fashing, P. J.** (2006).  
777 Black and white colobus monkey (*Colobus guereza*) roars as a source of both  
778 honest and exaggerated information about body mass. *Ethology* **112**, 911-920.

779       **Hattori, S. and Fujimura, O.** (1958). Nasalization of vowels in relation to  
780 nasals. *Journal of the Acoustical Society of America* **30**, 267-274.

781       **Kidjo, N., Cargnelutti, B., Charlton, B. D., Wilson, C. and Reby, D.**  
782 (2008). Vocal behaviour in the endangered Corsican deer: description and  
783 phylogenetic implications. *Bioacoustics* **18**, 159-181.

784       **Koda, H., Nishimura, T., Tokuda, I. T., Oyakawa, C., Nihonmatsu, T.**  
785 **and Masataka, N.** (2012). Soprano singing in gibbons. *American Journal of*  
786 *Physical Anthropology* **149**, 347-355.

787       **Lieberman, P., Crelin, E. S. and Klatt, D. H.** (1972). Phonetic ability and  
788 related anatomy of the newborn and adult human, Neanderthal man, and the  
789 chimpanzee. *American Anthropologist* **74**, 287-307.

790       **McElligott, A. G., Birrer, M. and Vannoni, E.** (2006). Retraction of the  
791 mobile descended larynx during groaning enables fallow bucks (*Dama dama*) to  
792 lower their formant frequencies. *Journal of Zoology* **270**, 340-345.

793       **McElligott, A. G. and Hayden, T. J.** (1999). Context-related vocalization  
794 rates of fallow bucks, *Dama dama*. *Animal Behaviour* **58**, 1095-1104.

795       **Ohala, J. J.** (1984). An ethological perspective on common cross-language  
796 utilization of F0 of voice. *Phonetica* **41**, 1-16.

797       **Passilongo, D., Reby, D., Carranza, J. and Apollonio, M.** (2013). Roaring  
798 high and low: composition and possible functions of the Iberian stag's vocal  
799 repertoire. *PLoS ONE* **8**, e63841.

800       **Pitcher, B. J., Briefer, E. F. and McElligott, A. G.** (2015). Intrasexual  
801 selection drives sensitivity to pitch, formants and duration in the competitive  
802 calls of fallow bucks. *BMC Evolutionary Biology* **15**, 149.

803       **Pruthi, T., Espy-Wilson, C. Y. and Story, B. H.** (2007). Simulation and  
804 analysis of nasalized vowels based on magnetic resonance imaging data. *Journal*  
805 *of the Acoustical Society of America* **121**, 3858-3873.

806       **Reby, D., Joachim, J., Lauga, J., Lek, S. and Aulagnier, S.** (1998).  
807 Individuality in the groans of fallow deer (*Dama dama*) bucks. *Journal of Zoology*  
808 **245**, 79-84.

809       **Reby, D. and McComb, K.** (2003). Anatomical constraints generate  
810 honesty: acoustic cues to age and weight in the roars of red deer stags. *Animal*  
811 *Behaviour* **65**, 519-530.

812       **Reby, D., McComb, K., Cargnelutti, B., Darwin, C., Fitch, W. T. and**  
813 **Clutton-Brock, T. H.** (2005). Red deer stags use formants as assessment cues  
814 during intrasexual agonistic interactions. *Proceedings of the Royal Society of*  
815 *London, Series B: Biological Sciences* **272**, 941-947.

816       **Reby, D., Wyman, M. T., Frey, R., Passilongo, D., Gilbert, J., Locatelli, Y.**  
817 **and Charlton, B. D.** (2016). Evidence of biphonation and source-filter  
818 interactions in the bugles of male North American wapiti (*Cervus canadensis*).  
819 *Journal of Experimental Biology* **219**, 1224-1236.

820       **Riede, T., Bronson, E., Hatzikirou, H. and Zuberbuhler, K.** (2005).  
821 Vocal production mechanisms in a non-human primate: morphological data and  
822 a model. *Journal of Human Evolution* **48**, 85-96.

823       **Riede, T. and Zuberbuhler, K.** (2003). The relationship between  
824 acoustic structure and semantic information in Diana monkey alarm  
825 vocalization. *Journal of the Acoustical Society of America* **114**, 1132-1142.

826           **Story, B. H.** (2005). A parametric model of the vocal tract area function  
827 for vowel and consonant simulation. *Journal of the Acoustical Society of America*  
828 **117**, 3231-3254.

829           **Taylor, A., Charlton, B. D. and Reby, D.** (2016). Vocal production by  
830 terrestrial mammals: source, filter and function. In *Vertebrate sound production*  
831 *and acoustic communication*, eds. R. A. Suthers W. T. Fitch R. R. Fay and A.  
832 Popper). Berlin: Springer International Publishing.

833           **Titze, I. R.** (1989). Physiologic and acoustic differences between male and  
834 female voices. *Journal of the Acoustical Society of America* **85**, 1699-1707.

835           **Titze, I. R.** (1994). Principles of Voice Production. Englewood Cliffs, New  
836 Jersey: Prentice Hall.


837           **Vannoni, E. and McElligott, A. G.** (2007). Individual acoustic variation in  
838 fallow deer (*Dama dama*) common and harsh groans: a source-filter theory  
839 perspective. *Ethology* **113**, 223-234.

840           **Vannoni, E. and McElligott, A. G.** (2008). Low frequency groans indicate  
841 larger and more dominant fallow deer (*Dama dama* ) males. *PLoS ONE* **3**, e3113.  
842
















A




B







(A) Male 1



(B) Male 2

