



**HAL**  
open science

# Energy (Im-)Balance in Frugivorous Lemurs in Southern Madagascar: A Preliminary Study in *Lemur catta* and *Eulemur rufifrons x collaris*

Bruno Simmen, Hanta Rasamimanana

## ► To cite this version:

Bruno Simmen, Hanta Rasamimanana. Energy (Im-)Balance in Frugivorous Lemurs in Southern Madagascar: A Preliminary Study in *Lemur catta* and *Eulemur rufifrons x collaris*. *Folia Primatologica*, 2018, 89 (6), pp.382-396. 10.1159/000492570 . hal-02349627

**HAL Id: hal-02349627**

**<https://hal.science/hal-02349627>**

Submitted on 5 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Energy (im-)balance in frugivorous lemurs in southern Madagascar:  
2 a preliminary study in *Lemur catta* and *Eulemur rufifrons x collaris*

3  
4  
5 Bruno Simmen<sup>1</sup> and Hanta Rasamimanana<sup>2</sup>

6  
7  
8 <sup>1</sup> Centre National de la Recherche Scientifique/Muséum national d'Histoire naturelle,  
9 UMR 7206 Eco-anthropologie et Ethnobiologie

10  
11 <sup>2</sup> Ecole Normale Supérieure, Université d'Antananarivo, Antananarivo, Madagascar

12  
13  
14  
15  
16 **Short title: Energy intake in wild lemurs**

17  
18  
19 **#words: 8332**

20  
21  
22  
23  
24  
25  
26  
27 **Corresponding author:** Bruno Simmen, Muséum national d'Histoire naturelle, 1 avenue du Petit  
28 Château, 91800 Brunoy, France

29 E-mail: simmen@mnhn.fr, Phone number: +33 (0)1 60 47 92 33

30           **Abstract**

31  
32   The reproductive cycle of several lemur species is synchronized by photoperiodic changes and is  
33   attuned to the seasonal fluctuations of food supply. Irregular periods of food shortage nevertheless  
34   occur in Madagascar, which can result in negative energy balance and reduced fitness, especially in  
35   frugivorous species. Here, we test if ring-tailed lemurs (*Lemur catta*) and brown lemurs (*Eulemur*  
36   *rufifrons x collaris*) use fat stores to maintain their energy balance in a gallery forest of Madagascar  
37   (Berenty Reserve). We assessed the energy intake from weight-based estimates of food intake  
38   (mouthful count) and macronutrient contents in the diet during dry and wet months. The  
39   metabolizable energy ingested by these frugivorous/folivorous species was high during the late wet  
40   season, but was insufficient to meet their energy requirements during the dry season. These  
41   preliminary data suggest that these lemurs must store sufficient amount of fat mass during the  
42   period of food abundance to meet their maintenance requirements and to cope with the energetic  
43   burden of the mating season. Estimates based on models assuming no fibre digestibility during the  
44   dry season are consistent with the prediction that brown lemurs should supplement their daytime  
45   diet with nighttime feeding in addition to the use of body fat stores.

46

47

48

49

50           **Key words**

51           metabolic costs, seasonal breeding, energy intake, nutrients, body fat, dry season, primates

52

53

54

55

56

57

58

**Introduction**

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

Species that primarily depend on food resources scarcely or irregularly distributed in time and space like fruit, flowers and animal matter, are at risk of energy imbalance [Oates, 1987; Wright, 1999]. In tropical habitats with unpredictable rainfall and/or a long dry season, frugivorous species have developed opportunistic foraging strategies and shift their diet toward lower quality or fallback food during the lean season [Oates, 1987; Lambert and Rothman, 2015]. In some cases, dietary shifts are accompanied by morphological and physiological adjustments such as changes in the digestive tract size [Munn et al., 2009], basal metabolic rate, and other energy-saving mechanisms such as body fat storage [Muroyama et al., 2006; Canale and Henry, 2010]. Apart from its consequences on survival during the lean season, the storage of body fat could also have evolved to sustain the energy cost of reproduction (e.g., in seasonal breeding mammals or in harsh environments [Brockman and van Schaik, 2005; Canale and Henry, 2010; Garcia et al., 2011; Wells, 2012]). Seasonal body weight changes have sometimes been recorded in wild primates (e.g. [Knott 1998; Richard et al., 2000; Lewis and Kappeler 2005]), but in studies that only measure body mass, it is unclear whether the weight gain reflects fat storage or increased muscle volume [Bercovitch 1987]. It has been argued that, due to their primarily arboreal lifestyle, primates may generally have not relied on body fat storage strategies throughout their evolutionary history [Dittus, 2013]. However, the available data are still too limited to invalidate the existence of fattening in many wild primates. The seasonal increase in body fat has been reported for the first time in semi-free ranging and wild male squirrel monkeys in relation to rainfall cycles and sexual selection [Du Mond & Hutchinson 1967]. High body fat content following periods of high food supply has been reported in Bornean orangutans (*Pongo p. pygmaeus*) living in unpredictable habitats and some strepsirrhine species, especially among the small heterotherm cheirogaleids that enter torpid states during the dry season of Madagascar [Knott, 1998; Fietz and Dausmann, 2006; Génin, 2008; Simmen et al., 2010]. Humans also accumulate fat, a characteristic that was

83 undoubtedly present in early hominins exposed to highly seasonal or volatile environments [Wells,  
84 2012]. Obviously, more data are needed to better understand the metabolic adaptations to seasonal  
85 energy shortages and to the costs of reproduction in primates, particularly the potential buffer role  
86 of fattening.

87 Lemurs provide an interesting case study with regard to the effect of seasonal changes in  
88 resource availability and dietary quality on energy balance. Within lemur communities, the scarcity  
89 of fruit specialists relative to their African or Asian counterparts has been related to the ecological  
90 particularities of Madagascar assumed by some authors, namely unusually low food productions,  
91 low protein content of fruits, and fruit scarcity for long periods during the dry season [Fleagle and  
92 Reed, 1996; Ganzhorn et al., 2009; Donati et al., 2017]. There is also some evidence that, due to  
93 climatic irregularities in some years, lemur communities may experience a fruit shortage during the  
94 rainy season beside the dry season [Wright, 1999]. Finally, stochastic and frequent dramatic  
95 climatic events such as cyclones occur in Madagascar, which affects individual survival through the  
96 loss of food resources [Gould et al., 1999; Dunham et al., 2011; Lewis and Rakotondranaivo, 2011].  
97 At an evolutionary time-scale, it would therefore make sense that a range of lemurs has evolved a  
98 combination of energy-saving mechanisms, behavioural flexibility and opportunistic feeding  
99 strategies. In this respect, a study of body composition and metabolism in sympatric ring-tailed  
100 lemurs (*Lemur catta*) and “brown lemurs” (*Eulemur rufifrons x collaris*) living in the subarid south  
101 of Madagascar (Berenty Reserve) showed that these seasonal breeding species had large amounts of  
102 body fat following the period of greatest food abundance [Simmen et al., 2010], consistent with the  
103 hypothesis of a well-defined fattening period just prior to the mating season and the long dry season  
104 (see also [LaFleur, 2012]).

105 Given the fragmented nature of available data on the metabolic ecology of these large-  
106 sized lemurs in the wild, one aim of the present study is to provide a more in-depth analysis of the  
107 risk of energy imbalance incurred by ring-tailed lemurs and brown lemurs in different seasons and  
108 food supply contexts. In particular, we analyze whether energy consumption varies according to the

109 timing of the reproductive cycle and is compatible with a fattening process. In order to identify  
110 possible periods of positive and negative energy balance, we estimate metabolizable energy intake  
111 in different seasons and contrast these estimates with total energy expenditure (TEE) measured  
112 previously with doubly labelled water at the same study site in both species (Berenty Reserve  
113 [Simmen et al., 2010]). Finally we discuss whether the cathemeral behaviour (i.e. significant  
114 nocturnal and diurnal activity) reported for these lemurs, one characteristic of the so-called “lemur  
115 syndrome” [Wright, 1999], may be a consequence of insufficient energy intake during daytime or  
116 low ability of these species to digest plant fibres [Engqvist and Richard, 1991; Donati et al., 2007,  
117 2009, 2013; LaFleur et al., 2014]. Based on the seasonal pattern of nocturnality reported for  
118 *Eulemur rufifrons x collaris* at Berenty [Donati et al., 2009] and some activity of *Lemur catta* at  
119 night (Berenty: [Donati et al., 2013]), we made the following predictions: during the dry season, the  
120 two lemurs are not able to meet their energy needs (with possible species differences) by simply  
121 feeding during the day, even accounting for body fat stored during the wet season. In contrast, we  
122 expect that during the period of fruit abundance prior to the mating period and long dry season,  
123 these lemurs maximize their caloric intake, with a large positive energy balance reached through  
124 daytime feeding.

125

## 126 **Methods**

127 We studied diet composition and seasonal dietary changes in the two lemur species in  
128 1998/1999 [Simmen et al., 2003]. Information on the study site, subjects, behavioural methods and  
129 diets can be found in the 2003 publication. However we give here an overview of the methodology  
130 used to determine diet as, in the present paper, we quantify the daily intake of macronutrients,  
131 secondary metabolites and metabolizable energy using the proportions of plant species and plant  
132 parts in the diet determined in the previous study.

133

134 *Study site and lemur groups*

135 Malaza gallery forest (97 ha) is part of Berenty Private Reserve, Southern Madagascar  
136 (25°0.29'S, 46°19.37'E). The climate is sub-arid with marked, irregular annual rainfall variation  
137 (300-900 mm) and a dry season lasting from April to October [Jolly et al., 2006]. Details on forest  
138 composition and habitat structure in Malaza can be found elsewhere [Jolly et al., 2006; Simmen et  
139 al., 2012, 2014]. At the forest edge, exotic trees species used as foods by some of the focal groups  
140 of lemurs are grown as ornamental plants [Simmen et al., 2003].

141 Ring-tailed lemurs (*Lemur catta*) and “brown” lemurs (*Eulemur rufifrons x collaris*) live in  
142 multi-male, multi-female troops and are primarily frugivore/folivores, supplementing their diet with  
143 flowers (Table 1). There is no significant sex difference of body mass across the populations studied  
144 in the forest areas investigated here (ring-tailed lemurs:  $2.3\pm 0.3\text{kg}$ ; brown lemurs:  $1.9\pm 0.2\text{kg}$ ;  
145 [Simmen et al., 2010]). Ring-tailed lemurs at Berenty defend relatively small, well-defined home  
146 ranges whereas groups of brown lemurs forage over comparatively larger areas and are less  
147 territorial (Jolly et al., 2006; Pinkus et al., 2006; Tanaka 2007; Simmen, pers. obs.). In the  
148 comparative analysis of diet that we carried out in 1998 and 1999 [Simmen et al., 2003], we studied  
149 3 to 5 groups per lemur species. We selected groups of brown lemurs that foraged within the areas  
150 used by ring-tailed lemur troops. Some focal groups of ring-tailed lemurs foraged inside Malaza  
151 forest, while others lived at the forest edge having access to ornamental plants. Brown lemur groups  
152 were more arboreal than ring-tailed lemurs and were more rarely observed in environments without  
153 tree cover at the forest edge [Simmen et al 2003].

154

#### 155 *Behavioural data collection*

156 Daytime is the only temporal sequence of the nycthemeral cycle during which we can  
157 reliably estimate food and energy intake on a quantitative basis. Therefore, in 1998/1999, we  
158 determined diets from mouthful counts (Hladik 2011; Zinner 1999; Simmen et al., 2017) during  
159 continuous observation of focal individuals (*continuous focal animal sampling*) followed from  
160 dawn to dusk during three periods, following a preliminary survey in December 1997: May-June

161 (early dry season), October-November (late dry season), February-March (late wet season). Due to  
162 the highly synchronized reproductive cycle related to photoperiod changes, breeding occurs  
163 approximately in April-May, gestation lasts until September-October, and lactation until February.  
164 Focal individuals could be approached within a few meters and the visibility generally allowed  
165 close observation of individual feeding activity. The observation sample consisted of  $4.0 \pm 0.6$  full  
166 days of observation on average per species per study period during which only food intake was  
167 recorded (311 hours total observation balanced between the two species and periods). Focal animals  
168 were adult males and females (excluding lactating females), with these identified from their natural  
169 markings or dye spots. Each full-day follow was devoted to one focal animal from one group, but  
170 when the animal was out of sight due to poor visibility conditions, we temporarily shifted to the  
171 nearest individual (adult or large subadult of the same sex). Observation days were spread as much  
172 as possible throughout each study period and alternated between groups of the two species within  
173 intervals generally  $\leq 5$  days. Daily observations of focal individuals followed during periods up to  
174 14 hours (depending on seasonal photoperiod changes) without major interruption constituted 40%  
175 of total records. We weighed the fresh mass of each food item eaten to convert mouthfuls into  
176 masses of ingested matter. We derived the amount of food eaten for each given day from the sum of  
177 foods consumed by the focal individual (supplemented by the food intake of the alternative  
178 individuals). We averaged daily food intake recorded for each study period and each lemur species.  
179 On this basis, we calculated the relative proportions of the different food items in the seasonal diets  
180 [Simmen et al 2003].

181           The data collected may not be representative of the full feeding repertoire since the  
182 observed sample was small. However, referring to the main studies carried out in the last decades  
183 on the feeding ecology of ring-tailed lemur and brown lemur at the study site, we believe this did  
184 not substantially affect recognition of major food plants in the diet: all studies point out to a  
185 consistently low dietary diversity in both species, while the plant species that contributed most to  
186 their diet in our study were consistent with the top plant species used as foods reported in

187 [Rasamimanana and Rafidinarivo, 1993; Jolly et al. 2006; Pinkus et al. 2006]. This, together with  
188 the observations made during our successive studies at Malaza from 1987 until the present,  
189 suggested that we captured the most prominent features of the dietary characteristics of these  
190 species at the study site. As a result,  $\leq 4$  plant species accounted for  $\geq 75\%$  of the food matter  
191 ingested by each species in each season from a total of 66 food plant spp. (87 food items) and 24  
192 spp. (38 items) consumed respectively by ring-tailed lemurs and by brown lemurs during the study  
193 [Simmen et al., 2003]. The two lemur species typically focused their food choices on *Tamarindus*  
194 *indica*, *Celtis* spp. and other dominant plant species of the gallery forest, supplemented by  
195 *Azadirachta indica*, one common ornamental plant available for groups ranging at the forest edge.  
196 Dietary overlap between the two species was the largest during the early dry season, when these  
197 lemurs used a restricted set of similar plant species as their staple, as found elsewhere [Sussman,  
198 1974; Rasamimanana and Rafidinarivo, 1993]. Table 1 provides an overview of major food  
199 categories in the diet, split by seasons.

200 [Table 1]

201

## 202 *Chemical assays*

203 We conducted plant sampling and chemical analyses at the same time as the initial  
204 comparative study of the feeding ecology of these species [Simmen et al., 2003; Rothman et al.  
205 2012]. We collected plant samples from feeding trees and dried major food plant samples (62 items;  
206 39 plant species) in an electric field oven (at 50°C) for botanical identification at the National  
207 Museum of Natural History in Paris by experts (Annette Hladik and colleagues) and for chemical  
208 analysis. We obtained samples of major foods from multiple food trees or lianas of the same species  
209 and mixed them to account for within-species variation in chemical content [Rothman et al., 2012].  
210 Since we were primarily interested in estimating the overall diet quality and average daily energy  
211 intake in each season, we blended fruits, flowers and leaves according to their proportion in the  
212 seasonal diets of each lemur species prior to performing the chemical analyses. The mixtures were

213 composed of plant parts that together accounted for 85-100% of the total dry matter consumed by  
214 ring-tailed lemurs or brown lemurs during each of the three seasons studied. Although the dietary-  
215 mixture method reduces the costs of the chemical analyses, one limitation of the method is that it  
216 does not allow for testing the statistical significance of seasonal differences in nutrient and energy  
217 intake. Sampling heterogeneous mixtures when the sample contains many different samples and the  
218 amount of sample used in each assay is low, also introduces a risk that the food mixture is not  
219 representative. To assess the consistency of the results, we performed additional chemical analyses  
220 on staple foods separately. We then calculated the weighted mean proportion of nutrients from  
221 results obtained on each staple food — that together accounted for  $\geq 75\%$  of the total dry matter  
222 consumed by the lemurs in each season — and we compared it with the nutrient composition of the  
223 dietary mixtures.

224         We determined humidity from portions of plant samples dried at 105°C in the laboratory.  
225 We carried out chemical analyses on macronutrients and fibre at the ISHA (Institut Scientifique  
226 d'Hygiène et d'Analyses, Longjumeau) following official analytical norms (AFNOR group as a  
227 member of ISO). The analyses on the dry powdered samples were performed in duplicate and the  
228 results were expressed as mean values with a  $\leq 5\%$  difference between the assays for each sample.  
229 We determined crude protein ( $N \times 6.25$ ; Kjeldahl method), crude lipids (HCl hydrolysis and  
230 extraction with petroleum-ether), soluble sugars (HPLC), neutral detergent fibre (NDF), and ash  
231 (see Simmen et al. [2014] for specific references). We also analyzed phenolic compounds (modified  
232 Prussian blue assay [Price and Butler 1977], condensed tannins with acid butanol assay [Porter et  
233 al., 1986]), and protein precipitation efficiency (bovine serum albumin-binding property of the plant  
234 extract [Asquith and Butler, 1985]) to provide a broader assessment of diet quality beyond  
235 macronutrient content. Results for phenolics and condensed tannin concentrations are expressed in  
236 relative units as equivalents of a chemical standard (tannic acid and purified quebracho tannin,  
237 respectively) and cannot be considered absolute [Rothman et al., 2012].

238

239 *Data analysis*

240 Metabolizable energy intake in each season was the sum of energy provided by readily  
241 digestible energy (i.e. soluble carbohydrates, fat, protein, pectin, and starch) and NDF corrected for  
242 digestibility in food eaten daily averaged for that season. Daily metabolizable energy intake was  
243 calculated in  $\text{kJ}\cdot\text{d}^{-1}$  by multiplying the summed  $\text{kCal}$  eaten daily by 4.18 [Rothman et al., 2012]. The  
244 HPLC assay for soluble sugars does not account for other readily available carbohydrates  
245 potentially present in primate foods, e.g. soluble fibres like pectin and storage compounds like  
246 starch, which could lead to an underestimation of the metabolizable energy value of the diet.  
247 Therefore, we assumed an additional average concentration of 5% of the total dry mass for pectin  
248 and starch, respectively, based on a review of the literature on primate fruits and other tropical fruits  
249 and leaves [Simmen et al., 2017].

250 Similarly wild lemurs' ability to digest NDF is not known but experiments on fibre  
251 digestion in captive *Lemur catta*, *Eulemur* spp. and another large frugivorous lemur, *Varecia*  
252 *variegata*, are helpful for calculating a range of energy intake values derived from microbial  
253 fermentation [Overdorff and Rasmussen, 1995; Campbell et al., 2004a; Schmidt et al., 2005].  
254 Specifically, the NDF digestibility coefficients we used varied according to the fibre content in the  
255 dry matter ingested by our study species. When NDF content in the natural diets was low ( $< 25\%$   
256 during the late wet season; table 2), we used a NDF digestibility coefficient at 41% (*Eulemur fulvus*  
257 [Campbell et al., 2004a]). When NDF content was high (up to 46% during the early dry and late dry  
258 season; table 2), we calculated two energy estimates that assume either low or no fibre digestibility  
259 as reported in captivity. In the first estimate, we used a figure at 12% for fibre digestibility, as found  
260 in *Eulemur macaco* fed diets with NDF at 31-35% [Schmidt et al., 2005]). In the second estimate,  
261 we assumed zero fibre digestibility (comparable to the results for *Eulemur fulvus* and *Lemur catta*  
262 fed with large-sized cellulose particles [Overdorff and Rasmussen 1995]) and we calculated  
263 metabolizable energy input based on readily digestible energy only.

264 We estimated the daily metabolizable energy input in each season by first converting the  
265 average dry matter ingested daily into caloric values ( $4/9/4$  kCal.g<sup>-1</sup> for the energy equivalents of  
266 soluble sugars, fat, and protein, respectively). We assumed an energy equivalent of NDF and pectin  
267 at 3 kCal.g<sup>-1</sup> to account for the energy diverted by gut microbial metabolism [Conklin et al., 2006].  
268 Basically, we used the same range of NDF digestibility coefficients for brown lemurs and ring-  
269 tailed lemurs given similarities in the range of cellulose digestibility in the study reported by  
270 Overdorff and Rasmussen ([1995]; 0 to 21% and 0 to 30%, respectively). Yet, we accounted for a  
271 possibly higher ability of *Lemur catta* to subsist on fibrous foods (see Introduction) when  
272 discussing the relationship between energy intake and energy expenditure. Indeed, *Lemur catta* is  
273 more closely related to the folivorous bamboo lemurs (*Hapalemur* spp.; [Markolf and Kappeler,  
274 2013], it possesses a large and haustrated caecum that may function to improve fibre fermentation  
275 [Campbell et al., 2000], and the overall foliage proportion in their diet is higher than that of brown  
276 lemurs which eat more fleshy fruits [Berenty; Simmen et al., 2003; Pinkus et al. 2006].

277

## 278 **Results**

279 Table 2 shows the nutrient composition of the seasonal diets (i.e. the dietary mixtures) as  
280 well as the weighted mean concentration of nutrients calculated from the chemical composition of  
281 each staple food. Both types of analyses produced broadly similar results even though the  
282 contribution of food items to the diet differed somewhat between the two analyses. Mature leaves  
283 and unripe fruits of *Tamarindus indica* (Caesalpinioideae), with their high concentrations of NDF  
284 and condensed tannins as well as large protein-precipitation effects, were the staple foods consumed  
285 during the dry months — during the early dry season, they accounted for 60 to 96% of the total  
286 amount of food ingested daily. Consequently, as for other macronutrients and secondary  
287 metabolites, the NDF content of the dry season diet of the two lemur species largely reflected the  
288 chemical composition of this tree species. Brown lemurs had lower dietary proportions of protein  
289 than ring-tailed lemurs during the dry months, while the diet of both species contained large

290 concentrations of nitrogen-digestibility reducers (bovine serum albumin-binding efficiency and  
291 phenolic content; Table 2).

292

293 [Table 2]

294

295 [Table 3]

296 At the beginning of the early dry season, both primate species had lower quality diets and  
297 reduced metabolizable energy intake compared with the wet season (Fig. 1). During these dry  
298 months, the physiological fuel value of the diet was similar between the two species (Table 3), but  
299 brown lemurs consumed digestible calories in greater quantity than ring-tailed lemurs because of  
300 their greater daily food intake.

301 During the late wet season when ripe fruit pulps were the staple, soluble carbohydrates  
302 accounted for a large proportion of the food matter consumed by both lemur species (table 2).  
303 During this period, daily metabolizable energy intake was the highest, with readily digestible  
304 energy accounting for a great proportion of total metabolizable energy intake (Fig. 1). As a result,  
305 wet season diets were characterized by a higher physiological value ( $\text{kJ}\cdot\text{g}^{-1}$  dry mass) relative to  
306 other seasons (Table 3). Interestingly, the contribution of readily digestible energy to overall energy  
307 intake was always remarkably high regardless of the digestibility coefficients used to estimate the  
308 contribution of fibre to energy input. We found no substantial difference between the two species  
309 with regard to this energy fraction (Fig. 1) or the NDF content of their diet at any season (Table 3).

310 [Figure 1]

311

## 312 **Discussion**

313 Although based on a limited sample of feeding observations, our data reveal a large  
314 seasonal variation of energy intake, with a broadly similar temporal pattern in the two lemur species  
315 (i.e. an increase during the wet season). The high energy consumption found during the wet season

316 is consistent with the hypothesis of a seasonal storage of body fat. In captivity, ring-tailed lemurs  
317 and brown lemurs store fat and undergo seasonal variation in hair metabolism, feeding and activity  
318 of metabolically-active hormones [Pereira and Pond, 1995; Pereira et al., 1999]. These biological  
319 changes are synchronized by the day-length variation, a predictable abiotic cue that allows them to  
320 anticipate the season of food shortage lasting up to 6 or 7 months in their habitats and to adjust their  
321 reproductive cycle accordingly. In our previous study of body composition in ring-tailed lemurs and  
322 brown lemurs at Malaza [Simmen et al., 2010], we observed a high body fat content at about 18%  
323 in both species, without sex differences, at the end of March, before the dry season. This is close to  
324 the body fat percentage found in captive orangutans (approximately 20% [Pontzer et al., 2016a])  
325 and much higher than that of other non-hibernating primates feeding on natural foods (2-5%  
326 [review in Dittus, 2013]). In addition, in the Simmen et al 2010 study, the few individuals captured  
327 from each species showed a large increase in body mass ( $17 \pm 4\%$ ) between the December pilot  
328 study (i.e. after the rains resumed and new leaves and flower buds were available) and when they  
329 were recaptured in March of the following year. Consequently, energy data from different years  
330 indicate a recurring trend in these lemurs of achieving a positive energy balance during the rainy  
331 season. This conclusion as well as the question of whether ring-tailed lemurs and brown lemurs  
332 present an energy imbalance during the dry season can be tested further by contrasting our estimates  
333 of energy intake with the metabolic data previously collected in March/April 2009 in these species  
334 [Simmen et al., 2010], namely total energy expenditure (TEE) and body composition.

335         Energy intake theoretically equates with TEE to sustain a balanced energy budget if  
336 animals do not store fat. TEE was formerly assessed in groups of lemurs that were foraging within  
337 the same forest areas as those studied here. Despite the discontinuity between the studies, energy  
338 intake can be contrasted to energy expenditure values on the basis that TEE is maintained within a  
339 narrow, species-specific physiological range, as discussed by Pontzer et al. [2015]. Indeed, recent  
340 findings in humans [Pontzer et al., 2016b] show that TEE during habitual physical activity cannot  
341 be considered as the sum of each metabolic cost incurred for body maintenance and behavioural

342 activity (referred to as the additive model). The constrained energy model [Pontzer et al., 2016b], in  
343 contrast, is supported by an increasing number of studies in mammals, including primates, which  
344 show that TEE (measured with doubly labelled water over an interval of a few days) does not differ  
345 substantially between seasons or between captive and wild individuals (TEE or mass-adjusted TEE  
346 [Kenagy et al., 1989; Nagy et al., 1999; Pontzer et al., 2014, 2016a; Pontzer, 2015]). In this model,  
347 TEE is maintained within a narrow metabolic range because metabolic and behavioural adjustments  
348 offset the costs induced by energy-demanding activities such as vigorous physical activity [Pontzer,  
349 2015; Pontzer et al., 2016b]. In *Microcebus murinus*, for example, TEE differs neither between the  
350 dry and the wet season in normothermic individuals [Schmid and Speakman, 2000]. Other studies  
351 show that gregarious lemurs reduce thermal stress by using behavioural thermoregulation like social  
352 huddles and peculiar resting postures and, according to some authors, by performing daylight and  
353 nocturnal activity ([Donati et al., 2011; LaFleur et al., 2014; Kelley et al., 2016; Eppley et al.,  
354 2017]; but see Fletcher et al. [2012]).

355         Assuming that the lemurs studied here also maintain their TEE within a narrow metabolic  
356 range, the daily energy intake of *Lemur catta* in the early dry season ( $\leq 443 \text{ kJ.d}^{-1}$ ) is less than the  
357 total energy expenditure previously measured at  $626 \pm 138 \text{ kJ.d}^{-1}$  with doubly labelled water. With  
358 large amounts of body fat reserves, however, ring-tailed lemurs could theoretically meet their  
359 maintenance energy requirements during most of the dry season: an average of 424 grams of body  
360 fat recorded at the end of the wet season [Simmen et al., 2010] is equivalent to a 25-day supply (or  
361 15,800 kJ) for an individual expending  $626 \text{ kJ.d}^{-1}$ . The energy contained in these fat reserves,  
362 combined with the calories obtained from the early dry season diet during daytime, would make it  
363 possible to balance the energy needs over a 3-month period. Foraging during the late dry season on  
364 the other hand yields energy intake well above requirements (Table 3). In brown lemurs, the  
365 calories ingested during daytime in the early dry season ( $576\text{-}659 \text{ kJ.d}^{-1}$ ) and late dry season ( $516\text{-}$   
366  $573 \text{ kJ.d}^{-1}$ ) are barely or not sufficient to meet the energy requirements (TEE:  $610 \pm 90 \text{ kJ.d}^{-1}$   
367 [Simmen et al., 2010]). However, the 325 g of fat stored at the end of the wet season [Simmen et al.,

368 2010] are equivalent to a 20-day energy supply (or 12,100 kJ) for an average individual expending  
369 610 kJ.d<sup>-1</sup>. In this case, the combination of calories ingested during the dry season with energy  
370 stored as body fat is theoretically sufficient to avoid negative energy balance during the whole dry  
371 season. Finally, crude protein intake in both species (Table 3) is far above the estimated protein  
372 requirements of omnivorous primates (<15% of metabolizable energy; [Oftedal 1991]), especially  
373 during the dry season.

374 We acknowledge that these analyses are solely predictive since our study spanned a short  
375 period and we assumed similar digestibility coefficients between wild and captive animals. Also,  
376 one potential limitation of our energy throughput analysis is that we compared energy intake with  
377 body fat content and energy expenditure measured from different years, using the constrained  
378 energy model of Pontzer et al. [2016b]. Further studies should be conducted to assess whether the  
379 energy expenditure of these lemurs may vary to a greater extent than this model suggests.

380 Assuming that our hypotheses are correct, the results are, at first glance, not in agreement  
381 with studies that suggest a link between diet and cathemerality (i.e. the nocturnal and diurnal  
382 activity). Cathemeral activity in *Eulemur*, it is argued, may be a consequence of the high proportion  
383 of fibrous foods in the diet and the inability to extract enough energy from the plant parts consumed  
384 during the day [Donati et al., 2007, 2009]. As a result, lemurs are not able to meet their energy  
385 needs if they do not supplement their diurnal feeding activity with night-time feeding activity  
386 [Engqvist and Richard, 1991].

387 In fact, our data can be reconciled with the dietary fibre-driven hypothesis on cathemerality  
388 to the extent that we probably overestimated energy available from food and adipose tissue. Not all  
389 protein ingested and fat substrates can be used as fuel for energy expenditure, and primates, like  
390 other mammals, require up to 2.8 g.kg BW<sup>-1</sup>.d<sup>-1</sup> of protein and at least 2% body fat to meet their  
391 maintenance costs (Oftedal 1991; NRC 2003; Dittus 2013). Accordingly, in ring-tailed lemurs, the  
392 pool of energy available after subtracting these proportions of nutrients ( $\leq 336$  kJ.d<sup>-1</sup> from the diet  
393 and 14,500 kJ as fat; Table 3) suggests that this species may avoid negative energy balance based

394 on daytime foraging during no more than 2 months at the beginning of the dry season (that is, until  
395 June-July). Brown lemurs, with  $490 \text{ kJ.d}^{-1}$  as readily digestible energy (assuming zero fibre  
396 digestibility from NDF-rich diets during the dry season; Table 3) plus 10,400 kJ as fat stores, would  
397 be able to balance their energy needs during no more than 3 months (that is, until July-August). In  
398 addition, calculations suggest that daytime energy intake is insufficient for this species at the end of  
399 the dry season, unlike ring-tailed lemurs (Table 3). Yet, these predictions do not account for other  
400 potentially limiting factors. For instance, protein bioavailability is likely diminished by the high  
401 concentrations of tannin and other phenolic compounds in the diet (both species; Table 2). The dry  
402 season diet of *Eulemur* contained a limited concentration of crude protein relative to estimated  
403 requirements of omnivorous primates (as low as 8.5% on a dry matter basis, Table 2, versus 7-16%,  
404 [Oftedal 1991]) — even though protein intake was far above the estimated requirements ( $>26\%$  as a  
405 proportion of metabolizable energy, Table 3, versus  $<15\%$ , [Oftedal 1991]). In addition, due to the  
406 timing of reproduction and high energy cost of the breeding season for both males and gestating  
407 females [May-June; Sauther, 1991; Pereira et al., 1999; Rasamimanana et al., 2006; Ostner et al.,  
408 2008], the fat reserves may in fact be depleted shortly after the end of the rainy season. This would  
409 force individuals to distribute their dry season feeding activity over a 24-hour period to balance  
410 their energy needs. In *Eulemur rufifrons x collaris* at Malaza, cathemeral activity is predominantly  
411 observed during the dry season, with a peak in June-July, when food is scarce and the diet is rich in  
412 fibrous foods [Pinkus et al., 2006; Donati et al., 2009]. Systematic observations of *Lemur catta*  
413 using camera traps and direct observation in the wild are limited to the wet season and reveal some  
414 limited activity at night (Berenty: Donati et al., 2013); Tsimananpetsotsa: [LaFleur et al., 2014]).  
415 The nycthemeral activity pattern of this species during the dry season suggests substantial nocturnal  
416 activity, consistent with the hypothesis of a risk of energy imbalance, but the data remain scant  
417 (April-June; [Traina 2001]).

418 In conclusion, our results are in agreement with the hypothesis that lemur species that  
419 primarily feed on scarce and irregular food resources (fruit or insects) have evolved energy-saving

420 mechanisms including fat storage [Sauther, 1998; Génin, 2008; Simmen et al., 2010]. The risk of  
421 nutritional deficiency in these lemurs and other frugivorous lemurs in Madagascar may vary each  
422 year due to the erratic climate and irregularities in the fruit supply [Sauther, 1998; Wright, 1999;  
423 Dewar and Richard, 2007]. Accordingly we expect that the fat storage process and energy intake  
424 may be compromised during unfavourable years and that any major irregularities in the food supply  
425 during the wet season will exert a strong selective pressure on the fitness of these lemurs. If the  
426 lemurs fail to store enough fat, they could in the short term also rely on the catabolism of muscle  
427 tissue, but this hypothesis needs to be explored further. Finally, given the differences in diet, gut  
428 morphology and physiology between the two species studied, and partitioning of their ecological  
429 niches under conditions of natural coexistence, we expect that ring-tailed lemurs and brown lemurs  
430 do not incur the same risk of energy deficiency. During the period of food shortage at Berenty,  
431 brown lemurs tend to eat larger quantities of food than ring-tailed lemurs, probably to compensate  
432 for the low amount of energy extracted from fibrous foods [Simmen et al., 2003]. Conversely,  
433 seasonal changes in the gut microbiota of wild *Lemur catta* [Fogel 2015] indicate possible  
434 compensation effect of the gut symbiotic microorganisms to extract metabolizable energy from  
435 leaves. The possibility that ring-tailed lemurs obtain more energy per unit weight of food consumed  
436 than brown lemurs must be explored further.

437

### 438 **Acknowledgements**

439 We thank the DeHeulme family for providing permission to study lemur ecology in the  
440 Berenty Reserve, and the DGDRF in Antananarivo (Ministère des Eaux et Forêts, Madagascar) who  
441 delivered the authorizations to carry out our research program. We thank André Marez for his help  
442 in the chemical assays of secondary metabolites. We are grateful to Shelly Masi who provided  
443 useful comments, to V. Winchester, Anne Mertl-Millhollen and James R. Tysell who edited the  
444 successive drafts of the text, and to three anonymous reviewers. We declare we have no competing  
445 interests.

446

447

**References**

448

Asquith TN, Butler LG (1985). Use of a dye-labeled protein as spectrophotometric assay for protein precipitants such as tannin. *Journal of Chemical Ecology* 11: 1535–1544.

449

450

Bercovitch FB (1987). Female weight and reproductive condition in a population of olive baboons (*Papio anubis*). *American Journal of Primatology* 12: 189–195.

451

452

Brockman DK, van Schaik CP (2005). Seasonality and reproductive function. In *Seasonality in Primates: Studies of Living and Extinct Human and Non-Human Primates* (Brockman DK, van Schaik CP, eds.), pp 269–305. New York, Cambridge University Press.

453

454

455

Campbell JL, Williams CV, Eisemann JH (2004a). Use of total dietary fibre across four lemur species (*Propithecus verreauxi coquereli*, *Haplemur griseus griseus*, *Varecia variegata*, and *Eulemur fulvus*): does fibre type affect digestive efficiency? *American Journal of Primatology* 64: 323–335.

456

457

458

459

Campbell JL, Williams CV, Eisemann JH (2004b). Characterizing gastrointestinal transit time in four lemur species using barium-impregnated polyethylene spheres (BIPS). *American Journal of Primatology* 64: 309–321.

460

461

462

Campbell JL, Eisemann JH, Williams CV, Glenn KM (2000). Description of the gastrointestinal tract of five lemur species: *Propithecus tattersalli*, *Propithecus verreauxi coquereli*, *Varecia variegata*, *Haplemur griseus*, and *Lemur catta*. *American Journal of Primatology* 52: 133–142.

463

464

465

466

Canale CI, Henry PY (2010). Adaptive phenotypic plasticity and resilience of vertebrates to increasing climatic unpredictability. *Climate Research* 43: 135–147.

467

468

Conklin-Brittain NL, Knott CD, Wrangham RW (2006). Energy intake by wild chimpanzees and orang-utans: methodological considerations and a preliminary comparison. In *Feeding Ecology in Apes and Other Primates. Ecological, Physical and Behavioral Aspects* (Hohmann G, Robbins MM, Boesch C, eds.), pp 445–471. Cambridge, Cambridge University Press.

469

470

471

- 472 Daniels HL (1984). Oxygen consumption in *Lemur fulvus*: deviation from the ideal model. *Journal*  
473 *of Mammalogy* 65: 584–592.
- 474 Dewar RE, Richard AF (2007). Evolution in the hypervariable environment of Madagascar.  
475 *Proceedings of the National Academy of Sciences of the United States of America* 104: 13723–  
476 13727.
- 477 Dittus WPJ (2013). Arboreal adaptations of body fat in wild Toque macaques (*Macaca sinica*) and  
478 the evolution of adiposity in primates. *American Journal of Primatology* 152: 333–344.
- 479 Donati G, Bollen A, Borgognini-Tarli SM, Ganzhorn JU (2007). Feeding over the 24-h cycle:  
480 dietary flexibility of cathemeral collared lemurs (*Eulemur collaris*) *Behavioral Ecology and*  
481 *Sociobiology* 61:1237–1251.
- 482 Donati G, Baldi N, Morelli V, Ganzhorn JU, Borgognini-Tarli SM (2009). Proximate and ultimate  
483 determinants of cathemeral activity in brown lemurs. *Animal Behaviour* 77: 317–325.
- 484 Donati G, Ricci E, Baldi N, Morelli V, Borgognini-Tarli SM (2011). Behavioral thermoregulation  
485 in a gregarious lemur, *Eulemur collaris*: effects of climatic and dietary-related factors.  
486 *American Journal of Physical Anthropology* 144: 355–364.
- 487 Donati G, Santini L, Razafindramanana J, Boitani L, Borgognini-Tarli S (2013). (Un-)expected  
488 nocturnal activity in “diurnal” *Lemur catta* supports cathemerality as one of the key  
489 adaptations of the lemurid radiation. *American Journal of Physical Anthropology* 150: 99–106.
- 490 Donati G, Santini L, Eppley TM, Arrigo-Nelson SJ, Balestri M, Boinski S, Bollen A, Bridgeman  
491 LL, Campera M, Carrai V, Chalise MK, Derby A, Hohmann G, Kinnaird MF, Koenig A,  
492 Kowalewski M, Lahann P, McLennan MR, Nekaris KAI, Nijman V, Norscia I, Ostner J,  
493 Polowinsky SY, Schülke O, Schwitzer C, Stevenson PR, Talebi MG, Tan C, Vogel ER, Wright  
494 PC, Ganzhorn JU (2017). Low levels of fruit nitrogen as drivers for the evolution of  
495 Madagascar’s primate communities. *Scientific Reports* 7: 14406.
- 496 Du Mond FV, Hutchinson TC (1967). Squirrel monkey reproduction: the "fatted" male  
497 phenomenon and seasonal spermatogenesis. *Science* 158: 1067–1070.

- 498 Dunham AE, Erhart EM, Wright PC (2011). Global climate cycles and cyclones: consequences for  
499 rainfall patterns and lemur reproduction in southeastern Madagascar. *Global Change Biology*  
500 17: 219–227.
- 501 Engqvist A, Richard A (1991). Diet as a possible determinant of cathemeral activity patterns in  
502 primates. *Folia Primatologica* 57:169–172.
- 503 Eppley TM, Watzek J, Dausmann KH, Ganzhorn JU, Donati D (2017). Huddling is more important  
504 than rest site selection for thermoregulation in southern bamboo lemurs *Animal Behaviour* 127:  
505 153–161.
- 506 Fietz J, Dausmann KH (2006). Big is beautiful – fat storage and hibernation as a strategy to cope  
507 with marked seasonality in the fat-tailed dwarf lemur (*Cheirogaleus medius*). In *Lemurs:  
508 Ecology and Adaptation* (Gould L, Sauther ML, eds), pp 97–111. New York, Springer.
- 509 Fleagle JG, Reed KE (1996). Comparing primate communities: a multivariate approach. *Journal of  
510 Human Evolution* 30: 489–510.
- 511 Fletcher QE, Speakman JR, Boutin S, McAdam AG, Woods SB, Humphries MM (2012). Seasonal  
512 stage differences overwhelm environmental and individual factors as determinants of energy  
513 expenditure in free-ranging red squirrels. *Functional Ecology* 26: 677–687.
- 514 Fogel AT (2015). The gut microbiome of wild lemurs: a comparison of sympatric *Lemur catta* and  
515 *Propithecus verreauxi*. *Folia Primatologica* 86: 85–95.
- 516 Ganzhorn JU, Arrigo-Nelson S, Boinski S, Bollen A, Carrai V, Derby A, Donati G, Koenig A,  
517 Kowalewski M, Lahann P, Norscia I, Polowinsky SY, Schwitzer C, Stevenson PR, Talebi MG,  
518 Tan C, Vogel ER, Wright PC (2009). Possible fruit protein effects on primate communities in  
519 Madagascar and the Neotropics. *PLoS ONE* 4(12), DOI: 10.1371/e8253.
- 520 Garcia C, Huffman MA, Shimizu K, Speakman JR (2011). Energetic consequences of seasonal  
521 breeding in female Japanese macaques (*Macaca fuscata*). *American Journal of Physical  
522 Anthropology* 146: 161–170.
- 523 Génin F (2008). Life in unpredictable environments: first investigation of the natural history of

- 524 *Microcebus griseorufus*. *International Journal of Primatology* 29: 303–321.
- 525 Gould L, Sussman RW, Sauther ML (1999). Natural disasters and primate populations. The effects  
526 of a two-year drought on a naturally occurring population of ring-tailed lemurs (*Lemur catta*) in  
527 Southwestern Madagascar. *International Journal of Primatology* 20: 69–84.
- 528 Hladik CM (2011). Considerations on field methods used to assess non-human primate feeding  
529 behaviour and human food intake in terms of nutritional requirements. In *Centralizing*  
530 *Fieldwork: Critical Perspectives from Primatology, Biological Anthropology and Social*  
531 *Anthropology* (MacClancy J, Fuentes A, eds.), pp 170–185. Oxford: Berghahn.
- 532 Irwin MT, Raharison JL, Raubenheimer D, Chapman CA, Rothman JM (2014). Nutritional  
533 correlates of the “lean season”: effects of seasonality and frugivory on the nutritional ecology  
534 of Diademed sifakas. *American Journal of Physical Anthropology* 153: 78–91.
- 535 Jolly A, Sussman RW, Koyama N, Rasamimanana H, editors (2006). *Ringtailed Lemur Biology.*  
536 *Developments in Primatology: Progress and Prospect*. Chicago, Springer.
- 537 Kelley EA, Jablonski NG, Chaplin G, Sussman RW, Kamilar JM (2016). Behavioral  
538 thermoregulation in *Lemur catta*: The significance of sunning and huddling behaviors.  
539 *American Journal of Primatology* 78: 745–754.
- 540 Kenagy GJ, Sharbaugh SM, Nagy KA (1989). Annual cycle of energy and time expenditure in a  
541 golden-mantled ground squirrel population. *Oecologia* 78: 269–282.
- 542 Knott CD (1998). Changes in orangutan caloric intake, energy balance, and ketones in response to  
543 fluctuating fruit availability. *International Journal of Primatology* 19: 1061–1079.
- 544 Koch F, Ganzhorn JU, Rothman JM, Chapman CA, Fichtel C (2016). Sex and seasonal differences  
545 in diet and nutrient intake in Verreaux's sifakas (*Propithecus verreauxi*). *American Journal of*  
546 *Primatology* DOI: 10.1002/ajp.22595.
- 547 LaFleur M (2012). *Ecology of ring-tailed lemurs (Lemur catta) at the Tsimanampetsotsa National*  
548 *Park, Madagascar: implications for female dominance and the evolution of lemur traits*. PhD  
549 thesis, University of Colorado at Boulder, USA.

- 550 LaFleur M, Sauther M, Cuzzo F, Yamashita N, Youssouf IAJ, Bender R (2014). Cathemerality in  
551 wild ring-tailed lemurs (*Lemur catta*) in the spiny forest of Tsimanampetsotsa National Park:  
552 camera trap data and preliminary behavioral observations. *Primates* 55: 207–217.
- 553 Lambert JE, Rothman JM (2015). Fallback foods, optimal diets, and nutritional targets: primate  
554 responses to varying food availability and quality. *Annual Review of Anthropology* 44: 493–  
555 512.
- 556 Lewis RJ, Kappeler PM (2005). Seasonality, body condition, and timing of reproduction in  
557 *Propithecus verreauxi verreauxi* in the Kirindy Forest. *American Journal of Primatology*  
558 67: 347–364.
- 559 Lewis RJ, Rakotonirainy F (2011). The impact of Cyclone Fanele on sifaka body condition and  
560 reproduction in the tropical dry forest of western Madagascar. *Journal of Tropical Ecology*  
561 27: 429–432.
- 562 Long E (2002). *The Feeding Ecology of Pteropus rufus in a Remnant Gallery Forest Surrounded by*  
563 *Sisal Plantations in South-East Madagascar*. PhD thesis, University of Aberdeen, UK.
- 564 Markolf M, Kappeler PM (2013). Phylogeographic analysis of the true lemurs (genus *Eulemur*)  
565 underlines the role of river catchments for the evolution of micro-endemism in Madagascar.  
566 *Frontiers in Zoology* 10:70.
- 567 Munn AJ, Clissold F, Tarszisz E, Kimpton K, Dickman C, Hume ID (2009). Hindgut plasticity in  
568 wallabies fed hay either unchopped or ground and pelleted: fiber is not the only factor.  
569 *Physiological and Biochemical Zoology* 82: 270–279.
- 570 Muroyama Y, Kanamori H, Kitahara E (2006). Seasonal variation and sex differences in the  
571 nutritional status in two local populations of wild Japanese macaques. *Primates* 47: 355–364.
- 572 Nagy KA, Girard IA, Brown TK (1999). Energetics of free-ranging mammals, reptiles, and birds.  
573 *Annual Review of Nutrition* 19: 247–277.
- 574 NRC (2003). *Nutrient Requirements of Nonhuman Primates: Second Revised Edition*. Washington  
575 DC, The National Academies Press.

- 576 Oates JF (1987). Food distribution and foraging behavior. In *Primate Societies* (Smuts BB, Cheney  
577 DL, Seyfarth RM, Wrangham RW, Struhsaker TT, eds.), pp 197-209. Chicago, The University  
578 of Chicago Press.
- 579 Oftedal OT (1991). The nutritional consequences of foraging in primates: the relationship of  
580 nutrient intakes to nutrient requirements. *Philosophical Transactions of the Royal Society of  
581 London B* 334: 161–170.
- 582 Ostner J, Kappeler P, Heistermann M (2008). Androgen and glucocorticoid level reflect seasonally  
583 occurring social challenges in male redfronted lemurs (*Eulemur fulvus rufus*). *Behavioral  
584 Ecology and Sociobiology* 62: 627–638.
- 585 Overdorff DJ, Rasmussen MA (1995). Determinants of nighttime activity in “diurnal” lemurid  
586 primates. In *Creatures of the Dark* (Alterman L, ed.), pp 61–84. New York, Plenum Press.
- 587 Pereira ME, Pond CM (1995). Organization of white adipose tissue in Lemuridae. *American  
588 Journal of Primatology* 35: 1–13.
- 589 Pereira ME, Strohecker RA, Cavigelli SA, Hughes CL, Pearson DD (1999). Metabolic strategy and  
590 social behavior in Lemuridae. In *New Directions in Lemur Studies* (Rakotosamimanana B,  
591 Rasamimanana H, Ganzhorn JU, Goodman S, eds.), pp 93–117. New York, Kluwer  
592 Academic/Plenum Publishers.
- 593 Pichon C, Simmen B (2015). Energy management in Crowned sifakas (*Propithecus coronatus*) and  
594 the timing of reproduction in a seasonal environment. *American Journal of Physical  
595 Anthropology* 158: 269–278.
- 596 Pinkus S, Smith, JNM, Jolly, A (2006). Feeding competition between introduced *Eulemur fulvus*  
597 and native *Lemur catta* during the birth season at Berenty reserve, Southern Madagascar. In  
598 *Ringtailed Lemur Biology. Developments in Primatology: Progress and Prospect* (Jolly A,  
599 Sussman RW, Koyama N, Rasamimanana H, eds.), pp 119–140. Springer, Chicago.
- 600 Pontzer H (2015). Energy expenditure in humans and other primates: a new synthesis. *Annual  
601 Review of Anthropology* 44: 169–187.

- 602 Pontzer H, Raichlen DA, Gordon AD, Schroepfer-Walkere KK, Hare B, O'Neill MC, Muldoong  
603 KM, Dunsworth HM, Wood BM, Isler K, Burkart J, Irwin M, Shumaker RW, Lonsdorf EV,  
604 Ross SR (2014). Primate energy expenditure and life history. *Proceedings of the National*  
605 *Academy of Sciences of the United States of America* 111: 1433–1437.
- 606 Pontzer H, Brown MH, Raichlen DA, Dunsworth H, Hare B, Walker K, Luke A, Dugas LR,  
607 Durazo-Arvizu R, Schoeller D, Plange-Rhule J, Bovet P, Forrester TE, Lambert EV, Emery  
608 Thompson M, Shumaker RW, Ross SR (2016a). Metabolic acceleration and the evolution of  
609 human brain size and life history. *Nature* 533: 390–392.
- 610 Pontzer H, Durazo-Arvizu R, Dugas LD, Plange-Rhule J, Bovet P, Forrester TE, Lambert EV,  
611 Cooper RS, Schoeller DA, Luke A (2016b). Constrained total energy expenditure and  
612 metabolic adaptation to physical activity in adult humans. *Current Biology* 26: 410–417.
- 613 Porter LJ, Hrstich LN, Chan BC (1986). The conversion of procyanidins and prodelphinidins to  
614 cyanidin and delphinidin. *Phytochemistry* 25: 223–230.
- 615 Price MP, Butler LG (1977). Rapid visual estimation and spectrophotometric determination of  
616 tannin content of *Sorghum* grain. *Journal of Agricultural and Food Chemistry* 25: 1268–1273.
- 617 Rasamimanana HR, Rafidinarivo E (1993). Feeding behavior of *Lemur catta* females in relation to  
618 their physiological state. In *Lemur Social Systems and Their Ecological Basis* (Kappeler PM,  
619 Ganzhorn JU, eds.), pp 123–133. New York, Plenum.
- 620 Rasamimanana H, Andrianome V, Rambeloarivony H, Pasquet P (2006). Male and female ring-  
621 tailed lemurs' energetic strategy does not explain female dominance. In *Ringtailed Lemur*  
622 *Biology. Developments in Primatology: Progress and Prospect* (Jolly A, Sussman RW,  
623 Koyama N, Rasamimanana H, eds.), pp 271–295. Springer, Chicago.
- 624 Richard AF, Dewar RE, Schwartz M, Ratsirarson J (2000). Mass change, environmental variability  
625 and female fertility in wild *Propithecus verreauxi*. *Journal of Human Evolution* 39: 381–391.
- 626 Rothman JM, Chapman CA, Van Soest PJ (2012). Methods in primate nutritional ecology: a user's  
627 guide. *International Journal of Primatology* 33: 542–566.

- 628 Sauther ML (1991). Reproductive behavior of free-ranging *Lemur catta* at Beza Mahafaly Special  
629 Reserve, Madagascar. *American Journal of Physical Anthropology* 84: 463–477.
- 630 Sauther ML (1998). Interplay of phenology and reproduction in ring-tailed lemurs: implications for  
631 ring-tailed lemur conservation. *Folia Primatologica* 69: 309–320.
- 632 Schmidt DA, Kerley MS, Porton IJ, Porter JH, Dempsey JL, Griffin ME, Ellersieck MR, Sadler  
633 WC (2005). Fiber digestibility by black lemurs (*Eulemur macaco macaco*). *Journal of Zoo and*  
634 *Wildlife Medicine* 36: 204–211.
- 635 Schmid J, Speakman JR (2000). Daily energy expenditure of the grey mouse lemur (*Microcebus*  
636 *murinus*): a small primate that uses torpor. *Journal of Comparative Physiology B* 170: 633–41.
- 637 Simmen B, Hladik A, Ramasiarisoa PL (2003). Food intake and dietary overlap in native *Lemur*  
638 *catta* and *Propithecus verreauxi* and introduced *Eulemur fulvus* at Berenty, Southern  
639 Madagascar. *International Journal of Primatology* 24: 949–968.
- 640 Simmen B, Tarnaud L, Hladik A (2012). Leaf nutritional quality as a predictor of primate biomass:  
641 further evidence of an ecological anomaly within prosimian communities in Madagascar.  
642 *Journal of Tropical Ecology* 28: 141–151.
- 643 Simmen B, Tarnaud L, Marez A, Hladik A (2014). Leaf chemistry as a predictor of primate biomass  
644 and the mediating role of food selection: a case study in a folivorous lemur (*Propithecus*  
645 *verreauxi*). *American Journal of Primatology* 76: 563–575.
- 646 Simmen B, Bayart F, Rasamimanana H, Zahariev A, Blanc S, Pasquet P (2010). Total energy  
647 expenditure and body composition in two free-living sympatric lemurs. *Plos ONE* 5: e9860.
- 648 Simmen B, Pasquet P, Masi S, Koppert GJA, Wells JCK, Hladik CM (2017). Primate energy input  
649 and the evolutionary transition to energy-dense diets in humans. *Proceedings of the Royal*  
650 *Society London B* DOI: 10.1098/20170577.
- 651 Sussman R (1974). Ecological distinctions in sympatric species of *Lemur*. In *Prosimian Biology*  
652 (Martin RD, Doyle GA, Walker AC, eds.), pp 75–108. London, Duckworth.
- 653 Tanaka M (2007). Habitat use and social structure of a brown lemur hybrid population in the

- 654 Berenty Reserve, Madagascar. *American Journal of Primatology* 69: 1189–1194.
- 655 Traina A (2001). Activity pattern and feeding behavior of ring-tailed lemurs (*Lemur catta*) at  
656 Berenty Reserve in Madagascar during the day and night. *Folia Primatologica* 72:188.
- 657 Wells JCK (2012). The capital economy in hominin evolution: how adipose tissue and social  
658 relationships confer phenotypic flexibility and resilience in stochastic environments. *Current*  
659 *Anthropology* 53: S466–S478.
- 660 Westerterp KR, Speakman JR (2008). Physical activity energy expenditure has not declined since  
661 the 1980s and matches energy expenditures of wild mammals. *International Journal of Obesity*  
662 32: 1256–1263.
- 663 Wright PC (1999). Lemur traits and Madagascar ecology: coping with an island environment.  
664 *Yearbook of Physical Anthropology* 42: 31–72.
- 665 Zinner D (1999). Relationship between feeding time and food intake in hamadryas baboons (*Papio*  
666 *hamadryas*) and the value of feeding time as predictor of food intake. *Zoo Biology* 18: 495–  
667 505.