Atmospheric nitrate export in streams along a montane to urban gradient

Ilann Bourgeois^{a,b,§}, Joel Savarino^a, Julien Némery^a, Nicolas Caillon^a, Sarah Albertin^a, Franck Delbart^c, Didier Voisin^a, and Jean-Christophe Clément^{b,d}

^aUniversité Grenoble Alpes, CNRS, IRD, Grenoble INP*, IGE, F-38000, Grenoble, France

^bUniversité Grenoble Alpes, CNRS, LECA, F-38000, Grenoble, France

^cUniversité Grenoble Alpes, CNRS, SAJF, F-38000, Grenoble, France

^dUniversité Savoie Mont Blanc, INRA, CARRTEL, F-74200, Thonon-Les Bains, France

* Institute of Engineering Univ. Grenoble Alpes

§ ilann.bourgeois@noaa.gov

Abstract

Nitrogen (N) emissions associated with urbanization exacerbate the atmospheric nitrogen influx to remote ecosystems – like mountains –, leading to well-documented detrimental effects on ecosystems (e.g., soil acidification, pollution of freshwaters). Here, the aim was to evaluate the importance and fate of N deposition in a watershed along a montane to urban gradient, using a multi-isotopic tracers approach ($\Delta^{17}\text{O}$, $\delta^{15}\text{N}$, $\delta^{18}\text{O}$ of nitrate, $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of water). In this setting, the montane streams had higher proportions of atmospheric nitrate compared to urban streams, and exported more atmospheric nitrate on a yearly basis (0.35 vs 0.10 kg-N ha⁻¹ yr⁻¹). In urban areas, nitrate exports were driven by groundwater, whereas in the catchment head nitrate exports were dominated by surface runoffs. The main sources of nitrate to the montane streams were microbial nitrification and atmospheric deposition whereas microbial nitrification and sewage leakages contributed most to urban streams. Based on the measurement of $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ -NO₃⁻, biological processes such as denitrification or N assimilation were not predominant in any streams in this study. The observed low $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ range of terrestrial nitrate (*i.e.*, nitrate not coming from atmospheric deposition) in surface water compared to literature suggests that atmospheric deposition may be underestimated as a direct source of N.

Keywords: Nitrogen, deposition, $\Delta^{17}\text{O}$, subalpine, isotope, Lautaret

Highlights

How is nitrogen deposition reflected in streams NO_3^- exports in the French Alps?

NO_3^- exports in seven streams along an altitude gradient were monitored for two years.

NO_3^- isotopic composition ($\Delta^{17}\text{O}$, $\delta^{15}\text{N}$, $\delta^{18}\text{O}$) was analyzed to apportion the sources.

Up to 21% of NO_3^- in montane streams and 5% in urban streams was unprocessed $\text{NO}_3^-_{\text{atm}}$.

Nitrification of atmospheric and mineralized NH_4^+ was the ubiquitous main NO_3^- source.

1. Introduction

Atmospheric nitrogen (N) deposition has increased 10-fold over the past century, increasingly contributing to the global N **availability** (Galloway et al., 2004). Anthropogenic activities such as fossil-fuel combustion, agriculture, and fertilizers use are responsible for this increase (Fowler et al., 2015; Galloway et al., 2008; Vitousek et al., 1997), with impacts **observed** in remote ecosystems (Hastings et al., 2009; Holtgrieve et al., 2011; Preunkert, 2003). **High N loading to the environment has been documented for many ecosystems** (Aber et al., 1989; Clark et al., 2017; Elser et al., 2009; Matson et al., 2002). To address this issue, global efforts are underway to alleviate N inputs into ecosystems, aiming at “minimizing the consequent harm to humans and the environment” (International Nitrogen Initiative, <http://www.initrogen.org>).

Nitrate concentrations, and fluxes in soils and streams have often been used to assess the N saturation status in watersheds (Aber et al., 1989; Baron and Campbell, 1997; Lovett and Goodale, 2011). However, N exports in stream depend on multiple parameters such as basin topography (Balestrini et al., 2013; Clow and Sueker, 2000), land-cover (Barnes et al., 2014; Williams et al., 2016) and land-management (Barnes and Raymond, 2010; Burns et al., 2009; Lefebvre et al., 2007). As streams integrate many processes at the watershed scale, **understanding the specific sources of NO_3^-** is important to (1) evaluate the respective contribution of biological and anthropogenic sources, (2) clarify the fate of deposited N in the environment, and (3) understand the origin and the development of N saturation.

Mountainous ecosystems are particularly sensitive to increased N inputs by atmospheric deposition (Baron et al., 2000, 2005, 2011), as they are historically N-limited (Kaye and Hart, 1997). Critical loads for these ecosystems are among the

lowest for pristine environments (Baron et al., 2011; Bowman et al., 2006; Nanus et al., 2017), making them vulnerable to long-range transport of atmospheric N emitted from distant sources (Mast et al., 2014; Wasiuta et al., 2015). Atmospheric deposition of N has been shown to contribute significantly, either directly or indirectly, to year-round NO_3^- exports from mountainous catchments (Hundey et al., 2016; Nanus et al., 2008), typically showing a pulse at spring as soils subsurface NO_3^- reservoirs are flushed by snowmelt water (Kendall et al., 1995; Williams et al., 2009; Williams and Melack, 1991).

Atmospheric deposition is also a major source of N to urban areas, which receive much higher loads than adjacent environments (Bettez and Groffman, 2013; Fang et al., 2011; Hall et al., 2014; Rao et al., 2014; Templer et al., 2015). Local and regional emissions and subsequent deposition of fuel-combustion derived NO_x and ammonia (NH_3) are responsible for this pattern (Galloway et al., 2004; Kean et al., 2000). NO_x compounds are oxidized into NO_3^- within hours (Beirle et al., 2011), then scavenged from the atmosphere by wet and dry deposition (Hertel et al., 2012). In the atmosphere, NH_3 is in equilibrium with ammonium (NH_4^+), the other primary component of bulk N deposition. Because of the particular topography of urban basins (extended impervious surface, rapid precipitation runoff), urbanization can lead to high NO_3^- exports in freshwater bodies (Groffman et al., 2004; Riha et al., 2014), with major ecological, economic and health consequences (Dodds et al., 2009).

This study aims at evaluating how atmospheric deposition of N is reflected in NO_3^- exports in several streams along a montane to urban gradient in the French Alps. Isotopic ($\Delta^{17}\text{O}$, $\delta^{15}\text{N}$, $\delta^{18}\text{O}$ of nitrate, $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of water) and hydro-chemical techniques were combined to evaluate the drivers of NO_3^- inputs and

removal in all streams. The first hypothesis was that due to a higher contribution from other sources (e.g., sewage, fertilizers) in urban areas compared to the mountains, coupled with higher local atmospheric N inputs, total NO_3^- and atmospheric nitrate ($\text{NO}_{3\text{ atm}}^-$) exports should increase along the gradient. The second hypothesis was that different $\text{NO}_{3\text{ atm}}^-$ export dynamics should be observed in urban streams compared to montane streams, due to different hydrological drivers that need to be determined. To test these hypotheses, (1) NO_3^- concentration and isotopic composition were determined in six streams and one lake, ranging from 2000m above sea level (a.s.l.) to 200m (a.s.l.) in the French Alps, (2) the annual export fluxes of $\text{NO}_{3\text{ atm}}^-$ and total NO_3^- were compared across sites along the gradient, (3) the hydrological drivers of the seasonal variability in $\text{NO}_{3\text{ atm}}^-$ proportion were identified and (4) the additional sources of NO_3^- in each watershed were determined.

2. Methods

2.1 Study site and selected streams

The Romanche Valley, located in the central French Alps, spreads from the Lautaret pass (2058m a.s.l.) down to Grenoble (250m a.s.l.) (Figure S1a and b). The Grenoble conurbation counts around 500 000 inhabitants, and is the biggest alpine metropolis in France.

Six streams and one lake were sampled from the Lautaret pass to Grenoble, draining watersheds of increasing size and with distinct geomorphic and biogeographic characteristics (Table 1). Two alpine streams were sampled at ~2000m a.s.l.. These drain the South exposed side of the Lautaret pass (S-upper montane, n = 157) and the North exposed side (N-upper montane, n=93), where for the latter the stream is mainly fed by glacier melt (Figure S1c). The mid montane stream (n=67) was sampled at ~1600 m a.s.l., 6km down from the Lautaret pass (Figure S1c). All three streams are tributaries of Romanche (lower montane stream, n=127) that was sampled 15km down from the Lautaret pass at ~1000m a.s.l., before ending in Chambon lake (Figure S1d). The lake (~ 900m a.s.l., n=29) is an artificial reservoir managed by the French national electricity company (EDF), 21km down from the Lautaret pass. The two urban streams (~210 m a.s.l.) were located, for one, upstream from Grenoble main urbanized area (upper urban stream, n=48) and, for the other, downstream (lower urban stream, n=45), and were separated by 8km (Figure S1e). The lower urban stream waters are derived from the lower montane stream (and the lake). The upper urban stream, Isère, has surface water from another alpine valley. The upper urban stream flows through residential areas, gardens, and some arable fields prior to the sampling location whereas the lower

urban site abuts industrial developments as well as residential neighborhoods
(Dutordoir, 2014).

2.2 Deposition and streams sampling, discharge and conductivity

Atmospheric wet and dry deposition samples were collected using a WADOS Kroneis collector (GmbH Austria) at the S-upper montane site from April 2016 to April 2017. A moving lid – connected to heated moisture sensor – ensured the separated collection by covering alternatively the dry funnel (under rain or snow conditions) and the wet funnel (the rest of the time). The deposition collector was installed on the roof of the Lautaret pass Alpine Research Station (~210 m a.s.l.), <https://www.jardinalpindulautaret.fr>) to reduce the impact of ground turbulence and input of soil particles. Samples were collected approximately every three weeks. Precipitation samples were kept at 4°C during the sampling period, then frozen until further analysis. After exposure to the atmosphere, the dry deposition funnel was rinsed with 500 mL of MilliQ water, and the sample was stored at -20°C.

Stream and lake water was sampled at all sites according to the Niwot Ridge LTER protocol (Williams and Melack, 1991). In short, samples (total n = 566) were collected manually in 1L Nalgene bottles, previously washed 3 times with MilliQ water (18.2 MΩ.cm⁻¹ at 25°C) and rinsed three times with stream water before collection. All samples were then kept at -20°C until further treatment. Water samples were collected on a weekly to monthly basis from January 2015 to December 2016, with increased sampling frequency in two of the upper tributaries from April to May 2016. During this intensive sampling period, water was collected using three automatic water samplers (Teledyne ISCO® 3700) every two or three hours at the S-upper and lower montane sites. Water conductivity, shown to be a good proxy of discharge

(Weijs et al., 2013), and water temperature were also intensively monitored (2 minutes stepwise) over that period using CTD-Diver[®] sensors. Spring sampling was sporadic in the N-upper montane site because of persistent snow cover. Winter sampling was not possible at the lake site as it was frozen.

Discharge data was obtained from the national discharge-monitoring network (Banque Hydro) or measured every hour for both lower montane and urban site. Stream stage at the upper urban site was monitored every 30 minutes using water level gauge sensor (OTT[®]) at Grenoble Campus station (more details in Mano, 2008; Némery et al., 2013). Discharge was estimated from stage measurements by using rating curve (water level vs discharge) (Némery et al., 2013). Continuous discharge could not be measured at the upper and mid montane sites **due to the torturous nature of the streams**. Precipitation data were obtained from the Alpine Mountain Meteorological Office Network (www.romma.fr) at the upper montane site and using a rain gauge (Précis Mécaniques[®]) on a building roof of the Grenoble Campus, close to the upper urban site.

2.3 Chemical and isotopic analysis

All stream and deposition samples were left to unfreeze at ambient temperature prior to being filtered using 0.45µm Whatman GD/X syringe filters linked to a peristaltic pump. They were subsequently analyzed for ion concentrations ([NH₄⁺], [NO₃⁻], [SO₄²⁻], [Cl⁻], [Ca²⁺]) using a colorimetric technique (Gallery Plus, Thermo Fisher Scientific, Waltham, Massachusetts, USA). **An analytical error <0.01 mg L⁻¹ was determined for all ions based on the standard deviation of commercial control solutions (ThermoFisher[®]) analyzed as unknowns.** Major ions such as sulfate and calcium and chloride are often used as hydro-chemical tracers of water

flowpaths or water sources (Briand, 2014; Devito et al., 2000; Hall et al., 2016; Stoewer et al., 2015), and were measured to identify the dominating hydrological regimes in the studied sites.

All samples were concentrated on an anionic resin (0.6mL AG 1-X8 resin, Cl^- -form, Bio-Rad) with recovery efficiency over 98.5% (Erbland, 2011) and eluted with 10mL of a 1M NaCl solution for isotopic analysis (Templer and Weathers, 2011). Isotopic analyses of NO_3^- ($\Delta^{17}\text{O}$, $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$) were conducted on an MAT 253 IRMS using an adapted version of the denitrifier method (Kaiser et al., 2007; Morin et al., 2008). More details about the experimental setup can be found elsewhere (SI Text). The analytical errors – calculated as the standard deviation of the residuals from the linear regression between raw data of the reference standards (USGS 32, USGS 34 and USGS 35) and their accepted values – were ± 0.4 , 1.5 and 0.3‰ for $\Delta^{17}\text{O}$, $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ of NO_3^- , respectively. Slopes and intercepts of the linear regressions are then used to convert raw sample data into calibrated values (VSMOW scale for O isotopes, and N_2 -air scale for N isotopes).

A subset of stream, rain, and snow samples (collected in winter 2015-2016) were sent for $\delta^2\text{H}$ and $\delta^{18}\text{O}$ analyses on a Picarro L2130-i at the Laboratoire des Sciences et du Climat (LSCE) in Paris, France. The analytical error – inferred from the replicate analysis of internal standards calibrated against V-SMOW2 et V-SLAP2 standards – was ± 0.7 and 0.2 ‰ for $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of H_2O , respectively.

2.4 NO_3^- sources apportionment

A number of previous studies have used the dual isotope approach ($\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ of NO_3^-) to track the spatio-temporal variability of sources contribution to NO_3^- pools in a large variety of environmental matrixes (Campbell et al., 2002; Durka et al.,

1994; Elliott et al., 2009; Yang and Toor, 2016). Biochemical processes such as denitrification or assimilation have also been shown in laboratory experiments to distinguishably enrich residual NO_3^- in heavier isotopes, here ^{15}N and ^{18}O (Granger et al., 2004, 2010; Treibergs and Granger, 2017), although this enrichment can be diluted by newly nitrified NO_3^- with low N and O isotopic values (Granger and Wankel, 2016; Mayer et al., 2002). Environmental studies also reported characteristic NO_3^- isotopic enrichment for denitrification (Clément et al., 2003; Fang et al., 2015; Wexler et al., 2014), assimilation (Emmerton et al., 2001; Estrada et al., 2017; X.-Y. Liu et al., 2013) or photolysis (Frey et al., 2009; Shi et al., 2015; Ye et al., 2016). However, the isotopic fingerprint of biological processes can lead to inaccurate NO_3^- source apportionment in some cases, especially in delineating the respective contribution of the microbial and the atmospheric sources (Michalski et al., 2004; Riha et al., 2014; Rose et al., 2015a).

In the past few years, a growing number of studies have used an isotopic particularity of $\text{NO}_3^-_{atm}$ to quantify the contribution of atmospheric deposition to terrestrial N pools (Costa et al., 2011; Hundey et al., 2016; Tsunogai et al., 2014). $\text{NO}_3^-_{atm}$ is enriched in ^{17}O due to its production pathways (*i.e.*, oxidation of NO_x by O_3), showing a deviation from the Terrestrial Fractionation Line (Thiemens, 2006). $\Delta^{17}\text{O}$ is a quantification of this deviation, calculated as $\Delta^{17}\text{O} = \delta^{17}\text{O} - 0.52 * \delta^{18}\text{O}$ in the present work. $\Delta^{17}\text{O}$ value of $\text{NO}_3^-_{atm}$ generally ranges between 20 and 35‰ in temperate latitudes (Morin et al., 2009; Savarino et al., 2007), whereas $\Delta^{17}\text{O}$ value of NO_3^- from all other sources (industrial fertilizers, nitrification) or of biologically processed $\text{NO}_3^-_{atm}$, is 0‰ (Michalski et al., 2004, 2015). In this study, these two distinct $\Delta^{17}\text{O}$ values were used as end-members in a simple mixing model to quantify unprocessed $\text{NO}_3^-_{atm}$ proportion (f_{atm}) in streams, according to Michalski et al.

(2004):

$$(Eq. 1) \quad f_{atm} = (\Delta^{17}\text{O-NO}_3^- \text{ sample} / \Delta^{17}\text{O-NO}_3^- \text{ atm})$$

The atmospheric fraction (f_{atm}) and the terrestrial fraction of NO_3^- ($f_{ter} = 1 - f_{atm}$) can be used to remove the isotopic influence of $\text{NO}_3^- \text{ atm}$ on samples, and allow for a better interpretation of biological processes that affect $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ (Dejwakh et al., 2012; Michalski et al., 2004; Riha et al., 2014; Tsunogai et al., 2010). $\delta^{18}\text{O}$ vs $\delta^{15}\text{N}$ plots have been intensively used to evaluate sources of N and potential processes (denitrification, assimilation) in the environment (Burns et al., 2009; Durka et al., 1994; Griffiths et al., 2016; Kendall et al., 1995; X.-Y. Liu et al., 2013). However, because of strongly distinct $\delta^{18}\text{O-NO}_3^- \text{ atm}$ values compared to $\delta^{18}\text{O-NO}_3^- \text{ ter}$, even a low f_{atm} can lead to scatter in a dual-isotope plot. Removing the atmospheric $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ components from environmental samples leads to an easier assessment of biological processes, such as assimilation or denitrification. This isotope correction was applied on the samples using the isotopic mass balances from Riha et al. (2014):

$$(Eq.2) \quad \delta^{18}\text{O}_{ter} = (\delta^{18}\text{O}_{sample} - \delta^{18}\text{O}_{atm} * f_{atm}) / f_{ter}$$

$$(Eq.3) \quad \delta^{15}\text{N}_{ter} = (\delta^{15}\text{N}_{sample} - \delta^{15}\text{N}_{atm} * f_{atm}) / f_{ter}$$

$\delta^{15}\text{N}_{atm}$ and $\delta^{18}\text{O}_{atm}$ were inferred from NO_3^- in wet and dry atmospheric deposition (see section 3.2). A formal propagation of uncertainty was performed to assess the error associated with individual values of f_{atm} , $\delta^{18}\text{O-NO}_3^- \text{ ter}$ and $\delta^{15}\text{N-NO}_3^- \text{ ter}$. Although

the error associated with individual values can be large (Figure S2), averaging across samples throughout the year results in a robust estimate of annual f_{atm} , $\delta^{18}\text{O}-\text{NO}_3^-$ and $\delta^{15}\text{N}-\text{NO}_3^-$ values (Dejwakh et al., 2012).

2.5 Fluxes of NO_3^- and $\text{NO}_3^-_{atm}$

NO_3^- yields (F_m in kg-N w^{-1}) were calculated at each site – when possible – on a weekly basis using the mean weekly discharge (Q_m in $\text{m}^3 \text{s}^{-1}$) and instantaneous concentrations ($[\text{NO}_3^-]_i$ in mg-N L^{-1}), as follows:

$$\text{(Eq.4)} \quad F_m = Q_m * [\text{NO}_3^-]_i$$

Total specific annual stream loads of NO_3^- (F_{tot} in $\text{kg-N ha}^{-1} \text{yr}^{-1}$) were calculated as the annual sum of weekly NO_3^- exports during the study period (2015 and 2016), as follows:

$$\text{(Eq.5)} \quad F_{tot} = (\sum_m F_m) / (\text{watershed area})$$

Finally, total annual $\text{NO}_3^-_{atm}$ loads (F_{atm}) were calculated for each year by multiplying the total annual stream yield of NO_3^- (F_{tot}) by the discharge-weighted annual mean f_{atm} in the stream (see Eq. 1), according to Tsunogai et al. (2014, 2010):

$$\text{(Eq. 6)} \quad F_{atm} = F_{tot} * f_{atm}$$

When $[\text{NO}_3^-]$ or $\Delta^{17}\text{O}$ values were not available for a given date ($n=4$, 2 and 2 for the lower montane, upper urban and lower urban streams, respectively), they were

replaced by the mean $[\text{NO}_3^-]$ or $\Delta^{17}\text{O}-\text{NO}_3^-$ values measured on the previous and the following dates, to capture temporal variations due to hydrological events. Note that the use of instantaneous $[\text{NO}_3^-]$ to calculate weekly streams NO_3^- loads might over or under estimate the real NO_3^- yield in streams.

2.6 Statistical analysis

A Mann-Whitney test was applied to surface water data to determine whether mean concentrations and isotopic values were significantly different between streams. A Spearman test was applied to evaluate the correlation between stream water parameters (typically $\Delta^{17}\text{O}$, $\delta^{18}\text{O}$, $\delta^{15}\text{N}$ and ion concentrations). Differences and correlations were considered significant when the p value was ≤ 0.05 . All statistical analyses were conducted using R software (v3.2.3). Unless specified otherwise, the uncertainty accompanying the mean values is the standard deviation.

3. Results

3.1 Precipitation and discharge

Cumulated precipitation was 537 and 609mm in 2015 and 804 and 740mm in 2016 at the upper montane and the urban sites, respectively. Discharge at the lower montane site reflects a snowmelt influenced hydrological regime. It peaked up to $30 \text{ m}^3 \text{ s}^{-1}$ in spring, and was significantly higher from April to October compared to the rest of the year. Discharge at both urban sites was consistent with a hydrological regime driven by snowmelt in spring-summer (main maximum), and rainfall in autumn (secondary maximum), a dual contribution similar to what was obtained in the Colorado Front Range (Cowie et al., 2017). At the upper urban site, discharge peaked at $900 \text{ m}^3 \text{ s}^{-1}$, whereas maximum discharge was $300 \text{ m}^3 \text{ s}^{-1}$ at the lower

urban site, a difference due to the streams draining two distinct watersheds (Figure S3).

3.2 Comparison of dissolved solutes across sites

Solutes concentration varied by up to 3 orders of magnitude among precipitation and streams samples (Table S2 and Figure 1). $[\text{NH}_4^+]$ was low in both dry and wet deposition at the S-upper montane site, and mostly (>90%) under the detection limit ($<0.02 \text{ mg L}^{-1}$) in all streams. $[\text{NO}_3^-]$ was significantly higher in rain samples compared to stream samples. The highest annual mean $[\text{NO}_3^-]$ was measured in the lower urban stream ($1.4 \pm 0.5 \text{ mg L}^{-1}$) and the lowest in the S-upper montane stream ($0.2 \pm 0.2 \text{ mg L}^{-1}$). $[\text{NO}_3^-]$ at the N-upper montane, mid montane, lower montane and upper urban sites were not statistically different (1.4 ± 1.3 , 1.1 ± 0.6 , 1.0 ± 0.6 and $1.1 \pm 0.4 \text{ mg L}^{-1}$, respectively). $[\text{SO}_4^{2-}]$ and $[\text{Ca}^{2+}]$ were both very low in precipitation relative to streams. The S-upper montane stream exhibited the highest annual mean $[\text{SO}_4^{2-}]$ and $[\text{Ca}^{2+}]$ (98.5 ± 39.5 and $44.5 \pm 18.2 \text{ mg L}^{-1}$, respectively), followed by the upper urban stream (74.1 ± 26.9 and $35.2 \pm 13.1 \text{ mg L}^{-1}$, respectively), and the lowest concentrations were measured at the N-upper montane site (22.1 ± 35.9 and $16.5 \pm 16.0 \text{ mg L}^{-1}$, respectively). $[\text{Cl}^-]$ was very low in precipitation samples and in most montane streams (mean values all $< 3 \text{ mg L}^{-1}$), respectively. Higher annual mean $[\text{Cl}^-]$ were measured in the urban streams (4.7 ± 1.8 and $16.6 \pm 12.3 \text{ mg L}^{-1}$ for the upper and lower urban stream, respectively).

3.3 Isotopic composition of NO_3^- in precipitation and streams

Mean annual $\text{NO}_3^-_{\text{atm}}$ isotopic values (Table S1) were non-significantly higher in dry deposition than in wet deposition (significant only for $\delta^{15}\text{N-NO}_3^-$), a pattern

widely monitored at temperate latitudes (Beyn et al., 2014; Freyer, 1991; Mara et al., 2009). Mean and associated standard deviation of $\Delta^{17}\text{O-NO}_3^-$ in precipitation and dry deposition were 26.4 ± 3.2 and 24.7 ± 3.5 ‰, respectively. These values are in the same range as previously reported data for $\Delta^{17}\text{O-NO}_3^-_{atm}$ (Costa et al., 2011; Michalski et al., 2004; Tsunogai et al., 2010, 2016). The mean $\Delta^{17}\text{O}$ value for total deposition was 25.6 ± 3.3 ‰, and was used to quantify the atmospheric component of NO_3^- pools in streams (see section 2.4, Eq. 1). Mean annual total $\delta^{18}\text{O-NO}_3^-_{atm}$ (70.8 ± 7.2 ‰) and $\delta^{15}\text{N-NO}_3^-_{atm}$ (-4.8 ± 5.9 ‰) are also consistent with a reservoir of exclusively $\text{NO}_3^-_{atm}$ (Kendall et al., 2007), and these values were used to correct samples from their atmospheric $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ components (see section 2.4, Eq.2 and 3).

Significantly higher annual mean $\Delta^{17}\text{O-NO}_3^-$ was measured in montane streams compared to urban streams, with the exception of the mid montane site (1.0 ± 0.7 ‰), which exhibited the lowest values (Figure 2a). Both upper and lower urban, along with the mid montane streams were also characterized by the highest annual mean $\delta^{15}\text{N-NO}_3^-$ (3.7 ± 1.0 ‰, 4.9 ± 5.6 ‰ and 4.0 ± 2.0 ‰, respectively) (Figure 2c). The lake, fueled by the lower montane stream, presented the highest annual mean $\Delta^{17}\text{O-NO}_3^-$ (5.1 ± 1.8 ‰) and $\delta^{18}\text{O-NO}_3^-$ (8.6 ± 6.2 ‰), and the lowest $\delta^{15}\text{N-NO}_3^-$ (-0.2 ± 2.4 ‰) (Figure 2a, b and c). It must be stressed out again that the lake was only sampled when unfrozen (*i.e.*, late spring, summer and autumn), potentially biasing NO_3^- mean isotopic values at this site.

3.4 Trends in isotope composition of NO_3^- across sites

Seasonal variations of stream NO_3^- isotopes and corresponding f_{atm} are shown in Figure 3. The three montane streams followed a similar pattern of seasonal variations in 2015 and 2016. At the S-upper montane site, $\Delta^{17}\text{O}-\text{NO}_3^-$ and $\delta^{15}\text{N}-\text{NO}_3^-$ were significantly negatively correlated, with brief but high $\Delta^{17}\text{O}-\text{NO}_3^-$ peaks in spring, followed by a slow but continuous increase peaking late summer/beginning of autumn. At the N-upper site, a saw tooth pattern in summer and autumn occurred right after a small but temporally wider increase at snowmelt, in early summer. $\Delta^{17}\text{O}-\text{NO}_3^-$ and $\delta^{15}\text{N}-\text{NO}_3^-$ in the mid montane stream presented little variations throughout the seasons, apart from a raise in $\delta^{15}\text{N}-\text{NO}_3^-$ in spring/summer 2016. $\Delta^{17}\text{O}-\text{NO}_3^-$ and $\delta^{15}\text{N}-\text{NO}_3^-$ were significantly negatively correlated in the lower montane stream and in the lake, with significantly higher $\Delta^{17}\text{O}-\text{NO}_3^-$ from April to October – relative to the rest of the year – in a very similar way to discharge (Figure S3). Finally, $\Delta^{17}\text{O}-\text{NO}_3^-$ and $\delta^{15}\text{N}-\text{NO}_3^-$ were not correlated and did not vary much in urban streams, except for some occasional high $\delta^{15}\text{N}$ peaks in spring and winter 2015 at the lower urban site.

4. Discussion

4.1 NO_3^- exports in streams

4.1.1 $[\text{NO}_3^-]$ across the altitude gradient

There were little differences in streams $[\text{NO}_3^-]$ along the elevation gradient despite an expected higher contribution from anthropogenic activities to streams N pool closer to urbanized areas (Groffman et al., 2004; Tsunogai et al., 2016). The 0.2-1.4 mg L^{-1} mean $[\text{NO}_3^-]$ range at the montane sites is consistent with the 0.6-1.6 mg L^{-1} range in other Alpine valleys (reviewed in Balestrini et al., 2013), and the 0.9-1.5 mg L^{-1} range in elevated sites of the Colorado Front Range (Baron and Campbell, 1997; Campbell et al., 1995). The N-upper montane site had a much wider $[\text{NO}_3^-]$

range than the S-upper montane site, which is attributed to glacier melt influence (Barnes et al., 2014; Williams et al., 2007). $[\text{NO}_3^-]$ in urban streams was low when compared to other concentrations measured in urban settings, such as the Tucson basin where values up to 62 mg L^{-1} were determined (Dejwakh et al., 2012) or up to 26 mg L^{-1} in the Gwynns Falls watershed (Groffman et al., 2004).

4.1.2 $\text{NO}_3^-_{atm}$ contribution to total NO_3^-

Atmospheric deposition contributed significantly more to NO_3^- pool in streams at the montane sites relative to the urban sites – exception being the mid montane site (Figure 2). The Chambon lake showed the highest f_{atm} , although it is possible that sampling of this reservoir exclusively in summer resulted in a biased estimate of mean annual $\text{NO}_3^-_{atm}$. High $\text{NO}_3^-_{atm}$ inputs in montane streams (up to 47%) clearly outline the influence of the melting snowpack releasing its load of atmospheric nitrate (see section 4.2.1), as widely reported elsewhere (Kendall et al., 1995; Ohte et al., 2004; Pellerin et al., 2012; Tsunogai et al., 2014, 2016). The mid montane stream exported the lowest proportion of $\text{NO}_3^-_{atm}$. Smoother topography gradient on this watershed (slopes are 17%, Table 1) can account for this by impeding direct runoff to stream during hydrological extremes (e.g., snowmelt), thus exerting a strong control over NO_3^- isotopic values in the stream (Balestrini et al., 2013; Clow and Sueker, 2000). In addition, a smoother topography gradient also drives a higher $\text{NO}_3^-_{atm}$ residence time in soils that would overwrite its $\Delta^{17}\text{O}$ signature when recycled by microorganisms. The low f_{atm} in urban streams is dissimilar to others results that have shown an increasing f_{atm} in streams along an urbanization gradient (Riha et al., 2014), or high $\text{NO}_3^-_{atm}$ inputs in urban runoff (Yang and Toor, 2016). In urban areas, the combination of more impervious surface area coupled to higher atmospheric

deposition inputs is a strong driver that was expected to promote a higher proportion of $\text{NO}_3^-_{\text{atm}}$ in urban streams, relative to montane sites. Indeed, NO_3^- atmospheric concentrations in Grenoble's urban area are 10 to 100 times higher than atmospheric concentrations observed at the Lautaret pass (Bourgeois, 2017; Tomaz et al., 2017), which should have promoted higher $\text{NO}_3^-_{\text{atm}}$ loads in urban streams. However, higher total $\text{NO}_3^-_{\text{atm}}$ loads are calculated at the lower montane site where $\text{NO}_3^-_{\text{atm}}$ represented 21% of the total annual NO_3^- exports (Table 2).

Elevated N deposition in the Alps – also illustrated by higher $[\text{NO}_3^-]$ in wet and dry deposition compared to streams (Table S2) – is a well-documented phenomenon (Kirchner et al., 2014; Rogora et al., 2006). Sustained nitrogen deposition in addition to the steep slopes of the lower montane watershed (Table 1) can account for the higher total $\text{NO}_3^-_{\text{atm}}$ flux at the montane site. Grenoble conurbation land cover is mainly constituted of forest and lower vegetation (64%) followed by agricultural fields (24%), and impervious surfaces (10%) are only a minor surface occupation (UE-SOeS, CORINE Land Cover, 2006). $\text{NO}_3^-_{\text{atm}}$ inputs are thus likely to move to multiple sinks, be biologically recycled or lost *via* different processes (e.g., denitrification, photolysis (Clément et al., 2003; Ye et al., 2016)) before being collected by the urban streams (Rao et al., 2014). Furthermore, a strong decoupling between atmospheric N deposition and urban soils N cycling has recently been proposed, pointing at a distinct urban biogeochemical N cycle – that entails that land-management practice, not N deposition, is the ultimate driver of N cycling – and that would need further investigation (Decina et al., 2017). Additionally, in France urban rainwater and domestic wastewater are traditionally collected in the same pipe system before being purified in a sewage treatment plant downstream of the cities. This would also limit direct rainwater runoff to the streams, as illustrated by the statistically similar $[\text{NO}_3^-]$ in

urban streams compared to montane streams (Figure 1). One, or a combination of several of these factors would lower $\text{NO}_3^-_{atm}$ exports in urban streams, even lower than montane streams $\text{NO}_3^-_{atm}$ exports in this case. Therefore, the low f_{atm} in urban streams certainly reflects the recycling and replacement of $\text{NO}_3^-_{atm}$ by terrestrial NO_3^- sources along the smoother topography gradient, in a similar way to the mid-montane stream.

4.2 Links between hydrology, topography and $\text{NO}_3^-_{atm}$ exports

4.2.1 Seasonal variations of $\text{NO}_3^-_{atm}$ proportion in streams

Seasonal variations of $\text{NO}_3^-_{atm}$ exports at the montane sites reflected the influence of hydrologic and topographic features of the basins, and are thoroughly discussed elsewhere (Bourgeois, 2017). The S-upper montane site exports are governed by quick runoff to streams in the early stages of spring, followed by a diffuse resurgence of infiltrated snowmelt water in summer and autumn. The period between these two event lasts about 5 months both in 2015 and in 2016, an indicator of either the water transit time in the calcareous substrate of the watershed or of the aquifer refill time before flushing to the stream. The N-upper montane stream is fueled by glacier melt, leading to higher $[\text{NO}_3^-]$ than its southern counterpart by 1 order of magnitude (Figure 1). This is a consequence of the permafrost thawing at that site (Bodin et al., 2009), thus increasing nitrification and/or mobilization of stored N from disturbed soils (Barnes et al., 2014; Hood and Scott, 2008; Louiseize et al., 2014). Occasional higher f_{atm} at the N-upper site are triggered by rainstorm events (Figure S3). The lower montane stream and the lake had very similar seasonal variations of $\Delta^{17}\text{O-NO}_3^-$, featuring the gradual melting of upstream snowpack throughout the growing season (May to October). The mid montane and the urban

streams showed no marked seasonal variation, **because of** the buffering effect of soils and the higher N cycling rates inherent to more intensive land management treatments **that prevented** $\text{NO}_3^-_{atm}$ inputs to streams (Table 1).

4.2.2 Groundwater and surface runoff respective influence $\text{NO}_3^-_{atm}$ proportion in streams

Significant correlations between rock-derived solutes (SO_4^{2-} and Ca^{2+}) in all streams (Figure S4) indicate groundwater contribution to streams as a unifying feature along the montane to urban gradient. Varying concentrations of rock-derived solutes are in good agreement with the geologic characteristics of each watershed (*i.e.*, highest concentrations in the S-upper montane and the upper urban streams draining calcareous basins, lowest concentrations in the N-upper montane stream draining a less erodible bedrock, see Table 1 and Figure 1). In this study, year-round detectable $\Delta^{17}\text{O}-\text{NO}_3^-$ translate the continuous $\text{NO}_3^-_{atm}$ exports in all streams (Figure 3), even during baseflow when streams are mostly fueled by groundwater. Most studies focusing on NO_3^- dynamics in runoff reported $\text{NO}_3^-_{atm}$ only during stormflow or snowmelt periods (Burns et al., 2009; Campbell et al., 2002; Pellerin et al., 2012). However, these studies used the dual isotope approach (*i.e.*, $\delta^{18}\text{O}$ and $\delta^{15}\text{N}$ of NO_3^-), which **does not** account accurately for $\text{NO}_3^-_{atm}$ in streams compared to the $\Delta^{17}\text{O}$ tracer (Michalski et al., 2004; Rose et al., 2015a). In a growing body of work using $\Delta^{17}\text{O}-\text{NO}_3^-$ in streams, $\text{NO}_3^-_{atm}$ presence during baseflow is a unifying feature (Rose et al., 2015a; Sabo et al., 2016; Tsunogai et al., 2016). In all these studies, f_{atm} ranged from 1-12% in baseflow, except for a coniferous watershed in West Virginia, which led to f_{atm} of 54% in stream, due to low sampling resolution ($n=4$). Here, f_{atm} ranged from 1-20% (with a median associated error of 3%, Figure S2) in all streams

during the dormant season (in grey, Figure 3), somewhat higher than the literature data range – because of the extended influence of snowmelt in mountainous sites during the growing season. The high $\Delta^{17}\text{O}-\text{NO}_3^-$ peaks at the S-upper montane site in March 2015 are due to a shorter winter this year.

However, even if groundwater contributed at all sites to streams water, its influence on streams ionic exports **had most impact** in the urban reaches. No or near static correlations between either NO_3^- isotopes or conservative ions and discharge (Figure 4) imply that a geochemically equilibrated groundwater reservoir dominates stream exports at the urban sites, instead of surface runoff (Hall et al., 2016).

Oppositely, $\Delta^{17}\text{O}-\text{NO}_3^-$ increased significantly with discharge at the lower montane site, highlighting the influence of snowmelt on $\text{NO}_3^-_{atm}$ exports (Figure 4a).

Decreasing weathering products concentrations with high flow also evidence the **decreasing contribution of groundwater towards more surficial water at snowmelt** (Figure 4d and f). No significant correlation of $[\text{NO}_3^-]$ with discharge in any stream implies a homogenized NO_3^- pool in groundwater reservoirs by infiltrated snowmelt inputs. In line with this hypothesis, a dual isotope plot of water **illustrates** that snowmelt more than rain fueled the streams, even down to the outskirts of Grenoble (Figure S5). Other studies also reported similar findings of underground reservoirs fed by snowmelt in mountains vicinity (Brooks et al., 1999; Hall et al., 2016; Williams et al., 2009).

4.2.3 A focus on snowmelt period in montane streams

The intensive sampling campaign in spring 2016 at the S-upper montane and the lower montane sites provided no evidence of diurnal pattern for any solutes in the streams (data not shown). However, $\Delta^{17}\text{O}-\text{NO}_3^-$ **did follow** a diurnal pattern at the S-

upper montane site, a pattern not seen in the lower montane stream (Figure S6a). Conductivity (at both sites) and discharge (lower montane site only) showed very pronounced diurnal variations, highlighting the hydrological influence of snowmelt during the day (Figure S6b and c), also consistent with previous studies in the same location (Mano et al., 2009). At the lower montane site, discharge and conductivity were negatively correlated (Figure 5d), as their diurnal variations were driven by distinct hydrological regime (*i.e.*, baseflow for conductivity, snowmelt for discharge). That conductivity is controlled by baseflow is also supported by significant positive correlations between weathering solutes and conductivity at the S-upper montane site (Figure 5c). Therefore, assuming a similar relationship between conductivity and discharge **is true** at the S-upper montane stream, and considering that conductivity and $\Delta^{17}\text{O-NO}_3^-$ were also negatively correlated (Figure 5a), snowmelt-induced discharge **in daytime** is certainly **the driver** of the diurnal variations of $\text{NO}_3^-_{\text{atm}}$ exports. Oppositely to another study showing in stream diurnal $[\text{NO}_3^-]$ variations at snowmelt in northern Vermont (Pellerin et al., 2012), **no significant temporal trend in any of the two streams was observed here. Yet, the lower montane stream was only on the rising limb of discharge when sampled**, at the very beginning of the snowmelt period. **Therefore**, the absence of observed diurnal cycle at this site at this time may not be representative of day-night variations later in the season.

4.3 Isotopic composition of $\text{NO}_3^-_{\text{ter}}$

4.3.1 NH_4^+ nitrification is the main source of $\text{NO}_3^-_{\text{ter}}$ in all streams

Isotopic values of $\text{NO}_3^-_{\text{ter}}$ (*i.e.*, NO_3^- corrected from the atmospheric component according to Eq. 2 and 3) are reported in a dual isotope ($\delta^{18}\text{O}$ and $\delta^{15}\text{N}$) plot in Figure 6. This allows the evaluation of the respective contribution of terrestrial

sources of NO_3^- to streams. Here, most $\delta^{15}\text{N}-\text{NO}_3^-$ in streams span a 0 to 7‰ range, delineating nitrification of NH_4^+ as the main source of stream NO_3^- contents (Figure 6). This is in line with watersheds land cover consisting principally in natural or semi-natural vegetation (Table 1), where NO_3^- fertilizers are rarely used. Only very few samples ($n = 4$) had both $\delta^{15}\text{N}-\text{NO}_3^-$ and $\delta^{18}\text{O}-\text{NO}_3^-$ characteristic of NO_3^- fertilizers. It is complicated to segregate the substrate of nitrification (*i.e.*, deposited atmospheric NH_4^+ ($\text{NH}_4^+_{\text{atm}}$) or mineralized soil NH_4^+ ($\text{NH}_4^+_{\text{org}}$)) that led to the production of NO_3^- in streams. Indeed, nitrification of $\text{NH}_4^+_{\text{atm}}$ and $\text{NH}_4^+_{\text{org}}$ produces NO_3^- with overlapping $\delta^{15}\text{N}$ ranges (Kendall et al., 2007). However, a fraction of samples, mostly from montane streams, had $\delta^{15}\text{N}-\text{NO}_3^-$ lower than 0‰, consistent with nitrification of $\text{NH}_4^+_{\text{atm}}$ (Kendall et al., 2007; T. Liu et al., 2013).

4.3.2 Biological removal of NO_3^- was not predominant in any stream

Weak negative correlations between $\delta^{18}\text{O}-\text{NO}_3^-$ and $\delta^{15}\text{N}-\text{NO}_3^-$ at all sites, exception made of the lower urban site, indicate that biological processes such as denitrification or assimilation do not control NO_3^- cycling in streams. Indeed, such processes, if dominant, should enrich residual $\delta^{18}\text{O}-\text{NO}_3^-$ and $\delta^{15}\text{N}-\text{NO}_3^-$ along a positively correlated line (Granger et al., 2008, 2010; Kendall et al., 2007), in a similar fashion to what is observed at the lower urban site. Yet, that $\delta^{15}\text{N}-\text{NO}_3^-$ was significantly correlated to $[\text{Cl}^-]$ in this stream (data not shown) rather proves that the high $\delta^{15}\text{N}-\text{NO}_3^-$ values are due to interception of sewage-originated wastewater (Hall et al., 2016). Decreasing $\delta^{15}\text{N}-\text{NO}_3^-$ and $[\text{Cl}^-]$ with discharge (Figure 4b and e) at this specific site also supports a groundwater origin of these nitrate pollution episodes.

4.3.3 Low $\delta^{18}\text{O}-\text{NO}_3^-$ calls for a more systematic use of the $\Delta^{17}\text{O}$ tracer

$\delta^{18}\text{O}-\text{NO}_3^-$ values as low as -20‰ were obtained after removing the atmospheric imprint (Figure 6 and Figure S7). The mean $\delta^{18}\text{O}-\text{NO}_3^-$ ranged between -6.3 and -8.0‰ in the mountainous streams, significantly lower than previously reported values of terrestrial $\delta^{18}\text{O}-\text{NO}_3^-$ in freshwater systems (from -2.9 to -4.2‰ in Tsunogai et al. (2016, 2014, 2010)) or produced through in vitro incubation experiments (> 0‰ in Burns and Kendall (2002), Mayer et al. (2001) and Spoelstra et al. (2007)). The mean $\delta^{18}\text{O}-\text{NO}_3^-$ range between -2.6 and -2.8‰ in the urban streams, far lower than values > 9‰ inferred in other urban areas (Riha et al., 2014) or mostly positive values in a semiarid urban setting (Dejwakh et al., 2012).

The theoretical range for $\delta^{18}\text{O}-\text{NO}_3^-_{nit}$ (i.e., theoretical $\delta^{18}\text{O}-\text{NO}_3^-$ after nitrification) is -15 to 15‰, assuming soil-water and O_2 respectively contribute to NO_3^- -O accordingly to $\delta^{18}\text{O}-\text{NO}_3^- = 1/3 (\delta^{18}\text{O}-\text{O}_2) + 2/3 (\delta^{18}\text{O}-\text{H}_2\text{O})$ (Kendall et al., 2007). Here, a subset of 5 snow-water samples and 2 rain samples were analyzed for $\delta^{18}\text{O}$, with respective mean values of -16.6 and -8.2‰. Soil-water isotopes were not directly measured, but assimilated to precipitation water isotopes. Assuming $\delta^{18}\text{O}$ values of air- O_2 of 23.88‰ (Barkan and Luz, 2005), we ended up with $\delta^{18}\text{O}-\text{NO}_3^- = -3.3\%$ (if the end-member is snow water) and +2.3‰ (if the end-member is rain water) for $\text{NO}_3^-_{nit}$. Even though this approach comes with a number of assumptions and caveats (Rose et al., 2015b; Snider et al., 2010), the calculated values are significantly higher – by ca 4‰ – than the mean annual $\delta^{18}\text{O}-\text{NO}_3^-_{ter}$ (Table 3). In a study focusing on an estuarine system in San Francisco Bay, it was postulated that rapid recycling of NO_3^- under low $[\text{NO}_3^-]$ conditions (<0.9 mg L⁻¹) could result in an increased incorporation of H_2O -O atoms into the NO_3^- molecule (Wankel et al., 2006). Another work focusing on nitrification in forests soils of central Japan attributed low $\delta^{18}\text{O}-\text{NO}_3^-$ to a combination of O atoms exchange between H_2O and NO_2^- and kinetic isotope fractionation

associated with oxygen incorporation throughout the nitrification process in acidic soils (Fang et al., 2012). The latter hypothesis could explain the generally acidic subalpine soils characterized by slow N cycling (Robson et al., 2007), but further investigation is warranted to understand the cause of ^{18}O depleted NO_3^- .

When calculating f_{atm} using the approach described by Barnes et al. (2008):

$$(Eq.7) \quad f_{atm} = (\delta^{18}\text{O}_{sample} - \delta^{18}\text{O}_{nit}) / (\delta^{18}\text{O}_{atm} - \delta^{18}\text{O}_{nit})$$

and by using either $\delta^{18}\text{O}-\text{NO}_3^-_{ter}$ or $\delta^{18}\text{O}-\text{NO}_3^-_{nit}$ as nitrification end-member, significant differences between the estimated f_{atm} are obtained (Table 3). These findings suggest that atmospheric NO_3^- inputs to an ecosystem are underestimated (e.g., up to 6% in this study, Table 3) when calculated using Eq.7 with $\delta^{18}\text{O}-\text{NO}_3^-_{nit}$ as nitrification end-member.

5. Conclusions

The results showing higher $\text{NO}_3^-_{atm}$ exports in a montane stream relative to urban streams have several key implications. First, it sheds light on the contribution of snowmelt-derived groundwater to year-round $\text{NO}_3^-_{atm}$ exports in all streams at baseflow. Contamination of groundwater by $\text{NO}_3^-_{atm}$ may be a widespread phenomenon, regardless of emitting sources proximity: other studies also reported ubiquitous presence of $\text{NO}_3^-_{atm}$ in groundwater (Dejwakh et al., 2012; Dietzel et al., 2014; Nakagawa et al., 2013; Stoewer et al., 2015). Second, hydrological events (e.g., snowmelt) are the main controls of upland $\text{NO}_3^-_{atm}$ exports, whereas a chemically equilibrated groundwater reservoir dominates discharge in urban streams. In a context of climate change, with temporally shorter and shallower snow coverage

of altitude catchments, decreasing exports of $\text{NO}_3^-_{atm}$ by snowmelt water could result in proportional $\text{NO}_3^-_{atm}$ enrichment of alpine soils. However, how this would correlate with elsewhere reported snow cover removal effects on microbial communities, N retention, plants biomass and soil respiration is a complex question yet to be addressed (Brooks et al., 2011; Gavazov et al., 2017; Sorensen et al., 2016; Vankoughnett and Henry, 2013). **Third, the association** of $\Delta^{17}\text{O}$, $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ is a potent tool to unravel the respective atmospheric and terrestrial sources contribution to NO_3^- pools. **To correct** the $\delta^{18}\text{O}-\text{NO}_3^-$ from its atmospheric composition enables the determination of the $\delta^{18}\text{O}-\text{NO}_3^-_{ter}$, which came significantly lower at all sites than the theoretically calculated $\delta^{18}\text{O}-\text{NO}_3^-_{nit}$. This resulted in significantly higher estimated f_{atm} when using a more robust $\delta^{18}\text{O}-\text{NO}_3^-_{ter}$ (compared to $\delta^{18}\text{O}-\text{NO}_3^-_{nit}$) as nitrification end-member in Eq.7. Studies that did not use $\delta^{18}\text{O}-\text{NO}_3^-_{ter}$ in Eq.7 to determine f_{atm} may have underestimated atmospheric nitrate inputs to the environment (e.g., Barnes et al. (2008), Durka et al. (1994), Griffiths et al. (2016), Pellerin et al. (2012), Sebestyen et al. (2014)). **It is recommended** that more work be conducted using the triple isotope technique to revise **the assessment of $\text{NO}_3^-_{atm}$ ubiquity** in the environment.

Acknowledgments

This study was supported by grants from the Labex OSUG@2020 (“Investissements d’avenir” - ANR10 LABX56), the ARC – Environnement Région Rhone-Alpes, the Grenoble-Chambéry DIPEE CNRS. This work also benefited from the National Research Agency supports (“Investissements d’avenir” - ANR11 INBS-0001AnaEE - Services and “FloodScale project” - ANR 2011 BS56 027) and from the SAJF research station (UMR 3370, UGA-CNRS) infrastructures and competences. The

study took place on a Long Term Ecological Research site of the ZAA (Zone Atelier Alpes) and the Isère Campus monitoring site is a CNRS-labeled site of the ZABR (Rhône Basin Long Term Ecological Research site). We would like to thank G. Nord, J.-L. Jaffrezo, F. Masson, V. Lucaire, E. Vince and C. Arnoldi for help with either laboratory or field work. We also want to acknowledge J. Renaud for help with SIG.

Appendix A

The supplementary information provides additional Text on the analytical methods, Figures S1-S7 and Tables S1-S2.

References

- Aber, J.D., Nadelhoffer, K.J., Steudler, P., Melillo, J.M., 1989. Nitrogen Saturation in Northern Forest Ecosystems. *BioScience* 39, 378–386. <https://doi.org/10.2307/1311067>
- Balestrini, R., Arese, C., Freppaz, M., Buffagni, A., 2013. Catchment features controlling nitrogen dynamics in running waters above the tree line (central Italian Alps). *Hydrol. Earth Syst. Sci.* 17, 989–1001. <https://doi.org/10.5194/hess-17-989-2013>
- Barkan, E., Luz, B., 2005. High precision measurements of 17O/16O and 18O/16O ratios in H₂O. *Rapid Commun. Mass Spectrom.* 19, 3737–3742. <https://doi.org/10.1002/rcm.2250>
- Barnes, R.T., Raymond, P.A., 2010. Land-use controls on sources and processing of nitrate in small watersheds: insights from dual isotopic analysis. *Ecol. Appl.* 20, 1961–1978. <https://doi.org/10.1890/08-1328.1>
- Barnes, R.T., Raymond, P.A., Casciotti, K.L., 2008. Dual isotope analyses indicate efficient processing of atmospheric nitrate by forested watersheds in the northeastern US. *Biogeochemistry* 90, 15–27.
- Barnes, R.T., Williams, M.W., Parman, J.N., Hill, K., Caine, N., 2014. Thawing glacial and permafrost features contribute to nitrogen export from Green Lakes Valley, Colorado Front Range, USA. *Biogeochemistry* 117, 413–430. <https://doi.org/10.1007/s10533-013-9886-5>
- Baron, J.S., Campbell, D.H., 1997. Nitrogen fluxes in a high elevation Colorado Rocky Mountain basin. *Hydrol. Process.* 11, 783–799.
- Baron, J.S., Driscoll, C.T., Stoddard, J.L., Richer, E.E., 2011. Empirical Critical Loads of Atmospheric Nitrogen Deposition for Nutrient Enrichment and Acidification of Sensitive US Lakes. *BioScience* 61, 602–613. <https://doi.org/10.1525/bio.2011.61.8.6>
- Baron, J.S., Nydick, K.R., Rueth, H.M., Lafrancois, B.M., Wolfe, A.P., 2005. High Elevation Ecosystem Responses to Atmospheric Deposition of Nitrogen in the Colorado Rocky Mountains, USA, in: Huber, U.M., Bugmann, H.K.M., Reasoner, M.A. (Eds.), *Global Change and Mountain Regions*. Springer Netherlands, Dordrecht, pp. 429–436.

678 Baron, J.S., Rueth, H.M., Wolfe, A.M., Nydick, K.R., Allstott, E.J., Minear, J.T., Moraska, B.,
 679 2000. Ecosystem Responses to Nitrogen Deposition in the Colorado Front Range.
 680 *Ecosystems* 3, 352–368. <https://doi.org/10.1007/s100210000032>
 681 Beirle, S., Boersma, K.F., Platt, U., Lawrence, M.G., Wagner, T., 2011. Megacity Emissions
 682 and Lifetimes of Nitrogen Oxides Probed from Space. *Science* 333, 1737–1739.
 683 <https://doi.org/10.1126/science.1207824>
 684 Bettez, N.D., Groffman, P.M., 2013. Nitrogen Deposition in and near an Urban Ecosystem.
 685 *Environ. Sci. Technol.* 47, 6047–6051. <https://doi.org/10.1021/es400664b>
 686 Beyn, F., Matthias, V., Dähnke, K., 2014. Changes in atmospheric nitrate deposition in
 687 Germany – An isotopic perspective. *Environ. Pollut.* 194, 1–10.
 688 <https://doi.org/10.1016/j.envpol.2014.06.043>
 689 Bodin, X., Thibert, E., Fabre, D., Ribolini, A., Schoeneich, P., Francou, B., Reynaud, L., Fort,
 690 M., 2009. Two decades of responses (1986–2006) to climate by the Laurichard rock
 691 glacier, French Alps. *Permafr. Periglac. Process.* 20, 331–344.
 692 <https://doi.org/10.1002/ppp.665>
 693 Bourgeois, I., 2017. Atmospheric nitrate deposition on subalpine meadows of the
 694 Lautaret pass (phdthesis). Université Grenoble Alpes.
 695 Bourgeois, I., Savarino, J., Caillon, N., Angot, H., Barbero, A., Delbart, F., Voisin, D.,
 696 Clement, J.-C., n.d. Tracing the fate of atmospheric nitrate in a subalpine watershed using
 697 $\Delta^{17}\text{O}$. *Environ. Sci. Technol.* in review.
 698 Bowman, W.D., Gartner, J.R., Holland, K., Wiedermann, M., 2006. Nitrogen Critical Loads
 699 For Alpine Vegetation And Terrestrial Ecosystem Response: Are We There Yet? *Ecol.*
 700 *Appl.* 16, 1183–1193. [https://doi.org/10.1890/1051-](https://doi.org/10.1890/1051-0761(2006)016[1183:NCLFAV]2.0.CO;2)
 701 [0761\(2006\)016\[1183:NCLFAV\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[1183:NCLFAV]2.0.CO;2)
 702 Briand, C., 2014. Approche multi-traceurs pour la détermination de l'origine des nitrates
 703 dans les eaux souterraines: exemple d'une source karstique dans les Landes. Université
 704 Pierre et Marie Curie-Paris VI.
 705 Brooks, P.D., Campbell, D.H., Tonnessen, K.A., Heuer, K., 1999. Natural variability in N
 706 export from headwater catchments: snow cover controls on ecosystem N retention.
 707 *Hydrol. Process.* 13, 2191–2201.
 708 Brooks, P.D., Grogan, P., Templer, P.H., Groffman, P., Öquist, M.G., Schimel, J., 2011.
 709 Carbon and Nitrogen Cycling in Snow-Covered Environments. *Geogr. Compass* 5, 682–
 710 699. <https://doi.org/10.1111/j.1749-8198.2011.00420.x>
 711 Burns, D.A., Boyer, E.W., Elliott, E.M., Kendall, C., 2009. Sources and Transformations of
 712 Nitrate from Streams Draining Varying Land Uses: Evidence from Dual Isotope Analysis.
 713 *J. Environ. Qual.* 38, 1149. <https://doi.org/10.2134/jeq2008.0371>
 714 Burns, D.A., Kendall, C., 2002. Analysis of $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ to differentiate NO_3^- sources
 715 in runoff at two watersheds in the Catskill Mountains of New York. *Water Resour. Res.*
 716 38, 9–1. <https://doi.org/10.1029/2001WR000292>
 717 Campbell, D.H., Clow, D.W., Ingersoll, G.P., Mast, M.A., Spahr, N.E., Turk, J.T., 1995.
 718 Nitrogen deposition and release in alpine watersheds, Loch Vale, Colorado, USA. *IAHS*
 719 *Publ.-Ser. Proc. Rep.-Intern Assoc Hydrol. Sci.* 228, 243–254.
 720 Campbell, D.H., Kendall, C., Chang, C.C.Y., Silva, S.R., Tonnessen, K.A., 2002. Pathways for
 721 nitrate release from an alpine watershed: Determination using $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$: ALPINE
 722 WATERSHED NITRATE $\delta^{15}\text{N}$ AND $\delta^{18}\text{O}$. *Water Resour. Res.* 38, 10-1-10-9.
 723 <https://doi.org/10.1029/2001WR000294>
 724 Clark, C.M., Bell, M.D., Boyd, J.W., Compton, J.E., Davidson, E.A., Davis, C., Fenn, M.E.,
 725 Geiser, L., Jones, L., Blett, T.F., 2017. Nitrogen-induced terrestrial eutrophication:
 726 cascading effects and impacts on ecosystem services. *Ecosphere* 8, e01877.

<https://doi.org/10.1002/ecs2.1877>
 Clément, J.-C., Holmes, R.M., Peterson, B.J., Pinay, G., 2003. Isotopic investigation of denitrification in a riparian ecosystem in western France. *J. Appl. Ecol.* 40, 1035–1048. <https://doi.org/10.1111/j.1365-2664.2003.00854.x>
 Clow, D.W., Sueker, J.K., 2000. Relations between basin characteristics and stream water chemistry in alpine/subalpine basins in Rocky Mountain National Park, Colorado. *Water Resour. Res.* 36, 49–61. <https://doi.org/10.1029/1999WR900294>
 Costa, A.W., Michalski, G., Schauer, A.J., Alexander, B., Steig, E.J., Shepson, P.B., 2011. Analysis of atmospheric inputs of nitrate to a temperate forest ecosystem from $\Delta^{17}\text{O}$ isotope ratio measurements: ATMOSPHERIC NITRATE INPUTS TO A FOREST. *Geophys. Res. Lett.* 38, n/a–n/a. <https://doi.org/10.1029/2011GL047539>
 Cowie, R.M., Knowles, J.F., Dailey, K.R., Williams, M.W., Mills, T.J., Molotch, N.P., 2017. Sources of streamflow along a headwater catchment elevational gradient. *J. Hydrol.* 549, 163–178. <https://doi.org/10.1016/j.jhydrol.2017.03.044>
 Decina, S.M., Templer, P.H., Hutyra, L.R., Gately, C.K., Rao, P., 2017. Variability, drivers, and effects of atmospheric nitrogen inputs across an urban area: Emerging patterns among human activities, the atmosphere, and soils. *Sci. Total Environ.* 609, 1524–1534. <https://doi.org/10.1016/j.scitotenv.2017.07.166>
 Deiwakh, N.R., Meixner, T., Michalski, G., McIntosh, J., 2012. Using ^{17}O to Investigate Nitrate Sources and Sinks in a Semi-Arid Groundwater System. *Environ. Sci. Technol.* 46, 745–751. <https://doi.org/10.1021/es203450z>
 Devito, K.J., Fitzgerald, D., Hill, A.R., Aravena, R., 2000. Nitrate dynamics in relation to lithology and hydrologic flow path in a river riparian zone. *J. Environ. Qual.* 29, 1075–1084.
 Dietzel, M., Leis, A., Abdalla, R., Savarino, J., Morin, S., Böttcher, M.E., Köhler, S., 2014. ^{17}O excess traces atmospheric nitrate in paleo-groundwater of the Saharan desert. *Biogeosciences* 11, 3149–3161. <https://doi.org/10.5194/bg-11-3149-2014>
 Dodds, W.K., Bouska, W.W., Eitzmann, J.L., Pilger, T.J., Pitts, K.L., Riley, A.J., Schloesser, J.T., Thornbrugh, D.J., 2009. Eutrophication of U.S. Freshwaters: Analysis of Potential Economic Damages. *Environ. Sci. Technol.* 43, 12–19. <https://doi.org/10.1021/es801217q>
 Durka, W., Schulze, E.-D., Gebauer, G., Voerkeliust, S., 1994. Effects of forest decline on uptake and leaching of deposited nitrate determined from ^{15}N and ^{18}O measurements. *Nature* 372, 765–767. <https://doi.org/10.1038/372765a0>
 Dutordoir, S., 2014. Bilan des flux de métaux, carbone organique et nutriments contenus dans une rivière alpine: part des rejets urbains de l'agglomération de Grenoble et apports amont (Isère et Drac). Université Grenoble Alpes.
 Elliott, E.M., Kendall, C., Boyer, E.W., Burns, D.A., Lear, G.G., Golden, H.E., Harlin, K., Bytnerowicz, A., Butler, T.J., Glatz, R., 2009. Dual nitrate isotopes in dry deposition: Utility for partitioning NO_x source contributions to landscape nitrogen deposition. *J. Geophys. Res.* 114. <https://doi.org/10.1029/2008JG000889>
 Elser, J.J., Andersen, T., Baron, J.S., Bergström, A.-K., Jansson, M., Kyle, M., Nydick, K.R., Steger, L., Hessen, D.O., 2009. Shifts in lake N:P stoichiometry and nutrient limitation driven by atmospheric nitrogen deposition. *Science* 326, 835–837. <https://doi.org/10.1126/science.1176199>
 Emmerton, K.S., Callaghan, T.V., Jones, H.E., Leake, J.R., Michelsen, A., Read, D.J., 2001. Assimilation and Isotopic Fractionation of Nitrogen by Mycorrhizal and Nonmycorrhizal Subarctic Plants. *New Phytol.* 151, 513–524. <https://doi.org/10.2307/1353802>

776 Erbland, J., 2011. Contraintes isotopiques sur l'interprétation de l'enregistrement en
 777 nitrate dans la carotte de glace de Vostok. Université de Grenoble.
 778 Estrada, N.L., Böhlke, J.K., Sturchio, N.C., Gu, B., Harvey, G., Burkey, K.O., Grantz, D.A.,
 779 McGrath, M.T., Anderson, T.A., Rao, B., Sevanthi, R., Hatzinger, P.B., Jackson, W.A., 2017.
 780 Stable isotopic composition of perchlorate and nitrate accumulated in plants:
 781 Hydroponic experiments and field data. *Sci. Total Environ.* 595, 556–566.
 782 <https://doi.org/10.1016/j.scitotenv.2017.03.223>
 783 Fang, Y., Koba, K., Makabe, A., Takahashi, C., Zhu, W., Hayashi, T., Hokari, A.A., Urakawa,
 784 R., Bai, E., Houlton, B.Z., Xi, D., Zhang, S., Matsushita, K., Tu, Y., Liu, D., Zhu, F., Wang, Z.,
 785 Zhou, G., Chen, D., Makita, T., Toda, H., Liu, X., Chen, Q., Zhang, D., Li, Y., Yoh, M., 2015.
 786 Microbial denitrification dominates nitrate losses from forest ecosystems. *Proc. Natl.*
 787 *Acad. Sci.* 112, 1470–1474. <https://doi.org/10.1073/pnas.1416776112>
 788 Fang, Y., Koba, K., Makabe, A., Zhu, F., Fan, S., Liu, X., Yoh, M., 2012. Low $\delta^{18}\text{O}$ Values of
 789 Nitrate Produced from Nitrification in Temperate Forest Soils. *Environ. Sci. Technol.* 46,
 790 8723–8730. <https://doi.org/10.1021/es300510r>
 791 Fang, Y., Yoh, M., Koba, K., Zhu, W., Takebayashi, Y., Xiao, Y., Lei, C., Mo, J., Zhang, W., Lu,
 792 X., 2011. Nitrogen deposition and forest nitrogen cycling along an urban-rural transect
 793 in southern China: FOREST N CYCLING IN SOUTHERN CHINA. *Glob. Change Biol.* 17,
 794 872–885. <https://doi.org/10.1111/j.1365-2486.2010.02283.x>
 795 Fowler, D., Steadman, C.E., Stevenson, D., Coyle, M., Rees, R.M., Skiba, U.M., Sutton, M.A.,
 796 Cape, J.N., Dore, A.J., Viero, M., Simpson, D., Zaehle, S., Stocker, B.D., Rinaldi, M., Facchini,
 797 M.C., Flechard, C.R., Nemitz, E., Twigg, M., Erisman, J.W., Butterbach-Bahl, K., Galloway,
 798 J.N., 2015. Effects of global change during the 21st century on the nitrogen cycle.
 799 *Atmospheric Chem. Phys.* 15, 13849–13893. [https://doi.org/10.5194/acp-15-13849-](https://doi.org/10.5194/acp-15-13849-2015)
 800 2015
 801 Frey, M.M., Savarino, J., Morin, S., Erbland, J., Martins, J.M.F., 2009. Photolysis imprint in
 802 the nitrate stable isotope signal in snow and atmosphere of East Antarctica and
 803 implications for reactive nitrogen cycling. *Atmospheric Chem. Phys.* 9, 8681–8696.
 804 Freyer, H.D., 1991. Seasonal variation of $^{15}\text{N}/^{14}\text{N}$ ratios in atmospheric nitrate species.
 805 *Tellus B* 43, 30–44. <https://doi.org/10.1034/j.1600-0889.1991.00003.x>
 806 Galloway, J.N., Dentener, F.J., Capone, D.G., Boyer, E.W., Howarth, R.W., Seitzinger, S.P.,
 807 Asner, G.P., Cleveland, C.C., Green, P.A., Holland, E.A., Karl, D.M., Michaels, A.F., Porter,
 808 J.H., Townsend, A.R., Voesmarty, C.J., 2004. Nitrogen Cycles: Past, Present, and Future.
 809 *Biogeochemistry* 70, 153–226. <https://doi.org/10.1007/s10533-004-0370-0>
 810 Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R., Martinelli,
 811 L.A., Seitzinger, S.P., Sutton, M.A., 2008. Transformation of the nitrogen cycle: recent
 812 trends, questions, and potential solutions. *Science* 320, 889–892.
 813 Gavazov, K., Ingrisch, J., Hasibeder, R., Mills, R.T.E., Buttler, A., Gleixner, G., Pumpanen, J.,
 814 Bahn, M., 2017. Winter ecology of a subalpine grassland: Effects of snow removal on soil
 815 respiration, microbial structure and function. *Sci. Total Environ.* 590–591, 316–324.
 816 <https://doi.org/10.1016/j.scitotenv.2017.03.010>
 817 Granger, J., Sigman, D.M., Lehmann, M.F., Tortell, P.D., 2008. Nitrogen and oxygen isotope
 818 fractionation during dissimilatory nitrate reduction by denitrifying bacteria. *Limnol.*
 819 *Oceanogr.* 53, 2533.
 820 Granger, J., Sigman, D.M., Needoba, J.A., Harrison, P.J., 2004. Coupled nitrogen and oxygen
 821 isotope fractionation of nitrate during assimilation by cultures of marine phytoplankton.
 822 *Limnol. Oceanogr.* 49, 1763–1773.
 823 Granger, J., Sigman, D.M., Rohde, M.M., Maldonado, M.T., Tortell, P.D., 2010. N and O
 824 isotope effects during nitrate assimilation by unicellular prokaryotic and eukaryotic

825 plankton cultures. *Geochim. Cosmochim. Acta* 74, 1030–1040.
826 <https://doi.org/10.1016/j.gca.2009.10.044>
827 Granger, J., Wankel, S.D., 2016. Isotopic overprinting of nitrification on denitrification as
828 a ubiquitous and unifying feature of environmental nitrogen cycling. *Proc. Natl. Acad.*
829 *Sci.* 113, E6391–E6400. <https://doi.org/10.1073/pnas.1601383113>
830 Griffiths, N.A., Jackson, C.R., McDonnell, J.J., Klaus, J., Du, E., Bitew, M.M., 2016. Dual
831 nitrate isotopes clarify the role of biological processing and hydrologic flow paths on
832 nitrogen cycling in subtropical low-gradient watersheds: N CYCLING IN LOW-RELIEF
833 WATERSHEDS. *J. Geophys. Res. Biogeosciences* 121, 422–437.
834 <https://doi.org/10.1002/2015JG003189>
835 Groffman, P.M., Law, N.L., Belt, K.T., Band, L.E., Fisher, G.T., 2004. Nitrogen Fluxes and
836 Retention in Urban Watershed Ecosystems. *Ecosystems* 7.
837 <https://doi.org/10.1007/s10021-003-0039-x>
838 Hall, S.J., Maurer, G., Hoch, S.W., Taylor, R., Bowling, D.R., 2014. Impacts of anthropogenic
839 emissions and cold air pools on urban to montane gradients of snowpack ion
840 concentrations in the Wasatch Mountains, Utah. *Atmos. Environ.* 98, 231–241.
841 <https://doi.org/10.1016/j.atmosenv.2014.08.076>
842 Hall, S.J., Weintraub, S.R., Eiriksson, D., Brooks, P.D., Baker, M.A., Bowen, G.J., Bowling,
843 D.R., 2016. Stream Nitrogen Inputs Reflect Groundwater Across a Snowmelt-Dominated
844 Montane to Urban Watershed. *Environ. Sci. Technol.* 50, 1137–1146.
845 <https://doi.org/10.1021/acs.est.5b04805>
846 Hastings, M.G., Jarvis, J.C., Steig, E.J., 2009. Anthropogenic Impacts on Nitrogen Isotopes
847 of Ice-Core Nitrate. *Science* 324, 1288–1288. <https://doi.org/10.1126/science.1170510>
848 Hertel, O., Skjøth, C.A., Reis, S., Bleeker, A., Harrison, R.M., Cape, J.N., Fowler, D., Skiba, U.,
849 Simpson, D., Jickells, T., Kulmala, M., Gyldenkerne, S., Sørensen, L.L., Erisman, J.W.,
850 Sutton, M.A., 2012. Governing processes for reactive nitrogen compounds in the
851 European atmosphere. *Biogeosciences* 9, 4921–4954. [https://doi.org/10.5194/bg-9-](https://doi.org/10.5194/bg-9-4921-2012)
852 [4921-2012](https://doi.org/10.5194/bg-9-4921-2012)
853 Holtgrieve, G.W., Schindler, D.E., Hobbs, W.O., Leavitt, P.R., Ward, E.J., Bunting, L., Chen,
854 G., Finney, B.P., Gregory-Eaves, I., Holmgren, S., Lisac, M.J., Lisi, P.J., Nydick, K., Rogers,
855 L.A., Saros, J.E., Selbie, D.T., Shapley, M.D., Walsh, P.B., Wolfe, A.P., 2011. A Coherent
856 Signature of Anthropogenic Nitrogen Deposition to Remote Watersheds of the Northern
857 Hemisphere. *Science* 334, 1545–1548. <https://doi.org/10.1126/science.1212267>
858 Hood, E., Scott, D., 2008. Riverine organic matter and nutrients in southeast Alaska
859 affected by glacial coverage. *Nat. Geosci.* 1, 583–587. <https://doi.org/10.1038/ngeo280>
860 Hundey, E.J., Russell, S.D., Longstaffe, F.J., Moser, K.A., 2016. Agriculture causes nitrate
861 fertilization of remote alpine lakes. *Nat. Commun.* 7, 10571.
862 <https://doi.org/10.1038/ncomms10571>
863 Kaiser, J., Hastings, M.G., Houlton, B.Z., Röckmann, T., Sigman, D.M., 2007. Triple Oxygen
864 Isotope Analysis of Nitrate Using the Denitrifier Method and Thermal Decomposition of
865 N₂O. *Anal. Chem.* 79, 599–607. <https://doi.org/10.1021/ac061022s>
866 Kaye, J.P., Hart, S.C., 1997. Competition for nitrogen between plants and soil
867 microorganisms. *Trends Ecol. Evol.* 12, 139–143. [https://doi.org/10.1016/S0169-](https://doi.org/10.1016/S0169-5347(97)01001-X)
868 [5347\(97\)01001-X](https://doi.org/10.1016/S0169-5347(97)01001-X)
869 Kean, A.J., Harley, R.A., Littlejohn, D., Kendall, G.R., 2000. On-Road Measurement of
870 Ammonia and Other Motor Vehicle Exhaust Emissions. *Environ. Sci. Technol.* 34, 3535–
871 3539. <https://doi.org/10.1021/es991451q>
872 Kendall, C., Campbell, D.H., Burns, D.A., Shanley, J.B., Silva, S.R., Chang, C.C., 1995. Tracing
873 sources of nitrate in snowmelt runoff using the oxygen and nitrogen isotopic

874 compositions of nitrate. IAHS Publ.-Ser. Proc. Rep.-Intern Assoc Hydrol. Sci. 228, 339–
875 348.

876 Kendall, C., Elliott, E.M., Wankel, S.D., 2007. Tracing anthropogenic inputs of nitrogen to
877 ecosystems. *Stable Isot. Ecol. Environ. Sci.* 2, 375–449.

878 Kirchner, M., Fegg, W., Römmelt, H., Leuchner, M., Ries, L., Zimmermann, R., Michalke, B.,
879 Wallasch, M., Maguhn, J., Faus-Kessler, T., Jakobi, G., 2014. Nitrogen deposition along
880 differently exposed slopes in the Bavarian Alps. *Sci. Total Environ.* 470–471, 895–906.
881 <https://doi.org/10.1016/j.scitotenv.2013.10.036>

882 Lefebvre, S., Clément, J.-C., Pinay, G., Thenail, C., Durand, P., Marmonier, P., 2007. 15n-
883 Nitrate Signature in Low-Order Streams: Effects of Land Cover and Agricultural
884 Practices. *Ecol. Appl.* 17, 2333–2346. <https://doi.org/10.1890/06-1496.1>

885 Liu, T., Wang, F., Michalski, G., Xia, X., Liu, S., 2013. Using ¹⁵ N, ¹⁷ O, and ¹⁸ O To Determine
886 Nitrate Sources in the Yellow River, China. *Environ. Sci. Technol.* 47, 13412–13421.
887 <https://doi.org/10.1021/es403357m>

888 Liu, X.-Y., Koba, K., Takebayashi, Y., Liu, C.-Q., Fang, Y.-T., Yoh, M., 2013. Dual N and O
889 isotopes of nitrate in natural plants: first insights into individual variability and organ-
890 specific patterns. *Biogeochemistry* 114, 399–411. [https://doi.org/10.1007/s10533-012-](https://doi.org/10.1007/s10533-012-9721-4)
891 [9721-4](https://doi.org/10.1007/s10533-012-9721-4)

892 Louiseize, N.L., Lafrenière, M.J., Hastings, M.G., 2014. Stable isotopic evidence of
893 enhanced export of microbially derived NO_3^- following active
894 layer slope disturbance in the Canadian High Arctic. *Biogeochemistry* 121, 565–580.
895 <https://doi.org/10.1007/s10533-014-0023-x>

896 Lovett, G.M., Goodale, C.L., 2011. A New Conceptual Model of Nitrogen Saturation Based
897 on Experimental Nitrogen Addition to an Oak Forest. *Ecosystems* 14, 615–631.
898 <https://doi.org/10.1007/s10021-011-9432-z>

899 Mano, V., 2008. Processus fondamentaux conditionnant les apports de sédiments fins
900 dans les retenues-Optimisation des méthodes de mesure et modélisation statistique.
901 Université Joseph-Fourier-Grenoble I.

902 Mano, V., Nemery, J., Belleudy, P., Poirel, A., 2009. Assessment of suspended sediment
903 transport in four alpine watersheds (France): influence of the climatic regime. *Hydrol.*
904 *Process.* 23, 777–792. <https://doi.org/10.1002/hyp.7178>

905 Mara, P., Mihalopoulos, N., Gogou, A., Daehnke, K., Schlarbaum, T., Emeis, K.-C., Krom, M.,
906 2009. Isotopic composition of nitrate in wet and dry atmospheric deposition on Crete in
907 the eastern Mediterranean Sea: ISOTOPIC COMPOSITION OF NITRATE IN DEPOSITION.
908 *Glob. Biogeochem. Cycles* 23, n/a-n/a. <https://doi.org/10.1029/2008GB003395>

909 Mast, M.A., Clow, D.W., Baron, J.S., Wetherbee, G.A., 2014. Links between N Deposition
910 and Nitrate Export from a High-Elevation Watershed in the Colorado Front Range.
911 *Environ. Sci. Technol.* 48, 14258–14265. <https://doi.org/10.1021/es502461k>

912 Matson, P., Lohse, K.A., Hall, S.J., 2002. The Globalization of Nitrogen Deposition:
913 Consequences for Terrestrial Ecosystems. *AMBIO J. Hum. Environ.* 31, 113.
914 [https://doi.org/10.1639/0044-7447\(2002\)031\[0113:TGONDC\]2.0.CO;2](https://doi.org/10.1639/0044-7447(2002)031[0113:TGONDC]2.0.CO;2)

915 Mayer, B., Bollwerk, S.M., Mansfeldt, T., Hütter, B., Veizer, J., 2001. The oxygen isotope
916 composition of nitrate generated by nitrification in acid forest floors. *Geochim.*
917 *Cosmochim. Acta* 65, 2743–2756. [https://doi.org/10.1016/S0016-7037\(01\)00612-3](https://doi.org/10.1016/S0016-7037(01)00612-3)

918 Mayer, B., Boyer, E.W., Goodale, C., Jaworski, N.A., Van Breemen, N., Howarth, R.W.,
919 Seitzinger, S., Billen, G., Lajtha, K., Nadelhoffer, K., others, 2002. Sources of nitrate in
920 rivers draining sixteen watersheds in the northeastern US: Isotopic constraints.
921 *Biogeochemistry* 57, 171–197.

922 Michalski, G., Kolanowski, M., Riha, K.M., 2015. Oxygen and nitrogen isotopic

composition of nitrate in commercial fertilizers, nitric acid, and reagent salts. *Isotopes Environ. Health Stud.* 51, 382–391. <https://doi.org/10.1080/10256016.2015.1054821>

Michalski, G., Meixner, T., Fenn, M., Hernandez, L., Sirulnik, A., Allen, E., Thiemens, M., 2004. Tracing Atmospheric Nitrate Deposition in a Complex Semiarid Ecosystem Using $\Delta^{17}\text{O}$. *Environ. Sci. Technol.* 38, 2175–2181. <https://doi.org/10.1021/es034980+>

Morin, S., Savarino, J., Frey, M.M., Domine, F., Jacobi, H.-W., Kaleschke, L., Martins, J.M.F., 2009. Comprehensive isotopic composition of atmospheric nitrate in the Atlantic Ocean boundary layer from 65°S to 79°N. *J. Geophys. Res.* 114. <https://doi.org/10.1029/2008JD010696>

Morin, S., Savarino, J., Frey, M.M., Yan, N., Bekki, S., Bottenheim, J.W., Martins, J.M.F., 2008. Tracing the Origin and Fate of NO_x in the Arctic Atmosphere Using Stable Isotopes in Nitrate. *Science* 322, 730–732. <https://doi.org/10.1126/science.1161910>

Nakagawa, F., Suzuki, A., Daita, S., Ohyama, T., Komatsu, D.D., Tsunogai, U., 2013. Tracing atmospheric nitrate in groundwater using triple oxygen isotopes: evaluation based on bottled drinking water. *Biogeosciences* 10, 3547–3558.

Nanus, L., McMurray, J.A., Clow, D.W., Saros, J.E., Blett, T., Gurdak, J.J., 2017. Spatial variation of atmospheric nitrogen deposition and critical loads for aquatic ecosystems in the Greater Yellowstone Area. *Environ. Pollut.* 223, 644–656. <https://doi.org/10.1016/j.envpol.2017.01.077>

Nanus, L., Williams, M.W., Campbell, D.H., Elliott, E.M., Kendall, C., 2008. Evaluating Regional Patterns in Nitrate Sources to Watersheds in National Parks of the Rocky Mountains using Nitrate Isotopes. *Environ. Sci. Technol.* 42, 6487–6493. <https://doi.org/10.1021/es800739e>

Némery, J., Mano, V., Coynel, A., Etcheber, H., Moatar, F., Meybeck, M., Belleudy, P., Poirol, A., 2013. Carbon and suspended sediment transport in an impounded alpine river (Isère, France). *Hydrol. Process.* 27, n/a–n/a. <https://doi.org/10.1002/hyp.9387>

Ohte, N., Sebestyen, S.D., Shanley, J.B., Doctor, D.H., Kendall, C., Wankel, S.D., Boyer, E.W., 2004. Tracing sources of nitrate in snowmelt runoff using a high-resolution isotopic technique: TRACING SOURCES OF NITRATE IN SNOWMELT RUNOFF. *Geophys. Res. Lett.* 31, n/a–n/a. <https://doi.org/10.1029/2004GL020908>

Pellerin, B.A., Saraceno, J.F., Shanley, J.B., Sebestyen, S.D., Aiken, G.R., Wollheim, W.M., Bergamaschi, B.A., 2012. Taking the pulse of snowmelt: in situ sensors reveal seasonal, event and diurnal patterns of nitrate and dissolved organic matter variability in an upland forest stream. *Biogeochemistry* 108, 183–198. <https://doi.org/10.1007/s10533-011-9589-8>

Preunkert, S., 2003. A seasonally resolved alpine ice core record of nitrate: Comparison with anthropogenic inventories and estimation of preindustrial emissions of NO in Europe. *J. Geophys. Res.* 108. <https://doi.org/10.1029/2003JD003475>

Rao, P., Hutyra, L.R., Raciti, S.M., Templer, P.H., 2014. Atmospheric nitrogen inputs and losses along an urbanization gradient from Boston to Harvard Forest, MA. *Biogeochemistry* 121, 229–245. <https://doi.org/10.1007/s10533-013-9861-1>

Riha, K.M., Michalski, G., Gallo, E.L., Lohse, K.A., Brooks, P.D., Meixner, T., 2014. High Atmospheric Nitrate Inputs and Nitrogen Turnover in Semi-arid Urban Catchments. *Ecosystems* 17, 1309–1325. <https://doi.org/10.1007/s10021-014-9797-x>

Robson, T., Lavorel, S., Clement, J., Roux, X., 2007. Neglect of mowing and manuring leads to slower nitrogen cycling in subalpine grasslands. *Soil Biol. Biochem.* 39, 930–941. <https://doi.org/10.1016/j.soilbio.2006.11.004>

Rogora, M., Mosello, R., Arisci, S., Brizzio, M.C., Barbieri, A., Balestrini, R., Waldner, P., Schmitt, M., Stähli, M., Thimonier, A., Kalina, M., Puxbaum, H., Nickus, U., Ulrich, E.,

972 Probst, A., 2006. An Overview of Atmospheric Deposition Chemistry over the Alps:
 973 Present Status and Long-term Trends. *Hydrobiologia* 562, 17–40.
 974 <https://doi.org/10.1007/s10750-005-1803-z>
 975 Rose, L.A., Elliott, E.M., Adams, M.B., 2015a. Triple Nitrate Isotopes Indicate Differing
 976 Nitrate Source Contributions to Streams Across a Nitrogen Saturation Gradient.
 977 *Ecosystems* 18, 1209–1223. <https://doi.org/10.1007/s10021-015-9891-8>
 978 Rose, L.A., Sebestyen, S.D., Elliott, E.M., Koba, K., 2015b. Drivers of atmospheric nitrate
 979 processing and export in forested catchments. *Water Resour. Res.* 51, 1333–1352.
 980 <https://doi.org/10.1002/2014WR015716>
 981 Sabo, R.D., Nelson, D.M., Eshleman, K.N., 2016. Episodic, seasonal, and annual export of
 982 atmospheric and microbial nitrate from a temperate forest: MICROBIAL AND
 983 ATMOSPHERIC NITRATE EXPORT. *Geophys. Res. Lett.* 43, 683–691.
 984 <https://doi.org/10.1002/2015GL066758>
 985 Savarino, J., Kaiser, J., Morin, S., Sigman, D.M., Thiemens, M.H., 2007. Nitrogen and oxygen
 986 isotopic constraints on the origin of atmospheric nitrate in coastal Antarctica.
 987 *Atmospheric Chem. Phys.* 7, 1925–1945. <https://doi.org/10.5194/acp-7-1925-2007>
 988 Sebestyen, S.D., Shanley, J.B., Boyer, E.W., Kendall, C., Doctor, D.H., 2014. Coupled
 989 hydrological and biogeochemical processes controlling variability of nitrogen species in
 990 streamflow during autumn in an upland forest: STREAM N DYNAMICS DURING
 991 AUTUMN. *Water Resour. Res.* 50, 1569–1591. <https://doi.org/10.1002/2013WR013670>
 992 Shi, G., Buffen, A.M., Hastings, M.G., Li, C., Ma, H., Li, Y., Sun, B., An, C., Jiang, S., 2015.
 993 Investigation of post-depositional processing of nitrate in East Antarctic snow: isotopic
 994 constraints on photolytic loss, re-oxidation, and source inputs. *Atmospheric Chem. Phys.*
 995 15, 9435–9453. <https://doi.org/10.5194/acp-15-9435-2015>
 996 Snider, D.M., Spoelstra, J., Schiff, S.L., Venkiteswaran, J.J., 2010. Stable Oxygen Isotope
 997 Ratios of Nitrate Produced from Nitrification: 18O-Labeled Water Incubations of
 998 Agricultural and Temperate Forest Soils. *Environ. Sci. Technol.* 44, 5358–5364.
 999 <https://doi.org/10.1021/es1002567>
 1000 Sorensen, P.O., Templer, P.H., Christenson, L., Duran, J., Fahey, T., Fisk, M.C., Groffman,
 1001 P.M., Morse, J.L., Finzi, A.C., 2016. Reduced snow cover alters root-microbe interactions
 1002 and decreases nitrification rates in a northern hardwood forest. *Ecology* 97, 3359–3368.
 1003 <https://doi.org/10.1002/ecy.1599>
 1004 Spoelstra, J., Schiff, S.L., Hazlett, P.W., Jeffries, D.S., Semkin, R.G., 2007. The isotopic
 1005 composition of nitrate produced from nitrification in a hardwood forest floor. *Geochim.*
 1006 *Cosmochim. Acta* 71, 3757–3771. <https://doi.org/10.1016/j.gca.2007.05.021>
 1007 Stoewer, M.M., Knöller, K., Stumpp, C., 2015. Tracing freshwater nitrate sources in pre-
 1008 alpine groundwater catchments using environmental tracers. *J. Hydrol.* 524, 753–767.
 1009 <https://doi.org/10.1016/j.jhydrol.2015.03.022>
 1010 Templer, P.H., Weathers, K.C., 2011. Use of mixed ion exchange resin and the denitrifier
 1011 method to determine isotopic values of nitrate in atmospheric deposition and canopy
 1012 throughfall. *Atmos. Environ.* 45, 2017–2020.
 1013 <https://doi.org/10.1016/j.atmosenv.2011.01.035>
 1014 Templer, P.H., Weathers, K.C., Lindsey, A., Lenoir, K., Scott, L., 2015. Atmospheric inputs
 1015 and nitrogen saturation status in and adjacent to Class I wilderness areas of the
 1016 northeastern US. *Oecologia* 177, 5–15. <https://doi.org/10.1007/s00442-014-3121-5>
 1017 Thiemens, M.H., 2006. History and applications of mass-independent isotope effects.
 1018 *Annu Rev Earth Planet Sci* 34, 217–262.
 1019 Tomaz, S., Jaffrezo, J.-L., Favez, O., Perraudin, E., Villenave, E., Albinet, A., 2017. Sources
 1020 and atmospheric chemistry of oxy- and nitro-PAHs in the ambient air of Grenoble

1021 (France). *Atmos. Environ.* 161, 144–154.
 1022 <https://doi.org/10.1016/j.atmosenv.2017.04.042>
 1023 Treibergs, L.A., Granger, J., 2017. Enzyme level N and O isotope effects of assimilatory
 1024 and dissimilatory nitrate reduction: Enzyme level N and O isotope effects. *Limnol.*
 1025 *Oceanogr.* 62, 272–288. <https://doi.org/10.1002/lno.10393>
 1026 Tsunogai, U., Komatsu, D.D., Daita, S., Kazemi, G.A., Nakagawa, F., Noguchi, I., Zhang, J.,
 1027 2010. Tracing the fate of atmospheric nitrate deposited onto a forest ecosystem in
 1028 Eastern Asia using $\Delta 17\text{O}$. *Atmospheric Chem. Phys.* 10, 1809–1820.
 1029 Tsunogai, U., Komatsu, D.D., Ohyama, T., Suzuki, A., Nakagawa, F., Noguchi, I., Takagi, K.,
 1030 Nomura, M., Fukuzawa, K., Shibata, H., 2014. Quantifying the effects of clear-cutting and
 1031 strip-cutting on nitrate dynamics in a forested watershed using triple oxygen isotopes as
 1032 tracers. *Biogeosciences* 11, 5411–5424. <https://doi.org/10.5194/bg-11-5411-2014>
 1033 Tsunogai, U., Miyauchi, T., Ohyama, T., Komatsu, D.D., Nakagawa, F., Obata, Y., Sato, K.,
 1034 Ohizumi, T., 2016. Accurate and precise quantification of atmospheric nitrate in streams
 1035 draining land of various uses by using triple oxygen isotopes as tracers. *Biogeosciences*
 1036 13, 3441–3459. <https://doi.org/10.5194/bg-13-3441-2016>
 1037 Vankoughnett, M.R., Henry, H.A.L., 2013. Combined Effects of Soil Freezing and N
 1038 Addition on Losses and Interception of N Over Winter and Summer. *Ecosystems* 16,
 1039 694–703. <https://doi.org/10.1007/s10021-013-9642-7>
 1040 Vitousek, P.M., Aber, J.D., Howarth, R.W., Likens, G.E., Matson, P.A., Schindler, D.W.,
 1041 Schlesinger, W.H., Tilman, D.G., 1997. HUMAN ALTERATION OF THE GLOBAL NITROGEN
 1042 CYCLE: SOURCES AND CONSEQUENCES. *Ecol. Appl.* 7, 737–750.
 1043 [https://doi.org/10.1890/1051-0761\(1997\)007\[0737:HAOTGN\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1997)007[0737:HAOTGN]2.0.CO;2)
 1044 Wankel, S.D., Kendall, C., Francis, C.A., Paytan, A., 2006. Nitrogen sources and cycling in
 1045 the San Francisco Bay Estuary: A nitrate dual isotopic composition approach. *Limnol.*
 1046 *Oceanogr.* 51, 1654–1664.
 1047 Wasiuta, V., Lafrenière, M.J., Norman, A.-L., Hastings, M.G., 2015. Summer deposition of
 1048 sulfate and reactive nitrogen to two alpine valleys in the Canadian Rocky Mountains.
 1049 *Atmos. Environ.* 101, 270–285. <https://doi.org/10.1016/j.atmosenv.2014.10.041>
 1050 Weijs, S.V., Mutzner, R., Parlange, M.B., 2013. Could electrical conductivity replace water
 1051 level in rating curves for alpine streams?: ELECTRICAL CONDUCTIVITY STREAMFLOW
 1052 RATING CURVES. *Water Resour. Res.* 49, 343–351.
 1053 <https://doi.org/10.1029/2012WR012181>
 1054 Wexler, S.K., Goodale, C.L., McGuire, K.J., Bailey, S.W., Groffman, P.M., 2014. Isotopic
 1055 signals of summer denitrification in a northern hardwood forested catchment. *Proc.*
 1056 *Natl. Acad. Sci.* 111, 16413–16418. <https://doi.org/10.1073/pnas.1404321111>
 1057 Williams, J.J., Nurse, A., Saros, J.E., Riedel, J., Beutel, M., 2016. Effects of glaciers on
 1058 nutrient concentrations and phytoplankton in lakes within the Northern Cascades
 1059 Mountains (USA). *Biogeochemistry* 131, 373–385. [https://doi.org/10.1007/s10533-](https://doi.org/10.1007/s10533-016-0264-y)
 1060 [016-0264-y](https://doi.org/10.1007/s10533-016-0264-y)
 1061 Williams, M.W., Knauf, M., Cory, R., Caine, N., Liu, F., 2007. Nitrate content and potential
 1062 microbial signature of rock glacier outflow, Colorado Front Range. *Earth Surf. Process.*
 1063 *Landf.* 32, 1032–1047. <https://doi.org/10.1002/esp.1455>
 1064 Williams, M.W., Melack, J.M., 1991. Solute chemistry of snowmelt and runoff in an Alpine
 1065 Basin, Sierra Nevada. *Water Resour. Res.* 27, 1575–1588.
 1066 <https://doi.org/10.1029/90WR02774>
 1067 Williams, M.W., Seibold, C., Chowanski, K., 2009. Storage and release of solutes from a
 1068 subalpine seasonal snowpack: soil and stream water response, Niwot Ridge, Colorado.
 1069 *Biogeochemistry* 95, 77–94. <https://doi.org/10.1007/s10533-009-9288-x>

1070 Yang, Y.-Y., Toor, G.S., 2016. $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ Reveal the Sources of Nitrate-Nitrogen in
1071 Urban Residential Stormwater Runoff. Environ. Sci. Technol.
1072 <https://doi.org/10.1021/acs.est.5b05353>
1073 Ye, C., Gao, H., Zhang, N., Zhou, X., 2016. Photolysis of Nitric Acid and Nitrate on Natural
1074 and Artificial Surfaces. Environ. Sci. Technol. 50, 3530–3536.
1075 <https://doi.org/10.1021/acs.est.5b05032>

Watershed	S-upper montane	N-upper montane	Mid montane	Lower montane	Upper urban	Lower Urban
Main regime	Snowmelt	Glacier melt	Snowmelt	Snowmelt	Snowmelt - Rainfall	Snowmelt - Rainfall
Altitude range (m a.s.l.)	1667-2725	1667-3155	1618-1980	1050-4088	210-2650	210-4088
Size (km ²)	3.4	5.3	0.5	220	5720	3600
Mean slope (%)	25	30	17	28	22	22
Geology	Calcareous	Granitic and ancient volcanic rocks	Calcareous	60% Marls and carbonates, 40% Metamorphic and crystalline rocks	77% Marls and carbonates, 15% Metamorphic and crystalline rocks, 8% glacial deposits	Crystalline rocks (granite, gneiss and amphibolite)
Land cover	30% alpine meadows (grazed), 27% uncovered, 26% abandoned grasslands, 13% terraced meadows (mown for hay), some habitations	Mostly uncovered and abandoned grasslands	80% terraced meadows (mown for hay), 13% alpine meadows, one village	60% Uncovered, 40% Forest or low vegetation, a few villages	74% Forest and semi-natural vegetation, 22% Agricultural fields, 4% Urbanized areas	81% Forest and semi-natural vegetation, 16% Agricultural fields, 3% urbanized areas

1076
 1077
 1078
 1079
 1080
 1081
 1082
 1083
 1084
 1085

1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114

Table 1 Characteristics of the Romanche Valley and Isere watersheds along the montane to urban gradient

Figure 1 Two years solute concentrations in streams, with a) [NO₃⁻] (mg L⁻¹), b) [SO₄²⁻] (mg L⁻¹), c) [Cl⁻] (mg L⁻¹) and d) [Ca²⁺] (mg L⁻¹). Different letters denote significant differences in NO₃⁻ isotopic composition across sites. (e.g., a site with *ab* letters is significantly different from sites with letters from *c* onwards, but not significantly different from sites with *a* or *b* letters).

Figure 2 Two years isotopic composition of NO_3^- in streams, with a) $\Delta^{17}\text{O}$ (‰) (and corresponding f_{atm} (%) on the red y-axis), b) $\delta^{18}\text{O}$ (‰) and c) $\delta^{15}\text{N}$ (‰). Different letters denote significant differences in NO_3^- isotopic composition among sites (e.g., a site with ab letters is significantly different from sites with letters from c onwards, but not significantly different from sites with a or b letters).

Figure 3 Seasonal variations of $\Delta^{17}\text{O-NO}_3^-$ and $\delta^{15}\text{N-NO}_3^-$ (‰) in streams. Point colors denote isotopes as indicated in legend. Range of corresponding f_{atm} (%) is indicated on the secondary y-axis. on the red y-axis). Highlighted in grey is the dormant season (October-April). Note the different y-axes scales across the different panels.

	Mean discharge-weighted annual $[\text{NO}_3^-]$ (mgL^{-1})		Mean discharge-weighted f_{atm} (%) based on $\Delta^{17}\text{O}$ (‰)		Total NO_3^- flux ($\text{kg N ha}^{-1} \text{yr}^{-1}$)		Total $\text{NO}_3^-_{atm}$ flux ($\text{kg N ha}^{-1} \text{yr}^{-1}$)	
	2015	2016	2015	2016	2015	2016	2015	2016
Lower montane	1.0	0.9	22	19	1.9	1.4	0.4	0.3
Upper urban	1.0	1.2	5	4	2.0	2.6	0.1	0.1
Lower urban	1.7	1.3	4	5	4.8	2.1	0.2	0.1

Table 2 Mean discharge-weighted annual $[\text{NO}_3^-]$ and associated (mgL^{-1}) and f_{atm} (%), annual fluxes of total NO_3^- and total $\text{NO}_3^-_{atm}$ in streams and deduced annual contribution of $\text{NO}_3^-_{atm}$ yield to the total NO_3^- flux at three sites along the montane to urban gradient for years 2015 and 2016.

Figure 4 Relationships of nitrate isotopes and solutes concentration with discharge at three sites along the montane to urban gradient. Linear correlations were plotted only if significant. Note the differences in the y-axis scales among the panels, and the specific x-axis for discharge at the lower montane site (in green). Point colors denote sites as indicated in legend.

Figure 5 Relationships of NO_3^- isotopes, solute concentration and discharge with conductivity during snowmelt (April 8 to April 24 in 2016) at the S-upper montane (in red) and the lower montane (in green) sites. Linear regressions were plotted only if significant. Full or open circles feature either a) different NO_3^- isotopes or b) and c) different solute concentrations.

Figure 6 Dual isotopes plot ($\delta^{18}\text{O}$ vs $\delta^{15}\text{N}$) illustrating the mixing between distinct sources of NO_3^- (non-atmospheric) in streams. All isotopic values were obtained by the $\Delta^{17}\text{O}$ transform of NO_3^- collected at all sites (see *Methods*). The colored boxes represent the reported range for NO_3^- terrestrial sources, featuring atmospheric ammonium ($\text{NH}_4^+_{\text{atm}}$) in grey, mineralized ammonium ($\text{NH}_4^+_{\text{org}}$) in turquoise, manure or sewage derived NH_4^+ in beige and fertilizers NO_3^- in violet (Kendall et al., 2007). Linear regressions were plotted only when significant. The blue diamond reflects the isotopic composition of $\text{NO}_3^-_{\text{atm}}$ as measured in this study. Point colors denote sites as indicated in legend.

	Mean annual $\delta^{18}\text{O}-\text{NO}_3^-$ (‰)	Mean annual $\delta^{18}\text{O}-\text{NO}_3^-_{\text{ter}}$ (‰) and deduced % $\text{NO}_3^-_{\text{atm}}$	Calculated $\delta^{18}\text{O}-\text{NO}_3^-_{\text{nit}}$ (‰) and deduced % $\text{NO}_3^-_{\text{atm}}$
S-upper montane	0.7 ± 6.6	-7.7 ± 4.8 11 ± 7^a	-3.3 ± 3.4 5 ± 9^b
N-uppermontane	2.8 ± 4.2	-6.3 ± 5.6 12 ± 7^a	-3.3 ± 3.4 8 ± 6^b
Mid-montane	-4.9 ± 4.1	-8.0 ± 3.7 4 ± 3^a	-3.3 ± 3.4 0 ± 5^b
Lower montane	4.1 ± 6.6	-6.8 ± 6.7 14 ± 8^a	-3.3 ± 3.4 10 ± 9^b
Lake	8.6 ± 6.2	-7.3 ± 4.8 20 ± 7^a	-3.3 ± 3.4 16 ± 8^a
Upper urban	0.9 ± 5.7	-2.8 ± 5.3 5 ± 2^a	2.3 ± 4.2 0 ± 4^b
Lower urban	1.1 ± 3.8	-2.6 ± 3.5 5 ± 3^a	2.3 ± 4.2 0 ± 6^b

Table 3 The second column gives the mean annual $\delta^{18}\text{O}-\text{NO}_3^-$ (‰) measured in streams. The third column shows the mean annual $\delta^{18}\text{O}-\text{NO}_3^-_{\text{ter}}$ (‰) after samples isotopic composition was corrected for the atmospheric imprint (see *Methods*), and the corresponding proportion of $\text{NO}_3^-_{\text{atm}}$ when using the corrected value as microbial end-member. Fourth column presents the calculated theoretical $\delta^{18}\text{O}-\text{NO}_3^-_{\text{nit}}$ when using the approach detailed in section 4.3, and the corresponding proportion of $\text{NO}_3^-_{\text{atm}}$ when using this theoretical value as microbial end-member. Superscript letters indicate significant differences in the proportion of $\text{NO}_3^-_{\text{atm}}$ inferred by both methods ($p < 0.05$).