Meiofauna versus macrofauna as a food resource in a tropical intertidal mudflat

Pierre-Yves Pascal, Pierrick Bocher, Christel Lefrançois, Hien Duy Nguyen, Johan Chevalier, Christine Dupuy

► To cite this version:

Pierre-Yves Pascal, Pierrick Bocher, Christel Lefrançois, Hien Duy Nguyen, Johan Chevalier, et al.. Meiofauna versus macrofauna as a food resource in a tropical intertidal mudflat. *Marine Biology*, 2019, 166 (11), pp.144. 10.1007/s00227-019-3588-z . hal-02355875

HAL Id: hal-02355875

<https://hal.science/hal-02355875>

Submitted on 8 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Meiofauna *versus* macrofauna as a food resource in a tropical intertidal mudflat

Pierre-Yves Pascal^{1*}, Pierrick Bocher², Christel Lefrançois²

Hien T. Nguyen^{2,3}, Johan Chevalier⁴, Christine Dupuy²

¹ Institut de Systématique, Evolution, Biodiversité (ISYEB) UMR 7205, Equipe Biologie de la Mangrove -
Université des Antilles, 97159 Pointe-à-Pitre, Guadeloupe, France

² Littoral, Environnement et Sociétés (LIENSs) UMR 7266 CNRS – La Rochelle University, 2 Rue Olympe de
Gouges, 17000, La Rochelle, France

³ University of Science and Technology of Hanoi (USTH) - Vietnam Academy of Science and Technology
(VAST), 8 Hoans Quoc Viet Street, 10000 Hanoi, Viet Nam

⁴ Réserve Naturelle Nationale de l’Amana, Awala-Yalimapo, French Guiana

*Corresponding author: pierre-yves.pascal@univ-antilles.fr

Key Words: foodweb, mudflat, goby, shrimp, crab, Tanaidacea, diet

21

22 **Abstract**

23 Evaluations of the functioning of benthic marine food webs could be improved by
24 quantifying organic matter fluxes from the meiofauna to higher trophic levels. In this study,
25 we measured the simultaneous ingestion of meiofauna and macrofauna by common dwellers
26 of a tropical intertidal mudflat on the coast of Amazonia. The meiofauna and macrofauna
27 (tanaid) communities of a tropical intertidal mudflat of French Guiana were separately
28 enriched with ^{15}N and ^{13}C , respectively. The enriched preys were then used as tracers during
29 feeding experiments with common predators of different sizes and feeding mechanisms: a
30 Portunidae crab (*Callinectes bocourti*), a Penaeidae shrimp (*Farfantepenaeus subtilis*) and a
31 Gobiidae fish (*Gobionellus oceanicus*). In feeding experiments with all predators except
32 crabs, feeding rates increased with the availability of meiofauna and macrofauna food
33 sources. The ability of consumers to ingest their food selectively was evaluated by calculating
34 the differences in the ratio of macrofauna to meiofauna between the *i*) ingested material and
35 *ii*) that available in the environment. Larger predators showed a higher degree of preferential
36 macrofauna ingestion than smaller predators, consistent with the optimal foraging theory. For
37 large predators, the meiofauna would be important only during early life or in the absence of
38 large food items.

39

40

41 **Introduction**

42 Marine sediments constitute one of the largest environments in the world, in terms of
43 the area covered. They form the habitat of the meiofauna, a discrete group of small (passing
44 through a 0.5 mm-mesh sieve), highly diverse and abundant organisms (Giere 2009).
45 Meiofauna biomass varies considerably between habitats. It is generally smaller than that of
46 other benthic components, but the meiofauna could plays a key role in this ecosystem, due to
47 its high levels of production and activity (Schwinghamer et al. 1986, Moens et al. 2013). The
48 meiofauna move through the sediment and construct burrows, thereby conveying oxygen and
49 nutrients vertically and increasing organic matter mineralization and nutrient cycling (Aller &
50 Aller 1992, Coull 1999, Bonaglia et al. 2014). The meiofauna also supplies ecosystem
51 services, such as nutrient cycling and waste processing, and it plays an important role in food
52 webs (Schratzberger & Ingels 2018). The meiofauna is highly diverse, with a large array of
53 feeding strategies, resulting in the consumption of a broad spectrum of food sources,
54 including microphytobenthos (Middelburg et al. 2000, Moens et al. 2002) and, to a lesser
55 extent, bacteria (van Oevelen et al. 2006, Pascal et al. 2009). The meiofauna may, therefore,
56 represents a major link between small food items and higher trophic levels (Coull 1999). The
57 meiofauna is a high-quality food source, as the animals in this population can biosynthesize
58 and accumulate highly unsaturated fatty acids (Fleeger 2007, Leduc et al. 2009, De Troch et
59 al. 2012, Braeckman et al. 2015) that most metazoans are unable to produce. Numerous
60 studies have demonstrated the importance of the meiofauna in the diet of epibenthic predators
61 (Coull 1990), such as fishes (Fitzhugh & Fleeger 1985, Henry & Jenkins 1995), shrimps (Bell
62 & Coull 1978, Nilsson et al. 1993) and crabs (Scherer & Reise 1981). Many predators present
63 developmental shifts in food preference, with the young preferentially ingesting meiofauna
64 and adults preferentially ingesting larger food items (Coull 1990, 1999, Nilsson et al. 1993).

65 The criteria governing the selection of meiofauna *versus* macrofauna by predators, as a
66 function of the respective availabilities of these two food sources, remain unclear

67 The simplest approach to determining trophic linkages between a consumer and its prey
68 is the direct observation of feeding behavior (Majdi et al. 2018). However, such observations
69 are particularly difficult in the fine sediment of the benthic environment, due to the high
70 turbidity of the overlying water, enhanced by from the suspension of sediment due to the
71 movements of foraging consumers (Kneib 1985). As an alternative approach, a visual analysis
72 can be performed on the gut contents of consumers, but such studies are subject to biases due
73 to differences in digestion rates between food items (Alheit & Scheibel 1982, Scholz et al.
74 1991). The overall impact of predation on the meiofauna can be evaluated by excluding
75 predators over small (Hall et al. 1990) or large scales (Carpenter et al. 1995, Deegan et al.
76 2007). However, the effects of predation are confounded with indirect effects, such as
77 depletion of the meiofaunal food source (Ólafsson et al. 1993) and/or physical disturbance
78 due to regular bioturbation by predators (Austen & Widdicombe 1998, Schratzberger &
79 Warwick 1999, Ólafsson 2003, Fleeger et al. 2008). The use of stable isotopes is considered
80 to be a powerful tool for determining nutrient sources for consumers, as stable carbon and
81 nitrogen ratios are largely determined by diet (Fry 2006). The full potential of stable isotopes
82 is closely linked to the discrimination of potential food sources in terms of isotopic
83 composition. Stable isotopes can also be used as tracers, after the artificial enrichment of
84 preys. This approach has been used to evaluate the ingestion of bacteria and
85 microphytobenthos by meiofauna (Moodley et al. 2002, Pascal et al. 2008b, Pascal et al.
86 2008c), macrofauna (Pascal et al. 2008a, Leroy et al. 2012) and fishes (Como et al. 2018). A
87 similar approach has also been used to measure the consumption of macroalgae (Pascal &
88 Fleeger 2013, Legrand et al. 2018). However, to our knowledge, this approach has never been
89 used to assess the ingestion of meiofauna and macrofauna by predators.

90 Improvements in our understanding of the complexity of benthic marine food webs
91 require both *i)* improvements in the quantification of the fraction of the meiofauna transferred
92 to higher trophic levels and *ii)* evaluations of potential variations of this trophic flux
93 according to the availability of other food sources. The aim of this study was to measure the
94 simultaneous ingestion of meiofauna and macrofauna by common dwellers of the tropical
95 intertidal mudflat of the coast of Amazonia.

96

97

Materials and methods

Study site

This study was performed in the intertidal mudflat of Awala-Yalimapo (05°44'44"N, 53°55'36"W), in Western French Guiana, 850 km away from the mouth of the Amazon River (Fig 1). The study site has a humid tropical climate, with a wet season from January to July and a dry season from August to the end of December. This coast is subject to a semidiurnal tidal cycle, with a tidal range of 0.8 m (neap tides) to 2.9 m (spring tides). Over the course of the year, mud temperature oscillates between 29 and 33°C, sediment pore water salinity varies between 12 and 46 and sediment organic content fluctuates between 5.6 and 6.8% (Nguyen et al. 2018). The sampling station was located in the upper area of the intertidal mudflat, in very fluid mud adjacent to an area of consolidated mud with young mangrove trees.

In this area, meiofaunal abundance fluctuates between 1.7×10^6 and 4.4×10^6 ind.m⁻², equivalent to dry weights of 3.3 and 8.3 g m⁻², respectively, and the community is dominated by nematodes, with harpacticoid copepods and other groups (Ostracods, Platyhelminthes and Molluscs) accounting for only 26% and less than 1%, respectively, of the benthic meiofauna (Dupuy et al. 2015). The term “macrofauna” is used here for animals with a length of 0.5 mm to 20.0 mm. The macrofauna is not very diverse and is dominated by a small number of few taxa, such as tanaids, with the species *Halmyrapseudes spaansi* accounting for 84% of macrofauna specimens, with mean abundances from 2,600 ind.m⁻² to up to 73,000 ind.m⁻² corresponding to a maximum dry weight of 12.1 g m⁻² (Jourde et al. 2017; Nguyen et al. 2018). The macrofauna also includes members of the Polychaetes and, to a lesser extent, Gastropods and Nemertea (Jourde et al. 2017). The macrofaunal community is dominated by small species, such as *H. spaansi*, which never exceeds 6.4 mm in length (Nguyen et al. 2018). In this study, organisms of more than 20 mm in length, with the potential to ingest meiofauna and macrofauna, are considered to be predators. We focused on the most common

123 predators at this site, collected locally: the crab *C. bocourti*, the shrimp *F. subtilis* and the
124 highfin goby *G. oceanicus*. The abundances of *C. bocourti* and *F. subtilis* have never been
125 evaluated in the study area. The abundance of the high fin goby *G. oceanicus* was previously
126 estimated from benthic cores, which showed this species to have a patchy distribution, with
127 densities of up to 50 ind. m⁻² (Jourde et al. 2017). All feeding experiments were performed
128 during the dry season, in November 2015.

129 *Predator sampling*

130 The uppermost centimeter of the sediment was collected from the mudflat at low tide,
131 with a large rectangular scoop. The sediment was sieved with 63 µm and 500 µm meshes, to
132 extract the meiofauna and macrofauna, respectively. *C. bocourti* and *F. subtilis* were collected
133 at high tide, with a 500 µm-mesh landing net, *G. oceanicus* specimens were collected by
134 hand, from their burrows, at low tide.

135 *Preparation of enriched preys*

136 The study site has two advantages for studies of this type: *i*) easy extraction of the
137 meiofauna and macrofauna by sieving, because the sediment consists primarily of particles of
138 less than 63 µm in diameter (Dupuy et al. 2015) and *ii*) ease of experimental manipulation of
139 the macrofaunal community, due to its low diversity, dominated by a species of small tanaids
140 (Jourde et al. 2017, Nguyen et al. 2018). Sediment was sampled as previously described. For
141 the accurate control of prey availability, sediment depleted of as many meiofauna and
142 macrofauna as possible was required. The sediment was therefore sieved, and the fraction
143 passing through a 63 µm mesh was allowed to settle overnight, after which the overlying
144 water was removed (Fig. 2). A 1 cm layer of this deposited sediment and 1 liter of local
145 seawater, filtered through a 50 µm-mesh sieve, were placed in each 200 cm² aquarium used
146 for feeding experiments (*n*=16 and control (*n*=1); Table 1). Enriched ¹³C glucose and ¹⁵N
147 ammonium were added separately to the sediment to label the bacteria and/or
148 microphytobenthos for the secondary labeling of their grazers: the meiofauna and

macrofauna. For the preparation of labeled meiofauna, the sediment was sieved and the fraction passing through a 500 μm mesh was allowed to settle overnight. This fraction of deposited sediment was mixed with $^{15}\text{NH}_4\text{Cl}$ (99% ^{15}N -enriched NH_4Cl ; Euriso-top) at a final concentration of 4 g m^{-2} and incubated. For the preparation of labeled macrofauna, the collected sediment was mixed directly with ^{13}C -glucose (99% ^{15}N -enriched glucose; Euriso-top) at a final concentration of 2.5 g m^{-2} and incubated. Each type of sediment was i) placed in a separate large plate, at a thickness of 0.5 cm, ii) incubated under local conditions of temperature (29-33°C (Nguyen 2018)) and irradiance (2000 $\text{kW m}^{-2} \text{y}^{-1}$ (Elana 2017)), and iii) regularly sprayed with freshwater to compensate for evaporation. After four days of incubation, each type of sediment was independently sieved through separate meshes to collect ^{15}N -enriched meiofauna (63 μm) and ^{13}C -enriched macrofauna (500 μm). Each of the 16 aquaria (200 cm^2) used for feeding experiments was filled with 1 L of seawater and a 1 cm layer of fauna-free sediment, mixed with variable amounts of enriched meiofauna (^{15}N) and/or macrofauna (^{13}C) and one type of target predator. Species community compositions of enriched preys (meiofauna and macrofauna) were similar in all experiments. First, we increased the abundance of one resource (meiofauna or macrofauna), whilst making the other resource unavailable. Secondly both preys were available simultaneously, in different proportions (table 1). One gram (dry weight) of prey corresponds to 4.6×10^6 meiofauna specimens and 6.1×10^3 macrofauna specimens.

Feeding experiments

Predators were sampled as described above. All predators were kept for less than six hours in a large tank containing oxygenated local seawater and sediment with associated fauna before feeding experiments, to prevent starvation. We released 6 *C. bocourti*, 15 *F. subtilis* or 15 *G. oceanicus* specimens into each of the experimental aquaria. In the control aquarium, containing the highest abundance of each enriched prey, 15 specimens of *G. oceanicus* were placed in a cage closed with a 63 μm mesh, to obtain control predators unable

175 to ingest enriched preys. All incubations with predators were performed in oxygenated
176 aquaria, for 2 hours, under local temperature and irradiance conditions. At the end of the
177 incubation period, the predators were collected in an aquarium fish net and immediately
178 frozen (-18°C).

179 *Isotope analyses and calculations*

180 For the meiofauna, each stable isotope sample consisted of 700 nematodes or 300
181 copepods picked at random, by hand, under a dissecting microscope, after extraction from
182 sediment by sieving through a 63 µm mesh and Ludox HS40 centrifugation, and pooled
183 (Jonge & Bouwman 1977). Macrofauna were extracted from sediment by sieving through a
184 500 µm mesh, and six tanaid specimens per sample were pooled. For each set of conditions,
185 three pools of two *C. bocourti*, five *F. subtilis* and five *G. oceanicus* specimens, selected at
186 random, were homogenized in an Ultra-Turrax blender, frozen and freeze-dried.

187 The nitrogen and carbon isotopic compositions of the various organisms sampled (preys
188 and predators) were determined by EA-IRMS (Isoprime, Micromass, UK). The nitrogen and
189 carbon isotope ratios are expressed in the delta notation $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$, as follows: $\delta X =$
190 $[(R_{\text{Reference}}/R_{\text{Sample}}) - 1] \times 1000$, where $X = \delta^{15}\text{N}$ or $\delta^{13}\text{C}$ and R is the ratio $^{15}\text{N}:^{14}\text{N}$ or $^{13}\text{C}:^{12}\text{C}$
191 in the sample and in the reference material. Results are expressed relative to atmospheric
192 nitrogen for N and to Vienna Pee Dee Belemnite (VPDB) for C, and are expressed in units of
193 ‰ $\pm$ standard deviation (SD).

194 An excess of ^{15}N , above background levels (natural ^{15}N of predator without enrichment)
195 was considered to constitute ^{15}N enrichment, and is expressed in terms of specific uptake (I). I
196 was calculated as the product of excess ^{15}N (E) and the previously measured N biomass per
197 predator (Table 2). I was converted into meiofaunal or macrofaunal carbon ingested, with the
198 C/N ratio of each prey. E was defined as the difference between the background ($F_{\text{background}}$)
199 and sample (F_{sample}) ^{15}N fractions: $E = F_{\text{sample}} - F_{\text{background}}$, with $F = ^{15}\text{N}/(^{15}\text{N} + ^{14}\text{N}) = R / (R +$
200 $2)$ and R = the nitrogen isotope ratio. For $F_{\text{background}}$, we used values measured with predators,

201 without enrichment. R was derived from the measured $\delta^{15}\text{N}$ values: $R = [(\delta^{15}\text{N}/1000) + 1] \times$
202 R_{airN_2} . Prey intake was calculated as $[I \times (\text{C/N ratio of enriched prey}) / (F_{\text{enriched prey}} \times$
203 incubation time)] (Pascal et al., 2008).

204 The incorporation of ^{13}C was calculated in a similar manner, with $F = ^{13}\text{C} / (^{13}\text{C} + ^{12}\text{C}) =$
205 $R / (R+1)$, R_{airN_2} replaced by $R_{\text{VPDB}} = 0.0112372$ and $\text{Uptake} = I / (F_{\text{enriched predator}} \times \text{incubation}$
206 time)

207 *Data analyses*

208 One-way analysis of variance (ANOVA) with Tukey post hoc tests was used to analyze
209 the differences between ingestion rates. The normality of the data was first checked in a
210 Shapiro-Wilk test, and homoscedasticity was checked with a Bartlett test. The nonparametric
211 Kruskal-Wallis test was used to assess the differences in isotopic composition ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$)
212 of preys and predators. The selectivity of prey ingestion was evaluated by fitting a linear
213 model to the data for the ratio of the biomasses of macrofauna to meiofauna ingested, with a
214 95% confidence interval. All statistical analyses were performed with R. Values are presented
215 as means $\pm$ standard deviation (SD) unless otherwise specified.

216

Results

The individual dry weight, carbon and nitrogen contents of each type of prey and predator are presented in table 2. Tanaids had a mean individual dry weight 750 times higher than that of nematodes and 300 times higher than that of copepods. The natural isotopic compositions of preys and predators are presented in figure 3. Before enrichment, $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ levels were similar in the meiofauna and macrofauna (Tanaidacea) (Kruskall-Wallis, $p > 0.05$). After enrichment, the $\delta^{15}\text{N}$ levels of the nematodes and copepods used during feeding experiments were not significantly different (Kruskall-Wallis, $p > 0.05$), with respective enrichment values (δ after enrichment minus δ at T0) of $261.0 \pm 44.5 \text{ ‰}$ ($n = 6$ samples of 700 specimens) and $268.0 \pm 24.9 \text{ ‰}$ ($n = 6$ samples of 300 specimens). Tanaids had a mean $\delta^{13}\text{C}$ content of $2789.5 \pm 65.6 \text{ ‰}$ ($n = 3$ samples of 6 specimens each).

C. bocourti was the predator with the lowest individual weight, followed by *F. subtilis* and *G. oceanicus* (Tab. 2). *C. bocourti* had a mean width of $1.3 \text{ cm} \pm 0.4$ ($n = 42$) whereas *F. subtilis* and *G. oceanicus* were $4.4 \text{ cm} \pm 0.9$ ($n = 210$) and $5.0 \text{ cm} \pm 2.0$ ($n = 180$) long, respectively. The natural isotopic compositions of all predators (Fig. 3) were similar for $\delta^{13}\text{C}$ (Kruskall-Wallis, $p > 0.05$), whereas *F. subtilis* and *G. oceanicus* had significantly different $\delta^{15}\text{N}$ levels (Kruskall-Wallis, $p < 0.01$).

G. oceanicus caged in the aquarium with the highest abundances of each of the enriched preys had significantly higher ^{13}C and ^{15}N levels than the natural isotopic composition samples (Kruskall-Wallis, $p < 0.05$), by 1.5 ± 0.5 and $2.1 \pm 0.7 \text{ ‰}$, respectively ($n=3$). These enrichment levels correspond to 2.7% for ^{13}C and 7.1% for ^{15}N relative to free specimens of *G. oceanicus* able to ingest enriched preys in similar incubation conditions.

Meiofauna ingestion by *C. bocourti* was not significantly affected by meiofauna abundance, and the values obtained ranged from 11.8 to $18.9 \text{ mgC}_{\text{prey}} \text{ gC}_{\text{predator}}^{-1} \text{ h}^{-1}$, which is

242 equivalent to 3584 to 5766 nematode ind⁻¹ h⁻¹ (Fig. 4). By contrast, the ingestion of
243 meiofauna by *F. subtilis* and *G. oceanicus* increased with meiofauna abundance and was
244 significantly higher at the highest meiofauna abundance tested (Fig. 4), at reaching 12.7 ± 1.0
245 $\text{mgC}_{\text{prey}} \text{gC}_{\text{predator}}^{-1} \text{h}^{-1}$ (6342 ± 516 nematode ind⁻¹ h⁻¹) and $4.6 \pm 0.5 \text{ mgC}_{\text{prey}} \text{gC}_{\text{predator}}^{-1} \text{h}^{-1}$
246 (3120 ± 363 nematodes ind⁻¹ h⁻¹), respectively.

247 The ingestion of macrofauna by *C. bocourti* was not significantly affected by prey
248 abundance and ranged from 1.2 to 4.5 $\text{mgC}_{\text{prey}} \text{gC}_{\text{predator}}^{-1} \text{h}^{-1}$, equivalent to 2.0 to 2.9 tanaid
249 ind⁻¹ h⁻¹ (Fig. 5). By contrast, the ingestion of macrofauna by *F. subtilis* and *G. oceanicus*
250 increased significantly with macrofauna abundance, reaching $5.6 \pm 0.2 \text{ mgC}_{\text{prey}} \text{gC}_{\text{predator}}^{-1} \text{h}^{-1}$
251 (5.7 ± 0.2 tanaid ind⁻¹ h⁻¹) and $3.8 \pm 0.4 \text{ mgC}_{\text{prey}} \text{gC}_{\text{predator}}^{-1} \text{h}^{-1}$ (5.4 ± 0.5 tanaid ind⁻¹ h⁻¹),
252 respectively (Fig. 5).

253 The ability of predators to ingest their food in a selective manner was evaluated by
254 calculating the differences between the ratios of macrofauna to meiofauna ingested and ratios
255 of macrofauna to meiofauna available in the environment (Fig. 6). If these two ratios were
256 similar, it was concluded that there was no selection, whereas differences between these ratios
257 were interpreted as indicating the preferential ingestion of meiofauna or macrofauna. For each
258 predator, we fitted a linear model with a 95% confidence interval to the data (Fig. 6). For *C.*
259 *bocourti*, this confidence interval included all theoretical values for non-selective ingestion,
260 reflecting an absence of selection in feeding behavior. For *F. subtilis* and *G. oceanicus*, the
261 lowest values of the 95% confidence interval were always higher than the theoretical values,
262 implying a preferential ingestion of macrofauna over meiofauna by both these predator
263 species.

264

265

266

Discussion

267

268

269

The aim of this study is improve our understanding of benthic marine food webs by the quantification of ingestion of meiofauna and macrofauna by higher trophic levels in a tropical intertidal mudflat on the coast of Amazonia

270

Meiofauna predation

271

272

273

274

275

276

277

278

279

The effect of top-down control on the meiofauna is unclear. Several studies have suggested that predation on the meiofauna is negligible due to the ability of the meiofauna to disperse (Giere 2009) and fast turnover times, whereas predators are comparatively rare (Gibbons 1988, Shaw & Jenkins 1992, Coull 1999). On the contrary, other studies have concluded that predation affects meiofauna abundance (Danovaro et al. 2007, Fleeger et al. 2008) and community composition (Li et al. 1996, O'Gorman et al. 2008). These discrepancies may reflect biases associated with the different methods used to evaluate trophic fluxes (see below in the part *Methodological considerations*). Differences in the conclusions drawn may also reflect differences in trophic links between study sites.

280

281

282

283

284

285

286

287

288

289

Estimations of the top-down effect of predators require knowledge of their abundance. In the study area, the abundances of *C. bocourti* and *F. subtilis* have unfortunately never been evaluated. However, abundances of the high fin goby *G. oceanicus* have been estimated from measurements in benthic cores, revealing a patchy distribution, with densities reaching 50 ind. m⁻² (Jourde et al. 2017). For evaluation of the maximum top-down effect, daily ingestion was calculated, taking into account the highest abundance of fish and lowest abundance of meiofauna (Dupuy et al. 2015), and the highest ingestion rate measured in this study. This area of the mudflat is submerged and exposed to fish predation for only six hours per day (24 h). This maximum daily ingestion would therefore reach 56% of the standing stock of meiofauna in Awala. However, this assumption must be viewed with caution, because *i*)

290 grazers, such as shrimps and crabs, were not considered, as their natural abundances remain
291 unknown and *ii*) gobies have a patchy distribution and are not realistically sampled by benthic
292 cores. However, our results nevertheless suggest that the meiofauna community could be
293 regulated by predation, at least during the limited periods during which predators are
294 abundant as previously suggested by other studies (Danovaro et al. 2007, Fleeger et al. 2008).

295 *Meiofauna predators*

296 Body size is an important determinant of many physiological processes, and maximal
297 ingestion rate is generally inversely correlated with body size (Peters, 1983). This assumption
298 was confirmed in this study, as maximum rates of meiofauna ingestion decreased with
299 increasing predator weight, from crab to fish. In shrimp, the maximum daily meiofauna
300 ingestion rate corresponded to 30% of the predator's body weight. This rate is higher than that
301 estimated for the shrimp *Crangon crangon*, which eats 8-16% (del Norte-Campos &
302 Temming 1994, Feller 2006) or 10-12% (Pihl & Rosenberg 1984) of its own body weight
303 daily. This higher level of ingestion activity may be due to different evaluation techniques or
304 to the higher temperature in the tropical environment of *F. subtilis* than in the habitat of *C.*
305 *crangon*.

306 Theoretically, food uptake by a consumer should increase with food abundance.
307 However, beyond a certain threshold prey density, uptake rates remain constant, according to
308 Holling's prey-dependent type II functional response (Holling 1959), as reported for shrimp
309 (Gregg & Fleeger 1998). The threshold value was not reached here for *F. subtilis* and *G.*
310 *oceanicus*. As the natural abundance of the meiofauna (Dupuy et al. 2015) and macrofauna
311 (Nguyen et al. 2018) can exceed the abundances tested here, ingestion rates could be higher
312 than the measured values, for both preys.

313 The meiofauna constitutes a potentially high-quality food source, due to the high
314 levels of unsaturated fatty acids in these animals (Fleeger 2007, Leduc et al. 2009, De Troch
315 et al. 2012, Braeckman et al. 2015). The meiofauna is strongly associated with sediment

316 particles. When consuming the meiofauna, predators can *i*) ingest bulk sediment, which
317 entails a high cost in terms of the rejection of non-digestible material or *ii*) selectively ingest
318 the meiofauna, which entails a high cost in food selection. In this study, the three predators
319 have different feeding mechanisms for selecting their food. Juvenile crabs of the genus
320 *Callinectes* selectively ingest microbes rather than bulk detritus (Fantle et al. 1999). Penaid
321 shrimps feed selectively, with their chelate pereopods grasping and transporting acceptable
322 morsels of food to the mouth (Hindley & Alexander 1978). Gobies graze more or less
323 indiscriminately on sediment, in addition to predating on larger preys on sight (Fitzhugh &
324 Fleeger 1985, Hamerlynck & Cattrijsse 1994). The predators studied have different weight
325 ranges, with the crab *C. bocourti* smaller than the goby *G. oceanicus* by a factor of 1.6.
326 According to optimal foraging theory, that the greatest rewards per unit effort are obtained by
327 predating on the largest graspable prey items (Krebs 1978). Development shifts in diet, from
328 smaller meiofauna to larger macrofauna, have been reported for the crab *Carcinus maenas*
329 (Reise 1978, Scherer & Reise 1981). The shrimp *Crangon crangon* displays a similar dietary
330 shift when its total body length exceeds 20 mm (Pihl & Rosenberg 1984, Gee 1987).
331 Similarly, in our study, the shrimp *F. subtilis*, with a mean length of 44 mm, also
332 preferentially ingested macrofauna. For fish, the switch from meiofauna to macrofauna
333 appears to occur at a threshold total length of 3-4 cm (Gee 1989), as previously reported for
334 several goby species from the Adriatic Sea (Kovačić & la Mesa 2008), Baltic Sea (Aarnio &
335 Bonsdorff 1993), North Sea (Schückel et al. 2013) and Atlantic Ocean (D'Aguillo et al. 2014).
336 The results reported here are consistent with this assumption, as gobies of 5 cm or more in
337 length preferentially ingested macrofauna.

338 The experiments reported here suggest an important role of the macrofauna and
339 meiofauna in the diet of *G. oceanicus*, but natural isotopic composition revealed limited $\delta^{15}\text{N}$
340 enrichment between the preys and the goby, suggesting that other food items not considered
341 here, such as detritus or microphytobenthos, might play a role in the diet of these fish. Despite

342 preferential macrofauna ingestion, large *G. oceanicus* specimens continued to ingest
343 meiofauna, unlike other goby species of similar size (de Morais & Bodiou 1984). Gobies can
344 be opportunist predators with a diet reflecting variations in the food available (D'Aguillo et al.
345 2014). In the Awala mudflat, the abundance of tanaids varies by a factor of 10^3 , depending on
346 the season (Nguyen et al. 2018). The meiofauna may, therefore, play an important role in the
347 diet of the predator in conditions of lower macrofauna availability. Such seasonal variations in
348 the trophic role of the meiofauna have already been suggested for fish (Schückel et al. 2013).
349 Gobies switch to the ingestion of smaller items when in competition for food resources or if
350 the availability of large preys decreases (Henry & Jenkins 1995). Fluxes from the meiofauna
351 to higher trophic levels appeared complexes and likely variable along the year suggesting that
352 a temporal survey would be useful to clarify the trophic role of meiofauna. Despite those
353 potential variations, the present study provides quantitative fluxes embeddable in ecological
354 models of benthic food webs.

355 *Methodological considerations*

356 The potential of stable isotopes for studies of this kind depends heavily on the
357 difference in isotopic composition between the food sources to be distinguished. In this study,
358 the meiofaunal community and tanaids had similar isotopic compositions, rendering
359 interpretation of the patterns obtained with natural isotopes difficult and necessitating the use
360 of enrichment experiments.

361 Predator abundance in the study area had been determined only for *G. oceanicus*
362 (Jourde et al. 2017). For the purpose of feeding, the predators studied perform tidal
363 migrations, following the rising tide by swimming or walking over the sediment on the
364 bottom (Gibson, 2003). They can reach very high densities during this migration, and the
365 shrimp (*F. subtilis*) and crab (*C. bocourti*) species studied were chosen on the basis of their
366 high abundance and availability at the time of the experiments. Incubations were performed

367 with high abundances of predators, potentially affecting predator interactions and feeding
368 behavior, resulting in a possible underestimation of ingestion rates.

369 Copepods have a caloric value 35% higher than those of most members of the
370 meiofauna (Gee 1989). Copepods generally dwell closer to the sediment surface than
371 nematodes, and display emergence behavior relative to the water column (Giere 2009). This
372 differential distribution would facilitate predation on copepods by shrimps (Pihl & Rosenberg
373 1984, Gee 1987) and fishes (Coull 1990, McCall & Fleeger 1995). However, it was not
374 possible to differentiate between copepods and nematodes in this study, because these two
375 groups were labeled similarly. The availability of sediment in the aquarium constituted
376 realistic incubation conditions, because the sediment serves as a refuge, reducing the risk of
377 predation for meiofauna and macrofauna (Arnold 1984, Lipcius & Hines 1986, Gregg &
378 Fleeger 1998). Feeding experiments were run over a short period of two hours and did not,
379 therefore, necessarily yield results representative of feeding behavior over longer time scales,
380 as the feeding activity of the predators studied is known to fluctuate over the day (Hagerman
381 & Østrup 1980, Ryer 1987, Kanou et al. 2005). However, short incubation times were used
382 here, to limit *i*) label exchange between meiofaunal and macrofaunal preys and *ii*) the
383 excretion of ingested label by predators. This excretion bias would be limited, as feeding
384 experiments lasted two hours and gut retention time would not exceed this duration for
385 Portunidae crabs (Serrano 2012) or Gobiidae fish (Kanou et al. 2005). However, this bias
386 could be larger for Penaidae shrimps, which have gut retention times of between one and four
387 hours (Hoyt et al. 2000).

388 The isotopic compositions of predators changed slightly following their transfer to a
389 mesh cage in an aquarium containing enriched preys. This finding suggests that any bias due
390 to the inefficient removal of unincorporated label by rinsing would be limited, and that the
391 enrichments observed with free predators were due principally to trophic processes. The mud
392 sediment of French Guiana is dominated by silt (70-85%) and clay (10-15%), and mean

393 particle size is below 63 μm (Dupuy et al. 2015). The fauna thus represents a large portion of
394 the material remaining on the 63 μm mesh after sieving. However, a small fraction of detritus
395 would also have been present and the associated bacteria would have taken up glucose- ^{13}C or
396 ammonium- ^{15}N during the enrichment process. The ingestion of this enriched detritus by
397 predators would then lead to an overestimation of the ingestion of meiofauna and macrofauna.
398 This bias should be borne in mind when interpreting the results. Despite this shortcoming, the
399 method used here has several advantages: *i*) the measurement of ingestion fluxes, *ii*) dual
400 labeling of preys, making it possible to evaluate feeding preferences and, *iii*) ease of use, with
401 incubations possible even at remote sites with limited technical resources.

402 **Conclusion**

403 The use of stable isotope-enriched meiofauna and macrofauna in feeding experiments
404 is an appropriate method for evaluating preferential ingestion. In these experiments, feeding
405 rates increasing with the availability of meiofauna and macrofauna food for all predators other
406 than the crabs, suggesting that rates of ingestion of both preys may actually be higher than
407 measured. All three predators studied ingested meiofauna, but the contribution of the
408 meiofauna to the diet decreased with increasing predator size (*C. bocourti* > *F. subtilis* > *G.*
409 *oceanicus*), whereas no such relationship to size was observed for the macrofauna. Larger
410 predators preferentially ingested macrofauna to a greater extent than smaller predators, in
411 accordance with optimal foraging theory. For larger predators, the meiofauna is an important
412 component of the diet only at early stages of development or in the absence of large food
413 items.

414

415

416 **Figure and table**

417 Table 1. Measured prey abundances (dry weights of meiofauna and macrofauna in g m⁻²) in
418 dual-prey feeding experiments in 16 independent aquariums. Each letter refers to a predator.
419 A: *C. bocourti* (*n* = 4), B: *F. subtilis* (*n* = 6) and C: *G. oceanicus* (*n* = 6)

		Meiofauna (g m ⁻²)		
		3.7	9.1	22
Macrofauna (g m ⁻²)	0.9		A, B, C	
	1.3	A, B	A, B	
	1.8	C	B, C	C
	3.7		A, B, C	
	7.3		B, C	

420

421

422

423 Table 2. Dry weight, carbon and nitrogen weights per specimen of prey (nematode, copepod,
424 Tanaidacea) or predator (*C. bocourti*, *F. subtilis*, *G. oceanicus*) (mean $\pm$ SD, *n*: number of
425 samples and number of specimens per sample used to evaluate these weights).

	Individual dry weight	Individual carbon weight	Individual nitrogen weight	n	Specimens per sample
Nematoda	217 $\pm$ 61 ng	94 $\pm$ 29 ng	23 $\pm$ 8 ng	6	700
Copepoda	534 $\pm$ 92 ng	209 $\pm$ 31 ng	47 $\pm$ 7 ng	6	300
Tanaidacea	162 $\pm$ 45 μ g	46 $\pm$ 11 μ g	10 $\pm$ 2 μ g	6	6
<i>C. bocourti</i>	95 $\pm$ 36 mg	29 $\pm$ 13 mg	6 $\pm$ 3 mg	45	2
<i>F. subtilis</i>	122 $\pm$ 41 mg	47 $\pm$ 18 mg	13 $\pm$ 5 mg	57	5
<i>G. oceanicus</i>	156 $\pm$ 92 mg	65 $\pm$ 41 mg	18 $\pm$ 11 mg	54	5

426

427

428

431 Figure 1. Location of the Awala-Yalimapo study site in French Guiana

433

434

Figure 2

435

436 Figure 2. Diagram of the protocol for enrichment of the meiofauna (I) and macrofauna (II):
437 the sediment was a) passed through a 500 µm mesh for I and left untreated for II, b) mixed
438 with $^{15}\text{NH}_4\text{Cl}$ for I and ^{13}C -glucose for II and incubated for four days, c) passed through
439 meshes of 63 µm for I and 500 µm for II, to collect the ^{15}N -enriched meiofauna and ^{13}C -
440 enriched macrofauna, respectively. The enriched fauna was mixed with non-enriched
441 sediment devoid of fauna obtained with a 63 µm-mesh sieve

442

443

444

445

446 Figure 3. Natural isotopic composition ($\delta^{15}\text{N}$ as a function of $\delta^{13}\text{C}$, $\pm$ SD, $n= 6$) of preys
447 (meiofauna and macrofauna) and predators (*C. bocourti*, *F. subtilis* and *G. oceanicus*)

448

450

451 Figure 4. Ingestion of meiofauna (mgC_{prey} gC_{predator}⁻¹ h⁻¹; ± SD, n= 3) by *C. bocourti* (A), *F.*

452 *subtilis* (B) and *G. oceanicus* (C) as a function of dry meiofauna biomass (mg m⁻²). Different

453 lower-case letters indicate significant differences between the tested abundances (ANOVA,

454 Tukey test, *p*<0.005)

455

457

458 Figure 5. Ingestion of macrofauna (mgC_{prey} mgC_{predator}⁻¹ h⁻¹; $\pm$ SD, $n=3$) by *C. bocourti* (A),
459 *F. subtilis* (B) and *G. oceanicus* (C) as a function of dry macrofauna biomass (mg m⁻²).
460 Different lower-case letters indicate significant differences between the tested abundances
461 (ANOVA, Tukey test, $p<0.005$).

462

464
465 Figure 6. Ratio of the biomasses of macrofauna and meiofauna ingested plotted against the
466 ratio of biomasses of macrofauna and meiofauna available ($\pm$ SD, $n=3$), for *C. bocourti* (A),
467 *F. subtilis* (B) and *G. oceanicus* (C). The dotted line corresponds to a theoretical absence of
468 selection. Points above (blue part) or below (red part) the dotted line indicate the selective
469 ingestion of macrofauna and meiofauna, respectively. The linear models fitted to the data are
470 presented as solid lines with 95% confidence intervals

472

473

474

Acknowledgements

475

476

477

478

479

480

481

Compliance with ethical standards

482

Conflict of interest: The authors declare that they have no conflict of interest

483

484

485

486

487

488

489

490

491

492

493

Human and animal rights statement: This article does not contain any studies with human participants performed by any of the authors. All applicable international, national and/or institutional guidelines for the care and use of animals were followed. All procedures performed in the study involving animals were in accordance with the ethical standards of French law regarding maintaining animal welfare in research.

495 **Reference**

- 496 Aarnio K, Bonsdorff E (1993) Seasonal variation in abundance and diet of the sand goby
 497 *Pomatoschistus minutus* (Pallas) in a northern Baltic Archipelago. *Ophelia* 37: 19-30
- 498 Alheit J, Scheibel A (1982) Benthic harpacticoids as a food source for fish. *Mar Biol* 70: 141-147
- 499 Aller RC, Aller JY (1992) Meiofauna and solute transport in marine muds. *Limnol Oceanogr* 37:
 500 1018-1033
- 501 Arnold WS (1984) The effects of prey size, predator size, and sediment composition on the rate
 502 of predation of the blue crab, *Callinectes sapidus* Rathbun, on the hard clam, *Mercenaria*
 503 *mercenaria* (Linné). *J Exp Mar Biol Ecol* 80: 207-219
- 504 Austen MC, Widdicombe S (1998) Experimental evidence of effects of the heart urchin
 505 *Brissopsis lyrifera* on associated subtidal meiobenthic nematode communities. *J Exp Mar*
 506 *Biol Ecol* 222: 219-238
- 507 Bell SS, Coull BC (1978) Field evidence that shrimp predation regulates meiofauna. *Oecologia*
 508 35: 141-148
- 509 Bonaglia S, Nascimento FJA, Bartoli M, Klawonn I, Brüchert V (2014) Meiofauna increases
 510 bacterial denitrification in marine sediments. *Nat Commun* 5: 5133
- 511 Braeckman U, Provoost P, Sabbe K, Soetaert K, Middelburg JJ, Vincx M, Vanaverbeke J (2015)
 512 Temporal dynamics in a shallow coastal benthic food web: insights from fatty acid
 513 biomarkers and their stable isotopes. *Mar Environ Res* 108: 55-68
- 514 Carpenter SR, Chisholm SW, Krebs CJ, Schindler DW, Wright RF (1995) Ecosystem experiments.
 515 *Science* 269: 324-327
- 516 Como S, Carpentier A, Rossi F, Dupuy C, Richard P, Feunten E, Lefrancois C (2018) Stable
 517 isotope as tracers can reveal resource allocation in juvenile golden gray mullets (*Liza*
 518 *aurata*, Risso, 1810). *J Exp Mar Biol Ecol* 503: 72-79
- 519 Coull BC (1990) Are members of the meiofauna food for higher trophic levels? *Trans Am*
 520 *Microsc Soc* 109: 233-246
- 521 Coull BC (1999) Role of meiofauna in estuarine soft bottom habitats. *Aust J Ecol* 24: 327-331
- 522 D'Aguillo MC, Harold AS, Darden TL (2014) Diet composition and feeding ecology of the naked
 523 goby *Gobiosoma bosc* (Gobiidae) from four western Atlantic estuaries. *J Fish Biol* 85: 355-
 524 373
- 525 Danovaro R, Scopa M, Gambi C, Fraschetti S (2007) Trophic importance of subtidal metazoan
 526 meiofauna: evidence from *in situ* exclusion experiments on soft and rocky substrates. *Mar*
 527 *Biol* 152: 339-350
- 528 de Morais T, Bodiou JY (1984) Predation on meiofauna by juvenile fish in a Western
 529 Mediterranean flatfish nursery ground. *Mar Biol* 82: 209-215

530 De Troch M, Boeckx P, Cnudde C, Van Gansbeke D, Vanreusel A, Vincx M, Caramujo M-J (2012)
 531 Bioconversion of fatty acids at the basis of marine food webs: insight from compound-
 532 specific stable isotope analysis. *Mar Ecol Prog Ser* 465: 53-67
 533 Deegan LA, Bowen JL, Drake D, Fleeger JW, Fiedrichs CT, Galván KA, Hobbie JE, Hopkins CS,
 534 Johnson DS, Johnson JM, LeMay LE, Miller E, Peterson BJ, Picard C, Sheldom S, Sutherland
 535 M, Vallino J, Warren RS (2007) Susceptibility of salt-marshes to nutrient enrichment and
 536 predator removal. *Ecol Appl* 17: S42-S63
 537 del Norte-Campos AGC, Temming A (1994) Daily activity, feeding and rations in gobies and
 538 brown shrimp in the northern Wadden Sea. *Mar Ecol Prog Ser* 115: 41-53
 539 Dupuy C, Nguyen TH, Mizrahi D, Jourde J, Bréret M, Agogué H, Beaugéard L, Bocher P (2015)
 540 Structure and functional characteristics of the meiofauna community in highly unstable
 541 intertidal mudbanks in Suriname and French Guiana (North Atlantic coast of South
 542 America). *Cont Shelf Res* 110: 39-47
 543 Elana U (2017) Projet de parc solaire de la Sarcelle sur la commune de Mana (Guyane). SUEZ
 544 Fantle MS, Dittel AI, Schwalm SM, Epifanio CE, Fogel ML (1999) A food web analysis of the
 545 juvenile blue crab, *Callinectes sapidus* using stable isotopes in whole animals and
 546 individual amino acids. *Oecologia* 120: 416-426
 547 Feller RJ (2006) Weak meiofaunal trophic linkages in *Crangon crangon* and *Carcinus maenas*. *J*
 548 *Exp Mar Biol Ecol* 330: 274-283
 549 Fitzhugh GR, Fleeger JW (1985) Goby (Pisces: Gobiidae) interactions with meiofauna and small
 550 macrofauna. *Bull Mar Sci* 36: 436-444
 551 Fleeger JW (2007) The potential to mass-culture harpacticoid copepods for use as food for
 552 larval fish. In: Lee CS, O'Bryen PJ, Marcus NH (eds) *Copepods in Aquaculture*. Blackwell
 553 Publishing Professional, Ames, Iowa, USA
 554 Fleeger JW, Johnson DS, Galván KA, Deegan LA (2008) Top-down and bottom-up control of
 555 infauna varies across the saltmarsh landscape. *J Exp Mar Biol Ecol* 357: 20-34
 556 Fry B (2006) *Stable isotope ecology*. Springer, New-York
 557 Gee JM (1987) Impact of epibenthic predators on estuarine intertidal harpacticoid copepod
 558 populations. *Mar Biol* 96: 497-510
 559 Gee JM (1989) An ecological economic review of meiofauna as food for fish. *Zool J Limn Soc*
 560 96: 243-261
 561 Gibbons MJ (1988) Impact of predation by juvenile *Clinus superciliosus* on phytal meiofauna:
 562 are fish important as predators? *Mar Ecol Prog Ser* 45: 13-22
 563 Gibson RN (2003) Go with the flow: tidal migration in marine animals. *Hydrobiologia* 503: 153-
 564 161
 565 Giere O (2009) *Meiobenthology: the microscopic motile fauna of aquatic sediments*. Springer,
 566 Berlin
 567 Gregg CS, Fleeger JW (1998) Grass shrimp *Palaemonetes pugio* predation on sediment- and

568 stem-dwelling meiofauna: field and laboratory experiments. Mar Ecol Prog Ser 175: 77-86
 569 Hagerman L, Østrup J (1980) Seasonal and diel activity variations in the shrimp *Palaemon*
 570 *adspersus* from a Brackish, non-tidal area. Mar Ecol Prog Ser 2: 329-335
 571 Hall SJ, Raffaelli D, Turell WR (1990) Predator-caging experiments in marine systems: a
 572 reexamination of their value. Am Nat 136: 657-672
 573 Hamerlynck O, Cattrijse A (1994) The food of *Pomatoschistus minutus* (Pisces, Gobiidae) in
 574 Belgian coastal waters, and a comparison with the food of its potential competitor *P.*
 575 *lozanoi*. J Fish Biol 44: 753-771
 576 Henry BA, Jenkins GP (1995) The impact of predation by the girded goby, *Nesogobius* sp. 1, on
 577 abundances of meiofauna and small macrofauna. J Exp Mar Biol Ecol 191: 223-238
 578 Hindley JPR, Alexander CG (1978) Structure and function of the chelate pereopods of the
 579 banana prawn *Penaeus merquiensis*. Mar Biol 48: 153-160
 580 Holling CS (1959) Some characteristics of simple types of predation and parasitism. Can
 581 Entomol 91: 385-398
 582 Hoyt M, Fleeger JW, Seibeling R, Feller RJ (2000) Serological estimation of prey-protein gut-
 583 residence time and quantification of meal size for grass shrimp consuming meiofaunal
 584 copepod. J Exp Mar Biol Ecol 248: 105-119
 585 Jonge VND, Bouwman L (1977) A simple density separation technique for quantitative isolation
 586 of meiobenthos using the colloidal silica Ludox TM. Mar Biol 42: 143-148
 587 Jourde J, Dupuy C, Nguyen HT, Mizrahi D, da Pracontal N, Bocher P (2017) Low benthic
 588 macrofauna diversity in dynamic tropical tidal mudflats: migrating banks on Guiana's
 589 coast, South America. Estuar Coast 40: 1159-1170
 590 Kanou K, Sano M, Kohno H (2005) Ontogenic diet shift, feeding rhythm, and daily ration of
 591 juvenile yellowfin goby *Acanthogobius flavimanus* on a tidal mudflat in the Tama River
 592 estuary, central Japan. Ichtyol Res 52: 319-324
 593 Kneib RT (1985) Predation and disturbance by grass shrimp, *Palaemonetes pugio*, Holthuis, in
 594 soft-substratum invertebrate benthic assemblages. J Exp Mar Biol Ecol 93: 91-102
 595 Kovačić M, la Mesa M (2008) Feeding ecology of De Buen's goby *Buenia affinis*, in the Kvarner
 596 area (Adriatic Sea). Vie Milieu 58: 249-256
 597 Krebs JR (1978) Optimal foraging: decision rules for predators. In: Krebs JR, Davies NB (eds)
 598 Behavioural ecology, an evolutionary approach. Blackwell, Oxford
 599 Leduc D, Probert PK, Duncan A (2009) A multi-method approach for identifying meiofaunal
 600 trophic connections. Mar Ecol Prog Ser 383: 95-111
 601 Legrand E, Martin S, Leroux C, Riera P (2018) Effect of temperature on an algal-grazer trophic
 602 transfer: a dual stable isotope (^{13}C , ^{15}N) labeling experiment. Mar Ecol 39: e12495
 603 Leroy F, Riera P, Jeanthon C, Edmond F, Leroux C, Comtet T (2012) Importance of bacterivory
 604 and preferential selection toward diatoms in larvae of *Crepidula fornicata* (L.) assessed by
 605 a dual stable isotope (^{13}C , ^{15}N) labeling approach. J Sea Res 70: 23-31

606 Li J, Vincx M, Herman PMJ (1996) A model of nematode dynamics in the Westerschelde
 607 Estuary. *Ecological Modelling* 90: 271-284
 608 Lipcius RN, Hines AH (1986) Variable functional responses of a marine predator in dissimilar
 609 homogeneous microhabitats. *Ecology* 67: 1361-1371
 610 Majdi N, Hette-Tronquart N, Auclair E, Bec A, Chouvelon T, Cognie B, Danger M, Decottignies
 611 P, Dessier A, Desvillettes C, Dubois S, Dupuy C, Fritsch C, Gauchere C, Hedde M, Jabot F,
 612 Lefebvre S, Marzloff MP, Pey B, Peyrard N, Powolny T, Sabbadin R, Thébault E, Perga ME
 613 (2018) There's no harm in having too much: a comprehensive toolbox of methods in
 614 trophic ecology. *Food Webs* 16: e00100
 615 McCall JN, Fleeger JW (1995) Predation by juvenile fish on hyperbenthic meiofauna: a review
 616 with data on post-larval *Leiosomus xanthurus*. *Vie Milieu* 45: 61-73
 617 Middelburg JJ, Barranguet C, Boschker HTS, Herman PMJ, Moens T, Heip CHR (2000) The fate
 618 of intertidal microphytobenthos carbon. An *in situ* ^{13}C labelling study. *Limnol Oceanogr*
 619 45: 1224-1234
 620 Moens T, Braeckman U, Derycke S, Fonseca G, Gallucci F, Gingold R, Guilini K, Ingels J, Leduc D,
 621 Vanaverbeke J, Van Colen C (2013) Ecology of free-living marine nematode. In: Schmidt-
 622 Rhaesa A (ed) *Handbook of zoology: gastrotricha, cycloneuralia and gnathifera*, Berlin, pp
 623 109-152
 624 Moens T, Luyten C, Middelburg JJ, Herman PMJ, Vincx M (2002) Tracing organic matter sources
 625 of estuarine tidal flat nematodes with stable carbon isotopes. *Mar Ecol Prog Ser* 234: 127-
 626 137
 627 Moodley L, Middelburg JJ, Boschker HTS, Duineveld R, Pel R, Herman PMJ, Heip CHR (2002)
 628 Bacteria and foraminifera: key players in a short-term deep sea benthic response to
 629 phytodetritus. *Mar Ecol Prog Ser* 236: 23-29
 630 Nguyen HT, Dupuy C, Jourde J, Lefrançois C, Pascal PY, Carpentier A, Chevalier J, Bocher P
 631 (2018) Persistent benthic communities in the extreme dynamic intertidal mudflats of
 632 Amazonian coast: An overview of the Tanaidacea (Crustacea, Peracarida). *Mar Biodiv* 48:
 633 1841-1853
 634 Nilsson P, Sundback K, Jonsson B (1993) Effect of the brown shrimp *Crangon crangon* on
 635 endobenthic macrofauna, meiofauna and meiofaunal grazing rates. *Neth J Sea Res* 31:
 636 95-106
 637 O'Gorman EJ, Enright RA, Emmerson MC (2008) Predator diversity enhances secondary
 638 production and decrease likelihood of trophic cascades. *Oecologia* 158: 557-567
 639 Ólafsson E (2003) Do macrofauna structure meiofauna assemblages in marine soft-bottoms?
 640 *Vie Milieu* 53: 249-265
 641 Ólafsson E, Elmgren R, Papakosta O (1993) Effects of the deposit-feeding benthic bivalve
 642 *Macoma balthica* on meiobenthos. *Oecologia* 93: 457-462
 643 Pascal PY, Dupuy C, Haubois AG, Richard P, Niquil N (2008a) Influence of environment factors

644 on bacterial ingestion rate of the deposit-feeder *Hydrobia ulvae* and comparison with
 645 meiofauna. J Sea Res 60: 151-156
 646 Pascal PY, Dupuy C, Mallet C, Richard P, Niquil N (2008b) Bacterivory by benthic organism in
 647 sediment: quantification using ¹⁵N-enriched bacteria. J Exp Mar Biol Ecol 355: 18-26
 648 Pascal PY, Dupuy C, Richard P, Mallet C, Armynot du Chatelet E, Niquil N (2009) Seasonal
 649 variation in consumption of benthic bacteria by meio- and macrofauna in an intertidal
 650 mudflat. Limnol Oceanogr 54: 1048-1059
 651 Pascal PY, Dupuy C, Richard P, Rzeznik-Orignac J, Niquil N (2008c) Bacterivory of a mudflat
 652 nematode community under different environmental conditions. Mar Biol 154: 671-682
 653 Pascal PY, Fleeger JW (2013) Diverse dietary responses by saltmarsh consumers to chronic
 654 nutrient enrichment. Estuar Coast 36: 1115-1124
 655 Peters RH (1983) The ecological implication of body size. Cambridge University Press
 656 Pihl L, Rosenberg R (1984) Food selection and consumption of the shrimp *Crangon crangon* in
 657 some shallow marine areas in western Sweden. Mar Ecol Prog Ser 15: 159-168
 658 Reise K (1978) Experiments on epibenthic predation in the Wadden Sea. Helgol Wiss
 659 Meeresunters 31: 55-101
 660 Ryer CH (1987) Temporal patterns of feeding by blue crabs (*Callinectes sapidus*) in a tidal-
 661 marsh creek and adjacent seagrass meadow in the lower Chesapeake Bay. Estuaries 10:
 662 136-140
 663 Scherer B, Reise K (1981) Significant predation on micro- and macrobenthos by the crab
 664 *Carcinus maenas* L. in the Wadden Sea Kieler Meeresforsch, pp 490-500
 665 Scholz DS, Matthews LL, Feller RJ (1991) Detecting selective digestion of meiobenthic prey in
 666 juvenile spot *Leiostomus xanthurus* (Pisces) using immunoassays. Mar Ecol Prog Ser 72:
 667 59-67
 668 Schratzberger M, Ingels J (2018) Meiofauna matters: the role of meiofauna in benthic
 669 ecosystems. J Exp Mar Biol Ecol 502: 12-25
 670 Schratzberger M, Warwick JT (1999) Differential effects of various type of disturbances on the
 671 structure of nematode assemblages: an experimental approach. Mar Ecol Prog Ser 181:
 672 227-236
 673 Schückel S, Sell AF, Kihara TC, Koeppen A, Kröncke I, Reiss H (2013) Meiofauna as food source
 674 for small-sized demersal fish in the southern North Sea. Helgol Mar Res 67: 203-218
 675 Schwinghamer P, Hargrave BT, Peer D, Hawkins CM (1986) Partitioning of production and
 676 respiration among size groups of organisms in an intertidal benthic community. Mar Ecol
 677 Prog Ser 31: 151-166
 678 Serrano A (2012) Changes in gut evacuation time for larval mud crab, *Scylla serrata* (Crustacea:
 679 Portunidae) fed artificial or live food. AACL Bioflux 5: 240-248
 680 Shaw M, Jenkins GP (1992) Spatial variation in feeding, prey distribution and food limitation of
 681 juvenile flounder *Rhombosolea tapirina* Günther. J Exp Mar Biol Ecol 165: 1-21

682 van Oevelen D, Moodley L, Soetaert K, Middelburg JJ (2006) The trophic significance of
683 bacterial carbon in a marine intertidal sediment: Results of an *in situ* stable isotope
684 labeling study. Limnol Oceanogr 51: 2349-2359

685

686

687