Letter to the Editor: 1 H, 15 N and 13 C chemical shift assignments of the Pleckstrin Homology domain of the Human Protein Kinase B (PKB/Akt)

Daniel Auguin, Philippe Barthe, Marie-Thérèse Augé-Sénégas, François Hoh, Masayuki Noguchi, Christian Roumestand

► To cite this version:

Daniel Auguin, Philippe Barthe, Marie-Thérèse Augé-Sénégas, François Hoh, Masayuki Noguchi, et al.. Letter to the Editor: 1 H, 15 N and 13 C chemical shift assignments of the Pleckstrin Homology domain of the Human Protein Kinase B (PKB/Akt). *Journal of Biomolecular NMR*, 2003, 27, pp.287 - 288. 10.1023/a:1025490109384 . hal-02359502

HAL Id: hal-02359502

<https://hal.science/hal-02359502>

Submitted on 14 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Letter to the Editor: ^1H , ^{15}N and ^{13}C chemical shift assignments of the Pleckstrin Homology domain of the Human Protein Kinase B (PKB/Akt)

Daniel Auguin^a, Philippe Barthe^a, Marie-Thérèse Augé-Sénégas^a, François Hoh, Masayuki Noguchi^b & Christian Roumestand^{a,*}

^aCentre de Biochimie Structurale, UMR 5548 CNRS/UMI – UMR 554 INSERM/UMI, Faculté de Pharmacie, BP14491, 15 Avenue Charles Flahault, 34093 Montpellier Cedex 5, France; ^bDivision of Cancer Biology, Institute for Genetic Medicine, Hokkaido University, N15 W7, Kita-ku, Sapporo 060-0815, Japan

Received 1 May 2003; Accepted 2 June 2003

Key words: Akt, apoptosis, NMR assignments, PH domain, PKB, proliferation

Biological context

Protein kinase (PKB), also named Akt (Staal, 1987), is a 57–60 kDa serine/threonine protein kinase that has emerged as a crucial regulator of widely divergent cellular processes including apoptosis, proliferation, differentiation and metabolism (Nicholson and Anderson, 2002). PKB is activated in a pathway induced by the lipid products of the phosphoinositide 3-kinase (PI3K) that bind to the N-terminal Pleckstrin Homology (PH) domain of the enzyme, thus recruiting the kinase to the plasma membrane where it is phosphorylated. It was previously shown that the PH domain of PKB interacts also with the TCL1/MTCP1 family of proto-oncogene (the proteins p14^{TCL1} and p13^{MTCP1}, respectively), mediating the formation of oligomeric TCL1/MTCP1-PKB high molecular weight protein complexes *in vivo* (Laine et al., 2000; Küntzle et al., 2002; French et al., 2002). Within these complexes, PKB is preferentially phosphorylated and activated. The structures of p14^{TCL1} (Hoh et al., 1998), of p13^{MTCP1} (Yang et al., 1998), and recently, the crystal structure of the isoform PKB α -PH (Thomas et al., 1998) have been determined. Here we report ^1H , ^{15}N and ^{13}C resonance assignments for PKB β -PH (73% homology) in the presence of Ins(1,4,5)P₃, a first step toward obtaining the solution structure as well as information on the interaction between PKB-PH and the TCL1/MTCP1 proto-oncogenes using chemical shift mapping studies.

*To whom correspondence should be addressed. E-mail: christian.roumestand@cbs.cnrs.fr

Methods and experiments

Protein expression and purification

The cDNA encoding for the 111 residues of the human PKB β /Akt2 PH domain was sub-cloned into the *EcoRI/BamHI* sites of the pGEX-2T plasmid (Amersham Biosciences). The construct was transformed for over-expression in *Escherichia coli* BL21 strain. Uniform ^{15}N and $^{15}\text{N}/^{13}\text{C}$ labelling was obtained by growing cells in ECPM1 medium containing $^{15}\text{NH}_4\text{Cl}$ and/or $^{13}\text{C}_6$ -glucose as the sole nitrogen and carbon sources. Protein expression was induced for 2 hours by addition of 0.5 mM IPTG. The cells were then harvested by centrifugation, and the pellets were sonicated in a lysis buffer. The supernatant was applied onto a Glutathione-Sepharose (Amersham Biosciences) column. After washing, the fusion protein was cleaved overnight at 4 °C by adding bovine thrombin (Sigma Aldrich) to the beads in the low salt buffer (50 mM Tris/HCl pH 7.4, 300 mM NaCl, 2.5 mM EDTA). The cleaved protein was further purified using size exclusion chromatography with an Sephadex-HR100 column (Amersham Biosciences).

NMR spectroscopy

NMR samples were 0.3–0.4 mM PKB β -PH in a buffer containing 10 mM Tris/HCl pH 7.4, 300 mM NaCl, 0.1 mM EDTA, 0.1 mM benzamidine and 4 mM Ins (1,4,5)P₃ as stabilizing reagent (5% D₂O for the lock). All NMR experiments were performed at 13 °C on Bruker AVANCE 500 MHz and 600 MHz equipped with 5 mm z-gradient ^1H - ^{13}C - ^{15}N . ^1H chem-

Figure 1. ^1H - ^{15}N HSQC spectrum of the PKB β PH domain recorded at 13°C. The assignment of peaks is indicated with their one-letter amino acid and number. Peaks belonging to the 'minor form' are indicated with an asterisk.

ical shifts were directly referenced to the resonance of cryogenic or standard probe, respectively DSS, while ^{13}C and ^{15}N chemical shifts were indirectly referenced. The following spectra were used for the ^1H , ^{15}N , $^{13}\text{C}\alpha$ and $^{13}\text{C}\beta$ and ^{13}CO resonance assignments: ^1H - ^{15}N HSQC, HNCA, HN(CO)CA, HNCACB, CBCA(CO)NH, HNCO, HN(CA)CO, essentially used for sequential assignments, and H(CC)H-TOCSY, ^{15}N - and/or ^{13}C -edited NOESY and TOCSY experiments, essentially used for side-chains assignments. All NMR experiments were processed with Gifa (Pons et al., 1996).

Extent of assignments and data deposition

The ^1H - ^{15}N HSQC spectrum of the PKB β PH domain is shown in Figure 1. About 85% of the expected backbone resonances, and 77% of the side-chain resonances have been assigned. Sequential assignment

failed for two segments (Glu⁵⁹-Gln⁶¹ and Arg⁷⁶-Thr⁸⁷: $\approx 15\%$ of the backbone resonances) linked by a disulfide bond (Cys⁶⁰-Cys⁷⁷), probably due to local conformational exchange on an intermediate timescale arising from the bridge isomerization. In addition, the residues belonging to the segment Pro⁴²-Pro⁵¹ exhibited 'minor form' peaks, ranging in intensity from 15–25% of the 'major form' peak. Potential sources for the minor peaks presumably involve proline isomerization states (4 proline residues in this segment), and modification of aromatic ring current effects in the close neighbourhood of this peptidic segment. The chemical shifts of the PKB β PH domain (major conformation) have been deposited in the BioMagRes-Bank under the accession number BMRB-5778. A full 3D structure determination is in progress.

Acknowledgements

This work was supported by a grant from the Association pour la Recherche sur le Cancer. The authors are specially grateful to M.P. Strub for her assistance in protein expression and labelling.

References

- French, S.W., Shen, R.R., Koh, P.J., Malone, C.S., Mallick, P. and Teitell, M.A. (2002) *Biochemistry*, **41**, 6376–6382.
- Hoh, F., Yang, Y.-S., Guignard, L., Padilla, A., Stern, M.H., Lhoste, J.M. and van Tilbeurgh, H. (1998) *Structure*, **6**, 147–155.
- Künstle, G., Laine, J., Pierron, G., Kagami, S., Nakajima, H., Hoh, F., Roumestand, C., Stern, M.-H. and Noguchi, M. (2002) *Mol. Cell. Biol.*, **22**, 1513–1525.
- Laine, J., Künstle, G., Obata, Y., Sha, M. and Noguchi, M. (2000) *Mol. Cell.*, **6**, 395–407.
- Nicholson, K.M. and Anderson, N.G. (2002) *Cell. Signal.*, **14**, 381–395.
- Pons, J.L., Malliavin, T.E. and Delsuc, M.A. (1996) *J. Biomol. NMR*, **8**, 445–452.
- Stall, S.P. (1987) *Proc. Natl. Acad. Sci. U.S.A.*, **84**, 5034–5037.
- Thomas, C.C., Deak, M., Alessi, D.R. and van Aalfen, D.M.F. (2002) *Curr. Biol.*, **12**, 1256–1262.
- Yang, Y.-S., Guignard, L., Padilla, A., Hoh, F., Strub, M.P., Stern, M.-H., Lhoste, J.M. and Roumestand, C. (1998) *J. Biomol. NMR*, **11**, 337–354.