On the Morita Reduced Versions of Skew Group Algebras of Path Algebras

Patrick Le Meur

► To cite this version:

Patrick Le Meur. On the Morita Reduced Versions of Skew Group Algebras of Path Algebras. Quarterly Journal of Mathematics, 2020, 71 (3), 10.1093/qmathj/haaa014 . hal-02361334v2

HAL Id: hal-02361334

<https://hal.science/hal-02361334v2>

Submitted on 16 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE MORITA REDUCED VERSIONS OF SKEW GROUP ALGEBRAS OF PATH ALGEBRAS

PATRICK LE MEUR

ABSTRACT. Let R be the skew group algebra of a finite group acting on the path algebra of a quiver. This article develops both theoretical and practical methods to do computations in the Morita reduced algebra associated to R . Reiten and Riedtmann proved that there exists an idempotent e of R such that the algebra eRe is both Morita equivalent to R and isomorphic to the path algebra of some quiver which was described by Demonet. This article gives explicit formulas for the decomposition of any element of eRe as a linear combination of paths in the quiver described by Demonet. This is done by expressing appropriate compositions and pairings in a suitable monoidal category which takes into account the representation theory of the finite group.

INTRODUCTION

In [13], Reiten and Riedtmann initiated the investigation of the skew group algebras of Artin algebras from the viewpoint of homological dimensions and representation theory. Their article was followed up by many research works illustrating the following principle: Artin algebras share many properties, whether of homological or of representation theoretic nature, with their associated skew group algebras. For instance, see [12] for interactions with Koszul duality; [4, 11] for interactions with piecewise hereditary algebras; and [14, 3] for interactions with preprojective algebras and McKay correspondence.

Let $\mathbb{k}$ be an algebraically closed field, Q be a finite quiver that may have oriented cycles, and G be a finite group with order not divisible by $\text{char}(\mathbb{k})$ and acting on $\mathbb{k}Q$ by algebra automorphisms in such a way that both the set of (primitive idempotents of) vertices and the vector subspace generated by the arrows are stabilised by this action. Following [13], the skew group algebra $\mathbb{k}Q * G$ is hereditary.

While Reiten and Riedtmann described a quiver whose path algebra is Morita equivalent to $\mathbb{k}Q * G$ when G is cyclic, the description of such a quiver in general is recent. In [3], Demonet described an idempotent $\tilde{e}$ of $\mathbb{k}Q * G$ and a quiver Q_G such that $\tilde{e} \cdot (\mathbb{k}Q * G) \cdot \tilde{e}$ is Morita equivalent to $\mathbb{k}Q * G$ and isomorphic to $\mathbb{k}Q_G$. This, however, does not yield a completely described isomorphism, which might be source of trouble. Here is a situation taken from Calabi-Yau algebras and where such trouble may occur. Following [10], if W is a G -invariant potential on Q , then G acts on the Ginzburg dg algebra $\mathcal{A}(Q, W)$, defined in [6], and $\mathcal{A}(Q, W) * G$ is Morita equivalent to $\mathcal{A}(Q_G, W_G)$, where W_G is the potential on Q_G which, as a linear combination of oriented cycles, corresponds to $\tilde{e} \cdot W \cdot \tilde{e}$ under any chosen isomorphism $\mathbb{k}Q_G \rightarrow \tilde{e} \cdot (\mathbb{k}Q * G) \cdot \tilde{e}$. The problem here is that describing that linear combination explicitly is not easy. Up to now, the only fairly general explicit descriptions of W_G are due to Amiot and Plamondon [1] when G is of order 2 and to Giovannini and Pasquali [8] when G is cyclic, the stabiliser of each vertex is either trivial or the whole group, and every cycle appearing in W goes through fixed

Date: June 30, 2020.

2010 Mathematics Subject Classification. Primary 16S35; Secondary 16W22, 20C15, 18D10.

Key words and phrases. Path algebra; skew group algebra; monoidal category; intertwiner.

vertices only (or, through vertices with trivial stabilisers only, respectively) as soon as it goes through two of them; during the revision of the present article, Giovannini, Pasquali and Plamondon presented [9] an extension of the latter description to the case where G is finite and abelian.

This raises the question of decomposing explicitly the elements of $\tilde{e} \cdot (\mathbb{k}Q * G) \cdot \tilde{e}$ as linear combinations of paths in Q_G under an isomorphism $\mathbb{k}Q_G \rightarrow \tilde{e} \cdot (\mathbb{k}Q * G) \cdot \tilde{e}$.

Demonet's description of Q_G involves the representation theory of the stabilisers of the vertices of Q . It is hence expectable that answering the above-mentioned question should involve representation theory. Answers of this kind already exist in particular cases where Q has only one vertex. Here is an example due to Ginzburg [6]. He proved that, if G is a finite subgroup of $\mathrm{SL}_3(\mathbb{C})$, then $\mathbb{C}[x, y, z] * G$ is Calabi-Yau in dimension 3. This was done by taking Q to be the quiver with one vertex and three loops x, y , and z , by taking $W = xyz - xzy$, and by expressing W_G in terms of the monoidal category of finite dimensional representations of G . See [2] for a generalisation to $\mathbb{C}[x_1, \dots, x_n] * G$ where G is a finite subgroup of $\mathrm{SL}_n(\mathbb{C})$, in which case Q is the quiver with one vertex and n loops. Note that this generalisation is (detailed and) successfully applied by Giovannini in [7] to find subgroups Γ of $\mathrm{SL}_{n+1}(\mathbb{C})$ such that $\mathbb{C}[x_1, \dots, x_{n+1}] * \Gamma$ is Morita equivalent to the $(n+1)$ -preprojective algebra of an n -representation finite algebra.

This article hence gives explicit formulas for the decomposition of the elements of $\tilde{e} \cdot (\mathbb{k}Q * G) \cdot \tilde{e}$ as linear combinations of paths in Q_G under the isomorphism $\mathbb{k}Q_G \rightarrow \tilde{e} \cdot (\mathbb{k}Q * G) \cdot \tilde{e}$. These formulas are obtained by translating explicitly the result of natural operations and pairings in the monoidal category $(\mathrm{mod}(A^e), \otimes_A)$ of finite dimensional A -bimodules, where A is the direct product of the group algebras of the stabilisers of the vertices of Q .

On one hand, these formulas provide an interpretation of the coefficients of these linear combinations in terms of the representation theory of the stabilisers of the vertices of Q . On the other hand, they are explicit enough to be implemented by a computer or a human being able to manipulate the irreducible representations of the involved groups. In specific cases, for instance when the action of the group elements transforms arrows to scalar multiples of arrows and the stabilisers are cyclic, these formulas simplify in a combinatorial way. Finally, these formulas give a complete solution to the problem of computing a potential W_G on Q_G such that the Ginzburg dg algebra $\mathcal{A}(Q_G, W_G)$ is Morita equivalent to $\mathcal{A}(Q, W) * G$, for any given G -invariant potential W on Q .

1. DEFINITIONS AND MAIN RESULTS

Throughout the article, $\mathbb{k}$ denotes an algebraically closed field. For all finite dimensional algebras Λ , their categories of finite dimensional left modules are denoted by $\mathrm{mod}(\Lambda)$.

This section presents the main results of this text as well as the needed definitions and notation. These definitions are illustrated with a running example throughout the section. A full application of the main results is presented on an example in section 2. The remaining sections are devoted to the proofs of these results.

Setting 1.1. Let Q_0 , S , G , and M be as follows.

- Q_0 is a finite set and S is the semi-simple $\mathbb{k}$ -algebra $\mathbb{k}Q_0$.
- G is a finite group with order not divisible by $\mathrm{char}(\mathbb{k})$ and acting on Q_0 .
- M is a finite dimensional S -bimodule, considered as a collection of vector spaces $({}_i M_j)_{(i,j) \in Q_0^2}$, endowed with an action of G written exponentially and such that ${}^g({}_i M_j) = {}_{g \cdot i} M_{g \cdot j}$ for all $i, j \in Q_0$ and $g \in G$.

The choice of a basis of ${}_iM_j$, for all $i, j \in Q_0$ determines a quiver Q such that $T_S(M) \simeq \mathbb{k}Q$ as $\mathbb{k}$ -algebras. All finite quivers arise in this way.

The actions of G on S and M induce an action on $T_S(M)$ by algebra automorphisms such that ${}^g(m_1 \otimes \cdots \otimes m_n) = {}^gm_1 \otimes \cdots \otimes {}^gm_n$ for all $g \in G$ and $m_1, \dots, m_n \in M$. Recall that the skew group algebra $T_S(M) * G$ is the $\mathbb{k}$ -algebra with underlying vector space $T_S(M) \otimes G$, where the tensor $u \otimes v$ is denoted by $u * v$ for all $u \in T_S(M)$ and $v \in \mathbb{k}G$, and with product such that $(u * g) \cdot (u' * g') = (u \cdot {}^gu') * gg'$ for all $u, u' \in T_S(M)$ and $g, g' \in G$. It is customary to consider $T_S(M)$ and $\mathbb{k}G$ as subalgebras of $T_S(M) * G$ in the canonical way.

Notation 1.2. The following data is used throughout the article.

- $[G \backslash Q_0]$ denotes a complete set of representatives of the orbits in Q_0 .
- For all $i \in Q_0$,
 - G_i denotes the stabiliser of i and e_i denotes the corresponding primitive idempotent of S ,
 - $[G/G_i]$ denotes a complete set of representatives of cosets modulo G_i ,
 - $\text{irr}(G_i)$ denotes a complete set of representatives of the isomorphism classes of the irreducible representations of G_i ,
 - and ε_U denotes a primitive idempotent of $\mathbb{k}G_i$ such that $U \simeq \mathbb{k}G_i \cdot \varepsilon_U$ in $\text{mod}(\mathbb{k}G_i)$, for all $U \in \text{irr}(G_i)$.
- $\tilde{e}$ denotes the idempotent $\sum_{(i,U)} e_i * \varepsilon_U$ of $T_S(M) * G$, where i runs through $[G \backslash Q_0]$ and U runs through $\text{irr}(G_i)$.

Example 1.3. Let Q be the quiver

where $\bar{0}$ and $\bar{1}$ denote the classes in $\mathbb{Z}/2\mathbb{Z}$. For all vertices i, j of Q , the arrow from i to j is denoted by $x_{i,j}$. Let G be the group $\langle \tau \mid \tau^2 \rangle$ of order 2. The actions of G on Q and on M are assumed to be the ones such that

$$\tau \cdot \bullet = \bullet, \quad \tau \cdot \bar{0} = \bar{1},$$

and, for all vertices i, j of Q ,

$${}^\tau x_{i,j} = -x_{\tau \cdot i, \tau \cdot j}.$$

- Let $[G \backslash Q_0]$ be $\{\bullet, \bar{0}\}$. Note that $G_\bullet = \{\text{Id}, \tau\}$ and $G_{\bar{0}} = \{\text{Id}\}$.
- Let $[G/G_\bullet]$ and $[G/G_{\bar{0}}]$ be $\{\text{Id}\}$ and $\{\text{Id}, \tau\}$, respectively.
- For all $k \in \mathbb{Z}/2\mathbb{Z}$, denote by ε_k the idempotent $\frac{1}{2}(\text{Id} + (-1)^k \tau)$; denote by ρ_k the irreducible representation $\mathbb{k} \cdot \varepsilon_k$ of G .
- Let $\text{irr}(G_\bullet)$ and $\text{irr}(G_{\bar{0}})$ be $\{\rho_k \mid k \in \mathbb{Z}/2\mathbb{Z}\}$ and $\{\mathbb{k} \cdot \text{Id}\}$, respectively.

In this situation, $\tilde{e} = e_\bullet * \varepsilon_{\bar{0}} + e_\bullet * \varepsilon_{\bar{1}} + e_{\bar{0}} * \text{Id}$.

1.1. Demonet's quiver of the Morita reduction. Demonet proved in [3] that the skew group algebra $T_S(M) * G$ is Morita equivalent to the path algebra of a quiver as follows.

Setting 1.4. Let Q_G be a quiver with the following properties.

- The vertices are the couples (i, U) , where $i \in [G \backslash Q_0]$ and $U \in \text{irr}(G_i)$.
- For all vertices (i, U) and (j, V) , the arrows $(i, U) \rightarrow (j, V)$ form a basis over $\mathbb{k}$ of $\text{Hom}_{\mathbb{k}G_i}(U, M(i, j; V))$, here $M(i, j; V) = \bigoplus_{y \in [G/G_j]} {}_iM_{y \cdot j} \otimes_{\mathbb{k}} yV$, where yV stands for $y \otimes_{\mathbb{k}} V \subset \mathbb{k}G \otimes_{\mathbb{k}} V$, see Definition 1.8 for details.

Example 1.5. *This is the continuation of Example 1.3. In order to describe a quiver Q_G such as in Setting 1.4, Table 1 describes the G_i -module $M(i, j; U)$ for all possible i, j, U .*

$M(i, j; U)$	Basis over $\mathbb{k}$	Action of G_i
$M(\bullet, \bullet; \rho_k)$	$x_{\bullet, \bullet} \otimes \varepsilon_k$	$\tau(x_{\bullet, \bullet} \otimes \varepsilon_k) = (-1)^{k+1} x_{\bullet, \bullet} \otimes \varepsilon_k$
$M(\bullet, \bar{0}; \mathbb{k} \cdot \text{Id})$	$(x_{\bullet, k} \otimes \tau^k)_{k \in \mathbb{Z}/2\mathbb{Z}}$	$\tau(x_{\bullet, k} \otimes \tau^k) = -x_{\bullet, k+\bar{1}} \otimes \tau^{k+\bar{1}}$
$M(\bar{0}, \bullet; \rho_k)$	$x_{\bar{0}, \bullet} \otimes \varepsilon_k$	Trivial
$M(\bar{0}, \bar{0}; \mathbb{k} \cdot \text{Id})$	$(x_{\bar{0}, k} \otimes \tau^k)_{k \in \mathbb{Z}/2\mathbb{Z}}$	Trivial

TABLE 1. The G_i -modules $M(i, j; U)$ ($k \in \mathbb{Z}/2\mathbb{Z}$).

For all $k \in \mathbb{Z}/2\mathbb{Z}$, denote by a_k and b_k the following respective morphisms of $G_\bullet$ -modules

$$\begin{aligned} \rho_{k+\bar{1}} &\longrightarrow M(\bullet, \bullet; \rho_k) & \rho_k &\longrightarrow M(\bullet, \bar{0}; \mathbb{k} \cdot \text{Id}) \\ \varepsilon_{k+\bar{1}} &\longmapsto x_{\bullet, \bullet} \otimes \varepsilon_k & \varepsilon_k &\longmapsto \frac{1}{2}(x_{\bullet, \bar{0}} \otimes \text{Id} + (-1)^{k+\bar{1}} x_{\bullet, \bar{1}} \otimes \tau) \end{aligned}$$

and denote by c_k and d_k the following respective morphisms of $G_{\bar{0}}$ -modules

$$\begin{aligned} \mathbb{k} \cdot \text{Id} &\longrightarrow M(\bar{0}, \bullet; \rho_k) & \mathbb{k} \cdot \text{Id} &\longrightarrow M(\bar{0}, \bar{0}; \mathbb{k} \cdot \text{Id}) \\ \text{Id} &\longmapsto x_{\bar{0}, \bullet} \otimes \varepsilon_k & \text{Id} &\longmapsto (-1)^k x_{\bar{0}, k} \otimes \tau^k. \end{aligned}$$

Then, Q_G may be taken equal to

$$\begin{array}{ccc} (\bullet, \rho_{\bar{0}}) & \begin{array}{c} \xleftarrow{b_{\bar{0}}} \\ \xrightarrow{c_{\bar{0}}} \end{array} & (0, \mathbb{k} \cdot \text{Id}), \\ \begin{array}{c} \uparrow a_{\bar{1}} \\ \downarrow a_{\bar{0}} \end{array} & & \begin{array}{c} \xleftarrow{c_{\bar{1}}} \\ \xrightarrow{b_{\bar{1}}} \end{array} \\ (\bullet, \rho_{\bar{1}}) & & \end{array}$$

(Note: The diagram shows a quiver with vertices $(\bullet, \rho_{\bar{0}})$, $(\bullet, \rho_{\bar{1}})$, and $(0, \mathbb{k} \cdot \text{Id})$. Arrows are labeled $a_{\bar{1}}, a_{\bar{0}}, b_{\bar{0}}, b_{\bar{1}}, c_{\bar{0}}, c_{\bar{1}}, d_{\bar{0}}, d_{\bar{1}}$. Specifically, $a_{\bar{1}}: (\bullet, \rho_{\bar{0}}) \rightarrow (\bullet, \rho_{\bar{1}})$, $a_{\bar{0}}: (\bullet, \rho_{\bar{1}}) \rightarrow (\bullet, \rho_{\bar{0}})$, $b_{\bar{0}}: (\bullet, \rho_{\bar{0}}) \rightarrow (0, \mathbb{k} \cdot \text{Id})$, $b_{\bar{1}}: (\bullet, \rho_{\bar{1}}) \rightarrow (0, \mathbb{k} \cdot \text{Id})$, $c_{\bar{0}}: (0, \mathbb{k} \cdot \text{Id}) \rightarrow (\bullet, \rho_{\bar{0}})$, $c_{\bar{1}}: (0, \mathbb{k} \cdot \text{Id}) \rightarrow (\bullet, \rho_{\bar{1}})$, $d_{\bar{0}}: (0, \mathbb{k} \cdot \text{Id}) \rightarrow (0, \mathbb{k} \cdot \text{Id})$, and $d_{\bar{1}}: (0, \mathbb{k} \cdot \text{Id}) \rightarrow (0, \mathbb{k} \cdot \text{Id})$.)

Demonet's result may be reformulated as follows, see Section 5 for more details.

Theorem 1.6 ([3]). *The $\mathbb{k}$ -algebras $\mathbb{k}Q_G$ and $T_S(M) * G$ are Morita equivalent. Besides, assuming that $U = \mathbb{k}G_i \cdot \varepsilon_U$ for all vertices (i, U) of Q_G , then there exists an isomorphism of algebras*

$$(1.1.1) \quad \mathbb{k}Q_G \xrightarrow{\sim} \tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$$

which maps every arrow $f: (i, U) \rightarrow (j, V)$ of Q_G to $f(\varepsilon_U)$.

1.2. Monoidal categories as a toolbox for intertwiners. The present article introduces a $\mathbb{k}$ -algebra, called the algebra of intertwiners relative to M and denoted by $\mathcal{I}ntw$, and an isomorphism of $\mathbb{k}$ -algebras $\mathbb{k}Q_G \rightarrow \mathcal{I}ntw^{\text{op}}$ together with explicit formulas decomposing any given element of $\mathcal{I}ntw$ in the basis consisting of the images of the paths in Q_G under $\mathbb{k}Q_G \rightarrow \mathcal{I}ntw^{\text{op}}$. See Theorem 1.25 for details. Composing this isomorphism with the inverse of (1.1.1) results in a $\mathbb{k}$ -algebra isomorphism $\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e} \rightarrow \mathcal{I}ntw^{\text{op}}$. A description of it is given in Theorem 1.27 hence providing a decomposition, under (1.1.1), of any element of $\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$ in the basis of $\mathbb{k}Q_G$ consisting of the paths in Q_G . These decompositions involve non-degenerate pairings between intertwiners relative to M and intertwiners relative to the dual vector space M^* . Both the product in $\mathcal{I}ntw$ and the pairings are explicit reformulations of natural operations and pairings in a suitable monoidal category.

Remark 1.7 ([5]). Let $(\mathcal{C}, \otimes)$ be a monoidal $\mathbb{k}$ -linear category with finite dimensional morphism spaces.

- (1) Given objects X, Y, U, V , and W of $\mathcal{C}$, and following the ideas presented in [6, Subsection 4.4] for the monoidal categories of finite subgroups of $SL_3(\mathbb{C})$, there is a natural operation between morphisms $U \rightarrow X \otimes V$ and $V \rightarrow Y \otimes W$,

$$\begin{aligned} \mathcal{C}(U, X \otimes V) \times \mathcal{C}(V, Y \otimes W) &\longrightarrow \mathcal{C}(U, X \otimes Y \otimes W) \\ (f, f') &\longmapsto (X \otimes f') \circ f. \end{aligned}$$

- (2) Given an object X of $\mathcal{C}$, recall that a *left dual* of X is an object X' of $\mathcal{C}$ together with an adjunction $(X \otimes -) \vdash (X' \otimes -)$ of endofunctors of $\mathcal{C}$ ([5, Definition 2.10.1, Proposition 2.10.8]). If X has a left dual, then there exists a non-degenerate pairing, for all objects U and V of $\mathcal{C}$ such that $\mathcal{C}(U, U) = \mathbb{k} \cdot \text{Id}_U$,

$$\begin{aligned} \langle - | - \rangle: \mathcal{C}(U, X \otimes V) \otimes \mathcal{C}(V, X' \otimes U) &\longrightarrow \mathbb{k} \\ f \otimes \phi &\longmapsto \langle f | \phi \rangle \end{aligned}$$

such that $\langle f | \phi \rangle \cdot \text{Id}_U = \bar{\phi} \circ f$, where $\bar{\phi} \in \mathcal{C}(X \otimes V, U)$ is adjoint to ϕ ; because $\mathcal{C}(U, U) = \mathbb{k} \cdot \text{Id}_U$, a scalar is associated to every endomorphism of U , and hence to $\bar{\phi} \circ f$, which makes it possible to define a pairing with values in $\mathbb{k}$. When $\mathcal{C}$ is a multitensor category this pairing is just a combination of the adjunction $(X \otimes -) \vdash (X' \otimes -)$ and the non-degenerate pairing $\mathcal{C}(U, X \otimes V) \otimes \mathcal{C}(X \otimes V, U) \rightarrow \mathbb{k}$ which takes into account U being simple in $\mathcal{C}$ (see [5, Example 7.9.14]).

The monoidal category $\mathcal{C}$ considered here is the category $\text{mod}(A^e)$ of finite dimensional A -bimodules with tensor product being $\otimes_A$, where A is as follows,

$$(1.2.1) \quad A = \mathbb{k} \times \prod_{i \in Q_0} \mathbb{k}G_i.$$

Note that A is a semi-simple $\mathbb{k}$ -algebra; accordingly, any object X of $\mathcal{C}$ has a left dual given by the dual vector space X^* . The isolated factor $\mathbb{k}$ in the definition of A is included so that $\mathcal{C}$ contains $\text{mod}(\mathbb{k}G_i)$ as a full additive subcategory, yet not as a monoidal subcategory, for all $i \in Q_0$. The reason for this inclusion is the following, at the same time it clarifies the role of $\mathcal{C}$. The text manipulates morphism spaces of the shape

$$(1.2.2) \quad \text{Hom}_{\mathbb{k}G_{i_0}}(U, e_{i_0}(M \otimes_{\mathbb{k}} \mathbb{k}G)e_{i_1} \otimes_{\mathbb{k}G_{i_1}} \cdots \otimes_{\mathbb{k}G_{i_{n-1}}} e_{i_{n-1}}(M \otimes_{\mathbb{k}} \mathbb{k}G)e_{i_n} \otimes_{\mathbb{k}G_{i_n}} V)$$

where $i_0, \dots, i_n$ are vertices of Q whereas $U \in \text{mod}(\mathbb{k}G_{i_0})$ and $V \in \text{mod}(\mathbb{k}G_{i_n})$; note that each one of $e_{i_{t-1}}(M \otimes_{\mathbb{k}} \mathbb{k}G)e_{i_t}$ (for $1 \leq t \leq n$) is an object of $\mathcal{C}$, and so are U and V because of the isolated factor $\mathbb{k}$ in A ; the above-mentioned morphism space is hence equal to

$$(1.2.3) \quad \mathcal{C}(U, e_{i_0}(M \otimes_{\mathbb{k}} \mathbb{k}G)e_{i_1} \otimes \cdots \otimes e_{i_{n-1}}(M \otimes_{\mathbb{k}} \mathbb{k}G)e_{i_n} \otimes V)$$

and it is manipulated in this text using the tools recalled in Remark 1.7.

1.3. The interwiners. Here are the base ingredients needed to define Intw .

Definition 1.8. Let n be a non-negative integer. Let $\underline{i} = i_0, \dots, i_n$ be a sequence in Q_0 . Let $V \in \text{mod}(\mathbb{k}G_{i_n})$.

- (1) For all sequences $\underline{y} = y_1, \dots, y_n$, where $y_t \in [G/G_{i_t}]$ for all $t \in \{1, \dots, n\}$, let $M_{\underline{y}}(\underline{i}; V)$ be the following vector space,

$$M_{\underline{y}}(\underline{i}; V) = {}_{i_0}M_{y_1 \cdot i_1} \otimes_{\mathbb{k}} {}_{y_1 \cdot i_1}M_{y_1 y_2 \cdot i_2} \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} {}_{y_1 \cdots y_{n-1} \cdot i_{n-1}}M_{y_1 \cdots y_n \cdot i_n} \otimes_{\mathbb{k}} y_1 \cdots y_n V,$$

where $y_1 \cdots y_n V$ stands for the vector subspace $y_1 \cdots y_n \otimes_{\mathbb{k}} V$ of $\mathbb{k}G \otimes_{\mathbb{k}} V$.

(2) Define the $\mathbb{k}G_{i_0}$ -module $M(\underline{i}; V)$ as follows. Its underlying vector space is

$$(1.3.1) \quad \bigoplus_{\underline{y}} M_{\underline{y}}(\underline{i}; V),$$

where $\underline{y}$ runs through all possible sequences such as in (1). For all $g \in G_{i_0}$, the action of g is written exponentially and defined as follows. For all $\underline{y}$, there exist a unique sequence $\underline{y}' = y'_1, \dots, y'_n$, where $y'_t \in [G/G_{i_t}]$ for all $t \in \{1, \dots, n\}$, and a unique $h \in G_{i_n}$, such that

- $gy_1 \cdots y_t G_{i_t} = y'_1 \cdots y'_t G_{i_t}$ for all $t \in \{1, \dots, n\}$,
- and $gy_1 \cdots y_n = y'_1 \cdots y'_n h$;

then, the action of g transforms $M_{\underline{y}}(\underline{i}; V)$ into $M_{\underline{y}'}(\underline{i}; V)$ and, for all $m_1 \otimes \cdots \otimes m_n \otimes y_1 \cdots y_n v \in M_{\underline{y}}(\underline{i}; V)$,

$${}^g(m_1 \otimes \cdots \otimes m_n \otimes y_1 \cdots y_n v) = {}^g m_1 \otimes \cdots \otimes {}^g m_n \otimes y'_1 \cdots y'_n {}^h v.$$

This does define a finite dimensional $\mathbb{k}G_{i_0}$ -module. If $n = 0$, then $M(\underline{i}; V) = V$. In the monoidal category $(\text{mod}(A^e), \otimes_A)$ considered previously, $M(\underline{i}; V)$ is isomorphic to $M(\underline{i}) \otimes_A V$ where $M(\underline{i})$ is a $\mathbb{k}G_{i_0} - \mathbb{k}G_{i_n}$ -bimodule defined by M and $\underline{i}$, see (4.1.1) for the definition of $M(\underline{i})$ and (4.1.7) for the isomorphism.

Note that, in the previous definition, if $U = \mathbb{k}G_{i_0} \cdot \varepsilon_U$ and $V = \mathbb{k}G_{i_n} \cdot \varepsilon_V$ for primitive idempotents $\varepsilon_U \in \mathbb{k}G_{i_0}$ and $\varepsilon_V \in \mathbb{k}G_{i_n}$, then $M(\underline{i}; U)$ is a $\mathbb{k}G_{i_0}$ -submodule of $T_S(M) * G$.

Example 1.9. *This is the continuation of Example 1.5. Let $k \in \mathbb{Z}/2\mathbb{Z}$. Then $M(\bullet, \bar{0}, \bullet; \rho_k)$ identifies with the $\mathbb{k}G_\bullet$ -submodule of $T_S(M) * G$ whose underlying vector space is generated by $(x_{\bullet, a} x_{a, b} x_{b, \bullet} * \varepsilon_k)_{a, b \in \mathbb{Z}/2\mathbb{Z}}$. On this submodule, the action of τ is such that ${}^\tau(x_{\bullet, a} x_{a, b} x_{b, \bullet} * \varepsilon_k) = (-1)^{k+\bar{1}} x_{\bullet, a+\bar{1}} x_{a+\bar{1}, b+\bar{1}} x_{b+\bar{1}, \bullet} * \varepsilon_k$.*

The main results of this article are based on a description of the quiver $\mathbb{k}Q_G$ in terms of intertwiners.

Definition 1.10. An *intertwiner* relative to M is a morphism of $\mathbb{k}G_{i_0}$ -modules $U \rightarrow M(\underline{i}; V)$, for some sequence $\underline{i} = i_0, \dots, i_n$ in Q_0 and some modules $U \in \text{mod}(\mathbb{k}G_{i_0})$ and $V \in \text{mod}(\mathbb{k}G_{i_n})$.

Note that, in the previous definition, if $U = \mathbb{k}G_{i_0} \cdot \varepsilon_U$ and $V = \mathbb{k}G_{i_n} \cdot \varepsilon_V$ for primitive idempotents $\varepsilon_U \in \mathbb{k}G_{i_0}$ and $\varepsilon_V \in \mathbb{k}G_{i_n}$, then all intertwiners $U \rightarrow M(\underline{i}; V)$ take their values in $T_S(M) * G$.

Example 1.11. *This is the continuation of Example 1.9. Let $s, t \in \mathbb{Z}/2\mathbb{Z}$. For all $a, b \in \mathbb{Z}/2\mathbb{Z}$, the following holds true in $M(\bullet, \bar{0}, \bar{0}, \bullet; \rho_t)$,*

$$\varepsilon_s \cdot (x_{\bullet, a} x_{a, b} x_{b, \bullet} \otimes \varepsilon_t) = \frac{1}{2} (x_{\bullet, a} x_{a, b} x_{b, \bullet} \otimes \varepsilon_t + (-1)^{s+\bar{1}+t} x_{\bullet, a+\bar{1}} x_{a+\bar{1}, b+\bar{1}} x_{b+\bar{1}, \bullet} \otimes \varepsilon_t).$$

Thus, $\text{Hom}_{\mathbb{k}G_\bullet}(\rho_s, M(\bullet, \bar{0}, \bar{0}, \bullet; \rho_t))$ is two-dimensional with basis $(F_a)_{a \in \mathbb{Z}/2\mathbb{Z}}$, where F_a denotes the following intertwiner relative to M , for all $a \in \mathbb{Z}/2\mathbb{Z}$,

$$\begin{aligned} \rho_s &\longrightarrow M(\bullet, \bar{0}, \bar{0}, \bullet; \rho_t) \\ \varepsilon_s &\longmapsto \frac{1}{2} (x_{\bullet, \bar{0}} x_{\bar{0}, a} x_{a, \bullet} \otimes \varepsilon_t + (-1)^{s+\bar{1}+t} x_{\bullet, \bar{1}} x_{\bar{1}, a+\bar{1}} x_{a+\bar{1}, \bullet} \otimes \varepsilon_t). \end{aligned}$$

The main purpose of this article is to provide explicit formulas in the main theorems. These formulas use the following notation.

Notation 1.12. Let $\underline{i} = i_0, \dots, i_n$ be a sequence in Q_0 where $n \geq 1$. Let $U \in \text{mod}(\mathbb{k}G_{i_0})$ and $V \in \text{mod}(\mathbb{k}G_{i_n})$. Let $f \in \text{Hom}_{\mathbb{k}G_{i_0}}(U, M(\underline{i}; V))$. For all $u \in U$, write the decomposition of $f(u)$ along (1.3.1) as follows

$$f(u) = \sum_{\underline{y}} f_{\underline{y}}(u);$$

and, for all sequences $\underline{y} = y_1, \dots, y_n$, where $y_t \in [G/G_{i_t}]$ for all $t \in \{1, \dots, n\}$, write symbolically the element $f_{\underline{y}}(u)$ of $M_{\underline{y}}(\underline{i}; V)$ as follows, with the sum sign omitted,

$$f_{\underline{y}}(u) = f_{\underline{y}}^{(1)}(u) \otimes \dots \otimes f_{\underline{y}}^{(n)}(u) \otimes y_1 \dots y_n f_{\underline{y}}^{(0)}(u);$$

hence, $f_{\underline{y}}^{(t)}(u)$ is meant to lie in $_{y_1 \dots y_{t-1} \cdot i_{t-1}} M_{y_1 \dots y_t \cdot i_t}$, for all $t \in \{1, \dots, n\}$, and $f_{\underline{y}}^{(0)}(u)$ is meant to lie in V ; as a whole,

$$(1.3.2) \quad f(u) = \sum_{\underline{y}} f_{\underline{y}}^{(1)}(u) \otimes \dots \otimes f_{\underline{y}}^{(n)}(u) \otimes y_1 \dots y_n f_{\underline{y}}^{(0)}(u).$$

1.4. The algebra of intertwiners. Now here is the operation to be used as the multiplication in $\mathcal{Z}ntw$.

Definition 1.13. Let $\underline{i}'' = i_0, \dots, i_{m+n}$ be a sequence in Q_0 , where $m, n \geq 1$. Denote the sequences $i_0, \dots, i_m$ and $i_m, i_{m+1}, \dots, i_{m+n}$ by $\underline{i}$ and $\underline{i}'$, respectively. Let $U \in \text{mod}(\mathbb{k}G_{i_0})$, $V \in \text{mod}(\mathbb{k}G_{i_m})$, and $W \in \text{mod}(\mathbb{k}G_{i_{m+n}})$. For all $f \in \text{Hom}_{\mathbb{k}G_{i_0}}(U, M(\underline{i}; V))$ and $f' \in \text{Hom}_{\mathbb{k}G_{i_m}}(V, M(\underline{i}'; W))$, define a $\mathbb{k}$ -linear mapping

$$f' \circledast f: U \rightarrow M(\underline{i}''; W)$$

by

$$(1.4.1) \quad f' \circledast f = \sum_{\underline{y}''} (f' \circledast f)_{\underline{y}''},$$

where

- $\underline{y}''$ runs through all sequences $y_1, \dots, y_{m+n}$ with $y_t \in [G/G_{i_t}]$ for all t
- and $(f' \circledast f)_{\underline{y}''}$ is the mapping $U \rightarrow M_{\underline{y}''}(\underline{i}''; W)$ such that, for all $u \in U$,

$$\begin{aligned} (f' \circledast f)_{\underline{y}''}(u) &= f_{\underline{y}}^{(1)}(u) \otimes \dots \otimes f_{\underline{y}}^{(m)}(u) \otimes \\ &\quad y_1 \dots y_m (f_{\underline{y}'}^{(1)}(f_{\underline{y}}^{(0)}(u))) \otimes \dots \otimes y_1 \dots y_m (f_{\underline{y}'}^{(n)}(f_{\underline{y}}^{(0)}(u))) \otimes \\ &\quad y_1 \dots y_{m+n} f_{\underline{y}'}^{(0)}(f_{\underline{y}}^{(0)}(u)), \end{aligned}$$

where $\underline{y} = y_1, \dots, y_m$ and $\underline{y}' = y_{m+1}, \dots, y_{m+n}$.

Note that, in the previous definition, if $U = \mathbb{k}G_{i_0} \cdot \varepsilon_U$, $V = \mathbb{k}G_{i_m} \cdot \varepsilon_V$, and $W = \mathbb{k}G_{i_{m+n}} \cdot \varepsilon_W$ for primitive idempotents $\varepsilon_U \in \mathbb{k}G_{i_0}$, $\varepsilon_V \in \mathbb{k}G_{i_m}$, and $\varepsilon_W \in \mathbb{k}G_{i_{m+n}}$, then

$$(1.4.2) \quad (f' \circledast f)(\varepsilon_U) = f(\varepsilon_U) \cdot f'(\varepsilon_V),$$

where the product on the right-hand side is taken in $T_S(M) * G$.

Remark 1.14. The definition of $\circledast$ is technical. Actually, it is an explicit reformulation of a simple operation in the monoidal category $(\text{mod}(A^e), \otimes_A)$. More precisely, using the comment that follows Definition 1.8,

- first, $M(\underline{i}'') \simeq M(\underline{i}) \otimes_A M(\underline{i}')$, see Lemma 4.2;
- next, if $M(\underline{i}; V)$, $M(\underline{i}'; W)$, and $M(\underline{i}''; W)$ are identified with $M(\underline{i}) \otimes_A V$, $M(\underline{i}') \otimes_A W$, and $M(\underline{i}'') \otimes_A W$, respectively, then $\circledast$ is an explicit reformulation of the operation introduced in part (1) of Remark 1.7, see Lemma 4.4 for a precise statement and proof.

In view of the previous remark, $\circledast$ does yield intertwiners, see Lemma 4.4, and it is associative, see Proposition 4.5.

Definition 1.15. The *algebra of intertwiners* relative to M is denoted by $\mathcal{I}ntw$ and defined by

$$(1.4.3) \quad \mathcal{I}ntw = \left(\bigoplus_{n, \underline{i}, U, V} \text{Hom}_{\mathbb{k}G_{i_0}}(U, M(\underline{i}; V)), \otimes \right),$$

where n runs through all non negative integers, $\underline{i} = i_0, \dots, i_n$ runs through all sequences in $[G \setminus Q_0]$, and U and V run through $\text{irr}(G_{i_0})$ and $\text{irr}(G_{i_n})$, respectively.

It is hence possible to associate an element of $\mathcal{I}ntw$ to every path in Q_G .

Notation 1.16. For all paths in Q_G

$$\gamma: (i_0, U_0) \xrightarrow{f_1} (i_1, U_1) \rightarrow \dots \rightarrow (i_{n-1}, U_{n-1}) \xrightarrow{f_n} (i_n, U_n),$$

denote by f_γ the following element of $\mathcal{I}ntw$ lying in $\text{Hom}_{\mathbb{k}G_{i_0}}(U_0, M(i_0, \dots, i_n; U_n))$,

$$(1.4.4) \quad f_\gamma = f_n \otimes f_{n-1} \otimes \dots \otimes f_1.$$

1.5. Dual intertwiners and their pairing with intertwiners. As stated below, see Theorem 1.25, assigning f_γ to γ for every path γ in Q_G yields an isomorphism of algebras from $\mathbb{k}Q_G$ to $\mathcal{I}ntw^{\text{op}}$. The purpose of this article is to explain how to decompose any element of $\mathcal{I}ntw$ as a linear combination of the f_γ 's. This involves non-degenerate pairings between certain spaces of intertwiners relative to M and to M^* , respectively. Here, M^* denotes the dual vector space $\text{Hom}_{\mathbb{k}}(M, \mathbb{k})$, it has a structure of S -bimodule such that ${}_i M_j^* = ({}_j M_i)^*$ for all $i, j \in Q_0$, and G acts on M^* in such a way that, for all $\varphi \in M^*$ and $g \in G$,

$${}^g \varphi = \varphi({}^{g^{-1}} \bullet).$$

This action is such that ${}^g({}_i M_j^*) = {}_{g \cdot i} M_{g \cdot j}^*$ for all $i, j \in Q_0$ and $g \in G$. Hence, the previous considerations may be applied to M^* instead of to M .

Example 1.17. This is the continuation of Example 1.11. For all vertices i, j of Q , the vector space ${}_i M_j^*$ is one-dimensional; denote by $x'_{i,j}$ its base element, dual to the base element $x_{j,i}$ of ${}_j M_i$; Hence, ${}^\tau x'_{i,j} = -x'_{\tau(i), \tau(j)}$. After replacing M by M^* , the considerations of Example 1.11 yield that $\text{Hom}_{\mathbb{k}G_\bullet}(\rho_t, M^*(\bullet, \bar{0}, \bar{0}, \bullet; \rho_s))$ is two-dimensional with basis $(\phi_a)_{a \in \mathbb{Z}/2\mathbb{Z}}$, where ϕ_a denotes the following intertwiner relative to M^* , for all $a \in \mathbb{Z}/2\mathbb{Z}$,

$$\begin{aligned} \rho_t &\longrightarrow M^*(\bullet, \bar{0}, \bar{0}, \bullet; \rho_s) \\ \varepsilon_t &\longmapsto x_{\bullet, \bar{0}} x_{\bar{0}, a} x_{a, \bullet} \otimes \varepsilon_s + (-1)^{t+\bar{1}+s} x_{\bullet, \bar{1}} x_{\bar{1}, a+\bar{1}} x_{a+\bar{1}, \bullet} \otimes \varepsilon_s. \end{aligned}$$

Notation 1.18. Let $\underline{i} = i_0, \dots, i_n$ be a sequence in Q_0 , where $n \geq 0$. Let $U \in \text{mod}(\mathbb{k}G_{i_0})$ and $V \in \text{mod}(\mathbb{k}G_{i_n})$.

- (1) Denote by $\underline{i}^o$ the sequence $i_n, i_{n-1}, \dots, i_0$.
- (2) Let $\varphi \in \text{Hom}_{\mathbb{k}G_{i_n}}(V, M^*(\underline{i}^o, U))$. Proceeding similarly as done in Notation 1.12,
 - denote by $\varphi^{\underline{x}}$ the composition of φ with the canonical projection $M^*(\underline{i}^o; U) \rightarrow M_{\underline{x}}^*(\underline{i}^o; U)$, for all sequences $\underline{x} = x_{n-1}, x_{n-2}, \dots, x_0$ where $x_t \in [G/G_{i_t}]$ for all $t \in \{0, \dots, n-1\}$,
 - and for all such $\underline{x}$ and all $v \in V$, write symbolically the element $\varphi^{\underline{x}}(v)$ of $M_{\underline{x}}^*(\underline{i}^o; U)$ as follows,

$$\varphi^{\underline{x}}(v) = \varphi_{(n)}^{\underline{x}}(v) \otimes \dots \otimes \varphi_{(1)}^{\underline{x}}(v) \otimes x_{n-1} \dots x_0 \varphi_{(0)}^{\underline{x}}(v).$$

Hence, $\varphi_{(t)}^{\underline{x}}(v)$ is meant to lie in ${}_{(x_{n-1}x_{n-2}\cdots x_t)\cdot i_t}M_{(x_{n-1}x_{n-2}\cdots x_{t-1})\cdot i_{t-1}}^*$ for all $t \in \{1, \dots, n\}$ and $\varphi_{(0)}^{\underline{x}}(v)$ is meant to lie in U . As a whole,

$$(1.5.1) \quad \varphi(v) = \sum_{\underline{x}} \varphi_{(n)}^{\underline{x}}(v) \otimes \cdots \otimes \varphi_{(1)}^{\underline{x}}(v) \otimes x_{n-1} \cdots x_1 \varphi_{(0)}^{\underline{x}}(v).$$

Here is an illustration of this notation.

Example 1.19. *This is the continuation of Example 1.17. Assume that $n = 2$ and $\underline{i} = \bar{0}, \bar{0}, \bullet$. Hence $\underline{i}^o = \bullet, \bar{0}, \bar{0}$. The underlying vector space of the $\mathbb{k}G_{\bullet}$ -module $M^*(\underline{i}^o; \mathbb{k} \cdot \text{Id})$ is freely generated by $(x'_{\bullet, a} x'_{a, b} \otimes \tau^b)_{a, b \in \mathbb{Z}/2\mathbb{Z}}$.*

Let $a, b, s \in \mathbb{Z}/2\mathbb{Z}$. Denote by φ the intertwiner $\rho_s \rightarrow M^(\underline{i}^o; \mathbb{k} \cdot \text{Id})$ such that $\varphi(\varepsilon_s) = \varepsilon_s \cdot (x'_{\bullet, a} x'_{a, b} \otimes \tau^b)$, that is $\varphi(\varepsilon_s) = \frac{1}{2} x'_{\bullet, a} x'_{a, b} \otimes \tau^b + \frac{(-1)^s}{2} x'_{\bullet, a+\bar{1}} x'_{a+\bar{1}, b+\bar{1}} \otimes \tau^{b+\bar{1}}$. On one hand, $x'_{\bullet, a} x'_{a, b} = x'_{\bullet, \tau^a \cdot \bar{0}} x'_{\tau^a \cdot \bar{0}, \tau^a \tau^{b-a} \cdot \bar{0}}$; Hence*

$$\begin{cases} \varphi_{(2)}^{\tau^a, \tau^{b-a}}(\varepsilon_s) \otimes \varphi_{(1)}^{\tau^a, \tau^{b-a}}(\varepsilon_s) \otimes \varphi_{(0)}^{\tau^a, \tau^{b-a}}(\varepsilon_s) &= \frac{1}{2} x'_{\bullet, a} x'_{a, b} \otimes \tau^b \\ \varphi_{(2)}^{\tau^a, \tau^{b-a}}(\varepsilon_s) \otimes \varphi_{(1)}^{\tau^a, \tau^{b-a}}(\varepsilon_s) \otimes \varphi_{(0)}^{\tau^a, \tau^{b-a}}(\varepsilon_s) &= \frac{1}{2} x'_{\bullet, a} \otimes x'_{a, b} \otimes \text{Id}. \end{cases}$$

On the other hand, $x'_{\bullet, a+\bar{1}} x'_{a+\bar{1}, b+\bar{1}} = x'_{\bullet, \tau^{a+\bar{1}} \cdot \bar{0}} x'_{\tau^{a+\bar{1}} \cdot \bar{0}, \tau^{a+\bar{1}} \tau^{b-a} \cdot \bar{0}}$; Hence

$$\begin{cases} \varphi_{(2)}^{\tau^{a+\bar{1}}, \tau^{b-a}}(\varepsilon_s) \otimes \varphi_{(1)}^{\tau^{a+\bar{1}}, \tau^{b-a}}(\varepsilon_s) \otimes \varphi_{(0)}^{\tau^{a+\bar{1}}, \tau^{b-a}}(\varepsilon_s) &= \frac{(-1)^s}{2} x'_{\bullet, a+\bar{1}} x'_{a+\bar{1}, b+\bar{1}} \otimes \tau^{b+\bar{1}} \\ \varphi_{(2)}^{\tau^{a+\bar{1}}, \tau^{b-a}}(\varepsilon_s) \otimes \varphi_{(1)}^{\tau^{a+\bar{1}}, \tau^{b-a}}(\varepsilon_s) \otimes \varphi_{(0)}^{\tau^{a+\bar{1}}, \tau^{b-a}}(\varepsilon_s) &= \frac{(-1)^s}{2} x'_{\bullet, a+\bar{1}} \otimes x'_{a+\bar{1}, b+\bar{1}} \otimes \text{Id}. \end{cases}$$

In view of description of $\varphi(\varepsilon_s)$ made initially, all the remaining terms of the sum of (1.5.1) are zero.

The above mentioned pairings are described explicitly as follows.

Proposition 1.20. *Let $\underline{i} = i_0, \dots, i_n$ be a sequence in Q_0 , where $n \geq 0$. Let $U \in \text{mod}(\mathbb{k}G_{i_0})$ and $V \in \text{mod}(\mathbb{k}G_{i_n})$. Assume that U is simple. There exists a non-degenerate pairing*

$$(-| -): \text{Hom}_{\mathbb{k}G_{i_0}}(U, M(\underline{i}; V)) \otimes_{\mathbb{k}} \text{Hom}_{\mathbb{k}G_{i_n}}(V, M^*(\underline{i}^o; U)) \rightarrow \mathbb{k}$$

with the following property. For all $f \in \text{Hom}_{\mathbb{k}G_{i_0}}(U, M(\underline{i}; V))$ and for all $\varphi \in \text{Hom}_{\mathbb{k}G_{i_n}}(V, M^(\underline{i}^o; U))$, the scalar $(f|\varphi)$ is such that, for all $u \in U$, the element $(f|\varphi)u$ of U is equal to*

$$(1.5.2) \quad \sum_{\underline{y}} \prod_{t=1}^n \varphi_{(t)}^{\underline{x}}(f_{\underline{y}}^{(0)}(u)) ({}^{(y_1 \cdots y_n)^{-1}} f_{\underline{y}}^{(t)}(u)) {}^{h_0} (\varphi_{(0)}^{\underline{x}}(f_{\underline{y}}^{(0)}(u))),$$

where $\underline{y} = y_1, \dots, y_n$ runs through all sequences with $y_t \in [G/G_{i_t}]$ for all t and $\underline{x} = x_{n-1}, \dots, x_0$ and h_0 are uniquely determined by the following conditions:

- (a) $x_t \in [G/G_{i_t}]$ for all t , and $h_0 \in G_{i_0}$.
- (b) $(y_t \cdots y_n)^{-1} \in x_{n-1}x_{n-2} \cdots x_{t-1}G_{i_{t-1}}$ for all $t \in \{1, \dots, n\}$.
- (c) $(y_1 \cdots y_n)^{-1} = x_{n-1}x_{n-2} \cdots x_0 h_0$.

In view of computations, here is a useful explanation regarding $\underline{x}$ and h_0 in the above proposition. For all sequences $\underline{y} = y_1, \dots, y_n$ in G such that $y_t \in [G/G_{i_t}]$ for all $t \in \{1, \dots, n\}$ and for all non zero tensors

$$m_1 \otimes \cdots \otimes m_n \in {}_{i_0}M_{y_1 \cdot i_1} \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} {}_{(y_1 \cdots y_{n-1}) \cdot i_{n-1}}M_{(y_1 \cdots y_n) \cdot i_n},$$

there exists a unique sequence $\underline{x} = x_{n-1}, x_{n-2}, \dots, x_0$ in G such that $x_t \in [G/G_{i_t}]$ for all $t \in \{0, \dots, n-1\}$ and

$${}^{(y_1 \cdots y_n)^{-1}} (m_1 \otimes \cdots \otimes m_n) \in {}_{(x_{n-1}x_{n-2} \cdots x_0) \cdot i_0}M_{(x_{n-1} \cdots x_1) \cdot i_1} \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} {}_{x_{n-1} \cdot i_{n-1}}M_{i_n};$$

besides, $(x_{n-1}x_{n-2} \cdots x_0)^{-1}(y_1 \cdots y_n)^{-1} \in G_{i_0}$. This sequence $\underline{x}$ is the one in the statement of Proposition 1.20 in which $h_0 = (x_{n-1}x_{n-2} \cdots x_0)^{-1}(y_1 \cdots y_n)^{-1}$.

When the action of G on M is such that arrows of Q are transformed into scalar multiples of arrows of Q , the expression (1.5.2) of $(-|-)$ simplifies to (2.1.3). The following example illustrates the computation of $(-|-)$ in this particular situation.

Example 1.21. *This is the continuation of Example 1.17. Let $s, t \in \mathbb{Z}/2\mathbb{Z}$. Let $a, b \in \mathbb{Z}/2\mathbb{Z}$. Consider $F_a: \rho_s \rightarrow M(\bullet, \bar{0}, \bar{0}, \bullet; \rho_t)$ and $\phi_b: \rho_t \rightarrow M(\bullet, \bar{0}, \bar{0}, \bullet; \rho_s)$ such as in Examples 1.11 and 1.17, respectively.*

In view of (2.1.3), $(F_a|\phi_b)$ is a sum of products of scalars, the sum is indexed by the paths in Q that appear in the expression of $F_a(\varepsilon_s)$. For all such paths γ , the scalars forming the corresponding product appear in Table 2 with bold face. For convenience, this table presents additional data, using the following notation, see Subsection 2.1 for more details. For a given path γ in Q :

- α_γ denotes the coefficient of $\gamma \otimes \varepsilon_t$ in $F_a(\varepsilon_s)$;
- y_1, y_2, y_3 is the sequence of G such that $y_1, y_2 \in [G/G_0]$, $y_3 \in [G/G_\bullet]$ and γ is a path of the shape $\bullet \rightarrow y_1 \cdot \bar{0} \rightarrow y_1 y_2 \cdot \bar{0} \rightarrow y_1 y_2 y_3 \cdot \bullet$;
- $(y_1 y_2 y_3)^{-1}(\gamma)$ denotes the path of Q and $\chi_{(y_1 y_2 y_3)^{-1}, \gamma}$ denotes the scalar such that $(y_1 y_2 y_3)^{-1} \gamma = \chi_{(y_1 y_2 y_3)^{-1}, \gamma} \cdot (y_1 y_2 y_3)^{-1}(\gamma)$;
- $(y_1 y_2 y_3)^{-1}(\gamma)^*$ denotes the element of the basis of $T_S(M^*)$ dual to the basis of $T_S(M)$ consisting of the paths of Q , associated to $(y_1 y_2 y_3)^{-1}(\gamma)$;
- $\beta_{(y_1 y_2 y_3)^{-1}(\gamma)^*}$ denotes the coefficient of $(y_1 y_2 y_3)^{-1}(\gamma)^*$ in $\phi_b(\varepsilon_t)$;
- χ_{ρ_s} denotes the character of ρ_s .

Thus,

$$(1.5.3) \quad (F_a|\phi_b) = \begin{cases} \frac{(-1)^{(s+\bar{1})a \cdot (1+(-1)^t)}}{2} & \text{if } a = b \\ 0 & \text{if not.} \end{cases}$$

γ	$x_{\bullet, \bar{0}} x_{\bar{0}, a} x_{a, \bullet}$	$x_{\bullet, \bar{1}} x_{\bar{1}, a+\bar{1}} x_{a+\bar{1}, \bullet}$
α_γ	$\frac{1}{2}$	$\frac{(-1)^{s+\bar{1}+t}}{2}$
y_1, y_2, y_3	$\text{Id}, \tau^a, \text{Id}$	τ, τ^a, Id
$(y_1 y_2 y_3)^{-1} \gamma$	$(-1)^a x_{\bullet, -a} x_{-a, \bar{0}} x_{\bar{0}, \bullet}$	$(-1)^{a+\bar{1}} x_{\bullet, -a} x_{-a, \bar{0}} x_{\bar{0}, \bullet}$
$\chi_{(y_1 y_2 y_3)^{-1}, \gamma}$	$(-1)^a$	$(-1)^{a+\bar{1}}$
$(y_1 y_2 y_3)^{-1}(\gamma)^*$	$x'_{\bullet, \bar{0}} x'_{\bar{0}, -a} x'_{-a, \bullet}$	$x'_{\bullet, \bar{0}} x'_{\bar{0}, -a} x'_{-a, \bullet}$
$\beta_{(y_1 y_2 y_3)^{-1}(\gamma)^*}$	$\begin{cases} 1 & \text{if } b = -a \\ 0 & \text{if not} \end{cases}$	$\begin{cases} 1 & \text{if } b = -a \\ 0 & \text{if not} \end{cases}$
x_2, x_1, x_0, h_0	$\text{Id}, \tau^{-a}, \text{Id}, \tau^{-a}$	$\text{Id}, \tau^{-a}, \text{Id}, \tau^{-a-\bar{1}}$
$\chi_{\rho_s}(\mathbf{h}_0)$	$(-1)^{sa}$	$(-1)^{s(a+\bar{1})}$
contribution of γ to $(F_a \phi_b)$	$\begin{cases} \frac{(-1)^{a(s+\bar{1})}}{2} & \text{if } b = -a \\ 0 & \text{if not} \end{cases}$	$\begin{cases} \frac{(-1)^{(s+\bar{1})a+t}}{2} & \text{if } b = -a \\ 0 & \text{if not} \end{cases}$

TABLE 2. The scalars appearing in the expression (2.1.3) of $(F_a|\phi_b)$

Remark 1.22. Like the operation $\otimes$, the pairing $(-|-)$ has a simple interpretation in the monoidal category $(\text{mod}(A^e), \otimes_A)$. Indeed,

- first, $M(\underline{i})$ admits $M^*(\underline{i}^o)$ as a left dual, see (4.1.5);
- next, if $M(\underline{i}; V)$ and $M^*(\underline{i}^o; U)$ are identified with $M(\underline{i}) \otimes_A V$ and $M^*(\underline{i}^o) \otimes_A U$, respectively, then $(-|-)$ is an explicit reformulation of $\langle -|- \rangle$ as introduced in part (2) of Remark 1.7, see (4.3.8).

Using $(-|-)$ yields intertwiners relative to M^* associated with the f_γ 's for paths γ in Q_G .

Notation 1.23. Let Q_G be such as in Setting 1.4.

- (1) Let $f \in (i, U) \rightarrow (j, V)$ be an arrow of Q_G . Using the basis of the vector space $\text{Hom}_{\mathbb{k}G_i}(U, M(i, j; V))$ consisting of the arrows $(i, U) \rightarrow (j, V)$ of Q_G and using the associated $(-|-)$ -dual basis of $\text{Hom}_{\mathbb{k}G_j}(V, M^*(j, i; U))$, denote by $f^\vee$ the dual element associated with f . In other words, $f^\vee$ lies in $\text{Hom}_{\mathbb{k}G_j}(V, M^*(j, i; U))$ and, for all arrows $f': (i, U) \rightarrow (j, V)$ of Q_G ,

$$(f'|f^\vee) = \begin{cases} 1 & \text{if } f' = f \\ 0 & \text{otherwise.} \end{cases}$$

- (2) For all paths γ in Q_G

$$\gamma: (i_0, U_0) \xrightarrow{f_1} (i_1, U_1) \rightarrow \cdots \rightarrow (i_{n-1}, U_{n-1}) \xrightarrow{f_n} (i_n, U_n),$$

denote by φ_γ the following intertwiner

$$\varphi_\gamma = f_1^\vee \otimes f_2^\vee \otimes \cdots \otimes f_n^\vee \in \text{Hom}_{\mathbb{k}G_{i_n}}(U_n, M^*(i_n, i_{n-1}, \dots, i_0; U_0)).$$

Example 1.24. This is the continuation of Example 1.21, it illustrates Notation 1.23.

Let $k \in \mathbb{Z}/2\mathbb{Z}$. Denote by a'_k and b'_k the following respective intertwiners relative to M^*

$$\begin{aligned} \rho_k &\longrightarrow M^*(\bullet, \bullet; \rho_{k+\bar{1}}) & \mathbb{k} \cdot \text{Id} &\longrightarrow M^*(\bar{0}, \bullet; \rho_k) \\ \varepsilon_k &\longmapsto x'_{\bullet, \bullet} \otimes \varepsilon_{k+\bar{1}} & \text{Id} &\longmapsto x'_{\bar{0}, \bullet} \otimes \varepsilon_k. \end{aligned}$$

Denote by c'_k and d'_k the following respective intertwiners relative to M^*

$$\begin{aligned} \rho_k &\longrightarrow M^*(\bullet, \bar{0}; \mathbb{k} \cdot \text{Id}) & \mathbb{k} \cdot \text{Id} &\longrightarrow M^*(\bar{0}, \bar{0}; \mathbb{k} \cdot \text{Id}) \\ \varepsilon_k &\longmapsto (x'_{\bullet, \bar{0}} \otimes \text{Id} + (-1)^{k+\bar{1}} x'_{\bullet, \bar{1}} \otimes \tau) & \text{Id} &\longmapsto x'_{\bar{0}, k} \otimes \tau^k. \end{aligned}$$

Then $(a_k|a'_k) = 1$, $(b_k|b'_k) = 1$, $(c_k|c'_k) = 1$ and $(d_k|d'_k) = \delta_{k, \ell}$ for all $\ell \in \mathbb{Z}/2\mathbb{Z}$, where $\delta_{k, \ell}$ stands for the Kronecker symbol. Using (2.1.3), the intermediate steps of the computation of these pairings are given in Tables 3, 4, 5 and 6 (see Example 1.21). In view of the shape of Q_G , these equalities entail that $a_k^\vee = a'_k$, $b_k^\vee = b'_k$, $c_k^\vee = c_k$ and $d_k^\vee = d'_k$.

γ	α_γ	y_1	$y_1^{-1} \gamma$	$\chi_{y_1^{-1}, \gamma}$	$y_1^{-1}(\gamma)^*$	$\beta_{y_1^{-1}(\gamma)^*}$	x_0, h_0	$\chi_{\rho_{k+\bar{1}}}(\mathbf{h}_0)$	contr. of γ
$x_{\bullet, \bullet}$	$\frac{1}{2}$	Id	$x_{\bullet, \bullet}$	1	$x'_{\bullet, \bullet}$	1	Id, Id	1	1

TABLE 3. $(a_k|a'_k) = 1$

γ	α_γ	y_1	$y_1^{-1} \gamma$	$\chi_{y_1^{-1}, \gamma}$	$y_1^{-1}(\gamma)^*$	$\beta_{y_1^{-1}(\gamma)^*}$	x_0, h_0	$\chi_{\rho_{\mathbf{k}}}(\mathbf{h}_0)$	contr. of γ
$x_{\bullet, \bar{0}}$	$\frac{1}{2}$	Id	$x_{\bullet, \bar{0}}$	1	$x'_{\bar{0}, \bullet}$	1	Id, Id	1	$\frac{1}{2}$
$x_{\bullet, \bar{1}}$	$\frac{(-1)^{k+\bar{1}}}{2}$	τ	$-x_{\bullet, \bar{0}}$	-1	$x'_{\bar{0}, \bullet}$	1	Id, τ	$(-1)^k$	$\frac{1}{2}$

TABLE 4. $(b_k|b'_k) = \frac{1}{2} + \frac{1}{2} = 1$

γ	α_γ	y_1	$y_1^{-1} \gamma$	$\chi_{y_1^{-1}, \gamma}$	$y_1^{-1}(\gamma)^*$	$\beta_{y_1^{-1}(\gamma)^*}$	x_0, h_0	$\chi_{\rho_{\mathbf{k}}}(\mathbf{h}_0)$	contr. of γ
$x_{\bar{0}, \bullet}$	1	Id	$x_{\bar{0}, \bullet}$	1	$x'_{\bullet, \bar{0}}$	1	Id, Id	1	$\frac{1}{2}$

TABLE 5. $(c_k|c'_k) = 1$

Now it is possible to give examples of intertwiners relative to M^* and associated to paths in Q_G . Let $s, t, a \in \mathbb{Z}/2\mathbb{Z}$. Denote by γ the following path of Q_G .

$$(\bullet, \rho_s) \xrightarrow{b_s} (\bar{0}, \mathbb{k} \cdot \text{Id}) \xrightarrow{d_a} (\bar{0}, \mathbb{k} \cdot \text{Id}) \xrightarrow{c_t} (\bullet, \rho_t).$$

γ	α_γ	y_1	$y_1^{-1} \gamma$	$\chi_{y_1^{-1}, \gamma}$	$y_1^{-1}(\gamma)^*$	$\beta_{y_1^{-1}(\gamma)^*}$	x_0, h_0	$\chi_{\mathbf{k} \cdot \text{Id}}(\mathbf{h}_0)$	contr. of γ
$x_{\bar{0}, k}$	$(-1)^{\mathbf{k}}$	$\tau^{\mathbf{k}}$	$(-1)^{\mathbf{k}} x_{k, \bar{0}}$	$(-1)^{\mathbf{k}}$	$x'_{\bar{0}, k}$	$\delta_{\mathbf{k}, \ell}$	$\tau^{-\mathbf{k}}, \text{Id}$	$\mathbf{1}$	$\delta_{\mathbf{k}, \ell}$

TABLE 6. $(d_k | d'_\ell) = \delta_{k, \ell}$

Then $\varphi_\gamma = b_s^\vee \otimes d_a^\vee \otimes c_t^\vee = b'_s \otimes d'_a \otimes c'_t$. This is the intertwiner $\rho_t \rightarrow M^*(\bullet, \bar{0}, \bar{0}, \bullet; \rho_s)$ such that (see (1.4.2))

$$\begin{aligned} \varphi_\gamma(\varepsilon_t) &= c'_t(\varepsilon_t) \cdot d'_a(\text{Id}) \cdot b'_s(\text{Id}) \\ &= (x'_{\bullet, \bar{0}} \otimes \text{Id} + (-1)^{t+\bar{1}} x'_{\bullet, \bar{1}} \otimes \tau) \cdot (x'_{\bar{0}, a} \otimes \tau^a) \cdot (x'_{\bar{0}, \bullet} \otimes \varepsilon_s) \\ &= (-1)^{a(s+\bar{1})} (x'_{\bullet, \bar{0}} x'_{\bar{0}, a} x'_{a, \bullet} \otimes \varepsilon_s + (-1)^{t+\bar{1}+s} x'_{\bullet, \bar{1}} x'_{\bar{1}, a+\bar{1}} x'_{a+\bar{1}, \bullet} \otimes \varepsilon_s). \end{aligned}$$

Thus, using the notation introduced in Example 1.17, $\varphi_\gamma = (-1)^{a(s+\bar{1})} \phi_a$.

1.6. The comparison of the algebra of intertwiners with the Morita reduction. Now, here is the result comparing $\mathbb{k}Q_G$ and $\mathcal{I}\text{ntw}$.

Theorem 1.25. *Keep Settings 1.1 and 1.4.*

(1) Assigning f_γ to every path γ in Q_G yields an isomorphism of algebras

$$(1.6.1) \quad \mathbb{k}Q_G \xrightarrow{\sim} \mathcal{I}\text{ntw}^{\text{op}}.$$

(2) For all $f \in \mathcal{I}\text{ntw}$, say lying in $\text{Hom}_{\mathbb{k}G_{i_0}}(U_0, M(i_0, \dots, i_n; U_n))$,

$$(1.6.2) \quad f = \sum_{\gamma} (f | \varphi_\gamma) f_\gamma,$$

where γ runs through all paths of Q_G of the shape $(i_0, U_0) \rightarrow (i_1, \bullet) \rightarrow \dots \rightarrow (i_{n-1}, \bullet) \rightarrow (i_n, U_n)$.

It is now possible to explain how to decompose any element of $\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$ along the paths in Q_G via the isomorphism $\mathbb{k}Q_G \xrightarrow{(1.1.1)} \tilde{e}(T_S(M) * G) \cdot \tilde{e}$. This is done with the following construction and assuming that $U = \mathbb{k}G_i \cdot \varepsilon_U$ for all vertices (i, U) of Q_G . Given a tensor $m_1 \otimes \dots \otimes m_n * g \in e_{i_0} \cdot (M^{\otimes n} * G) \cdot e_{i_n}$, where $n \geq 0$ and $i_0, i_n \in [G \setminus Q_0]$, then

$m_1 \otimes \dots \otimes m_n * g \in ({}_{i_0} M_{y_1 \cdot i_1} \otimes_{\mathbb{k}} \dots \otimes_{\mathbb{k}} ({}_{y_1 \dots y_{n-1}} M_{(y_1 \dots y_n) \cdot i_n}) * y_1 \dots y_n \mathbb{k}G_{i_n}$ for unique $i_1, \dots, i_{n-1} \in [G \setminus Q_0]$ and $y_t \in [G/G_{i_t}]$, for all $t \in \{1, \dots, n\}$; this defines an intertwiner, for all $U \in \text{irr}(G_{i_0})$ and $V \in \text{irr}(G_{i_n})$,

$$\begin{aligned} \mathbb{k}G_{i_0} \cdot \varepsilon_U &\longrightarrow M(i_0, \dots, i_n; \mathbb{k}G_{i_n} \cdot \varepsilon_V) \\ u \cdot \varepsilon_U &\longmapsto (u * \varepsilon_U) \cdot (m_1 \otimes \dots \otimes m_n * g) \cdot (e_{i_n} * \varepsilon_V), \end{aligned}$$

where the products on the right-hand side are meant in $T_S(M) * G$; summing these intertwiners over all $U \in \text{irr}(G_{i_0})$ and $V \in \text{irr}(G_{i_n})$ defines an element of $\mathcal{I}\text{ntw}$. This construction may be extended linearly to a linear mapping

$$\hat{e} \cdot (T_S(M) * G) \cdot \hat{e} \longrightarrow \mathcal{I}\text{ntw}^{\text{op}},$$

where $\hat{e} = \sum_{i \in [G \setminus Q_0]} e_i$. Denote by Ξ the restriction of this mapping to $\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$,

$$(1.6.3) \quad \Xi: \tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e} \longrightarrow \mathcal{I}\text{ntw}^{\text{op}}.$$

Remark 1.26. Here is how $\hat{e}$ and $\tilde{e}$ are related. Given idempotents a and b of an algebra Λ , say that b is a *reduction* of a whenever Λb is a direct summand of Λa in $\text{Mod}(\Lambda)$ and the algebras $\text{End}_\Lambda(\Lambda b)$ and $\text{End}_\Lambda(\Lambda a)$ are Morita equivalent. Now $\tilde{e}$ ($= \sum_{i \in [G \setminus Q_0]} e_i * (\sum_{U \in \text{irr}(G_i)} \varepsilon_U)$) is a reduction of $\hat{e} * 1_{\mathbb{k}G}$ ($= \sum_{i \in [G \setminus Q_0]} e_i * 1_{\mathbb{k}G_i}$) in the algebra $T_S(M) * G$.

Theorem 1.27. *Let Q_0 , S , G , and M be as in Setting 1.1. Let Q_G be as in Setting 1.4. Assume that $U = \mathbb{k}G_i \cdot \varepsilon_U$ for all vertices (i, U) of Q_G . Then, Ξ , as introduced in (1.6.3), is an isomorphism of $\mathbb{k}$ -algebras such that the following diagram is commutative,*

$$(1.6.4) \quad \begin{array}{ccc} & \tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e} & \\ \nearrow (1.1.1) & \downarrow \Xi & \\ \mathbb{k}Q_G & & \mathcal{Intw}^{\text{op}} \\ \searrow (1.6.1) & & \end{array}$$

In particular, if (1.1.1) is used as an identification, then, for all $\theta \in \tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$,

$$(1.6.5) \quad \theta = \sum_{\gamma \text{ path in } Q_G} (\Xi(\theta)|\varphi_\gamma) \cdot \gamma.$$

Example 1.28. *This is the continuation of Example 1.24. Denote by w the path $x_{\bullet, \bar{0}} x_{\bar{0}, \bar{1}} x_{\bar{1}, \bullet}$ in Q , this is an oriented cycle of length three going clockwise in the picture of Q in Example 1.3. The present example describes the intertwiner $\Xi(\tilde{e} \cdot (w * \text{Id}) \cdot \tilde{e})$ as well as its preimage under the isomorphism $\mathbb{k}Q_G \xrightarrow{(1.6.1)} \mathcal{Intw}^{\text{op}}$.*

Because of the definition,

$$\Xi(\tilde{e} \cdot (w * \text{Id}) \cdot \tilde{e}) = \sum_{s, t \in \mathbb{Z}/2\mathbb{Z}} \Xi(\tilde{e} \cdot (w * \text{Id}) \cdot \tilde{e})_{s, t},$$

where, for all $s, t \in \mathbb{Z}/2\mathbb{Z}$, the piece of notation $\Xi(\tilde{e} \cdot (w * \text{Id}) \cdot \tilde{e})_{s, t}$ stands for the intertwiner $\rho_s \rightarrow M(\bullet, \bar{0}, \bar{0}, \bullet; \rho_t)$ which maps ε_s to $(e_\bullet \otimes \varepsilon_s) \cdot \tilde{e} \cdot (w * \text{Id}) \cdot \tilde{e} \cdot (e_\bullet \otimes \varepsilon_t)$. For all $s, t \in \mathbb{Z}/2\mathbb{Z}$, the latter expression is equal to $(1_{\mathbb{k}Q} \otimes \varepsilon_s) \cdot (w \otimes \varepsilon_t)$, which in turn is equal to $\frac{1}{2}(x_{\bullet, \bar{0}} x_{\bar{0}, \bar{1}} x_{\bar{1}, \bullet} \otimes \varepsilon_t + (-1)^{s+\bar{1}+t} x_{\bullet, \bar{1}} x_{\bar{1}, \bar{0}} x_{\bar{0}, \bullet} \otimes \varepsilon_t)$; therefore, using the notation of Example 1.11,

$$\Xi(\tilde{e} \cdot (w * \text{Id}) \cdot \tilde{e})_{s, t} = F_{\bar{0}}.$$

Now, let $s, t, a \in \mathbb{Z}/2\mathbb{Z}$; using the computation of the pairings of the shape $(F_a|\phi_b)$ made in Example 1.21 as well as the description of the intertwiners relative to M^* associated to paths in Q_G made in Example 1.24, and denoting by γ the path $(\bullet, \rho_s) \xrightarrow{b_s} (\bar{0}, \mathbb{k} \cdot \text{Id}) \xrightarrow{d_a} (\bar{0}, \mathbb{k} \cdot \text{Id}) \xrightarrow{c_t} (\bullet, \rho_t)$ of Q_G ,

$$(\Xi(\tilde{e}(w \otimes \text{Id}) \cdot \tilde{e})_{s, t}|\varphi_\gamma) = \begin{cases} \frac{1+(-1)^t}{2} & \text{if } a = \bar{0} \\ 0 & \text{if not;} \end{cases}$$

Therefore $\Xi(\tilde{e} \cdot (w \otimes \text{Id}) \cdot \tilde{e})_{s, t} = \frac{1+(-1)^t}{2} \cdot b_s d_{\bar{0}} c_t$ (see (1.6.5)). As a whole, $\Xi(\tilde{e} \cdot (w \otimes \text{Id}) \cdot \tilde{e}) = b_{\bar{0}} d_{\bar{0}} c_{\bar{0}} + b_{\bar{1}} d_{\bar{0}} c_{\bar{0}}$.

2. PRACTICAL ASPECTS

Let Q_0 , S , G and M be as in Setting 1.1.

Applying Theorem 1.27 for computations requires to compute Q_G first. Actually, rather than computing explicitly the intertwiners relative to M which form the arrows of Q_G , it is simpler to

- (1) compute a basis of $\text{Hom}_{\mathbb{k}G_j}(\tau, M^*(j, i; \rho))$ for all pairs (i, ρ) and (j, τ) such that $i, j \in [G \setminus Q_0]$, $\rho \in \text{irr}(G_i)$, and $\tau \in \text{irr}(G_j)$.

Because of the pairing $(-|-)$, this yields a quiver Q_G such as in Setting 1.4 as well as the intertwiners φ_α for all arrows α of Q_G . This avoids computing explicitly the intertwiners corresponding to the arrows of Q , which are not used in the formulas of Theorem 1.27.

Given an element θ of $\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$, say homogeneous of degree n for the grading induced by the tensor powers of M , the decomposition into a linear combination of paths in Q_G of the inverse image of θ under the isomorphism $\mathbb{k}Q_G \xrightarrow{(1.1.1)} \tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$ may be computed as follows using Theorem 1.27,

- (2) compute $\Xi(\theta)$, which is a sum of intertwiners relative to M , see (1.6.3);
- (3) compute the intertwiners φ_γ for all paths γ of length n in Q_G , see Notation 1.23;
- (4) compute the pairing $(\Xi(\theta)|\varphi_\gamma)$ for all such paths γ , see Proposition 1.20.

The desired linear combination of paths is hence $\sum_\gamma (\Xi(\theta)|\varphi_\gamma) \cdot \gamma$ (see (1.6.5)).

These computations may be performed by a computer that is able to determine the irreducible representations of $\text{irr}(G_i)$ for all $i \in [G \setminus Q_0]$. This section illustrates these computations.

2.1. How to pair intertwiners? In view of (1.4.2), the main difficulty in the computations listed in the introduction of Section 2 lies in computing pairings for $(-|-)$. This section explains how to compute (1.5.2) in combinatorial terms in the following specific setting.

Setting 2.1. Let Q_0 , S , G and M be as in Setting 1.1. Let Q be a quiver with vertex set being Q_0 and such that M is equal to the vector space with basis elements being the arrows of Q . Let Q_G be as in Setting 1.4. Assume that

- the stabiliser G_i is abelian for all $i \in Q_0$,
- $U = \mathbb{k}G_i \cdot \varepsilon_U$ for all vertices (i, U) of Q_G , denote by χ_U the character $G_i \rightarrow \mathbb{k}^\times$ of U ,
- and, for all $g \in G$ and all paths γ in Q , there exists a scalar $\chi_{g,\gamma} \in \mathbb{k}^\times$ and a path $g(\gamma)$ such that ${}^g\gamma = \chi_{g,\gamma} g(\gamma)$.

If G is abelian, then it is possible to find a quiver Q and to choose irreducible representations U of G_i , for all $i \in [G \setminus Q_0]$, fitting in this setting.

Notation 2.2. The following notation relative to M is useful.

- For all arrows $a: i \rightarrow j$ of Q , denote by a^* the element of M^* such that $a^*(a) = 1$ and $a^*(b) = 0$ for all arrows b of Q distinct from a .
- Denote by Q^* the quiver with vertex set being Q_0 and whose arrows are the linear forms a^* introduced just before. This quiver is isomorphic to the opposite quiver of Q .
- For all paths $\gamma: t_0 \xrightarrow{a_1} t_1 \rightarrow \cdots \rightarrow t_{n-1} \xrightarrow{a_n} t_n$ in Q , denote by γ^* the path $t_n \xrightarrow{a_n^*} t_{n-1} \rightarrow \cdots \rightarrow t_1 \xrightarrow{a_1^*} t_0$ in Q^* .

Note that every path in Q starting in some vertex i_0 lying in $[G \setminus Q_0]$ is of the shape $i_0 \rightarrow y_1 \cdot i_1 \rightarrow \cdots \rightarrow (y_1 \cdots y_n) \cdot i_n$ for a unique sequence $i_1, \dots, i_n$ in $[G \setminus Q_0]$ and a unique sequence $y_1, \dots, y_n$ in G such that $y_t \in [G/G_{i_t}]$ for all $t \in \{1, \dots, n\}$.

Let $\underline{i} = i_0, \dots, i_n$ be a sequence in $[G \setminus Q_0]$, consider $U \in \text{irr}(G_{i_0})$ and $V \in \text{irr}(G_{i_n})$, and let $f: U \rightarrow M(\underline{i}; V)$ and $\varphi: V \rightarrow M^*(\underline{i}^\circ; U)$ be intertwiners. Since the representation V is one dimensional, then,

$$(2.1.1) \quad f(\varepsilon_u) = \sum_{\gamma} \alpha_\gamma \gamma \otimes y_1 y_2 \cdots y_n \varepsilon_v,$$

where γ runs through all paths in Q of the shape

$$i_0 \rightarrow y_1 \cdot i_1 \rightarrow y_1 y_2 \cdot i_2 \rightarrow \cdots \rightarrow (y_1 \cdots y_n) \cdot i_n$$

for some (unique) sequence $\underline{y} = y_1, \dots, y_n$ such that $y_t \in [G/G_{i_t}]$ for all $t \in \{1, \dots, n\}$, and $\alpha_\gamma \in \mathbb{k}$. Note that the considered paths are precisely the ones of the

shape $i_0 \rightarrow j_1 \rightarrow \cdots \rightarrow j_n$ such that j_t lies in the orbit of i_t for all $t \in \{1, \dots, n\}$. Similarly,

$$(2.1.2) \quad \varphi(\varepsilon_V) = \sum_{\gamma'} \beta_{\gamma'} \gamma' \otimes x_{n-1} \cdots x_0 \varepsilon_U,$$

where γ' runs through all paths in Q of the shape

$$i_n \rightarrow x_{n-1} \cdot i_{n-1} \rightarrow x_{n-1}x_{n-2} \cdot i_{n-2} \rightarrow \cdots \rightarrow (x_{n-1}x_{n-2} \cdots x_0) \cdot i_0$$

for some (unique) sequence $\underline{x} = x_{n-1}, \dots, x_0$ such that $x_t \in [G/G_{i_t}]$ for all $t \in \{0, \dots, n-1\}$, and $\beta_{\gamma'} \in \mathbb{k}$.

Lemma 2.3. *Using the decompositions (2.1.1) and (2.1.2), then*

$$(2.1.3) \quad (f|\varphi) = \sum_{\gamma} \alpha_{\gamma} \cdot \beta_{(y_1 \cdots y_n)^{-1}(\gamma)^*} \cdot \chi_{(y_1 \cdots y_n)^{-1}, \gamma} \cdot \chi_U(h_0),$$

where γ runs through all paths in Q of the shape $i_0 \rightarrow j_1 \rightarrow \cdots \rightarrow j_n$ such that j_t lies in the orbit of i_t for all $t \in \{1, \dots, n\}$, where $\alpha_{\gamma} \in \mathbb{k}$, and where, for all such paths γ ,

- $\underline{y} = y_1, \dots, y_n$ denotes the sequence in G such that $y_t \in [G/G_{i_t}]$, for all $t \in \{1, \dots, n\}$ and γ is a path of the shape $i_0 \rightarrow y_1 \cdot i_1 \rightarrow \cdots \rightarrow y_1 \cdots y_n \cdot i_n$,
- h_0 is the element of G_{i_0} such as in (c) in Proposition 1.20,
- $\chi_{(y_1 \cdots y_n)^{-1}, \gamma}$ is the element of $\mathbb{k}^{\times}$ and $(y_1 \cdots y_n)^{-1}(\gamma)$ is the path in Q such that $(y_1 \cdots y_n)^{-1} \gamma = \chi_{(y_1 \cdots y_n)^{-1}, \gamma} (y_1 \cdots y_n)^{-1}(\gamma)$,
- and $((y_1 \cdots y_n)^{-1}(\gamma))^*$ is the reverse path in Q^* associated to the path $(y_1 \cdots y_n)^{-1}(\gamma)$ in Q .

Proof. For all sequences $\underline{y} = y_1, \dots, y_n$ and $\underline{x} = x_{n-1}, \dots, x_0$ such that $x_t \in [G/G_{i_t}]$ and $y_t \in [G/G_{i_t}]$ for all t , denote by $\mathcal{C}(\underline{y})$ and $\mathcal{C}'(\underline{x})$ the sets of paths in Q and in Q^* of the shape $i_0 \rightarrow y_1 \cdot i_1 \rightarrow \cdots \rightarrow y_1 \cdots y_n \cdot i_n$ and $i_n \rightarrow x_{n-1} \cdot i_{n-1} \rightarrow \cdots \rightarrow x_{n-1} \cdots x_0 \cdot i_0$, respectively. For all $\underline{y}$,

$$f_{\underline{y}}(\varepsilon_U) = \sum_{\gamma \in \mathcal{C}(\underline{y})} \alpha_{\gamma} \gamma \otimes y_1 \cdots y_n \varepsilon_U;$$

hence, the following term that serves in the definition of $(f|\varphi)$, see (1.5.2),

$$(y_1 \cdots y_n)^{-1} f_{\underline{y}}^{(1)}(\varepsilon_U) \otimes \cdots \otimes (y_1 \cdots y_n)^{-1} f_{\underline{y}}^{(n)}(\varepsilon_U) \otimes f_{\underline{y}}^{(0)}(\varepsilon_U),$$

is equal to

$$\sum_{\gamma \in \mathcal{C}(\underline{y})} \alpha_{\gamma} \cdot \chi_{(y_1 \cdots y_n)^{-1}, \gamma} \cdot \underbrace{(y_1 \cdots y_n)^{-1}(\gamma)}_{\in M^{\otimes S^n}} \otimes \varepsilon_U.$$

Similarly, for all $\underline{x}$,

$$\varphi^{\underline{x}}(\varepsilon_V) = \sum_{\gamma' \in \mathcal{C}(\underline{x})} \beta_{\gamma'} \gamma' \otimes x_{n-1} \cdots x_0 \varepsilon_U.$$

Therefore, given $\underline{y}$, the “ Π_t ”-term in (1.5.2) is equal to

$$(2.1.4) \quad \sum_{\gamma \in \mathcal{C}(\underline{y}), \gamma' \in \mathcal{C}(\underline{x})} \alpha_{\gamma} \cdot \beta_{\gamma'} \chi_{(y_1 \cdots y_n)^{-1}, \gamma} \cdot b_n^*(c_n) \cdots b_2^*(c_2) \cdot b_1^*(c_1) \cdot \chi_U(h_0) \cdot \varepsilon_U,$$

where

- $\underline{x} = x_{n-1}, \dots, x_0$ and h_0 are the elements of G determined by $\underline{y}$ and by (a), (b), and (c) in Proposition 1.20,
- $b_1^*, \dots, b_n^*$ are the arrows of Q^* such that γ' is $\cdot \xrightarrow{b_n^*} \cdot \xrightarrow{b_{n-1}^*} \cdots \xrightarrow{b_1^*} \cdot$,
- and $c_1, \dots, c_n$ are the arrows of Q such that $(y_1 \cdots y_n)^{-1}(\gamma)$ is $\cdot \xrightarrow{c_1} \cdot \xrightarrow{c_2} \cdots \xrightarrow{c_n} \cdot$.

By definition of the arrows $b_1^*, \dots, b_n^*$, the summand of (2.1.4) with index γ is non-zero only if the paths $\cdot \xrightarrow{b_1} \cdot \xrightarrow{b_2} \dots \xrightarrow{b_n} \cdot$ and $(y_1 \dots y_n)^{-1}(\gamma)$ in Q are equal, that is, only if $\gamma' = ((y_1 \dots y_n)^{-1}(\gamma))^*$. Summing (2.1.4) over all possible $\underline{y}$ yields that $(f|\varphi)\varepsilon_U$ is equal to

$$\sum_{\gamma} \alpha_{\gamma} \beta_{((y_1 \dots y_n)^{-1}(\gamma))^*} \chi_{(y_1 \dots y_n)^{-1}, \gamma} \chi_U(h_0) \varepsilon_U.$$

This proves (2.1.3). $\square$

2.2. An example. In this example, Q is the following quiver whose vertices are the elements of $\mathbb{Z}/4\mathbb{Z}$

(2.2.1)

and G is the dihedral group of order 10

$$G = \langle c, \tau \mid c^5, \tau^2, \tau c \tau c \rangle.$$

Denote by ε the group homomorphism $G \rightarrow \{-1, 1\}$ such that $\varepsilon(c) = 1$ and $\varepsilon(\tau) = -1$. Denote the arrows of Q by $x_{i,j}: i \rightarrow j$. For convenience, denote the arrows of Q^* by $x'_{i,j}: i \rightarrow j$, hence $(x_{j,i})^* = x'_{i,j}$. The actions of G on Q_0 and M are assumed to be the ones such that, for all $i \in Q_0$ and $g \in G$, and for all arrows $x_{i,j}$ of Q ,

$$c \cdot i = i + \bar{1}, \quad \tau \cdot i = -i, \quad {}^g x_{i,j} = \varepsilon(g) x_{g \cdot i, g \cdot j}.$$

The purpose of this section is to make a full illustration of Theorem 1.27, that is, for all $w \in T_S(M) * G$,

- to describe the intertwiner $\Xi(\theta)$, where θ denotes $\tilde{e} \cdot w \cdot \tilde{e}$;
- and to describe the preimage of θ under the isomorphism $\mathbb{k}Q_G \xrightarrow{(1.1.1)} \tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$.

Let $[G \backslash Q_0]$ be $\{\bar{0}\}$. Note that $G_{\bar{0}} = \{\text{Id}, \tau\}$. Let $[G/G_{\bar{0}}]$ be $\{\text{Id}, c, c^2, c^3, c^4\}$. For all $s \in \{0, 1\}$, denote by ε_s the following primitive idempotent of $\mathbb{k}G_{\bar{0}}$,

$$\varepsilon_s = \frac{1}{2}(\text{Id} + (-1)^s \tau)$$

and denote $\varepsilon_s \cdot \mathbb{k}G_{\bar{0}}$ by ρ_s . Finally, let $\text{irr}(G_{\bar{0}})$ be equal to $\{\rho_0, \rho_1\}$. In particular,

$$\tilde{e} = e_{\bar{0}} * \varepsilon_0 + e_{\bar{0}} * \varepsilon_1.$$

The setting presented below is assumed until the end of the subsection. In view of the purpose stated previously, this is not a loss of generality in making this assumption because

- the elements of the shape $p * g$, where p is a path in Q and $g \in G$, form a basis of $T_S(M) * G$ as a vector space;
- every $g \in G$ is equal to $c^\ell \tau^r$ for a unique couple $(\ell, r) \in \{0, 1, 2, 3, 4\} \times \{0, 1\}$;
- for all paths p of Q and for all $(\ell, r) \in \{0, 1, 2, 3, 4\} \times \{0, 1\}$, the product $\tilde{e} \cdot (p * g) \cdot \tilde{e}$ vanishes as soon as p is not a path from $\bar{0}$ to $-\bar{\ell}$.

Setting 2.4. Assume that w is equal to $p * c^\ell \tau^r$, where $(\ell, r) \in \{0, 1, 2, 3, 4\} \times \{0, 1\}$ and p is a path in Q from $\bar{0}$ to $-\bar{\ell}$. Denote the length of p by n . Denote by $i_0, \dots, i_n$ the sequence of vertices of Q such that p has the shape $i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_n$. Thus,

$p = x_{i_0, i_1} x_{i_1, i_2} \cdots x_{i_{n-1}, i_n}$. In particular, $i_0 = \bar{0}$, $i_n = -\bar{\ell}$ and $i_k - i_{k-1} \in \{-\bar{1}, \bar{1}\}$ for all $k \in \{1, \dots, n\}$.

2.2.1. *Computation of Q_G .* By definition, for all $t \in \{0, 1\}$,

$$M^*(\bar{0}, \bar{0}; \rho_t) = \text{Span}(x'_{\bar{0}, \bar{1}} \otimes c\varepsilon_t, x'_{\bar{0}, \bar{4}} \otimes c^4\varepsilon_t),$$

where the action of τ is given as follows

$$\begin{cases} x'_{\bar{0}, \bar{1}} \otimes c\varepsilon_t & \mapsto (-1)^{1+t} x'_{\bar{0}, \bar{4}} \otimes c^4\varepsilon_t \\ x'_{\bar{0}, \bar{4}} \otimes c^4\varepsilon_t & \mapsto (-1)^{1+t} x'_{\bar{0}, \bar{1}} \otimes c\varepsilon_t. \end{cases}$$

Hence, for all $s, t \in \{0, 1\}$,

$$\text{Hom}_{\mathbb{k}G_{\bar{0}}}(\rho_s, M^*(\bar{0}, \bar{0}; \rho_t)) = \text{Span}(\varphi_{s,t}),$$

where $\varphi_{s,t}$ is the intertwiner such that $\varphi_{s,t}(\varepsilon_s) = \varepsilon_s \cdot (x'_{\bar{0}, \bar{1}} \otimes c\varepsilon_t) (= \varepsilon_s \cdot (x'_{\bar{0}, \bar{1}} \otimes c) \cdot \varepsilon_t)$, that is,

$$\varphi_{s,t}(\varepsilon_s) = x'_{\bar{0}, \bar{1}} \otimes c\varepsilon_t + (-1)^{s+t+1} x'_{\bar{0}, \bar{4}} \otimes c^4\varepsilon_t.$$

Denote by $f_{s,t}$ the intertwiner $\rho_s \rightarrow M(\bar{0}, \bar{0}; \rho_t)$ such that $(f_{s,t}|\varphi_{t,s}) = 1$, for all $s, t \in \{0, 1\}$. Then Q_G may be taken equal to the following quiver

$$\begin{array}{ccc} \begin{array}{c} \text{\scriptsize } f_{0,0} \\ \curvearrowright \\ (1, \rho_0) \end{array} & \begin{array}{c} \xrightarrow{\text{\scriptsize } f_{0,1}} \\ \xleftarrow{\text{\scriptsize } f_{1,0}} \end{array} & \begin{array}{c} \text{\scriptsize } f_{1,1} \\ \curvearrowright \\ (1, \rho_1) \end{array} \end{array}$$

2.2.2. *The intertwiner $\Xi(\theta)$.* By definition, $\Xi(\theta)$ decomposes as

$$(2.2.2) \quad \Xi(\theta) = \sum_{s,t \in \{0,1\}} \Xi(\theta)_{s,t},$$

where, for all $s, t \in \{0, 1\}$, the intertwiner $\Xi(\theta)_{s,t}: \rho_s \rightarrow M(\underbrace{0, \dots, 0}_{n+1}; \rho_t)$ is such that

$\Xi(\theta)_{s,t}(\varepsilon_s) = \varepsilon_s \cdot \tilde{e} \cdot w \cdot \tilde{e} \cdot \varepsilon_t$, see (1.6.3). Note that

$$\begin{aligned} \Xi(\theta)_{s,t}(\varepsilon_s) &= \varepsilon_s \cdot \tilde{e} \cdot w \cdot \tilde{e} \cdot \varepsilon_t \\ &= \frac{1}{2}(\text{Id} + (-1)^s \tau) \cdot (p \otimes c^\ell \tau^r) \cdot \varepsilon_t \\ (2.2.3) \quad &= \frac{(-1)^{rt}}{2}(\text{Id} + (-1)^s \tau) \cdot (p \otimes c^\ell \varepsilon_t) \\ &= \frac{(-1)^{rt}}{2} (p \otimes c^\ell \varepsilon_t + (-1)^{s+t} (\tau p) \otimes c^{-\ell} \varepsilon_t), \end{aligned}$$

where $\tau p = (-1)^n x_{-i_0, -i_1} x_{-i_1, -i_2} \cdots x_{-i_{n-1}, -i_n}$.

2.2.3. *Intertwiners relative to M^* associated to paths of Q_G .* Consider a path of length denoted by m in Q_G , say $q: (0, \rho_{s_0}) \rightarrow (0, \rho_{s_1}) \rightarrow \cdots \rightarrow (0, \rho_{s_m})$, where $s_0, \dots, s_m \in \{-1, 1\}$. Then φ_q is the following intertwiner $\rho_{s_m} \rightarrow M^*(\underbrace{0, \dots, 0}_{n+1}; \rho_{s_0})$,

$$\varphi_q = \varphi_{s_1, s_0} \otimes \varphi_{s_2, s_1} \otimes \cdots \otimes \varphi_{s_m, s_{m-1}}.$$

Consider $\varphi_q(\varepsilon_{s_m})$ as an element of $\mathbb{k}Q^* * G$; following (1.4.2) it equals the product $(x'_{\bar{0}, \bar{1}} \otimes (-1)^{s_m + s_{m-1} + 1} x'_{\bar{0}, \bar{4}} \otimes c^4) \cdots (x'_{\bar{0}, \bar{1}} \otimes (-1)^{s_1 + s_0 + 1} x'_{\bar{0}, \bar{4}} \otimes c^4)$. Hence

$$(2.2.4) \quad \varphi_q(\varepsilon_{s_m}) = \sum (-1)^{\sum s_k + s_{k-1} + 1} x'_{\bar{0}, j_{m-1}} x'_{j_{m-1} j_{m-2}} \cdots x'_{j_1, j_0} \otimes c^{j_0},$$

where

- the first sum runs over all sequences $j_{m-1}, \dots, j_0 \in \mathbb{Z}/4\mathbb{Z}$ such that the expression $x'_{\bar{0}, j_{m-1}} \cdots x'_{j_1, j_0}$ is a path in Q' , which means that $j_t - j_{t-1} \in \{-\bar{1}, \bar{1}\}$ for all $t \in \{1, \dots, m\}$ (taking $j_m = \bar{0}$),

- and the second sum (in the exponent) runs over all indices $k \in \{1, \dots, m\}$ such that $x'_{j_k, j_{k-1}}$ of Q' goes anticlockwise when Q' is drawn like Q in (2.2.1), which means that $j_k - j_{k-1} = \bar{1}$.

2.2.4. *Computation of the pairing $(\Xi(\theta)|\varphi_q)$.* Let $q: (0, \rho_{s_0}) \rightarrow (0, \rho_{s_1}) \rightarrow \dots \rightarrow (0, \rho_{s_m})$ be a path in Q_G like in 2.2.3. Because of the definition of the action of G on $\mathbb{k}Q$, the pairing $(\Xi(\theta)|\varphi_q) (= (\Xi(\theta)_{s_0, s_m}|\varphi_q))$ is given by (2.1.3). In particular, because of (2.2.3) and (2.2.4),

- this pairing vanishes if $m \neq n$, that is, p and q have different lengths,
- if $m = n$ then, for all paths γ distinct from both $x_{i_0, i_1} x_{i_1, i_2} \dots x_{i_{n-1}, i_n}$ and $x_{i_0, -i_1} x_{-i_1, -i_2} \dots x_{-i_{n-1}, -i_n}$, the term of index γ in (2.2.3) vanishes.

Using (2.2.3) and (2.2.4), the scalars appearing in (2.1.3) may be computed directly for these two possibilities for γ , assuming that $m = n$. Table 7 describes these scalars, they appear in bold face. Recall that $i_0 = \bar{0}$.

γ	$x_{i_0, i_1} x_{i_1, i_2} \dots x_{i_{n-1}, i_n}$	$x_{i_0, -i_1} x_{-i_1, -i_2} \dots x_{-i_{n-1}, -i_n}$
α_γ	$\frac{(-1)^{rsn}}{2}$	$\frac{(-1)^{rsn+s_0+s_n+n}}{2}$
$y_1, \dots, y_n$	$c^{i_1-i_0}, c^{i_2-i_1}, \dots, c^{i_n-i_{n-1}}$	$c^{-i_1+i_0}, c^{-i_2+i_1}, \dots, c^{-i_n+i_{n-1}}$
$y_1 \dots y_n$	c^{in}	c^{-in}
$(y_1 \dots y_n)^{-1} \gamma$	$x_{-i_n, i_1-i_n} x_{i_1-i_n, i_2-i_n} \dots x_{i_{n-1}-i_n, \bar{0}}$	$x_{i_n, i_n-i_1} x_{i_n-i_1, i_n-i_2} \dots x_{i_n-i_{n-1}, \bar{0}}$
$\chi_{(y_1 \dots y_n)^{-1} \gamma}$	1	1
$(y_1 \dots y_n)^{-1} (\gamma)^*$	$x'_{\bar{0}, i_{n-1}-i_n} \dots x'_{i_2-i_n, i_1-i_n} x'_{i_1-i_n, -i_n}$	$x'_{\bar{0}, i_n-i_{n-1}} \dots x'_{i_n-i_2, i_n-i_1} x'_{i_n-i_1, i_n}$
$\beta_{(y_1 \dots y_n)^{-1} (\gamma)^*}$	$(-1)^{\mathbf{x}}$ where x is $\sum_{1 \leq k \leq n} s_k + s_{k-1} + 1$ s.t. $i_k - i_{k-1} = \bar{1}$	$(-1)^{\mathbf{x}}$, where x is $\sum_{1 \leq k \leq n} s_k + s_{k-1} + 1$ s.t. $i_k - i_{k-1} = -\bar{1}$
h_0	Id	Id
$\chi_{\rho_{s_0}}(h_0)$	1	1

TABLE 7. The coefficients in the expression (2.1.3) of $(\Xi(\theta)_{s_0, s_m}|\varphi_q)$ according to γ

Thus

$$\begin{aligned}
 (\Xi(\theta)|\varphi_q) &= \frac{(-1)^{rs_0}}{2} (-1)^{\sum_{i_k - i_{k-1} = \bar{1}} (s_k + s_{k-1} + 1)} \\
 &+ \frac{(-1)^{rs_0 + s_0 + s_n + n}}{2} (-1)^{\sum_{i_k - i_{k-1} = -\bar{1}} (s_k + s_{k-1} + 1)}.
 \end{aligned}$$

As whole (after factoring and simplifying),

$$(2.2.5) \quad (\Xi(\theta)|\varphi_q) = \begin{cases} \frac{(-1)^{rs_0}}{2} (-1)^{\sum_{i_k - i_{k-1} = \bar{1}} (s_k + s_{k-1} + 1)} & \text{if } m = n \\ 0 & \text{otherwise.} \end{cases}$$

2.2.5. *Decomposition of θ into paths in Q_G .* Because of Theorem 1.27, the preimage of θ under the isomorphism $\mathbb{k}Q_G \xrightarrow{(1.1.1)} \tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$ is equal to

$$\sum_{q \text{ path in } Q_G} (\Xi(\theta)|\varphi_q) \cdot q.$$

In view of (2.2.5), this preimage is hence equal to

$$\sum_{s_0, \dots, s_n \in \{0, 1\}} (-1)^{rs_0} (-1)^{\sum_{i_k - i_{k-1} = \bar{1}} s_k + s_{k-1} + 1} \cdot ((0, \rho_{s_0}) \rightarrow (0, \rho_{s_1}) \rightarrow \dots \rightarrow (0, \rho_{s_n})).$$

2.2.6. *Illustration.* Assume that $w = x_{\bar{0}, \bar{1}} x_{\bar{1}, \bar{2}} x_{\bar{2}, \bar{3}} x_{\bar{3}, \bar{4}} x_{\bar{4}, \bar{0}} * \text{Id} - x_{\bar{0}, \bar{4}} x_{\bar{4}, \bar{3}} x_{\bar{3}, \bar{2}} x_{\bar{2}, \bar{1}} x_{\bar{1}, \bar{0}} * \text{Id}$. Applying 2.2.5 yields that the preimage of $\tilde{e} \cdot w \cdot \tilde{e}$ under the isomorphism $\mathbb{k}Q_G \xrightarrow{(1.1.1)} \tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$ is equal to $-2 \sum_{\gamma} \gamma$, where γ runs through all oriented cycles of length 5 in Q_G .

3. DETAILS ON THE MONOIDAL CATEGORY OF BIMODULES OVER GROUPS

This section details some isomorphisms and properties in the monoidal category $(\text{mod}(A^e), \otimes_A)$ and related to the operation, say $\times$, and the pairing $\langle - | - \rangle$ considered in Remark 1.7.

- Section 3.1 describes the left dual of a tensor product of objects in $\text{mod}(A^e)$.
- Section 3.2 details the adjunction $(X \otimes_A -) \vdash (X^* \otimes_A -)$ for a given $X \in \text{mod}(A^e)$.
- Section 3.3 details the compatibility of $\langle - | - \rangle$.

The existence of the isomorphisms and adjunctions discussed in these sections may be part of the folklore (see [5, Exercise 2.10.16]). However, the proof of Theorem 1.25 is based on their description, which is the reason for detailing these here. For the sake of simplicity, the presentation is made independently of the monoidal category $(\text{mod}(A^e), \otimes_A)$ and of the framework of Theorem 1.25.

Here are some conventions. In any group, e denotes the neutral element; and $\text{Hom}_{\mathbb{k}G}$ and $\otimes_{\mathbb{k}G}$ are denoted by Hom_G and $\otimes_G$, respectively, for all finite groups G .

Given finite groups G and H , by “a module ${}_G X_H$ ” is meant a $\mathbb{k}G - \mathbb{k}H$ -bimodule, by “a module ${}_G X$ ” is meant a left $\mathbb{k}G$ -module, and by “by a module X_H ” is meant a right $\mathbb{k}H$ -module.

Given finite groups G and H and given a $\mathbb{k}G - \mathbb{k}H$ -bimodule X , the dual vector space X^* is a $\mathbb{k}H - \mathbb{k}G$ -bimodule in a natural way ($h \cdot \phi \cdot g = \phi(g \cdot \bullet \cdot h)$ for all $g \in G$, $h \in H$ and $\phi \in X^*$). In particular, taking H to be the trivial group, this defines a functor $X \mapsto X^*$ from left $\mathbb{k}G$ -modules to right $\mathbb{k}G$ -modules.

3.1. Tensor product of duals and dual of a tensor product.

Lemma 3.1. *Let H be a finite group with order not divisible by $\text{char}(\mathbb{k})$. For all finite dimensional $\mathbb{k}H$ -modules X_H and ${}_H Y$, there is a functorial bijective mapping*

$$(3.1.1) \quad \begin{aligned} Y^* \otimes_H X^* &\rightarrow (X \otimes_H Y)^* \\ \phi_2 \otimes \phi_1 &\mapsto (x \otimes y \mapsto \sum_{h \in H} \phi_1(x \cdot h) \phi_2(h^{-1} \cdot y)). \end{aligned}$$

Proof. The mapping is well defined. When $X = \mathbb{k}H_H$ and $Y = {}_H \mathbb{k}H$, it identifies with the following one, where $\{\delta_h\}_{h \in H}$ denotes the canonical basis of $(\mathbb{k}H)^*$,

$$\begin{aligned} (\mathbb{k}H)^* \otimes_H (\mathbb{k}H)^* &\rightarrow (\mathbb{k}H)^* \\ \delta_{h_2} \otimes \delta_{h_1} &\mapsto \delta_{h_1 h_2}; \end{aligned}$$

and the latter is bijective. Since X_H and ${}_H Y$ are finite dimensional and projective, then (3.1.1) is bijective. $\square$

The previous lemma generalises as follows. Given an integer $n \geq 2$, a sequence of groups $G_0, \dots, G_n$ and a finite dimensional $\mathbb{k}G_{i-1} - \mathbb{k}G_i$ -bimodule U_i for every $i \in \{1, \dots, n\}$, there is a functorial bijective mapping

$$(3.1.2) \quad X_n^* \otimes_{G_{n-1}} \cdots \otimes_{G_1} X_1^* \rightarrow (X_1 \otimes_{G_1} \cdots \otimes_{G_{n-1}} X_n)^*,$$

which maps any $\phi_n \otimes \cdots \otimes \phi_1$ to the following linear form on $X_1 \otimes_{G_1} \cdots \otimes_{G_{n-1}} X_n$,

$$x_1 \otimes \cdots \otimes x_n \mapsto \sum_{g_1, \dots, g_{n-1}} \prod_{t=1}^n \phi_t(g_{t-1}^{-1} \cdot x_t \cdot g_t),$$

where g_t runs through G_t ($1 \leq t \leq n-1$) and $g_0 = g_n = e$.

3.2. Adjunctions.

Lemma 3.2. *Let G and H be finite groups with orders not divisible by $\text{char}(\mathbb{k})$. Let ${}_G X_H$, ${}_H V$ and ${}_G U$ be (bi)modules over $\mathbb{k}G$ and $\mathbb{k}H$ such that X and V are finite dimensional. Then, there exists a functorial isomorphism*

$$\begin{aligned} & \Psi_X^{V,U} : \text{Hom}_H(V, X^* \otimes_G U) \rightarrow \text{Hom}_G(X \otimes_H V, U), \\ & \text{such that, for all } \phi \in \text{Hom}_H(V, X^* \otimes_G U), x \in X \text{ and } v \in V, \\ (3.2.1) \quad & (\Psi_X^{V,U}(\phi))(x \otimes v) = \sum_{g \in G} (\phi^{(2)}(v))(g^{-1} \cdot x)g \cdot \phi^{(1)}(v), \end{aligned}$$

where the following notation is used with sum sign omitted, for all $v \in V$,

$$\phi(v) = \phi^{(2)}(v) \otimes \phi^{(1)}(v) \in X^* \otimes_G U.$$

In particular, the pair of functors $(X \otimes_H -, X^* \otimes_G -)$ between finite dimensional left $\mathbb{k}H$ -modules and finite dimensional left $\mathbb{k}G$ -modules is adjoint.

Proof. Using the definition of the structure of H - G -bimodule of X^* , simple changes of variable in the sum " $\sum_{g \in G}$ " show that $\Psi_X^{V,U}$ is indeed well defined.

Since ${}_H V$ is finitely generated and projective, it suffices to prove the lemma assuming that $V = \mathbb{k}H$. In this case, $\Psi_X^{V,U}$ identifies with the mapping $\lambda_{X,U}$ defined (for all left $\mathbb{k}G$ -modules X and U) by

$$(3.2.2) \quad \begin{aligned} X^* \otimes_G U & \rightarrow \text{Hom}_G(X, U) \\ \phi \otimes u & \mapsto (x \mapsto \sum_{g \in G} \phi(g^{-1} \cdot x)g \cdot u). \end{aligned}$$

Now, $\lambda_{\mathbb{k}G, \mathbb{k}G}$ identifies with the following bijective mapping (recall that $|G| \in \mathbb{k}^\times$).

$$\begin{aligned} \mathbb{k}G^* \otimes_G \mathbb{k}G & \rightarrow \mathbb{k}G \\ \phi \otimes e & \mapsto \sum_{g \in G} \phi(g^{-1})g. \end{aligned}$$

Since ${}_G X$ is finite dimensional and projective, and since ${}_G U$ is projective, it follows that $\lambda_{X,U}$ is bijective. Thus, $\Psi_X^{V,U}$ is bijective. $\square$

Lemma 3.3. *Let G, H, K be finite groups with orders not divisible by $\text{char}(\mathbb{k})$. Let ${}_G X_H$, ${}_H Y_K$, ${}_K W$ and ${}_G U$ be finite dimensional modules. Then, the following diagram commutes*

$$(3.2.3) \quad \begin{array}{ccc} \text{Hom}_K(W, Y^* \otimes_H X^* \otimes_G U) & \xrightarrow{\quad} & \text{Hom}_K(W, (X \otimes_H Y)^* \otimes_G U) \\ \Psi_Y^{W, X^* \otimes_G U} \downarrow & & \downarrow \Psi_{X \otimes_H Y}^{W, U} \\ \text{Hom}_H(Y \otimes_K W, X^* \otimes_G U) & \xrightarrow[\Psi_X^{Y \otimes_K W, U}]{} & \text{Hom}_G(X \otimes_H Y \otimes_K W, U), \end{array}$$

where the top horizontal arrow is obtained upon applying $\text{Hom}_K(W, - \otimes_G U)$ to (3.1.1).

Proof. In view of the compatibility of the involved mappings with direct sum decompositions of the finite dimensional projective modules ${}_K W$ and ${}_G U$, it is sufficient to prove the lemma assuming that $W = \mathbb{k}K$ and $U = \mathbb{k}G$. In this case, the given diagram identifies with the following one

$$(3.2.4) \quad \begin{array}{ccc} Y^* \otimes_H X^* & \xrightarrow{(3.1.1)} & (X \otimes_H Y)^* \\ \beta \downarrow & & \downarrow \alpha \\ \text{Hom}_H(Y, X^*) & \xrightarrow{\gamma} & \text{Hom}_G(X \otimes_H Y, \mathbb{k}G), \end{array}$$

where, after using (3.2.2),

- α is given by $\phi \mapsto (x \otimes y \mapsto \sum_{g \in G} \phi(g^{-1} \cdot x \otimes y)g)$,
- β is given by $\phi_2 \otimes \phi_1 \mapsto (y \mapsto \sum_{h \in H} \phi_2(h^{-1} \cdot y) \underbrace{h \cdot \phi_1}_{\phi_1(\bullet h)})$ and
- γ is given by $f \mapsto (x \otimes y \mapsto \sum_{g \in G} (f(y))(g^{-1} \cdot x)g)$.

It is direct to check that (3.2.4) commutes, which proves the lemma. $\square$

3.3. Pairing of morphisms.

Definition 3.4. Let G and H be finite groups whose orders are not divisible by $\text{char}(\mathbb{k})$. Let ${}_G X_H$, ${}_H V$, and ${}_G U$ be (bi)modules such that X and V are finite dimensional. Assume that U is simple. Define a pairing

$$(3.3.1) \quad \langle - | - \rangle : \text{Hom}_G(U, X \otimes_H V) \otimes_{\mathbb{k}} \text{Hom}_H(V, X^* \otimes_G U) \rightarrow \mathbb{k}$$

as follows. For all $f \in \text{Hom}_G(U, X \otimes_H V)$ and $\phi \in \text{Hom}_H(V, X^* \otimes_G U)$,

$$(3.3.2) \quad \langle f | \phi \rangle \cdot \text{Id}_U = \Psi_X^{V,U}(\phi) \circ f$$

(following Lemma 3.2, the pairing $\langle - | - \rangle$ is non-degenerate).

This pairing is compatible with the following operation on morphisms.

Definition 3.5. Let G , H and K be finite groups whose orders are not divisible by $\text{char}(\mathbb{k})$. Let ${}_G X_H$, ${}_H Y_K$, ${}_G U$, ${}_H V$ and ${}_K W$ be (bi)modules. For all $f_1 \in \text{Hom}_G(U, X \otimes_H V)$ and $f_2 \in \text{Hom}_H(V, Y \otimes_K W)$, define $f_2 \times f_1 \in \text{Hom}_G(U, X \otimes_H Y \otimes_K W)$ as the following composite morphism

$$(3.3.3) \quad f_2 \times f_1 : U \xrightarrow{f_1} X \otimes_H V \xrightarrow{(X \otimes_H f_2)} X \otimes_H Y \otimes_K W.$$

Here is the above-mentioned compatibility.

Lemma 3.6. *Let*

- G , H , and K be finite groups with orders not divisible by $\text{char}(\mathbb{k})$,
- ${}_G X_H$, ${}_H Y_K$, ${}_G U$, ${}_H V$ and ${}_K W$ be (bi)modules such that X , Y , and W are finite dimensional and U and V are simple,
- $f_1 \in \text{Hom}_G(U, X \otimes_H V)$ and $f_2 \in \text{Hom}_H(V, Y \otimes_K W)$ and
- $\phi_1 \in \text{Hom}_H(V, X^* \otimes_G U)$ and $\phi_2 \in \text{Hom}_K(W, Y^* \otimes_H V)$.

Denote by λ the following isomorphism obtained upon applying $- \otimes_G U$ to (3.1.1),

$$Y^* \otimes_H X^* \otimes_G U \xrightarrow{\sim} (X \otimes_H Y)^* \otimes_G U.$$

Then, $\langle f_2 \times f_1 | \lambda \circ (\phi_1 \times \phi_2) \rangle = \langle f_2 | \phi_2 \rangle \cdot \langle f_1 | \phi_1 \rangle$.

Proof. Using (3.2.3) yields that

$$\Psi_{X \otimes_H Y}^{W,U}(\lambda \circ (\phi_1 \times \phi_2)) = \Psi_X^{Y \otimes_K W, U} \circ \Psi_Y^{W, X^* \otimes_G U}(\phi_1 \times \phi_2).$$

Recall that $\phi_1 \times \phi_2$ is the following composite morphism

$$W \xrightarrow{\phi_2} Y^* \otimes_H V \xrightarrow{Y^* \otimes_H \phi_1} Y^* \otimes_H X^* \otimes_G U.$$

Since $\Psi_X^{\bullet, \bullet}$ and $\Psi_Y^{\bullet, \bullet}$ are adjunction isomorphisms, then

$$\begin{aligned} \Psi_X^{Y \otimes_K W, U} \circ \Psi_Y^{W, X^* \otimes_G U}(\phi_1 \times \phi_2) &= \Psi_X^{Y \otimes_K W, U}(\phi_1 \circ \Psi_Y^{W, V}(\phi_2)) \\ &= \Psi_X^{V, U}(\phi_1) \circ (X \otimes_H \Psi_Y^{W, V}(\phi_2)). \end{aligned}$$

Therefore,

$$\begin{aligned} \Psi_{X \otimes_H Y}^{W,U}(\lambda \circ (\phi_1 \times \phi_2)) \circ (f_2 \times f_1) &= \Psi_X^{V, U}(\phi_1) \circ (X \otimes_H \Psi_Y^{W, V}(\phi_2)) \circ \\ &\quad (X \otimes_H f_2) \circ f_1 \\ &= \Psi_X^{V, U}(\phi_1) \circ (X \otimes_H \langle f_2 | \phi_2 \rangle \cdot \text{Id}_V) \circ f_1 \\ &= \langle f_1 | \phi_1 \rangle \cdot \langle f_2 | \phi_2 \rangle \cdot \text{Id}_U. \end{aligned}$$

The conclusion of the lemma then follows from the definition of $\langle - | - \rangle$. $\square$

The previous lemma has the following generalisation. Consider

- an integer $n \geq 2$,
- finite groups $G_0, \dots, G_n$ with orders not divisible by $\text{char}(\mathbb{k})$,
- a finite dimensional left $\mathbb{k}G_i$ -module U_i for every $i \in \{0, \dots, n\}$, such that $U_0, \dots, U_{n-1}$ are simple,
- a finite dimensional $\mathbb{k}G_{i-1} - \mathbb{k}G_i$ -bimodule X_i for every $i \in \{1, \dots, n\}$,
- $f_i \in \text{Hom}_{G_{i-1}}(U_{i-1}, X_i \otimes_{G_i} U_i)$ and $\phi_i \in \text{Hom}_{G_i}(U_i, X_i^* \otimes_{G_i} U_{i-1})$ for every $i \in \{1, \dots, n\}$.

Denote by $\lambda^{(n)}$ the isomorphism obtained upon applying $- \otimes_{G_0} U_0$ to (3.1.2),

$$\lambda^{(n)}: X_n^* \otimes_{G_{n-1}} \cdots \otimes_{G_1} X_1^* \otimes_{G_0} U_0 \xrightarrow{\sim} (X_1 \otimes_{G_1} \cdots \otimes_{G_{n-1}} X_n)^* \otimes_{G_0} U_0.$$

Then,

$$(3.3.4) \quad \langle f_n \times \cdots \times f_1 | \lambda^{(n)} \circ (\phi_1 \times \cdots \times \phi_n) \rangle = \prod_{t=1}^n \langle f_t | \phi_t \rangle.$$

4. INTERTWINERS

Let Q_0 , S , G and M be as in Setting 1.1. This section establishes properties on intertwiners relative to M which are needed to prove Theorem 1.25. This includes the properties of $\otimes$ mentioned in Section 1 as well as Proposition 1.20 on the existence of the pairing $\langle - | - \rangle$. This is done in several steps,

- first, by proving that, in Definition 1.8, the $\mathbb{k}G_{i_0}$ -module $M(\underline{i}; V)$ is isomorphic to $M(\underline{i}) \otimes_{G_{i_n}} V$ for a suitable $\mathbb{k}G_{i_0} - \mathbb{k}G_{i_n}$ -bimodule $M(\underline{i})$;
- next, by proving that, under this isomorphism, $\otimes$ corresponds to the operation $\times$ of Definition 3.5;
- and finally by defining $\langle - | - \rangle$ so that, under this isomorphism, it corresponds to the pairing $\langle - | - \rangle$ of Definition 3.4.

These steps are proceeded in turn in Sections 4.1, 4.2, and 4.3, respectively. They use the bimodule MG defined as follows. Note that the action of G on S yields the skew group algebra $S * G$. Like in Section 3, for all groups H , the pieces of notation e , $\otimes_H$ and Hom_H stand for the neutral element of H , for $\otimes_{\mathbb{k}H}$, and for $\text{Hom}_{\mathbb{k}H}$, respectively.

Definition 4.1. Define MG as the following $S * G - S * G$ -bimodule.

- Its underlying vector space is $M \otimes_{\mathbb{k}} \mathbb{k}G$, a tensor $m \otimes g$ is denoted by $m * g$ for all $m \in M$ and $g \in G$.
- The actions of S and G on the left and on the right are such that, for all $m \in M$, $g, h, k \in G$ and $s, s' \in S$,

$$\begin{aligned} s \cdot (m * g) \cdot s' &= sm^g s' * g \\ h \cdot (m * g) \cdot k &= {}^h m * h g k. \end{aligned}$$

The $S * G - S * G$ -bimodule M^*G is defined similarly after replacing M by M^* .

4.1. The space $M(\underline{i})$. Let n be a positive integer. Let $\underline{i} = i_0, \dots, i_n$ a sequence in Q_0 . Denote by $M(\underline{i})$ the following $\mathbb{k}G_{i_0} - \mathbb{k}G_{i_n}$ -bimodule,

$$(4.1.1) \quad M(\underline{i}) = e_{i_0} M G e_{i_1} \otimes_{G_{i_1}} \cdots \otimes_{G_{i_{n-1}}} e_{i_{n-1}} M G e_{i_n}.$$

Note that this yields the $\mathbb{k}G_{i_n} - \mathbb{k}G_{i_0}$ -bimodule $M^*(\underline{i}^o)$. These bimodules are dual to each other. Here is an explicit isomorphism used later. Consider the isomorphism (3.1.2) taking $X_t = e_{i_{t-1}} M^* G e_{i_t}$ for all $t \in \{1, \dots, n\}$,

$$(4.1.2) \quad e_{i_n} (MG)^* e_{i_{n-1}} \otimes_{G_{i_{n-1}}} \cdots \otimes_{G_{i_1}} e_{i_1} (MG)^* e_{i_0} \xrightarrow{(3.1.2)} M(\underline{i})^*.$$

Since the following mapping is a non-degenerate pairing

$$\begin{aligned} [-|-]: \quad MG \otimes_{\mathbb{k}} M^*G &\rightarrow \mathbb{k} \\ (m * g) \otimes (\varphi * h) &\mapsto \varphi(g^{-1}m)\delta_{e,hg}, \end{aligned}$$

it induces an isomorphism (of $S * G - S * G$ -bimodules)

$$(4.1.3) \quad \begin{aligned} M^*G &\rightarrow (MG)^* \\ \bullet &\mapsto [-|\bullet] \end{aligned}$$

the n -th tensor power of which induces an isomorphism of $\mathbb{k}G_{i_n} - \mathbb{k}G_{i_0}$ -bimodules,

$$(4.1.4) \quad M^*(\underline{i}^o) \xrightarrow{\sim} e_{i_n}(MG)^* e_{i_{n-1}} \otimes_{G_{i_{n-1}}} \cdots \otimes_{G_{i_1}} e_{i_1}(MG)^* e_{i_0}.$$

Composing (4.1.2) and (4.1.4) yields an isomorphism of $\mathbb{k}G_{i_n} - \mathbb{k}G_{i_0}$ -bimodules

$$(4.1.5) \quad \Lambda_{\underline{i}}: M^*(\underline{i}^o) \xrightarrow{\sim} M(\underline{i})^*,$$

which is induced by the mapping from $(MG)^{\otimes_k n}$ to $((MG)^{\otimes_k n})^*$ which assigns to any tensor $(\varphi_n * g_n) \otimes \cdots \otimes (\varphi_1 * g_1)$ the following linear form on $(MG)^{\otimes_k n}$,

$$(4.1.6) \quad (m_1 * g'_1) \otimes \cdots \otimes (m_n * g'_n) \mapsto \sum_{k_1, \dots, k_{n-1}} \prod_{t=1}^n \varphi_t(k_t^{-1} g'^{-1}_{t-1} m_t) \delta_{e, g_t k_{t-1}^{-1} g'_t k_t},$$

where δ denotes the Kronecker symbol, k_t runs through G_{i_t} and $k_n = k_0 = e$.

The bimodules $M(\underline{i})$ and $M^*(\underline{i}^o)$ are hence objects of the monoidal category $(\text{mod}(A^e), \otimes_A)$ in which they are left dual to each other. The following lemma describes tensor products of these objects in this monoidal category. The same statements are valid after replacing M by M^* .

Lemma 4.2. *Let m, n be positive integers. Let $\underline{i}'' = i_0, \dots, i_{m+n}$ be a sequence in Q_0 . Denote the sequences $i_0, \dots, i_m$ and $i_m, \dots, i_{m+n}$ by $\underline{i}$ and $\underline{i}'$, respectively.*

- (1) $M(\underline{i}'') = M(\underline{i}) \otimes_{G_{i_m}} M(\underline{i}')$.
- (2) *The following diagram is commutative*

$$\begin{array}{ccc} M^*(\underline{i}'^o) \otimes_{G_{i_m}} M^*(\underline{i}^o) & \xrightarrow{\Lambda_{\underline{i}'} \otimes_{G_{i_m}} \Lambda_{\underline{i}}} & M(\underline{i}')^* \otimes_{G_{i_m}} M(\underline{i})^* \\ \parallel & & \downarrow (3.1.1) \\ M^*(\underline{i}''^o) & \xrightarrow{\Lambda_{\underline{i}''}} & M(\underline{i}'')^*. \end{array}$$

Proof. (1) follows from the definition of the spaces $M(\bullet)$ and (2) follows from the definition of the morphisms $\Lambda_{\bullet}$ in terms of (3.1.2) and (4.1.3). $\square$

The following result is the link between the bimodules $M(\bullet)$ and the modules of Definition 1.8. This link makes it possible to translate the results of Section 3 in terms of intertwiners. From now on, the piece of notation $*$ used for tensors in MG is often dropped in order to avoid overloaded notation.

Lemma 4.3. *Let $\underline{i} = i_0, \dots, i_n$ be a sequence in Q_0 , where $n \geq 1$. Let $U_n \in \text{mod}(\mathbb{k}G_{i_n})$. There is an isomorphism in $\text{mod}(\mathbb{k}G_{i_0})$,*

$$(4.1.7) \quad \Theta_{\underline{i}; U_n}^M: M(\underline{i}; U_n) \xrightarrow{\sim} M(\underline{i}) \otimes_{G_{i_n}} U_n$$

given by

$$m_1 \otimes \cdots \otimes m_n \otimes y_1 \cdots y_n u \mapsto m_1 y_1 \otimes y_1^{-1} m_2 y_2 \otimes \cdots \otimes (y_1 \cdots y_{n-1})^{-1} m_n y_n \otimes u.$$

When there is no risk of confusion, $\Theta_{\underline{i}; U_n}^M$ is denoted by $\Theta_{\underline{i}; U_n}$.

Proof. This is straightforward to check. $\square$

The following natural transformation of functors from $\text{mod}(\mathbb{k}G_{i_0})$ to $\text{mod}(\mathbb{k})$, defined by $\Theta_{\underline{i}; U_n}$, is hence an isomorphism of functors,

$$(4.1.8) \quad \begin{array}{ccc} \text{Hom}_{G_{i_0}}(-, M(\underline{i}; U_n)) & \xrightarrow{\sim} & \text{Hom}_{G_{i_0}}(-, M(\underline{i}) \otimes_{G_{i_n}} U_n) \\ - & \mapsto & \Theta_{\underline{i}; U_n} \circ - \end{array}$$

4.2. Compositions of intertwiners. It is now possible to compare the operation $\circledast$ on intertwiners, see Definition 1.13, and the operation $\times$ of Definition 3.5.

Lemma 4.4. *Let $\underline{i}'' = i_0, \dots, i_{m+n}$ be a sequence in Q_0 , where $m, n \geq 1$. Denote the sequences $i_0, \dots, i_m$ and $i_m, i_{m+1}, \dots, i_{m+n}$ by $\underline{i}$ and $\underline{i}'$, respectively. Let $U_0 \in \text{mod}(\mathbb{k}G_{i_0})$, $U_m \in \text{mod}(\mathbb{k}G_{i_m})$ and $U_{m+n} \in \text{mod}(\mathbb{k}G_{i_{m+n}})$. Finally, let $f \in \text{Hom}_{\mathbb{k}G_{i_0}}(U_0, M(\underline{i}; U_m))$ and $f' \in \text{Hom}_{\mathbb{k}G_{i_m}}(U_m, M(\underline{i}'; U_{m+n}))$. Then,*

$$(4.2.1) \quad (\Theta_{\underline{i}'; U_{m+n}} \circ f') \times (\Theta_{\underline{i}; U_m} \circ f) = \Theta_{\underline{i}''; U_{m+n}} \circ (f' \circledast f).$$

As a consequence, $f' \circledast f \in \text{Hom}_{G_{i_0}}(U_0, M(\underline{i}''; U_{m+n}))$.

Proof. Let $u \in U_0$. Since $\Theta_{\underline{i}'; U_{m+n}} \circ f'$ lies in $\text{Hom}_{G_{i_m}}(U_m, M(\underline{i}''; U_{m+n}) \otimes_{G_{i_{m+n}}} U_{m+n})$ and $\Theta_{\underline{i}; U_m} \circ f$ lies in $\text{Hom}_{G_{i_0}}(U_0, M(\underline{i}) \otimes_{G_{i_m}} U_m)$, then

$$((\Theta_{\underline{i}'; U_{m+n}} \circ f') \times (\Theta_{\underline{i}; U_m} \circ f))(u) \stackrel{(3.3.3)}{=} (M(\underline{i}) \otimes_{\mathbb{k}G_{i_m}} (\Theta_{\underline{i}'; U_{m+n}} \circ f')) \circ (\Theta_{\underline{i}; U_m} \circ f)(u).$$

Using the definition of $\Theta_{\underline{i}; U_m}$, see (4.1.7), the right-hand side is equal to

$$\begin{aligned} & (M(\underline{i}) \otimes_{\mathbb{k}G_{i_m}} (\Theta_{\underline{i}'; U_{m+n}} \circ f')) (\sum_{\underline{y}} f_{\underline{y}}^{(1)}(u) y_1 \otimes y_1^{-1} f_{\underline{y}}^{(2)}(u) y_2 \otimes \dots \\ & \dots \otimes (y_1 \dots y_{m-1})^{-1} (f_{\underline{y}}^{(m)}(u)) y_m \otimes f_{\underline{y}}^{(0)}(u)), \end{aligned}$$

where $\underline{y}$ runs through all sequences $y_1, \dots, y_m$ such that $y_t \in [G/G_{i_t}]$ for all $t \in \{1, \dots, m\}$. Using the definition of $\Theta_{\underline{i}'; U_{m+n}}$, see (4.1.7), this is equal to

$$\begin{aligned} & \sum_{\underline{y}, \underline{y}'} f_{\underline{y}}^{(1)}(u) y_1 \otimes \dots \otimes (y_1 \dots y_{m-1})^{-1} (f_{\underline{y}}^{(m)}(u)) y_m \otimes \\ & f_{\underline{y}'}^{(1)}(f_{\underline{y}}^{(0)}(u)) y_{m+1} \otimes y_{m+1}^{-1} (f_{\underline{y}'}^{(2)}(f_{\underline{y}}^{(0)}(u))) y_{m+2} \otimes \dots \\ & \dots \otimes (y_{m+1} \dots y_{m+n})^{-1} (f_{\underline{y}'}^{(n)}(f_{\underline{y}}^{(0)}(u))) y_{m+n} \otimes f_{\underline{y}'}^{(0)}(f_{\underline{y}}^{(0)}(u)), \end{aligned}$$

where $\underline{y}$ is as before and $\underline{y}'$ runs through all sequences $y_{m+1}, \dots, y_{m+n}$ such that $y_{m+t} \in [G/G_{i_{m+t}}]$ for all $t \in \{1, \dots, n\}$. Using the definition of $\Theta_{\underline{i}''; U_{m+n}}$, see (4.1.7), this is equal to

$$\begin{aligned} & \Theta_{\underline{i}''; U_{m+n}} (\sum_{\underline{y}''} f_{\underline{y}''}^{(1)}(u) \otimes \dots \otimes f_{\underline{y}''}^{(m)}(u) \otimes y_1 \dots y_m (f_{\underline{y}'}^{(1)}(f_{\underline{y}}^{(0)}(u))) \otimes \dots \\ & \dots \otimes y_1 \dots y_m (f_{\underline{y}'}^{(n)}(f_{\underline{y}}^{(0)}(u))) \otimes y_1 \dots y_{m+n} f_{\underline{y}'}^{(0)}(f_{\underline{y}}^{(0)}(u))), \end{aligned}$$

where $\underline{y}''$ runs through all sequences $y_1, \dots, y_{m+n}$ such that $y_t \in [G/G_{i_t}]$ for all $t \in \{1, \dots, m+n\}$ and where $\underline{y}$ and $\underline{y}'$ stand for the sequences $y_1, \dots, y_m$ and $y_{m+1}, \dots, y_{m+n}$, respectively. Finally, using the definition of $\circledast$, see (1.4.1), this is equal to $(\Theta_{\underline{i}''; U_{m+n}} \circ (f' \circledast f))(u)$. Whence (4.2.1). Note that $f' \circledast f$ is a morphism of $\mathbb{k}G_{i_0}$ -modules because

- $\Theta_{\underline{i}''; U_{m+n}}$ is an isomorphism of $\mathbb{k}G_{i_0}$ -modules,
- f and $\Theta_{\underline{i}; U_m}$ are morphisms of $\mathbb{k}G_{i_0}$ -modules,
- f' and $\Theta_{\underline{i}'; U_{m+n}}$ are morphisms of $\mathbb{k}G_{i_m}$ -modules,
- and the operation “ $\times$ ” yields morphisms of $\mathbb{k}G_{i_0}$ -modules (see (3.3.3)).

□

Lemma 4.4 shows that Definition 1.15 yields a, possibly non-associative, algebra.

Proposition 4.5. *Intw is an associative and unital algebra.*

Proof. Note that $\mathcal{I}ntw$ is unital. Indeed, for all $i \in [G \setminus Q_0]$ and $U \in \text{irr}(G_i)$, denote by $\varepsilon_{i,U}$ the intertwiner $\text{Id}_U \in \text{Hom}_{\mathbb{k}G_i}(U, M(i; U))$, recall that $M(i; U) = U$. Then $\sum_{i,U} \varepsilon_{i,U}$ is a unity for $\otimes$.

In the framework of Definition 3.5, the equality $f_3 \times (f_2 \times f_1) = (f_3 \times f_2) \times f_1$ holds whenever both sides make sense because of the associativity of the tensor product. In view of Lemma 4.4, this entails that $\otimes$ is associative. $\square$

4.3. The pairing on intertwiners. Let n be a positive integer, let $\underline{i} = i_0, \dots, i_n$ be a sequence in Q_0 and let U_0 and U_n be left modules over $\mathbb{k}G_{i_0}$ and $\mathbb{k}G_{i_n}$, respectively. To every $\varphi \in \text{Hom}_{G_{i_n}}(U_n, M^*(\underline{i}^o; U_0))$ is associated a $\mathbb{k}$ -linear mapping

$$\Phi(\varphi): M(\underline{i}; U_n) \rightarrow U_0$$

defined as follows. For every sequence of representatives $y_1, \dots, y_n$ ($y_t \in [G/G_{i_t}]$) there exist unique sequences

$$\underline{x} = x_{n-1}, \dots, x_0 \quad (x_t \in [G/G_{i_t}]), \text{ and } h_{n-1}, \dots, h_0 \quad (h_t \in G_{i_t})$$

such that, for all $t \in \{1, \dots, n\}$,

$$(4.3.1) \quad (y_t \cdots y_n)^{-1} = x_{n-1} \cdots x_{t-1} h_{t-1};$$

these sequences are determined by a decreasing induction: $y_n^{-1} = x_{n-1} h_{n-1}$ and $h_t y_t^{-1} = x_{t-1} h_{t-1}$ for all $t \in \{1, \dots, n-1\}$; then, for every $m_1 \otimes \cdots \otimes m_t \otimes y_1 \cdots y_n u$ lying in $M(\underline{i}; U_n)$, define the element $\Phi(\varphi)(m_1 \otimes \cdots \otimes m_t \otimes y_1 \cdots y_n u)$ of U_0 , using Notation 1.18, as

$$(4.3.2) \quad \prod_{t=1}^n \varphi_{(t)}^{\underline{x}}(u) \left((y_1 \cdots y_n)^{-1} m_t \right)^{h_0^{-1}} (\varphi_{(0)}^{\underline{x}}(u)).$$

This does make sense, because $m_t \in {}_{y_1 \cdots y_{t-1} \cdot i_{t-1}} M_{y_1 \cdots y_t \cdot i_t}$, because $\varphi_{(t)}^{\underline{x}}$ is a linear form on ${}_{x_{n-1} \cdots x_{t-1} \cdot i_{t-1}} M_{x_{n-1} \cdots x_t \cdot i_t}$, and because of (4.3.1).

After appropriate identifications, Φ is related to the adjunction isomorphism $\Psi_{M(\underline{i})}^{\bullet, \bullet}$ of section 3.2 as explained in the following result. Recall that $\Psi_{M(\underline{i})}^{U_n, U_0}$, $\Theta_{\underline{i}^o; U_0}^{M^*}$ and $\Lambda_{\underline{i}}$ are defined in (3.2.1), (4.1.7), and (4.1.5), respectively.

Lemma 4.6. *Keep the previous setting. Denote by φ' the following composite morphism of left $\mathbb{k}G_{i_n}$ -modules,*

$$U_n \xrightarrow{\varphi} M^*(\underline{i}^o; U_0) \xrightarrow{\Theta_{\underline{i}^o; U_0}^{M^*}} M^*(\underline{i}^o) \otimes_{G_{i_0}} U_0 \xrightarrow{\Lambda_{\underline{i}} \otimes_{G_{i_0}} U_0} M(\underline{i})^* \otimes_{G_{i_0}} U_0.$$

Then,

$$(4.3.3) \quad \Phi(\varphi) = \Psi_{M(\underline{i})}^{U_n, U_0}(\varphi') \circ \Theta_{\underline{i}; U_n}^M.$$

In particular, $\Phi(\varphi) \in \text{Hom}_{G_{i_0}}(M(\underline{i}; U_n), U_0)$.

Proof. The composition $\Theta_{\underline{i}; U_0}^{M^*} \circ \varphi$ is given by (see (4.1.7))

$$u \mapsto \sum_{\underline{x}} \varphi_{(n)}^{\underline{x}}(u) x_{n-1} \otimes {}^{x_{n-1}^{-1}} \varphi_{(n-1)}^{\underline{x}}(u) x_{n-2} \otimes \cdots \otimes ({}^{x_{n-1} \cdots x_1})^{-1} \varphi_{(1)}^{\underline{x}}(u) x_0 \otimes \varphi_{(0)}^{\underline{x}}(u),$$

where $\underline{x} = x_{n-1}, \dots, x_0$ runs through all sequences of representatives ($x_t \in [G/G_{i_t}]$) for all $0 \leq t \leq n-1$. Hence, following (4.1.6), the morphism φ' is given by

$$u \mapsto \sum_{\underline{x}} \phi^{\underline{x}}(u) \otimes \varphi_{(0)}^{\underline{x}}(u)$$

where $\phi^{\underline{x}}(u)$ is the linear form on $M(\underline{i})$, which is a quotient of $(MG)^{\otimes_{\mathbb{k}} n}$, induced by the linear form on $(MG)^{\otimes_{\mathbb{k}} n}$ mapping any $(m_1 * g_1) \otimes \cdots \otimes (m_n * g_n)$ to

$$(4.3.4) \quad \sum_{k_1, \dots, k_{n-1}} \prod_{t=1}^n \left((x_{n-1} x_{n-2} \cdots x_t)^{-1} (\varphi_{(t)}^{\underline{x}}(u)) \right) \left(k_t^{-1} g_t^{-1} m_t \right) \cdot \delta_{e, x_{t-1} k_{t-1}^{-1} g_t k_t},$$

where δ denotes the Kronecker symbol, $k_0 = k_n = e$ and k_t runs through G_{i_t} for $1 \leq t \leq n-1$.

In order to simplify the expression of (4.3.4), consider a tensor $(m_1 * g_1) \otimes \cdots \otimes (m_n * g_n)$ in $(MG)^{\otimes_{\mathbb{k}} n}$; for all $k_1, \dots, k_{n-1}$ in G , the following assertions are equivalent, keeping $k_0 = k_n = e$,

- (i) $(\forall t \in \{1, \dots, n-1\}) e = x_{t-1} k_{t-1}^{-1} g_t k_t$,
- (ii) $k_t = (x_{t-1} \cdots x_1 x_0 g_1 g_2 \cdots g_t)^{-1}$ for all $t \in \{1, \dots, n\}$, and $g_1 \cdots g_n$ is the inverse of $x_{n-1} \cdots x_1 x_0$;

given that each $(x_{n-1} x_{n-2} \cdots x_t)^{-1} (\varphi_{(t)}^{\underline{x}}(u))$ in (4.3.4) is a linear form on $x_{t-1} i_{t-1} M_{i_t}$ and each k_t is constrained to lie in G_{i_t} , then (4.3.4) is equal to

$$\prod_{t=1}^n (\varphi_{(t)}^{\underline{x}}(u)) (x_{n-1} \cdots x_1 x_0 g_1 g_2 \cdots g_{t-1} m_t) \delta_{e, g_1 g_2 \cdots g_n x_{n-1} \cdots x_1 x_0},$$

which simplifies to

$$(4.3.5) \quad \prod_{t=1}^n (\varphi_{(t)}^{\underline{x}}(u)) \left((g_t g_{t+1} \cdots g_n)^{-1} m_t \right) \delta_{(g_1 g_2 \cdots g_n)^{-1}, x_{n-1} \cdots x_1 x_0}.$$

Thus, the morphism φ' is given by $u \mapsto \sum_{\underline{x}} \phi^{\underline{x}}(u) \otimes \varphi_{(0)}^{\underline{x}}(u)$, where $\phi^{\underline{x}}(u)$ is the linear form on $M(\underline{i})$ induced by the mapping assigning (4.3.5) to any element $(m_1 * g_1) \otimes \cdots \otimes (m_n * g_n)$ of $(MG)^{\otimes_{\mathbb{k}} n}$.

Hence (see Lemma 3.2), $\Psi_{M(\underline{i})}^{U_n, U_0}(\varphi')$ is the morphism from $M(\underline{i}) \otimes_{G_{i_n}} U_n$ to U_0 induced by the mapping from $(MG)^{\otimes_{\mathbb{k}} n} \otimes_{\mathbb{k}} U_n$ to U_0 which maps any $(m_1 * g_1) \otimes \cdots \otimes (m_n * g_n) \otimes u$ to

$$\begin{aligned} & \sum_{\underline{x}} \sum_{g \in G_{i_0}} \phi^{\underline{x}}(u) (g^{-1} m_1 g_1 \otimes m_2 g_2 \otimes \cdots \otimes m_n g_n) {}^g(\varphi_{(0)}^{\underline{x}}(u)) \\ &= \sum_{g, \underline{x}} \prod_{t=1}^n \varphi_{(t)}^{\underline{x}}(u) ((g_t \cdots g_n)^{-1} m_t) \delta_{(g_1 \cdots g_n)^{-1} g, x_{n-1} \cdots x_0} {}^g(\varphi_{(0)}^{\underline{x}}(u)). \end{aligned}$$

Now, let $m_1 \otimes \cdots \otimes m_n \otimes y_1 \cdots y_n u \in M(\underline{i}; U_n)$. Recall that its image under $\Theta_{\underline{i}; U_n}^M$ is $m_1 y_1 \otimes y_1^{-1} m_2 y_2 \otimes \cdots \otimes (y_1 \cdots y_{n-1})^{-1} m_n y_n \otimes u$ (see (4.1.7)). Hence, its image under $\Psi_{M(\underline{i})}^{U_n, U_0}(\varphi') \circ \Theta_{\underline{i}; U_n}^M$ is

$$(4.3.6) \quad \sum_{g, \underline{x}} \prod_{t=1}^n \varphi_{(t)}^{\underline{x}}(u) ((y_1 \cdots y_n)^{-1} m_t) \delta_{(y_1 \cdots y_n)^{-1} g, x_{n-1} \cdots x_0} {}^g(\varphi_{(0)}^{\underline{x}}(u)).$$

Note that, given $g \in G_{i_0}$ and $\underline{x}$, then, for all $t \in \{1, \dots, n\}$,

- $\varphi_{(t)}^{\underline{x}}(u) \in (x_{n-1} \cdots x_{t-1} i_{t-1} M_{x_{n-1} \cdots x_t i_t})^*$ and
- $(y_1 \cdots y_n)^{-1} m_t \in (y_t \cdots y_n)^{-1} i_{t-1} M_{(y_{t+1} \cdots y_n)^{-1} i_t}$;

in particular, if for a given index $(g, \underline{x})$ in (4.3.6) the product “ $\prod_{t=1}^n$ ” is non-zero then this product is the unique non-zero term of the sum (4.3.6). More precisely, let $\{x_t, h_t\}_{0 \leq t \leq n-1}$ be the unique collection of elements in G such that, for all $t \in \{0, \dots, n-1\}$,

- $x_t \in [G/G_{i_t}]$,
- $h_t \in G_{i_t}$ and
- $(y_{t+1} \cdots y_n)^{-1} = x_{n-1} \cdots x_t h_t$;

then, the only possibly non-zero term of the sum (4.3.6) is the one with index $(h_0^{-1}, \underline{x})$. Therefore, (4.3.6) equals

$$\prod_{t=1}^n \varphi_{(t)}^{\underline{x}}(u) (y_1 \cdots y_n)^{-1} m_t)^{h_0^{-1}} (\varphi_{(0)}^{\underline{x}}(u)),$$

which is equal to $\Phi(\varphi)(m_1 \otimes \cdots \otimes m_n \otimes y_1 \cdots y_n u)$ (see (4.3.2)). Thus, $\Phi(\varphi) = \Psi_{M(\underline{i})}^{U_n, U_0}(\varphi') \circ \Theta_{\underline{i}; U_n}^M$. This proves (4.3.3). The second assertion of the lemma is a direct consequence of (4.3.3). $\square$

It follows from Lemma 4.6 that the following mapping is bijective,

$$\Phi: \text{Hom}_{G_{i_n}}(U_n, M^*(\underline{i}^o; U_0)) \rightarrow \text{Hom}_{G_{i_0}}(M(\underline{i}; U_n), U_0).$$

This yields a non-degenerate pairing defined as follows.

Definition 4.7. Let $\underline{i} = i_0, \dots, i_n$ be a sequence in Q_0 , where $n \geq 1$. Let $U_0 \in \text{mod}(\mathbb{k}G_{i_0})$ be simple. Let $U_n \in \text{mod}(\mathbb{k}G_{i_n})$. Define $(-|-)$ to be the non-degenerate pairing

$$(-|-): \text{Hom}_{G_{i_0}}(U_0, M(\underline{i}; U_n)) \otimes_{\mathbb{k}} \text{Hom}_{G_{i_n}}(U_n, M^*(\underline{i}^o; U_0)) \rightarrow \mathbb{k},$$

such that, for all $f \in \text{Hom}_{G_{i_0}}(U_0, M(\underline{i}; U_n))$ and $\varphi \in \text{Hom}_{G_{i_n}}(U_n, M^*(\underline{i}^o; U_0))$,

$$(4.3.7) \quad (f|\varphi) \cdot \text{Id}_{U_0} = \Phi(\varphi) \circ f;$$

With this definition, it follows from (3.3.2) and (4.3.3) that

$$(4.3.8) \quad (f|\varphi) = \langle \Theta_{\underline{i}; U_n}^M \circ f | (\Lambda_{\underline{i}} \otimes_{G_{i_0}} U_0) \circ \Theta_{\underline{i}^o; U_0}^{M^*} \circ \varphi \rangle.$$

It is now possible to prove Proposition 1.20.

Proof of Proposition 1.20. Note that (1.5.2) is equal to $(\Phi(\varphi))(f(u))$, see (4.3.2). Hence, the pairing $(-|-)$ claimed by the proposition is the one of Definition 4.7. $\square$

The pairing $(-|-)$ is compatible with the composition of (dual) intertwiners in the following sense.

Lemma 4.8. Let m, n be positive integers, let $\underline{i}'' = i_0, \dots, i_{m+n}$ be a sequence in Q_0 , let U_0 and U_m be simple left modules over $\mathbb{k}G_{i_0}$ and $\mathbb{k}G_{i_m}$, respectively, and let U_{m+n} be a left $\mathbb{k}G_{i_{m+n}}$ -module. Then, for all intertwiners

$$\begin{aligned} f_1: U_0 &\rightarrow M(\underline{i}; U_m), & \varphi_1: U_m &\rightarrow M^*(\underline{i}^o; U_0), \\ f_2: U_m &\rightarrow M(\underline{i}'; U_{m+n}), & \varphi_2: U_{m+n} &\rightarrow M^*(\underline{i}'^o; U_m), \end{aligned}$$

where $\underline{i} = i_0, \dots, i_m$ and $\underline{i}' = i_m, \dots, i_{m+n}$, the following holds

$$(4.3.9) \quad (f_2 \otimes f_1 | \varphi_1 \otimes \varphi_2) = (f_2 | \varphi_2) \cdot (f_1 | \varphi_1).$$

Proof. In order to avoid overloaded expressions, the following shorthand notation is used in this proof: $\Theta_{\underline{i}; U_0}$ for $\Theta_{\underline{i}; U_0}^M$ and $\Theta'_{\underline{i}; U_0}$ for $\Theta_{\underline{i}^o; U_0}^{M^*}$. Define $f'_1, f'_2, \varphi'_1, \varphi'_2$ as follows

$$\begin{aligned} f'_1 &= \Theta_{\underline{i}; U_m} \circ f_1, & \varphi'_1 &= (\Lambda_{\underline{i}} \otimes_{G_{i_0}} U_0) \circ \Theta'_{\underline{i}; U_0} \circ \varphi_1, \\ f'_2 &= \Theta_{\underline{i}'; U_{m+n}} \circ f_2, & \varphi'_2 &= (\Lambda_{\underline{i}'} \otimes_{G_{i_m}} U_m) \circ \Theta'_{\underline{i}'; U_m} \circ \varphi_2. \end{aligned}$$

It follows from (4.3.8) that $(f_2 \otimes f_1 | \varphi_1 \otimes \varphi_2)$ is equal to

$$\langle \Theta_{\underline{i}'', U_{m+n}} \circ (f_2 \otimes f_1) | (\Lambda_{\underline{i}''} \otimes_{G_{i_0}} U_0) \circ \Theta'_{\underline{i}'', U_0} \circ (\varphi_1 \otimes \varphi_2) \rangle;$$

applying Lemma 4.4 twice, first for M and next for M^* , yields that the previous expression is equal to

$$\underbrace{\langle \Theta_{\underline{i}', U_{m+n}} \circ f_2 \rangle}_{f'_1} \times \underbrace{\langle \Theta_{\underline{i}, U_m} \circ f_1 \rangle}_{f'_2} | (\Lambda_{\underline{i}''} \otimes_{G_{i_0}} U_0) \circ \left[(\Theta'_{\underline{i}, U_0} \circ \varphi_1) \times (\Theta'_{\underline{i}', U_m} \circ \varphi_2) \right];$$

according to Lemma 4.2 this is equal to

$$(4.3.10) \quad \langle f'_1 \times f'_2 | \lambda \circ (\Lambda_{\underline{i}'} \otimes_{G_{i_m}} \Lambda_{\underline{i}} \otimes_{G_{i_0}} U_0) \circ [(\Theta'_{\underline{i}, U_0} \circ \varphi_1) \times (\Theta'_{\underline{i}', U_m} \circ \varphi_2)] \rangle,$$

where λ is as in Lemma 3.6. Now, since the following composite morphisms are equal due to the associativity of the tensor product

$$M^*(\underline{i}'^o)_{G_{i_m}} \otimes U_m \xrightarrow{1 \otimes (\Theta'_{\underline{i}, U_0} \circ \varphi_1)} M^*(\underline{i}'^o)_{G_{i_m}} \otimes M^*(\underline{i}^o)_{G_{i_0}} \otimes U_0 \xrightarrow{\Lambda_{\underline{i}'} \otimes \Lambda_{\underline{i}} \otimes 1} M(\underline{i}')^*_{G_{i_m}} \otimes M(\underline{i})^*_{G_{i_0}} \otimes U_0$$

and

$$M^*(\underline{i}'^o)_{G_{i_m}} \otimes U_m \xrightarrow{\Lambda_{\underline{i}'} \otimes 1} M(\underline{i}')^*_{G_{i_m}} \otimes U_m \xrightarrow{1 \otimes (\Theta'_{\underline{i}, U_0} \circ \varphi_1)} M(\underline{i}')^*_{G_{i_m}} \otimes M^*(\underline{i}^o)_{G_{i_0}} \otimes U_0 \xrightarrow{1 \otimes \Lambda_{\underline{i}} \otimes 1} M(\underline{i}')^*_{G_{i_m}} \otimes M(\underline{i})^*_{G_{i_0}} \otimes U_0,$$

then $(\Lambda_{\underline{i}'} \otimes_{G_{i_m}} \Lambda_{\underline{i}} \otimes_{G_{i_0}} U_0) \circ [(\Theta'_{\underline{i}, U_0} \circ \varphi_1) \times (\Theta'_{\underline{i}', U_m} \circ \varphi_2)]$ is equal to

$$\underbrace{[(\Lambda_{\underline{i}} \otimes_{G_{i_0}} U_0) \circ \Theta'_{\underline{i}, U_0} \circ \varphi_1]}_{\varphi'_1} \times \underbrace{[(\Lambda_{\underline{i}'} \otimes_{G_{i_m}} U_m) \circ \Theta'_{\underline{i}', U_m} \circ \varphi_1]}_{\varphi'_2}.$$

Therefore,

$$(4.3.10) \quad \begin{aligned} &= \langle f'_2 \times f'_1 | \lambda \circ (\varphi'_1 \times \varphi'_2) \rangle \\ &\stackrel{\text{Lemma 3.6}}{=} \langle f'_2 | \varphi'_2 \rangle \cdot \langle f'_1 | \varphi'_1 \rangle \\ &\stackrel{(4.3.8)}{=} (f_2 | \varphi_2) \cdot (f_1 | \varphi_1). \end{aligned}$$

Thus, $(f_2 \otimes f_1 | \varphi_1 \otimes \varphi_2) = (f_1 | \varphi_1) \cdot (f_2 | \varphi_2)$. $\square$

5. PROOF OF THE MAIN RESULTS

This section proves Theorems 1.25 and 1.27. Keeping Q_0 , S , G and M as in Setting 1.1 and Q_G as in Setting 1.4, the proofs use the following notation:

- denote by $\hat{e}$ the idempotent $\sum_{i \in [G \setminus Q_0]} e_i$ of S , this idempotent is not to be confused with $\tilde{e}$ which is equal to $\sum_{(i, U)} e_i * \varepsilon_U$, where (i, U) runs through the vertices of Q_G ;
- denote by S_1 the $\mathbb{k}$ -algebra $\Pi_{i \in [G \setminus Q_0]} \mathbb{k}G_i$.

First, it is necessary to check that the quiver Q_G corresponds to the one given by Demonet in [3].

Proof of Theorem 1.6. Following [3], see p. 1057, the algebra $T_S(M) * G$ is Morita equivalent to $\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$, which is isomorphic to the path algebra of a quiver having the same vertices as Q_G and such that the arrows from a vertex (i, U) to a vertex (j, V) form a basis of the vector space $(e_i * \varepsilon_U) \cdot (MG) \cdot (e_j * \varepsilon_V)$. Now, considering e_i , e_j , ε_U , and ε_V as elements of $S * G$,

$$\begin{aligned} (e_i * \varepsilon_U) \cdot (MG) \cdot (e_j * \varepsilon_V) &= \varepsilon_U \cdot (e_i \cdot (MG) \cdot e_j) \cdot \varepsilon_V \\ &\stackrel{(4.1.1)}{=} \varepsilon_U \cdot M(i, j) \cdot \varepsilon_V \\ &\simeq \text{Hom}_{\mathbb{k}G_i}(U, M(i, j) \otimes_{G_j} V) \\ &\stackrel{(4.1.7)}{\simeq} \text{Hom}_{\mathbb{k}G_i}(U, M(i, j; V)). \end{aligned}$$

Whence the first statement of the theorem. Now, if $U = \mathbb{k}G_i \cdot \varepsilon_U$ for all $(i, U) \in Q_G$, then $f(\varepsilon_U)$ does belong to $(e_i * \varepsilon_U) \cdot (MG) \cdot (e_j * \varepsilon_V)$, which is contained in $\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$, for all arrows $f: (i, U) \rightarrow (j, V)$ of Q_G . Whence the isomorphism (1.1.1). $\square$

The proof of Theorem 1.25 uses Morita equivalences in the graded sense. Two $(\mathbb{N})$ -graded finite dimensional $\mathbb{k}$ -algebras Λ_1 and Λ_2 are Morita equivalent in the graded sense if $\Lambda_2 \simeq \text{End}_{\Lambda_1}(P)^{\text{op}}$ for some graded projective Λ_1 -module P which is a generator of the category of graded Λ_1 -modules, and hence of $\text{mod}(\Lambda_1)$. Both

Λ_1 and Λ_2 are isomorphic as graded algebras to basic graded finite dimensional algebras; moreover Λ_1 and Λ_2 are Morita equivalent in the graded sense if and only if their basic versions are isomorphic as graded algebras.

Proof of Theorem 1.25. Denote by Φ the algebra homomorphism $\mathbb{k}Q_G \rightarrow \mathcal{I}ntw^{\text{op}}$ given in the statement of the theorem. Hence, $\Phi(\gamma) = f_\gamma$ for all paths γ in Q_G .

(1) Let γ and γ' be parallel paths in Q_G . It follows from the definition of $\varphi_{\gamma'}$, see Notation 1.23, and from (4.3.9) that

$$(5.0.1) \quad (f_\gamma | \varphi_{\gamma'}) = \begin{cases} 1 & \text{if } \gamma = \gamma', \\ 0 & \text{otherwise.} \end{cases}$$

Since Φ maps non parallel paths in Q_G into distinct components of $\mathcal{I}ntw$ in the decomposition (1.4.3), then Φ is injective. In order to prove that Φ is an isomorphism, it is hence sufficient to prove that, for all integers n , the homogeneous components of degree n of the graded algebras $\mathbb{k}Q_G$ and $\mathcal{I}ntw$ have the same dimensions. Note that $\mathbb{k}Q_G$ is graded by the length of the paths in Q_G and $\mathcal{I}ntw$ is graded so that every intertwiner lying in a space of the shape $\text{Hom}_{\mathbb{k}G_{i_0}}(U, M(i_0, \dots, i_n; V))$ is homogeneous of degree n .

First, following [3], see Theorem 1.6, the algebras $\mathbb{k}Q_G$ and $\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$ are isomorphic $\mathbb{k}$ -algebras both Morita equivalent to $T_S(M) * G$. Actually, the proof given in [3] works in the graded setting.

Next, $\hat{e}$ may be considered as an idempotent of $T_S(M) * G$. In this sense,

$$\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e} = \tilde{e} \cdot \hat{e}(T_S(M) * G) \cdot \hat{e} \cdot \tilde{e}.$$

Following [3], see p. 1057, the algebra $\hat{e} \cdot (T_S(M) * G) \cdot \hat{e}$ is Morita equivalent to $T_{S_1}(\hat{e} \cdot (MG) \cdot \hat{e})$. Again, this is still true in the graded sense. Hence, $\tilde{e} \cdot (T_S(M) * G) \cdot \tilde{e}$, which is basic, is isomorphic to $\tilde{e} \cdot T_{S_1}(\hat{e} \cdot (MG) \cdot \hat{e}) \cdot \tilde{e}$ as a graded algebra.

Finally, by definition of $\tilde{e}$ and $\hat{e}$, the homogeneous component of degree n of $\tilde{e} \cdot T_{S_1}(\hat{e} \cdot (MG) \cdot \hat{e}) \cdot \tilde{e}$ is equal to

$$\bigoplus_{i_0, i_n \in [G \setminus Q_0], U_0 \in \text{irr}(G_{i_0}), U_n \in \text{irr}(G_{i_n})} (e_{i_0} * \varepsilon_{U_0}) \cdot (MG)^{\otimes_{S_1} n} \cdot (e_{i_n} * \varepsilon_{U_n}).$$

This is equal to

$$\bigoplus_{i_0, \dots, i_n \in [G \setminus Q_0], U_0 \in \text{irr}(G_{i_0}), U_n \in \text{irr}(G_{i_n})} \varepsilon_{U_0} \cdot M(i_0, \dots, i_n) \cdot \varepsilon_{U_n}.$$

This is isomorphic to

$$\bigoplus_{i_0, \dots, i_n, U_0, U_n} \text{Hom}_{\mathbb{k}G_{i_0}}(U_0, \underbrace{M(i_0, \dots, i_n) \otimes_{G_{i_n}} V}_{\simeq M(i_0, \dots, i_n; V)}).$$

These considerations prove that the homogeneous components of degree n of $\mathbb{k}Q_G$ and $\mathcal{I}ntw$ are isomorphic. Thus, Φ is an isomorphism.

(2) follows from (5.0.1) and from the fact that Φ is surjective. $\square$

Proof of Theorem 1.27. First, Ξ is bijective. Indeed, on one hand, there is an equality of vector subspaces of $T_S(M) * G$,

$$\hat{e} \cdot (T_S(M) * G) \cdot \hat{e} = \bigoplus_{\underline{i}, \underline{y}} ({}_{i_0} M_{y_1 \cdot i_1} \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} {}_{y_1 \cdots y_{n-1} \cdot i_{n-1}} M_{y_1 \cdots y_n \cdot i_n}) * y_1 \cdots y_n \mathbb{k}G,$$

where $\underline{i}$ runs through all sequences $i_0, \dots, i_n$ of $[G \setminus Q_0]$ and $\underline{y}$ runs through all sequences $y_1, \dots, y_n$ such that $y_t \in [G/G_{i_t}]$ for all $t \in \{1, \dots, n\}$. On the other

hand, for all vertices (i, U) and (j, V) of Q_G and for all $\mathbb{k}G_i - \mathbb{k}G_j$ -bimodules N , the following mapping is bijective,

$$\begin{array}{ccc} \text{Hom}_{\mathbb{k}G_i}(U, N \cdot \varepsilon_V) & \longrightarrow & \varepsilon_U \cdot N \cdot \varepsilon_V \\ f & \mapsto & f(\varepsilon_U). \end{array}$$

Hence, Ξ is bijective and Ξ^{-1} is given by $\Xi^{-1}(f) = f(\varepsilon_U)$ for all $f \in \text{Intw}$ with domain denoted by U .

Next, Ξ is a morphism of $\mathbb{k}$ -algebras. Indeed, for all $f \in \text{Hom}_{\mathbb{k}G_{i_0}}(U, M(\underline{i}; V))$ lying in Intw , then $\Xi^{-1}(f) = f(\varepsilon_U)$. Therefore, with the setting of Definition 1.13, then $(f' \otimes f)(\varepsilon_U)$ is equal to the product $f(\varepsilon_V) \cdot f'(\varepsilon_U)$ in $T_S(M) * G$, see (1.4.1). Accordingly, Ξ^{-1} , and hence Ξ , is a $\mathbb{k}$ -algebra homomorphism.

Finally, the diagram (1.6.4) is commutative. Indeed, all its arrows are $\mathbb{k}$ -algebra homomorphisms. Hence, it suffices to prove the commutativity on the arrows of Q_G , which follows from the definitions of Ξ , (1.1.1), and (1.6.1). Note that (1.6.5) follows from (1.6.2). $\square$

6. ACKNOWLEDGEMENTS

I thank Edson Ribeiro Alvares for helpful comments on previous versions of this article. I also thank the referee for several suggestions that improved the presentation of the article.

REFERENCES

- [1] C. Amiot and P.-G. Plamondon. The cluster category of a surface with punctures via group actions. arXiv:1707.01834 [math.RT], 2017.
- [2] R. Bocklandt, T. Schedler, and M. Wemyss. Superpotentials and higher order derivations. *J. Pure Appl. Algebra*, 214(9):1501–1522, 2010.
- [3] L. Demonet. Skew group algebras of path algebras and preprojective algebras. *J. Algebra*, 323(4):1052–1059, 2010.
- [4] J. Dionne, M. Lanzilotta, and D. Smith. Skew group algebras of piecewise hereditary algebras are piecewise hereditary. *J. Pure Appl. Algebra*, 213(2):241–249, 2009.
- [5] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. *Tensor categories*, volume 205 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.
- [6] V. Ginzburg. Calabi-Yau algebras. arXiv:math/0612139 v3, 2006.
- [7] S. Giovannini. Higher representation infinite algebras from McKay quivers of metacyclic groups. *Comm. Algebra*, 47(9):3672–3711, 2019.
- [8] S. Giovannini and A. Pasquali. Skew group algebras of Jacobian algebras. *J. Algebra*, 526:112–165, 2019.
- [9] S. Giovannini, A. Pasquali, and P.-G. Plamondon. Quivers with potentials and actions of finite abelian groups. arXiv:1912.11284 [math.RT], 2019.
- [10] P. Le Meur. Crossed-products of Calabi-Yau algebras by finite groups. *Journal of Pure and Applied Algebra*, 224(10):106394, 2020.
- [11] L. Li. Finitistic dimensions and piecewise hereditary property of skew group algebras. *Glasg. Math. J.*, 57(3):509–517, 2015.
- [12] R. Martínez-Villa. Skew group algebras and their Yoneda algebras. *Math. J. Okayama Univ.*, 43:1–16, 2001.
- [13] I. Reiten and C. Riedtmann. Skew group algebras in the representation theory of Artin algebras. *J. Algebra*, 92(1):224–282, 1985.
- [14] I. Reiten and M. Van den Bergh. Two-dimensional tame and maximal orders of finite representation type. *Mem. Amer. Math. Soc.*, 80(408):viii+72, 1989.

UNIVERSITÉ DE PARIS, SORBONNE UNIVERSITÉ, CNRS, INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, F-75013 PARIS, FRANCE

Email address: patrick.le-meur@imj-prg.fr