Northern Puna Plateau-scale survey of Li brine-type deposits in the Andes of NW Argentina

R. Lucrecia López Steinmetz, Stefano Salvi, M. Gabriela García, Didier Béziat, Gabriela Franco, Ornela Constantini, Francisco Córdoba, Pablo Caffè

► To cite this version:

R. Lucrecia López Steinmetz, Stefano Salvi, M. Gabriela García, Didier Béziat, Gabriela Franco, et al.. Northern Puna Plateau-scale survey of Li brine-type deposits in the Andes of NW Argentina. *Journal of Geochemical Exploration*, 2018, 190 (2), pp.26-38. 10.1016/j.gexplo.2018.02.013 . hal-02363093

HAL Id: hal-02363093

<https://hal.science/hal-02363093>

Submitted on 6 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Northern Puna Plateau-scale survey of Li brine-type deposits in the Andes of NW Argentina

R. Lucrecia López Steinmetz ^{1,2,*}, Stefano Salvi ³, M. Gabriela García ⁴, Didier Beziat ³, Gabriela Franco ⁵, Ornella Constantini ⁵, Francisco E. Córdoba ^{5,6}, Pablo J. Caffè ^{5,6}

¹ Instituto de Geología y Minería, Universidad Nacional de Jujuy, Av. Bolivia 1661, S.S. de Jujuy 4600, Argentina

² CIT Jujuy CONICET, Universidad Nacional de Jujuy - CONICET, S.S. de Jujuy 4600, Argentina

³ Université de Toulouse, CNRS, GET, IRD, OMP, 14 Av. Edouard Belin, Toulouse 31400, France

⁴ CICTERRA-CONICET-Universidad Nacional de Córdoba, Córdoba 5016, Argentina

⁵ Instituto de Ecorregiones Andinas (INECOA), Universidad Nacional de Jujuy - CONICET, S.S. de Jujuy 4600, Argentina

⁶ Facultad de Ingeniería, Universidad Nacional de Jujuy, Argentina

* corresponding author: lucrecialopezsteinmetz@hotmail.com

Abstract

Salars of the Andean Plateau in the Central Andes are the largest lithium deposits on Earth. The most notorious are the Chilean Salar de Atacama, and Uyuni in the Bolivian Altiplano. Despite the relevance of the region concerning the lithium resources, there is still scarce scientific literature on the hydrochemistry of lithium deposits in the Argentine portion of the Andean Plateau. In this article we present new hydrochemical data from the first regional-scale reconnaissance exploration of the four major salars in the northernmost Argentine Andes. Data revealed that brines in the studied salars are characterized by mean Li⁺ concentrations ranging between 82 and 1,014 mg L⁻¹, and mean Li:Mg ratios from 0.92 to 0.54. The size of the study salars becomes a potential limitation for the whole Li resources in comparison with the giant Atacama and Uyuni. Nonetheless, when considering the Li grade and the Li:Mg ratio of brines, the Northern Puna salars turn out to be very remarkable lithium prospects. Data emerged from this survey represent a valuable tool for: 1) private investment projects by defining Li mining targets, 2) for the administration of natural resources and the definition of the State's politics, and 3) for scientific purposes, especially in investigations meant to better understand the processes involved in the formation of Li brine deposits, salars, and endorheic basins.

Keywords: lithium, salars, Central Andes, Altiplano-Puna Plateau, hydrochemistry

Introduction

Salars of the Central Andes are the largest lithium deposits on Earth (Warren 2010; Houston et al. 2011; Kesler et al. 2012; Munk et al. 2016). Particularly, the largest global resources concentrate in numerous internally drained basins of Chile, Bolivia and Argentina, where the lithium (Li) extraction has already become an important source of income in these South American countries. The Andean salars of Uyuni in Bolivia and Atacama in Chile (Fig. 1) contain the most important known Li resources in the world (Moraga et al. 1974; Gruber and Medina 2010; Mohr et al. 2010; Kunasz 2006; Kesler et al. 2012; Munk et al. 2016).

Uyuni is a spectacularly giant salar and with a surface of more than 10,500 km² it is the largest salt pan of the Andes and on Earth (Risacher and Fritz 1991; Kesler et al. 2012). This salt pan is located in the Altiplano at 3,663 m a.s.l. (Fig. 1). The mean Li content in the near-surface brines of Uyuni ranges from 316 to 602 mg L⁻¹ and the Li concentration pattern across the salar increases towards the south (Ericksen et al. 1976; Ericksen et al. 1977; Rettig et al. 1980; Risacher and Fritz 1991; Schmidt 2010). The estimated Li resources of Uyuni are over 10.2 Mt, which represent ~12 times the estimated resources of Greenbushes in Australia, the world famous Li pegmatite deposit (Kesler et al. 2012).

Although Uyuni is probably the most important Li resources worldwide, this salar is not in production yet. Much of the Li being produced globally at present is obtained from the Salar de Atacama. With estimated resources reaching 6.3 Mt, Atacama is thus the most important Li mine in the world (Moraga et al. 1974; Ide and Kunasz 1989; Kunasz 2006; Lowenstein and Risacher 2009; Gruber and Medina 2010; Mohr et al. 2010; Kesler et al. 2012; COCHILCO 2013).

The Salar de Atacama (Fig. 1) is a 3,000 km² salt pan located at 2,300 m a.s.l., whose halite nucleus spreads 1,100 km² and ~1,000 m depth (Moraga et al. 1974; Stoertz and Ericksen 1974; Bevacqua 1992; Alonso and Risacher 1996; Carmona et al. 2000; Boschetti et al. 2007; Jordan et al. 2007; Boutt et al. 2016; Corenthal et al. 2016). The average Li concentration of Atacama is ~1,400 mg L⁻¹, ranging between 1,000 and 6,400 mg L⁻¹ (Moraga et al. 1974; Ide and Kunasz 1989; Kunasz 2006), representing the maximum Li concentration in a brine-type deposit reported to date worldwide. The Li concentration pattern across the Salar de Atacama is well known: five concentric Li grade zones were defined, with the higher Li grade zone located in the southern part of the salar (Ide and Kunasz 1989).

In the Argentine Puna region, in the southernmost part of the Andean Plateau (Fig. 1), there are also some remarkable Li prospects. The most prominent is the Salar de Hombre Muerto (Fig. 1), which became the

first Li mine operating in Argentina since the 90's (Kesler et al. 2012; Fornillo et al. 2015). Hombre Muerto comprises a 300 km² salt pan placed at 3,970 m a.s.l., with estimated Li resources of 0.8 Mt, and mean Li contents of ~520 mg L⁻¹; the concentration pattern across the salar increases towards the south (Garrett 2004; Kesler et al. 2012; Godfrey et al. 2013).

As mentioned, the Li resources of salars in the Chilean Desierto de Atacama and the Bolivian Altiplano are largely known (Ericksen et al. 1977; Ide and Kunasz 1989; Risacher and Fritz 1991; Alonso and Risacher 1996; Risacher et al. 1999; Carmona et al. 2000; Risacher et al. 2003; Kunasz 2006; Risacher and Fritz 2009). On the contrary, and despite the relevance of this strategic mineral resource and the outstanding prospect of the Andean salars, there are still scarce scientific works on the hydrochemistry of Li deposits in the Argentinean Puna (Fig. 1). This region is becoming an increasing focus of exploration activity. Most of the available information of Puna salars is based on reports provided by mining companies and regional data reported in a number of scientific papers (Yaksic and Tilton 2009; Gruber and Medina 2010; King 2010; Houston et al. 2011; Fornillo et al. 2015; Munk et al. 2016). On the contrary, local works aiming to characterize the hydrochemistry of Li deposits in this Andean region are restricted to the Salar de Hombre Muerto (Godfrey et al. 2013; Fig. 1) and the Salar de Centenario-Ratones (Orberger et al. 2015) in the Southern Puna (Alonso et al. 1984), and the Guayatayoc playa lake in the Northern Puna (López Steinmetz 2017; Fig. 2). There are still many other salars in the Puna region that are promising Li prospects, and that – to date – were not object of scientific surveys, even of a reconnaissance nature. For instance, in the Northern Puna little is known about the hydrochemistry of the Cauchari and the Salinas Grandes salt pans, whereas the Salar de Jama remains completely unexplored (Fig. 2).

In this work we report the results of a regional Li exploration campaign covering an area of 9,000 km² in the endorheic Northern Puna (Fig. 2) that includes the four major salars which represent promising Li prospects. The major chemical composition and Li content of brine samples collected from the Salar de Jama, Olaroz, Cauchari and Salinas Grandes are analysed in order to contribute to the knowledge of the Andean salars hydrochemistry from a regional perspective. This research is the first scientific report on Li resources in the Northern Puna Plateau, and, along with the information previously reported by López Steinmetz (2017), completes the first broad survey of Li-bearing salars on the northern Argentine Plateau. We finally propose to discuss the Li resource perspectives of the Northern Puna region in the context of the most notorious Andean prospects.

The study area

The salars analysed in this study are all set in the endorheic Northern Puna region (Fig. 2). The Salar de Jama is the smallest one, and is located on the Argentina-Chile border, at the foothills of the Jama volcano. Eastward, the Salar de Olaroz and Cauchari occupy the same north-south elongate orographic depression, superficially detached one from the other by prograding alluvial fans (the composite Archibarca alluvial cone). Finally, the Salinas Grandes covers the easternmost reaches of the lower Northern Puna, limiting to the north with the Guayatayoc salar basin.

There is an altitudinal gradient across the Northern Puna, which determines that western salars are placed at higher altitude than those located to east (Fig. 2). These four salars extend between 23° - 24°S. Northward, the endorheic basins of the Northern Puna are characterized by the presence of lakes that only occasionally dry out (e.g., Laguna de Pozuelos, McGlue et al. 2012; Laguna de Pululos, Lupo et al. 2006).

Orographic depressions that host salars are defined by north-south trending ranges formed during Andean thrusting events. These ranges expose the basement, which consist of Lower to Middle Ordovician (Santa Victoria Group, Turner 1960 a, b) and Precambrian - Eocambrian (Puncoviscana Formation, Turner 1960 a, b) metasedimentary rocks in the Puna Plateau and the Eastern Cordillera, respectively. The latter unit, together with Upper Cambrian sedimentary rocks (Meson Group, Turner 1960 a, b) are only present along the eastern border of the study area (Fig. 2). Middle to Upper Ordovician peraluminous, plutonic complexes and Jurassic to Cretaceous alkaline granites (Méndez et al. 1973; Zappettini 1989; Coira et al. 1999; Cristiani et al. 1999; Kirschbaum et al. 2006) are widely distributed in ranges along the western side of the Salinas Grandes and, in a lesser extent, along the eastern margin of the Salar de Olaroz (Fig. 2).

Cretaceous to Paleocene rift-related sedimentary rocks (Salta Group, Turner 1959) frequently crop out overlain by Upper Eocene to Upper Miocene synorogenic deposits (Turner 1972; Schwab 1973; Schwab and Lippolt 1974; Arias and Viera 1982; Seggiaro 2013). These units are widespread in the southern portions of the Tanque and Pircas ranges, and in the Alta range, in the southern border of the Salinas Grandes.

Paleogene to Lower Miocene continental red beds (Log Log, Vizcachera and Peña Colorada formations; see Coira et al. 2004; Seggiaro 2013; and references therein) were deposited afterwards in a broken foreland environment (e.g., del Papa et al. 2013), Fluvial to lacustrine deposits intercalated with air fall tuffs and pyroclastic flow deposits (Tiomayo, Trincheras and Pastos Chicos formations) characterise the Middle to Upper Miocene (16 to 10 Ma) sedimentary/volcanic record (Schwab and Lippolt 1974; Coira et al. 2004). Towards the Upper Miocene-Pliocene transition similar sedimentary sequences (Loma Blanca/Sijes Formation; Alonso 1986) locally involved deposition of evaporite facies (Olaroz-Coranzulí).

Large areas of the Northern Puna are covered by Upper Miocene-Pliocene and Pleistocene ignimbrites, dacite to rhyolite lavas and subvolcanic intrusions, and mafic andesite volcanic centres (Coira et al. 1993; Seggiaro 1994; Coira et al. 2004; Seggiaro 2013; Maro and Caffè 2016; and references therein). Quaternary alluvial fan deposits extend between the foot of the mountains and the flat areas, which are covered by fine-grained sediments and salt pans. Salar mining in the Northern Puna was traditionally related to the extraction of halite and ulexite from these salt pans, while Li mining started recently (2014), the Salar de Olaroz and Cauchari being the only active brine prospects (Fornillo et al. 2015).

The environmental conditions that prevail in these high-altitude endorheic basins are typically extreme, with low effective precipitation and wide temperature fluctuations. Thus, rainfall is markedly seasonal, with ~70% of the yearly total falling during the austral summer (December-March), while the dry season spans over the rest of the year. High rates of evaporation and low annual rainfall, averaging 150 - 400 mm and increasing from SW to NE, are responsible for a negative water balance. As a consequence, fluvial systems mainly consist of ephemeral streams. Despite most tributaries infiltrate before reaching the salars, some major fluvial collectors feed the systems over the year (Fig. 2). In addition, groundwater discharge probably represents an important source of water to the salar systems (Peralta Arnold et al. 2017).

Methods

For the Li reconnaissance exploration and hydrochemical assessment of the Northern Puna salars, residual brines were sampled across the Salar de Jama (JA), Olaroz (OL), Cauchari (CA) and Salinas Grandes (SG, Fig. 3). The detailed location of samples is presented in Table A of the electronic supplementary material. Brine samples were collected and analysed during the austral spring 2010 (from September until December). Samples were collected at 40 to 90 cm depth in pits dug in salars, except for samples JA10, OL10, CA1, CA2 and SG10 that were collected from brine pools. Samples were stored in pre-cleaned polyethylene bottles and preserved refrigerated during transport.

Brine samples were analysed at the Agua de los Andes S. A. laboratory. Analytical results are summarized in Table 1. Physicochemical parameters such as pH and electrical conductivity were determined with a Hanna HI2314 multi-range conductivity meter. The major ion composition was determined following standard recommendations (Clesceri et al. 1998). Calcium (Ca^{2+}) was determined by titration using a 0.01 M EDTA solution, while magnesium (Mg^{2+}) was calculated as the difference between total hardness and calcium concentration. Chloride (Cl^-) was determined by titration using the argentometric method. Sulphate (SO_4^{2-}) was

precipitated in an acetic acid medium with BaCl_2 and its concentration was determined by turbidimetry. Bicarbonate and carbonate (HCO_3^- and CO_3^{2-}) were determined by titration using a 0.01 N sulfuric acid solution and methyl-orange and phenolphthalein were used as end-point indicators, respectively. Boron (B) was determined by colorimetry using carmine reactive. Sodium (Na^+), potassium (K^+) and lithium (Li^+) were determined by Atomic Emission Spectroscopy. Ion balances were typically lower than 5% in all cases. Saturation index (SI) was calculated by using the PHREEQC 3.3.8 public domain and simulation program of the USGS, and employing the Pitzer database.

Li-bearing salars of the Northern Puna

The Salar de Jama

Jama is a north-south elongated salt pan located at 4,080 m a.s.l. The salar surface is 25 km², spreading between 67°02' - 67°04'W and 23°15' - 23°21'S (Fig. 3A). Main tributaries are the Coruro, Lina and Pircas Rivers. Local lithology involves the Ordovician metasedimentary basement and Miocene - Pliocene ignimbrites (the Mucar, Atana and Toconao ignimbrites), dacite lavas as well as small mafic andesite volcanic centres (the Jama and Archibarca volcanoes). The salar overlies Quaternary alluvial deposits that interdigitate with a muddy salt shoreline along the south-eastern border of the salar.

Jama's brines are alkaline, with pH values ranging between 8.2 and 8.8 (Table 1), and moderately saline, with electrical conductivity values ranging between 12.33 and 183.63 mS cm⁻¹ (Table 1). According to their major chemical composition, brines are of the Na/Cl⁻ - SO₄²⁻ type, with relatively abundant Ca²⁺ and K⁺ (Fig. 4).

When the mean ionic concentration of Jama's brines is compared with those of the other Northern Puna salars, the Jama brines result the most dilute (Fig. 5). The Na:Cl ratios in the Jama brines are ~1 (Fig. 6A). The SO₄²⁻ and Ca²⁺ concentration pattern characterizes by a SO₄²⁻>Ca²⁺ linear trend (Fig. 6B), except for the samples JA6, JA9 and JA10 that depict much higher SO₄²⁻ concentrations.

Ratios near to 1 evidence salt recycling process, including halite and anhydrite/gypsum solubilisation and precipitation. According to saturation indexes (SI), these brines are unsaturated in anhydrite (SI from -2.19 to -1.41), gypsum (SI from -1.97 to -1.11), sylvite (SI from -4.29 to -2.11) and halite (SI from -3.60 to -1.43). According to this, the excess of SO₄²⁻ over Ca²⁺ (Fig. 6B) evidences the contribution of sulphate originated from the solubilisation of non-Ca sulphates and/or the scavenging of Ca²⁺ by cation exchange and/or by

dolomite/calcite precipitation. Because brines are supersaturated with dolomite (IS from 0.33 to 1.07) and calcite (IS from 0.05 to 0.60) it seems that the latter should be the dominant process controlling Ca^{2+} concentrations in Jama's brines.

The Li^+ concentration in the Jama's brines ranges between 1.5 and 262 mg L^{-1} , with the highest Li grade zone along the north-western border of the salar (Fig. 7A). The mean Li^+ concentration in the shallowest brines of the salar is 82 mg L^{-1} while the mean Li:Mg ratio is 0.63 (Table 2). Borate concentrations in these brines range between 236 and 3,069 mg L^{-1} ; no linear correlation between Li and B has been observed, being $\text{Li} < \text{B}$ with the Li/B ratio up to 0.29 (Table 1).

The Salar de Olaroz and Cauchari

These two salars are located in a N-S elongated basin at an average altitude of 3,903 m a.s.l. The Salar de Olaroz is a 130 km^2 salt pan spanning between $66^\circ 39' - 66^\circ 45' \text{W}$ and $23^\circ 24' - 23^\circ 33' \text{S}$ (Fig. 3B). The northern catchment area includes numerous hydrothermal springs, where the main fluvial collector is the Rosario River that discharges into the north of the salar. The other main tributary to the salar is the Archibarca River that reaches the salt pan from the west. This river forms a huge prograding alluvial fan that dissects the Olaroz - Cauchari topographic depression and acts as a barrier between the Olaroz (to the north) and the Cauchari salt pans.

The Salar the Cauchari is located at the southern part of the basin, between $23^\circ 39' - 23^\circ 50' \text{S}$, covering an area of $\sim 80 \text{ km}^2$ (Fig. 3B). The dominant lithologies in the basin are sedimentary sequences and volcanic rocks. The Tocomar River is the major fluvial collector and reaches the salar from the south (Fig. 2). The southern catchment area includes numerous hydrothermal springs.

Shallow brines in both salars are circumneutral to alkaline, with pH values ranging between 8.1 and 8.5 in Olaroz and from 7.2 to 9.3 in Cauchari (Table 1). The brines' major chemical composition is dominated by Na/Cl^- type waters (Fig. 4), with electrical conductivity values and ionic contents that are up to one order of magnitude higher than those measured in the Jama's brines (from 184.85 to 320.15 mS cm^{-1} , Fig. 5B). These values represent the highest ionic concentrations measured in the Northern Puna salars.

The Na:Cl ratios are ~ 1 in Cauchari and ≤ 1 in Olaroz (Figure 6A). Despite this, brines are undersaturated with halite in both the Cauchari and the Olaroz salars (SI ranging from -1.44 to -0.91). Like Jama, the concentration of SO_4^{2-} is higher than Ca^{2+} in the Olaroz and Cauchari brines (Fig. 6B). This pattern could be the result of coupled reactions involving dissolution of gypsum/anhydrite and subsequent scavenging of Ca^{2+} by

dolomite/aragonite/calcite precipitation. In agreement, saturation indexes for anhydrite and gypsum vary from -3.29 to -2.11 and -3.16 to -1.92, respectively. However, brines are supersaturated with carbonates only in the northern Cauchari area (the $SI_{dolomite}$ ranges from 0.44 to 1.40 in samples CA1, CA2 and CA3, the $SI_{aragonite}$ is 0.24 in the CA2, and the $SI_{calcite}$ varies from 0.08 to 0.38 in samples CA2 and CA3), but are undersaturated in the southern Cauchari and the entire Olaroz ($SI_{dolomite}$: -1.56 to -0.16; $SI_{aragonite}$: -1.37 to -0.07; $SI_{calcite}$: -1.22 to -0.22). These values evidence thermodynamically optimum conditions for locally constrained carbonate precipitation and sulphate dissolution, leading to the concomitant existence of sulphated-rich inputs.

The Li^+ concentration in shallow brines ranges from 771 – 1,213 mg L⁻¹ in Olaroz and 442 – 1,739 mg L⁻¹ in Cauchari. The highest Li grade zones occupy the central area of these two salars (Fig. 7B and C). The mean Li^+ concentration is 860 mg L⁻¹ in Cauchari and 1,014 mg L⁻¹ in Olaroz, the Li:Mg ratios are 0.86 and 0.89, respectively (Table 2). The borate concentration in these brines varies between 1,336 and 3,175 mg L⁻¹ (Table 1) and do not show a linear correlation with Li contents, with $Li < B$ and the Li/B ratio up to 0.96 in Cauchari and 0.29 in Olaroz.

The Salinas Grandes

Salinas Grandes is a large salt pan of 280 km² placed at 3,410 m a.s.l. between 23°33' - 23°45'S and 65°51' - 66°12'W (Fig. 3C). This salar developed south of the Las Burras alluvial fan, which separates the Salinas Grandes salt pan from the Guayatayoc playa lake to the north. Permanent streams feeding the Salinas Grandes include the San Antonio de los Cobres and the El Moreno Rivers. The southern catchment area of the San Antonio de los Cobres River includes numerous hydrothermal springs, while the headwater area of the El Moreno basin involves seasonal periglacial processes. Muddy salt shorelines associated to the mouth of the El Moreno River are mainly developed along the south-eastern border of the salt pan. Quaternary alluvial deposits cover the low lying areas and reach the foot of mountains. The metasedimentary basement and the Cretaceous to Paleocene sedimentary sequences are the most widespread lithologies along the eastern side of the basin, while Paleozoic and Mesozoic plutonic rocks outcrop in the west-boundary ranges.

Similar to other salars of the study area, shallow brines in the Salinas Grandes are circumneutral to slightly alkaline, with pH ranging from 7.1 and 8.0 (Table 1). Electrical conductivity show intermediate values compared with the remaining salars of the Northern Puna, typically ranging between 232.18 and 241.08 mS cm⁻¹ (Fig. 5B). Major ionic concentrations are dominated by Na and Cl ions (Fig. 4). Interestingly, these brines show the largest variability of the ionic concentrations compared with the neighbour salars (Fig. 5). The sample SG1 is

the most diluted (conductivity = 233.85 mS cm⁻¹) and characterizes by the lowest Ca²⁺ concentration compared with the remaining samples collected from the Salinas Grandes (Fig. 5), and a Na:Cl ratio ~1 (Fig. 6A). Conversely, the sample SG7 contains high Mg²⁺ and K⁺ concentrations, even when compared with the other salars of the Northern Puna (Fig. 5A). The Salinas Grandes brines also show nearly equivalent proportions of Ca²⁺, Mg²⁺ and K⁺ with Ca:Mg ~1 (Fig. 4), and typically have Ca²⁺ > SO₄²⁻ (Fig. 6B). Saturation indexes indicate that brines are undersaturated in calcite, aragonite, and dolomite (SI_{dolomite}: -2.07 and -1.55, SI_{aragonite}: -1.30 to -1.11 and the SI_{calcite}: -1.15 and -0.97).

The equivalent concentrations of Li and borate through the salar are similar (Fig. 5A), with the Li:B ratio close to 1 in samples SG4, SG6 and SG9 (Li/B of 0.95, 1.10 and 0.99, respectively), Li>B in samples 5, 8 and 10 (from 1.17 to 1.32), and Li<B in the remaining samples (SG1 = 0.07, SG2 = 0.67 and SG3 = 0.70). The concentration of borate is from 309 and up to 542 mg L⁻¹, and Li⁺ ranges between 25 and 1,018 mg L⁻¹, with the highest Li grade zone located at the western margin of the salar (Fig. 7D). The mean Li⁺ concentration is 332 mg L⁻¹ and the mean Li:Mg ratio is 0.54 (Table2).

DISCUSSION

The major chemical composition of the studied brines is rather constant; most samples are of the Na/Cl type with some subordinated samples of the Na/Cl-SO₄²⁻ type. Thus, the highly concentrated brines are typically rich in Na⁺ and Cl⁻, and relatively poor in Ca²⁺, Mg²⁺ and HCO₃⁻ + CO₃²⁻. An additional common feature in the Northern Puna brines is that K⁺ concentrations are from 2 to 6 times higher than those of Mg²⁺. However, the most conspicuous feature is the high Li grades, with average concentrations from 82 to 1,014 mg L⁻¹, the highest ones found in the Cauchari and Olaroz salt pans.

Hydrochemical features of the Andean salars are controlled by a number of variables acting at different scales. In a recent work, Munk et al. (2016) have defined six major features shared by most Li-rich brines deposits at a global scale: (1) arid climate; (2) closed basin containing a salar (salt crust), a salt lake, or both; (3) associated igneous and/or hydrothermal activity; (4) tectonically driven subsidence; (5) suitable Li sources; and (6) sufficient time to concentrate Li in the brine. However, even though all these features converge across the Puna Plateau, brines from the studied salars have an uneven distribution of Li grades. Therefore, there must be some local factors constraining the higher concentrations of Li to some particular systems.

Some *in situ* processes that are taking place within the salars are conditioned by the ionic content and salinity (Fig. 5) of the study brines. In the four studied salars, average salinity is the highest in Olaroz and the

lowest in Jama, while intermediate values were measured in Cauchari and Salinas Grandes. In addition to salinity, the concentration pattern $OL \geq CA \geq SG \geq JA$ is valid for Mg^{2+} , K^+ , Li^+ and BO_3^{3-} concentrations, but not for Ca^{2+} , whose regional concentration pattern is $SG \geq JA \geq CA \geq OL$. The lower Ca^{2+} concentrations observed in Olaroz and Cauchari are the consequence of local geochemical processes that involve precipitation of calcite/aragonite but also of some other Ca-scavenging process such as cation exchange. The latter mechanism might occur during the interaction between Ca-rich solutions and clay minerals accumulated in the salt pan.

Residual brines in Northern Puna salars are characterized by a final major composition dominated by Na^+ - Cl^- and SO_4^{2-} ions, which is one typical end member in the chemical divide trend proposed by Hardie and Eugster (1970). The behaviour of trace elements such as Li and borate in the context of this chemical divide has been scarcely assessed. For example, Witherow and Lyons (2011) determined on the basis of geochemical modelling that Li is largely conservative in water samples of the Great Basin (USA), Saskatchewan, and the McMurdo Dry Valleys of Antarctica and, thus, its concentration is the result of long-term solute input and concentration through evaporation and/or sublimation processes. According to this, Li behaves as an incompatible element, not only during the stages of magmatic differentiation but also during weathering (Munk et al. 2016). Therefore, its concentration during brine evolution should follow an increasing trend with increasing water salinity. However, we have observed no unequivocal correlation between the Li grade of brines and conductivity nor other conservative solutes such as Na and Cl. For instance, as shown in Figure 8A, the Jama and Cauchari samples have a remarkable Li/Conductivity correlation, but brines from the Salinas Grandes and Olaroz show Li concentrations that increase independently from salinity (Fig. 8A). This suggests that during the sediment-water interaction, Li ions may in some extent co-precipitate as an impurity in the minerals present in the saline crusts and efflorescens or it could be adsorbed onto clay minerals due to the capacity of Li^+ to be incorporated into the smectite interlayers (e.g., Zhang et al. 1998; William and Herving 2002).

The hydrothermal activity is considered as one of the most significant factors for the formation of Li and boron rich brines in the Central Andes (Risacher et al. 2003; Godfrey et al. 2013; López Steinmetz 2017; and references therein). Indeed, a number of authors indicated that the main sources of Li in Uyuni and the nearby salars of Coipasa and Empexa in the southern part of the Bolivian Altiplano are thermal springs associated with Quaternary rhyolitic volcanism (Shcherbakov and Dvorov 1970; Ericksen et al. 1977; Campbell 2009). Since hydrothermal activity on the region is largely related to volcanism and the formation of the plateau (i.e., Isacks 1988; de Silva 1989; Allmendinger et al. 1997; McQuarrie et al. 2002; Kay et al. 2010; Perkins et al. 2016; and

references therein), it could be considered that this regionally pervasive and large scale process is most likely a primary condition for the formation of exceptional Li brine-type deposits on the Argentine Puna region.

The data that we present in this contribution clearly reveal that in the Norther Puna region, the highest concentrations of Li are restricted to the Olaroz and Cauchari salt pans (Fig. 7). These two salars are located in the same tectonically N-S elongated depression. Therefore, the distinctive Li enrichment in Olaroz and Cauchari is being locally influenced by basin-scale processes, such as the characteristics and density (in terms of the quantity of springs that exist in the basin) of geothermal fields, the weathering of Li-rich lithologies, and/or geochemical processes taking place within the salars. An additional local driving factor on increasing the Li grade in the Olaroz and Cauchari's brines should likely be associated to the role of major thrusts on transporting fluids from the deepest zones of the upper crust and generating large ionic remobilizations. For example, Franco et al. (2016) reported the presence of thermal springs discharging at the location of the basin bounding structures along the marginal zones of the Salar de Olaroz. Additionally, the basin involves several hydrothermal fields directly associated with volcanic centers (Peralta Arnold et al. 2016). Therefore, hydrothermal springs in the Salar de Olaroz and Cauchari are likely related to the presence of volcanic centers, which include isolated edifices along oblique volcanic belts (e.g., Presta and Caffè 2014; Maro et al. 2016) and large volcanic complex calderas (Seggiaro 1994; Seggiaro 2013).

The lower Li grade in brines from the Salinas Grandes relative to Olaroz and Cauchari could be associated with several characteristics of the basin: 1) the catchment lithology is dominated by the Eastern Cordillera basement, which is different from that of the Northern Puna, and would, therefore, have a different geochemical signature; 2) the lack of large hydrothermal fields and volcanic rock outcrops; and 3) the less arid conditions in the basin's catchments located at the easternmost edge of the Norther Puna region. Consequently, even if the hydrothermal activity - which represents just one of the multiple expressions related to the intrinsic nature of the plateau - is considered as a major source for Li supply, additional geological, climatic, and geochemical aspects locally generating distinctive Li grades and brine compositions in every salt pan and/or basin should also be considered.

In order to compare the relative abundance of ionic species in different salars, or between samples from the same salar with uneven salinities, it is necessary to remove the effect of the evaporative concentration. The relative abundance of Ca^{2+} , Mg^{2+} , K^{+} , Li^{+} and BO_3^{3-} in salars from the Northern Puna is presented in Figure 9. In this diagram, the evaporative effect was removed by employing TDS (total dissolved solids, for TDS method see caption of Fig. 9) normalized values. This diagram is substantially different from Figure 5 where plotted data

involve absolute ionic concentrations (the amount of ionic species that could be obtained from one litre of brine). Normalized concentrations revealed that the four salars contain similar proportions of Ca^{2+} , the Salinas Grandes brines are proportionally richer in K^+ and BO_3^{3-} , and the Salar de Cauchari contains proportionally higher Li^+ concentrations than its neighbour Northern Puna salars (Fig.8). This suggests that if a brine sample from the salar de Jama (which is less saline than brines from Cauchari) is hypothetically evaporated in order to increase its salinity up to the values of Cauchari, then, one litre of the Jama brine would still have a lower Li grade than the same volume of the Cauchari brine. Consequently, the evaporative concentration process of such brines would be more efficient for Li recovering at Cauchari than Jama. This statement seems an evident matter when comparing Cauchari and Jama's brines because the mean Li^+ concentration of Cauchari is 860 mg L^{-1} and that of Jama is 82 mg L^{-1} (Table 2). However, the extrapolation of this rule to the other salars of the Northern Puna (Fig. 9) allows stating that the evaporative concentration process in Cauchari is more efficient than in Olaroz (even if the mean Li^+ concentration of the latter reaches $1,014 \text{ mg L}^{-1}$ versus 860 mg L^{-1} of the former).

The integrated analysis of the Li-rich brine chemistry from Northern Puna provides important information to evaluate the potentially economic viability of these brines with respect to Li extraction. In addition to the Li grade of brines, the substrate porosity-permeability, the aquifer water balance, and the volume of the host aquifer (Houston et al. 2011), the magnesium content is an additional issue regarding the Li-mining process because Mg^{2+} causes chemical interferences in the brine purification process and limits the Li^+ recovering (e.g., Crespo et al. 1987; An et al. 2012; Kesler et al. 2012; An et al. 2012). The Uyuni's brines typically have high grade Mg^{2+} (Ericksen et al. 1978; Risacher and Fritz 1991) while the Salar de Hombre Muerto represents the opposite situation, with low magnesium contents (Kesler et al. 2012; Godfrey et al. 2013; Table 2, Fig. 8B). The Li:Mg ratios in the Northern Puna range from 0.54 to 0.92 (Table 2), which in the regional context make them closer to Hombre Muerto than to Uyuni (Fig. 8B).

Economically viable Li-rich brines in the Central Andes contain varying amounts of the major cations and anions (Na^+ , K^+ , Mg^{2+} , Ca^{2+} , Cl^- , SO_4^{2-} , and CO_3^{2-}), which can form a wide range of ionic brine compositions. In general, the Northern Puna's brines contain less Mg^{2+} than those accumulated in Atacama and Uyuni (Fig. 8B). Nonetheless, the extremely high Li^+ concentration of Atacama remains the most important difference between all these salars (Table 2). An additional and substantial difference is the oversized surface of Uyuni and Atacama compared with the dimension of the Northern Puna salars. These two aspects, the Li^+ concentration and the salar size, are potentially limiting the whole resources of the Northern Puna salt pans.

However, when considering the Li grade and the Li:Mg ratio of brines, the Northern Puna salt pans are revealed as very remarkable Li prospects.

Conclusion

This study reports the results of a regional Li exploration campaign covering an area of 9,000 km² in the Northern Puna that included the four major salars in the region. These evaporite systems represent remarkable Li prospects with mean Li⁺ concentrations varying between 82 and 1,014 mg L⁻¹, and mean Li:Mg ratios ranging from 0.92 to 0.54.

The major chemical composition of the studied brines is rather constant; most samples are of the Na/Cl type with some subordinated samples with a major ionic composition of the Na/Cl-SO₄²⁻ type, and with K>Mg. Evaporative concentration, dissolution/precipitation of halite and gypsum/anhydrite, and the precipitation of calcite/aragonite/dolomite are important processes controlling the major chemical composition of brines. Unlike major ions, the concentrations of minor elements such as Mg²⁺, Li⁺ and BO₃³⁻ do not show clear correlation with salinity, which suggest that some other factors are controlling their spatial distribution in the shallowest brines of the studied salars. The absence of a geographical/geomorphological pattern for the ionic concentrations suggests that the hydrochemistry of salars is being partially conditioned by local and regional intrinsic characteristics of each basin, as well as by regional features related to the geological/tectonic nature of the Andean Plateau. These characteristics are likely responsible for the higher Li concentrations in the N-S depression occupied by the Olaroz and Cauchari salars compared with the neighbour salars located towards the margins of the northern Puna region.

The relationship between the Li grade and brine salinity indicates that the evaporative Li recovering should be more efficient in Cauchari than in the remaining salars of the Northern Puna. Despite that the salar dimension is a factor potentially limiting the whole Li resources in the Northern Puna, especially when comparing with Atacama and Uyuni, the Li grade and the Li:Mg ratio in the Northern Puna brines are defining outstanding perspectives for the salt pans of this Andean region.

ACKNOWLEDGMENTS

This study was founded by the PRH Red 101 ANPCyT of the Argentine National Government and the National University of Jujuy. R.L. López Steinmetz thanks the grant E/B003 of the SECTER - UNJu. Other funds came from projects PIO CONICET-UNJu 14020140100010CO, and FONARSEC-FITR Industria 2013 N°

391 9 (ANPCyT). The laboratory of Aguas de los Andes S. A. provided lab facilities and the Instituto de Geología y
392 Minería (Universidad Nacional de Jujuy) complemented field logistics. The authors thank Ing. Maria Silvina
393 Muhana Senn and Mr. Luis Omar Villegas from the Agua de los Andes laboratory for analytical advising, and
394 also inhabitants from the communities of the Jujuy's Puna for their hospitality. Special thanks to Captain
395 Christian Meuric for large field logistical support.

396

397 **References**

398 Allmendinger W, Jordan E, Kay M, Isacks BL (1997) The evolution of the Altiplano-Puna plateau of the Central
399 Andes. *Annu. Rev. Earth Planet. Sci.*, 25, 139-174.

400 Alonso RN (1986) Ocurrencia, Posición Estratigráfica y Génesis de los Depósitos de Boratos de la Puna
401 Argentina. PhD Thesis, Universidad Nacional de Salta, 196 p.

402 Alonso H, Risacher F (1996) Geoquímica del Salar de Atacama, Parte 1: origen de los componentes y balance
403 salino. *Revista Geológica de Chile*, 23, 127-136.

404 Alonso R, Gutiérrez R, Viramonte J (1984) Puna Austral - Bases para el subprovincialismo geológico de la Puna
405 Argentina. 9th Congreso Geológico Argentino, 1, 43-63.

406 An JW, Kang DJ, Tran KT, Kim MJ, Lim T, Tran T (2012) Recovery of lithium from Uyuni salar brine.
407 *Hydrometallurgy*, 64-70, 117-118.

408 Arias J, Viera O (1982) Estratigrafía y tectónica de la comarca Olacapato - Tuzgle, provincias de Jujuy y Salta,
409 República Argentina. *Revista del Instituto de Ciencias Geológicas*, 5, 71-86, Jujuy.

410 Bevacqua PSJ (1992) Geomorfología del Salar de Atacama y estratigrafía de su núcleo y delta, Segunda Región de
411 Antofagasta, Chile. Undergraduate thesis (unpublished). Universidad Católica del Norte, Antofagasta,
412 284 p.

413 Boschetti T, Cortecci G, Barbieri M, Mussi M (2007) New and past geochemical data on fresh to brine waters of
414 the Salar de Atacama and Andean Altiplano, northern Chile. *Geofluids*, 7, 33-50.

415 [Boutt DF, Hynke SA, Munk LA, Corenthal LG \(2016\). Rapid recharge of fresh water to the halite-hosted brine
416 aquifer of Salar de Atacama, Chile. *Hydrological Processes*, 30 \(25\), 4720-4740.](#)

417 Campbell, MG (2009) Battery lithium could come from geothermal waters. *The New Scientist*, 204 (2738), 23.

418 Carmona V, Pueyo JJ, Taberner C, Chong, G, Thirlwall M (2000) Solute inputs in the Salar de Atacama (N.
419 Chile). *Journal of Geochemical Exploration*, 69-70, 449-452.

420 Clesceri LS, Greenberg AE, Eaton AD (1998) Standard Methods for the Examination of Water and Waste Water.
 421 Amer. Public. Health. Assoc., 20th Edition, USA.

422 COCHILCO (2013) Compilación de informes sobre mercado internacional del litio y el potencial del litio en
 423 salares del norte de Chile, <http://www.cochilco.cl/estudios/info-litio.asp>

424 Coira B, Kay S, Viramonte JG (1993) Upper Cenozoic magmatic evolution of the Argentine Puna. A model for
 425 changing subduction geometry. *International Geology Review*, 35, 677-720.

426 Coira B, Kay S, Pérez B, Woll B, Hanning M, Flores P (1999) Magmatic sources and tectonic setting of
 427 Gondwana margin Ordovician magmas, northern Puna of Argentina and Chile, in: Ramos VA, Keppie D
 428 (Eds.), *Laurentia Gondwana Connections before Pangea*. Geological Society of America, Special Paper
 429 336, 145-171.

430 Coira B, Caffè PJ, Ramirez A, Chayle W, Diaz A, Rosas S, Perez B, Orozco O, Martinez M (2004) Hoja Geológica
 431 2366-I/2166-III: Mina Pirquitas, Provincia de Jujuy, 1:250,000. Servicio Geológico Minero Argentino,
 432 Buenos Aires.

433 [Corenthal LG, Boutt DF, Hynek SA, Munk LA \(2016\). Regional groundwater flow and accumulation of a massive](#)
 434 [evaporate deposit at the margin of the Chilean Altiplano. *Geophysical Research Letters*, 43 \(15\), 8017-](#)
 435 [8025.](#)

436 Crespo P, Palma H, Quintanilla J, Quispe L (1987) Tratamiento químico de salmueras del Salar de Uyuni, Potosí.
 437 UMSa-ORSTOM, Informe 7.

438 Cristiani C, Del Moro A, Matteini M, Mazzuoli R, Omarini R (1999) The Magmatism linked to the Jurassic –
 439 Cretaceous rift of NW Argentina: the Tusaquillas plutonic complex (Central Andes). 14th Congreso
 440 Geológico Argentino, 2, 190-193.

441 de Silva SL (1989) Altiplano-Puna volcanic complex of the central Andes. *Geology* 17, 1102-1106.

442 del Papa C, Hongn F, Powell J, Payrola P, Do Campo M, Strecker MR, Petrinovic I, Schmitt AK, Pereyra R
 443 (2013) Middle Eocene-Oligocene broken foreland evolution in the Andean Calchaquí Valley, NW
 444 Argentina: insights from stratigraphic, structural and provenance studies. *Basin Research*, 25, 574-593.

445 Erickson GE, Chong G, Vila T (1976) Lithium resources of salars in the Central Andes. In: Vine JD (ed.), *Lithium*
 446 *Resources and requirements by the Year 2000*. Geological Survey Professional Paper, 1005, 66-74.

447 Erickson GE, Salas R (1977) *Geology and resources of salars in the central Andes*. U.S. Geological Survey, Open
 448 File Repository 88-210, 51 p.

449 Ericksen GE, Vine JD, Ballon RA (1978) Chemical composition and distribution of lithium-rich brines in salar de
 450 Uyuni and nearby salars in southwestern Bolivia. *Energy*, 3 (3), 355-363.

451 Fornillo B, Zicari J, Slipak AM, Puente F, Argento M (2015) Geopolítica del litio: industria, ciencia y energía en
 452 Argentina. Ed. El Colectivo CLACSO, Buenos Aires, 212 p.

453 Franco MG, Borda L, García MG, López Steinmetz RL, Flores P, Córdoba F (2016) Geochemical and
 454 sedimentological characterization of the Salar de Olaroz, northern Argentinean Puna, Central Andes. 3rd
 455 International Workshop on Lithium, Industrial Minerals and Energy, Jujuy, Argentina.

456 Garrett DE (2004) Handbook of lithium and natural calcium chloride: their deposits, processing, uses and
 457 properties. Elsevier Academic Press, 1st edition.

458 Godfrey LV, Chan L-H, Alonso RN, Lowenstein TK, McDonough WF, Houston J, Li J, Bobst A, Jordan TE
 459 (2013) The role of climate in the accumulation of lithium-rich brine in the Central Andes. *Applied*
 460 *Geochemistry*, 38, 92-102.

461 González MA, Tchilinguirian P, Pereyra F, Ramallo E (2000) Hoja geológica 2366-IV: Ciudad del Libertador
 462 General San Martín, 1:250,000. Servicio Geológico Minero Argentino, Buenos Aires.

463 Gruber P, Medina P (2010) Global lithium availability: a constraint for electric vehicles? Master thesis, University
 464 of Michigan, 76 p.

465 Hardie L, Eugster H (1970) The evolution of closed-basin brines. Fiftieth Anniversary Symposia, Mineralogy and
 466 Geochemistry of Non-Marine Evaporites. Mineralogical Society of America Special Publication, 273-
 467 290.

468 Houston J, Butcher A, Ehren P, Evans K, Godfrey L (2011) The evaluation of brine prospects and the requirement
 469 for modifications to filing standards. *Economic Geology*, 106, 1225-1239.

470 Ide F, Kunasz IA (1989) Origin of lithium in Salar de Atacama, northern Chile, in: Ericksen GE, Cañas Pinochet
 471 MT, Reinemund JA (Eds.), *Geology of the Andes and Its Relation to Hydrocarbon and Mineral*
 472 *Resources*. Circum-Pacific Council for Energy and Mineral Resources, Earth Sciences Series, 11, 165-
 473 172.

474 Isacks BL (1988) Uplift of the Central Andean Plateau and bending of the Bolivian Orocline. *J. Geophys. Res.* 93,
 475 3211-3231.

476 [Jordan TE, Mpodozis C, Munoz N, Blanco N, Pananont P, Gardeweg M \(2007\). Cenozoic subsurface stratigraphy](#)
 477 [and structure of the Salar de Atacama Basin, northern Chile. *Journal of South American Earth Sciences*,](#)
 478 [23, 122-146.](#)

479 Kay SM, Coira B, Caffè PJ, Chen C-H (2010) Regional chemical diversity, crustal and mantle sources and
 480 evolution of central Andean Puna plateau ignimbrites. *Journal of Volcanology and Geothermal Research*,
 481 198 (1-2), 81-111.

482 Kesler SE, Gruber PW, Medina PA, Keoleian GA, Everson MP, Wallington TJ (2012) Global lithium resources:
 483 relative importance of pegmatites, brine and other deposits. *Ore Geology Review*, 48, 55-69.

484 King M (2010) Amended inferred resource estimation of lithium and potassium at the Cauchari and Olaroz salars
 485 Jujuy province, Argentina. Technical report, Groundwater Insight, Inc.

486 Kirschbaum A, Hongn F, Menegatti N (2006) The Cobres Plutonic Complex, eastern Puna (NW Argentina):
 487 Petrological and structural constraints for Lower Paleozoic magmatism. *Journal of South American Earth*
 488 *Sciences*, 21, 252-266.

489 Ide F, Kunasz IA (1989) Origin of lithium in Salar de Atacama, northern Chile, in: Ericksen, GE, Cañas Pinochet
 490 MT, Reinemund JA (Eds.), *Geology of the Andes and its relation to hydrocarbon and mineral resources*.
 491 Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, 11, 165-172.

492 Kunasz IA (2006) Lithium resources. In *Industrial minerals and rocks, commodities, markets and uses*. 7th edition.

493 López Steinmetz RL (2017) Lithium- and boron-bearing brines in the Central Andes: exploring hydrofacies on the
 494 eastern Puna plateau between 23° and 23°30'S. *Mineralium Deposita*, 52, 35-50.

495 Lowenstein T, Risacher F (2009) Closed basin brine evolution and the influence of Ca-Cl inflow waters. Death
 496 Valley and Bristol Dry Lake, California, Qaidam Basin, China, and Salar de Atacama, Chile. *Aquatic*
 497 *Geochemistry*, 15, 71-94.

498 Lupo L, Morales M, Maldonado A, Grosjean M (2006) A high-resolution pollen and diatom record from Laguna
 499 Los Pululos (22°36'S/66°44'W/4500 m asl), NW Argentinean Puna, since *ca.* 800 AD. *Reconstructing*
 500 *Past Regional Climate Variations in South America over the Late Holocene: A New PAGES Initiative*.
 501 International Symposium, Malargüe, Argentina, 24.

502 Maro G, Caffè PJ (2016) The Cerro Bitiche Andesitic Field: petrological diversity and implications for magmatic
 503 evolution of mafic centers from the northern Puna. *Bulletin of Volcanology*, 78-51, doi: 10.1007/s00445-
 504 016-1039-y

505 Maro G, Caffè PJ, Jofré C (2016) Xenolitos ultramáficos en andesitas máficas neógenas de la Puna Norte. *Revista*
 506 *de la Asociación Geológica Argentina*, 73, in press.

507 McGlue MM, Ellis G, Cohen A, Swarzenski P (2012) Playalake sedimentation and organic matter accumulation in
 508 an Andean piggyback basin: the recent record from the Cuenca de Pozuelos, NW
 509 Argentina. *Sedimentology*, 59,1237-1256.

510 McQuarrie N, Horton BK, Zandt G, Beck S, DeCelles PG (2002) Lithospheric evolution of the Andean fold-thrust
 511 belt, Bolivia, and the origin of the central Andean plateau. *Tectonophysics* 399 (1e4), 15e37.
 512 <http://dx.doi.org/10.1016/j.tecto.2004.12.013>.

513 Méndez V, Navarini A, Plaza D, Viera O (1973) Faja Eruptiva de la Puna oriental. 5th Congreso Geológico
 514 Argentino, 4, 89-100.

515 Mohr S, Mudd G, Giurco D (2010) Lithium Resources and Production: a critical global assessment. Prepared for
 516 CSIRO Minerals Down Under Flagship, by the Institute for Sustainable Futures, University of
 517 Technology, Sydney, and Department of Civil Engineering, Monash University, Final Report, 107 p.

518 Moraga A, Chong G, Fortt MA, Henríquez H (1974) Estudio geológico del Salar de Atacama. Provincia de
 519 Antofagasta. *Boletín del Instituto de Investigaciones Geológicas*, 29, Santiago, Chile, 56 p.

520 Munk LA, Hynek SA, Bradley D, Boutt D, Labay K, Jochens H (2016) Lithium brines: A Global Perspective.
 521 *Reviews in Economic Geology*, 18, 339-365.

522 Orberger B, Rojas W, Millot R, Flehoc C (2015) Stable isotopes (Li, O, H) combined with chemistry: powerful
 523 tracers for Li origins in Salar deposits from the Puna region, Argentina. *Procedia Earth and Planetary*
 524 *Science*, 13, 307-311.

525 Peralta Arnold YJ, Cabssi J, Tassi F, Caffè JP, Vaselli O (2017). Fluid geochemistry of a deep-seated geothermal
 526 resource in the Puna plateau (Jujuy Province, Argentina). *Journal of Volcanology and Geothermal*
 527 *Research*, doi: [10.1016/j.jvolgeores.2017.03.030](https://doi.org/10.1016/j.jvolgeores.2017.03.030).

528 Peralta Arnold YJ, Tassi F, Caffè PJ (2016) Hydrothermal systems and lithium deposits. Northern Puna, Jujuy
 529 Argentina. 3rd International Workshop on Lithium, Industrial Minerals and Energy, Jujuy, Argentina.

530 Perkins JP, Ward KM, de Silva SL, Zandt G, Beck SL, Finnegan NJ (2016) Surface uplift in the Central Andes
 531 driven by growth of the Altiplano Puna Magma Body. *Nature Communications*, doi:
 532 [10.1038/ncomms13185](https://doi.org/10.1038/ncomms13185)

533 Presta JF, Caffè PJ (2014) Historia eruptiva de los volcanes monogenéticos de El Toro (23°05'S–66°42'O), Puna
 534 norte, Argentina. *Andean Geology* 41:142-173

535 Rettig SL, Jones BF, Risacher F (1980) Geochemical evolution of brines in the Salar de Uyuni, Bolivia. *Chemical*
 536 *Geology*, 30, 57-79.

537 Risacher F, Fritz B (1991) Quaternary geochemical evolution of the Salar of Uyuni and Coipasa, Central
 538 Altiplano, Bolivia. *Chemical Geology*, 90, 211-231.
 539 Risacher F, Fritz B (2009) Origin of salt and brine evolution of Bolivian and Chilean Salars. *Aquat. Geochemistry*,
 540 15, 123-157.
 541 Risacher F, Alonso H, Salazar C (1999) Geoquímica de aguas en cuencas cerradas: I, II y III Regiones – Chile.
 542 Convenio de Cooperación ORSTOM – DGA – UCN – IRD, SIT 51, Santiago de Chile, 781 p.
 543 Risacher F, Alonso H, Salazar C (2003) The origin of brines and salts in Chilean Salars: a hydrochemical review.
 544 *Earth Sciences Revue*, 63, 249-292.
 545 Seggiaro RE (1994) Petrología, geoquímica y mecanismos de erupción del complejo volcánico Coranzulí. Ph.D.
 546 thesis, Universidad Nacional de Salta.
 547 Seggiaro RE (2013) Hoja geológica 2366-III: Susques, 1:250,000. Servicio Geológico Minero Argentino, Buenos
 548 Aires.
 549 Schmidt N (2010) Hydrological and hydrochemical investigations at the Salar de Uyuni (Bolivia) with regard to
 550 the extraction of lithium. *FOG*, 26.
 551 Schwab K (1973) Die Stratigraphie in der umgebung des salar Cauchari. *Geotekt. Forsch.*, Germany, 43, 168 p.
 552 Schwab K, Lippolt H (1974) K-Ar mineral ages and Late Cenozoic history of the Salar de Cauchari area
 553 (Argentine Puna). *International Association of Volcanic Chemistry, Earth's Interior Symposium, Andean*
 554 *and Antarctic Volcanological Problems*, Chile, 698-714.
 555 Shcherbakov AV, Dvorov VI (1970) Thermal waters as a source for extraction of chemicals. *Geothermics* 2 (2),
 556 1636–1639.
 557 Stoertz GE, Erickson GE (1974) Geology of salars in northern Chile. *Geological Survey professional paper* 811, 65
 558 p.
 559 Turner JC (1959) Estratigrafía del Cordón de Escaya y la Sierra de Rinconada, Jujuy. *Revista de la Asociación*
 560 *Geológica Argentina*, 15 (1), 15-39.
 561 Turner JC (1960a) Estratigrafía de la Sierra de Santa Victoria y adyacencias. *Boletín de la Academia Nacional de*
 562 *Ciencias*, 41 (2), 163-169, Córdoba.
 563 Turner JC (1960b) Estratigrafía del Nevado de Cachi y sector al oeste (Salta). *Acta Geológica Lilloana*, 3, 191-
 564 226.
 565 Turner JC (1972) Puna. *Geología Regional Argentina*, 1, 13-56. Buenos Aires.

566 Warren JK (2010) Evaporites through time: Tectonic, climatic and eustatic controls in marine and nonmarine
 567 deposits. *Earth Sciences Reviews*, 98 (3), 217-268.

568 Williams LB, Hervig RL (2002) Exploring intracrystalline boron isotope variations in mixed-layer illite/smectite.
 569 *Am. Miner.* 87.

570 Witherow RA, Lyons WB (2011) The fate of minor alkali elements in the chemical evolution of salt lakes. *Saline*
 571 *Systems*, 7 (2), 2-14.

572 Yaksic A, Tilton JE (2009) Using the cumulative availability curve to assess the threat of mineral depletion: The
 573 case of lithium. *Resources Policy*, 34, 185-194.

574 Zhang L, Chan LH, Gieskes JM (1998) Lithium isotope geochemistry of pore waters, Ocean Drilling Program
 575 Sites 918 and 919, Irminger Basin. *Geochim. Cosmochim. Acta*, 62.

576 Zappettini E (1989) Geología y metalogénesis de la región comprendida entre las localidades de Santa Ana y
 577 Cobres, Provincias de Jujuy y Salta. República Argentina. Ph.D. thesis, Universidad de Buenos Aires, 189
 578 p.

579

580

581 **Figure captions**

582

583 Fig.1. Map of the Andean Plateau (dark green area), showing the location of the Salar de Atacama, Uyuni and
 584 Hombre Muerto (white areas). Thick dashed grey lines indicate the internal Plateau boundaries between the
 585 Bolivian Altiplano, the Argentinean Northern and Southern Puna (Alonso et al. 1984). There are many other
 586 salars in the Chilean Desierto de Atacama, in the Bolivian Altiplano and in the Argentinean Puna Plateau that are
 587 not shown in this map.

588

589 Fig. 2. Simplified geologic map of the Northern Puna Plateau between 23°06'S and 23°56'S, modified from
 590 González et al. (2000), Coira et al. (2004), and Seggiaro (2013). The map shows the location of the Northern
 591 Puna's salars. A-A' is the altitudinal transect of the Northern Puna Plateau at 23°30'S. Lithological legend: 1)
 592 modern salar; 2) Quaternary alluvial deposits, including piedmonts, alluvial fans and alluvial plains; 3)
 593 sedimentary units including the Cretaceous-Paleocene rift related and the Upper Eocene to Upper Miocene
 594 synorogenic deposits; 4) Plio-Pleistocene ignimbrite; 5) Miocene-Pliocene ignimbrite; 6) Miocene dacite; 7)
 595 Jurassic-Cretaceous granite; 8) Ordovician granite; 9) Ordovician metasedimentary basement of the Northern

Puna; 10) Precambrian - Eocambrian metasedimentary basement of the Eastern Cordillera and Upper Cambrian sedimentary units.

Fig. 3. Maps showing the location of shallow brine sampling points in the Salar de Jama (A), Olaroz, Cauchari (B) and Salinas Grandes (C). Precise latitude and longitude of the samples are given in Table A of the electronic supplementary material.

Fig. 4. Ternary diagrams showing the relative ionic concentrations of brines from the Salar de Jama (JA), Olaroz (OL), Cauchari (CA) and Salinas Grandes (SG). Ionic concentrations are expressed in meq L⁻¹.

Fig. 5. Comparative concentrations of Ca²⁺, Mg²⁺, K⁺, Li⁺ and BO₃³⁻ (A) and electrical conductivity (B) of samples collected from the Salar de Jama (JA), Olaroz (OL), Cauchari (CA) and Salinas Grandes (SG).

Fig. 6. Scatter diagrams showing the variation of Na⁺ versus Cl⁻ (A) and Ca²⁺ versus SO₄²⁻ (B and C) in the studied brines. Abbreviations are: Jama (JA), Olaroz (OL), Cauchari (CA), Salinas Grandes (SG), Guayatayoc (G), Hombre Muerto (HM), Uyuni (UYU) and Atacama (ATA). The HM data were taken from Godfrey et al. (2013). The UYU samples UB and UE and the ATA data set are from Risacher and Fritz (2009). The UYU samples 109, 128 and 120 were taken from Ericksen et al. (1977). The G samples are from Lopez Steinmetz (2017) and samples JA, OL, CA and SG are from this contribution. The HM samples 10, 11 and 12 are lagoon brines, sample 13 is a brine collected at 200 cm depth in a hole drilled into the salar (Godfrey et al. 2013). The UYU samples UB and UE correspond to brines collected at 100 cm depth (Risacher and Fritz 2009), sample 109 was collected in a brine pool and samples 120 and 128 are brines collected from holes drilled 40 to 50 cm into the salt crust (Ericksen et al. 1977). Samples GG₃, GG₄ and GQ₀ were collected from pits dug in the playa lake sediments (Lopez Steinmetz 2017).

Fig. 7. Li grade zones in the four major salars of the Northern Puna Plateau. Brine Li concentrations are in mg L⁻¹.

Fig. 8. Scatter diagrams showing the variation of (Na⁺+Cl⁻) and Conductivity with Li⁺ (A) and Li⁺ versus Mg²⁺ (B and C) in the studied brines. Jama (JA), Olaroz (OL), Cauchari (CA), Salinas Grandes (SG). Data

corresponding to Guayatayoc (G), Hombre Muerto (HM), Uyuni (UYU) and Atacama (ATA) are plotted for comparison. Details concerning the HM, UYU, ATA and G plotted data are presented in caption of Fig. 6.

Fig. 9. Comparative Ca^{2+} , Mg^{2+} , K^+ , Li^+ and BO_3^{3-} concentration ranges of samples collected from the Salar de Jama (JA), Olaroz (OL), Cauchari (CA) and Salinas Grandes (SG). The TDS value was obtained from electrical conductivity, by employing a conversion factor of 1 ($\text{TDS} = \text{Conductivity}$) based on the relation between the concentration of all the analysed ions relative to conductivity (Fig. A of the electronic supplementary material).

Table 1. Chemical composition of brines samples collected in the Salar de Jama (JA), Olaroz (OL), Cauchari (CA) and Salinas Grandes (SG).

Table 2. Comparative attributes of the Northern Puna's salars and Atacama, Uyuni and Hombre Muerto. The mean Li^+ concentration and Li:Mg ratios of Jama, Olaroz, Cauchari and Salinas Grandes were settled based on original data from this contribution (Table 1). The mean Li^+ concentration and Li:Mg of Guayatayoc correspond to the mean value of the 3 brine samples (samples G3, G4 and Q0) reported in Lopez Steinmetz (2017). The mean Li^+ concentration reported for Atacama are from Ide and Kunasz (1989) and Kunasz (2006), and the mean Li:Mg corresponds to values reported by COCHILCO (2013). The mean Li^{+a} concentration and Li:Mg^a ratio in Uyuni corresponds to the mean values of 38 samples (from sample 100 to sample 133 and samples 146, 147 and 147A) reported by Ericksen et al. (1977). The mean Li^{+b} concentration and Li:Mg^b ratio corresponds to the mean values of the 164 samples (from sample UA to sample RZ) reported by Risacher and Fritz (1991). The mean Li^{+c} concentration and Li:Mg ratio of Hombre Muerto correspond to the mean value of the 9 samples (samples 4, 5, 6 and 9 to 14) reported by Godfrey et al. (2013). The mean Li^{+d} concentration in Hombre Muerto corresponds to the mean Li^+ concentration of 100 cm-deep brines reported by Garrett (2004) and Kesler et al. (2012).