First record of Sivameryx (Cetartiodactyla, Anthracotheriidae) from the lower Miocene of Israel highlights the importance of the Levantine Corridor as a dispersal route between Eurasia and Africa

Ari Grossman, Ran Calvo, Raquel López-Antoñanzas, Fabien Knoll, Gideon Hartman, Rivka Rabinovich

► **To cite this version:**

Ari Grossman, Ran Calvo, Raquel López-Antoñanzas, Fabien Knoll, Gideon Hartman, et al.. First record of Sivameryx (Cetartiodactyla, Anthracotheriidae) from the lower Miocene of Israel highlights the importance of the Levantine Corridor as a dispersal route between Eurasia and Africa. *Journal of Vertebrate Paleontology*, 2019, 39 (2), pp.e1599901. 10.1080/02724634.2019.1599901 . hal-02365607

HAL Id: hal-02365607

<https://hal.science/hal-02365607>

Submitted on 16 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First description of *Sivameryx* (Cetartiodactyla: Anthracotheriidae) from the Early Miocene of Israel highlights the importance of the Levantine Corridor as a dispersal route between Eurasia and Africa

Journal:	<i>Journal of Vertebrate Paleontology</i>
Manuscript ID	JVP-2018-0083.R3
Manuscript Type:	Article
Date Submitted by the Author:	11-Mar-2019
Complete List of Authors:	Grossman, Aryeh; Midwestern University, Anatomy; Midwestern University, College of Veterinary Medicine; Midwestern University, Arizona College of Osteopathic Medicine Calvo, Ran; Geological Survey of Israel López-Antoñanzas, Raquel; Institut des Sciences de l'Évolution de Montpellier, Université Montpellier, CNRS, IRD, EPHE. Knoll, Fabien; Fundación Conjunto Paleontológico de Teruel-Dinópolis, ; University of Manchester, School of Earth & Environmental Sciences, Hartman, Gideon; University of Connecticut, Anthropology Rabinovich, Rivka; The Hebrew University of Jerusalem, National Natural History Collections, Institute of Earth Sciences, Institute of Archaeology,
Key Words:	Israel, early Miocene, anthracotheriidae, cetartiodactyla, <i>Sivameryx</i> , Levantine corridor

SCHOLARONE™
Manuscripts

1 First **description** of *Sivameryx* (Cetartiodactyla, Anthracotheriidae) from the Early Miocene of
2
3
4
5
6 2 Israel highlights the importance of the Levantine Corridor as a dispersal route between Eurasia
7
8 3 and Africa
9

10
11 4 ARI GROSSMAN, *,^{1,2,3} RAN CALVO,⁴ RAQUEL LOPEZ-ANTONANZAS, ⁵ FABIEN KNOL, ^{6,7} GIDEON
12
13
14 5 HARTMAN, ^{8,9} and RIVKA RABINOVICH¹⁰
15

16
17 6 ¹Department of Anatomy, College of Graduate Studies, Midwestern University, Glendale, AZ
18
19 7 85308, USA, agross@midwestern.edu;
20
21

22 8 ²College of Veterinary Medicine, Midwestern University, Glendale, AZ 85308, USA;
23
24

25 9 ³Arizona College of Osteopathic Medicine, Midwestern University, Glendale, AZ 85308, USA;
26
27

28 10 ⁴Geological Survey of Israel, Jerusalem, Israel;
29
30

31 11 ⁵Institut des Sciences de l'Évolution de Montpellier (ISE - M, UMR 5554, CNRS/UM/IRD/EPHE),
32
33 12 c.c. 64, Université de Montpellier, Place Eugène Bataillon, F -34095 Montpellier Cedex 05,
34
35 13 France;
36
37

38 14 ⁶ARAID—Fundación Conjunto Paleontológico de Teruel-Dinopolis, 44002, Teruel, Spain;
39
40

41
42 15 ⁷School of Earth and Environmental Sciences, University of Manchester, Manchester, M13 9PL,
43
44 16 United Kingdom;
45
46

47
48 17 ⁸Department of Anthropology, University of Connecticut, Unit 1176, 354 Mansfield Road,
49
50 18 Storrs, CT 06269, USA;
51
52
53
54
55

56 * Corresponding author
57
58
59
60

19 ⁹Center for Environmental Sciences and Engineering, University of Connecticut, 3107 Horsebarn
20 Hill Road, Building 4 room 10, Storrs, CT 06269, USA;

21 ¹⁰National Natural History Collections, Institute of Earth Sciences, Institute of Archaeology, The
22 Hebrew University of Jerusalem, Berman Building, Edmond J. Safra Campus, Givat Ram,
23 Jerusalem 91904, Israel

25 RH: GROSSMAN ET AL.—FIRST ANTHRACOTHERE FROM ISRAEL

26

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT—The genus *Sivameryx* (Cetartiodactyla: Anthracotheriidae), found both in Asia and Africa, is considered of Asian origin. Recent excavations in the Negev region of southern Israel led to the discovery of a new Early Miocene site called Kamus Junction. Among the fossils recovered at Kamus Junction is an upper molar of *Sivameryx palaeindicus*. Although known species of *Sivameryx* have often been distinguished by size, comparisons of the new specimen with known *Sivameryx* teeth from Asia and Africa emphasize the need for caution when assigning *Sivameryx* fossils to species based on size alone. This record of *Sivameryx* highlights the importance of the Levant as a corridor connecting Eurasia and Africa. The new find, along with other recent finds, demonstrates that the Levantine Corridor facilitated faunal dispersal events that shaped modern biotas as early as the Early Miocene.

INTRODUCTION

Multiple mammalian orders and families appear to have migrated from Africa to Eurasia during the Early and Middle Miocene in a series of relatively rapid migration events (e.g. Cote et al., 2018; Leakey et al., 2011). An uncertain but substantial number of African taxa such as proboscideans migrated out of Africa roughly simultaneously (e.g. Sanders et al., 2010; Sanders and Miller, 2002). Direct fossil evidence for these migrations from sites that lie between Africa and Eurasia are rare but crucial for reconstructing biogeographic patterns of dispersal. The Levantine Corridor, located between Africa and Eurasia, provides the best case for studying such patterns (Tchernov et al., 1987). Preliminary descriptions recognized 19 fossil Early

1
2
3 48 Miocene mammal taxa from the Negev of Israel (Goldsmith et al., 1982; Savage and Tchernov,
4
5 49 1968; Tchernov et al., 1987). These taxa were primarily of African affinities but demonstrated
6
7
8 50 high levels of regional endemism, which, the authors argued, were probably indicative of
9
10
11 51 environmental and not temporal differences with other sites in Africa and Asia (Tchernov et al.,
12
13 52 1987).

14
15
16 53 Anthracotheres are relatively common in the Early Miocene fossil record of East and
17
18 54 North Africa (Evans et al., 1981; Leakey et al., 2011; Rowan et al., 2015) as well as Eurasia
19
20
21 55 (Antoine et al., 2013; Ducrocq and Lihoreau, 2006; Lihoreau and Ducrocq, 2007; Patnaik, 2013;
22
23 56 Savage et al., 1977; Sehgal and Bhandari, 2014) . Because anthracotheres are so widely
24
25
26 57 dispersed, paleontologists often regard them as useful indicators of early phases of
27
28 58 intercontinental dispersal events (e.g. Lihoreau and Ducrocq, 1995; Holroyd et al., 2010).
29
30
31 59 Anthracotheres appear in the early reports of mammals from the Miocene of Israel (Goldsmith
32
33 60 et al., 1982; Savage and Tchernov, 1968), and Goldsmith even named the main Early Miocene
34
35
36 61 locality from the Negev "Anthracothere Hill" (Goldsmith et al., 1988). However, the only formal
37
38 62 description of the Miocene mammals of the Negev in Israel, Tchernov et al. (1987: Table 7) does
39
40
41 63 not mention anthracotheres. Here we formally describe anthracothere remains from Israel for
42
43 64 the first time, specifically the material found at Kamus Junction.

44
45
46 65 In 2012, we began surveying and excavating new localities in Israel to reconstruct the
47
48 66 landscape and biota of the Levantine Corridor during the Miocene (López-Antoñanzas et al.,
49
50
51 67 2014; López-Antoñanzas et al., 2016; Rabinovich et al., 2014). During excavation of Kamus
52
53
54 68 Junction, a new Early Miocene locality, new specimens were recovered including

proboscideans, a giraffid (*Canthumeryx* sp.), and one anthracothere tooth, which we describe here.

Location and Geological Setting of Kamus Junction Locality

The Kamus Junction (KJ) locality is located at the southwestern side of the Yamin Plain, in the northern Negev desert of Israel. The KJ sediments were deposited in the Miocene Hazeva Formation. During most of the Early and Middle Miocene, the Neo-Tethys shoreline was located in Be'er Sheva area (Gvirtzman and Buchbinder, 1969), with the site some 30km southeast (Fig. 1).

During the Early Miocene large drainage systems flowed from deep-inland areas of the Arabian Plate, in the southeast, toward the Neo-Tethys, in the northwest. The Hazeva Formation consists mostly of fluvial, fine to coarse, sandstones, with some shale and conglomerates. Episodically, lacustrine environments prevailed, and marls and limestone were also deposited (Calvo and Bartov, 2001). Remnants of the Miocene fluvial-lacustrine Hazeva Formation are preserved in the northern Negev in Israel in synclinal basins within the folds of the 'Syrian Arc' folds system (Zilberman and Calvo, 2013) (Fig. 1). Several large unconformities separate the members in the formation (Fig. 2). In the Yamin Plain, the upper parts of the Zefa Member coat the basin edge, and are covered by cycles of layers of the Rotem Member (Fig. 1C, E). The Rotem Member, with a thickness of up to 1,000 m in the Arava Valley, contains low- as well as high-energy alluvial facies represented by conglomerate-sand-silt-clay cycles. Due to unconformities, during- and post-Hazeva time, only ~50m of the Rotem Member are preserved in Yamin Plain. In

KJ locality the lower part of the Rotem Member begins with 10m of coarse sandstone layers with cross-bedding, followed by 15m of alternating sandstone and shale-marls layers (Fig. 1E). The upper section begins with 1m thick gritstone, which continues into friable sandstone with petrified tree-trunks. HUJI KJ 31 was found eroding out of the sandstone along with fossils of other mammals, crocodilians and turtles.

Age of Kamus Junction (KJ)

Due to the absence of radiometric dateable markers within the Hazeva Formation, only its base and top ages are known. In central Sinai (Egypt) conglomerate of the base of the formation overlay dikes of ~25 Ma (K-Ar ages) (Steinitz et al., 1978), while in Jordan (east of the Yamin Plain) a ~20 Ma dike cuts the same stratigraphic unit (Calvo, 2000), indicating that the lower parts of the Hazeva Formation were likely deposited during the late Oligocene or Early Miocene (Calvo and Bartov, 2001). Based on geomorphological relationship between marine invasion and terrestrial erosional episodes in Be'er Sheva area, Bar and Zilberman (2016) concluded that the Hazeva Formation deposition ended before ~16-14 Ma.

Dental terminology—mostly follows Boisserie et al. 2010 and Bärmann and Rössner. 2011.

SYSTEMATIC PALEONTOLOGY

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

109

110 CETARTIODACTYLA Montgelard et al. 1997

111 ANTHRACOTHERIIDAE Leidy 1869

112 BOTHRIODONTINAE Scott, 1940

113 MERYCOPOTAMINI Lydekker, 1883

114 SIVAMERYX Lydekker, 1877

115 Type-species *Sivameryx palaeindicus* Lydekker, 1877

116 (included species: *S. africanus* Andrews 1914; *S. moneyi* Fourteau 1918)

117 Stratigraphic range: Early Miocene

118 **Diagnosis**—Following Pickford (1991); Lihoreau (2003), Holroyd et al. (2010), Lihoreau
119 et al. (2016). Medium to small in size; canines sexually dimorphic; p1 double rooted; lower
120 molars selenodont with anterior crests of labial cusps reaching lingual surface of crown, often
121 ending in a small cuspule; four crests from the metaconid; lingual cusps of lower molars
122 mediolaterally compressed; talonid of m3 loop-like and strongly obliquely oriented; upper
123 molars with four main cusps and a paraconule almost fused with protocone and of similar
124 height (=quasi-pentacuspitate); looplike parastyles and mesostyles; two distal crests from
125 protocone; symphysis reaches back to level of p1; no genial spine; postcanine diastema long,
126 with a flange-like protuberance leaning laterally; enamel microstructure is a mono-zonal radial
127 Schmelzmuster (not equivalent to the cetartiodactyl plesiomorph condition).

1
2
3 128 **New Material**—HUJI KJ 31: Right second molar (M2) (Fig. 2)
4
5

6 129 **Comparative Description**—the tooth is a nearly complete crown with a slightly broken
7
8
9 130 anterior (=mesial) cingulum and the anterior part of the roots also sheared. The molar is bucco-
10
11 131 lingually (21.8mm) wider than it is mesio-distally long (19.8mm). The crown is brachyodont and
12
13
14 132 the cusps are selenodont and high, giving the crown a tall appearance. There are four main
15
16 133 cusps (paracone, protocone, metacone, metaconule), and a fifth very small paraconule, a
17
18 134 distinctive character of *Sivameryx* referred to as quasi-pentacuspitate by some authors
19
20 135 (Holroyd et al., 2010; Pickford, 1991). The tips of the metacone and the paracone of HUJI KJ 31
21
22
23 136 pinch-in so that the tips of the cusps are close together. This pinching exaggerates the sloping
24
25
26 137 of the buccal sides of each buccal cusp contrasting with the less sloped lingual sides of the
27
28 138 lingual cusps. This is especially visible in the metacone. A looping mesostyle connects the
29
30 139 two buccal cusps. The parastyle is partially broken but was clearly also looping and prominent.
31
32
33 140 The metastyle of HUJI KJ 31 is relatively smaller than the mesostyle or parastyle. The
34
35
36 141 preparacristule is broad but slightly abraded. It wraps around the mesiolingual side of the
37
38 142 paracone to merge with the parastyle. Both the metacone and the paracone have distinct
39
40 143 bulbous ribs descending buccally; that of the paracone appears larger although this could be an
41
42
43 144 artefact of the degree of erosion of the metacone. A median transverse valley separates the
44
45
46 145 mesial and distal portions of the tooth. The narrow base of the valley is approximately at the
47
48 146 level of the cervix of the tooth. The median transverse valley is enclosed buccally by the
49
50
51 147 mesostyle, and continues lingually to terminate at the lingual cingulum. The mesial cingulum
52
53 148 begins at the mesial end of the parastyle where the preparacristule merges with the paracone.
54
55 149 The mesial cingulum is continuous with a very prominent and arcuate cingulum that surrounds
56
57
58
59
60

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

the protocone lingually. This protocone cingulum merges with a second arcuate cingulum that surrounds the metaconule, and the two cingula meet at the termination of the median transverse valley. The metaconule cingulum is continuous with a distal cingulum that extends to the distal part of the metastyle. Thus, the cingula form a single, shelf-like cingulum covering all but the buccal aspect of the tooth.

HUJI KJ 31 shares with other bothriodontine anthracotheres an upper molar mesostyle invaded by the median transverse valley and selenodont crescentric cusps. Anthracotheriinae differ by their accessory protostyle on the mesial cingulum, and relatively bunodont upper molar cusps. Micorbunodontinae differ because they lack the median transverse valley invading the mesostyle,

Members of the tribe Merycopotamini (*Afromeryx*, *Sivameryx*, *Hemimeryx*, *Merycopotamus*, *Libycosaurus*) are distinguished in their upper molars from other bothriodontine anthracotheres by a preparacrista that connects labially with the parastyle, lack of ectocristyle, and development of the molar entostyle (Lihoreau et al., 2016). All characters observed in HUJI KJ 31. We thus restrict further comparisons primarily to Merycopotamini with a few additions.

We place HUJI KJ 31 in *Sivameryx* because it is a five-cusped upper molar with a reduced paraconule (i.e. quasi-pentascuspidate). It also shares with *Sivameryx* a loop-like mesostyle and a strongly developed lingual cingulum of the protocone.

1
2
3 169 *Brachyodus* (Early Miocene of Africa and Europe, Middle Miocene of Thailand) differs
4
5
6 170 from HUJI KJ 31 because it has pinched rather than looping styles on its upper molars.
7
8 171 *Brachyodus* is also much bigger than HUJI KJ 31.
9

10
11 172 *Afromeryx* (Early Miocene of East and North Africa), *Hemimeryx* (Late Oligocene to late
12
13 173 Early Miocene of Pakistan), *Telmatodon* (Early Miocene of Palistan), *Gonotelma* (*Libycosaurus*
14
15 174 (Late Miocene of North Africa and Uganda), and *Merycompotamus* (late Middle Miocene to
16
17
18 175 Late Miocene of India and Pakistan) differ from HUJI KJ 31 because they are tetracuspitate and
19
20
21 176 their upper molars lack a paraconule.
22

23
24 177 *Elomeryx*, a bothriodontine genus with a widespread Eurasian and North American
25
26 178 distribution differs from HUJI KJ 31, by its much more developed paraconule, which is located
27
28
29 179 substantially farther from the protocone than seen in *Sivameryx*.
30
31

32
33 180 We recognize HUJI KJ 31 as a second upper molar (M2). The tooth has a distal wear
34
35 181 facet indicating that it is not an M3. Furthermore, the marked pinching of the tips of the
36
37 182 metacone and the paracone causing exaggerated sloping of the buccal sides of each buccal cusp
38
39 183 is particularly exaggerated on the metacone of HUJI KJ 31, a condition typically more marked on
40
41
42 184 M1 and M2. To determine if it is a first or second upper molar, we collected all available mesio-
43
44 185 distal length and bucco-lingual width measurements of upper molars assigned to *Sivameryx*
46
47 186 (Table 1). We used a simple bi-variate plot to compare HUJI KJ 31 with the other upper molars
48
49 187 belonging to *Sivameryx* species (Fig. 3). Recent discoveries of skulls of *S. africanus* (Rowan et al.,
50
51 188 2015) and *S. moneyi* (Miller et al., 2014) demonstrate that in both *S. africanus* and *S. moneyi*
52
53
54 189 the M1 is smaller than M2 and M3. The M2 and M3 overlap in size although the M3 is generally
55
56
57
58
59
60

larger than M2 (Fig. 3). HUJI KJ 31 falls within the cluster that includes M2 of *S. africanus*, *S. moneyi*, and *S. palaeindicus* (Fig. 3), and thus we recognize it as an M2.

DISCUSSION

Four Early Miocene genera of anthracotheres are known from Africa: *Brachyodus*, *Sivameryx*, *Afromeryx*, and *Jaggermeryx*. The distribution of the species of these genera suggests that African anthracotheres dispersed from Africa to Eurasia, while Eurasian anthracotheres dispersed to Africa. However, the exact pathways enabling such dispersals are unknown. *Jaggermeryx naida* is a monospecific genus identified only from dentaries and lower dentition found at Wadi Moghra in Egypt (Miller et al., 2014). As such, direct comparisons with HUJI KJ 31 are impossible. However, *J. naida* differs from known lower dentition of *Sivameryx* by its bunodont rather than selenodont dentition and overall greater size (Miller et al., 2014). *J. naida* is an African genus. Miller et al. (2014) suggested that this genus is derived from older African anthracotheres found in Oligocene deposits of the Fayum in Egypt.

Brachyodus is considered a descendant of *Bothriogenys* from the Oligocene of the Fayum in Egypt and thus of African origin (Holroyd et al., 2010). *Brachyodus* currently includes three species in Africa; *B. depreteti* from Wadi Moghara and Siwa in Egypt (Miller et al., 2014; Pickford, 1991), and from Sperrgebiet in Namibia (Pickford, 2008); *B. mogharensis* found only at Wadi Moghra (Miller et al., 2014; Pickford, 1991); and *B. aequatorialis* in various localities in East Africa (Kalodirr, Rusinga, Loperot, and Locherangan, Kenya, and Moroto, Uganda) (Holroyd et al., 2010). Records of *Brachyodus* in Eurasia include: *B. onoideus* from Early Miocene deposits

in southern Europe (Portugal, Spain, France; (e.g. [Antunes and Ginsburg, 2010](#); [Orliac et al., 2013](#)) and from Middle Miocene deposits in Thailand (Ducroq et al., 2003). The exact pathway and timing of the dispersal by *Brachyodus* from Africa into Eurasia is as yet undetermined.

Afromeryx zelteni is recognized in North Africa (Jebel Zelten, Libya), East Africa (Loperot, Buluk, Ombo, Nachola and Wayondo, Kenya), and in Ghaba, Sultanate of Oman ([Holroyd et al., 2010](#); [Pickford, 1991](#); [Thomas et al., 1999](#)). A second species *A. africanus* is known only from Jebel Zelten in Libya. Two new species *A. grex* and *A. palustris* were recently described from Wadi Moghara ([Miller et al., 2014](#)). No records of *Afromeryx* are known from Eurasia. Although there is disagreement about its exact ancestral relationships (e.g. [Pickford, 1991](#) vs. [Lihoreau and Ducroq, 2007](#)), some researchers argue that it is of Eurasian origin ([Holroyd et al., 2010](#); [Lihoreau and Ducroq, 2007](#); [Pickford, 1991](#)).

Three species, one Asian and two African, are currently included in the genus *Sivameryx*. The type species *S. palaeindicus*, known from several sites in India and Pakistan, is similar in size and morphology to *S. africanus*, found in East and North Africa (e.g. [Holroyd et al., 2010](#); [Lihoreau and Ducroq, 2007](#)). While some prefer to retain them as two geographically distinct species ([Holroyd et al., 2010](#)), others suggest that there is insufficient morphological evidence to distinguish between the two species and the earlier name *Sivameryx palaeindicus* should take precedence (Lihoreau and Ducroq, 2007). In addition to *Sivameryx africanus* identified in East Africa (Karungu, Rusinga, Chianda Uyoma, Kalodirr, [Holroyd et al., 2010](#); [Pickford, 1991](#); [Rowan et al., 2015](#)), and North Africa (Gebel Zelten, [Pickford, 1991](#)), and possibly in Tunisia (Oued Bazina, [Lihoreau, 2003](#)), the third species, *S. moneyi*, is known only from Wadi Moghara

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

in Egypt (Holroyd et al., 2010; Miller et al., 2014). The two African species are regarded as morphologically indistinct and are only separated by size differences expressed in the length of the mandibular molar row (~80mm in *S. africanus* vs. ~59mm in *S. moneyi*) (Pickford, 1991). *Sivameryx africanus* and *S. moneyi* are considered migrants from Eurasia to Africa (Holroyd et al., 2010; Lihoreau and Ducroq, 2007; Miller et al., 2014; Pickford, 1991).

In order to assign HUJI KJ 31 to a species we wanted to determine if it is similar in size to *S. moneyi* from Wadi Moghra or the larger species *S. palaeindicus* and *S. africanus*. HUJI KJ 31 is an upper molar, but we expect that the maxillary molars will mirror differences in the size of mandibular molars. However, our analysis of the upper molar dimensions of *Sivameryx* from Asia, East Africa, and North Africa (Fig. 3) does not support such size distinction. In our analysis, the smallest individuals are specimens assigned to *S. africanus* from Kalodirr (Holroyd et al., 2010; Rowan et al., 2015). At the same time, the *S. moneyi* specimens from Wadi Moghara have a size range as great as the range of upper molars from India, Pakistan, or Gebel Zelten, attributed to either *S. palaeindicus* or *S. africanus* (Fig. 3). Alternatively, the overlap in size of the upper molars of *S. moneyi*, *S. africanus*, and *S. palaeindicus* combined with very little morphological variation may indicate that *Sivameryx* is a monospecific genus. A complete phylogenetic analysis required to examine this alternative is outside the scope of this paper. We suggest that size alone, at least as expressed via molar size, is not sufficient to distinguish different species of *Sivameryx* without additional material from well-dated localities. We therefore assign the specimen from Israel as *S. palaeindicus* because we cannot separate any of the species by upper molar size or morphology and the name *S. palaeindicus* takes precedence in the literature.

Regardless whether the genus *Sivameryx* has one or three species, all authors agree that the genus is an Asian immigrant into Africa (Holroyd et al., 2010; Lihoreau and Ducrocq, 2007; Miller et al., 2014; Pickford, 1991; Rowan et al., 2015). HUJI KJ 31 is found in the Levantine Corridor, an area that physically connects the two continents. This discovery of *Sivameryx* in Israel is part of a growing body of evidence (e.g. ctenodactylid rodents: López-Antoñanzas et al., 2016) that mammals exploited this route to migrate between Asia and Africa as early as the Early Miocene.

ACKNOWLEDGMENTS

This research is supported by ISF Grant #925/16 (RR and RC). AG is supported by a grant from Midwestern University. We are very grateful to the two anonymous reviewers for their thorough and helpful comments and comments from the editor, Dr. M. Sanchez-Villagra, which greatly improved the final manuscript. We wish to thank G. Beiner for preparing the material and V. Gutkin for SEM of the specimen. The faunal collection is deposited at the National Natural History Collection of the Hebrew University of Jerusalem.

REFERENCES

1

2

3273 Andrews, C. W. 1914. On the Lower Miocene Vertebrates from British East Africa, collected by Dr. Felix

4

5274 Oswald. Quarterly Journal of the Geological Society 70:163-186.

6

7275 Antoine, P.-O., G. Métais, M. J. Orliac, J. Crochet, L. J. Flynn, L. Marivaux, A. R. Rajpar, G. Roohi, and J.-L.

8

9276 Welcomme. 2013. Mammalian Neogene biostratigraphy of the Sulaiman Province, Pakistan.

10

11277 Fossil mammals of Asia: Neogene Biostratigraphy and Chronology. Columbia University Press,

12

13278 New York:400 – 422.

14

15

16279 Antunes, M. T., and L. Ginsburg. 2010. The last Anthracothere Brachyodus onoideus (Mammalia,

17

18280 Artiodactyla) from westernmost Europe and its extinction.

19

20

21281 Bar, O., and E. Zilberman. 2016. Subsidence and conversion of the Dead Sea basin to an inland erosion

22

23282 base level in the early middle Miocene as inferred from geomorphological analysis of its ancient

24

25283 western fluvial outlet. Geomorphology 261:147 – 161.

26

27

28284 Bärmann, E. V., and G. E. Rössner. 2011. Dental nomenclature in Ruminantia: Towards a standard

29

30285 terminological framework. Mammalian Biology 76:762–768.

31

32286 Boisserie, J.-R., F. Lihoreau, M. Orliac, R. E. Fisher, E. M. Weston, and S. Ducrocq. 2010. Morphology and

33

34287 phylogenetic relationships of the earliest known hippopotamids (Cetartiodactyla,

35

36288 Hippopotamidae, Kenyapotaminae). Zoological Journal of the Linnean Society 158:325-366.

37

38

39289 Calvo, R. 2000. Stratigraphy and petrology of the Hazeva Formation in the Arava and the Negev:

40

41290 Implications for the development of the sedimentary basins and the morphotectonics of the

42

43291 Dead Sea Rift Valley: In *Geology*, Vol. PhD, pp. 113 (in Hebrew, with English abstract). The

44

45292 Hebrew University in Jerusalem.

46

47

48293 Calvo, R., and Y. Bartov. 2001. Hazeva Group, southern Israel: New observations, and their implications

49

50294 for its stratigraphy, paleogeography, and tectono-sedimentary regime. Israel Journal of Earth

51

52295 Sciences 50:71 – 99.

53

54

55

56

57

58

59

60

- 296 Cote, S., J. Kingston, A. Deino, A. Winkler, R. Kityo, and L. MacLatchy. 2018. Evidence for rapid faunal
297 change in the early Miocene of East Africa based on revised biostratigraphic and radiometric
298 dating of Bukwa, Uganda. *Journal of Human Evolution* 116:95 – 107.
- 299 Ducrocq, S., Y. Chaimanee, V. Suteethorn, and J.-J. Jaeger. 2003. Occurrence of the anthracotheriid
300 *Brachyodus* (Artiodactyla, Mammalia) in the early Middle Miocene of Thailand. *Comptes Rendus*
301 *Palevol* 2:261-268.
- 302 Ducrocq, S., and F. Lihoreau. 2006. The occurrence of bothriodontines (Artiodactyla, Mammalia) in the
303 Paleogene of Asia with special reference to *Elomeryx*: Paleobiogeographical implications.
304 *Journal of Asian Earth Sciences* 27:885 – 891.
- 305 Evans, E. N., J. A. Van Couvering, and P. Andrews. 1981. Palaeoecology of Miocene sites in western
306 Kenya. *Journal of Human Evolution* 10:99 – 116.
- 307 Fourteau, R. 1918. Contribution a l'etude des vertebres miocenes de l'Egypte. Geological Survey
308 Department, Ministry of Finance, Egypt, Cairo. 120p.
- 309 Goldsmith, N., E. Tchernov, L. Ginsburg, P. Tassy, and J. Van Couvering. 1982. Ctenodactylid rodents in
310 the Miocene Negev fauna of Israel. *Nature* 296:645 – 647.
- 311 Goldsmith, N. F., F. Hirsch, G. M. Friedman, E. Tchernov, B. Derin, E. Gerry, A. Horowitz, and G.
312 Weinberger. 1988. Rotem mammals and Yeroham crassostreids: stratigraphy of the Hazeva
313 Formation (Israel) and the paleogeography of Miocene Africa. *Newsletters on Stratigraphy*:73 –
314 90.
- 315 Gvirtzman, G., and B. Buchbinder. 1969. Outcrops of Neogene Formation in the Central and Southern
316 Coastal Plain, Hashephela and Beer Sheva'Regions, Israel. *Geological Survey of Israel Bulletin*
317 50:1 – 73.
- 318 Holroyd, P., F. Lihoreau, G. Gunnell, E. Miller, L. Werdelin, and W. Sanders. 2010. Anthracotheriidae.
319 *Cenozoic Mammals of Africa*. University of California Press, Berkeley:843 – 851.

1
2
3 320 Leakey, M., A. Grossman, M. Gutiérrez, and J. G. Fleagle. 2011. Faunal change in the Turkana Basin
4
5 321 during the late Oligocene and Miocene. *Evolutionary Anthropology: Issues, News, and Reviews*
6
7 322 20:238 – 253.
8
9
10 323 Leidy, J. 1869. The extinct mammalian fauna of Dakota and Nebraska including an account of some allied
11
12 324 forms from other localities, together with a synopsis of the mammalian remains of North
13
14 325 America. *Journal of the Academy of Natural Sciences of Philadelphia*, series 2 7:1-472.
15
16
17 326 Lihoreau, F. 2003. Systématique et paléoécologie des Anthracotheriidae [Artiodactyla; Suiformes] du
18
19 327 mio-pliocène de l'ancien monde: implications paléobiogéographiques. Ph.D. Thesis, Université
20
21 328 de Poitiers, Poitiers, France, 395pp.
22
23 329 Lihoreau, F., L. Alloing-Séguier, P.-O. Antoine, J.-R. Boisserie, L. Marivaux, G. Métais, and J.-L.
24
25 330 Welcomme. 2016. Enamel microstructure defines a major Paleogene hippopotamoid clade: the
26
27 331 Merycopotamini (Cetartiodactyla, Hippopotamoidea). *Historical Biology*:1 – 11.
28
29
30 332 Lihoreau, F., and S. Ducrocq. 2007. Family Anthracotheriidae; pp. 89 – 105 in D. R. Prothero, and S. E.
31
32 333 Foss (eds.), *The evolution of artiodactyls*. The Johns Hopkins University Press, Baltimore, MD.
33
34 334 López-Antoñanzas, R., V. Gutkin, R. Rabinovich, R. Calvo, and A. Grossman. 2014. The rodent fauna from
35
36 335 the Early Miocene of the Rotem Basin (Israel): African, Asian, both or neither. *Journal of*
37
38 336 *Vertebrate Paleontology* 34 (suppl.):170.
39
40
41 337 López-Antoñanzas, R., V. Gutkin, R. Rabinovich, R. Calvo, and A. Grossman. 2016. A Transitional Gundi
42
43 338 (Rodentia: Ctenodactylidae) from the Miocene of Israel. *PLoS One* 11:e0151804.
44
45
46 339 Lydekker, R. 1877. Notices of new and rare Mammals from the Siwaliks. *Records of the geological Survey*
47
48 340 of India 10:76-83.
49
50 341 Miller, E. R., G. F. Gunnell, M. A. Gawad, M. Hamdan, A. N. El-Barkooky, M. T. Clementz, and S. M.
51
52 342 Hassan. 2014. Anthracotheres from Wadi Moghra, early Miocene, Egypt. *Journal of Paleontology*
53
54 343 88:967 – 981.
55
56
57
58
59
60

- 344 Montgelard, C., F. M. Catzeflis, and E. Douzery. 1997. Phylogenetic relationships of artiodactyls and
345 cetaceans as deduced from the comparison of cytochrome b and 12S rRNA mitochondrial
346 sequences. *Molecular Biology and Evolution* 14:550-559.
- 347 Orliac, M. J., P.-O. Antoine, A.-L. Charruault, S. Hervet, F. Prodeo, and F. Duranthon. 2013. Specialization
348 for amphibiosis in *Brachyodus onoideus* (Artiodactyla, Hippopotamoidea) from the Early
349 Miocene of France. *Swiss Journal of Geosciences* 106:265-278.
- 350 Patnaik, R. 2013. Indian Neogene Siwalik mammalian biostratigraphy: an overview. *Fossil mammals of*
351 *Asia: Neogene Biostratigraphy and Chronology*. Columbia University Press, New York:423 – 444.
- 352 Pickford, M. 1991. Revision of the Neogene Anthracotheriidae of Africa. *The Geology of Libya*:1491 –
353 1525.
- 354 Pickford, M. 2008. Anthracotheriidae from the Early Miocene deposits of the northern Sperrgebiet,
355 Namibia. *Geology and palaeobiology of the Northern Sperrgebiet, Namibia*. Geological Survey of
356 Namibia, Memoir 20:343-348.
- 357 Rabinovich, R., A. Grossman, H. Ginat, Y. Avni, and R. Calvo. 2014. Newly discovered Miocene
358 proboscideans in the southern Levant. *Journal of Vertebrate Paleontology* 34 (suppl.):210.
- 359 Rowan, J., B. Adrian, and A. Grossman. 2015. The first skull of *Sivameryx africanus* (Anthracotheriidae,
360 Bothriodontinae) from the early Miocene of East Africa. *Journal of Vertebrate Paleontology* 35:3
361 e928305.
- 362 Sanders, W. j., E. Gheerbrant, J. M. Harris, H. Saegusa, and C. Delmer. 2010. Proboscidea; pp. 161 – 251
363 in L. Werdelin, and W. J. Sanders (eds.), *Cenozoic Mammals of Africa*. University of California
364 Press.
- 365 Sanders, W. J., and E. R. Miller. 2002. New proboscideans from the early Miocene of Wadi Moghara,
366 Egypt. *Journal of Vertebrate Paleontology* 22:388 – 404.
- 367 Savage, R., and E. Tchernov. 1968. Miocene mammals of Israel. *Proc Geol Soc London* 1648:98 – 101.

1
2
3 368 Savage, R. J., P. C. Dixit, and D. Murty. 1977. On an anthracothere upper molar from Ladakh, Kashmir.
4
5 369 Journal of the Palaeontological Society of India 20:219 – 223.
6
7 370 Sehgal, R., and A. Bhandari. 2014. Miocene mammals from India: present status and future prospects.
8
9 371 Indian Miocene: a geodynamic and chronological framework for palaeobiota, sedimentary
10
11 372 environments and palaeoclimates. Palaeontol. Soc. India Spec. Publ 5:199 – 212.
12
13 373 Steinitz, G., Y. Bartov, and J. Hunziker. 1978. K-Ar age determinations of some Miocene–Pliocene basalts
14
15 374 in Israel: their significance to the tectonics of the Rift Valley. Geological Magazine 115:329 – 340.
16
17 375 Tchernov, E., L. Ginsburg, P. Tassy, and N. Goldsmith. 1987. Miocene mammals of the Negev (Israel).
18
19 376 Journal of Vertebrate Paleontology 7:284 – 310.
20
21 377 Thomas, H., R. Jack, S. Sen, M. Pickford, E. Gheerbrant, Z. Al-Sulaimani, and S. Al-Busaidi. 1999.
22
23 378 Oligocene and Miocene terrestrial vertebrates in the Southern Arabian peninsula (Sultanate of
24
25 379 Oman) and their geodynamic and paleogeographic settings; pp. 430-442 in P. J. Whybrow, and
26
27 380 A. Hill (eds.), Fossil vertebrates of Arabia. Yale University Press, New Haven, CT.
28
29 381 Zilberman, E., and R. Calvo. 2013. Remnants of Miocene fluvial sediments in the Negev Desert, Israel,
30
31 382 and the Jordanian Plateau: Evidence for an extensive subsiding basin in the northwestern
32
33 383 margins of the Arabian plate. Journal of African Earth Sciences 82:33 – 53.
34
35 384
36
37
38
39
40
41
42
43 385 Submitted July 14, 2018; accepted Month DD, YYYY
44
45
46
47
48
49
50
51
52
53 388

FIGURE CAPTIONS

FIGURE 1. Location map of the Kamus Junction (KJ) locality in the Yamin Plain, northern Negev, Israel.

A. General location of northern Negev, Israel.

B. Hazeva Formation outcrops in the northern Negev. Kamus Junction area is marked by a blue star.

C. The Hazeva Formation fills the Yamin Plain with all of its members. In the Kamus Junction area the upper parts of Zefa Member are overlain by clastic units of the Rotem Member. Small unconformity separates the lower and upper parts of the Rotem Member. Green dot marks the location of the upper level of KJ site where HUJI KJ 31 was found.

D. Topographic elevation of KJ site where HUJI KJ 31 was found (Green dot).

E. Columnar stratigraphic section in the Kamus Junction locality.

[Intended for Full page width]

FIGURE 2. SEM and schematic drawing of HUJI KJ 31—M2 of *Sivameryx palaeindicus* in occlusal view. Arrow on schematic indicates the paraconule.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

408 [Intended for 2/3 page width]

409

410 FIGURE 3. Scatter plot of upper molars of *Sivameryx* spp.

411 KEY: Triangles = M1, Squares = M2, Trapezoid = M3, Cross = HUJI KJ 31

412 Grey fill = *S. moneyi*, Dark fill = *S. africanus*, White fill = *S. palaeindicus*

413 Measurements listed in Table 1

414 [Intended for 2/3 page width]

415

416

TABLE 1. Measurements of *Sivameryx* spp. Upper molars

Specimen No.	MD	BL	Tooth position	Reference	Site
KJ 31	19.8	21.8	M2	This paper	Kamus Junction
<i>Sivameryx moneyi</i>					
CUWM 64AB	19.92	18.27	M1	Miller et al. 2014	Wadi Moghara
CUWM 64AB	23.08	21.18	M2	Miller et al. 2014	Wadi Moghara
CUWM 64AB	23.44	22.46	M3	Miller et al. 2014	Wadi Moghara
CUWM 70	17.5	17.5	M1	Miller et al. 2014	Wadi Moghara
CUWM 70	22.11	21.55	M2	Miller et al. 2014	Wadi Moghara
DPC 4066	21.47	22.34	M2	Miller et al. 2014	Wadi Moghara
DPC 4066	24.04	24.86	M3	Miller et al. 2014	Wadi Moghara
DPC 5979	20.71	20.89	M2	Miller et al. 2014	Wadi Moghara
DPC 6243	15.5	16.35	M1	Miller et al. 2014	Wadi Moghara
DPC 6243	23.12	20.81	M2	Miller et al. 2014	Wadi Moghara
DPC 6289	21.1	21.32	M3	Miller et al. 2014	Wadi Moghara
DPC 7659	25.35	22.89	M3	Miller et al. 2014	Wadi Moghara

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLE 1. (Continued)

DPC 17685	20.69	20.08	M2	Miller et al. 2014	Wadi Moghara
DPC 17685	22.23	22.12	M3	Miller et al. 2014	Wadi Moghara
DPC 21506	27.49	30.03	M3	Miller et al. 2014	Wadi Moghara

Sivameryx africanus

WK 18129	12.4	14.7	M1	Rowan et al. 2015	Kalodirr
WK 18129	11.6	15.5	M1	Rowan et al. 2015	Kalodirr
WK 18129	13.6	18.3	M2	Rowan et al. 2015	Kalodirr
WK 18129	14.6	18.7	M2	Rowan et al. 2015	Kalodirr
WK 18129	19.9	20.3	M3	Rowan et al. 2015	Kalodirr
WK 18129	16.6	18.9	M3	Rowan et al. 2015	Kalodirr
WK 17109	10.9	11.9	M1	Holroyd et al. 2010	Kalodirr

<i>S. africanus</i>	22.9	20.8	M2	Lihoreau et al. 2016	Gebel Zelten, or Rusinga or Karungu
---------------------	------	------	----	-------------------------	---

Sivameryx palaeindicus

Ladakh Molar	24	26	M2	Savage et al. 1977	Ladakh
B82	22	25	?M3	Savage et al. 1977	Laki Hills
B82	20.5	22.5	?M2	Savage et al. 1977	Laki Hills
TABLE 1. (Continued)					
M12738	24	25	M3	Savage et al. 1977	Dera Bugti
B482	19	20	?M1	Savage et al. 1977	Bugti Hills
B482a	21	22.5	?M2	Savage et al. 1977	Bugti Hills
Y238	19	18	M1	Lihoreau 2003	Potwar Y27926

FIGURE 1. Location map of the Kamus Junction (KJ) locality in the Yamin Plain, northern Negev, Israel.

A. General location of northern Negev, Israel.

B. Hazeva Formation outcrops in the northern Negev. Kamus Junction area is marked by a blue star.

C. The Hazeva Formation fills the Yamin Plain with all of its members. In the Kamus Junction area the upper parts of Zefa Member are overlain by clastic units of the Rotem Member. Small unconformity separates the lower and upper parts of the Rotem Member. Green dot marks the location of the upper level of KJ site where HUJI KJ 31 was found.

D. Topographic elevation of KJ site where HUJI KJ 31 was found (Green dot).

E. Columnar stratigraphic section in the Kamus Junction locality.

103x58mm (300 x 300 DPI)

FIGURE 1. Location map of the Kamus Junction (KJ) locality in the Yamin Plain, northern Negev, Israel.

A. General location of northern Negev, Israel.

B. Hazeva Formation outcrops in the northern Negev. Kamus Junction area is marked by a blue star.

C. The Hazeva Formation fills the Yamin Plain with all of its members. In the Kamus Junction area the upper parts of Zefa Member are overlain by clastic units of the Rotem Member. Small unconformity separates the lower and upper parts of the Rotem Member. Green dot marks the location of the upper level of KJ site where HUJI KJ 31 was found.

D. Topographic elevation of KJ site where HUJI KJ 31 was found (Green dot).

E. Columnar stratigraphic section in the Kamus Junction locality.

103x58mm (300 x 300 DPI)

FIGURE 2. SEM and schematic drawing of HUJI KJ 31—M2 of *Sivameryx palaeindicus* in occlusal view. Arrow on schematic drawing indicates the paraconule.

119x64mm (300 x 300 DPI)