Control of the mobilization of arsenic and other natural pollutants in groundwater by calcium carbonate concretions in the Pampean Aquifer, southeast of the Buenos Aires province, Argentina.

M. Vital^{1*}, D. E. Martinez¹, P. Babay³, S. Quiroga⁴, A. Clement², D. Daval²

¹Instituto de Geología de Costas y Cuaternario (UNMDP-CIC) – Instituto de Investigaciones Marinas y Costeras (CONICET-UNMDP). Mar del Plata, Argentina.

²Université de Strasbourg/EOST, CNRS, Laboratoire d'Hydrologie et de Géochimie de Strasbourg, 1 rue Blessig, F-67084 Strasbourg Cedex, France

³Comisión Nacional de Energía Atómica. Centro Atómico Constituyentes. Av. Gral. Paz 1499, B1650 Villa Maipú, Pcia de Buenos Aires

⁴ Departamento de Química, Universidad Nacional de Mar del Plata, Argentina

* corresponding autor : ingagr.melanie.vital@gmail.com

Submitted to Science of the Total Environment

GRAPHICAL ABSTRACT

HIGHLIGHTS

- The origin of As, F and other pollutants in groundwater is still misunderstood.
- Consumption of water with high content of As is a major threat for human health.
- Time-resolved experiments were used to study elemental release in water.
- As, V, Ba, Sr are probably coprecipitated with calcite and F adsorbed on calcite.
- The mobility of pollutants in the Pampean Aquifer is controlled by calcrete.

ABSTRACT

The water supply for human consumption in the Chaco-Pampean region in Argentina is restricted by the low quality of groundwater due to elevated concentrations of arsenic and other trace elements. Previous studies indicated a complex concurrence of factors and processes that are believed responsible to control the distribution of arsenic in groundwater. For a better understanding of the origin of trace elements in the Pampean aquifer, flow-through experiments with loess and calcrete samples representative of the sediments that constitute the aquifer were carried out in continuous flow reactors. The aqueous solutions were collected and the concentrations of $\text{SiO}_2(\text{aq})$, Ca^{2+} , SO_4^{2-} , Na^+ , Cl^- , F^- and trace elements (Ba, Sr, V, and As) were measured by inductively coupled plasma atomic emission spectroscopy (ICP-AES), high performance liquid chromatography (HPLC) and capillary electrophoresis. The experiments showed differences in the release rate of elements to the solution according to the type of sediment. The highest concentrations of V, Ba, and As were measured in experiments conducted with loess, and these elements were released quickly to the solution in the first minute of the test. In the case of loess, V and As are suggested to be adsorbed on the solid particles surface. Conversely, the experiments conducted with calcrete showed a lower but continuous release of those elements. This last result may indicate that the trace elements were coprecipitated in the calcite. In addition, it was demonstrated that F did not come from the dissolution of minerals such as fluorapatite, but both desorption from solid surface and dissolution from calcite minerals account for the release of F. This study support that both dissolution and adsorption-desorption processes can control the mobility of trace elements, with an emphasis on the role of calcrete in the retention and the mobilization of trace elements in the Pampean aquifer.

Keywords: Arsenic - carbonate concretions - loess - groundwater contamination –kinetic modeling – Pampean aquifer

INTRODUCTION

Arsenic dissolved in water is one of the most important concerns regarding drinking uses in many areas of the world. Naturally-occurring high arsenic (As) groundwater concentrations are widely distributed in some large regions of different countries, including the USA (Scanlon et al., 2009; Chappells et al., 2014), Mexico (Wurl et al., 2014), Argentina (Nicolli et al., 1989; Gomez et al., 2009), China (Guo et al., 2015), Chile (Sancha and O’Ryan, 2008), Vietnam (Erban et al., 2013), India (Chakraborti et al., 2016) and Bangladesh (Fendorf et al., 2010). The consumption of water with high concentration of As is related to arsenicosis, being an important threat for human health (Agusa et al., 2014; Joseph et al., 2015). About 140 million people in the world are exposed to As contaminated drinking waters. Other trace elements such as fluoride and vanadium are also of concern for human health. High fluoride concentrations in waters are associated with dental fluorosis and sometimes bone fluorosis (Sota de la et al., 1997), while vanadium can cause aneugenic effects and irritation of the respiratory tract (M. Costigan, 2001). Because of their important impact on human health, it is necessary to improve the understanding of the processes responsible for trace elements release and mobilization.

The mobility of arsenic in sedimentary environments has been studied in previous investigations, because high As concentration in drinking water is particularly common in arid or semi-arid regions (Robertson, 1989; Wyatt et al., 1998; Levy et al., 1999; Mählknecht et al., 2004, Farooqi et al., 2007; Currell et al., 2011, Wang et al., 2018; Yang et al., 2018). The desorption from Fe-(hydr)oxides was identified as the main process for As release in these areas under oxidizing environments. In the Pampean aquifer, the origin of As, F, Ba, Sr, and V, has been widely studied. Many authors attributed the high concentration of these elements with the presence of volcanic glasses (Nicolli et al., 1989; Smedley et al., 2002; Kruse and Ainchil, 2003; García et al., 2012; Auge, 2014). Other studies attributed the elevated

concentration of trace elements to the process of desorption from Al/Fe/Mn oxides (Smedley and Kinniburgh, 2002; Bhattacharya et al., 2006; Gomez et al., 2009). Nevertheless, volcanic glass shards are generally present in a homogeneous proportion in the Pampean sediments, whereas the concentrations of As, F and other trace elements show high variability throughout the territory (Blanco et al., 2012). For this reason, it is believed that other mechanisms could control the mobility of those elements in the groundwater.

The ability of calcite (CaCO_3) to retain inorganic contaminants has been described by many authors (Comans and Middelburg, 1987; Reeder, 1996; Tesoriero and Pankow, 1996; Cheng et al., 1999; Pokrovsky and Schott, 2002; Roman-Ross et al., 2003; Stipp et al., 2003; Renard et al., 2015). The uptake of contaminants in calcium carbonates can occur via adsorption on solid surface or coprecipitation reactions. Calcite has been shown to be capable of retaining such trace elements, retarding their transport, and thus plays an important role in the migration of contaminants in natural environments (Zachara et al., 1991; Stipp et al., 1992; Paquette and Reeder, 1995; Elzinga et al., 2002; Rouff et al., 2004). As banks of calcrete are extensively present in the Pampean sediments, adsorption or coprecipitation with calcite could explain most, or part, of the variability of As, F, V, Sr and Ba in the Pampean aquifer.

In the Chaco-Pampean plain of Argentina, the shallowest groundwater is contained in a sedimentary deposit of aeolian origin, which is one the most important sources of water supply for water to the population, agriculture and industry, and contributes to more than 60% of the country's gross national product (Schulz and Castro, 2003). The quality of the water from this reservoir depends on the hydrological balance and the interaction with the materials that constitute this aquifer. The quality of groundwater for drinking purposes varies greatly due to the presence of inorganic contaminants such as arsenic, fluoride, and other trace elements such as vanadium, strontium, and barium (Nicolli et al., 1989; Smedley et al., 2002). The negative effect of the consumption of water with high content of arsenic in this region is

well known (Hopenhayn-Rich et al., 1996; Bardach et al., 2015). In the province of Buenos Aires, arsenic concentration ranges from 23 to 289 $\mu\text{g.l}^{-1}$ (Al Rawahi et al., 2015) and the fluoride concentration is approximately 1.0 mg.l^{-1} (Kruse and Ainchil, 2003; Martínez et al., 2012; Calvi et al., 2016). In the central arid area of the country, the concentrations of arsenic and fluoride can be as high as 4.8 mg.l^{-1} (Bundschuh et al., 2004), and 28 mg.l^{-1} respectively (Smedley et al., 2002). In this area, the barium and vanadium concentrations are approximately 45 $\mu\text{g.l}^{-1}$ and 50 $\mu\text{g.l}^{-1}$, respectively (Smedley et al., 2002). The concentrations of these elements exceed the WHO guidelines values in many places (As: 10 $\mu\text{g.l}^{-1}$; F: 1.5 $\mu\text{g.l}^{-1}$), presenting risks for human health and limiting the use of groundwater resources (Nicolli et al., 1989; Bundschuh et al., 2000; Smedley et al., 2000; Fiorentino et al., 2007; Smedley et al., 2008; Nicolli et al., 2012a; Nicolli et al., 2012b). Whereas barium and strontium are present in solution as Ba^{2+} and Sr^{2+} cations, vanadium is generally found in solution as vanadate anions (HVO_4^{2-} , H_2VO_4^-), (Al Rawahi, 2016). Arsenic can be present in four oxidation states (-III, 0, + III, + V) but in the studied hydrological system, it is most commonly found as As (III) (H_3AsO_3 or H_2AsO_3^-) and As (V), (H_2AsO_4^- o HAsO_4^{2-}). The As (V) form is predominant at neutral and alkaline pH (Smedley et al., 2001). Those inorganic forms of arsenic (As(III) and As(V)) are highly toxic and mobile in the environment (Smedley and Kinniburgh, 2001; Jang et al., 2016).

Since the groundwater resources of the Pampean aquifer are threatened by the high toxicity of trace elements, it is important to improve the knowledge of their mobilization in this hydrologic system. In the present study, the putatively important role played by calcrete for the mobility of trace elements is investigated. Though the chemical composition of the water of the Pampean aquifer is commonly considered based on a thermodynamic equilibrium approach (Garrels and Christ, 1965; Glok Galli et al., 2014), it has been demonstrated in a previous paper that this assumption is questionable for some specific solid phases (Vital et al., 2018). To go beyond the thermodynamic equilibrium approach, flow-through experiments of interactions of loess and calcrete with water have been performed and coupled to reactive

transport simulations to identify the processes involved in the release of trace elements into the solution. In addition, while our previous study focused on the release of major elements, the present study aims at unraveling the source of trace elements released during the flow-through experiment with calcrete and loess samples.

DESCRIPTION OF THE STUDY SITE

The association of sediments occupying a surface area of more than one million square kilometers is known as the Pampean aquifer (Auge et al., 2002; Auge, 2008). The term "Pampean sediments" is an informal term and includes the Quaternary aeolian deposits of the Pampean region, consisting mainly of sand, silt, clay (Frenguelli and Teruggi, 1955). The sediments of the Pampean aquifer are considered as a formation of loess type. Unlike the "typical" loess, loess in Argentina has high contents of aluminosilicates and volcanic glass (Tricart, 1973; Pye, 1995; Zárate, 2003). In general, its composition is homogeneous, and most of its components originate from volcanic eruptions (Teruggi, 1957; Tricart, 1973). Calcareous formations occur as concretions and / or banks. Where the calcification is intense, banks of calcium carbonates result in sedimentary structures of the type of calcrete, locally denominated "tosca" (Sayago et al., 2001). The Pampean loess composition reveals 2 to 4% of calcium carbonate as calcite (Teruggi, 1993), mostly located in the upper part of the aquifer. Calcretes are widely abundant and common in the Pampean sediments, being well distributed superficially and at different depths. Calcrete is typical of the study site, forming continuous layers that have been related to paleo-surfaces (Teruggi et al., 1973), occupying the top of the hills. The calcretes may have formed over periods of several thousand years and are an indicator of semi-arid conditions (Zárate and Fasano, 1989). Calcrete layers are particularly abundant in the East of the Buenos Aires province, typically at a depth of around 1 m below ground level. This has resulted in the development of a largely flat-lying plain. This has been well described by Zárate and Folguera (2009) quoting original observations done by Charles

Darwin (Darwin, 1846). The hydrological conditions in the region are characterized by average annual precipitations around 900 mm, and a groundwater recharge estimated at 15% of total precipitation (Quiroz Londoño et al., 2012). The mean pH of the Pampean groundwater in the studied area is 7.15 (Quiroz-Londoño et al., 2008).

MATERIALS AND METHODS

1. Sample collection and mineralogical characterization

Samples of loess and calcrete were collected at an outcrop in the area of Mar del Plata, in the province of Buenos Aires, Argentina (38°05'12.7"S 57°38'57.1"W), (Figure 1). The area of sampling was chosen as being a representative area of the mineralogical composition of the Pampean aquifer in this sector of the Pampa plain. The outcrop includes both loess-like and calcrete sediments which were collected from a non-saturated zone with the same sedimentary composition as the Pampean aquifer. The samples were characterized in previous studies (Vital et al., 2016 ; Vital et al., 2018) using X-ray diffraction (XRD) coupled to Rietveld refinement and scanning electron microscopy coupled to energy dispersive X-ray spectroscopy (SEM/EDS). The analysis of the particle size distribution (PSD) by sieving revealed that loess is composed of 23 % of medium sand (0.25–0.5 mm), 15 % of fine sand (125–250 µm), 25% of very fine sand (62.5–125 µm) and 37 % of pelites (silt and clay, 0.997–62.5 µm). Calcrete samples were also collected in other areas of the province of Buenos Aires (Otamendi, Tandil and Tres Arroyos) in order to check the representativeness of the results obtained for As.

2. Flow-through experiments

The experiments were designed to monitor and measure the release of major and trace elements in the outflowing solution, as a function of reaction time with the solid samples of loess and calcrete (Figure 2). Continuous flow-through reactors simulate the conditions of an open geochemical system. The experiments were carried out using the set-up designed by

Cama et al. (2000). The Teflon-based continuous flow reactor of known volume of 21.2 cm³ was placed on a magnetic stir plate at room temperature. The solid samples of either loess or calcrete powders were introduced in the reactor. Through this reactor, deionized miliQ water (pH = 5.7) was continuously injected at a determined flow rate of 5 ml.min⁻¹. In this study, the objective of the experiment was to monitor the total release of the trace elements by the calcrete and the loess samples. Thus, pure milliQ water was used as a stock solution, to ensure that the input water is free of the trace elements that are under study. Importantly, the use of milliQ water is not intended to mimic rainwater or groundwater. This simplification was necessary to unravel the processes responsible for the release of the considered elements. Once such process are correctly singled out, the extrapolation to different real conditions (i.e. different rainwater or groundwater compositions) can be easily achieved by using geochemical modeling software. The solution inside the reactor was continuously stirred so that the concentrations and the temperature were theoretically homogeneous. The solution was maintained at far-from-equilibrium conditions with respect to the dissolving phases because of the continuous replenishment of the solution.

The experiment was designed to evidence the differences in elemental release between samples of loess and calcrete. It was divided into two steps: the first test T1 and the second test T2, in order to observe if the targeted elements are quickly and uniquely released to the solution over T1 (corresponding either to desorption, or to the disappearance of accessory minerals) or if they are continuously released over T1 and T2 (corresponding to the congruent dissolution of trace elements that were coprecipitated in major rock-forming minerals and/or minerals with slow dissolution kinetics). Drying the samples after collecting them is required to stop the dissolution reaction of the samples, and limit the possible precipitation of undesired secondary phases formed prior to the onset of the second experiment. In that way, it was possible to decipher whether the trace elements are adsorbed on solid particles or coprecipitated as impurities in some minerals. For this purpose, the

samples of loess and calcrete were introduced two times in the reactor. First, 2 grams of powder (loess or calcrete from the various above-mentioned locations) were introduced in the reactor for 2 hours. Sampling of the outflowing aqueous solution was carried out at the following duration steps: 1, 3, 5, 10, 15, 20, 30, 45, 60, 90 and 120 minutes. Those collected aqueous samples correspond to the first test T1. Afterwards, the solid samples were recovered at the end of the experiments, dried in an oven for 24 hours at 60°C. The dried solid samples were then put back in the flow-through reactor for two additional hours. The collection of aqueous samples in the second part of the experiment was carried out using the same previous time steps. The collected aqueous samples correspond to the second test T2. The experiments were repeated 3 times for each samples and each location. In all cases, the aqueous samples were filtered through a 0.45 µm filter and stored at -4°C prior to their analyses.

3. Aqueous sample analyses

The aqueous solution samples recovered at the reactor outlet, from the experiments realized with the loess and calcrete samples from the area of Mar del Plata, were analyzed to quantify the concentration of calcium, sodium, potassium and magnesium by capillary electrophoresis at the National Atomic Energy Commission (CNEA) laboratory (Buenos Aires, Argentina) with a Beckman Coulter brand equipment using a 40 cm-long fused silica capillary with a diameter of 75 µm (25 KV). The background electrolyte (BGE) was composed of a deimidazole solution of 10 mmol.l⁻¹, and acetic acid (40 mmol.l⁻¹). BGE was used as a buffer and is essential to stabilize the electro-osmotic flow and the migration time of the analytes. Calcium, sodium, potassium and magnesium standard solutions of 0.1, 0.5, 1, 5, 10, and 15 ppm were injected to determine the time after which each element appeared and to obtain the calibration curve of the peak area vs. the concentration. Sulfate, chloride and fluoride concentrations of the aqueous samples were measured by high performance liquid

chromatography (HPLC) with a Thermo Separation Products, CA, USA team, with a Spectra SERIES P200 binary pump, and a UV-Vis UV100 detector. An HPLC anion exchange column, PRP-X100, HAMILTON® and a mobile phase of potassium biphthalate ($4,5 \text{ mmol.l}^{-1}$) were used at the CNEA laboratory. Standard solutions of 0.5, 1, 2, 5, 10 and 15 ppm of fluorine, sulfate, and chloride were prepared to establish pattern curves in order to calibrate the concentration of each anion as a function of the area of the peak. The barium, vanadium, strontium, arsenic and silica concentrations of the aqueous samples were measured by inductively coupled plasma atomic emission spectroscopy (ICP-AES) with a Thermo ICAP 6000 brand equipment at the laboratory of hydrology and geochemistry of Strasbourg (France). In all analyses, pure milliQ water was used as a blank standard.

Arsenic concentrations of the aqueous samples obtained from the experiments T1 and T2 realized with the calcrete samples from the area of Tandil and Tres Arroyos were determined by Total reflection X-Ray Fluorescence (TXRF), using a TXRF spectrometer S2 Picofox (Bruker, Germany).

4. KIRMAT Modeling

Fluorapatite was identified in the loess and calcrete samples (Vital et al, 2018) and fluorine has been detected as an impurity in the calcrete with concentrations ranging between approximately $220 \text{ }\mu\text{g.g}^{-1}$ (García et al., 2006) and up to 1500 mg.kg^{-1} (Limbozzi, 2015). According to those observations, kinetic modeling was performed to assess the contribution of fluorapatite and fluoride-bearing calcite to the concentration of fluoride in the aqueous samples. Kinetic simulations coupled to mass transport were realized with the hydrochemical and thermo-kinetic code KIRMAT (Kinetic Reaction and Mass Transfer), (Gérard et al., 1998). A phase of calcite with 0.1% of fluoride as an impurity with a solubility identical to that of pure calcite was added to the KIRMAT thermodynamic database. The concentrations of fluoride obtained in the flow-through experiments test 1 (T1) and test 2 (T2) with the samples of loess

from Mar del Plata, and the samples of calcrete from the four locations, were compared to dissolution simulations of loess and calcrete containing fluorapatite or F-bearing calcite (0.1% of fluoride). The BET surface area of each sample from (Vital et al., 2018) was used as well as kinetic rate laws gathered in Palandri and Kharaka, (2004). Other input parameters for the simulations include deionized water in equilibrium with atmospheric CO_2 (corresponding to a starting pH of 5.7), a temperature of 25°C, and a flow rate of 5 $\text{ml}\cdot\text{min}^{-1}$. The thermodynamic database used in KIRMAT comes from the THERMODDEM database of the Bureau de Recherche Geologique et Minière (BRGM - <http://www.brgm.fr/>).

RESULTS

To determine the source of trace elements and their relations to major ions, the evolution of the output concentrations of major and minor elements obtained from the flow-through experiments conducted with loess and calcrete samples from the area of Mar del Plata are presented in Table 1 and Figure 3.

1. Experiments conducted on loess samples

In the experiment conducted on loess samples, the major ions $\text{SiO}_2(\text{aq})$, Ca^{2+} , Na^+ , SO_4^{2-} , Cl^- and K^+ are released in the solution following each of them a similar tendency, which consists of an initial peak of ion incorporation after the very first minutes of the experiment in the test T1. Then, the concentrations of $\text{SiO}_2(\text{aq})$, Ca^{2+} , Na^+ , SO_4^{2-} , Cl^- and K^+ decrease to reach a constant final value after 120 minutes (TF1). A lower concentration of those ions is measured in T2 but also starting with an initial peak of release. In detail, the highest concentration of silica (0.24 $\text{mmol}\cdot\text{l}^{-1}$) was measured in the sample collected during the first minute of contact with loess (table 1). Subsequently, the silica concentration dramatically decreases to reach 0.04 $\text{mmol}\cdot\text{l}^{-1}$ all over T1. In the second test(T2), a significant release of silica was observed in the first minute (0.21 $\text{mmol}\cdot\text{l}^{-1}$) followed by a very significant decrease. Sodium and chloride ions are released

very quickly to the solution, with the highest concentrations reached during the first minute of T1 (3.3 and 21 mmol.l⁻¹, respectively). A very significant decrease in the concentrations of these elements is observed to reach stable values in the first hour of T1 (0.2 and 0.01 mmol.l⁻¹). Over the course of T2, the values measured in the samples in the first minute are significantly lower than the values measured in the first minute of T1 (0.09 and 0.1 mmol.l⁻¹). The same trend is observed for calcium and sulfate ions, with the highest concentrations reached in the first minutes of T1 of 4.8 mmol.l⁻¹ and 0.7 mmol.l⁻¹.

The pH of the outflowing solution follows the patterns observed for the released ions. The pH of the initial input solution is 5.7. Initially, a fast increase from 5.7 to 6.6 is observed, but then pH decreases slowly to the initial input value after 180 min. The same behavior was observed for the TDS.

The behavior of trace elements is more complex (Figure3): while the concentration of some elements such as Sr and V follows a trend that is similar to that of major ions (i.e., initial peak after one minute in T1 and T2 followed by a decrease and the attainment of a steady-state concentration), the concentrations of other elements (e.g., Ba, F, As) steadily decrease throughout T1 and T2. The initial concentration of Sr²⁺ reached 8.72 µmol.l⁻¹ in T1, with a decrease down to 0.2 µmol.l⁻¹, followed by another peak in T2 at 0.3 µmol.l⁻¹. The concentration of vanadium is 0.2 µmol.l⁻¹ at the beginning of T1 and then, after an initial peak in T2 (0.07 µmol.l⁻¹), the final concentration of V is far below 0.1 µmol.l⁻¹. On the other hand, trace elements such as F⁻, Ba²⁺, and As start with a peak concentration in T1 (0.03 mmol.l⁻¹, 0.51 µmol.l⁻¹ and 0.027 µmol.l⁻¹ respectively) to decrease in all T1 and T2, and As concentration is even below the detection limit in T2. These results provide a first indirect indication that these elements were either released from specific minor phases enriched in these elements that were completely dissolved by the end of T2, or that they have been rapidly desorbed from the solid samples.

Further treatment of the data was carried out for the determination of the origin of other trace elements. While only Sr^{2+} is linearly correlated with Ca^{2+} release ($r^2 = 1$) (Figure 4), the concentrations of Sr^{2+} and Ba^{2+} in the aqueous samples are linearly correlated with that of Na^+ ($r^2_{\text{Na/Sr}} = 0.999$ and $r^2_{\text{Na/Ba}} = 0.96$) and K^+ ($r^2_{\text{K/Sr}} = 0.93$ and $r^2_{\text{K/Ba}} = 0.97$); (Figure 5). A lack of linear correlations between V and As concentrations and Na^+ or K^+ concentrations can be observed in Figure 5, as well as no linear correlation between Ca^{2+} and F^- in T1 ($r^2 = 0.69$) (Figure 6). On the contrary, a linear correlation is observed between the release of F^- and Ca^{2+} in the test T2 ($r^2 = 0.95$)

2. Experiments conducted on calcrete samples

In these experiments, the initial pH (5.7) quickly increased to stabilize at values comprised between 7.6 and 7.9, because of the CaCO_3 dissolution. These are the typical values of groundwater at the southeast of the province of Buenos Aires. The variation in concentrations in T1 and T2 of major ions are presented in Table 1. The highest concentration of calcium (0.64 mmol.l^{-1}) was obtained in the first minute of the test T1, and decreased down to 0.2 mmol.l^{-1} after 20 minutes of reaction. In the second test, a peak of Ca^{2+} appears in the first minute of reaction (0.37 mmol.l^{-1}). A peak of sodium (3.3 mmol.l^{-1}) and chloride (21 mmol.l^{-1}) is observed in the first minute of the experiment and then the concentrations stabilized below 0.5 mmol.l^{-1} and 0.1 mmol.l^{-1} respectively. In the test T2, a very similar trend is observed for the two elements. Similar trends were observed for $\text{SiO}_2(\text{aq})$ and SO_4^{2-} .

The experiments conducted with calcrete exhibited large release of strontium and barium during the first minutes of reaction in the first test (T1) and second test (T2), followed by a progressive decrease in the release rate of these ions during the subsequent hours (Figure 3). Of note, the maximum concentrations of Sr^{2+} and Ba^{2+} at the beginning of T2 are equivalent to those reached at the beginning of T1. Conversely, vanadium concentration peaks to a higher value in the first test ($0.9 \text{ } \mu\text{mol.l}^{-1}$) compared to the second test ($0.1 \text{ } \mu\text{mol.l}^{-1}$). Arsenic is

detected in T1 ($0.027 \mu\text{mol.l}^{-1}$) and only at the beginning of T2 ($0.013 \mu\text{mol.l}^{-1}$). A linear relation was evidenced between the concentration of calcium and the trace elements Ba^{2+} ($r^2 = 0.97$), As ($r^2 = 0.93$), and Sr^{2+} ($r^2 = 0.99$), (Figure 7a), and to a lesser extent, V ($r^2 = 0.84$). No linear relationship between Ca^{2+} and F^- concentrations was evidenced in T1. Conversely, the linear relationship was retrieved in T2 ($r^2 = 0.86$), (Figure 7b).

3. Confirmation of the control of As mobilization by calcite contained in calcrete

Flow-through experiments were repeated with calcrete from three other places of the province of Buenos Aires: Otamendi, Tres Arroyos, and Tandil. Arsenic concentrations were measured by TXRF, which is a method less sensitive than ICP-AES. Semi-quantitative values were obtained, indicating that arsenic was detected in almost all the aqueous samples (Table 2). In the first and second tests, the concentration ranged from 0.002 to 0.005 mg.l^{-1} ($0.026 \mu\text{mol.l}^{-1}$ to $0.067 \mu\text{mol.l}^{-1}$). This experiment showed that As is most certainly incorporated into calcite, as supported by the linear correlation between Ca^{2+} and As previously reported in Figure 7a.

4. Unravelling the source of fluoride using reactive transport modeling

The fluoride concentrations measured in T1 and T2 conducted with loess and calcrete were compared with the outputs of simulations of sediment (calcrete or loess) dissolution containing F-calcite and fluorapatite (Figure 8). Fluorapatite was previously detected by SEM observations reported in Vital et al. (2018) but could not be identified by XRD. Because it was estimated that the detection limit of our XRD was $\sim 2 \text{ wt. \%}$, we ran the kinetic simulations using this value as an upper bound for fluorapatite content in the solid sample. Also, calcite containing $0.1 \text{ wt. \%}$ of F was added (calcite represent $>90\%$ of the weight of the calcrete sample, and $1-2\%$ of the loess sample). Following Kitano and Okumura (1973), $0.1 \text{ wt. \%}$ is the

maximum concentration of F that can be incorporated into calcite. We used this value as an upper bound for the F content of calcite.

Figure 8a shows the results of the fluoride concentrations obtained in T1 and T2 flow-through experiments for the loess sample. As can be seen, the model successfully predicts the concentration of fluoride measured by the end of experiment T2. It further reveals that calcite is the main contributor to this agreement, as testified from the simulation performed with 0 wt. % of F-calcite and 2 wt. % of fluorapatite. Nevertheless, the first peak in T1 could not be fitted, which is an indirect evidence for a two-step process: (i) desorption of F at the beginning of the experiment T1 (not taken into account in the simulation), and (ii) continuous release from F-bearing phases.

A similar approach was used for calcrete from Mar del Plata, Otamendi, Tandil and Tres Arroyos flow-through experiments. The simulations were conducted with 2 wt. % of fluorapatite and considering F-bearing calcite instead of pure calcite (Figure 8b). The outputs of the simulations are in good agreement with the concentration plateau observed in T2, but, again, the initial peak of fluoride release at the beginning of T1 could not be fitted.

DISCUSSION

1. Experiments conducted on loess samples

An initial peak of major ion such as $\text{SiO}_2(\text{aq})$, Na^+ , Cl^- , Ca^{2+} is systematically observed in the flow-through experiments of loess in test 1. This may result from the rapid dissolution of the finest particles of the primary rock forming minerals (e.g., Schott et al., 1981). In T2 series, these elements are still released to the solution, with a steady-state concentration close to the concentration reached at the end of T1. Not surprisingly, these results indicate that these elements must have been released by mineral phases that were not completely consumed by the end of the T1, i.e., most likely the dissolution of primary rock-forming minerals. In a

previous study (Vital et al., 2018), we showed that the release of major ions from loess is related to the presence of minerals such as quartz, albite and accessory minerals such as halite, gypsum and calcite. An initial peak of strontium, vanadium, barium, fluoride and arsenic is also observed in T1. The concentrations of those elements decrease all along T1 and T2.

The presence of vanadium, strontium, barium and arsenic in groundwater has been related to loess deposits (Farías et al., 2003; Fiorentino et al., 2007; Nicolli et al., 2012a), mainly due to the release from volcanic glass shards and Al/Fe oxides (Steinhauser and Bichler, 2008; Nicolli et al., 2010; Nicolli et al., 2012a; Nicolli et al., 2012b; Bia et al., 2015; Barranquero et al., 2017; Alvarez and Carol, 2018), which could explain the behavior of the trace elements in the flow-through experiments T1 and T2 conducted on loess. The desorption of As and F from Fe/Mn oxides has been mentioned in many other studies focused on aquifers in China (Currell et al., 2011), or Korea (Kim et al., 2012), and Argentina (Smedley et al., 2002; Smedley et al., 2005; Bhattacharya et al., 2006; Borgnino et al., 2013). Currell et al., 2011 showed that increasing the duration of the interaction between water and loess samples generally did not result in an increase in F or As concentrations in solution and indicated that these ions are most likely mobilized by de-sorption rather than dissolution of F- or As-bearing minerals. Our results with the loess sample of the area of Mar del Plata are consistent with this previous study, suggesting that the rapid release of As could be the consequence of the desorption from those solid surfaces.

The results also show that only Sr is linearly correlated with Ca release. Vital et al., 2018 showed that the loess samples could contain up to 2% of calcite. Therefore, the strontium was likely coprecipitated with calcite, and its release to the solution results from calcite dissolution. In addition, the concentrations of Sr^{2+} and Ba^{2+} in the aqueous samples are linearly correlated with that of Na^+ and K^+ . The correlation of Ba^{2+} and Sr^{2+} with Na^+ and K^+ could also suggest that these trace elements were incorporated into the main rock-forming minerals of the loess

sample (feldspars). Conversely, the absence of linear correlations between V and As concentrations and Na^+ or K^+ concentrations supports the assumption that these trace elements were not released from the congruent dissolution of feldspars. Instead, it is suggested that they were carried by minor phases that have almost disappeared by the end of the experiment T2. Furthermore, Nicolli et al., (1989), Bhattacharya et al. (2006), Fiorentino et al. (2007), Heredia and Cirelli (2009), Puntoriero et al. (2015), showed that V is strongly correlated with As in groundwaters in the Pampean Region. This result indicates that those elements may have similar sources, mobilization and transport processes and could be mainly desorbed from Al/Mn oxides.

Finally, no linear correlation between Ca and F was evidenced in the T1 test, questioning the possibility that F was released congruently from a putative F-bearing calcite. Instead, the mechanism of fluoride release may be the rapid desorption from the particle's surface. Conversely, a linear correlation can be observed between Ca and F concentrations in the T2 test. Therefore, it is likely that the loess sample contains a second pool of fluoride, which might have been released through the congruent dissolution of a Ca-rich phase, such as the putative F-bearing calcite. The ability of calcite to be either a substrate for fluoride adsorption/desorption, or to incorporate fluoride during the course of its precipitation has been supported by the works of Kitano and Okumura (1973); Turner et al. (2005); Budyanto et al. (2015); Padhi and Tokunaga (2015).

2. Experiments conducted on calcrete samples

Calcium and trace elements were released all over the experiments conducted on calcrete, with an initial peak at the beginning of each test T1 and T2. The observed linear correlation between the concentration of calcium and the trace elements Ba, As, Sr and V, suggests that these elements were coprecipitated in calcite and their release results from calcite dissolution. This suggestion is supported by the work of Dietrich et al. (2016), who showed that carbonates

were the main As-bearing minerals in the carbonate-rich unsaturated zone from a large plain environment (Pampean plain, Argentina). Moreover, Renard et al. (2015) showed that As(V) was mostly incorporated in the calcite structure. When calcite dissolves, these elements are released stoichiometrically to the solution. In numerous studies, it has been demonstrated that calcite is able to retain trace elements, delaying their transport (Zachara et al., 1991; Stipp et al., 1992; Paquette and Reeder, 1995; Elzinga and Reeder, 2002; Rouff et al., 2004). The results obtained in the present work confirm the conclusions of these previous studies. An absence of linear relationship between Ca and F concentrations in the T1 test conducted with calcrete was observed, while this correlation was evidenced in the second T2 test. These latter results are very similar to those previously discussed for loess: At the beginning of T1, the main mechanism is F desorption from minerals surface (Kumar et al., 2016). The second mechanism, which prevails after the F on particles surface is totally desorbed, could be the release of fluoride that was incorporated in calcite.

3. Confirmation of the control of trace elements mobilization by calcite contained in calcrete

A significant difference was observed in the behavior of the trace elements released from loess or calcrete samples. The highest concentrations of vanadium, barium and arsenic have been measured for the experiments conducted with loess. These elements were released quickly to the solution in the first minute of T1 before decreasing sharply, while for calcrete, a continuous release is observed, and these trace elements are still released to the solution over the course of T2. This behavior was observed in four location of the province of Buenos Aires. The mechanisms of trace elements release from volcanic glass shards or Fe/Al/Mn oxides mentioned above could explain the higher concentration of trace elements in the aqueous samples collected for the experiments conducted with loess.

Conversely, trace elements may have been coprecipitated in calcite, which would explain their continuous release to the solution throughout the experiments conducted with calcrete. The experiments conducted with calcrete samples from different location (Tandil, Otamendi, Tres Arroyos) show that the release of trace elements from calcrete may apply throughout the Buenos Aires province. Importantly, the present results demonstrate that calcium carbonate may play a prevailing role in controlling the trace elements distribution for the study site. To the best of our knowledge, this aspect had never been proposed before in this area. Depending on the solution composition, calcium carbonates can therefore act as a sink for As if precipitation occurs, or as a source during dissolution. Alexandratos et al. (2007) suggested that arsenates are substituted in carbonate sites of calcite in most of the calcrete that is present in the Pampean aquifer. Sørensen et al. (2008) measured that the adsorption of arsenate increases with decreasing alkalinity, indicating a competition between arsenate and (bi)carbonate for sorption sites. These authors also observed that the mobility of arsenate in calcite bearing aquifers is greatly reduced due to sorption on calcite, whereas sorption on calcite in aquifers with low calcite content affects the mobility of arsenate to a lesser extent. Another study (Romero et al., 2004) focused on the carbonate-rich aquifer at Zimapán, México showed that carbonates play an important role in As mobility, and proved the high capacity of As(V) retention from this material. It was also shown that at alkaline pH (between 7 and 9), the carbonate-rich sample had a higher retention capacity of As.

Considering the presence of thick layers of calcrete and groundwater pHs around 7.5 to 8.0, it is possible to conclude that the coprecipitation and desorption from calcretes are factors controlling the As and other trace elements mobility and distribution in the aquifer. The experimental results showed that calcrete dissolution can be the most important source for As over long time periods, when the solution is undersaturated with respect to calcite. On the contrary, when the solution is supersaturated with respect to calcite, it can precipitate with trace-elements and act as a sink for these elements. It is expected that the calcite saturation

index of water samples could be considered as a proxy of As contents for some zones, as shown by Calvi et al. (2016), who reported a correlation between calcite supersaturation areas and low fluoride concentrations.

4. Unravelling the source of fluoride using reactive transport modeling

The results of the KIRMAT modeling showed that the neither fluorapatite nor F-calcite dissolution can account for the initial peak of F observed in T1 for loess and calcrete experiments, leaving F desorption as the best explanation for the observed F release when the sample is put in contact with water for the first time. Of note, several studies indicated that fluorine comes from surface desorption, as it was found adsorbed on calcite particles (Turner et al., 2005), iron oxides (Borgnino et al., 2013), or associated with volcanic glasses (Bundschuh et al., 2004; Bhattacharya et al., 2006; Nicolli et al., 2012a; García et al., 2014). These observations are in good agreement with the very fast release of fluorine to the solution and further suggest that when the sample was put in contact with water for the first time, the initial mechanism of F release is surface desorption.

Interestingly, the concentration of fluoride measured in the T2 for both loess and calcrete samples match the simulations of KIRMAT. This result confirms that in T2, a second mechanism of F release occurs and is most probably the release from F-bearing minerals such as fluorapatite and F coprecipitated in calcite. Therefore, we suggest that the two mechanisms of F release (desorption and dissolution from calcite and fluorapatite) may superimpose. Again, this confirms that calcrete plays an important role in the release of trace elements.

CONCLUSION

In this study, flow-through experiments associated to kinetic modeling were used in order to assess the specific role of calcite in the control of the mobility of trace elements in the Pampean aquifer. Samples of loess and calcrete from different areas of the province of Buenos

Aires (Mar del Plata, Tandil, and Tres Arroyos) were collected. In the experiments, the solid samples were first reacted with water in a flow-through set-up, dried, and then put again in contact with water. The chemical composition of the aqueous solution varied with the type of solid sample reacting (loess or calcrete). Regarding experiments conducted with loess, arsenic and other trace elements (V, Ba, Sr) were rapidly released during the first minute of reaction, likely due to their desorption from solid particles in loess. These observations agree with the results of other studies suggesting that the original source of those trace elements could be the volcanic glass or the Al/Fe/Mn oxides shards present in the loess sample. On the contrary, regarding experiments conducted with calcrete, those trace elements were released to the solution for several minutes, but also in the second experiment performed with the dried samples recovered from the first experiment. It has been shown in this work that arsenic, barium, vanadium and strontium have a strong correlation with Ca during the first part of the experiment, suggesting that they are released to the solution as a result of calcite dissolution. This work shows that calcite plays an important role in the migration of inorganic contaminants such as arsenic in the environment. The mobilization of contaminants from calcium carbonates can occur by desorption from the solid surface or by dissolution of the coprecipitated elements. In the Pampean aquifer, those trace elements can be coprecipitated in the calcrete banks and can be re-mobilized by calcite dissolution as a result of variation in physicochemical changes in the groundwater (i.e. pH, saturation index). For both loess and calcrete samples, fluorine release occurred through two different mechanisms. In a first step, the fluoride that is most likely adsorbed to the surface of the particles is released to the solution. Afterwards, fluorine coprecipitated in calcite is released to the solution. The use of reactive transport modeling allowed validating this model. In the future, it would be interesting to expand this study on loess and calcrete from different locations and different depths of the very great territory of Argentina. Because it was demonstrated in this study that trace elements adsorption/desorption and precipitation/dissolution from calcrete are

important processes impacting the groundwater chemistry, it would be necessary to include them in the geochemical simulations. This would allow to improve the knowledge of the mobilization and distribution of As and other trace elements in groundwater.

Acknowledgements:

The authors are thankful to the National Agency for Science and Technology Promotion (ANPCyT, PICT 2014 N°1529) which supported this study financially and to the National Scientific and Technical Research Council (CONICET). We also acknowledge the Mincyt-ECOS project for supporting international collaboration between the ICGyC/IIMyC in Argentina and the LHyGes in France. The authors are also thankful to Dr. Carlos Ayora and Dr. Jordi Cama (IDÆA, CSIC) for providing the Flow-through reactor and to Mr. G. Bernava for the chemical analyses. The authors thank Gilles Morvan (LHyGeS), and René Boutin (LHyGeS) for performing MEB, providing supplementary data and helping with ICP-AES measurements, respectively. This work has benefited from the constructive comments and insightful suggestions of 3 anonymous reviewers who are warmly thanked.

References

- Agusa, T., Trang, P. T. K., Lan, V. M., Anh, D. H., Tanabe, S., Viet, P. H., & Berg, M. (2014). Human exposure to arsenic from drinking water in Vietnam. *Science of the total Environment*, 488, 562-569. doi: <https://doi.org/10.1016/j.scitotenv.2013.10.039>
- Al Rawahi, W., Marcilla, A., & Ward, N. (2015). Arsenic Speciation Analysis of Ground Waters in Southern La Pampa and Buenos Aires Proviinces, Argentina.
- Al Rawahi, W. A. (2016). Vanadium, arsenic and fluoride in natural waters from Argentina and possible impact on human health (Doctoral dissertation, University of Surrey (United Kingdom)).
- Alexandratos, V. G., Elzinga, E. J., & Reeder, R. J. (2007). Arsenate uptake by calcite: macroscopic and spectroscopic characterization of adsorption and incorporation mechanisms. *Geochimica et Cosmochimica Acta*, 71(17), 4172-4187. doi: <https://doi.org/10.1016/j.gca.2007.06.055>
- Alvarez, M. d. P., & Carol, E. (2018). Geochemical occurrence of arsenic, vanadium and fluoride in groundwater of Patagonia, Argentina: Sources and mobilization processes. *Journal of South American Earth Sciences*. doi: <https://doi.org/10.1016/j.jsames.2018.10.006>
- Auge, M. (2008). Agua subterránea, deterioro de calidad y reserva. *Facultad de Ciencias Exactas y Naturales (UBA)*. http://sedici.unlp.edu.ar/bitstream/handle/10915/15908/Documento_completo.pdf?sequence=1&isAllowed=y
- Auge, M. (2014). *Arsenic in the groundwater of the Buenos Aires province, Argentina*. Paper presented at the One Century of the Discovery of Arsenicosis in Latin America (1914-2014) As2014: Proceedings of the 5th International Congress on Arsenic in the Environment, May 11-16, 2014, Buenos Aires, Argentina.
- Auge, M. P., Hernández, M. A., & Hernández, L. (2002). *Actualización del conocimiento del acuífero semiconfinado Puelche en la provincia de Buenos Aires, Argentina*. Paper presented at the XXXII IAH & VI ALHSUD Congress. Actas.
- Bardach, A. E., Ciapponi, A., Soto, N., Chaparro, M. R., Calderon, M., Briatore, A., Cadoppi, N., Tassara, R., & Litter, M. I. (2015). Epidemiology of chronic disease related to arsenic in Argentina: A systematic review. *Science of the total Environment*, 538, 802-816. doi: <https://doi.org/10.1016/j.scitotenv.2015.08.070>
- Barranquero, R. S., Varni, M., Vega, M., Pardi, R., & Ruiz de Galarreta, A. (2017). Arsenic, fluoride and other trace elements in the Argentina Pampean plain. *Geologica acta*, 15(3), 0187-0200. doi: 10.1344/GeologicaActa2017.15.3.3
- Bhattacharya, P., Claesson, M., Bundschuh, J., Sracek, O., Fagerberg, J., Jacks, G., Martin, R. A., Storniolo, A. d. R., & Thir, J. M. (2006). Distribution and mobility of arsenic in the Rio Dulce alluvial aquifers in Santiago del Estero Province, Argentina. *Science of the total Environment*, 358(1), 97-120. doi: <https://doi.org/10.1016/j.scitotenv.2005.04.048>
- Bia, G., Borgnino, L., Gaiero, D., & García, M. G. (2015). Arsenic-bearing phases in South Andean volcanic ashes: Implications for As mobility in aquatic environments. *Chemical Geology*, 393, 26-35. doi: <https://doi.org/10.1016/j.chemgeo.2014.10.007>
- Blanco, M. d. C., Paoloni, J. D., Morrás, H., Fiorentino, C., Sequeira, M. E., Amiotti, N. N., Bravo, O., Diaz, S., & Espósito, M. (2012). Partition of arsenic in soils sediments and the origin of naturally elevated concentrations in groundwater of the southern pampa region (Argentina). *Environmental Earth Sciences*, 66(7), 2075-2084. doi: <https://doi.org/10.1007/s12665-011-1433-x>
- Borgnino, L., Garcia, M., Bia, G., Stupar, Y., Le Coustumer, P., & Depetris, P. (2013). Mechanisms of fluoride release in sediments of Argentina's central region. *Science of*

- the total Environment, 443, 245-255. doi:
<https://doi.org/10.1016/j.scitotenv.2012.10.093>
- Budyanto, S., Kuo, Y.-L., & Liu, J. (2015). Adsorption and precipitation of fluoride on calcite nanoparticles: a spectroscopic study. *Separation and Purification Technology*, 150, 325-331. doi: <https://doi.org/10.1016/j.seppur.2015.07.016>
- Bundschuh, J., Bonorino, G., Viero, A., Albouy, R., & Fuertes, A. (2000). *Arsenic and other trace elements in sedimentary aquifers in the Chaco-Pampean Plain, Argentina: Origin, distribution, speciation, social and economic consequences*. Paper presented at the Arsenic in Groundwater of Sedimentary Aquifers, Pre-Congress Workshop, 31st Internat. Geol. Cong., Rio de Janeiro, Brazil.
- Bundschuh, J., Farias, B., Martin, R., Storniolo, A., Bhattacharya, P., Cortes, J., Bonorino, G., & Albouy, R. (2004). Groundwater arsenic in the Chaco-Pampean plain, Argentina: case study from Robles county, Santiago del Estero province. *Applied Geochemistry*, 19(2), 231-243. doi: <https://doi.org/10.1016/j.apgeochem.2003.09.009>
- Calvi, C., Martinez, D., Dapeña, C., & Gutheim, F. (2016). Abundance and distribution of fluoride concentrations in groundwater: La Ballenera catchment, southeast of Buenos Aires Province, Argentina. *Environmental Earth Sciences*, 75(6), 534. doi: <https://doi.org/10.1007/s12665-015-4972-8>
- Cama, J., Ganor, J., Ayora, C., & Lasaga, C. A. (2000). Smectite dissolution kinetics at 80 C and pH 8.8. *Geochimica et Cosmochimica Acta*, 64(15), 2701-2717. DOI: 10.1016/S0016-7037(00)00378-1
- Chakraborti, D., Rahman, M. M., Ahamed, S., Dutta, R. N., Pati, S., & Mukherjee, S. C. (2016). Arsenic groundwater contamination and its health effects in Patna district (capital of Bihar) in the middle Ganga plain, India. *Chemosphere*, 152, 520-529. doi: <https://doi.org/10.1016/j.chemosphere.2016.02.119>
- Chappells, H., Parker, L., Fernandez, C. V., Conrad, C., Drage, J., O'Toole, G., Campbell, N., & Dummer, T. J. (2014). Arsenic in private drinking water wells: an assessment of jurisdictional regulations and guidelines for risk remediation in North America. *Journal of water and health*, 12(3), 372-392. doi: <https://doi.org/10.2166/wh.2014.054>
- Cheng, L., Fenter, P., Sturchio, N. C., Zhong, Z., & Bedzyk, M. J. (1999). X-ray standing wave study of arsenite incorporation at the calcite surface. *Geochimica et Cosmochimica Acta*, 63(19-20), 3153-3157. doi: [https://doi.org/10.1016/S0016-7037\(99\)00242-2](https://doi.org/10.1016/S0016-7037(99)00242-2)
- Comans, R. N., & Middelburg, J. J. (1987). Sorption of trace metals on calcite: Applicability of the surface precipitation model. *Geochimica et Cosmochimica Acta*, 51(9), 2587-2591.
- Costigan, R. C., S. Dobson. (2001). Vanadium Pentoxide and other Inorganic Vanadium Compounds. Concise International Chemical Assessment Document 29. Geneva: World Health Organization, World Health Organization, 1-53.
- Currell, M., Cartwright, I., Ravaggi, M., & Han, D. (2011). Controls on elevated fluoride and arsenic concentrations in groundwater from the Yuncheng Basin, China. *Applied Geochemistry*, 26(4), 540-552. <https://doi.org/10.1016/j.apgeochem.2011.01.012>
- Dietrich, S., Bea, S. A., Weinzettel, P., Torres, E., & Ayora, C. (2016). Occurrence and distribution of arsenic in the sediments of a carbonate-rich unsaturated zone. *Environmental Earth Sciences*, 75(2), 90. doi: <https://doi.org/10.1007/s12665-015-4892-7>
- Elzinga, E., & Reeder, R. (2002). X-ray absorption spectroscopy study of Cu²⁺ and Zn²⁺ adsorption complexes at the calcite surface: Implications for site-specific metal incorporation preferences during calcite crystal growth. *Geochimica et Cosmochimica Acta*, 66(22), 3943-3954. doi: [https://doi.org/10.1016/S0016-7037\(02\)00971-7](https://doi.org/10.1016/S0016-7037(02)00971-7)
- Elzinga, E., Reeder, R., Withers, S., Peale, R. E., Mason, R., Beck, K. M., & Hess, W. P. (2002). EXAFS study of rare-earth element coordination in calcite. *Geochimica et Cosmochimica Acta*, 66(16), 2875-2885.

- Erban, L. E., Gorelick, S. M., Zebker, H. A., & Fendorf, S. (2013). Release of arsenic to deep groundwater in the Mekong Delta, Vietnam, linked to pumping-induced land subsidence. *Proceedings of the National Academy of Sciences*, 110(34), 13751-13756. doi: <https://doi.org/10.1073/pnas.1300503110>
- Farías, S. S., Casa, V. A., Vázquez, C., Ferpozzi, L., Pucci, G. N., & Cohen, I. M. (2003). Natural contamination with arsenic and other trace elements in ground waters of Argentine Pampean Plain. *Science of the total Environment*, 309(1), 187-199. doi: [https://doi.org/10.1016/S0048-9697\(03\)00056-1](https://doi.org/10.1016/S0048-9697(03)00056-1)
- Farooqi, A., Masuda, H., & Firdous, N. (2007). Toxic fluoride and arsenic contaminated groundwater in the Lahore and Kasur districts, Punjab, Pakistan and possible contaminant sources. *Environmental Pollution*, 145(3), 839-849. <https://doi.org/10.1016/j.envpol.2006.05.007>
- Fendorf, S., Michael, H. A., & van Geen, A. (2010). Spatial and temporal variations of groundwater arsenic in South and Southeast Asia. *Science*, 328(5982), 1123-1127. doi: DOI: 10.1126/science.1172974
- Fiorentino, C. E., Paoloni, J. D., Sequeira, M. E., & Arosteguy, P. (2007). The presence of vanadium in groundwater of southeastern extreme the pampean region Argentina: Relationship with other chemical elements. *Journal of contaminant hydrology*, 93(1-4), 122-129. doi: <https://doi.org/10.1016/j.jconhyd.2007.02.001>
- Frenguelli, J., & Teruggi, M. E. (1955). *Loess y limos pampeanos*: Olivieri y Dominguez.
- García, M., Borgnino, L., Bia, G., & Depetris, P. (2014). Mechanisms of arsenic and fluoride release from Chacopampean sediments (Argentina). *International Journal of Environment and Health*, 7(1), 41-57.
- García, M. G., Lecomte, K. L., Stupar, Y., Formica, S., Barrionuevo, M., Vesco, M., Gallará, R., & Ponce, R. (2012). Geochemistry and health aspects of F-rich mountainous streams and groundwaters from sierras Pampeanas de Cordoba, Argentina. *Environmental Earth Sciences*, 65(2), 535-545. doi: <https://doi.org/10.1007/s12665-011-1006-z>
- Garrels RM, Christ CL (1965) Solutions, minerals, and equilibrium. Harper & Row. Harper's geoscience series, New York
- Gérard, F., Clement, A., & Fritz, B. (1998). Numerical validation of a Eulerian hydrochemical code using a 1D multisolute mass transport system involving heterogeneous kinetically controlled reactions. *Journal of contaminant hydrology*, 30(3-4), 201-216. doi: [https://doi.org/10.1016/S0169-7722\(97\)00047-8](https://doi.org/10.1016/S0169-7722(97)00047-8)
- Glok Galli, M. G., Martínez, D. E., Kruse, E. E., Grondona, S. I., & Lima, M. L. (2014). Hydrochemical and isotopic characterization of the hydrological budget of a MAB Reserve: Mar Chiquita lagoon, province of Buenos Aires, Argentina. *Environmental earth sciences*, 72(8), 2821-2835. <https://doi.org/10.1007/s12665-014-3187-8>
- Gomez, M., Blarasin, M., & Martínez, D. (2009). Arsenic and fluoride in a loess aquifer in the central area of Argentina. *Environmental Geology*, 57(1), 143-155. doi: <https://doi.org/10.1007/s00254-008-1290-4>
- Guo, H., Liu, Z., Ding, S., Hao, C., Xiu, W., & Hou, W. (2015). Arsenate reduction and mobilization in the presence of indigenous aerobic bacteria obtained from high arsenic aquifers of the Hetao basin, Inner Mongolia. *Environmental pollution*, 203, 50-59. doi: <https://doi.org/10.1016/j.envpol.2015.03.034>
- Heredia, O. S., & Cirelli, A. F. (2009). Trace elements distribution in soil, pore water and groundwater in Buenos Aires, Argentina. *Geoderma*, 149(3-4), 409-414. doi: <https://doi.org/10.1016/j.geoderma.2008.12.020>
- Hopenhayn-Rich, C., Biggs, M. L., Fuchs, A., Bergoglio, R., Tello, E. E., Nicolli, H., & Smith, A. H. (1996). Bladder cancer mortality associated with arsenic in drinking water in Argentina. *Epidemiology*, 117-124.
- Jang, Y. C., Somanna, Y., & Kim, H. (2016). Source, distribution, toxicity and remediation of arsenic in the environment—a review. *Int. J. Appl. Environ. Sci*, 11(2), 559-581.

- Joseph, T., Dubey, B., & McBean, E. A. (2015). Human health risk assessment from arsenic exposures in Bangladesh. *Science of the total Environment*, 527, 552-560. doi: <https://doi.org/10.1016/j.scitotenv.2015.05.053>
- Kruse, E., & Ainchil, J. (2003). Fluoride variations in groundwater of an area in Buenos Aires Province, Argentina. *Environmental Geology*, 44(1), 86-89. doi: <https://doi.org/10.1007/s00254-002-0702-0>
- Kumar, M., Das, N., Goswami, R., Sarma, K. P., Bhattacharya, P., & Ramanathan, A. (2016). Coupling fractionation and batch desorption to understand arsenic and fluoride co-contamination in the aquifer system. *Chemosphere*, 164, 657-667. doi: <https://doi.org/10.1016/j.chemosphere.2016.08.075>
- Levy, D. B., Schramke, J. A., Esposito, K. J., Erickson, T. A., & Moore, J. C. (1999). The shallow ground water chemistry of arsenic, fluorine, and major elements: Eastern Owens Lake, California. *Applied Geochemistry*, 14(1), 53-65. [https://doi.org/10.1016/S0883-2927\(98\)00038-9](https://doi.org/10.1016/S0883-2927(98)00038-9)
- Logan, W. S., & Rudolph, D. L. (1997). Microdepression-focused recharge in a coastal wetland, La Plata, Argentina. *Journal of Hydrology*, 194(1), 221-238. doi: [https://doi.org/10.1016/S0022-1694\(96\)03205-2](https://doi.org/10.1016/S0022-1694(96)03205-2)
- Londoño, O. Q., Martínez, D. E., Dapeña, C., & Massone, H. (2008). Hydrogeochemistry and isotope analyses used to determine groundwater recharge and flow in low-gradient catchments of the province of Buenos Aires, Argentina. *Hydrogeology Journal*, 16(6), 1113-1127. <https://doi.org/10.1007/s10040-008-0289-y>
- Mahlknecht, J., Steinich, B., & De León, I. N. (2004). Groundwater chemistry and mass transfers in the Independence aquifer, central Mexico, by using multivariate statistics and mass-balance models. *Environmental Geology*, 45(6), 781-795. <https://doi.org/10.1007/s00254-003-0938-3>
- Martínez, D., & Bocanegra, E. (2002). Hydrogeochemistry and cation-exchange processes in the coastal aquifer of Mar Del Plata, Argentina. *Hydrogeology Journal*, 10(3), 393-408. doi: <https://doi.org/10.1007/s10040-002-0195-7>
- Martínez, D., Londoño, O. Q., Massone, H., Buitrago, P. P., & Lima, L. (2012). Hydrogeochemistry of fluoride in the Quequen river basin: natural pollutants distribution in the argentine pampa. *Environmental Earth Sciences*, 65(2), 411-420. doi: <https://doi.org/10.1007/s12665-011-0988-x>
- Nicolli, H., Bundschuh, J., Blanco, M., Tujchneider, O., Panarello, H., Dapeña, C., & Rusansky, J. (2012a). Arsenic and associated trace-elements in groundwater from the Chaco-Pampean plain, Argentina: results from 100 years of research. *Science of the total Environment*, 429, 36-56. doi: <https://doi.org/10.1016/j.scitotenv.2012.04.048>
- Nicolli, H., García, J., Falcón, C., & Smedley, P. (2012b). Mobilization of arsenic and other trace elements of health concern in groundwater from the Salí River Basin, Tucumán Province, Argentina. *Environmental geochemistry and health*, 34(2), 251-262. doi: <https://doi.org/10.1007/s10653-011-9429-8>
- Nicolli, H., Suriano, J., Gomez Peral, M., Ferpozzi, L., & Baleani, O. (1989). Groundwater contamination with arsenic and other trace elements in an area of the Pampa, Province of Córdoba, Argentina. *Environmental Geology and Water Sciences*, 14(1), 3-16. doi: <https://doi.org/10.1007/BF01740581>
- Nicolli, H. B., Bundschuh, J., García, J. W., Falcón, C. M., & Jean, J.-S. (2010). Sources and controls for the mobility of arsenic in oxidizing groundwaters from loess-type sediments in arid/semi-arid dry climates—Evidence from the Chaco—Pampean plain (Argentina). *Water research*, 44(19), 5589-5604. doi: <https://doi.org/10.1016/j.watres.2010.09.029>
- Padhi, S., & Tokunaga, T. (2015). Surface complexation modeling of fluoride sorption onto calcite. *Journal of environmental chemical engineering*, 3(3), 1892-1900. doi: <https://doi.org/10.1016/j.jece.2015.06.027>

- Palandri, J. L., & Kharaka, Y. K. (2004). A compilation of rate parameters of water-mineral interaction kinetics for application to geochemical modeling: GEOLOGICAL SURVEY MENLO PARK CA.
- Paquette, J., & Reeder, R. J. (1995). Relationship between surface structure, growth mechanism, and trace element incorporation in calcite. *Geochimica et Cosmochimica Acta*, 59(4), 735-749. doi: [https://doi.org/10.1016/0016-7037\(95\)00004-J](https://doi.org/10.1016/0016-7037(95)00004-J)
- Pokrovsky, O., & Schott, J. (2002). Surface chemistry and dissolution kinetics of divalent metal carbonates. *Environmental science & technology*, 36(3), 426-432. doi: 10.1021/es010925u
- Puntoriero, M. L., Cirelli, A. F., & Volpedo, A. V. (2015). Geochemical mechanisms controlling the chemical composition of groundwater and surface water in the southwest of the Pampean plain (Argentina). *Journal of Geochemical Exploration*, 150, 64-72. doi: <https://doi.org/10.1016/j.gexplo.2014.12.011>
- Pye, K. (1995). The nature, origin and accumulation of loess. *Quaternary Science Reviews*, 14(7), 653-667. doi: [https://doi.org/10.1016/0277-3791\(95\)00047-X](https://doi.org/10.1016/0277-3791(95)00047-X)
- Reeder, R. J. (1996). Interaction of divalent cobalt, zinc, cadmium, and barium with the calcite surface during layer growth. *Geochimica et Cosmochimica Acta*, 60(9), 1543-1552. doi: [https://doi.org/10.1016/0016-7037\(96\)00034-8](https://doi.org/10.1016/0016-7037(96)00034-8)
- Renard, F., Putnis, C. V., Montes-Hernandez, G., Ruiz-Agudo, E., Hovelmann, J., & Sarret, G. (2015). Interactions of arsenic with calcite surfaces revealed by in situ nanoscale imaging. *Geochimica et Cosmochimica Acta*, 159, 61-79. doi: <https://doi.org/10.1016/j.gca.2015.03.025>
- Robertson, F. N. (1989). Arsenic in ground-water under oxidizing conditions, south-west United States. *Environmental geochemistry and Health*, 11(3-4), 171-185. doi: <https://doi.org/10.1007/BF01758668>
- Romero, F. M., Armienta, M. A., & Carrillo-Chavez, A. (2004). Arsenic sorption by carbonate-rich aquifer material, a control on arsenic mobility at Zimapan, Mexico. *Archives of environmental contamination and toxicology*, 47(1), 1-13. doi: <https://doi.org/10.1007/s00244-004-3009-1>
- Roman-Ross, G., Charlet, L., Cuello, G., & Tisserand, D. (2003). *Arsenic removal by gypsum and calcite in lacustrine environments*. Paper presented at the Journal de Physique IV (Proceedings).
- Rouff, A. A., Elzinga, E. J., Reeder, R. J., & Fisher, N. S. (2004). X-ray Absorption Spectroscopic Evidence for the Formation of Pb (II) Inner-Sphere Adsorption Complexes and Precipitates at the Calcite– Water Interface. *Environmental science & technology*, 38(6), 1700-1707. doi: 10.1021/es0345625
- Sancha, A. M., & O’Ryan, R. (2008). Managing hazardous pollutants in Chile: arsenic *Reviews of Environmental Contamination and Toxicology Vol 196* (pp. 123-146): Springer.
- Sayago, J., Collantes, M., Karlson, A., & Sanabria, J. (2001). Genesis and distribution of the Late Pleistocene and Holocene loess of Argentina: a regional approximation. *Quaternary International*, 76, 247-257. doi: [https://doi.org/10.1016/S1040-6182\(00\)00107-5](https://doi.org/10.1016/S1040-6182(00)00107-5)
- Scanlon, B. R., Nicot, J., Reedy, R., Kurtzman, D., Mukherjee, A., & Nordstrom, D. K. (2009). Elevated naturally occurring arsenic in a semiarid oxidizing system, Southern High Plains aquifer, Texas, USA. *Applied Geochemistry*, 24(11), 2061-2071. doi: <https://doi.org/10.1016/j.apgeochem.2009.08.004>
- Schulz, C., & Castro, E. (2003). *Estudio, planificación y explotación del agua subterránea: una trilogía utópica en la República Argentina [Study, planning and exploitation of groundwater: a utopian trilogy in the Argentina Republic]*. Paper presented at the Proceedings of III Congreso Argentino de Hidrogeología, Rosario, Argentina, Edit. Universidad Nacional de Rosario.
- Smedley, P., Macdonald, D., Nicolli, H., Barros, A., Tullio, J., & Pearce, J. (2000). Arsenic and other quality problems in groundwater from northern La Pampa Province, Argentina.

- Smedley, P. L., & Kinniburgh, D. G. (2001). Source and behavior of arsenic in natural waters. *United Nations synthesis report on arsenic in drinking water*. World Health Organization, Geneva, Switzerland. http://www.who.int/water_sanitation_health/dwq/arsenicun1.pdf, 1-61.
- Smedley, P., Nicolli, H., Macdonald, D., Barros, A., & Tullio, J. (2002). Hydrogeochemistry of arsenic and other inorganic constituents in groundwaters from La Pampa, Argentina. *Applied Geochemistry*, 17(3), 259-284. doi: [https://doi.org/10.1016/S0883-2927\(01\)00082-8](https://doi.org/10.1016/S0883-2927(01)00082-8)
- Smedley, P. L., Kinniburgh, D. G., Macdonald, D. M. J., Nicolli, H. B., Barros, A. J., Tullio, J. O., ... & Alonso, M. S. (2005). Arsenic associations in sediments from the loess aquifer of La Pampa, Argentina. *Applied geochemistry*, 20(5), 989-1016.
- Smedley, P., Nicolli, H., Macdonald, D., & Kinniburgh, D. (2008). Arsenic in groundwater and sediments from La Pampa province, Argentina. *Bundschuh J.; Armienta MA; Birkle P.; Bhattacharya P*, 35-45.
- Sø, H. U., Postma, D., Jakobsen, R., & Larsen, F. (2008). Sorption and desorption of arsenate and arsenite on calcite. *Geochimica et Cosmochimica Acta*, 72(24), 5871-5884. doi: <https://doi.org/10.1016/j.gca.2008.09.023>
- Sota de la, M., Puche, R., Rigalli, A., Fernandez, L., Benassatti, S., & Boland, R. (1997). Modificaciones en la masa o sea en la homeostasis de la glucosa en residentes de la zona de Bahía Blanca con alta ingesta espontánea de Fluor. *MEDICINA*, 57:417-420.
- Steinhauser, G., & Bichler, M. (2008). Adsorption of ions onto high silica volcanic glass. *Applied Radiation and Isotopes*, 66(1), 1-8. doi: <https://doi.org/10.1016/j.apradiso.2007.07.010>
- Stipp, S., Lakshtanov, L., Jensen, J., & Baker, J. (2003). Eu³⁺ uptake by calcite: Preliminary results from coprecipitation experiments and observations with surface-sensitive techniques. *Journal of contaminant hydrology*, 61(1-4), 33-43. doi: [https://doi.org/10.1016/S0169-7722\(02\)00111-0](https://doi.org/10.1016/S0169-7722(02)00111-0)
- Stipp, S. L., Hochella Jr, M. F., Parks, G. A., & Leckie, J. O. (1992). CT²⁺ uptake by calcite, solid-state diffusion, and the formation of solid-solution: Interface processes observed with near-surface sensitive techniques (XPS, LEED, and AES). *Geochimica et Cosmochimica Acta*, 56(5), 1941-1954. doi: [https://doi.org/10.1016/0016-7037\(92\)90321-9](https://doi.org/10.1016/0016-7037(92)90321-9)
- Tesoriero, A. J., & Pankow, J. F. (1996). Solid solution partitioning of Sr²⁺, Ba²⁺, and CT²⁺ to calcite. *Geochimica et Cosmochimica Acta*, 60(6), 1053-1063. doi: [https://doi.org/10.1016/0016-7037\(95\)00449-1](https://doi.org/10.1016/0016-7037(95)00449-1)
- Tricart, J. (1973). Geomorfología de la Pampa Deprimida: base para los estudios edafológicos y agronómicos: Secretaría de Estado de Agricultura y Ganadería de la Nación, Instituto Nacional de Tecnología Agropecuaria.
- Turner, B. D., Binning, P., & Stipp, S. (2005). Fluoride removal by calcite: evidence for fluorite precipitation and surface adsorption. *Environmental science & technology*, 39(24), 9561-9568. doi: 10.1021/es0505090
- Vital, M., Martinez, D. E., Borrelli, N., & Quiroga, S. (2016). Kinetics of dissolution processes in loess-like sediments and carbonate concretions in the southeast of the province of Buenos Aires, Argentina. *Environmental Earth Sciences*, 75(17), 1231. doi: <https://doi.org/10.1007/s12665-016-6011-9>
- Vital, M., Daval, D., Clément, A., Quiroga, S., Fritz, B., & Martinez, D. E. (2018). Importance of accessory minerals for the control of water chemistry of the Pampean aquifer, province of Buenos Aires, Argentina. *CATENA*, 160, 112-123. doi: <https://doi.org/10.1016/j.catena.2017.09.005>
- Wang, Z., Guo, H., Xiu, W., Wang, J., & Shen, M. (2018). High arsenic groundwater in the Guide basin, northwestern China: Distribution and genesis mechanisms. *Science of the total Environment*, 640, 194-206. doi: <https://doi.org/10.1016/j.scitotenv.2018.05.255>
- Wurl, J., Mendez-Rodriguez, L., & Acosta-Vargas, B. (2014). Arsenic content in groundwater from the southern part of the San Antonio-El Triunfo mining district, Baja California

- Sur, Mexico. *Journal of Hydrology*, 518, 447-459. doi:
<https://doi.org/10.1016/j.jhydrol.2014.05.009>
- Wyatt, C. J., Quiroga, V. L., Acosta, R. T. O., & Méndez, R. O. (1998). Excretion of arsenic (As) in urine of children, 7–11 years, exposed to elevated levels of As in the city water supply in Hermosillo, Sonora, Mexico. *Environmental research*, 78(1), 19-24. <https://doi.org/10.1006/enrs.1998.3844>
- Yang, K., Lu, H., Yue, S., Zhang, G., Lei, Y., La, Z., & Wang, W. (2018). Quantifying recent precipitation change and predicting lake expansion in the Inner Tibetan Plateau. *Climatic Change*, 147(1-2), 149-163. doi: <https://doi.org/10.1007/s10584-017-2127-5>
- Zabala, M., Manzano, M., & Vives, L. (2015). The origin of groundwater composition in the Pampeano Aquifer underlying the Del Azul Creek basin, Argentina. *Science of the total Environment*, 518, 168-188. doi: <https://doi.org/10.1016/j.scitotenv.2015.02.065>
- Zachara, J., Cowan, C., & Resch, C. (1991). Sorption of divalent metals on calcite. *Geochimica et Cosmochimica Acta*, 55(6), 1549-1562. doi: [https://doi.org/10.1016/0016-7037\(91\)90127-Q](https://doi.org/10.1016/0016-7037(91)90127-Q)
- Zárate, M. A. (2003). Loess of southern South America. *Quaternary Science Reviews*, 22(18), 1987-2006. doi: [https://doi.org/10.1016/S0277-3791\(03\)00165-3](https://doi.org/10.1016/S0277-3791(03)00165-3)

Table 1. Highest concentrations of major elements in the first test (T1) and second test (T2) of the flow-through and final concentration of the first test (TF1) conducted with the samples of loess and calcrete collected in the area of Mar del Plata.

	LOESS			CALCRETE		
	T1	TF1	T2	T1	TF1	T2
Ca ²⁺	4.8	0.06	0.16	0.64	0,2	0,37
Na ⁺	3.3	0.2	0.09	3.2	0,5	0,58
SiO ₂	0.24	0.04	0.21	0,29	0,20	0,16
SO ₄ ⁻²	0.7	0.02	0.01	0,40	n/d	0,05
K ⁺	0.14	0.05	0.015	0.1	0.03	0.08
Cl ⁻	21	0.01	0.1	7,99	0,1	0,1

* mmol.l⁻¹

Table 2. Arsenic content of solutions in contact with calcrete in flow-through experiments with tests T1 and T2 from 3 locations of the Buenos Aires Province.

	T1				T2			
	t1	t5	t30	t60	t1	t5	t30	t60
Otamendi	det <0.005	det <0.005	no det <0.002	no det <0.002	det <0.005	det <0.005	det <0.005	det <0.005
Tandil	det <0.005	det <0.005	det <0.005	no det <0.002	det <0.005	det <0.005	no det <0.002	no det <0.002
Tres arroyos	det <0.005	det <0.005	det <0.005	no det <0.002	det <0.005	det <0.005	no det <0.002	no det <0.002

T1: Test 1 - T2: Test 2
t: time in minutes
Det: detected - No det: not detected
* $\mu\text{mol.l}^{-1}$

Figure 1. Location of the sampling points (indicate by red dots on the map)

Figure 2. Schematic diagrams of the flow-through experiment and reactor. The deionized water (pH 5.7) is continuously pumped through the reactor at a flow rate of 5 ml.min⁻¹ where the loess or calcrete samples were previously introduced.

Figure 3. Release of trace elements in the flow-through experiment test 1 (T1) and test 2 (T2) conducted with the loess and calcrete samples from the area of Mar del Plata.

Figure 4. Correlation between Ca and trace elements (Ba, Sr, V, As) in the flow-through experiment test T1 conducted with loess.

Figure 5. Correlation between Na^+ , K^+ and trace elements (Ba^{2+} , Sr^{2+}) in the flow through experiment
Test T1 conducted with loess.

Figure 6. Correlation between Ca^{2+} and F^- release in T1 and T2 flow-through experiments conducted with loess.

Figure7
Click here to download Figure: figure7.docx

Figure7. Correlations between Ca^{2+} and Ba^{2+} , Sr^{2+} , As, V in the flow through experiments T1 (a) and between Ca^{2+} and F^- (b) in the test T1 and T2 conducted on calcrete samples.

Figure 8. Simulations of fluoride release for loess (a) and calcrete (b) experiments including accessory minerals. The kirmat simulation for loess and calcrete were first realized with 2% of fluorapatite. Then, another kirmat simulation for loess was realized with 2% Calcite containing 0.1% impurity of fluoride and 2% of fluorapatite. Another kirmat simulation for calcrete was realized with 95% Calcite containing 0.1% impurity of fluoride and 2% of fluorapatite. The Kirmat simulations were compared to the flow-through experiments T1 and T2 on the loess sample from Mar del Plata (MdP) and the calcrete samples from Mar del Plata (MdP), Otamendi, Tandil and Tres Arroyos (TresA), in the province of Buenos Aires.