



HAL
open science

Set-membership computation of integrals with uncertain endpoints

Olivier Mullier, Julien Alexandre Dit Sandretto

► **To cite this version:**

Olivier Mullier, Julien Alexandre Dit Sandretto. Set-membership computation of integrals with uncertain endpoints. NUMTA 2019, Jun 2019, Le Castella village, Italy. hal-02372387

HAL Id: hal-02372387

<https://hal.science/hal-02372387>

Submitted on 20 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Set-membership computation of integrals with uncertain endpoints

Olivier Mullier, Julien Alexandre dit Sandretto

ENSTA ParisTech, 828 bd des maréchaux 91120 Palaiseau, France
mullier@ensta.fr, alexandre@ensta.fr

Keywords. Integral; Set-membership Computation; Interval Methods.

Numerical integration is one of the fundamental tool of scientific computation. Providing a reliable result to such problem is important for validated simulation [1] or for global optimization with continuous objective function [2]. An important work on inclusion methods for integral equations can be found in [3]. We propose an efficient **guaranteed** method for the computation of the **integral** of a nonlinear continuous function f between two **interval endpoints** $[x_1]$ and $[x_2]$, define by:

$$\int_{[x_1]}^{[x_2]} f(x)dx = \left\{ \int_{x_1}^{x_2} f(x)dx : x_1 \in [x_1], x_2 \in [x_2] \right\} \quad (1)$$

Two cases can occur whether the intervals $[x_1]$ and $[x_2]$ intersect or not. If the two intervals do not intersect, we produce an outer approximation of the set of possible values for the integral proving the following proposed theorem:

Theorem 1. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function. For $[x_1] = [x_1, \bar{x}_1]$ and $[x_2] = [x_2, \bar{x}_2]$ two intervals in \mathbb{IR} . Then the minimum and the maximum of set (1) can be defined by*

$$\int_{[x_1]}^{[x_2]} f(x)dx = \left[\min \int_{\mathcal{X}_{1-}^* \cup \{\bar{x}_1\}}^{\mathcal{X}_{2+}^* \cup \{x_2\}} f(x)dx, \max \int_{\mathcal{X}_{1+}^* \cup \{\bar{x}_1\}}^{\mathcal{X}_{2-}^* \cup \{x_2\}} f(x)dx \right] \quad (2)$$

using

$\mathcal{X}_{i-}^* = \{x \in [x_i] : f(x) = 0, f'(x) < 0\}$, $\mathcal{X}_{i+}^* = \{x \in [x_i] : f(x) = 0, f'(x) > 0\}$.

When the two intervals intersect, Second Fundamental Theorem of Calculus is applied. An implementation of the computation of the bounds (2) is introduced using interval analysis [4] and tested on several examples.

References

- [1] Alexandre dit Sandretto, J. and Chapoutot, A. (2016) *Validated Explicit and Implicit Runge-Kutta Methods*. Reliable Computing.
- [2] Hansen, Eldon R. (2003) *Global Optimization Using Interval Analysis*. CRC Press.
- [3] Corliss, G. (1987) *Computing narrow inclusions for definite integrals*. Computer-arithmetic (eds. E. Kaucher et al.), pp. 150—179. Wiley-Teubner, Stuttgart.
- [4] Moore, Ramon E. (1966) *Interval analysis*. Prentice-Hall Englewood Cliffs, NJ.