Vocal expression of emotional valence in pigs across multiple call types and contexts

Elodie Briefer, Pavel Linhart, Richard Policht, Lisette M C Leliveld, Sandra Döpjan, Birger Puppe, Mónica Padilla de La Torre, Andrew M Janczak, Cécile Bourguet, Véronique Deiss, et al.

► To cite this version:

Elodie Briefer, Pavel Linhart, Richard Policht, Lisette M C Leliveld, Sandra Döpjan, et al.. Vocal expression of emotional valence in pigs across multiple call types and contexts. 2019. hal-02375173

HAL Id: hal-02375173

<https://hal.science/hal-02375173>

Preprint submitted on 22 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Vocal expression of emotional valence in pigs across multiple call types and contexts

Elodie F. Briefer¹, Pavel Linhart², Richard Policht², Marek Špinka², Lisette M. C. Leliveld³, Sandra Döpjan³, Birger Puppe³, Mónica Padilla de la Torre⁴, Andrew M. Janczak⁴, Cécile Bourguet⁵, Véronique Deiss⁶, Alain Boissy⁶, Carole Guérin⁷, Eva Read⁷, Marjorie Coulon⁸, Edna Hillmann⁹, Céline Tallet⁷

¹ Institute of Agricultural Sciences, ETH Zürich, Switzerland

² Institute of Animal Science, Czechia

³ Leibniz Institute for Farm Animal Biology, Germany

⁴ Department of Production Animal Clinical Sciences, Norwegian University of Life Sciences, Norway

⁵ Bureau E.T.R.E., France

⁶ INRA UMR 1213 Herbivores, France

⁷ INRA UMR 1348 PEGASE, France

⁸ Cabinet EASIER, France

⁹ Humboldt-Universität zu Berlin, Germany

Corresponding Author:

Elodie F Briefer¹

Present address: Behavioural Ecology Group, Section for Ecology and Evolution, Department of Biology, University of Copenhagen, Denmark

Email address: elodie.briefer@bio.ku.dk

Introduction

Emotions, unlike mood, are short-lived reactions associated with specific events. They can be characterized by two main dimensions; their arousal (bodily activation) and valence (negative versus positive) (Mendl et al. 2010). Knowledge of the valence of emotions experienced by domestic and captive animals is crucial for assessing and improving their welfare, as it enables us to minimize the negative emotions that they might experience and to promote positive ones. Emotions can affect vocalizations directly or indirectly through the brain, lungs, larynx or vocal tract. As a result, vocal expression of emotions has been observed across species (Briefer 2012), and could serve as a non-invasive and potentially very reliable tool to assess animal emotions. In pigs (*Sus scrofa*), vocal expression of emotions has been relatively well studied (e.g. Leliveld et al. 2016; Briefer et al. 2019). However, it is not known if the vocal indicators revealed in previous studies are valid across call types and contexts. To find this out, we conducted an analysis of the effect of emotional valence on a large database of pig vocalizations, including calls recorded in the most common emotional situations encountered by pigs throughout their lives, from birth to slaughter.

Materials & Methods

Recordings

Pigs of various ages (piglets to finishing pigs) were recorded in 22 contexts triggering both negative emotions (e.g. crushing, missed nursing, castration, fear conditioning, isolation, restraint, barren environment, and slaughter), and positive emotions (e.g. nursing, huddling, social reunion, exposition to an enriched arena, and running) (for more details see Tallet et al. 2013; Linhart et al. 2015; Leliveld et al. 2016; Leliveld et al. 2017; Briefer et al. 2019). The putative valence of the various contexts was based on the function of emotions to trigger avoidance (negative emotions) or approach (positive emotions) and the behavior of the pigs (Mendl et al. 2010). Experiments included in this analysis were approved by the respective authorities for each country (Germany: Federal State of Mecklenburg-Western Pomerania (AZ:7221.3-2-045/13); Switzerland: Swiss Cantonal Veterinary Office (TG02/2014); Czechia: Institutional Animal Care and Use Committee of the Institute of Animal Science and the Czech Central Committee for Protection of Animals, Ministry of Agriculture (dMZe 1244 and 44248/2007–17210); Norway: National animal research authority (FOTS id 12021)).

Vocal analyses

In order to exclude very short sounds, in which parameters might not be accurately measured, only high quality calls with a duration > 0.05 s were selected for the acoustic analysis ($n = 7392$ calls). We used the acoustic features of the calls to classify them as low-frequency stable, modulated or tonal calls, high-frequency stable or modulated calls, or mixed calls (6 types), based on Tallet et al. (2013). Then, depending on the call type, we extracted 11 to 18 vocal parameters using a custom-built script in Praat, which batch-processed the analyses and the exporting of output data. The measured parameters belonged to the six following categories: source-related (fundamental frequency, "F0"), energy spectrum distribution, duration, amplitude modulation ("AM"), noise, filter-related (vocal tract resonances).

Statistical analyses

To eliminate redundancy, we used a principal component analysis to select one vocal parameter within each category, which explained most of the variance in the data across all call types, for further analyses. Since the minimum formant dispersion ("DFmin"), originally categorized along with the linear predictive coding ("LPC") coefficients never associated (i.e. loaded highly ($r \geq 0.5$) on the same PC) with these parameters, it was analyzed separately. These selected seven parameters (i.e. one for each of the six categories and DFmin; Table 1) were then used as outcome variables in linear mixed-effects models (lmer function in R software), to assess if they were affected by the valence of the contexts (positive or negative; fixed factor). The models included as control factors the age category and the call type. The context of production nested within the identity of the pig, nested within the experiment number, nested within the team who performed the recording was added as a random factor to control for repeated measurements and dependencies. The p-values were calculated with parametric bootstrap tests.

Results and Discussion

Five of the seven tested vocal parameters were affected by the valence of the context (Table 1). After controlling for the type of call and the age category (control factors), our analyses revealed that pigs

produced calls characterized by a higher center of gravity, a shorter duration, less noise (lower Wiener entropy), lower formants (measured using the formant dispersion) and LPC coefficients in positive compared to negative contexts.

Table 1. Model estimates, lower (lo.ci) and upper (up.ci) 95% confidence intervals for the vocal parameters included in the linear mixed-effect models, as a function of the valence of the contexts (* $p < 0.05$; ** $p < 0.01$; "NS" Non significant).

Parameter	Valence	estim	lo.ci	up.ci	P value
Mean F0 (Hz)	Pos	132.91	124.81	141.13	NS
	Neg	138.19	130.49	146.26	
Spectral centre of gravity (Hz)	Pos	967.70	877.57	1084.53	*
	Neg	895.54	806.46	996.58	
Duration (s)	Pos	0.17	0.14	0.21	**
	Neg	0.42	0.34	0.51	
AM extent (dB)	Pos	5.77	4.60	7.30	NS
	Neg	5.67	4.50	7.24	
Wiener entropy	Pos	-1.63	-1.84	-1.44	**
	Neg	-1.52	-1.71	-1.33	
DFmin (Hz)	Pos	846.64	778.01	921.82	**
	Neg	964.72	899.40	1035.78	
4th LPC coefficient (Hz)	Pos	3913.22	3742.58	4069.90	**
	Neg	4185.93	4020.41	4334.90	

Some of these changes are in line with previous findings (e.g. spectral center of gravity, Leliveld et al. 2016; duration, Briefer et al. 2019). In particular, shorter durations in positive contexts have been observed across multiple species and could be a feature conserved throughout evolution (Briefer 2012). Overall, our results suggest that some parameters change with the valence experienced by pigs in a similar way across call types. These vocal parameters could be very useful for developing automated methods to monitor pig welfare on-farm.

References

- Briefer, E. F. (2012). Vocal expression of emotions in mammals: mechanisms of production and evidence. *Journal of Zoology*, 288:1–20
- Briefer, E. F., Vizier, E., Gygax, L., Hillmann, E. (2019). Expression of emotional valence in pig closed-mouth grunts: Involvement of both source- and filter-related parameters. *Journal of the Acoustical Society of America*, 145:2895–2908.
- Leliveld, L. M. C., Düpjan, S., Tuchscherer, A., Puppe, B. (2016). Behavioural and physiological measures indicate subtle variations in the emotional valence of young pigs. *Physiology & Behavior*, 157:116–124.
- Leliveld, L. M. C., Düpjan, S., Tuchscherer, A., Puppe, B. (2017). Vocal correlates of emotional reactivity within and across contexts in domestic pigs (*Sus scrofa*). *Physiology & Behavior*, 181:117–126.
- Linhart, P., Ratcliffe, V. F., Reby, D., Špinká, M. (2015). Expression of emotional arousal in two different piglet call types. *PLoS ONE*, 10:e0135414.
- Mendl, M., Burman, O. H. P., Paul, E. S. (2010). An integrative and functional framework for the study of animal emotion and mood. *Proceedings of the Royal Society B*, 277:2895–2904.
- Tallet, C., Linhart, P., Policht, R., et al (2013). Encoding of situations in the vocal repertoire of piglets (*Sus scrofa*): A comparison of discrete and graded classifications. *PLoS ONE*, 8:e71841.