Ecological assessment of groundwater ecosystems disturbed by recharge systems using organic matter quality, biofilm characteristics and bacterial diversity

Jérémy Voisin, B. Cournoyer, L. Marjolet, Antonin Vienney, Florian Mermillod-Blondin

► To cite this version:

Jérémy Voisin, B. Cournoyer, L. Marjolet, Antonin Vienney, Florian Mermillod-Blondin. Ecological assessment of groundwater ecosystems disturbed by recharge systems using organic matter quality, biofilm characteristics and bacterial diversity. *Environmental Science and Pollution Research*, 2020, 27 (3), pp.3295-3308. 10.1007/s11356-019-06971-5 . hal-02386310

HAL Id: hal-02386310

<https://hal.science/hal-02386310>

Submitted on 29 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Ecological assessment of groundwater ecosystems disturbed by recharge systems using organic matter quality, biofilm characteristics and bacterial diversity

Jérémy VOISIN^{1,2}, Benoit COURNOYER², Laurence MARJOLET², Antonin VIENNEY¹,
Florian MERMILLOD-BLONDIN^{1*}

1- Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR5023, Écologie des
Hydrosystèmes Naturels et Anthropisés, Villeurbanne, F-69622, France

2- Univ Lyon, Université Claude Bernard Lyon 1, VetAgro Sup, CNRS, UMR5557, INRA
UMR1418, Ecologie Microbienne, Research Team "Bacterial Opportunistic Pathogens and
Environment", Marcy L'Etoile, F-69280, France

* Corresponding author:

Florian Mermillod-Blondin

Phone: 00 33 4 72 43 13 64

E-mail: florian.mermillod-blondin@univ-lyon1.fr

19 Abstract

20 Recharge of aquifers by urban stormwater may trigger significant ecological changes that can be
21 detrimental to the biodiversity and functioning of groundwater ecosystems. Here, the effects of
22 aquifer recharge (AR) on three levels of parameters were investigated: dissolved organic carbon
23 (DOC) quantity and quality, global biofilm characteristics, and diversity changes of bacterial
24 communities. As DOC enrichment by AR can be mitigated by vadose zone (VZ) thickness, 3 AR sites
25 with thin VZ (< 3 m) and 3 sites with thick VZ (> 10 m) were selected. For each AR site, clay beads
26 were incubated over a 10 days-long rainy period through wells in recharged and non-recharged
27 groundwaters. Total proteins, dehydrogenase and hydrolytic activities were monitored from clay
28 beads to assess biofilm development. Bacterial richness on beads was estimated by 16S rRNA-based
29 metabarcoding. AR was found to significantly increase DOC and biodegradable DOC (BDOC)
30 concentrations, biofilm development and bacterial richness especially in sites with thin VZ. VZ
31 thickness was inversely related to microbial growth indicators and bacterial richness in groundwater,
32 through a control of DOC availability. The proportion of *Bacteroidetes* 16S rRNA gene reads was
33 higher in recharged groundwater than in non-recharged groundwater, suggesting that this phylum
34 could be used as an indicator of DOC enrichment associated with AR. Quantitative PCR assays for
35 *Bacteroides* DNA confirmed these trends, and showed an enrichment of this bacterial group in DOC-
36 rich aquifer waters. The positive linear relationships between BDOC concentrations and biofilm
37 variables highlighted a strong C-limitation of groundwater impacting bacterial species sorting and
38 activity.

40 Keywords

41 Nutrient-driven diversity changes; vadose zone thickness; stormwater infiltration; clay bead biofilms
42 bacterial communities; aquifer 16S rRNA metabarcoding profiling

44 Introduction

45 In ecology, the species-energy theory (Wright 1983, Evans et al. 2005; Fuhrman 2009) is considered
46 as a universal law stating that species richness correlates positively with energy availability. Smith
47 (2007) reviewed 70 published studies in aquatic ecosystems and concluded that diversity-productivity
48 patterns are similar between micro- and macro-organisms. Nonetheless, the shape of the diversity-
49 productivity relationship was variable depending on the studied micro-organisms and environments
50 (Smith 2007; Prosser et al. 2007; Bernstein et al. 2017). Although these relationships have been
51 reported from many aquatic ecosystems including natural lakes, ponds, and oceans, few studies have
52 investigated the influence of energy availability on the diversity of microbial communities in
53 groundwater ecosystems (but see Hershey et al. 2018 for a deep karstic aquifer).

54 Groundwater ecosystems are among the most stable and energy-limited environments on Earth
55 because of the lack of photosynthesis and low inputs of organic matter infiltrating from surface
56 ecosystems (Gibert 1994; Griebler and Lueders, 2009). Nonetheless, connectivity and dissolved
57 organic matter (DOM) transfers between surfaces and subterranean systems can be enhanced by
58 aquifer recharge (AR) practices (e.g., Kolehmainen et al. 2007). In urban environments, these
59 practices were developed to collect stormwater runoff to recharge aquifers by infiltration through
60 porous media (Marsalek and Chocat, 2002; Hunt et al. 2010). AR practices have been shown to
61 increase DOM in groundwaters over the levels of those that were not influenced by recharge (Datry
62 et al. 2004; Foulquier et al. 2010; Mermillod-Blondin et al. 2013). DOM is the main source of food
63 for micro-organisms in groundwater ecosystems (Goldscheider et al. 2006), and its increase can lead
64 to denser and more active microbial communities (Foulquier et al. 2011, Landa et al. 2013). For
65 example, Mermillod-Blondin et al. (2013) showed that the growth, and the esterase and
66 dehydrogenase activities of biofilms were stimulated by DOM enrichment associated with aquifer
67 recharge in an urban context. Nevertheless, the influence of DOM supplied by AR systems on
68 microbial diversity is still poorly defined.

69 The aim of the present work was to evaluate the relative efficiencies of three levels of descriptors to

detect the effect of stormwater inputs on groundwater: DOM quality, biofilm characteristics and bacterial diversity. A field experiment was conducted during a rainy period in 6 AR sites that were divided into two groups: (1) those with a thin vadose zone (< 3 m) and (2) those with a thick vadose zone (> 10 m). The vadose zone plays a key role on DOM retention during its transfer from surface to groundwater (Pabich et al. 2001; Shen et al. 2015). Selection of sites with different vadose zone thicknesses should lead to significant differences in groundwater disturbance. Dissolved organic carbon (DOC) and biodegradable (BDOC) concentrations were measured together with the incubation of artificial substrates (i.e., clay beads) in recharged and non-recharged groundwaters. Clay beads were previously shown to be efficient artificial substrates that can allow reliable bacterial diversity analyses of groundwaters (Voisin et al. 2016; Mermillod-Blondin et al. 2019). Microbial productivity and activities of the biofilms on the clay beads could be monitored from biomass measurements (total proteins), and from esterase and dehydrogenase activities. Bacterial diversity was investigated through a next-generation sequencing (NGS) dataset of PCR amplified 16S rRNA gene segments generated from biofilm DNA extracts. According to Datry et al. (2005), we expected that DOM concentrations would be higher in AR sites with a thin vadose zone in comparison with those with a thick vadose zone. Consequently, the hypothesis was that the influence of AR practices on microbial biomass, activity and diversity should be highest in AR sites with the thinnest vadose zone (< 3 m). In addition, positive correlations between DOM concentrations over the 6 sites and bacterial richness of the clay bead biofilms developing in groundwaters should be observed.

Material and methods

Sites

Aquifer recharge (AR) sites were located on the eastern aquifer of the Lyon metropolitan area, France (Fig. 1A). This aquifer (catchment area: 314 km²) is characterized by three corridors of highly permeable glacio-fluvial sediments (hydraulic conductivity 10⁻³-10⁻² m/s) (Foulquier et al. 2009). The aquifer is artificially recharged with stormwater runoff at multiple AR sites to compensate for the

reduced recharge caused by the sealing of urban surfaces. The AR sites of this study corresponded to retention/infiltration basins that collected stormwater from residential, commercial, and industrial areas.

Experimental design

Six AR sites (Figure 1AB) were selected: 3 with vadose zone thicknesses < 3 meters (IUT, Grange Blanche (GB) and Minerve (MIN) basins) and 3 with vadose zone thicknesses > 10 meters (Django Reinhardt (DJR), Raquin (RAQ) and Feyzin (FEY) basins). Each AR site is equipped with two wells: a well located outside the area of influence of the stormwater plume induced by infiltration which was used as a control well, and a well located in the immediate downstream vicinity of the infiltration basin used as a recharge well because it intersected the stormwater plume induced by the infiltration.. Clay beads (expanded clay aggregates, Botanic®, Saint-Julien en Genevois, France) of a diameter of 8 mm were incubated in these wells to trap micro-organisms. These clay beads favor biofilm developments from the groundwater microbiota (Voisin et al. 2016; Mermillod-Blondin et al. 2019), and enable a monitoring of parameters such as microbial productivity (total proteins), and esterase and dehydrogenase activities. Before incubation, clay beads were heated to 550°C to eliminate organic matter traces and then autoclaved at 121°C for 20 min. They were incubated in wells using small polypropylene bags (110 × 40 mm) with a mesh size of 3.4 mm, previously treated with ethanol (70%). A total of 30 clay beads were inserted per bag, and 3 bags were incubated per well after well purging with a submerged pump (see below). From each bag, clay beads were used to measure all microbial variables (total proteins, esterase and dehydrogenase activities, bacterial community structure, and qPCR assays on specific genes) in order to obtain mean biofilm parameters per well calculated from 3 values (corresponding to batches of clay beads from the three bags incubated in the same well). According to previous laboratory and field experiments (Voisin et al. 2016; Mermillod-Blondin et al. 2019), clay bead incubation lasted 10 days during a rainy period to obtain (1) a significant microbial colonization of clay beads submerged in groundwaters and (2) a time-integrated monitoring

of micro-organisms in recharged groundwater during a rainy event. The experiment was performed between September 12 and September 22 in 2015. During this period, precipitations were of 26.6 mm on the 12, 34.6 mm on the 13, 1.6 mm on the 14, 49.6 mm on the 17, 2.8 mm on the 18, 3.2 mm on the 19, and 1 mm on the 20. For each AR site, groundwaters (recharged and non-recharged) were also collected from the same wells but at three dates over the clay bead incubation period (Day 0, Day 5 and Day 10) to obtain mean values of DOC and BDOC per well during the incubation period of clay beads. Groundwaters were sampled in wells using a submerged pump (PP36 inox, SDEC, Reignac-sur-Indre, France). The first 50 pumped liters were used to rinse the sampling equipment. The following liter was collected for analyses of DOC and BDOC analyses. Samples were stored in pre-combusted glass bottles and brought back to the laboratory in an isotherm box at 4°C.

Water physico-chemistry

DOC (dissolved organic carbon) was determined using a carbon analyzer (multi N/C® 3100, Analytik Jena, Jena, Germany) based on thermocatalytic oxidation (850°C) of organic carbon and infrared detection of CO₂, after removal of dissolved inorganic C with HCl. Biodegradable DOC (BDOC) was determined by the method of Servais et al. (1987, 1989) following Mermillod-Blondin et al. (2015). Briefly, water samples (100 mL) were filtered through a pre-washed 0.2 µm polycarbonate membrane and incubated with a bacterial inoculum at 20°C for 30 days in the dark. DOC concentrations were measured from filtered (0.2 µm) water samples at the start and the end of the incubation period to determine the initial DOC concentration and the remaining DOC concentration after 30 days of incubation (representing the refractory dissolved organic carbon, RDOC), respectively. BDOC concentration was then calculated as the difference between initial DOC concentration and RDOC concentration.

Microbial analysis

Microbial biomass

148 Microbial biomasses on clay beads were assessed from Lowry's total protein assay using the Sigma
 149 Protein Assay Kit P-5656 (Sigma Diagnostics, St Louis, MO, USA) according to Peterson (1977).
 150 For each incubated bag, 5 randomly-sampled clay beads were treated with 4 mL of a 50 % solution
 151 of Lowry's reagent for 20 min. At the start of the reaction, the contact between proteins on beads and
 152 the reagent was improved by vortexing at 1000 rpm for 1 minute. After the 20-min long reaction, 0.5
 153 mL of Folin reagent was added, followed by a 30 min color reaction. The absorbance of the solution
 154 was then measured at 750 nm with a spectrophotometer (Aquamate, ThermoSpectronic, Cambridge,
 155 UK) and total protein content on beads was determined using a calibration curve performed with
 156 bovine serum albumin. Total proteins were expressed in μg of proteins per cm^2 of clay bead, after
 157 subtracting values measured from 3 batches of 5 sterile clay beads.

159 Microbial activity

160 Hydrolytic activity of biofilms was estimated using the fluorescein diacetate (FDA) hydrolysis
 161 method (Fontvieille et al. 1992) following Voisin et al. (2016). For each incubated bag, 5 randomly-
 162 sampled clay beads were incubated into 3 ml of a pH 7.6 phosphate buffer solution with 0.15 ml of
 163 4.8 mM FDA solution. Incubation was performed at 15°C for 3 h. The reaction was stopped by
 164 addition of 3 ml of acetone in the sample and storage at -20°C (Battin 1997). Fluorescein
 165 concentration was estimated by measuring the absorbance at 490 nm of the filtered supernatant (0.45
 166 μm , HAWP, Millipore, Billerica, MA, U.S.A.). Then, the hydrolytic activity of biofilms was
 167 expressed in μmol of FDA hydrolyzed per hour and per cm^2 of clay bead, after subtracting values
 168 measured from 3 batches of 5 sterile clay beads.

169 The 2-(p-iodophenyl)-3-(p-nitrophenyl)-5-phenyl tetrazolium chloride (INT) was used to measure
 170 dehydrogenase activity (respiratory activity) as modified from Houri-Davignon et al. (1989)
 171 following Mermillod-Blondin et al. (2019). For each incubated bag, 5 randomly-sampled clay beads
 172 were incubated in a 0.02% INT solution for 3 h at 15°C and the supernatant was then filtered on a
 173 nylon membrane (0.22 μm , Millipore, Billerica, MA, U.S.A.) to retrieve floating microbial cells.

174 Extraction of INT-formazan from clay beads and nylon membrane containing microbial cells was
175 performed with 5 ml of methanol during 12 h. The INT-formazan concentration was determined by
176 absorbance measurement at 480 nm. Then, the dehydrogenase activity of biofilms was expressed in
177 μmol of INT-formazan produced per hour and per cm^2 of clay bead, after subtracting values measured
178 from 3 batches of 5 sterile clay beads.

16S rRNA gene bacterial community profiling

DNA extraction and qPCR assays

182 DNA extraction from clay beads biofilms was performed following Voisin et al. (2016). For each
183 incubated bag, microbial cells were detached from 15 randomly-selected clay beads by shaking at
184 2500 rpm for 2 minutes in 10 mL of 0.8 % NaCl. Cells and aggregates were retrieved from the NaCl
185 solution by a 0.22 μm filtration. DNA was extracted from filters using the FastDNA spin kit for soil
186 (MP biomedical). Then, the amount of extracted DNA was quantified spectrophotometrically using
187 a Nanodrop ND1000 (Thermo Fisher Scientific, USA).

188 qPCR assays were performed on a Bio-Rad CFX96 real-time PCR instrument with Bio-Rad CFX
189 Manager software, version 3.0 (Marnes-la-Coquette, France). Total *Bacteroides* DNA targets were
190 PCR amplified according to Layton et al. (2006) using the Brilliant II SYBR Green low ROX qPCR
191 master mix for TaqMan qPCR (Agilent, Vénissieux, France). The human-specific HF183 *Bacteroides*
192 qPCR assay was performed according to Seurinck et al. (2005), and the assay for the 16S rRNA gene
193 segment for total bacteria was performed according to Park and Crowley (2006) using primers 338F
194 and 518R, and using the Brilliant II SYBR green low ROX qPCR master mix for SYBR Green qPCR.
195 Melting T° was 60°C for all assays. Linearized plasmid DNA containing 16S rRNA genes from the
196 targeted DNAs were run as standards using 10-fold dilutions of the plasmids. These plasmids were
197 obtained from Marti et al. (2017). Presence of inhibitors was checked by spiking known amount of
198 plasmid harboring *int2* with 10 times dilution of sample DNA extracts (10^7 copies of plasmid per μL
199 of DNA diluted solution). Number of cycles needed to have a significant signal was compared with

wells where only plasmid harboring *int2* was added to the qPCR mix. When a higher number of cycles was needed to observe a signal, a dilution by 5 or 10 fold was done and another run of tests was performed to confirm the absence of PCR inhibitions. Negative controls without template DNA were run in triplicate. Each assay was triplicated on distinct DNA extracts, and technical triplicates were performed.

V5-V6 analyses of 16S rRNA gene PCR products

Sequencing of the V5-V6 16S rRNA gene segments were performed at MrDNA (Shallowater, TX, USA) with the Illumina MiSeq technology. Primers 799F (5'-ACCMGGATTAGATACCKG-3') and 1193R (5'-CRTCCMCACCTTCCTC-3') were used, and 20,000 reads (i.e. 2 x 300-bp paired-end reads) were generated per sample. Data processing was realized with the MOTHUR software version 1.35.1 (Schloss et al. (2009) following Voisin et al. (2018)). Briefly, raw sequences were filtered by discarding those having (1) read sizes < 200 bp and > 400 pb, (2) reads containing homopolymers > 8 bp, (3) more than one sequence mismatch to barcode DNA tags and more than two mismatches with the forward sequencing primer, (4) reads containing ambiguous bases, and (5) average sequencing quality score < 35 using a 50-bp sliding window (Schloss et al. 2011). After this filtering step, reads were aligned against the SILVA database release 123 (Quast et al. 2013) and reads that could not be reliably assigned to the V5-V6 region were discarded. A pre-cluster step was applied following Huse et al. (2010). Potential chimeric sequences were detected and removed using CHIMERA UCHIME (Edgar et al. 2011). Then, reads were clustered into operational taxonomic units (OTUs) by a clustering at 97% similarity (Schloss and Westcott 2011). OTUs were affiliated to taxonomic groups according to the SILVA database using an 80 % minimum bootstrap cutoff (Wang et al. 2007). DNA sequences classified as *Archaea*, *Eukaryota*, chloroplasts or mitochondria were discarded.

Data deposition

Sequence data are available at the European Nucleotide Archive (<https://www.ebi.ac.uk/ena>) under the project accession #PRJEB29925. Sample accession #ERS2912777 (SAMEA5128392) = run2. See Table S1 for barcode information.

Data analysis

The effects of artificial recharge (i.e., infiltration) and vadose zone thickness (VZT) on DOC and BDOC concentrations, biofilm characteristics (i.e., proteins, FDA and INT assays) and diversity (number of OTUs, Shannon diversity index, and Pielou's evenness index) were tested using repeated-measure analyses of variance (2-way RM-ANOVA) with VZT as fixed factor and recharge as repeated factor to take into account the non-independence of recharged and non-recharged wells in each site. Pearson's correlations were computed to test the significance of the relations between DOM (i.e., DOC and BDOC) and clay bead biofilm variables (i.e. biomass, activities, OTU richness, and diversity indices). Most variables were logit transformed to satisfy the normality assumption verified by Shapiro's test. Sphericity assumption for RM-ANOVA was verified using Mauchly's sphericity test. As Log transformed qPCR datasets did not satisfy the assumption of normality, Kruskal-Wallis (KW) and Spearman correlation tests were applied on the qPCR datasets.

A non-metric multidimensional scaling (NMDS) was performed based on the Bray-Curtis dissimilarity distances. Differences in bacterial community structures among biofilms from recharged and non-recharged groundwaters in AR sites with thin and thick vadose zones (4 groups taking into account 2 VZT conditions * 2 recharge conditions) were tested using permutational multivariate analysis of variance (PERMANOVA; Anderson 2001). Statistical tests were based on 999 permutations of the Bray-Curtis matrix and significance was accepted at p-value < 0.05. All statistical tests were performed using the R software (R Development Core Team 2008). The ade4 and Vegan packages were used for nMDS and PERMANOVA analyses (Oksanen et al. 2007).

Results

DOM concentrations in groundwater of AR sites

DOC and BDOC concentrations in groundwater averaged $1.19 \pm 0.55 \text{ mg L}^{-1}$ and $0.56 \pm 0.27 \text{ mg L}^{-1}$, respectively. Both DOC and BDOC concentrations were significantly higher in recharged groundwaters than in non-recharged groundwaters (Fig. 2AB, 2-way RM-ANOVA, recharge effect, $p < 0.05$ for DOC and BDOC concentrations). DOC concentrations in recharged groundwater averaged $1.52 \pm 0.54 \text{ mg L}^{-1}$ whereas measured concentrations in non-recharged groundwater were $0.87 \pm 0.33 \text{ mg L}^{-1}$. The same pattern was observed for BDOC concentrations with higher values measured in recharged groundwater ($0.69 \pm 0.26 \text{ mg L}^{-1}$) in comparison with concentrations from non-recharged groundwater ($0.42 \pm 0.22 \text{ mg L}^{-1}$). The mean enrichment in DOM concentrations induced by groundwater recharge (recharged versus non-recharged groundwater) was more important in AR sites with thin vadose zones (mean enrichment of 0.89 mg L^{-1} and 0.37 mg L^{-1} for DOC and BDOC, respectively) than in AR sites with thick vadose zones (mean enrichment of 0.40 mg L^{-1} and 0.17 mg L^{-1} for DOC and BDOC, respectively). Consequently, DOC and BDOC concentrations in groundwaters were significantly higher in AR site with a thin vadose zone than in AR sites with a thick vadose zone (2-way RM-ANOVA, VZT effect, $p < 0.03$).

Influence of groundwater recharge on biofilm biomass and activities

Biofilm growth parameters were significantly influenced by recharge of groundwater with stormwater runoffs (2-way RM-ANOVA, recharge effect, $p < 0.05$). Total proteins measured on clay beads were significantly higher in biofilms developed in recharged groundwaters (mean total proteins: $7.34 \pm 5.09 \text{ } \mu\text{g per cm}^2$ of clay beads) than in biofilms recovered from non-recharged groundwaters (mean total proteins: $4.23 \pm 1.79 \text{ } \mu\text{g per cm}^2$ of clay beads) (Fig. 2C). The same trends were observed for hydrolytic and dehydrogenase activities measured on biofilms (Fig. 2DE) although no significant statistical results were obtained due to the high variability among sites (2-way RM-ANOVA, recharge effect, $0.2 > p > 0.15$ for hydrolytic activity and $0.1 > p > 0.05$ for dehydrogenase activity). On average, hydrolytic activity was more than 4-fold higher for biofilms developed in recharged

groundwaters compared to those developed in non-recharged groundwaters (Fig. 2D). Groundwater recharge increased by more than 1.5-fold the dehydrogenase activity of biofilms over the activities obtained from non-recharged groundwaters (Fig. 2E). Quantitative PCR assays used to determine bacterial concentrations over the bead surfaces followed similar trends. High bacterial contents among the clay bead biofilms going from 2×10^6 to 1.88×10^7 16S rRNA gene copies per cm^2 of clay beads incubated in non-recharged groundwater, and 2.35×10^6 to 6×10^8 copies per cm^2 of clay beads incubated in recharged groundwater (Table S2). 16S rRNA qPCR DNA counts were significantly higher in clay bead biofilms from recharged wells with a thin vadose zone rather than those with a thick vadose zone (Kruskal-Wallis tests, $p < 0.01$). This VZT effect was confirmed with the other biofilm biomass and activities parameters also giving higher values in AR sites with thin vadose zones rather than thick ones (2-way RM-ANOVA, VZT effect, $p < 0.05$ for the three variables). Although statistical analyses were close to significant level (2-way RM-ANOVA, “recharge x VZT” interaction, $0.1 > p > 0.05$ for dehydrogenase activity), biomass and activities tended to be more stimulated by recharge in AR sites with vadose zone thicknesses < 3 m in AR sites with vadose zone thicknesses > 10 m.

Influence of groundwater recharge on biofilm 16S rRNA gene diversity indices and bacterial community structure

The OTU numbers were estimated from 16S rRNA V5-V6 DNA segments generated from the clay bead biofilm DNA extracts (after a re-sampling step set at 32,000 sequences per sample, after the trimming-cleaning processes). These DNA reads were grouped into OTU when showing 3 % or less divergence. Accordingly, the DNA extracts from recharged groundwaters were more diversified than biofilms from non-recharged ones (2-way RM-ANOVA, recharge effect, $p < 0.05$). Numbers of detected OTUs were 1224 ± 472 in recharged groundwater and 870 ± 202 in non-recharged groundwaters (Fig. 2F). OTU abundance also depended on VZT (2-way RM-ANOVA, VZT effect, $p < 0.02$). OTU increases were mainly observed in AR sites with vadose zone thicknesses < 3 meters

(mean increase of 607 OTUs). The other three sites with thicker vadose zones showed no clear changes in OTU numbers associated with AR (mean decrease of 102 OTUs). Shannon diversity indices computed from OTU numbers were higher in biofilms from recharged groundwaters than non-recharged ones (Fig. 2G, 2-way RM-ANOVA, recharge effect, $p < 0.05$). Pielou's evenness index indicated that these differences in OTU were associated with an increase in evenness among OTUs from biofilms of the recharged sites (Fig. 2H, 2-way RM-ANOVA, recharge effect, $p < 0.05$). In addition, the NMDS ordination of OTUs from clay beads showed significant differences in bacterial community structure between biofilms developing in groundwater for the four tested conditions (2 recharge conditions * 2 VZT condition, Fig. 3, PERMANOVA, $p < 0.001$).

Core bacterial taxa inferred from aquifer biofilm 16S rRNA gene sequences

Only 130 OTUs which corresponded to 1.38 % of the total number of OTUs (9435) were shared between the 36 clay bead biofilm samples. Nonetheless, these 130 OTUs represented 71.66 % of all sequences. The largest groups of core OTUs were allocated to the following phyla: *Proteobacteria* (109 OTUs with 87 % of sequences), followed by *Bacteroidetes* (13 OTUs with 11 % of sequences), *Actinobacteria* (5 OTUs with 1.6 % of sequences) and *Nitrospirae* (3 OTUs with only 0.05 % of sequences). *Proteobacteria* were mainly represented by *Gamma-Proteobacteria* (28 OTUs with 44 % of sequences), *Beta-Proteobacteria* (50 OTUs with 32 % of sequences) and *Alpha-Proteobacteria* (27 OTUs with 10% of sequences).

Core clay bead biofilm communities of non-recharged groundwaters and recharged groundwaters were similar with *Proteobacteria* being predominant according to number of OTUs and read numbers (Fig. 4). *Proteobacteria* was followed by *Bacteroidetes*, and *Actinobacteria*. Other phyla such as *Nitrospirae* and *Elusimicrobia* were found but in smaller proportions in both core communities. *Chlorobi* and *Chloroflexi* were only detected in core communities of recharged groundwaters, while *Firmicutes* were only found in core communities of non-recharged groundwaters. Except these rare phyla, the main difference between the core communities of recharged and non-recharged

groundwaters was observed on the relative proportions of *Bacteroidetes* (Fig. 4): *Bacteroidetes* were represented more than 3-fold in the core community of recharged groundwaters (17.1 % of sequences) than in core community of non-recharged (4.9 % of sequences). Quantitative PCR assays confirmed these trends (Table S2). Concentrations of total *Bacteroides* were found significantly higher by qPCR in clay bead biofilm DNA extracts from recharged wells than those recovered from non-recharged wells (Table S2) (for each VZT category, Kruskal-Wallis, p-value <0.01). Furthermore, biofilms from shallow aquifers were always found to have significantly higher total *Bacteroides* counts than those from aquifers with higher VZT (KW p-value <0.01). Total *Bacteroides* DNA could not be detected among most clay bead biofilms recovered from non-recharged wells from sites with a thick VZT except for two samples in the RAQ site (Table S2).

Relations between DOM and biofilm biomass, activities and diversity

Biofilm characteristics such as qPCR datasets, microbial biomass estimated from Lowry's total proteins estimations or microbial activities estimated from FDA and INT were positively and significantly correlated with the concentrations of DOC and BDOC associated with stormwater infiltration (e. g. Table 1), with the exception of the relation between FDA activity and DOC that was not significant. Nevertheless, stronger correlations between these three microbial parameters with BDOC were observed rather than with DOC. Number of bacterial cells among aquifer biofilms as estimated by qPCR were positively correlated with DOC and BDOC (Spearman's rho factor > 0.6, p < 0.01). A positive and significant correlation was also detected between OTU number and BDOC (r = 0.79; p < 0.05) and between OTU number and DOC (r = 0.74, p < 0.05). When considering the main bacterial phyla, a positive correlation was observed between the number of OTUs belonging to *Bacteroidetes* and BDOC (r = 0.71; p < 0.01) and between *Bacteroidetes* and DOC (r = 0.67; p < 0.05) (Fig. 5). This correlation held true using total *Bacteroides* qPCR counts (Spearman's rho factor > 0.7, p < 0.01) (according to Table S2). Furthermore, the total *Bacteroides* qPCR counts were found correlated with qPCR counts for the HF183 *Bacteroides* target which is specific for species associated

with the human intestinal tract (Table S2) (Spearman's rho factor > 0.7, $p < 0.01$). HF183 *Bacteroides* qPCR counts were also correlated with the DOC and BDOC values (Spearman's rho factor > 0.56, $p < 0.01$). HF183 were only detected in biofilms from AR sites with a thin vadose zone (all biofilm samples from recharged groundwater of GB site and one biofilm sample from recharged groundwater of IUT site, Table S2). The number of OTUs belonging to *Alpha-proteobacteria* and *Beta-proteobacteria* also significantly increased with DOC ($r = 0.81$; $p < 0.01$ for *Alpha-proteobacteria* and $r = 0.71$; $p < 0.05$ for *Beta-proteobacteria*) and BDOC concentrations ($r = 0.79$; $p < 0.01$ for *Alpha-proteobacteria* and $r = 0.65$; $p < 0.05$ for *Beta-proteobacteria*) (Fig. 5). In the dataset, one other phylum, *Chlorobi*, showed a significant and positive correlation between OTU numbers and BDOC concentrations ($r = 0.66$ and $p < 0.05$) (Fig. 5).

Discussion

The influence of AR practices was very well illustrated by the DOC contents of groundwaters. The relationships between DOC and vadose zone thicknesses confirmed the hypothesis that DOM enrichment of groundwaters decreases with VZT (Pabich et al. 2001; Datry et al. 2006). Concentrations of total dissolved organic carbon (DOC) and biodegradable dissolved organic carbon (BDOC) were significantly higher in recharged than non-recharged groundwaters but mainly for AR sites with thin vadose zones (< 3 m). This effect was not significant for AR sites with a thick vadose zone (> 10 m). The influence of VZT on DOM reaching groundwater was thus linked to the DOM soil retention capacity. Biotic (microbial consumption of BDOC, Marschner and Kalbitz 2003; Li et al. 2013; Shen et al. 2015) and abiotic (sorption of DOM on clay particles, soil organic matter, or/and Fe and Al oxides/hydroxides, McKnight et al. 1992; Kalbitz et al. 2000; Saidy et al. 2013) soil components are likely involved in this VZT-related effect. Consequently, and according to several studies (e.g., Pabich et al. 2001; Shen et al. 2015), the thicker the VZ, the more efficient is its ability to limit groundwater contamination with DOM.

Based on groundwater samples from Cape Cod (USA), Pabich et al. (2001) determined a statistical relationship between VZT (in m) and DOC concentration (in mg L⁻¹) in groundwater: DOC concentration = DOC concentration in surface water * e^{-0.7*VZT}. By applying the same relationship using a mean DOC concentration of 5 mg L⁻¹ in surface water (according to Voisin et al. 2018), we calculated DOC concentrations ranging between 0.9 and 3.05 mg L⁻¹ for groundwaters influenced by recharge in sites with vadose zones < 3 meters. Interestingly, the measured DOC concentrations at these sites were in the same range (i.e., 1.6 – 2.3 mg L⁻¹). One would have expected that AR would have produced higher DOC concentrations than those expected from the model of Pabich et al. (2001) because of the stronger hydrological connectivity between the surface and the aquifer in these AR sites (Foulquier et al. 2011). Two mechanisms could explain the concordance between our measurements and simulations by the model of Pabich and colleagues: (1) soil and VZ of AR systems were very efficient in DOC retention (Zhang et al. 2012; Mermillod-Blondin et al. 2015) and (2) DOC from infiltrating waters was significantly diluted by groundwater (poor in DOC) in recharge zones of the aquifer (Foulquier et al. 2010). In some of our AR sites, two studies (Mermillod-Blondin et al. 2015; Voisin et al. 2018) tried to determine the respective influences of dilution and retention in organic matter dynamics by using chloride as conservative tracer to evaluate the role of dilution. They found that the dilution of surface water with groundwater could not explain the decrease of DOC concentrations from surface waters to the aquifer. Then, it is expected that abiotic and biotic retention of DOC during water infiltration through the soil and VZ was the main mechanism involved in DOC dynamics in AR sites. Nevertheless, this retention process was not efficient enough to limit significant DOC and BDOC enrichments of groundwater in AR sites with thin VZ. The concentration of organic carbon was thus a good indicator of the connectivity between surface and groundwater ecosystems.

As most groundwater ecosystems experience strong carbon limitation that severely limit growth of microbial communities (Bengtsson 1989; Kazumi and Capone 1994; Baker et al. 2000; Goldscheider 2006), the biomass (total proteins), 16S rRNA gene counts, and activities of biofilm that grew on

artificial substrates (clay beads) introduced in the aquifer were positively linked with DOC enrichment. The vadose zone thickness was also significantly related to the observed changes in microbial activities that were measured: artificial recharge significantly stimulated micro-organisms for AR sites with a thin vadose zone whereas it was not the case in AR sites with a thick vadose. To our knowledge, these results are the first to demonstrate that VZT influences microbial growth in groundwaters through its control over DOM availability in AR systems. Indeed, Datry et al. (2005) demonstrated the influence of VZT on DOC concentrations and groundwater invertebrates but they did not evaluate the response of the micro-organisms which are known to mediate energy fluxes from DOM towards invertebrates (Cooney and Simon 2009). Foulquier et al. (2011) estimated bacterial cell numbers (i.e., DAPI staining counts and Lowry's total proteins assays) and microbial activities (i.e., esterase and dehydrogenase activities) on sediments collected from three MAR systems with a thin VZ but they did not consider systems with thick VZ.

The scarcity of studies on microbial biofilms from deep alluvial aquifers (i.e., when the vadose zone thickness exceeds 10 m) results from technical difficulties and costs associated with pumping or coring for sediment collection (Chapelle 2001). Thus, the use of artificial substrates (clay beads) appeared an efficient strategy to compare biofilm growth development and activities among sites of contrasted VZT including deep aquifers. The positive and significant correlations obtained between microbial datasets from biofilms developing on clay beads and DOC and/or BDOC concentrations (Table 1) confirmed the efficacy of this approach for monitoring trophic conditions in groundwater ecosystems (Williamson et al. 2012; Mermillod-Blondin et al. 2013; Voisin et al. 2016).

The influence of AR practices on the OTU richness of biofilms collected from artificial substrates (clay beads) was also related to the VZT. Indeed, OTU number was significantly higher in recharged groundwaters than in non-recharged ones for AR sites with thin VZ. In contrast, no significant effect of recharge was measured on OTU numbers in AR sites with thick VZ. These increasing OTU numbers were found to match the increase of 16S rRNA Bacterial qPCR counts observed in recharged wells of AR sites with a thin vadose zone in comparison with recharged wells from AR sites with a

thick vadose zone. In addition, Shannon diversity and Pielou's evenness indices indicated that the higher OTU richness of biofilms from recharged wells in comparison with biofilms from non-recharged wells was not due to the addition of relatively rare or relatively abundant taxa in the community. It rather seemed to be associated with an increasing in numbers of less frequent OTU at the non-recharged sites, leading to a greater evenness in the datasets between the OTUs of the recharged sites. This confirms that bacterial community structures were significantly influenced by recharge (more OTUs and higher evenness in the bacterial community). NMDS ordination of OTUs and PERMANOVA test also highlighted that bacterial community structures of biofilms incubated in groundwaters were significantly different among the four tested conditions (2 VZT conditions * 2 recharge conditions). According to the observations of Hug et al. (2015) on groundwater microorganisms, the spatial distribution of the six AR sites (Fig. 1) would have led to high differences in bacterial community structures among these sites, reducing the ability to detect a significant influence of VZT and recharge conditions on bacterial communities using AR sites as replicates. As it was not the case, VZT and recharge could be considered as main factors structuring the bacterial communities of groundwater biofilms in the present study.

The V5-V6 16S rRNA gene-inferred richness indices were found coherent with changes in DOC availability in groundwaters associated with recharge, and positive significant relations were obtained between OTU numbers and BDOC concentrations. These trends were further confirmed by qPCR assays targeting *Bacteroides* species. Biofilms from shallow aquifers were found to have significantly higher total *Bacteroides* counts than those from aquifers with higher VZT. Furthermore, the application of a more specific PCR screening for *Bacteroides* species specialized for the human intestinal tract confirmed a strong relation between their counts in an aquifer biofilm and DOC/BDOC. Thus, the hypothesis of an increased richness of bacterial "species" (inferred from OTU numbers) with the availability of nutrient (i.e., BDOC being the biodegradable fraction of DOC for bacteria) in groundwater ecosystems is further supported by these datasets. However, DOC transfers were also found to point out probable transfers of allochthonous bacteria associated with organic matters such as

those coming from human wastes.

We also showed that biomass (concentrations of total proteins), activities (fluorescein diacetate (FDA) hydrolysis, INT-formazan reduction) and 16S rRNA genetic diversity of biofilms growing on clay beads (in groundwaters) were all correlated positively with DOC and BDOC concentrations. These results are in line with the “more individuals” hypothesis (Srivastava and Lawton 1998) of the species–energy theory (Wright 1983; Cardinale et al. 2009) which indicates that resource availability positively influenced species richness by increasing population sizes and reducing the probability of extinction. This positive response of bacterial diversity with resource availability is more likely expected in ecosystems experiencing strong nutrient limitations (see Bienhold et al. 2012 for an example in Arctic deep-sea sediments). Given the oligotrophic conditions encountered in groundwater environments, it is thus probable that increased DOC and BDOC concentrations associated with AR practices would enable the maintenance of copiotrophic bacteria that were not adapted to live in the more oligotrophic areas of aquifers (e.g., Fierer et al. 2007; Lever et al. 2015). These diversity changes were related to changes in the proportions of bacterial phyla among the core communities. The relative proportion of *Bacteroidetes* increased in the recharged groundwaters as observed from the OTU profiling and qPCR datasets. These observations are consistent with an increase of copiotrophic bacteria with increasing energy availability because members of the phylum *Bacteroidetes* are often abundant in nutrient-rich aquatic environments (e.g., Gomez-Pereira et al. 2010; Landa et al. 2013; Marti et al. 2017). Indeed, this phylum has been detected in most aquatic habitats (e.g., Alonso et al. 2007; Diez-vives et al. 2014; Schauer et al. 2010) but bacteria of this group which are known to degrade polymeric organic matter (Cottrell and Kirchman 2000) are often associated with blooms of phytoplankton (e.g., Pinhassi et al. 2004; Teeling et al. 2012; Tada et al. 2011) and fecal contaminations (e. g. Marti et al. 2017; Bernardin-Souibgui et al. 2018). The latter origin was confirmed for some species of this phylum which are specific of the human intestinal tract (HF183 DNA target). The HF183 DNA target was found in the aquifer of AR sites with a shallow aquifer (but not for those with a thick vadose zone), suggesting a potential transfer from stormwater

runoff to the aquifer. It is also to be noted that qPCR screenings for *Bacteroides* species appeared good indicators of the ability of a vadose zone at preventing bacterial contamination of the aquifer with surface bacteria. The total *Bacteroides* PCR screening could only detect this bacterial group among biofilm samples recovered from recharged wells of AR sites. In addition to the effects observed on *Bacteroidetes*, the diversities of most phyla in terms of OTU numbers showed positive linear relations with BDOC concentrations. The fact that dominant bacterial groups increased in diversity with nutrient (BDOC) availability in groundwater environment might have resulted from the harsh trophic conditions encountered by micro-organisms in subterranean ecosystems (Chapelle and Lovley 1990). Indeed, we measured relatively low BDOC concentrations (below 0.9 mg L⁻¹) in groundwater samples. However, Li et al. (2013) showed that an increase in BDOC concentration from 0.5 mg L⁻¹ to 1.1 mg L⁻¹ was high enough to reduce 16S rRNA gene bacterial diversity of biofilms developing in sediment columns (under laboratory conditions). We can thus not exclude that a reduction of the diversity of some bacterial groups could occur due to BDOC enrichments of groundwater associated with AR practices if these enrichments lead to BDOC concentrations exceeding 1.1 mg L⁻¹. Here, coupling measurements of DOC availability, biofilm variables, bacterial richness and proportion of particular taxa among bacterial communities (e. g. all *Bacteroides* or particular species) appeared sufficient to infer the relative incidence of AR practices on aquifers. This approach based on functional indicators appears promising for water manager to evaluate groundwater ecosystem health. Moreover, the results of the present study clearly showed that the implementation of AR sites in porous aquifers would be recommended in areas with thick VZ (> 10 m) to prevent groundwater contamination.

Acknowledgments The authors thank Léa Dantony, Mathilde Pigneret and Felix Vallier for their support and advice during field and laboratory work and Pierre Marmonier for editing an earlier draft of this manuscript.

Funding The authors thank the Auvergne-Rhône-Alpes region for Jérémy Voisin's Ph.D grant. This work was supported by l'Agence Nationale de la Recherche [ANR-16-CE32-0006 FROG], Lyon Metropole and Agence de l'Eau Rhône-Méditerranée-Corse within the framework of the experimental observatory for urban hydrology (OTHU, <http://www.graie.org/othu>) and Rhône Basin LTER (ZABR), and the French national research program for environmental and occupational health of Anses under the terms of project "Iouqmer" EST 2016/1/120.

Conflicts of interest. None declared.

References

- Alonso C, Warnecke F, Amann R, Pernthaler J (2007) High local and global diversity of Flavobacteria in marine plankton. *Environ Microbiol* 9:1253–1266.
- Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance. *Austral Ecol* 26:32–46.
- Baker MA, Valett HM, Dahm CN (2000) Organic carbon supply and metabolism in a shallow groundwater ecosystem. *Ecology* 81:3133–3148.
- Battin TJ (1997) Assessment of fluorescein diacetate hydrolysis as a measure of total esterase activity in natural stream sediment biofilms. *Sci Total Environ* 198:51–60.
- Bengtsson G (1989) Growth and metabolic flexibility in groundwater bacteria. *Microb Ecol* 18:235–248.
- Bernardin-Souibgui C, Barraud S, Bourgeois E, Aubin JB, Becouze-Lareure C, Wiest L, Marjolet L, Colinon C, Lipeme Kouyi G, Cournoyer B, Blaha D (2018). Incidence of hydrological, chemical, and physical constraints on bacterial pathogens, *Nocardia* cells, and fecal indicator bacteria trapped in an urban stormwater detention basin in Chassieu, France. *Env Sci Poll Res*, 25:24860-24881.
- Bernstein HC, Brislawn C, Renslow RS, Dana K, Morton B, Lindemann SR, Song H-S, Atci E,

- Beyenal H, Fredrickson JK, Jansson JK, Morand JJ (2017) Trade-offs between microbiome diversity and productivity in a stratified microbial mat. *ISME J* 11:405-414.
- Bienhold C, Boetius A, Ramette A (2012) The energy–diversity relationship of complex bacterial communities in Arctic deep-sea sediments. *ISME J* 6:724–732.
- Cardinale BJ, Hillebrand H, Harpole WS, Gross K, Ptacnik R (2009) Separating the influence of resource "availability" from resource "imbalance" on productivity-diversity relationships. *Ecol Lett* 12:475–487.
- Chapelle FH (2001) *Groundwater microbiology and geochemistry*. Wiley Press, New York.
- Chapelle FH, Lovley DR (1990) Rates of microbial metabolism in deep coastal plain aquifers. *Appl Environ Microbiol* 56:1865–1874.
- Cooney TJ, Simon KS (2009) Influence of dissolved organic matter and invertebrates on the function of microbial films in groundwater. *Microb Ecol* 58:599–610.
- Cottrell MT, Kirchman DL (2000) Community composition of marine bacterioplankton determined by 16S rRNA gene clone libraries and fluorescence in situ hybridization. *Appl Environ Microbiol* 66:5116–5122.
- Datry T, Malard F, Gibert J (2004) Dynamics of solutes and dissolved oxygen in shallow urban groundwater below a stormwater infiltration basin. *Sci Total Environ* 329:215–229.
- Datry T, Malard F, Gibert J (2005) Response of invertebrate assemblages to increased groundwater recharge rates in a phreatic aquifer. *J North Am Benthol Soc* 24:461-477.
- Datry T, Malard F, Gibert J (2006) Effect of artificial stormwater infiltration on urban groundwater ecosystems. In: Tellam JH, Rivett MO, Israfilov RG, Herringshaw LG (Eds). *Urban Groundwater Management and Sustainability*. Springer, Dordrecht, 331-345.
- Díez-Vives C, Gasol JM, Acinas SG (2014) Spatial and temporal variability among marine Bacteroidetes populations in the NW Mediterranean Sea. *Syst Appl Microbiol* 37:68–78.
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27:2194–200.

562 Evans KL, Warren PH, Gaston KJ (2005) Species–energy relationships at the macroecological scale:
563 a review of the mechanisms. *Biol Rev* 80:1–25.

564 Fierer N, Bradford MA, Jackson RB (2007) Toward an ecological classification of soil bacteria.
565 *Ecology* 88:1354–64.

566 Fontvieille DA, Outaguerouine A, Thevenot DR (1992) Fluorescein diacetate hydrolysis as a measure
567 of microbial activity in aquatic systems: Application to activated sludges. *Environ Technol*
568 13:531–40.

569 Foulquier A, Malard F, Barraud S, Gibert J (2009). Thermal influence of urban groundwater recharge
570 from stormwater infiltration basins. *Hydrol Proc* 23: 1701-13.

571 Foulquier A, Malard F, Mermillod-Blondin F, Datry T, Simon L, Montuelle B, Gibert J (2010) Vertical
572 change in dissolved organic carbon and oxygen at the water table region of an aquifer recharged
573 with stormwater: biological uptake or mixing? *Biogeochemistry* 99:31–47.

574 Foulquier A, Malard F, Mermillod-Blondin F, Montuelle B, Dolédec S, Volat B, Gibert J (2011)
575 Surface water linkages regulate trophic interactions in a groundwater food web. *Ecosystems*
576 14:1339–53.

577 Fuhrman JA (2009) Microbial community structure and its functional implications. *Nature* 459:193–
578 199.

579 Gibert J, Danielopol DL, Stanford JA (1994) *Groundwater ecology*. Academic Press, San Diego,
580 California, USA.

581 Goldscheider N, Hunkeler D, Rossi P (2006) Review: Microbial biocenoses in pristine aquifers and
582 an assessment of investigative methods. *Hydrogeol J* 14:926–941.

583 Gómez-Pereira PR, Fuchs BM, Alonso C, Oliver MJ, Van Beusekom JE, Amann R (2010) Distinct
584 flavobacterial communities in contrasting water masses of the north Atlantic Ocean. *ISME J*
585 4:472–487.

586 Griebler C, Lueders T (2009) Microbial biodiversity in groundwater ecosystems. *Freshw Biol*
587 54:649–677.

588 Hershey OS, Kallmeyer J, Wallace A, Barton MD, Barton HA (2018) High microbial diversity despite
589 extremely low biomass in a deep karst aquifer. *Frontiers Microbiol* 9.
590 <https://www.frontiersin.org/articles/10.3389/fmicb.2018.02823/full>

591 Houri-Davignon J-C, Etcheber HC (1989) Measurement of actual electron transport system (ETS)
592 activity in marine sediments by incubation with INT. *Environ Technol* 10:91–100.

593 Hug LA, Thomas BC, Brown CT, Frischkorn KR, Williams KH, Tringe SG, Banfield JF (2015)
594 Aquifer environment selects for microbial species cohorts in sediment and groundwater. *ISME*
595 *J* 9:1846–1856.

596 Hunt W, Traver R, Davis A, Emerson CH, Collins KA, Stagge JH (2010) Low impact development
597 practices: designing to infiltrate in urban environments. In: Chang NB (Ed). *Effects of*
598 *urbanization on groundwater*. American Society of Civil Engineers, 308–343.

599 Huse SM, Welch DM, Morrison HG, Sogin ML (2010) Ironing out the wrinkles in the rare biosphere
600 through improved OTU clustering. *Environ Microbiol* 12:1889–1898.

601 Kalbitz K, Solinger S, Park J-H, Michalzik B, Matzner E (2000) Controls on the dynamics of
602 dissolved organic matter in soils: a review. *Soil Sci* 165:277–304.

603 Kazumi J, Capone D (1994) Heterotrophic microbial activity in shallow aquifer sediments of Long
604 Island, New York. *Microb Ecol* 28:19–37.

605 Kolehmainen RE, Langwaldt JH, Puhakka JA (2007) Natural organic matter (NOM) removal and
606 structural changes in the bacterial community during artificial groundwater recharge with humic
607 lake water. *Water Res* 41:2715–2725.

608 Landa M, Cottrell MT, Kirchman DL, Blain S, Obernosterer I (2013) Changes in bacterial diversity
609 in response to dissolved organic matter supply in a continuous culture experiment. *Aquat Microb*
610 *Ecol* 69:157–168.

611 Layton A, McKay L, Williams D, Garrett V, Gentry R, Sayler G (2006) Development of *Bacteroides*
612 16S rRNA gene TaqMan-based real-time PCR assays for estimation of total, human, and
613 bovine fecal pollution in water. *Appl Environ Microbiol* 72:4214–4224.

- Lever MA, Rogers KL, Lloyd KG, Overmann J, Schink B, Thauer RK, Hoehler TM, Jørgensen BB (2015) Life under extreme energy limitation: A synthesis of laboratory and field-based investigations. *FEMS Microbiol Rev* 39:688–728.
- Li D, Alidina M, Ouf M, Sharp JO, Saikaly P, Drewes JE (2013) Microbial community evolution during simulated managed aquifer recharge in response to different biodegradable dissolved organic carbon (BDOC) concentrations. *Water Res* 47:2421–30.
- Marsalek J, Chocat B (2002) International report: Stormwater management. *Water Sci Technol* 46:1–17.
- Marschner B, Kalbitz K (2003) Controls of bioavailability and biodegradability of dissolved organic matter in soils. *Geoderma* 113:211–235.
- Marti R, Ribun S, Aubin J-B, Colinon C, Petit S, Marjolet L, Gourmelon M, Schmitt L, Breil P, Cottet M, Cournoyer B (2017) Human-driven microbiological contamination of benthic and hyporheic sediments of an intermittent peri-urban river assessed from MST and 16S rRNA genetic structure analyses. *Front. Microbiol.* 8:19.
- McKnight DM, Bencaia KE, Zeeiweger GW, Aiken GR, Feder GL, Thorn KA (1992) Sorption of dissolved organic carbon by hydrous aluminum and iron oxides occurring at the confluence of Deer Creek with the Snake River, Summit County, Colorado. *Environ Sci Technol* 26:1388–1396.
- Mermillod-Blondin F, Foulquier A, Maazouzi C, Navel S, Negrutiu Y, Vienney A, Simon L, Marmonier P (2013) Ecological assessment of groundwater trophic status by using artificial substrates to monitor biofilm growth and activity. *Ecol Indic* 25:230–238.
- Mermillod-Blondin F, Simon L, Maazouzi C, Foulquier A, Delolme C, Marmonier P (2015) Dynamics of dissolved organic carbon (DOC) through stormwater basins designed for groundwater recharge in urban area: Assessment of retention efficiency. *Water Res* 81:27–37.
- Mermillod-Blondin F, Voisin J, Marjolet L, Marmonier P, Cournoyer B (2019) Clay beads as artificial trapping matrices for monitoring bacterial distribution among urban stormwater infiltration

- systems and their connected aquifers. *Env Monitor Assess* 191:58. <https://link-springer-com.docelec.univ-lyon1.fr/article/10.1007/s10661-019-7190-0>
- Oksanen J, Kindt R, Legendre P, O'Hara BV (2007) The vegan package. Community ecology package, R package version 1.
- Pabich WJ, Valiela I, Hemond HF (2001) Relationship between DOC concentration and vadose zone thickness and depth below water table in groundwater of Cape Cod, USA. *Biogeochemistry* 55:247–68.
- Park J-W, Crowley DE (2006) Dynamic changes in *nahAc* gene copy numbers during degradation of naphthalene in PAH-contaminated soils. *Appl Microbiol Biotechnol* 72:1322–1329.
- Peterson GL (1977) A simplification of the protein assay method of Lowry et al. which is more generally applicable. *Anal Biochem* 83:346–356.
- Pinhassi J, Sala MM, Havskum H, Peters F, Guadayol O, Malits A, Marrasé C (2004) Changes in bacterioplankton composition under different phytoplankton regimens. *Appl Environ Microbiol* 70:6753–6766.
- Prosser JJ, Bohannon BJ, Curtis TP, Ellis RJ, Firestone MK, Freckleton RP, Green JL, Green LE, Killham K, Lennon JJ, Osborn AM, Solan M, van der Gast CJ, Young JPW (2007) The role of ecological theory in microbial ecology. *Nature Rev Microbiol* 5:384–392.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO (2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* 41, D590–6.
- Saidy AR, Smernik RJ, Baldock JA, Kaiser K, Sanderman J (2013) The sorption of organic carbon onto differing clay minerals in the presence and absence of hydrous iron oxide. *Geoderma* 209–210:15–21.
- Schauer R, Bienhold C, Ramette A, Harder J (2010) Bacterial diversity and biogeography in deep-sea surface sediments of the South Atlantic Ocean. *ISME J* 4:159–170.
- Schloss PD, Westcott SL (2011) Assessing and improving methods used in operational taxonomic

unit-based approaches for 16S rRNA gene sequence analysis. Appl Environ Microbiol 77:3219–3226.

Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF (2009) Introducing mothur: Open-source, platform-independent, community supported software for describing and comparing microbial communities. Appl Environ Microbiol 75:7537–7541.

Schloss PD, Gevers D, Westcott SL (2011) Reducing the effects of PCR amplification and sequencing artifacts on 16S rRNA-based studies. PLoS One, 6, e27310. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0027310>

Servais P, Anzil A, Ventresque C (1989) Simple method for determination of biodegradable dissolved organic carbon in water. Appl Environ Microbiol 55:2732–2734.

Servais P, Billen G, Hascoët MC (1987) Determination of the biodegradable fraction of dissolved organic matter in waters. Water Res 21:445–450.

Seurinck S, Defoirdt T, Verstraete W, Siciliano SD (2005) Detection and quantification of the human-specific HF183 *Bacteroides* 16S rRNA genetic marker with real-time PCR for assessment of human faecal pollution in freshwater. Environ Microbiol 7:249-259.

Shen Y, Chapelle FH, Strom EW, Benner R (2015) Origins and bioavailability of dissolved organic matter in groundwater. Biogeochemistry 122:61–78.

Smith VH (2007) Microbial diversity-productivity relationships in aquatic ecosystems. FEMS Microbiol Ecol 62:181–186.

Tada Y, Taniguchi A, Nagao I, Miki T, Uematsu M, Tsuda A, Hamasaki K (2011) Differing growth responses of major phylogenetic groups of marine bacteria to natural phytoplankton blooms in the Western North Pacific Ocean. Appl Environ Microbiol 77:4055–4065.

Voisin J, Cournoyer B, Mermillod-Blondin F (2016) Assessment of artificial substrates for evaluating groundwater microbial quality. Ecol Indic 71:577–586.

Voisin J, Cournoyer B, Vienney A, Mermillod-Blondin F (2018) Aquifer recharge with stormwater

runoff in urban areas: Influence of vadose zone thickness on nutrient and bacterial transfers from the surface of infiltration basins to groundwater. *Sci Total Env* 637:1496-1507.

Wang Q (2007) Naïve Bayesian Classifier for Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Appl Environ Microbiol* 73:5261–5267.

Williamson WM, Close ME, Leonard MM, Webber JB, Lin S (2012) Groundwater biofilm dynamics grown in situ along a nutrient gradient. *Ground Water* 50:690–703.

Wright DH (1983) Species–energy theory: an extension of species–area theory. *Oikos* 41:496–506.

Zhang X, Zhao X, Zhang M (2012) Functional diversity changes of microbial communities along a soil aquifer for reclaimed water recharge. *FEMS Microbiol Ecol* 80:9–18.

703 Figure captions

704
2

705 Figure 1: Selected aquifer recharge (AR) sites in the eastern aquifer of the Lyon metropolitan area,
4
5
706 France. (a) Location of AR sites and (b) their main characteristics.

707
9

10
708 Figure 2: (a) Concentrations of dissolved organic carbon (DOC) and (b) biodegradable dissolved
11
12 organic carbon (BDOC) in non-recharged (NR, white) and recharged (R, gray) groundwater (GW).
13
14 (c) Total proteins (i.e., proxy of biofilm biomass), (d) hydrolytic (FDA), (e) dehydrogenase (INT)
15
16 (f) activities, (f) OTU numbers, (g) Shannon diversity index, and (h) Pielou's evenness index of biofilms
17
18 developed on artificial substrates (clay beads) incubated in non-recharged (NR, white) and recharged
19
20 (R, gray) groundwater (GW). The AR sites are used as replicates (n=3) for the two conditions of
21
22 vadose zone (VZ) thickness: thick VZ (> 10 meters) on the left of the panel and thin VZ (< 3 meters)
23
24 on the right of the panel.

25
26
27
28
29
30
31
32

33 Figure 3: NMDS ordination of V5-V6 16S rRNA gene OTU profiles from biofilms incubated in two
34
35 zones (recharged (gray centroid) and non-recharged (white centroid) groundwaters –GW-) over three
36
37 AR sites with a thin vadose zone (VZ < 3 m) and three AR sites with a thick vadose zone (> 10 m).
38
39 Stress test = 0.103

40
41
42
43

44
45 Figure 4: Relative numbers of core phyla among clay bead biofilms which developed over a 10 day
46
47 period among non-recharged (a and c) and recharged (b and d) groundwater. Panels a and b indicate
48
49 the relative numbers of OTU per phylum, and, c and d, the relative numbers of reads per phylum.

50
51
52
53

54
55 Figure 5: Linear regression analyses of BDOC values against number of OTU among selected phyla.
56
57 (a) total number of bacterial OTUs (b) *Bacteroidetes*, (c) *Chlorobi*, (d) *Alpha-proteobacteria* and (e)
58
59 *Beta-proteobacteria*. Pearson's r factors are indicated with their respective p-value. For *Bacteroidetes*
60

729 (panel b), the relationship between richness and BDOC values remains significant ($r = 0.7$, $p < 0.05$)
730 after suppression of the lowest and highest values of *Bacteroidetes* richness.

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig 1

a

- Moraine hill (glacial till)
- Groundwater flow direction
- AR sites

b

Main characteristics of AR sites

AR sites	Catchment area (ha)	Infiltration basin surface (m ²)	Vadose zone thickness (m)
DJR	185	8750	11
FEY	315	2550	12
RAQ	26	285	17
MIN	165	4200	2
IUT	2.5	850	2.5
GB	300	3200	0.7

Fig 1

Fig 2

Fig 2

Fig 3

Fig 3

Fig 4

Fig 4

Fig 5

Fig 5

Table 1 Pearson’s correlation coefficients between pairs of datasets produced in this study. DOC: dissolved organic carbon, BDOC: biodegradable organic carbon, FDA: Fluorescein diacetate (i.e., hydrolytic activity) and INT: 2-(p-iodophenyl)-3-(p-nitrophenyl)-5- phenyl tetrazolium chloride (i.e., dehydrogenase activity).

Variables	DOC	BDOC	Proteins	FDA	INT
BDOC	0.92***				
Proteins	0.65*	0.75**			
FDA	0.53	0.64*	0.97***		
INT	0.59*	0.70*	0.96***	0.95***	
OTU number	0.74**	0.79**	0.80**	0.78**	0.77**

*p < 0.05 **p < 0.01 ***p < 0.001

[Click here to access/download](#)
Supplementary Material
Table S1.xlsx

[Click here to access/download](#)
Supplementary Material
Table S2 qPCR assays bc1.xlsx