Desulfurization of Thioketals into Methylene and Methyl Derivatives: Nickel or not Nickel?

Guangkuan Zhao,^[a] Ling-Zhi Yuan,^[a] Mouad Alami,^[a] and Olivier Provot^{*[a]}

Abstract: In this review, the desulfurization of thioacetals and thioketals leading to methyl and methylene derivatives is summarized through metal and metal-free methodologies. In the section concerning the use of metals for desulfurization reactions, we will be successively presenting and discussing all methods requiring Ni and Ni derivatives as reducing agents and other examples in which metals were used to generate solvating

electrons as reducing species. The following section will concern the use of other desulfurizing agents as metal-hydrides (Bu_3SnH , Et_3SiH) and metal-free desulfurization reactions of thioketals. The functional group tolerance of the presented methods with functionalized and sensitive substrates will be discussed.

1. Introduction

2. Desulfurization of thioacetals using metals

- 2.1 Raney-Ni
- 2.2 Other Nickel derivatives
 - 2.2.1 NiCRAs from $\text{Ni}(\text{OAc})_2$, NaH and *t*-amylalcohol
 - 2.2.2 “ Ni_2B ” from NiCl_2 and NaBH_4
 - 2.2.3 $\text{NiBr}_2\cdot\text{DME}$, PPh_3 and LiAlH_4
 - 2.2.4 Nickelocene- LiAlH_4 hydride
- 2.3 Other metals
 - 2.3.1 Na and Li in liquid NH_3
 - 2.3.2 Li in EtNH_2
 - 2.3.3 LiAlH_4 associated with $\text{CuCl}_2/\text{ZnCl}_2$ or with TiCl_4
 - 2.3.4 Miscellaneous metal processes using $\text{Mo}(\text{CO})_6$ and Sml_2

3. Other desulfurizing reagents

- 3.1 Metal hydrides
 - 3.1.1 Bu_3SnH with AIBN
 - 3.1.2 $\text{B}(\text{C}_6\text{F}_5)_3$ catalyzed desulfurization of thioketals in the presence of Et_3SiH
- 3.2 Metal-free desulfurizations
 - 3.2.1 Hydrated hydrazine
 - 3.2.2 P_2I_4 in CH_2Cl_2
 - 3.2.3 Photoreduction of dithianes and dithiolanes with $t\text{BuOK}$ in DMSO
 - 3.2.4 EtSnA in MeOH

4. Conclusion

1. Introduction

Desulfurization of dithioacetal, thiolane and dithiane function into a methyl or methylene group is an important chemical transformation, which can be applied almost under neutral conditions. The desulfurizing process of thioacetal derivatives is often preferable to Clemmensen or Wolff-Kishner reductions of carbonyl compounds, particularly in the case of acidic or basic sensitive substrates. The Ni-promoted desulfurization transformation^[1a] of thioketals, known as “Mozingo reaction”, was firstly described in 1944 by Wolfrom and Karabinos, which used Raney-nickel to reduce a thioacetal function into a methylene group (Scheme 1).^[1b]

Scheme 1. First thioacetals hydrogenolysis by Raney-Ni.

In the present review, a compilation of the existing methods reported to achieve this transformation from various thioacetals of aldehyde and ketone derivatives will be highlighted. In the section 2 of this review, we will also present in a non-exhaustive manner, some selected examples using metals as reducing agents in which Raney-Ni^[2] and Ni-derivatives predominate beside other Raney-Ni of special activity (W1 to W8). The use of solvated electrons generated by other metals to perform these desulfurization reactions will be discussed afterwards. Section 3 will deal with desulfurization reactions performed in the presence of metal hydrides and the use of non-metallic processes will be explored.

[a] BioCIS, Univ. Paris-Sud, CNRS, Université Paris-Saclay, 92290, Châtenay-Malabry, France.
E-mail: olivier.provot@u-psud.fr

Supporting information for this article is given via a link at the end of the document. ~~((Please delete this text if not appropriate))~~

2. Desulfurization of thioacetals using metals

2.1. Raney-Ni

From all the methods of thioacetals desulfurization, the use of Raney-Ni is by far the most used method since the pioneer work of Wolfrom and Karabinos.^[1b] This is probably due to the easy access to this catalyst and to the fact that Raney-Ni tolerates a large number of functional groups.^[2] The preparation of Raney-Ni^[3] has been largely reported by Raney^[4] and others,^[5] who mixed finely powdered Nickel-Aluminum alloy in a hot alkali solution. One can note that during this process, a large quantity of molecular hydrogen (H_2) was produced with considerable amounts of H_2 remained on the metal. For this reason, a large number of desulfurization reactions were achieved without the need of an external source of H_2 . However, side-reactions such as debenzylation,^[6] carbon-carbon double bond reduction,^[7] hydrogenolysis of a carbon-halogen bonds^[8] and **deoxygenation of an epoxide**^[9] sometimes occurred, greatly lowering the yield of the process. In Scheme 2, we have selected some recent representative examples using Raney-Ni to desulfurize poly-functionalized molecules in total or complex syntheses.^[9,10]

It was noted that, alcohols like MeOH and EtOH are the solvents of choice in the great majority of these reducing transformations. A great functional group tolerance was observed in the selected examples requiring Raney-Ni among which heterocycles, ketones, quinones, alcohols, OTIPS, OBn,

NTs, esters, lactones and multiple bonds remained unchanged. It should be added that in some cases, the use of additional H_2 source is needed (e.g., reduction of compounds **3**, **5** and **11**).

Scheme 2. Application of desulfurization reactions by Raney-Ni in total syntheses

Guangkuan Zhao was born in 1988 in Shandong Province, China. He spent four years (2008–2012) in Ludong University pursuing bachelor degree of basic chemistry. After, he moved to Zhengzhou University where he obtained his master degree (organic chemistry, 2012–2015) under the supervision of Professor Dapeng Zou. He is currently a PhD student and works on chemical methodology and antitumor molecules with Dr. Olivier Provot in the Faculty of Pharmacy, University Paris-Saclay.

Dr. Ling-Zhi YUAN was born in 1987 in Huang-shi (Hubei Province, China) and granted with National Scholarship (2007–2008) during undergraduate period. After he received his master degree in 2013 with Prof. Jing-Yuan XU at Tianjin Medical University, he received his PhD in 2017 on selective metal-free deoxygenation of unsymmetrical 1,2-dicarbonyl compounds under the supervision of Dr. Olivier Provot in the Faculty of Pharmacy, University Paris-Saclay.

Dr. Mouad Alami, 58 years is a Research Director at the CNRS (Faculty of Pharmacy, BioCIS, France). He did his PhD at UPMC in Paris (1987) and after two years of post-doctoral position in collaboration with the Elf-Atochem company, he joined in 1989 the CNRS at the ENS-Paris (Dr. G. Linstrumelle). In 1993, he spent one year at the University of Santa-Barbara (Pr. B. H. Lipshutz). In 1999, he became CNRS Research Director at the Faculty of Pharmacy (University Paris-Saclay). His fields of specialization includes: (i) development of novel selective synthetic methodologies involving organometallic chemistry and (ii) synthesis of structurally challenging and biologically relevant products with antitumoral activity as vascular disrupting agents (VDAs) and their application in the context of antibody-drug conjugates.

Dr. Olivier Provot was born in 1963 in the beautiful city of Caen (Normandy, France) and received his PhD in total synthesis of ant venom alkaloids in the Lab of Pr. G. Lhommet in 1991 (UPMC, Paris VI). After a post-doctoral position in industry for two years and then for one year in the Lab of Medicinal Chemistry of Pr. R. Milcent (University Paris VII), he moved in the Lab of Pr. H. Moskowitz as assistant Professor in 1993 (University of Paris-Sud) and worked on the synthesis of natural peroxides having antimalarial activities. In 2003, he joined the group of Pr. J.-D. Brion and Dr. M. Alami (University Paris-Saclay) to develop novel organometallic methodologies to prepare novel anti-cancer drugs related to Combretastatin A-4.

2.2. Other Nickel derivatives

Even if Raney-Ni are efficient reagents in desulfurization reactions and very often compatible with functional groups, some experiments failed to give the expected products. Otherwise, the desulfurization properties of Raney nickels were attributed to the large excess of H_2 finely adsorbed on the catalyst's surface during its preparation,^[5a, 11] which can be deleterious with sensitive functionalized molecules. Moreover, Raney-Ni is highly pyrophoric and difficult to handle and it is also difficult to exactly determine the amount of nickel used for these transformations. To overcome these difficulties, a variety of non-pyrophoric nickel-containing reducing agents (NiCRAs) were prepared and used for the desulfurization of thioacetals and thioketals.

2.2.1. NiCRAs from $Ni(OAc)_2$, NaH and *t*-amylalcohol

In the 1980s, Caubère and co-workers developed easy to handle and to weigh nickel complexes from the reaction of NaH, *t*-amylalcohol and $Ni(OAc)_2$ with or without ligands. The NiCRAs (Ni-containing reducing agents) were able to reduce a wide variety of functions,^[12] and were also used as catalysts for coupling reactions.^[13] These NiCRAs complexes were also efficiently used to desulfurize a wide variety of sulfur-containing substrates including thioacetals^[14] because these nickel reagents possessed a high single electron transfer (SET) ability.^[15] Some representative examples are depicted in Scheme 3.

Scheme 3. Reagent and conditions: ^aNiCRA was prepared with *t*-AmOH (20 mmol) in THF, NaH (70 mmol) and $Ni(OAc)_2$ (10 mmol). Reaction performed on 1 mmol scale with NiCRA in THF at 65 °C with a ratio Ni/S = 30. ^bNiCRA-bpy was prepared with *t*-AmOH (20 mmol) in DME, NaH (60 mmol), $Ni(OAc)_2$ (10 mmol) and 2,2'-bipyridine (20 mmol).^[14]

NiCRA, prepared from $Ni(OAc)_2$, NaH and *t*-AmOH in THF totally desulfurized the thioacetals shown in Scheme 3 with excellent yields (GC). Caubère also demonstrated that a weaker reducing complex (e.g., NiCRA-bpy),^[15] prepared by adding 2,2'-bi-pyridine to NiCRA, was able to open only one of the two C-S bonds by a half-desulfurization process, which was not easy to control with other Ni complexes.^[14a] It should be noted that *t*-AmOH can be replaced by *N*-methyl-3-hydroxypiperidine to obtain the reduced compounds from thioacetals with no loss in yields.^[14c] Moreover, in this study, the authors demonstrated that NiCRA totally desulfurized 3,3-ethylene dithiocholestone (99%) whereas, this reaction gave a mixture of compounds with Ni_2B (prepared from Ni(II) salts and $NaBH_4$, see the next section).^[16] In our opinion, the main problem associated with NiCRAs is their use in large excess for desulfurization reactions (atom ratio Ni/S = 5 to 30).

2.2.2. “ Ni_2B ” from $NiCl_2$ and $NaBH_4$

Ni_2B is a weak reducing agent prepared from the reduction of $NiCl_2$ by $NaBH_4$ to give an easy to handle black non-pyrophoric precipitate. However, Ni_2B can also be used directly in the reaction flask from $NiCl_2$ and $NaBH_4$ in the presence of the sulfur compounds without prior preparation. The advantages of Ni_2B compared to Raney-Ni are easy handling, the possible determination of the Ni amount used, and lower reactivity. Some representative examples are depicted in Scheme 4.

Scheme 4. Reagent and conditions: ^athioacetal (1 equiv), $NiCl_2 \cdot 6H_2O$ (5 to 10 equiv), $NaBH_4$ (10 to 30 mmol) in MeOH-THF, 0 °C to rt.

1,2-Thiolanes,^[17a] 1,3-dithianes^[17b] and *bis*-phenylsulfanes^[17b] were rapidly desulfurized by the *in situ* prepared Ni₂B. Reactions occurred at room temperature in few hours, using a very simple workup to provide methyl and methylene derivatives in good to excellent yields. As observed with NiCRAs, the quantities of Ni₂B are important (atom Ni/atom S from 5 to 10). Some functional groups (*e.g.*, halogen, OMe, ester or lactam) were well tolerated under the reaction conditions. One can note that, due to the low reactivity of Ni₂B, the total desulfurization of thioacetals can be complicated and two successive reduction steps with half-desulfurized compounds as intermediates were needed.^[18]

2.2.3. NiBr₂·DME, PPh₃ and LiAlH₄

Luh and co-workers studied the reduction of sulfoxides and sulfones by nickel complexes and also found that the use of NiBr₂·DME, PPh₃ and LiAlH₄ combination was able to totally desulfurize a dithioacetal and a *bis*-dithioacetal function. In the examples depicted in Scheme 5,^[19] the nickel complex was prepared by mixing NiBr₂·DME with 2 equiv of PPh₃ and 2 equiv of LiAlH₄ in THF at room temperature to give a dark solution in which was syringed thioacetals. Unfortunately, no comparison with Raney-Ni or NiCRAs was described for the desulfurization of fluorenonedithioacetal and *bis*-dithioacetal. It would be also interesting to consider if this Ni-complex is compatible with sensitive functional groups.

Scheme 5. Desulfurization of thioacetals by NiBr₂·DME, PPh₃ and LiAlH₄. Reaction and conditions: ^aNiBr₂, PPh₃ (2 equiv) and LiAlH₄ (2 equiv) in THF at rt were mixed with dithioacetal.

2.2.4. Nickelocene-LiAlH₄ hydride^[20]

Luh also reported that nickelocene reacted with mercaptans to give the corresponding binuclear complexes^[21] in which the carbon-sulfur bond was activated and can be reduced by LiAlH₄.^[22] Keeping this idea in mind, Luh studied the reductive properties of a complex nickelocene-LiAlH₄ for the desulfurization of thiols, sulfoxides, sulfones, and thioethers including three examples of thioacetals (Scheme 6). As it can be observed, by using this homogeneous organonickel reagent, isolated yields in reduced compounds were poor by comparison with other Ni-reducing reagents. It has been showed with other substrates that this nickel complex was compatible with several functional groups (halogen, MeO, CF₃, carbonyl, ester,...) but a NO₂ function was unstable under the reaction conditions.

Scheme 6. Desulfurization of dithiolanes with nickelocene and LiAlH₄. Reaction and conditions: ^aNickelocene (0.5 equiv) was mixed with LiAlH₄ (0.5 equiv) and dithiolanes in THF at rt.

2.3. Other metals

2.3.1. Na and Li in liquid NH₃

The first desulfurization of a 1,2-thiolane using solvated electrons generated from Na in liquid ammonia was published by Ireland and co-workers in 1958.^[23] The authors reported that the ethylene thioketal of cholest-4-en-3-one was efficiently reduced to give the expected cholest-4-ene (89%) with no reduction of the internal double bond (Scheme 7).

Scheme 7. First desulfurization of a thioketal using solvated electrons generated from Na in liquid ammonia.

This protocol was efficiently used few years later by Fochi^[24] and Clive^[25] with 1,3-benzodithiol derivatives. The results of these studies are compacted in Scheme 8.

Scheme 8. Selection of desulfurization of 1,3-benzodithiolanes by solvated electrons generated from Na in liquid ammonia.

The authors indicated that, beside high yields, another advantage was to possibly recover a large amount of the starting benzene-1,2-dithiol.^[24] In liquid ammonia, solvated electrons can also be generated from Li as showed by Zeelen during the total synthesis of dl-19-Nor-4-pregnen-20-one (Scheme 9).^[26]

Scheme 9. Desulfurization of a thioketal by solvated electrons generated from Li in liquid ammonia.

Authors pointed out that a keto group on C-20 was not compatible with this desulfurizing process on the contrary to a secondary alcohol function, which was subsequently oxidized by chromic acid.^[26]

2.3.2. Li in EtNH₂

Crossley and Henbest reported that Li/liquid ammonia can be replaced by Li/EtNH₂ for the electron-mediated desulfurization of thioketals of perhydroanthracenes.^[27a] This protocol produced the methylene compounds in good yields did not affect the stereochemistry of the bridge head positions. The desulfurization process was slow and not selective since double bonds present within the structures were also reduced (Scheme 10). Replacement of Li in EtNH₂ at -20 °C by Raney-Ni did not improve the selectivity and the yields of this process. This protocol was successfully transposed during the synthesis of Lasiol, an olfactory compound with no-reduction of the tri-substituted double bond.^[27b]

Scheme 10. Desulfurization of thioketals by solvated electrons generated from Li in EtNH₂.

2.3.3. LiAlH₄ associated with CuCl₂/ZnCl₂ or with TiCl₄

During his work concerning the reducing properties of CuCl₂ with LiAlH₄, Mukaiyama observed that the yield in diphenylmethane was doubled (84%) when ZnCl₂ was added to the reducing LiAlH₄-CuCl₂ combination^[28a] (Scheme 11). However, 4 equiv of ZnCl₂ and CuCl₂ together with 16 equiv of LiAlH₄ were needed to achieve this transformation, which was not studied with functionalized substrates. Two years after, in 1973, Mukaiyama successfully replaced the couple ZnCl₂/CuCl₂ by TiCl₄ in the presence of LiAlH₄ to desulfurize thioketals.^[28b] Best conditions were found by using 4 equiv of LiAlH₄ with 2 equiv of TiCl₄ to provide the expected methylene products in yields ranging from 80 to 97%.

Scheme 11. LiAlH₄ associated to CuCl₂/ZnCl₂ or TiCl₄.

Unfortunately, these experimental conditions were not reported with functionalized dithioketals in which reducible functional groups would not be probably tolerated.

2.3.4. Miscellaneous metal processes using Mo(CO)₆ and Sml₂

Alper and Blais reported in 1980 that Mo(CO)₆ adsorbed on silica gel was able to reduce dibenzothiophene^[29] into biphenyl. Based on this observation, Luh studied the desulfurization of a series of compounds containing sulfur and only one example of a 1,2-thiolane reduction was reported^[30] (Scheme 12).

Scheme 12. Desulfurization of 9-fluorenone thioketal by Mo(CO)₆. Reaction and conditions: Mo(CO)₆ (1 equiv) was mixed with dibenzothiophene in THF at rt.

However, fluorene was obtained in a poor yield of 42% together with its dimer (34%) which was formed from a molybdenum-carbene complex, making this process unattractive for desulfurization of thioketals.

To close this section dedicated to desulfurization reactions by metals, Hilmersson reported in 2003 that Sml₂/H₂O/*n*BuNH₂ was able to transform a dithioacetal into its methyl derivative^[31] through a single example depicted in Scheme 13.

Scheme 13. Desulfurization of a dithioacetal by Sml₂/H₂O/*n*BuNH₂. Reaction and conditions: Sml₂ (2.5 equiv), *n*BuNH₂ (7.5 equiv) and H₂O (12.5 equiv) were mixed with dithioacetal at rt.

After 30 seconds of reaction, Sml₂ associated to H₂O and *n*BuNH₂ promoted the intramolecular coupling reaction but also the hydrogenolysis of one of the carbon-sulfur bonds. By adding an excess of Sml₂ and after a prolonged reaction time (between 1 to 2 h), a second hydrogenolysis reaction

occurred to provide the methyl compound (85%). As it can be observed in this part dedicated to the desulfurization by other metals, protocols using Na and Li in liquid ammonia or amines are possible alternatives to the use of nickel reducing reagents for desulfurization reactions of dithioketals and dithioacetals but these methodologies probably cannot be applied to highly functionalized or fragile molecules.

3. Other desulfurizing reagents

3.1. Metal hydrides

3.1.1. Bu_3SnH with AIBN

Gutierrez found in 1980, that in the presence of a radical initiator, Bu_3SnH was an effective reducing agent for complete or partial desulfurization of dithioketals and dithioacetals.^[32] In the presence of 4 equiv of Bu_3SnH and AIBN, 1,3-dithiolanes were totally reduced to give their methylene derivatives in good yields (Scheme 14).

Scheme 14. Desulfurization of dithiolanes and dithianes with Bu_3SnH .

1,3-Dithiolanes were half desulfurized using 1 equiv of Bu_3SnH to give quantitatively tri-*n*-butyltin sulfide products, which were destannylated on silica gel column. Gutierrez also demonstrated that 1,3-dithianes were easily desulfurized requiring only 2 equiv of Bu_3SnH to give the totally reduced compounds. This reducing process was successfully used by Zhai during the total synthesis of aphanorphine,^[33] and also by Nagamitsu during the synthesis of Citridone A^[34] (Scheme 15).

Scheme 15. Desulfurization of dithiolanes by Bu_3SnH in total syntheses.

3.1.2. $\text{B}(\text{C}_6\text{F}_5)_3$ catalyzed desulfurization of thioketals in the presence of Et_3SiH

During a study dedicated to the reducing properties of hydrosilanes towards sulfides in the presence of a catalytic amount of $\text{B}(\text{C}_6\text{F}_5)_3$, Akiyama observed that this combination was able to totally desulfurize 2-naphtylthioketals in nearly quantitative yields (Scheme 16).^[35] Et_3SiH (3 to 6 equiv) which was preferable to Ph_3SiH or $(\text{EtO})_3\text{SiH}$ reacted with thioketals and 2 mol% of $\text{B}(\text{C}_6\text{F}_5)_3$ in CHCl_3 to provide the totally reduced compounds.

Scheme 16. Desulfurization and half desulfurization of thioketals using Et_3SiH in the presence of $\text{B}(\text{C}_6\text{F}_5)_3$.

It is interesting to note that dithiolane and dithiane derived from 2-acetyl naphthalene were totally desulfurized to give their methylene derivatives, whereas dithiane and dithiolane prepared from 2-naphthalenecarbaldehyde were quantitatively half-desulfurized under the same experimental conditions, indicating that a methyl group close to the sulfur atom facilitated its coordination to the boron atom and then permitted the second reduction process.

3.2. Metal-free desulfurizations

3.2.1. Hydrated hydrazine

Giorgian and co-workers reported first in 1959 that hydrated hydrazine was able to desulfurize cyclic or acyclic thioketals in the presence of KOH.^[36] They also showed that KOH was useful but not essential in this reducing process but lowered considerably the effective temperature of reactions. Some representative examples are reported in Scheme 17.

Scheme 17. Desulfurization of thioketals by NH_2NH_2 and KOH.

Five years after the pioneer work of Giorgian,^[36] Corey described the total synthesis of Longifolene and studied the desulfurization of the dithiolane described in Scheme 18.^[37] The authors reported that Raney-Ni was ineffective in achieving this transformation, on the contrary to hydrazine

associated with Na, which quantitatively desulfurized the dithiolane into the expected methylene compound.

Scheme 18. Desulfurization of a dithioketal by NH_2NH_2 and Na.

3.2.2. P_2I_4 in CH_2Cl_2

Shigemasa studied the desulfurization of dithiolanes by P_2I_4 and observed the fundamental role of solvents in these reactions.^[38] In Ac_2O , dithiolanes were deprotected by P_2I_4 (1 equiv) to provide their parent ketones, whereas in methylene chloride and water, methylene compounds were obtained as majors products (Scheme 19).

Scheme 19. Reduction of dithiolanes by P_2I_4 in CH_2Cl_2 .

3.2.3. Photoreduction of dithianes and dithiolanes with $t\text{BuOK}$ in DMSO

Recently, an Argentinian group reported an elegant desulfurization process of dithioketals using a photoinduced electron transfer from $t\text{BuOK}$ in the presence of 1,4-cyclohexadiene as hydrogen atom donor (Scheme 20).^[39]

Scheme 20. Photoreduction of dithianes and dithiolanes with $t\text{BuOK}$ in DMSO.

The procedure seems to be compatible with both EDGs and EWGs on aromatic rings (e.g. CN, 83%). The mechanism of the photochemical reduction of thioketals in DMSO was well investigated by the authors and others^[40] through calculations of quantum yield for products and for the initiation step.

3.2.4. EtSNa in MeOH

The last example described in this review deals with the desulfurization of various 9,9-bis(ethylthio)fluorenes by EtSNa in MeOH (Scheme 21).^[41] Effects of neighboring ester groups (on *o*, *m* and *p*-positions of the aromatic ring) on the reaction rate are discussed and compared. For the derivative having an *ortho*-ester function close to the sulfur atoms, steric hindrance and stabilization of the benzylic anion formed after the first C-S bond cleavage lowered the rate of the process.

Scheme 21. Sodium ethanethiolate promoted the desulfurization of 9,9-bis(ethylthio)fluorenes.

4. Conclusion

The reducing desulfurization processes of thioketals into methylene or methyl derivatives were reviewed from the pioneer's works of Wolform and Karabinos until 2017. Raney-Ni is the most used reagent to achieve this transformation and was used in many total syntheses, demonstrating its efficacy and a certain functional group tolerance. However, (i) the pyrophoric properties of Raney-Ni, (ii) the hazardous manipulations and difficulty to calculate the exact quantity of Ni in experiments, (iii) the presence or not of H_2 in catalysts, (iv) the toxicity associated with the use of Ni and Ni-derivatives,^[42] and (v) the use of Ni reducing reagent in more than stoichiometric amounts, limit the use from an industrial point of view and for large scale preparations. Other metals or metal salts can successfully replace Ni and Ni salts as Na and Li in NH_3 or amines, Mo or ZnCl_2 , CuCl_2 and TiCl_4 . However, for these metals and metal salts, no complete studies were achieved to examine the scope and limitation of such methods. A similar comment can be applied to all other methodologies presented in this review, such as the use of metallic hydrides (Bu_3SnH and Et_3SiH) as well as metal-free processes and particularly the photochemical reduction of dithioketals, which is a promising method if applicable in highly functionalized molecules. There is an urgent need to find a soft metal-free methodology to desulfurize dithioketals, which is easy to handle, tolerant towards functional groups and provides the expected reduced products in high yields.

Acknowledgments

Authors gratefully acknowledge support of this project by CNRS, Univ. Paris-Saclay, and by La Ligue Nationale Contre le Cancer through an Equipe Labellisée 2014 grant. Guangkuan Zhao and Ling-Zhi Yuan thank the CSC (Chinese Scholarship Council) for a Ph.D. funding. Our laboratory (BioCIS UMR 8076) is a member of the laboratory of excellence LERMIT supported by a grant from ANR (ANR-10-LABX-33).

Conflict of Interest

The authors declare no conflict of interest.

Keywords: desulfurization · Ni derivatives · reduction · thioacetal · thioketal

- [1] a) J. Bougault, E. Cattelain, P. Chabrier, *Bull. Chem. Soc. Chim.* **1938**, 5, 1699-1712; b) L. Wolfrom, J. V. Karabinos, *J. Am. Chem. Soc.* **1944**, 66, 909-911.
- [2] G. R. Pettit, E. E. van Tamelen, *Org. React.* **1962**, 12, 356-529.
- [3] Inexpensive Raney-Ni is commercially available.
- [4] a) M. Raney, U.S. Patent 1,563,587 (Dec. 1, 1925); *Chem. Abstracts* **1926**, 20, 515; b) M. Raney, U.S. Patent 1,628,190 (May 10, 1927); *Chem. Abstracts* **1927**, 21, 2116; c) M. Raney, U.S. Patent 1,915,473 (Jun. 27, 1933); *Chem. Abstracts* **1933**, 27, 4361.
- [5] a) R. Mazingo, D. E. Wolf, S. A. Harris, K. Folkers, *J. Am. Chem. Soc.* **1943**, 65, 1013-1016; b) R. Mazingo, *Org. Syntheses Collective Vol. 3*, 181 (1955); c) P. Ruggli, E. Preiswerk, *Helv. Chim. Acta* **1939**, 22, 478-495.
- [6] E. Eich, H. Pertz, M. Kaloga, J. Schulz, M. R. Fesen, A. Mazumder, Y. Pommier, *J. Med. Chem.* **1996**, 39, 86-95.
- [7] Y. Unoh, K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2015**, 17, 704-707.
- [8] N. Chandrasoma, N. Brown, A. Brassfield, A. Nerurkar, S. Suarez, K. R. Buszek, *Tetrahedron Lett.* **2013**, 54, 913-917.
- [9] A. L. Hurski, M. V. Barysevich, T. S. Dalidovich, M. V. Irkryk, N. U. Kolasava, V. N. Zhabinskii, V. A. Khripach, *Chem. Eur. J.* **2016**, 22, 14171-14174.
- [10] For an excellent review on reductive desulfurization of complex molecules by Raney-Ni please see: a) J. Rentner, M. Kljajic, L. Offner, R. Breinbauer, *Tetrahedron* **2014**, 70, 8983-9027; b) S. H. Yu, W. Zhu, D. W. Ma, *J. Org. Chem.* **2005**, 70, 7364-7370; c) M. De Paolis, H. Rosso, M. Henrot, C. Prandi, F. d'Herouville, J. Maddaluno, *Chem. Eur. J.* **2010**, 16, 11229-11232; d) H. Suginome, S. Yamada, J. B. Wang, *J. Org. Chem.* **1990**, 55, 2170-2176; e) J. Shiina, M. Oikawa, K. Nakamura, R. Obata, S. Nishiyama, *Eur. J. Org. Chem.* **2007**, 5190-5197; f) E. Eich, H. Pertz, M. Kaloga, J. Schulz, M. R. Fesen, A. Mazumder, Y. Pommier, *J. Med. Chem.* **1996**, 39, 86-95. g) N. Rehnberg, G. Magnusson, *J. Org. Chem.* **1990**, 55, 4340-4349; h) C. Liu, Q. Yin, L. X. Dai, S. L. You, *Chem. Commun.* **2015**, 51, 5971-5974; i) J. M. Ren, P. Zhao, X. Xiao, T. Q. Chen, B. B. Zeng, *Synthesis* **2016**, 48, 4161-4166; j) S. A. Snyder, E. J. Corey, *J. Am. Chem. Soc.* **2006**, 128, 740-742; k) Y. Q. Feng, M. M. Majireck, S. M. Weinreb, *J. Org. Chem.* **2014**, 79, 7-24; l) A. A. Momin, T. D. Urmode, S. M. Bhosale, R. S. Kusurkar, *Synth. Commun.* **2016**, 46, 1292-1298.
- [11] H. Hauptmann, W. F. Walter, *Chem. Rev.* **1962**, 62, 347-404.
- [12] a) L. Mordenti, J. J. Brunet, P. Caubère, *J. Org. Chem.* **1979**, 44, 2203-2205; b) J. J. Brunet, P. Gallois, P. Caubère, *J. Org. Chem.* **1980**, 45, 1937-1945; c) P. Gallois, J. J. Brunet, P. Caubère, *J. Org. Chem.* **1980**, 45, 1946-1950; d) R. Vanderesse, J. J. Brunet, P. Caubère, *J. Org. Chem.* **1981**, 46, 1270-1277; e) J. J. Brunet, D. Besozzi, A. Courtois, P. Caubère, *J. Am. Chem. Soc.* **1982**, 104, 7130-7135; f) Y. Fort, R. Vanderesse, P. Caubère, *Tetrahedron Lett.* **1985**, 26, 3111-3114; g) Y. Fort, R. Vanderesse, P. Caubère, *Tetrahedron Lett.* **1986**, 27, 5487-5490; h) G. Feghouli, R. Vanderesse, Y. Fort, P. Caubère, *Tetrahedron Lett.* **1988**, 29, 1383-1384.
- [13] a) R. Vanderesse, J. J. Brunet, P. Caubère, *J. Organomet. Chem.* **1984**, 264, 263-271; b) R. Vanderesse, Y. Fort, S. Becker, P. Caubère, *Tetrahedron Lett.* **1986**, 27, 3517-3520; c) R. Vanderesse, M. Lourak, Y. Fort, P. Caubère, *Tetrahedron Lett.* **1986**, 27, 5483-5486; d) M. Lourak, R. Vanderesse, Y. Fort, P. Caubère, *Tetrahedron Lett.* **1988**, 29, 545-548; e) M. Lourak, R. Vanderesse, Y. Fort, P. Caubère, *J. Org. Chem.* **1989**, 54, 4840-4844; f) M. Lourak, R. Vanderesse, Y. Fort, P. Caubère, *J. Org. Chem.* **1989**, 54, 4844-4848.
- [14] a) S. Becker, Y. Fort, R. Vanderesse, P. Caubère, *Tetrahedron Lett.* **1988**, 29, 2963-2966; b) S. Becker, Y. Fort, P. Caubère, *J. Org. Chem.* **1990**, 55, 6194-6198; c) C. Kuehm-Caubère, A. Guilmar, S. Adach-Becker, Y. Fort, P. Caubère, *Tetrahedron Lett.* **1998**, 39, 8987-8990; d) half desulfurization of thioketals was also achieved using Ca in ammonia see: B. C. Newman, E. L. Eliel, *J. Org. Chem.* **1970**, 35, 3641-3646.
- [15] P. Caubère, *Angew. Chem. Int. Ed. Engl.* **1983**, 22, 599-613.
- [16] R. R. Boar, D. W. Hawkins, J. F. McGhie, D. H. R. Barton, *J. Chem. Soc. Perkin Trans. 1* **1973**, 654-657.
- [17] a) J. M. Khurana, D. Magoo, *Synth. Commun.* **2010**, 40, 2908-2913; b) M. Amat, M. Pérez, N. Llor, C. Escolano, F. J. Luque, E. Molins, J. Bosch, *J. Org. Chem.* **2004**, 69, 8681-8693.
- [18] W. E. Truce, F. E. Roberts, *J. Org. Chem.* **1963**, 28, 961-964.
- [19] K. M. Ho, C. H. Lam, T. Y. Luh, *J. Org. Chem.* **1989**, 54, 4474-4476.
- [20] M. C. Chan, K. M. Cheng, K. M. Ho, C. T. Ng, T. M. Yam, B. S. L. Wang, T. Y. Luh, *J. Org. Chem.* **1988**, 53, 4466-4471.
- [21] a) W. K. Schroop, *J. Inorg. Nucl. Chem.* **1962**, 24, 1688-1690. b) P. C. Ellgen, C. D. Gregory, *Inorg. Chem.* **1971**, 10, 980-984.
- [22] N. F. Ho, T. C. W. Mak, T. Y. Luh, *Organomet. Chem.* **1986**, 317, C28-C30.
- [23] R. E. Ireland, T. I. Wrigley, W. G. Young, *J. Am. Chem. Soc.* **1958**, 80, 4604-4606.
- [24] I. Degani, R. Fochi, *J. Chem. Soc. Perkin Trans. 1* **1978**, 1133-1136.
- [25] J. S. Prasad, D. L. J. Clive, G. V. J. da Silva, *J. Org. Chem.* **1986**, 51, 2717-2721.
- [26] J. A. M. Peters, T. A. P. Posthumus, N. P. van Vliet, F. J. Zeelen, *J. Org. Chem.* **1980**, 45, 2208-2214.
- [27] a) N. S. Crossley, H. B. Henbest, *J. Chem. Soc.* **1960**, 4413-441; b) S. Kuwahara, Shibata, A. Hiramatsu, *Liebigs Ann. Chem.* **1992**, 993-995.
- [28] a) T. Mukaiyama, K. Narasaka, K. Maekawa, M. Furusato, *Bull. Chem. Soc. Jpn* **1971**, 44, 2285; b) T. Mukaiyama, M. Hayashi, K. Narasaka, *Chem. Lett.* **1973**, 291-293.
- [29] H. Alper, C. Blais, *Fuel*, **1980**, 59, 670.
- [30] T. Y. Luh, C. S. Wong, *J. Org. Chem.* **1985**, 50, 5413-5415.
- [31] A. Dahlén, A. Petersson, G. Hillmersson, *Org. Biomol. Chem.* **2003**, 1, 2423-2426.
- [32] a) C. G. Gutierrez, R. A. Stringham, T. Nitasaka, K. G. Glasscock, *J. Org. Chem.* **1980**, 45, 3393-3395; b) K. Schmidt, S. O'Neal, T. C. Chan, C. P. Alexis, J. M. Uribe, K. Lossener, C. G. Gutierrez, *Tetrahedron Lett.* **1989**, 30, 7301-7304; c) C. G. Gutierrez, C. P. Alexis, J. M. Uribe, *J. Chem. Soc., Chem. Commun.* **1984**, 124-125.
- [33] Z. Ma, H. Hu, W. Xiong, H. Zhai, *Tetrahedron* **2007**, 63, 7523-7531.
- [34] T. Miyagawa, K. Nagai, A. Yamada, Y. Sugihara, T. Fukuda, R. Uchida, H. Tomoda, S. Omura, T. Nagamitsu, *Org. Lett.* **2011**, 13, 1158-1161.
- [35] K. Saito, K. Kondo, T. Akiyama, *Org. Lett.* **2015**, 17, 3366-3369.
- [36] V. Georgian, R. Harrisson, N. Gubisch, *J. Am. Chem. Soc.* **1959**, 81, 5834-5835.
- [37] E. J. Corey, M. Ohno, R. B. Mitra, P. A. Vatakencherry, *J. Am. Chem. Soc.* **1964**, 86, 478-485.
- [38] Y. Shigemasa, M. Ogawa, H. Sashiwa, H. Saimoto, *Tetrahedron Lett.* **1989**, 30, 1277-1278.
- [39] G. Oksdath-Mansilla, J. Argüello, A. B. Pénenory, *Tetrahedron Lett.* **2013**, 54, 1515-1518.
- [40] G. Goulthard, K. J. Emery, E. Doni, F. Cumine, G. Nocera, M. P. John, L. E. A. Berlouis, T. McGuire, T. Tuttle, J. A. Murphy, *J. Am. Chem. Soc.* **2016**, 138, 7402-7410.
- [41] W. Nakanishi, H. Nakanishi, Y. Yanagawa, Y. Ikeda, M. Oki, *Chem. Lett.* **1983**, 105-108.
- [42] For a review on toxicity and carcinogenicity of nickel compounds see: T. P. Coogan, D. M. Latta, E. T. Snow, M. Costa, *Crit. Rev. Toxicol.* **1989**, 19, 341-384.

Entry for the Table of Contents

Insert graphic for Table of Contents here. ((Please ensure your graphic is in **one** of following formats))

This review reports the total desulfurization of thioacetals and thioketals, leading to methyl and methylene derivatives, through metals, metal-hydrides and metal-free methodologies used between 1944 until 2017. Applications to functionalized molecules in a variety of complex or total syntheses is also presented.