One-pot hydrosilylation–protodesilylation of functionalized diarylalkynes: a highly selective access to Z-stilbenes. Application to the synthesis of combretastatin A-4

Anne Giraud, Olivier Provot, Abdallah Hamze, Jean-Daniel Brion, Mouad Alami

► To cite this version:

Anne Giraud, Olivier Provot, Abdallah Hamze, Jean-Daniel Brion, Mouad Alami. One-pot hydrosilylation–protodesilylation of functionalized diarylalkynes: a highly selective access to Z-stilbenes. Application to the synthesis of combretastatin A-4. *Tetrahedron Letters*, 2008, 49 (7), pp.1107-1110. 10.1016/j.tetlet.2007.12.057 . hal-02394604

HAL Id: hal-02394604

<https://hal.science/hal-02394604>

Submitted on 4 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

One-pot hydrosilylation-protodesilylation of functionalized diarylalkynes: a highly selective access to *Z*-stilbenes. Application to the synthesis of combretastatin A-4.

Anne Giraud¹, Olivier Provot,^{1,2*} Abdallah Hamzé^{1,2}, Jean-Daniel Brion^{1,2} and Mouâd Alami^{2,1*}

¹ Univ Paris-Sud, BioCIS, UMR 8076, Laboratoire de Chimie Thérapeutique ; ² CNRS, BioCIS, UMR 8076, Laboratoire de Chimie Thérapeutique, Faculté de Pharmacie, rue J.B. Clément, Châtenay-Malabry, F-92296, France

Abstract— An efficient stereoselective synthesis of *Z*-stilbenes has been developed from diarylalkynes *via* a new hydrosilylation-protodesilylation process. Scope and limitation of this method to prepare stereoselectively a wide range of (*Z*)-stilbenes in a one-pot way is presented. A concise application to the preparation of combretastatin A-4 (**CA-4**), a vascular targeting agent inhibitor of tubulin polymerisation is described.

A large number of stilbene (1,2-diphenylethylene) derivatives have been isolated from various plant species and exhibit a large panel of biological activities such as, antineoplastic, antiangiogenesis, cytotoxic and inhibitory of cell proliferation.¹ Beside *trans* stilbene derivatives and one of their leader, Resveratrol, a large number of their *cis* homologues were isolated, synthesized and evaluated. The most promising of these stilbenes thus far are combretastatins and particularly, the combretastatin A-4 (**CA-4**), due to its biological activities combined to its structural simplicity (Scheme 1).²

Scheme 1

If the access of *trans* stilbenes is well-documented,³ there are only few methodologies that give access to their *cis* counterparts. The most popular approach is based on the *Z* selective-Lindlar⁴ semi reduction of diarylalkynes but suffer from several drawbacks as (*Z*) to (*E*) isomerisation, over reduction to the alkane making purifications tedious and problems with reproducibility.

As a part of our research devoted to anticancer agents,⁵ that target tubulin,⁶ we wished to synthesize the natural product CA-4 as a reference molecule since, to our

knowledge; CA-4 is not commercially available anymore. The chemical synthesis of CA-4 has been tackled in a variety of ways.^{1b,7} while these reactions are suitable methods, many of them either display low stereoselectivity or do not tolerate sensitive functionalities. Therefore, alternative routes for the synthesis of CA-4 are welcome. A recent work describing the synthesis of CA-4 and analogues by hydrolysis of Ti(II)-alkynes complexes,⁸ prompted us to publish our preliminary results in the *Z*-semi reduction of alkynes field.

Recently, we showed that heterogeneous platinum oxide (PtO₂) is a very efficient catalyst for the hydrosilylation of *para* and *ortho* substituted diarylalkynes. The H–Si bond addition proceeds in a stereoselective *cis*-fashion⁹ and the regioselectivity of the reaction was found to be governed by *ortho*-directing effects (ODE)¹⁰ rather than the nature of the platinum catalysts. We reasoned that this efficient hydrosilylation of internal arylalkynes could be efficiently exploited to give, after linkage of the C–Si bond, *Z*-stilbenes of high biological interest. We present in this communication a convenient one-pot procedure for the *Z*-selective semi reduction of diarylalkynes *via* a hydrosilylation-protodesilylation sequence.

To define optimal reaction conditions, we have studied initially the hydrosilylation of diarylalkynes **1a** and **1b** as

*Corresponding authors. olivier.provot@u-psud.fr and/or mouad.alami@u-psud.fr

Keywords : combretastatin A4, (*Z*)-stilbenes, hydrosilylation, desilylation, platinum

Table 1. Hydrosilylation-protodesilylation sequence; access to Z-stilbenes **3**

entry	diarylalkyne 1	H-[Si]	catalyst	yield ^a of 2 ^b (%)	yield ^a of 3 (%)	Z/E ratio ^c
1	1a R = 4-OMe, R ¹ = H	1 : HSiEt ₃	PtO ₂	2a-1 97	3a 58	85/15
2	"	2 : HSiPh ₂ Me	"	2a-2 93	" 49	100/0
3	"	3 : HSi(OEt) ₃	"	2a-3 70	" 80	100/0
4	"	4 : HSiOEtMe ₂	PtO ₂	2a-4 86	" 90	100/0
5	"	"	PtCl ₂	2a-4 44	" nd ^d	nd ^d
6	"	"	Pt/C	2a-4 38	" -	-
7	"	"	PtCl ₄	2a-4 56	" -	-
8	"	"	H ₂ PtCl ₆	2a-4 mixture	" -	-
9	1b R = 2,3,4-OMe, R ¹ = H	4 : HSiOEtMe ₂	PtO ₂	2b-4 87	3b 77	98/2
10	"	3 : HSi(OEt) ₃	"	2b-3 78	84	98/2
11	"	4 : HSiOEtMe ₂	"	2b-4 87	88 ^e	100/0
12	1c R = 2,3,4-OMe, R ¹ = 3-OH, 4-OMe	"	"	2c-4 70 ^f	CA-4 69	100/0

^a Isolated yield.^b Mixture of regioisomers (ratio not determined).^c Z:E Ratios quoted as a percentage composition of total yield as estimated from ¹H NMR spectra.^d Not determined.^e Protodesilylation was achieved at RT.^f 4 eq of HSiOEtMe₂ were required.

model substrates and their subsequent desilylation. The results are summarized in Table 1. Preliminary experiment was carried out using HSiEt₃ (1.5 eq), PtO₂ (7 mol%) at 60 °C in the absence of solvent as we previously reported.⁹ In this case, a mixture of vinylsilanes was obtained with a quantitative yield after purification on silica gel. However, the desilylation of the vinylsilanes mixture using TBAF (1.5 eq) at 60 °C afforded a moderate yield of Z-**3a** together with significant amounts of E-**3a** (entry 1). A survey of the sequence hydrosilylation-protodesilylation sequence of alkyne **1a** under the same conditions in the presence of other representative silanes, including HSiPh₂Me, HSi(OEt)₃ and HSiOEtMe₂ was conducted (entries 2-4). In these cases, we were pleased to observe the formation of a single Z-**3a** stereoisomer whatever the nature of the silane used and the better overall yield was obtained with the non-toxic¹¹ HSiOEtMe₂ (77% for the two steps, entry 4). Encouraged by this result, the effect of PtCl₄, PtCl₂ and Pt/C on the outcome of the hydrosilylation reaction was investigated. None of these platinum catalysts led to better results than those achieved with PtO₂ (entries 5-7). One can note that H₂PtCl₆ (Speier's catalyst), which is considered to be the catalyst of choice for *cis*-

hydrosilylation of internal alkynes¹² was not efficient and led to a complex mixture (entry 7). Having defined the best conditions [PtO₂ (7 mol%), HSiOEtMe₂ (1.5 eq), 60 °C then TBAF (1.5 eq), 0 °C] Z-stilbene **3b** bearing a 3,4,5-trimethoxyaryl moiety was prepared and isolated with good yield (entry 9). Careful inspection of the ¹H NMR spectrum of the crude reaction mixture indicated that the stilbene **3b** was formed in a 98:2/Z:E ratio. A similar result was obtained when using HSi(OEt)₃ instead of HSiOEtMe₂ (entry 10). Finally, the best result in terms of yield and Z-stereoselectivity was observed when the desilylation step of the vinylsilane **2b-4** was achieved at room temperature (entry 11). In this case, Z-stilbene **3b** was obtained as a single Z stereoisomer in a 77% overall isolated yield (based on alkyne **1b**). We then applied this two stages process to prepare the target natural product **CA-4**¹³ from diarylalkyne **1c**. We were delighted to observe that the hydrosilylation-protodesilylation proceeded extremely efficiently to give the desired **CA-4** as a single Z-isomer with good yield (entry 12).

From a practical point of view of the synthetic chemist, we explored the process in a one-pot manner. Thus, after the

hydrosilylation step, the excess of the volatile HSiOEtMe_2 was removed to achieve the protodesilylation step with TBAF (3 eq) at 0 °C for a better *Z*-stereoselectivity. Scope and limitations of this one-pot process using various functionalized diarylalkynes **1** are summarized in Table 2.

Table 2. One-pot access to *Z*-stilbenes **3** from diarylalkynes **1**

entry	alkynes	stilbenes	<i>Z/E</i> ratio ^a	yield ^b
1	1a		3a 100/0	77
2	1d		3d 100/0	90
3	1e		3e 100/0	75
4	1f		3f 100/0	51
5	1g		3g 100/0	49 ^c
6	1h		3h 100/0	65
7	1i		3i 100/0	79
8	1j		3j >95/5	51
9	1k		3k 100/0	95
10	1l		3l >95/5	62
11	1m		3m >95/5	75
12	1n		3n 90/10	73
13	1o		3o 88/12	78
14	1c		CA-4 90/10	67

^a *Z/E* Ratios quoted as a percentage composition of total yield as estimated from ^1H NMR spectra.

^b Isolated yield.

^c Unoptimized yield.

Diarylalkynes **1a**, **1d-f** substituted with an electron donating group (OMe, Me) have been transformed successfully to their corresponding stilbenes **3** in good

yields (entries 1-4). It should be noted that the position of the substituent on the aromatic ring had no influence on the *Z*-stereochemistry of the stilbenes formed (entries 1-3). A similar result (total *Z*-stereocontrol) has been obtained from **1g** bearing a pivaloyl ester function but with a moderate unoptimized 49% yield (entry 5). We were also pleased to observe that halogenated substituted arylalkynes **1h** and **1i** afforded the expected stilbenes **3h** and **3i** respectively with acceptable yields and again as single *Z*-stereoisomers (entries 6 and 7). When the push-pull diarylalkynes **1j** was employed as a substrate, it was found that the *Z*-stereochemistry of this one-pot process was slightly reduced. In this case, traces of the *E*-stilbene were detected by ^1H NMR spectroscopy. This PtO_2 -catalyzed hydrosilylation-protodesilylation one-pot process can be performed also with alkyne **1k** bearing a methoxycarbonyl group as illustrated in entry 9. In this case again a total *Z*-stereochemistry associated with a nearly quantitative yield was observed (95%).

We next embarked on the utilization of this process for the conversion of 3,4,5-trimethoxyarylalkynes to their corresponding *Z*-stilbenes, analogues of CA-4. Starting from 3,4,5-trimethoxyarylalkynes **1l** and **1m** bearing an ortho or a para methoxy group respectively, it was noted that the 3,4,5 trimethoxyaryl unit smoothly affected the *Z*-stereoselectivity (> 95%, entries 10 and 11) providing **3l** and **3m**¹⁴ in good yields. Similarly, we succeeded in the one-pot preparation of known biologically active analogues of CA-4, **3n**¹⁵ and **3o**¹⁶ within good yields and high *Z*-stereocontrol (entries 12 and 13).

On account of the large scope of this one-pot procedure, we then attempted to synthesize CA-4 from the corresponding **1c** having a free phenolic function. Because of the side silylation of the phenol moiety, the desilylation step was conducted at 60 °C. As showed in entry 14, CA-4 was obtained with fortunately an excellent *Z*-stereoselectivity (*Z:E*/90:10) and a good yield. Careful separation on silica gel afforded pure CA-4 which could be used as reference for biological evaluation of analogues that would be reported in due course.

In summary, we have developed a new mild and efficient method for the synthesis of *Z*-stilbenes from diarylalkynes. This method is complementary to the existing procedures and sometimes could be the method of choice because of its chemoselectivity, simplicity and its excellent *Z*-stereoselectivity. The ease of which several *Z*-stilbenes were obtained with a total *Z*-stereocontrol and good yields encouraged us to prepare efficiently by this way the natural stilbene CA-4. Moreover, we demonstrated too that this hydrosilylation-protodesilylation sequence could be achieved in a one-pot way from various diarylalkynes.

Acknowledgments

The CNRS is gratefully acknowledged for financial support of this research and the MNSER for a doctoral fellowship to A.G.

References

- ¹ (a) Jang, M.; Cai, L.; Udeani, G. O.; Slowing, K. V.; Thomas, C. F.; Beecher, C. W. W.; Fong, H. H. S.; Farnworth, R. N.; Kinghorn, A. D.; Metha, R. G.; Moon, R. C.; Pezzuto, J. M. *Science* **1997**, 275, 218. (b) Pettit, G. R.; Rhodes, M. R.; Herald, D. L.; Hamel, E.; Schmidt, J. M.; Pettit, R. K. *J. Med. Chem.* **2005**, 48, 4087 and references cited therein.
- ² (a) Pettit, G. R.; Cragg, G. M.; Singh, S. B. *J. Nat. Prod.* **1987**, 50, 386. (b) Pettit, G. R.; Singh, S. B.; Hamel, E.; Lin, C. M.; Alberts, D. S.; Garcia-Kendall, D.; *Experientia* **1989**, 45, 209. (c) Pettit, G. R.; Singh, S. B.; Boyd, M. R.; Hamel, E.; Pettit, R. K.; Schmidt, J. M.; Hogan, F. *J. Med. Chem.* **1995**, 38, 1666.
- ³ Ferré-Filmon, K.; Delaude, L.; Demonceau, A.; Noels, A. F. *Coord. Chem. Rev.* **2004**, 248, 2323.
- ⁴ (a) Lindlar, H. *Helv. Chim. Acta* **1952**, 35, 446. (b) Lindlar, H.; Dubuis, R. *Org. Synth.* **1973**, 5, 880.
- ⁵ Le Bras, G.; Radanyi, C.; Peyrat, J.-F.; Brion, J.-D.; Alami, M.; Marsaud, V.; Stella, B.; Renoir, J.-M. *J. Med. Chem.* **2007**, 50, <http://dx.doi.org/10.1021/jm0707774>.
- ⁶ Provot, O.; Giraud, A.; Peyrat, J.-F.; Alami, M.; Brion, J.-D. *Tetrahedron Lett.* **2005**, 46, 8547.
- ⁷ (a) Robinson, J. E.; Taylor, R. J. K. *Chem. Commun.* **2007**, 1617. (b) Harrowven, D. C.; Guy, I. L.; Howell, M.; Packham, G. *Synlett* **2006**, 2977. (c) Gaukroger, K.; Hadfield, J. A.; Hepworth, L. A.; Lawrence, N. J.; McGown, A. T. *J. Org. Chem.* **2001**, 66, 8135. (d) Lawrence, N. J.; Abdul Ghani, F.; Hepworth, L. A.; Hadfield, J. A.; McGown, A. T.; Pritchard, R. G. *Synthesis* **1999**, 1656. (e) Fürstner, A.; Nikolakis, K. *Liebigs, Ann.* **1996**, 2107.
- ⁸ Lara-Ochoa, F.; Espinosa-Pérez, G. *Tetrahedron Lett.* **2007**, 48, 7007.
- ⁹ Hamze, A.; Provot, O.; Brion, J.-D.; Alami, M. *Synthesis*, **2007**, 2025. (b) Hamze, A.; Provot, O.; Alami, M.; Brion, J.-D. *Org. Lett.* **2005**, 7, 5625.
- ¹⁰ (a) Liron, F.; Le Garrec, P.; Alami, M. *Synlett* **1999**, 246. (b) Alami, M.; Liron, F.; Gervais, M.; Peyrat, J.-F.; Brion, J.-D. *Angew. Chem. Int. Ed.* **2002**, 41, 1578. (c) Liron, F.; Gervais, M.; Peyrat, J.-F.; Alami, M.; Brion, J.-D. *Tetrahedron Lett.* **2003**, 44, 2789. (d) Bujard, M.; Ferri, F.; Alami, M. *Tetrahedron Lett.* **1998**, 39, 4243. (e) Hamze, A.; Provot, O.; Brion, J.-D.; Alami, M. *J. Org. Chem.* **2007**, 72, 3868.
- ¹¹ Dodd, D. E.; Stuart, B. O.; Rothenberg, S. J.; Kershaw, M.; Mann, P. C.; James, J. T.; Lam, C. W. *Inhal. Toxicol.* **1994**, 6, 151.
- ¹² Speier's catalyst is well-known to provide *cis*-addition processes for internal and terminal alkynes; see: Tsipis, C. A. *J. Organomet. Chem.* **1980**, 187, 427.
- ¹³ Procedure for the synthesis of **CA-4**: In a 10 mL flask, PtO₂ (10 mg, 0.0035 mmol) and arylalkyne **1c** (0.5 mmol) were placed under nitrogen atmosphere. Dimethylethoxysilane (304 μ L, 2 mmol) was introduced via syringe and the mixture was stirred at 60 °C in an oil bath for 1 h. The residue was concentrated and then purified by column chromatography over silica gel to yield the vinylsilanes as a mixture of regioisomers (147 mg; 70%). The mixture of vinylsilanes (147 mg, 0.35 mmol) was treated with TBAF in THF (1 mL, 1 N) at 60 °C for 1 h. After concentration in *vacuo*, the crude product was purified by column chromatography (SiO₂, Cyclohexane/ EtOAc : 70/30) to afford **CA-4** as a single *Z*-stereoisomer (75 mg; 69%). TLC: R_f 0.46 (Cyclohexane/EtOAc : 70/30, 1/1, SiO₂). ¹H NMR (300 MHz, CDCl₃) δ 3.70 (s, 6H), 3.84 (s, 3H), 3.86 (s, 3H), 5.50 (brs, 1H, OH), 6.40 (d, 1H, *J* = 12.1 Hz), 6.47 (d, 1H, *J* = 12.1 Hz), 6.53 (s, 2H), 6.72 (d, 1H, *J* = 8.2 Hz), 6.80 (dd, 1H, *J* = 8.2 Hz, *J* = 1.8 Hz), 6.92 (d, 1H, *J* = 1.8 Hz). ¹³C NMR (75 MHz, CDCl₃) δ 55.9 (2 C), 56.1, 60.9, 106.0 (2 C), 110.3, 115.0, 121.1, 129.0, 129.5, 130.6, 132.7, 137.1, 145.2, 145.8, 152.8 (2 C).
- ¹⁴ Cushman, M.; Nagarathnam, D.; Gopal, D.; Chakraboti, A. K.; Lin, C. M.; Hamel, E. *J. Med. Chem.* **1991**, 34, 2579.
- ¹⁵ Lin, C. M.; Singh, S. B.; Chu, P. S.; Dempsey, R. O.; Schmidt, J. M.; Pettit, G. R.; Hamel, E. *Mol. Pharmacol.* **1988**, 34, 200.
- ¹⁶ Maya, A. B.; Del Rey, B.; De Clairac, R. P.; Caballero, E.; Barasoain, I.; Andreu, J. M.; Medarde, M. *Bioorg. Med. Chem. Lett.* **2000**, 10, 2549.