A reassessment of the iron isotope composition of the Moon and its implications for accretion and differentiation of terrestrial planets

Franck Poitrasson^{a,*}, Thomas Zambardi^a, Tomas Magna^b and Clive R. Neal^c

^a Laboratoire Géosciences Environnement Toulouse,

Centre National de la Recherche Scientifique UMR 5563 – UPS – IRD – CNES, 14-16,
avenue Edouard Belin, 31400 Toulouse, France.

^b Czech Geological Survey, Klarov 3, CZ-11821 Prague 1, Czech Republic.

^c Department of Civil and Environmental Engineering and Earth Sciences, University of
Notre Dame, 156 Fitzpatrick Hall, Notre Dame, IN 46556, USA.

*Corresponding author; e_mail: Franck.Poitrasson@get.omp.eu

Submitted to *Geochimica et Cosmochimica Acta*

Second revised version 20th September 2019

Abstract

The Fe isotope composition of planetary bodies may provide constraints on their accretion modes and/or differentiation processes, but to do so, the Fe isotope systematics of key planetary reservoirs needs to be determined. To investigate this for the Moon, we measured the Fe isotope compositions for a suite of 33 bulk lunar mare basalts and highland rocks. Combined with published data, a compendium of 73 different lunar bulk rocks reveals a statistically significant Fe isotope difference between low-Ti and high-Ti mare basalts, yielding average $\delta^{57}\text{Fe} = 0.127 \pm 0.012\text{‰}$ (2SE; $n = 27$) and $\delta^{57}\text{Fe} = 0.274 \pm 0.020\text{‰}$ (2 SE; $n = 25$), respectively, relative to the IRMM-14 isotopic reference material. As lunar basalts are thought to reflect the Fe isotope composition of their respective mantle sources, the estimated relative proportion of the low-Ti and high-Ti source mantle suggests that the lunar upper mantle $\delta^{57}\text{Fe}$ value should be close to $0.142 \pm 0.026\text{‰}$. Whilst the composition of highland rocks (ferroan anorthosites and Mg-suite rocks) should provide a more global view of the Moon, the calculation of the mean $\delta^{57}\text{Fe}$ value of 15 available highland rock analyses yields $\delta^{57}\text{Fe} = 0.078 \pm 0.124\text{‰}$. Such a value is not defined precisely enough to be of critical use for comparative planetology. Ferroan anorthosites and Mg-suite rocks also give unresolvable means. It appears that Fe isotope heterogeneity among the lunar highland rocks is caused by non-representatively too small sample aliquots of coarse-grained rocks. It can also be the result of mixed lithologies for some. When the (kinetic) effect of olivine tending towards low $\delta^{57}\text{Fe}$ and feldspar with predominantly high $\delta^{57}\text{Fe}$ is cancelled, a more precise $\delta^{57}\text{Fe}$ value of $0.094 \pm 0.035\text{‰}$ is calculated. It is indistinguishable from the mean $\delta^{57}\text{Fe}$ of impact melts and is also similar to the upper lunar mantle estimate obtained from mare basalts. Collectively, this newly determined Fe isotope composition of the bulk Moon is indistinguishable from that of the Earth, and heavier than those reported for other planetary bodies. This planetary isotope

relationship is only observed for silicon given the currently available mass-dependent stable isotope database. Because both iron and silicon reside in the Earth's metallic core in significant quantities, this may point to the involvement of metallic cores of the Earth and Moon in the interplanetary Fe and Si isotope fractionation. Rather than via high-pressure metal–silicate fractionation at the core–mantle boundary, this would more likely be achieved by partial vaporization of the liquid outer metallic core in the aftermath of a Moon-forming giant impact.

Keywords: iron isotopes; Moon; mare basalts; anorthosites; Mg-suite; highland rocks.

1. Introduction

Investigation of the stable isotope compositions of planetary bodies other than the Earth has been facilitated through the return of lunar samples by the Apollo missions, as well as the recognition of Martian meteorites in existing collections (McSween Jr et al., 1979), and meteorites coming from the differentiated asteroid Vesta (Consolmagno and Drake, 1977). Following pioneering studies of oxygen, silicon and potassium isotope compositions (e.g. Clayton et al., 1971; Clayton and Mayeda, 1983; Molini-Velsko et al., 1986; Humayun and Clayton, 1995), the past 15 years witnessed rapid improvements in our knowledge of stable isotope compositions of a number of major and trace elements in lunar and other planetary bodies (e.g., Mg: (Wiechert and Halliday, 2007; Bourdon et al., 2010; Teng et al., 2010; Hin et al., 2017); Si: (Georg et al., 2007; Fitoussi et al., 2009; Ziegler et al., 2010; Zambardi et al., 2013); Ca: (Simon and DePaolo, 2010); Fe: (Poitrasson et al., 2004; Weyer et al., 2005; Schoenberg and von Blanckenburg, 2006; Wang et al., 2015; Sossi and Moynier, 2017); Zn: (Moynier et al., 2006; Paniello et al., 2012; Kato et al., 2015). These results were sometimes

controversial, either due to analytical difficulties and/or the use of older generation of mass spectrometers, less precise and accurate (e.g., Mg: (Wiechert and Halliday, 2007; Bourdon et al., 2010; Chakrabarti and Jacobsen, 2010; Teng et al., 2010); Si: (Georg et al., 2007; Fitoussi et al., 2009; Savage et al., 2010; Ziegler et al., 2010; Zambardi et al., 2013), or resulting from different interpretations of mass-dependent stable isotope variations (e.g., Fe: (Poitrasson et al., 2004; Weyer et al., 2005; Schoenberg and von Blanckenburg, 2006; Beard and Johnson, 2007; Poitrasson, 2007; Weyer et al., 2007; Poitrasson, 2009; Polyakov, 2009; Rustad and Yin, 2009; Wang et al., 2015; Elardo and Shahar, 2017; Elardo et al., 2019). For iron, this resulted in debates on estimating the bulk stable isotope composition of planets (see Fig. 1, for the Earth and the Moon). However, the interplanetary, mass dependent stable isotope variations hold great promise in unraveling early Solar System processes, such as planetary accretion mechanisms or mantle–core differentiation (Poitrasson et al., 2004; Georg et al., 2007; Polyakov, 2009).

Ideally, these stable isotope tracers should be little affected by common geological processes to allow easy interplanetary comparison (Humayun and Clayton, 1995). However, for those stable isotopes affected by processes such as fractional crystallization or element diffusion in solids, their fractionation mechanisms should be well understood and fully quantified to be useful for planet formation studies. Unfortunately, this is not yet the case for many of the elements under consideration. Beyond general rules common to all stable isotopes, such as the effect of temperature on the magnitude of isotopic fractionation or the stiffness of inter-atomic bonds (e.g. Schauble, 2004), initial studies have shown that elements such as Fe, Mg or O will not fractionate in the same way because they are located in different crystallographic coordination configurations or may be differently affected by redox processes. For this reason, it is important to determine and understand the systematics of the stable isotope fractionation during geological processes for all elements of interest.

Among these, iron has attracted considerable attention given its importance and large abundances in terrestrial planets, and the discovery of different planetary Fe isotope signatures (Poitrasson et al., 2004). However, with the expanding database, more complexities than originally anticipated have been highlighted. For example, Liu et al. (2010) clearly showed that there is a difference in the Fe isotope composition of lunar low-Ti and high-Ti mare basalts that may directly mirror that of their respective mantle sources. This supports the observations of Snyder et al. (1992) and Beard et al. (1998) who implied distinct mantle sources of low-Ti and high-Ti mare basalts on the basis of petrogenetic modeling and radiogenic isotope systematics, respectively. If true, this may eventually lead to a different estimate for the bulk lunar Fe isotope composition. This finding is of notable importance as it may provide a test for interplanetary Fe isotope variations imparted by either mantle–core differentiation or partial vaporization in the aftermath of a Moon-forming giant impact (Poitrasson, 2009).

In order to provide new constraints on Fe isotope fractionation during magmatic evolution of the lunar magma ocean, we have determined the Fe isotope composition in a suite of mare basalts and highland rocks. This new data set should enable a better evaluation of potential differences between various lunar reservoirs and their significance in creating bulk planetary signatures previously debated in the literature (Poitrasson et al., 2004; Weyer et al., 2005; Beard and Johnson, 2007; Poitrasson, 2007; Weyer et al., 2007; Weyer and Ionov, 2007; Liu et al., 2010; Wang et al., 2015; Elardo and Shahr, 2017; Sossi and Moynier, 2017; Elardo et al., 2019). Furthermore, the new Fe isotope data for several specimens were collected in order to (i) verify earlier Fe isotope determinations by means of low mass resolution MC-ICP-MS (Wiesli et al., 2003; Poitrasson et al., 2004), and (ii) constrain potential sample heterogeneity problems (e.g., for olivine-normative basalt 15555).

2. Samples

The samples were selected to reflect a wide range in petrological and geochemical types (Neal and Taylor, 1992; Warren, 1993; Papike et al., 1998; Neal, 2001; Meyer, 2011), as well as in their location on the Moon as sampled by the Apollo missions. Nine low-Ti and 12 high-Ti mare basalts, one KREEP basalt and one KREEP-rich impact melt, five ferroan anorthosites, two Mg-suite norites, one Mg-suite troctolite, one dunite, and one Mg-suite noritic anorthosite were analyzed. Only pristine highland rocks were selected for the new analyses reported in the present work according to criteria given in Warren (1993) defined on the basis of siderophile element contents and textural characteristics. This is important when considering that meteorite bombardment affects Fe isotope signatures on the lunar surface (Wiesli et al., 2003; Moynier et al., 2006; Poitrasson, 2007). These new data are compared to all previously published bulk lunar rock Fe isotope determinations, making up a Fe isotope database of 73 lunar rock samples. Meteorites of lunar origin (see their Fe isotope data in Wang et al., 2015) were not included in this compilation. This is due to an uncertain location of their origin on the Moon, strong shock modification and potential weathering on the Earth's surface. Most lunar meteorites were recovered from hot deserts and it has previously been shown that weathering under these conditions can significantly affect the Fe isotope signatures of bulk meteorites (Saunier et al., 2010).

3. Methods

Between 3 and 30 mg of bulk-rock powder prepared from 0.1 to one gram of the original sample chip allocation were digested (See Table 1). Samples processed at GET-CNRS, Toulouse were dissolved with a mixture of concentrated HF–HNO₃–HCl in closed

high-pressure Teflon vessels in an oven at 135°C for 6 days. For samples processed at the University of Lausanne, the same acid mixture was used and beakers were placed on a hotplate at 120°C for 3 days. The resulting solutions were subsequently evaporated to dryness and re-dissolved in 6M HCl on a hotplate at 120°C for 48 hours. Dried residues were thereafter dissolved using 6M HCl prior to Fe purification by conventional anion-exchange chromatography following established analytical protocols (Poitrasson et al., 2004, 2005). The total procedural blanks were between 3 and 7 ng, which is more than three orders of magnitude lower than Fe levels in samples and is thus considered negligible. Iron isotope compositions were determined using a Neptune (ThermoFisher Scientific, Bremen, Germany) high mass resolution multiple-collector inductively-coupled-plasma mass spectrometer (MC-ICP-MS) housed at CNRS Toulouse, following the procedures detailed in Poitrasson and Freydier (2005), which included mass bias correction of the purified Fe samples by Ni doping. This approach accurately corrects for mass bias shifts caused by residual matrix effects and results in a superior reliability of the measurements over the conventional standard–sample bracketing method (see Poitrasson and Freydier, 2005).

The new Fe isotope compositions are reported in Table 1 using the standard delta notation, in per mil (‰), relative to the European isotopic IRMM-14 reference material. Analytical reproducibility was estimated on the basis of 80 analyses of an in-house hematite standard from Milhas, Pyrénées (sometimes referred to in the literature as “ETH hematite standard”), conducted over 3 years and in the same analytical sequences as the sample analyses reported in Table 1. Every sample was typically analyzed six times; the long term reproducibility of such pooled measurements can be estimated on the basis of the hematite reference material analyses pooled in groups of six individual measurements. This yielded $\delta^{57}\text{Fe} = 0.758 \pm 0.067\text{‰}$ and $\delta^{56}\text{Fe} = 0.511 \pm 0.046\text{‰}$ (2SD). Additionally, BHVO-2 and JB-2 basalt reference materials analyzed during this study yielded $\delta^{57}\text{Fe}$ of $0.182 \pm 0.062\text{‰}$ and

0.095 ± 0.082‰ (2SD), respectively, in good agreement with literature values (Weyer et al., 2005; Craddock and Dauphas, 2011).

In the reminder of this paper, the uncertainties of the sample set mean values are calculated as two standard errors of the mean (2SE) rather than two standard deviations (2SD) in order to indicate how well the mean values are constrained rather than to give a measure of the spread of a set of individual measurements around the mean (see e.g. Miller and Miller, 1993). The 2SE calculation involves the use of the t-correcting factor that takes into account the number of samples used in the calculation (Platzner, 1997). Furthermore, these 2SE uncertainties represent a direct proxy of the Student's t-test (e.g., Poitrasson et al., 2004, 2005; Zambardi et al., 2013). All literature data have been recalculated in this way to be mutually comparable. However, irrespective of whether calculated as either 2SE or 2SD, it is important to note that these estimated uncertainties only include analytical random errors but not systematic ones potentially associated with analytical or sampling biases.

4. Results and comparison with previous data

Thirty three new Fe isotope analyses of bulk lunar rock samples are listed in Table 1. The low-Ti basalts and high-Ti basalts show well-defined but mutually distinct $\delta^{57}\text{Fe}$ ranges from $0.043 \pm 0.023\text{‰}$ to $0.154 \pm 0.071\text{‰}$ and from $0.167 \pm 0.118\text{‰}$ to $0.354 \pm 0.118\text{‰}$, respectively. The new Fe isotope data for low-Ti mare basalts confirm previous observations of homogeneous $\delta^{57}\text{Fe}$ systematics, similar to Earth (Liu et al., 2010). Two high-K mare basalts (10049, 10057) from the Apollo 11 high-Ti group show systematically lower $\delta^{57}\text{Fe}$ values ($\leq 0.187\text{‰}$) compared with the rest of the Apollo 11 suite ($\geq 0.290\text{‰}$) and the entire Apollo 17 suite ($\geq 0.213\text{‰}$). KREEP basalt 15386 has $\delta^{57}\text{Fe} = 0.18 \pm 0.07\text{‰}$, intermediate between low-Ti and high-Ti mare basalts.

The new highland data reveal a bulk rock $\delta^{57}\text{Fe}$ range from $-0.73 \pm 0.06\text{‰}$ to $0.50 \pm 0.06\text{‰}$ (Table 1). This $>1.2\text{‰}$ variation is surprisingly large for igneous rocks, and is only approached by terrestrial mantle peridotites (see Williams et al., 2005; Weyer and Ionov, 2007; Zhao et al., 2010; Poitrasson et al., 2013). Ferroan anorthosites span a $\delta^{57}\text{Fe}$ range from $-0.39 \pm 0.05\text{‰}$ to $0.50 \pm 0.06\text{‰}$ although the negative $\delta^{57}\text{Fe}$ datum has only been measured for sample 60025. Two Mg-suite norites 15455 and 77215 exhibit $\delta^{57}\text{Fe}$ values of $0.07 \pm 0.02\text{‰}$ and $0.05 \pm 0.05\text{‰}$, respectively. Dunite 72415 yielded an extremely light $\delta^{57}\text{Fe}$ value of $-0.73 \pm 0.06\text{‰}$ (Table 1 and Fig. 2). Mg-suite troctolite 76335 has $\delta^{57}\text{Fe} = -0.01 \pm 0.06\text{‰}$. The two most negative values, from dunite 72415 and Fe-anorthosite 60025, were fully duplicated by new sample powder dissolutions, iron purifications by ion exchange chromatography and MC-ICP-MS measurements. In both cases, identical results were obtained so the figures reported in Table 1 represent the average of these duplicate determinations. The KREEP-rich basaltic impact melt 14310 yielded a $\delta^{57}\text{Fe} = 0.13 \pm 0.13\text{‰}$, similar to the value for pristine KREEP basalt 15386.

Some new analyses reported in Table 1 represent replicate data obtained on a new sample aliquot from the same Apollo rock compared to previously published values. Among these, the new $\delta^{57}\text{Fe}$ values for several samples (low-Ti mare basalts 12045 and 14053, high-Ti mare basalts 10057, 70135 and 74275, KREEP-rich basaltic impact melt 14310, KREEP basalt 15386, Mg-suite norite 15455, anorthosites 15455 and 62255) are indistinguishable within uncertainties from those reported elsewhere (Wiesli et al., 2003; Poitrasson et al., 2004; Weyer et al., 2005). In contrast, new $\delta^{57}\text{Fe}$ values for Mg-suite norite 77215 are significantly lower than the value published by Poitrasson et al. (2004), whereas the new $\delta^{57}\text{Fe}$ value for high-Ti basalt 10003 is higher than that of Sossi and Moynier (2017). A full replicate analysis of high-alumina basalt 14053 (Table 1), for which different chips were obtained in two different CAPTEM allocations (to CRN and FP), have indistinguishable Fe

isotope compositions; yet the recent $\delta^{57}\text{Fe}$ value by Sossi and Moynier (2017) is higher. This could be caused by modest sample heterogeneity because high-Al basalt 14053 was subject to thermal metamorphism in an impact melt sheet that caused native Fe to form in the outer portion of the sample (Taylor et al., 2004). The new measurement of Mg-suite dunite 72415 yielded $\delta^{57}\text{Fe} = -0.73 \pm 0.06\text{‰}$, which is statistically significantly lighter than the already very light previously reported values of $-0.53 \pm 0.03\text{‰}$ (Wang et al., 2015), and $-0.60 \pm 0.05\text{‰}$ (Sossi and Moynier, 2017). Therefore, the very light Fe isotope signature of 72415 is confirmed (Table 1 and Fig. 2).

Data reported for olivine-normative low-Ti basalt 15555 by Poitrasson et al. (2004) and Weyer et al. (2005) were differing by more than 0.16‰, and therefore well outside the quoted analytical uncertainties. This possibly results from the medium- to coarse-grained nature of this rock with variable olivine modal contents between different rock chips (Ryder, 1985). Our new Fe isotope determination for basalt 15555 yields a $\delta^{57}\text{Fe}$ value intermediate to those published elsewhere (Poitrasson et al., 2004; Weyer et al., 2005; Liu et al., 2010; Wang et al., 2015; Sossi and Moynier, 2017). There are no published $\delta^{57}\text{Fe}$ values yet of olivines from low-Ti basalt 15555. Taking the $\delta^{57}\text{Fe} = -0.117\text{‰}$ for olivine from low-Ti ilmenite basalt 12045 (Poitrasson et al., 2004) and assuming that the variable modal content of olivine in 15555 will not be accompanied by distinctive olivine Fe isotope compositions, we can compute a bulk rock $\delta^{57}\text{Fe}$ ranging from 0.141 to 0.101‰ with olivine modal content varying respectively from 5% to 20%, depending on the 15555 sample allocation (Meyer, 2011). This 0.040‰ range is four times smaller than the range of 15555 bulk rock values published so far, which is 0.164‰ from six different rock allocations (Table 1). Taking the other low-Ti basalt (sample 12021) olivine value of 0.08‰ published by Wang et al. (2015) would make the range among calculated bulk rock $\delta^{57}\text{Fe}$ values for 15555 even smaller since it is closer to the reported bulk rock values. This suggests that other causes than solely olivine modal contents

are responsible for the scatter among the published bulk rock $\delta^{57}\text{Fe}$ values for basalt 15555. As discussed below, it is likely that unrepresentative aliquot allocations of coarse-grained and mineralogically heterogeneous rocks such as 15555 is the main reason for this discrepancy. The most extreme discrepancy with previous results is observed for ferroan anorthosite 60025. The new $\delta^{57}\text{Fe}$ result is $>0.6\text{‰}$ lighter than the value reported in Poitrasson et al. (2004). A full replicate of the entire analytical protocol at CNRS Toulouse yielded an identical result confirming the reliability of the analytical and instrumental procedures. Petrologic heterogeneity within 60025 with various modal content of felsic and mafic minerals depending on the sample fragment has been previously documented (e.g. James et al., 1991; Torcivia and Neal, 2017, 2018). This will be discussed in more detail below.

5. Discussion

5.1 The key role of the sample size for representative Fe isotope analysis

The inter-laboratory comparison of Fe isotope measurement procedures on well-homogenized silicate rock reference materials commonly shows good agreement (Beard and Johnson, 2006; Poitrasson, 2006, 2007; Craddock and Dauphas, 2011). Hence, the analytically significant $\delta^{57}\text{Fe}$ differences between new data and published values for lunar samples 77215, 10003, 14053, 15555, 60025 and 72415 appear to reflect sample heterogeneity rather than analytical issues. Literature data for high-Ti mare basalt 70035 also yield discrepant $\delta^{57}\text{Fe}$ values (Fig. 2 and Supplementary Table). Given that the entire replicated lunar sample data were obtained on powder aliquots produced from different rock chips, it is possible that for at least these seven lunar rocks, the sample mass used to prepare the powders was too small to be representative of the bulk rock $\delta^{57}\text{Fe}$ given the variable mineral grain size. These sample sizes typically range from ~ 50 mg to ~ 1 g, but some studies

report Fe isotope analyses based on the direct dissolution of rock chips of “a few milligrams” size (Wang et al., 2015). For the smaller sample sizes, Fe isotope variations probably reflect the relative differences in abundance of the major mineral phases sampled (e.g. olivine in basalt 15555, Ryder, 1985), which may carry distinctly different Fe isotope signature (Poitrasson et al., 2004; Craddock et al., 2010; Wang et al., 2015). This would translate into rock chips used for the analyses having olivine modal contents well outside the range of 5 to 20% reported for 15555 (Meyer, 2011). In contrast, the Fe isotope analysis of bulk terrestrial rocks would be performed with an aliquot of powder typically produced from the crushing and homogenization of at least several hundred grams of rock (see general rock sample preparation methods in Potts, 1987). Though often underestimated, such a sampling issue was previously pointed out for lunar Apollo rocks. For example, trace and even major element concentration heterogeneity issues were noted for basalt 15555 when comparing analyses based on powders made from rock aliquots of less than 1g (Ryder and Steele, 1988; Ryder and Schuraytz, 2001). This was also reported to bias the oxygen isotope compositions of basalts 75075 and 15555 (Spicuzza et al., 2007; Liu et al., 2010), and, more recently, the silicon isotope composition of ferroan anorthosites 65035 and 60015 (Poitrasson and Zambardi, 2015), when powders were produced from rock chips weighing less than ~100 mg (Armytage et al., 2012). Moreover, for basalt 15555, an anomalously light Mg isotope signature reported recently (Sedaghatpour and Jacobsen, 2019), combined with significant Fe–Mg inter-mineral diffusion affecting Mg isotope systematics in olivine crystals of 15555 (Richter et al., 2016), provide an additional clue for possible sample Fe isotope heterogeneity issue (see Table 1).

Our new results (Table 1 and Fig. 2) reveal an extended range of Fe isotope compositions for lunar bulk rock samples (−0.73 to 0.50‰) relative to what has been published so far (Wiesli et al., 2003; Poitrasson et al., 2004; Weyer et al., 2005; Liu et al.,

2010; Wang et al., 2015; Sossi and Moynier, 2017) with the most extreme values observed among the highland rocks (Fig. 2 and Table 1). Most ferroan anorthosites show distinctly heavy Fe isotope composition ($\delta^{57}\text{Fe} > 0.18\text{‰}$), similar to high-Ti basalts (Figure 2 and see also Poitrasson, 2007), or even higher with ferroan anorthosite 15415 yielding $\delta^{57}\text{Fe} = 0.50\text{‰}$ (This study). This is consistent with the results of Fe isotope investigations of mineral separates performed by Craddock et al. (2010) on a suite of low-Ti and high-Ti mare basalts, clearly identifying ilmenite and plagioclase as ^{57}Fe -enriched phases.

Two different CAPTEM allocations of 60025 deviate by $>0.6\text{‰}$ in $\delta^{57}\text{Fe}$ (Table 1). Iron is predominantly hosted by pyroxene and olivine in lunar rocks (Papike et al., 1998) and if these phases are isotopically fractionated as is the case for mare basalts (Poitrasson et al., 2004; Craddock et al., 2010), an effect on the Fe isotope signature of bulk anorthosites may be expected since the occurrence of an isotopically fractionated Fe-rich mafic mineral phase may modify the $\delta^{57}\text{Fe}$ values. These pilot studies on mineral separates from lunar basalts also indicated that plagioclase is isotopically heavier than coexisting olivine and pyroxene. This could indeed explain the heavy Fe isotope signature of anorthosites compared to some other mafic highland rocks. In the case of 60025, Poitrasson et al. (2004) reported Fe content of 0.50 wt.%, whereas the analysis of the allocated sub-sample 60025,874 revealed 4.84 wt.% Fe, nearly one order of magnitude higher abundance (Supplementary Table). Such a composition approaches those of mafic portions of 60025 (Ryder, 1982; James et al., 1991; Meyer, 2011) and confirms the larger occurrence of olivine and pyroxene in our sample allocation, most probably leading to notably lower $\delta^{57}\text{Fe}$ signature relative to the other $\delta^{57}\text{Fe}$ measurements of 60025 and other anorthosites. In addition, Ryder (1982) concluded that systematic variations in the mineral chemistry of 60025 were probably a reflection that this sample was a mixture of closely related materials. Future work requires rigorous investigations by in situ isotopic work on anorthosite mineral phases (e.g., Oeser et al., 2015)

to fully evaluate this. The influence of olivine is well illustrated at the bulk-rock scale with dunite 72415 having ~93% modal olivine (Dymek et al., 1975) and with reported $\delta^{57}\text{Fe}$ values ranging from -0.5 to -0.73‰ depending on the rock chip analyzed (Table 1 and Fig. 2). Such isotopically light values in olivine could result from chemical diffusion-driven kinetic disequilibrium between olivine and the last melt olivine was in contact with (Wang et al., 2015). The implications of these findings are further discussed below.

5.2 Iron isotope composition of different lunar reservoirs

Bulk lunar samples may be grouped into four different sets given their petrologic characteristics (e.g., Papike et al., 1998): Low-Ti basalts, high-Ti basalts, highland rocks and impact melts. Nine new low-Ti basalts yield a mean $\delta^{57}\text{Fe} = 0.113 \pm 0.023\text{‰}$ (2SE) and 12 new high-Ti basalts analyses give $\delta^{57}\text{Fe} = 0.273 \pm 0.039\text{‰}$ (2SE). These values are indistinguishable from mean $\delta^{57}\text{Fe}$ values of $0.118 \pm 0.028\text{‰}$ (2SE, $n = 10$) and $0.291 \pm 0.051\text{‰}$ (2SE, $n = 7$), reported by Liu et al. (2010) for low- and high-Ti basalts, respectively. It should be noted that our new data include a full replicate analysis of high-alumina basalt 14053 (Table 1) for which different chips were obtained in two different CAPTEM allocations. Given that these replicate analyses are indistinguishable within uncertainty, their mean value is used for the low-Ti basalt average $\delta^{57}\text{Fe}$ reported above. The new data from this study together with previously published results by Wiesli et al. (2003), Poitrasson et al. (2004), Weyer et al. (2005), Liu et al. (2010), Wang et al. (2015) and Sossi and Moynier (2017) yield a cumulative mean $\delta^{57}\text{Fe}_{\text{low-Ti}} = 0.127 \pm 0.012\text{‰}$ (2SE, $n = 27$; Table 1). For replicate analyses, the average $\delta^{57}\text{Fe}$ of independent determinations is used to avoid overweighting of samples analyzed multiple times in the calculated mean. Using the same approach, we obtain a mean $\delta^{57}\text{Fe}_{\text{high-Ti}} = 0.274 \pm 0.020\text{‰}$ (2SE, $n = 25$; Table 1). Our eleven

new determinations of $\delta^{57}\text{Fe}$ in highland rocks lead to a less well-defined mean because of the significant Fe isotope variations among the samples. With the literature data included, the mean $\delta^{57}\text{Fe}_{\text{highland}} = 0.078 \pm 0.124\text{‰}$ (2SE, $n=15$; Table 1) is calculated. Within this group, Fe-anorthosites yield a mean $\delta^{57}\text{Fe}$ composition of $0.192 \pm 0.207\text{‰}$ (2SE; $n = 6$) whereas a $\delta^{57}\text{Fe}$ value of $0.002 \pm 0.179\text{‰}$ (2SE; $n = 9$) is obtained for the Mg-suite rocks. With literature data included, impact melts yield a mean $\delta^{57}\text{Fe}_{\text{impact}} = 0.093 \pm 0.055\text{‰}$ (2SE, $n=5$).

The new data imply that the Fe isotope systematics of the Moon is more complex than initially thought. As highlighted by Liu et al. (2010), low-Ti and high-Ti mare basalts possess distinctive Fe isotope signatures (Fig. 2). With the new and published mare basalt data included, a *t*-test allows us to reject the null hypothesis of a common origin of low-Ti and high-Ti basalts with a level of confidence well above 99% (See Supplementary Table). Furthermore, the *t*-tests show that both the highland rocks and the impact melts have Fe isotope composition indistinguishable from that of the low-Ti basalts, whereas their mean $\delta^{57}\text{Fe}$ values are statistically different from that of high-Ti basalts with a degree of confidence higher than 99% (Supplementary Table). Hence, from the viewpoint of the observed Fe isotope systematics, the different rock types may be arranged as two isotopically distinct groups: a ^{57}Fe -enriched end-member defined by high-Ti basalts, and a ^{57}Fe -depleted end-member best defined by low-Ti basalts, but also including impact melts and highland rocks, though the latter do not define a well-constrained mean $\delta^{57}\text{Fe}$ value (Tables 1 and Supplementary Table).

Impact melts do not represent a deep lunar reservoir since their origin in meteoritic bombardment events occur at the lunar surface. Based on the review of Shearer et al. (2006), low-Ti and high-Ti basalts make up a significantly larger component of the Moon in that they cover 17% of lunar surface, but they represent just ~1% of the lunar crust volume (Head, 1976). The largest reservoir for which we have direct samples are highland rocks,

representing the remaining 99% of the lunar crust, that is, ~10% of the volume of the Moon (Papike et al., 1998). However, the present work reveals that this reservoir is more heterogeneous than previously anticipated (Poitrasson et al., 2004), with a total $\delta^{57}\text{Fe}$ range of ~1.2‰. Subdividing these highland rocks into two groups consisting of ferroan anorthosites and Mg-suite rocks makes no difference since both groups are also very heterogeneous isotopically, as computed above. As discussed above, this likely stems from sample aliquot allocations that are too small to produce representative $\delta^{57}\text{Fe}$ analyses of bulk rocks given the frequently coarse-grained nature of highland rocks. Whether calculated as a highland rock group or separated as a ferroan anorthosite and Mg-suite groups, the uncertainties of the means are too high for the precision required for comparative planetology studies (i.e., <0.1‰ level; see Dauphas et al., 2009 and Poitrasson et al., 2013).

The next step is to determine whether the rock groups sampled at the surface of the Moon (i.e., low-Ti, high-Ti mare basalts, highland rocks...) are actually representative of larger-scale lunar reservoirs. Liu et al. (2010) have shown that low-Ti and high-Ti mare basalts do not exhibit Fe isotope variations as a function of magmatic differentiation. This feature has also been observed for most terrestrial igneous rocks, except for highly evolved silicate compositions, such as granites and rhyolites (Poitrasson and Freydier, 2005; Schoenberg and von Blanckenburg, 2006; Heimann et al., 2008; Schuessler et al., 2009; Telus et al., 2012; Foden et al., 2015). Iron isotope variations in mafic rocks have been observed in the closed-system Kilauea Iki lava lake, Hawaii (Teng et al., 2008), that were subsequently shown to be caused by kinetic effects (Sio et al., 2013). Therefore, the question arises whether or not mare basalts directly reflect the Fe isotope composition of their respective mantle sources. The importance of partial melting in the Earth's mantle to generate systematic Fe isotope differences in derivative melts has been debated for some time. Whereas some consider this process to play a key role (Weyer et al., 2005; Weyer and Ionov, 2007; Williams

et al., 2005; 2009; Dauphas et al., 2009), others suggested that partial melting of pristine peridotitic lithologies generates only minor $\delta^{57}\text{Fe}$ variations in the basaltic melts relative to their peridotitic protoliths (Beard and Johnson, 2007; Poitrasson, 2007; Zhao et al., 2010; Poitrasson et al., 2013). Accordingly, using Nuclear Resonant Inelastic X-ray Scattering (NRIXS) measurements of synthetic glass and associated partial melting calculations, Dauphas et al. (2014a) could explain only $\sim 1/3$ of the observed Fe isotopic difference between measured MORBs and their estimated bulk silicate mantle. The issue is essentially linked to their incorrect estimate of the bulk silicate mantle $\delta^{57}\text{Fe}$. If, instead of chondritic ($\sim 0\text{‰}$; Dauphas et al., 2009; 2014a) a $\delta^{57}\text{Fe}$ value of $\sim 0.1\text{‰}$ is adopted for the Earth's mantle (Poitrasson et al., 2004; 2013), then the discrepancy noted by Dauphas et al. (2014a) between their experimental results and natural observations disappears. Hence, this issue remains debated for the Earth and further experimental work is required. The putative mechanism to explain Fe isotope fractionation during partial melting proposed by Dauphas et al. (2009; 2014a) hinges on the preferential incorporation of isotopically heavy Fe^{3+} in melts relative to solid residuum. However, Liu et al. (2010) recognized that ferric iron does not occur in the lunar mantle (Sato et al., 1973; Sato, 1976) and that experimental studies have shown the lack of equilibrium Fe isotope fractionation between Fe metal (occurring in the lunar mantle) and the prevalent Fe^{2+} species at mantle conditions (Roskosz et al., 2006; Poitrasson et al., 2009). As a result, Liu et al. (2010) concluded that: 1) partial melting in the lunar mantle probably did not produce Fe isotope fractionation, and 2) Fe isotope compositions of low-Ti and high-Ti mare basalts directly mirror those of their corresponding lunar mantle sources. These conclusions have recently been challenged by Sedaghatpour and Jacobsen (2019) on the basis of a fractional crystallization model. However, these models are frequently underconstrained, with many of the unknown parameters adjusted to fit the observed bulk-rock analyses (see review by Dauphas et al., 2017). As a result, these models do not provide a strong constraint

on the actual process at play to explain the observed bulk-rock isotopic compositions measured. For instance, Sedaghatpour and Jacobsen (2019) used an estimated $\Delta^{57}\text{Fe}_{\text{olivine-melt}}$ fractionation factor of -0.21‰ whereas it was shown experimentally to be at $\sim 0\text{‰}$ for mafic melts, including those having the chemistry of mare basalts (Prissel et al., 2018). We thus take this as a strong support for the conclusions derived by Liu et al. (2010).

Many authors have estimated the mineralogy of mantle sources of low-Ti and high-Ti basalts. For example, Snyder et al. (1992) and Beard et al. (1998) inferred an important role of ilmenite in the genesis of high-Ti basalts, although physical constraints of the lunar magma ocean evolution must also be considered (e.g. Elkins-Tanton et al., 2011). Poitrasson (2007) discussed that ilmenite, although potentially explaining elevated Ti contents of high-Ti basalts (Snyder et al., 1992), would be incapable of shifting $\delta^{57}\text{Fe}$ towards high values given that theoretical $\log\beta$ factors (a reduced isotopic partition function factor) calculated for ilmenite were among the lowest for silicate and oxide minerals (compare Figures 1 and 2 from Polyakov and Mineev, 2000). As a consequence, melting of an ilmenite-rich source should not result in basaltic melts with elevated $\delta^{57}\text{Fe}$ values. However, more recent Fe isotope analyses of mineral phases of mare basalts have shown that ilmenite is in most cases isotopically heavier than coexisting pyroxene and plagioclase (Craddock et al., 2010). To directly compare Fe isotope data for mineral separates with theoretical calculations, we have to assume that the mineral fractions analyzed by Craddock et al. (2010) are pure and represent equilibrium mineral isotope compositions. This might not be the case for all samples in their study given the observed scatter in inter-mineral Fe isotopic fractionation, however. With this caveat in mind, it would indicate that, similar to pyrite (Blanchard et al., 2009, 2012; Polyakov et al., 2013), the Fe isotope fractionation factor for ilmenite estimated by Polyakov and Mineev (2000) appears to be grossly underestimated. If correct, the heavy Fe isotope signature of high-Ti mare basalts may be explained by partial melting of ilmenite in their

source mantle as proposed by Liu et al. (2010). They based their model on the inferences from Snyder et al. (1992) and Elkins-Tanton et al. (2011) suggesting that >78% of the early lunar magma ocean crystallized olivine and orthopyroxene, whereas an ilmenite-rich cumulate formed only after 95% lunar magma ocean crystallization. Combined with the remote sensing study of Giguere et al. (2000) indicating that high-Ti basalts should represent ~10% of all lunar maria, Liu et al. (2010) estimated that only ~10% of the lunar upper mantle is akin to the source of the high-Ti mare basalts, whereas the remaining is similar to the source of the low-Ti mare basalts. Given that mare basalts likely mirror the Fe isotope composition of their source (see above), we can estimate the $\delta^{57}\text{Fe}$ of the lunar upper mantle at $0.142 \pm 0.026\%$ (Fig. 2) from the newly derived mean $\delta^{57}\text{Fe}$ values of low-Ti and high-Ti basalts, combined with the relative proportion of their mantle protoliths (of respectively 90 and 10%, with an estimated uncertainty of 10% by Liu et al., 2010), using a simple mass balance calculation. This value would apply to the upper ~250 km of the lunar mantle considering mare basalt petrogenesis (Longhi, 1992). The estimated size of this reservoir from seismology extends down to 500 km, where the limit between the upper and lower lunar mantle was set (Nakamura, 1983).

It is interesting to note that lunar volcanic picritic glasses, thought to have been derived from the melting of protoliths located 360–520 km deep in the lunar mantle (Delano, 1979; Elkins-Tanton et al., 2003), show somewhat lighter Fe isotope signatures. However, the $\delta^{57}\text{Fe}$ values scatter by over ~0.5‰, even for the same samples (Poitrasson et al., 2004; Weyer et al., 2005; Moynier et al., 2006; Sossi and Moynier, 2017). Whether this reflects Fe isotope stratification intrinsic to the lunar mantle, an effect of volcanism through fire fountaining (Poitrasson et al., 2004; Sossi and Moynier, 2017) or a sample heterogeneity issue remains to be tested.

The lunar highland rocks, and notably the anorthosites that make up ~80% of the crust (Papike et al., 1998), may potentially give a more comprehensive view of the lunar mantle composition given that it was produced by magma ocean differentiation. Estimates for depth of the early lunar magma ocean may vary from 500 to 1200 km, and up to the entire Moon, depending again on whether seismic or petrological constraints are preferred (Shearer et al., 2006; Wiczorek et al., 2006). However, as discussed above, the majority of the highland samples for which Fe isotope data exist may show a potential effect of their coarse-grained, essentially mono-mineral nature. This makes the determination of the Fe isotope composition of the bulk lunar highlands reservoir based solely on mean $\delta^{57}\text{Fe}$ values of these rocks rather imprecise ($0.078 \pm 0.124\text{‰}$; Table 1) to provide useful constraints. Even if highland rocks are split into ferroan anorthosites and Mg-suite rocks, the large uncertainties remaining do not change this conclusion. As a parallel to the isotopically heterogeneous Earth's mantle possibly linked to partial melting events (Williams et al., 2005; Weyer and Ionov, 2007; Williams and Bizimis, 2014), but most importantly due to metasomatic effects (Weyer et al., 2007; Zhao et al., 2010; Poitrasson et al., 2013), a more precise $\delta^{57}\text{Fe}$ estimate of this reservoir can only be provided by a careful consideration of the petrology of these rocks, even though metasomatic processes are less likely for the Moon (Wang et al., 2015). It has been shown above that plagioclase leads to heavy bulk-rock Fe isotope compositions whereas olivine tends towards isotopically light values in case of diffusion. Besides sample aliquot representativeness issues, such kinetic effects in olivine may make the bulk rock $\delta^{57}\text{Fe}$ lighter (Collinet et al., 2017). In addition, these rocks may also represent mixed lithologies (Ryder, 1982; James et al., 1991). Plotting the bulk-rock $\delta^{57}\text{Fe}$ versus the modal content of the other major phases from these samples showing less variable Fe isotope fractionation, i.e., pyroxenes, leads to a more precise Fe isotope composition for highland rocks (Fig. 3). By using samples with >10 modal % pyroxene, a better-constrained mean $\delta^{57}\text{Fe}$ of $0.094 \pm 0.035\text{‰}$ is obtained for lunar

highlands. This value essentially corresponds to the mean of the six Mg-suite norites for which Fe isotope compositions are available (Supplementary Table), although three anorthosites and one Mg-suite troctolite also fall on this mean value (Fig. 3).

5.3 Implications for the Fe isotope composition of bulk Moon

Overall, it appears that the $\delta^{57}\text{Fe}$ estimate of the lunar upper mantle sampled by mare basalts ($0.142 \pm 0.026\text{‰}$) is indistinguishable from the value obtained from highland rocks ($0.094 \pm 0.035\text{‰}$; Fig. 3) that likely sample a larger portion of the lunar mantle. If partial melting of the lunar mantle imparted no measurable Fe isotope fractionation (Liu et al., 2010; Sossi and Moynier, 2017), it is likely that the highland $\delta^{57}\text{Fe}$ value of $0.094 \pm 0.035\text{‰}$ is a relevant estimate of the silicate portion of the Moon, considering the mass balance of these reservoirs.

Schoenberg and von Blanckenburg (2006) and Elardo and Shahar (2017) have hypothesized that Fe isotope fractionation between silicate portions of terrestrial planets and their metallic cores could occur. In contrast, the investigation of Fe isotope compositions in various meteorite classes led to the conclusion that this mechanism is unlikely for planetary bodies (Poitrasson et al., 2005), an inference also supported by subsequent experimental investigations (Roskosz et al., 2006; Poitrasson et al., 2009; Hin et al., 2012). This conclusion was challenged using the three-isotope experimental approach (Shahar et al., 2015), although four out of the five more recent experimental results of Elardo et al. (2019) showed no statistically significant Fe isotope fractionation between metal and silicate. In practice, this three-isotope methodology involves, at high temperature, a first thorough chemical and isotopic mixing between the liquid metallic alloy and the silicic melt followed by “un-mixing” along a secondary fractionation line according to Shahar et al. (2017, see their Fig. 2B). As a

result, the remainder of the system evolution until isotopic equilibrium is very similar to time-series experiments conducted without spike in which losses are monitored by mass balance. Besides time series, the experiments of Poitrasson et al. (2009) were reversed, and the lack of Fe isotope fractionation between metal and silicate at equilibrium under planetary core formation conditions was also concluded in studies using a completely different experimental set ups (Roskosz et al., 2006; Hin et al., 2012; and see also results of Elardo et al., 2019), as well as in studies using meteorite samples (Poitrasson et al., 2005; Chernonozhkin et al., 2016, 2017; Jordan et al., 2019). Further, this three-isotope experimental methodology could also be affected by possible kinetic biases (Bourdon et al., 2018), so results obtained by this method should be interpreted with caution.

On a more theoretical side based on NRIXS measurements of high-pressure mineral phases, Polyakov (2009) proposed that a very high pressure-induced phase change in the lower mantle, at pressures above ~ 100 GPa, may generate Fe isotope differences between the silicate Earth and metallic core. This hypothesis does not apply to the lunar core-mantle differentiation, though, given the too low pressures occurring in the lunar interior. The origin of the lunar material in the aftermath of a Moon-forming giant impact is debated (see review by Dauphas et al., 2014b), with models involving various proportions of the impactor material and the proto-Earth mantle. However, the Moon forming material is unlikely to be inherited from the silicate portion of the Earth from a giant impact occurring at a point when the Earth was already big enough to reach 100 GPa since only a very small portion of the deep Earth mantle would show this heavy Fe isotope signature in the framework of Polyakov (2009) theory. Moreover, subsequent studies using similar NRIXS approaches did not support Polyakov (2009) inferences since they have concluded that metal-silicate at high pressure and high temperature conditions should result in undetectably small Fe isotope fractionation (Dauphas et al., 2014a; Liu et al., 2017; Yang et al., 2019). Therefore, the estimate of Fe

isotope composition of the lunar silicate mantle also involves its small metallic core and it is thus valid for the bulk Moon (Fig. 2). Collectively, at the current level of knowledge, the bulk lunar $\delta^{57}\text{Fe}$ composition should be close to $0.094 \pm 0.035\text{‰}$, which is indistinguishable from the Earth's value of $0.10 \pm 0.03\text{‰}$ (Fig. 2).

This lunar estimate agrees well with the recent value proposed by Sossi and Moynier (2017), though on the basis of a more limited set of lunar samples (five Mg-suite rocks consisting mostly of norites) and using different lines of reasoning. It is significantly heavier than the other recent estimate of Wang et al. (2015) that was based on a sole sample only (dunite 72415); that particular sample has been shown to yield rather heterogeneous $\delta^{57}\text{Fe}$ values depending on the sample allocation (Fig. 2) and it was based on a corrected isotopic Fe composition using isotopic diffusion parameters in olivine that have been revised since (Oeser et al., 2015).

5.4 Implications for the formation of the Moon

Given the previously described experimental and theoretical disagreements noted in the literature as to whether the mantle–core differentiation did generate isotopically distinct silicate and metallic reservoirs in the interior of the Earth, the bulk lunar Fe isotopic composition can constitute a good test of this idea. If correct, we should expect a silicate Earth with a heavier Fe isotope composition than that of the silicate Moon since the high pressures required for the phase transition to impart additional Fe isotope fractionation were not reached in the lunar interior. Realizing that no observation supports a Moon isotopically lighter than the silicate fraction of the Earth, Polyakov (2009) and Rustad and Yin (2009) concluded that a Moon-forming giant impact is required to enable the Moon to acquire the heavy Fe isotope composition of the Earth through a homogenization process such as that

proposed by Pahlevan and Stevenson (2007). Such a scenario implies that a large fraction of the Moon-forming material came from the proto-Earth mantle, which is not easy to achieve even using high resolution numerical simulations (Canup et al., 2013). Furthermore, and as discussed above, another constraint to this scenario is that the giant impact should have occurred sufficiently late in the proto-Earth accretion history to achieve a mantle-core boundary pressure well beyond ~ 100 GPa pressure at which the Fe isotope metal–silicate fractionation process proposed by Polyakov (2009) can operate. However, the current Earth’s mantle–core boundary is only ~ 32 GPa higher than this limit, so the proportion of mantle having the phase change and associated isotopic effect proposed by Polyakov (2009) is limited. Therefore, the Earth should have nearly reached its current size and enough time should be allowed to lead the deep mantle to impart its Fe isotope composition to the shallower portions by convection before the impact. However, it appears that terrestrial mantle homogenization was not achieved ~ 1 Ga after Earth’s formation (Bennett et al., 2007; Touboul et al., 2012; Debaille et al., 2013), so well after the putative Moon-forming giant impact timeframe. Furthermore, the Moon forming material should essentially come from the proto-Earth to make this scenario tenable, which seems unlikely (Canup et al., 2013; Dauphas et al., 2014b). Hence, our best estimate for the Moon $\delta^{57}\text{Fe}$ value, which is indistinguishable from that of the Earth (Fig. 2), therefore also provides a constraint against terrestrial mantle–core Fe isotope fractionation.

However, whereas meteorite and experimental studies conclude the absence of Fe isotope fractionation during mantle–core differentiation of planetary bodies (Poitrasson et al., 2005; Roskosz et al., 2006; Poitrasson et al., 2009; Hin et al., 2012; Chernonozkhin et al., 2016, 2017), some recent experimental studies propose an opposite sense of fractionation during mantle–core differentiation, that is, with a metallic portion becoming isotopically heavier (Shahar et al., 2015; Elardo and Shahar, 2017; Elardo et al., 2019). Such a sense of Fe

isotope fractionation would yield planetary mantles isotopically lighter than chondrites. This finds neither evidence from the lunar samples analyzed so far apart from the extremely heterogeneous volcanic glasses (see above), nor is this supported by the latest planetary $\delta^{57}\text{Fe}$ estimates (see Wang et al., 2012; Sossi et al., 2016; Dauphas et al., 2017).

A simpler scenario can explain the heavy Fe isotope composition of the Earth, Moon and angrite parent body through the increased volatility of light Fe isotopes during the accretion of the protoplanetary disk (Sossi et al., 2016) and/or during accretionary impacts (see Poitrasson et al., 2004). Sossi et al. (2016) proposed an effect linked to differential condensation of solids in the protoplanetary accretion disk on the basis of a positive correlation between $\delta^{57}\text{Fe}$ and Fe/Mn estimates of planetary bodies (Fig. 4a), except chondrites. However, correlations between elemental ratios of contrasted volatilities (Rb/Sr or Mn/Na) and $\delta^{57}\text{Fe}$ would rather suggest two different histories for the Earth, the Moon and the angrite parent body on one side and Mars, Vesta and the chondrite (CI) parent body on the other (Fig. 4b and c). Since there are no reasons to consider that Mars, Vesta and the chondrite parent body chemistries were also unaffected by the effect of differential solid condensation of the protoplanetary accretion disk, the dichotomy observed in Figs. 4b and c suggests that Fe isotopes record different accretion mechanisms. The lack of Fe isotope variation for Mars and Vesta relative to the chondrite (CI) parent body would imply accretionary processes through runaway growth, whereas the Earth, Moon and angrite parent body would be additionally affected by high-energy impacts towards the end of their accretion history (Poitrasson et al., 2004; Wang et al., 2012). This would produce planetary bodies enriched in heavy Fe isotopes through the loss of light Fe isotopes that do not necessarily require percent-level Fe loss, as computed by Poitrasson et al., (2004), potentially leading to measurable Fe isotope variations. This is in contrast to the effect of differential condensation in the protoplanetary accretion disk that would primarily generate changes in elemental

abundances, potentially accompanied by Fe isotope variations (Sossi et al., 2016). According to the dynamic and thermodynamic calculation of Dauphas et al. (2015), such a loss of light isotopes to space in the aftermath of interplanetary impacts should be difficult for small parent bodies of the size of the angrite parent body. However, more recent calculations involving the track motion of Jupiter generating gas drag from recently impacted bodies make this hypothesis feasible (Hin et al., 2017), although the calculations were conducted down to bodies only having twice the size of the angrite parent body (<500 km). More observational constraints are thus required to refine and reconcile these models.

5.5 Evaluation of the model with other stable isotope systems

There has been a growing body of mass-dependent stable isotope planetary studies in the recent years that should be compared to the Fe isotope systematics to produce, as far as possible, a scenario taking into account all available observations. Humayun and Clayton (1995) failed to find K isotope fractionation among planetary bodies at the then-achievable levels of precision of $\pm\sim 0.5\%$ at the 95% confidence level. Two decades later and using advanced plasma source mass spectrometry techniques, Wang and Jacobsen (2016) determined that lunar rocks are higher by $\sim 0.4\%$ in $\delta^{41}\text{K}$ relative to terrestrial samples and chondrites. They interpreted this new observation as tracking the effect of a high-energy, high angular momentum giant impact in a thick gas atmosphere surrounding the impacted proto-Earth (Lock et al., 2018) rather than a low-energy disk equilibration model (Pahlevan and Stevenson, 2007).

The stable isotope compositions of volatile elements Ga (half-mass condensation temperature T_c of 968K, Lodders, 2003), Rb ($T_c = 800\text{K}$, Lodders, 2003) and Zn ($T_c = 726\text{K}$; Lodders, 2003) also yielded isotopically heavier values consistent with a giant impact Moon-

forming scenario, associated with volatile loss (Paniello et al., 2012; Kato et al., 2015; Kato et al., 2017; Pringle and Moynier, 2017). However, as the database grows, the lunar rocks appear to be extremely heterogeneous, notably anorthosites, yielding for example a $\delta^{66}\text{Zn}$ range of more than 15‰ (Paniello et al., 2012; Kato et al., 2015). By comparing the stable isotope composition of the variably volatile elements K, Rb and Zn, Pringle and Moynier (2017) noted that there is no clear systematics between the planetary isotope differences and the geochemical properties of these elements. This makes a detailed discussion, besides the general idea of a volatility effect explaining the high $\delta^{41}\text{K}$, $\delta^{71}\text{Ga}$, $\delta^{87}\text{Rb}$ and $\delta^{66}\text{Zn}$ for some lunar rocks relative to other planetary samples rather difficult. Clearly, more work is needed to understand the igneous systematics of these stable isotope systems, as already shown with the comparatively more studied Fe isotope systematics (see above).

There is no consensus yet as to whether the majority of terrestrial bodies have a chondritic $\delta^{26}\text{Mg}$ signature (see recent review by Teng, 2017, and see also Sedaghatpour and Jacobsen, 2019), or if the Earth is isotopically heavier (Wiechert and Halliday, 2007; Hin et al., 2017). This partly results from analytical difficulties to determine accurate $\delta^{26}\text{Mg}$ values of silicate rock samples. Furthermore, lunar samples appear to yield heterogeneous $\delta^{26}\text{Mg}$ values, with high-Ti basalts tending towards lighter isotopic compositions according to Sedaghatpour et al. (2013) and Sedaghatpour and Jacobsen (2019) whereas Wiechert and Halliday (2007) found the opposite. Given these controversies, it is still difficult at this point to use Mg isotope compositions to discuss planet formation and differentiation processes.

Silicon was another element that initially generated debates for analytical reasons (Georg et al., 2007; Fitoussi et al., 2009). Further, magma differentiation or the nature of protoliths may affect $\delta^{30}\text{Si}$ values of igneous rocks (Savage et al., 2014), including on the Moon (Poitrasson and Zambardi, 2015), although no difference between low-Ti and high-Ti

674 mare basalts was observed (Armytage et al., 2012; Fitoussi and Bourdon, 2012; Zambardi et
675 al., 2013). However, most authors now agree on the relative $\delta^{30}\text{Si}$ differences between
676 meteorite types. They also concur that the Earth and the Moon are isotopically
677 indistinguishable, with $\delta^{30}\text{Si}$ close to -0.29‰ relative to NBS-28 reference material). These
678 two bodies are also isotopically heavier than Mars, Vesta and chondrites, but lighter than
679 angrites (see recent review in Poitrasson, 2017).

680 Lithium is an element seemingly not easy to use for interplanetary comparisons as it
681 may potentially show strong $\delta^7\text{Li}$ variations at the intra-mineral scale caused by thermal
682 diffusion or fluid circulations (see recent reviews in Tomascak et al., 2016 and Penniston-
683 Dorland et al., 2017). Yet resolved $\delta^7\text{Li}$ variations were observed for low-Ti versus high-Ti
684 mare basalts (Magna et al., 2006; Seitz et al., 2006; Day et al., 2016). Low-Ti basalts, the
685 Earth's mantle (Jeffcoate et al., 2007; Magna et al., 2008; Tomascak et al., 2008; Marschall et
686 al., 2017), Mars (Magna et al., 2015) and Vesta (Magna et al., 2014) all carry broadly similar
687 $\delta^7\text{Li}$ signatures at ~3–4‰.

688 Pilot studies of mass-dependent isotopic fractionations of $\delta^{87}\text{Sr}$ (Moynier et al., 2010)
689 and $\delta^{44}\text{Ca}$ (Simon and DePaolo, 2010) do not seem to reveal significant planetary differences
690 either. This could be observed besides large stable isotope variations of chondritic samples
691 (Simon and DePaolo, 2010; Charlier et al., 2012) and an effect of differentiation for igneous
692 terrestrial rocks for $\delta^{87}\text{Sr}$ (Charlier et al., 2012). This effect of igneous differentiation was
693 also observed for mass-dependent $\delta^{49}\text{Ti}$ and $\delta^{53}\text{Cr}$ compositions on Earth, along with a
694 dichotomy between high- and low-Ti lunar basalts (Bonnand et al., 2016a; Millet et al., 2016).
695 Yet, the authors of these studies found similar Earth-Moon stable isotope compositions. More
696 recently, Sossi et al. (2018) produced more lunar sample $\delta^{53}\text{Cr}$ values that led to an even more
697 scattered data set. After data filtering, they found an isotopically lighter Cr signature of the
698 Moon relative to the Earth, potentially reflecting volatile loss of oxygenated Cr species from

the Moon towards the end of the lunar accretion history, at the lunar magma ocean stage. Chondrites and achondrites $\delta^{53}\text{Cr}$ are heterogeneous, possibly as a result of magmatic differentiation for the latter (Bonnand et al., 2016b). Nevertheless, they seem to share the $\delta^{53}\text{Cr}$ signature of the Earth overall (Schoenberg et al., 2016).

To summarize, only iron and silicon have so far been found to display a mass-dependent enrichment towards heavy isotope composition of the Earth–Moon system relative to other planetary bodies. Importantly, and in contrast to the other elements considered, Fe and Si are likely the most abundant elements of the Earth’s core, besides Ni (e.g., Allègre et al., 2001; McDonough, 2014), even though estimates for silicon remain more uncertain (e.g., Hirose et al., 2013; Zambardi et al., 2013; Dauphas et al., 2015). As discussed above, metal–silicate partitioning during mantle core–differentiation is unlikely to be the major cause to explain the observed planetary Fe isotope compositions. Hence, it would be preferable, until proof of the contrary is provided, if all those isotopic observations were explained by a single process that some authors (Poitrasson et al., 2004; Zambardi et al., 2013) have proposed to interpret the estimated Fe and Si isotopic composition of planets: the partial planetary melting and vaporization associated with the Moon-forming giant impact. Experimental work has shown that the vaporization rate of a metal species from liquid metal is much larger compared to that of an oxide from a silicic melt (Wang et al., 1994). Therefore, at a given temperature of an interplanetary impact-induced vaporization, elemental Fe and Si evaporation from a metallic melt will be much more efficient than evaporation of respective oxides from the coexisting silicic melt, and thus more likely to leave a stable isotope imprint through the loss of lighter Fe and Si isotopes in space (Poitrasson et al., 2004; Zambardi et al., 2013). Numerical simulation of a Moon-forming giant impact tend to exclude models yielding a too Fe-rich outer disk to take into account the estimated lower bulk Fe content of the Moon relative to the Earth (Canup, 2004; Cuk and Stewart, 2012; Reufer et al., 2012; Canup et al., 2013).

However, as mentioned above, the amount of Fe in the lunar-forming disk does not need to be high to account for the observed mass-dependent Fe and Si isotope systematics since previous quantitative estimates showed that less than 1% of iron and silicon loss from the Earth–Moon system into space is required in this context to explain their observed heavy isotopic compositions relative to other planetary bodies and chondrites (Poitrasson et al., 2004; Zambardi et al., 2013).

6. Conclusions

The new Fe isotope data for various lunar rocks clearly show isotopically distinct reservoirs. This is apparent from data for low-Ti and high-Ti mare basalts that have means significantly different to a degree of confidence well beyond 99%. Partial melting in the lunar mantle and subsequent magmatic evolution of these mafic melts probably did not induce Fe isotope fractionation. Rather, this sharp difference in Fe isotope composition is likely due to lunar mantle sources having different mineralogical compositions, with heavy Fe isotope composition of high-Ti basalts likely reflecting the occurrence of ilmenite. The highland rocks, that include ferroan anorthosites and Mg-suite rocks, also reveal a much larger isotopic scatter than previously found, of about 1.2‰. The finding of large $\delta^{57}\text{Fe}$ differences among different CAPTEM aliquot allocations of the same samples suggest that the common 0.1–1 g sample size allocations is insufficient to produce a powder representative of these coarse-grained rocks for Fe isotope measurements. It seems that feldspars tend towards heavy $\delta^{57}\text{Fe}$ whereas olivines tend towards light $\delta^{57}\text{Fe}$ values, possibly as a result of diffusive processes for the latter. Excluding these effects with highland rocks having more than 10% pyroxene modal content leads to a more precise highland $\delta^{57}\text{Fe}$ of $0.094 \pm 0.035\text{‰}$. This value, which is based on six Mg-suite rocks but that is also within uncertainty reflected by three anorthosites

and one troctolite, is indistinguishable from the mean upper lunar mantle value defined using the Fe isotope systematics of mare basalts. It likely represents the bulk Moon $\delta^{57}\text{Fe}$, which also agrees within uncertainties with previous estimates for the bulk Earth.

Combined with literature data, this Earth-like Fe isotope composition of the Moon, heavier by $\sim 0.1\%$ relative to chondrite parent bodies, Mars and Vesta, can be best explained as fingerprinting the Moon-forming giant impact. This process alone provides the most straightforward scenario to explain the observed isotopic pattern among planets, without necessarily requiring a high-pressure, high-temperature metal-silicate stable isotope fractionation event.

Acknowledgements – This study was made possible through three separate allocations of lunar samples to CRN, FP and TM by CAPTEM. Carole Boucayrand, Jonathan Prunier and Manuel Henry are thanked for maintaining the clean lab in excellent condition, as well as Rémi Freydier and Jérôme Chmeleff for their efforts to have the Neptune MC-ICP-MS up running. This manuscript was substantially revised and expanded following detailed comments from Steve Elardo, two anonymous referees and GCA AE Munir Humayun. This work was funded by a grant from the Programme National de Planétologie (PNP) of CNRS-INSU to FP. CRN was supported by NASA grant NASA-NNX15AH76G.

References

Allègre, C.J., Manhès, G. and Lewin, E. (2001) Chemical composition of the Earth and the volatility control on planetary genetics. *Earth Planet. Sci. Lett.* 185, 49-69.

773 Armytage, R.M.G., Georg, R.B., Williams, H.M. and Halliday, A.N. (2012) Silicon isotopes
 774 in lunar rocks: Implications for the Moon's formation and the early history of the Earth.
 775 *Geochim. Cosmochim. Acta* 77, 504-514.

776 Beard, B.L. and Johnson, C.M. (2004) Inter-mineral Fe isotope variations in mantle-derived
 777 rocks and implications for the Fe geochemical cycle. *Geochim. Cosmochim. Acta* 68,
 778 4727-4743.

779 Beard, B.L. and Johnson, C.M. (2006) Comment on "Heavy iron isotope composition of
 780 granites determined by high resolution MC-ICP-MS" by F. Poitrasson and R. Freydier,
 781 *Chemical Geology*, volume 222, pages 132-147. *Chem. Geol.* 235, 201-204.

782 Beard, B.L. and Johnson, C.M. (2007) Comment on "Iron isotope fractionation during
 783 planetary differentiation" by S. Weyer et al., *Earth Planet. Sci. Lett.* V240, pages 251-264.
 784 *Earth Planet. Sci. Lett.* 256, 633-637.

785 Beard, B.L., Taylor, L.A., Scherer, E.E., Johnson, C.M. and Snyder, G.A. (1998) The source
 786 region and melting mineralogy of high-titanium and low-titanium lunar basalts deduced
 787 from Lu-Hf isotope data. *Geochim. Cosmochim. Acta* 62, 525-544.

788 Bennett, V.C., Brandon, A.D. and Nutman, A.P. (2007) Coupled Nd-142-Nd-143 isotopic
 789 evidence for Hadean mantle dynamics. *Science* 318, 1907-1910.

790 Blanchard, M., Poitrasson, F., Méheut, M., Lazzeri, M., Mauri, F. and Balan, E. (2009) Iron
 791 isotope fractionation between pyrite (FeS₂), hematite (Fe₂O₃) and siderite (FeCO₃): A first-
 792 principles density functional theory study. *Geochim. Cosmochim. Acta* 73, 6565-6578.

793 Blanchard, M., Poitrasson, F., Méheut, M., Lazzeri, M., Mauri, F. and Balan, E. (2012)
 794 Comment on "New data on equilibrium iron isotope fractionation among sulfides:
 795 Constraints on the mechanisms of sulfide formation in hydrothermal and igneous systems"
 796 by V.B. Polyakov and D.M. Soultanov. *Geochim. Cosmochim. Acta* 87, 356-359.

797 Bonnard, P., Parkinson, I.J., Anand, M., 2016a. Mass dependent fractionation of stable
798 chromium isotopes in mare basalts: Implications for the formation and the differentiation
799 of the Moon. *Geochimica Et Cosmochimica Acta*, 175: 208-221.

800 Bonnard, P., Williams, H.M., Parkinson, I.J., Wood, B.J., Halliday, A.N., 2016b. Stable
801 chromium isotopic composition of meteorites and metal-silicate experiments: Implications
802 for fractionation during core formation. *Earth and Planetary Science Letters*, 435: 14-21.

803 Bourdon, B., Roskosz, M. and Hin, R.C. (2018) Isotope tracers of core formation. *Earth-Sci.*
804 *Rev.* 181, 61-81.

805 Bourdon, B., Tipper, E.T., Fitoussi, C. and Stracke, A. (2010) Chondritic Mg isotope
806 composition of the Earth. *Geochim. Cosmochim. Acta* 74, 5069-5083.

807 Canup, R.M., 2004. Simulations of a late lunar-forming impact. *Icarus*, 168(2): 433-456.

808 Canup, R.M., Barr, A.C., Crawford, D.A., 2013. Lunar-forming impacts: High-resolution
809 SPH and AMR-CTH simulations. *Icarus*, 222(1): 200-219.

810 Chakrabarti, R. and Jacobsen, S.B. (2010) The isotopic composition of magnesium in the
811 inner Solar System. *Earth Planet. Sci. Lett.* 293, 349-358.

812 Charlier, B.L.A., Nowell, G.M., Parkinson, I.J., Kelley, S.P., Pearson, D.G. and Burton, K.W.
813 (2012) High temperature strontium stable isotope behaviour in the early solar system and
814 planetary bodies. *Earth Planet. Sci. Lett.* 329, 31-40.

815 Chernozhukhin, S.M., Goderis, S., Costas-Rodriguez, M., Claeys, P. and Vanhaecke, F.
816 (2016) Effect of parent body evolution on equilibrium and kinetic isotope fractionation: a
817 combined Ni and Fe isotope study of iron and stony-iron meteorites. *Geochim.*
818 *Cosmochim. Acta* 186, 168-188.

819 Chernozhukhin, S.M., Weyrauch, M., Goderis, S., Oeser, M., McKibbin, S.J., Horn, I., Hecht,
820 L., Weyer, S., Claeys, P. and Vanhaecke, F. (2017) Thermal equilibration of iron meteorite

821 and pallasite parent bodies recorded at the mineral scale by Fe and Ni isotope systematics.
822 *Geochim. Cosmochim. Acta* 217, 95-111.

823 Clayton, R.N., Hurd, J.M. and Mayeda, T.K. (1971) Oxygen isotopic compositions of Apollo
824 15, 16 and 17 samples, and their bearing on lunar origin and petrogenesis. *Proceedings of*
825 *the Fourth Lunar and Planetary Science Conference* 4, 1417-1420.

826 Clayton, R.N. and Mayeda, T.K. (1983) Oxygen isotopes in eucrites, shergottites, nakhlites,
827 and chassignites. *Earth Planet. Sci. Lett.* 62, 1-6.

828 Collinet, M., Charlier, B., Namur, O., Oeser, M., Medard, E. and Weyer, S. (2017)
829 Crystallization history of enriched shergottites from Fe and Mg isotope fractionation in
830 olivine megacrysts. *Geochim. Cosmochim. Acta* 207, 277-297.

831 Consolmagno, G.J. and Drake, M.J. (1977) Composition and evolution of the eucrite parent
832 body: Evidence from rare earth elements. *Geochim. Cosmochim. Acta* 41, 1271-1282.

833 Craddock, P.R. and Dauphas, N. (2011) Iron isotope compositions of reference material,
834 geostandards and chondrites. *Geostandards and Geoanalytical Research* 35, 101-123.

835 Craddock, P.R., Dauphas, N. and Clayton, R.N. (2010) Mineralogical control on iron isotopic
836 fractionation during lunar differentiation and magmatism, *LPSC XXXXI*, Houston, p.
837 1230.pdf.

838 Craddock, P.R., Warren, J.M. and Dauphas, N. (2013) Abyssal peridotites reveal the near-
839 chondritic Fe isotopic composition of the Earth. *Earth Planet. Sci. Lett.* 365, 63-76.

840 Cuk, M., Stewart, S.T., 2012. Making the Moon from a Fast-Spinning Earth: A Giant Impact
841 Followed by Resonant Despinning. *Science*, 338(6110): 1047-1052.

842 Dauphas, N., Craddock, P.R., Asimow, P.D., Bennett, V.C., Nutman, A.P. and Ohnenstetter,
843 D. (2009) Iron isotopes may reveal the redox conditions of mantle melting from Archean
844 to Present. *Earth Planet. Sci. Lett.* 288, 255-267.

845 Dauphas, N., John, S.G. and Rouxel, O. (2017) Iron Isotope Systematics, in: Teng, F.Z.,
846 Watkins, J., Dauphas, N. (Eds.), Non-Traditional Stable Isotopes. Mineralogical Soc Amer
847 & Geochemical Soc, Chantilly, pp. 415-510.

848 Dauphas, N., Poitrasson, F., Burkhardt, C., Kobayashi, H. and Kurosawa, K. (2015) Planetary
849 and meteoritic Mg/Si and delta Si-30 variations inherited from solar nebula chemistry.
850 Earth Planet. Sci. Lett. 427, 236-248.

851 Dauphas, N., Burkhardt, C., Warren, P.H. and Teng, F.Z. (2014b) Geochemical arguments for
852 an Earth-like Moon-forming impactor. Philosophical Transactions of the Royal Society A-
853 Mathematical, Physical & Engineering Science 372, 20130244.

854 Dauphas, N., Roskosz, M., Alp, E.E., Neuville, D.R., Hu, M.Y., Sio, C.K., Tissot, F.L.H.,
855 Zhao, J., Tissandiere, L., Medard, E. and Cordier, C. (2014a) Magma redox and structural
856 controls on iron isotope variations in Earth's mantle and crust. Earth Planet. Sci. Lett. 398,
857 127-140.

858 Day, J.M.D., Qiu, L., Ash, R.D., McDonough, W.F., Teng, F.Z., Rudnick, R.L. and Taylor,
859 L.A. (2016) Evidence for high-temperature fractionation of lithium isotopes during
860 differentiation of the Moon. Meteoritics & Planetary Science 51, 1046-1062.

861 Debaille, V., O'Neill, C., Brandon, A.D., Haenecour, P., Yin, Q.Z., Mattielli, N. and Treiman,
862 A.H. (2013) Stagnant-lid tectonics in early Earth revealed by Nd-142 variations in late
863 Archean rocks. Earth Planet. Sci. Lett. 373, 83-92.

864 Dymek, R.F., Albee, A.L. and Chodos, A.A. (1975) Comparative petrology of lunar cumulate
865 rocks of possible primary origin: Dunite 72415, troctolite 76535, norite 78235 and
866 anorthosite 62237, Lunar and Planetary Science Conference, 6th. LPI, Houston, pp. 301-
867 341.

868 Elardo, S.M. and Shahr, A. (2017) Non-chondritic iron isotope ratios in planetary mantles as
869 a result of core formation. Nat. Geosci. 10, 317-321.

870 Elardo, S.M., Shahar, A., Mock, T.D. and Sio, C.K. (2019) The effect of core composition on
 871 iron isotope fractionation between planetary cores and mantles. *Earth Planet. Sci. Lett.*
 872 513, 124-134.

873 Elkins-Tanton, L.T., Burgess, S. and Yin, Q.Z. (2011) The lunar magma ocean: Reconciling
 874 the solidification process with lunar petrology and geochronology. *Earth Planet. Sci. Lett.*
 875 304, 326-336.

876 Fitoussi, C., Bourdon, B., Kleine, T., Oberli, F. and Reynolds, B.C. (2009) Si isotope
 877 systematics of meteorites and terrestrial peridotites: implications for Mg/Si fractionation in
 878 the solar nebula and for Si in the Earth's core. *Earth Planet. Sci. Lett.* 287, 77-85.

879 Foden, J., Sossi, P.A. and Wawryk, C.M. (2015) Fe isotopes and the contrasting petrogenesis
 880 of A-, I- and S-type granite. *Lithos* 212, 32-44.

881 Georg, R.B., Halliday, A.N., Schauble, E.A. and Reynolds, B.C. (2007) Silicon in the Earth's
 882 core. *Nature* 447, 1102-1106.

883 Giguere, T.A., Taylor, G.J., Hawke, B.R. and Lucey, P.G. (2000) The titanium contents of
 884 lunar mare basalts. *Meteoritics & Planetary Science* 35, 193-200.

885 Head, J.W. (1976) Lunar volcanism in space and time. *Reviews of Geophysics* 14, 265-300.

886 Heimann, A., Beard, B.L. and Johnson, C.M. (2008) The role of volatile exsolution and sub-
 887 solidus fluid/rock interactions in producing high Fe-56/Fe-54 ratios in siliceous igneous
 888 rocks. *Geochim. Cosmochim. Acta* 72, 4379-4396.

889 Hin, R.C., Coath, C.D., Carter, P.J., Nimmo, F., Lai, Y.J., von Strandmann, P., Willbold, M.,
 890 Leinhardt, Z.M., Walter, M.J. and Elliott, T. (2017) Magnesium isotope evidence that
 891 accretional vapour loss shapes planetary compositions. *Nature* 549, 511.

892 Hin, R.C., Schmidt, M.W. and Bourdon, B. (2012) Experimental evidence for the absence of
 893 iron isotope fractionation between metal and silicate liquids at 1 GPa and 1250-1300°C
 894 and its cosmochemical consequences. *Geochim. Cosmochim. Acta* 93, 164-181.

895 Hirose, K., Labrosse, S. and Hernlund, J. (2013) Composition and State of the Core. Annual
 896 Review of Earth and Planetary Sciences 41, 657-691.

897 Humayun, M. and Clayton, R.N. (1995) Potassium isotope cosmochemistry: Genetic
 898 implications of volatile element depletion. *Geochim. Cosmochim. Acta* 59, 2131-2148.

899 James, O.B., Lindstrom, M.M. and McGee, J.J. (1991) Lunar ferroan anorthosite-60025 -
 900 petrology and chemistry of mafic lithologies. *Proceedings of Lunar and Planetary Science*
 901 21, 63-87.

902 Jeffcoate, A.B., Elliott, T., Kasemann, S.A., Ionov, D., Cooper, K. and Brooker, R. (2007) Li
 903 isotope fractionation in peridotites and mafic melts. *Geochim. Cosmochim. Acta* 71, 202-
 904 218.

905 Jerde, E.A., Snyder, G.A., Taylor, L.A., Liu, Y.G. and Schmitt, R.A. (1994) The origin and
 906 evolution of lunar High-Ti basalts - periodic melting of a single-source at Mare
 907 Tranquillitatis. *Geochim. Cosmochim. Acta* 58, 515-527.

908 Jordan, M.K., Tang, H.L., Kohl, I.E. and Young, E.D. (2019) Iron isotope constraints on
 909 planetesimal core formation in the early solar system. *Geochim. Cosmochim. Acta* 246,
 910 461-477.

911 Kato, C., Moynier, F., Valdes, M.C., Dhaliwal, J.K. and Day, J.M.D. (2015) Extensive
 912 volatile loss during formation and differentiation of the Moon. *Nat. Commun.* 6, 4.

913 Kato, C. and Moynier, F. (2017) Gallium isotopic evidence for extensive volatile loss from
 914 the Moon during its formation. *Sci. Adv.* 3, 5.

915 Liu, J., Dauphas, N., Roskosz, M., Hu, M.Y., Yang, H., Bi, W.L., Zhao, J.Y., Alp, E.E., Hu,
 916 J.Y. and Lin, J.F. (2017) Iron isotopic fractionation between silicate mantle and metallic
 917 core at high pressure. *Nat. Commun.* 8, 6.

918 Liu, Y., Spicuzza, M.J., Craddock, P.R., Day, J.M.D., Valley, J.W., Dauphas, N. and Taylor,
 919 L.A. (2010) Oxygen and iron isotope constraints on near-surface fractionation effects and

920 the composition of lunar mare basalt source regions. *Geochim. Cosmochim. Acta* 74,
 921 6249-6262.

922 Lodders, K., 2003. Solar system abundances and condensation temperatures of the elements.
 923 *Astrophysical Journal*, 591(2): 1220-1247.

924 Lock, S.J., Stewart, S.T., Petaev, M.I., Leinhardt, Z., Mace, M.T., Jacobsen, S.B., Cuk, M.,
 925 2018. The Origin of the Moon Within a Terrestrial Synestia. *Journal of Geophysical*
 926 *Research-Planets*, 123(4): 910-951.

927 Longhi, J. (1992) Experimental Petrology and Petrogenesis of Mare Volcanics. *Geochim.*
 928 *Cosmochim. Acta* 56, 2235-2251.

929 Magna, T., Wiechert, U. and Halliday, A.N. (2006) New constraints on the lithium isotope
 930 compositions of the Moon and terrestrial planets. *Earth Planet. Sci. Lett.* 243, 336-353.

931 Magna, T., Ionov, D.A., Oberli, F. and Wiechert, U. (2008) Links between mantle
 932 metasomatism and lithium isotopes: Evidence from glass-bearing and cryptically
 933 metasomatized xenoliths from Mongolia. *Earth Planet. Sci. Lett.* 276, 214-222.

934 Magna, T., Simcikova, M. and Moynier, F. (2014) Lithium systematics in howardite-eucrite-
 935 diogenite meteorites: Implications for crust-mantle evolution of planetary embryos.
 936 *Geochim. Cosmochim. Acta* 125, 131-145.

937 Magna, T., Day, J.M.D., Mezger, K., Fehr, M.A., Dohmen, R., Aoudjehane, H.C. and Agee,
 938 C.B. (2015) Lithium isotope constraints on crust-mantle interactions and surface processes
 939 on Mars. *Geochim. Cosmochim. Acta* 162, 46-65.

940 Marschall, H.R., Wanless, V.D., Shimizu, N., von Strandmann, P., Elliott, T. and Monteleone,
 941 B.D. (2017) The boron and lithium isotopic composition of mid-ocean ridge basalts and
 942 the mantle. *Geochim. Cosmochim. Acta* 207, 102-138.

943 McDonough, W.F. (2014) Compositional model for the Earth's core, in: Carlson, R.W. (Ed.),
 944 *The mantle and core*, 2nd ed. Elsevier, Amsterdam, pp. 559-576.

945 McSween Jr, H.Y., Taylor, L.A. and Stolper, E.M. (1979) Allan Hills 77005: A new meteorite
 946 type found in Antarctica. *Science* 204, 1201-1203.

947 Meyer, C. (2011) Lunar sample compendium. NASA, Houston, p.
 948 <http://curator.jsc.nasa.gov/lunar/compendium.cfm>.

949 Miller, J.C. and Miller, J.N. (1993) *Statistics for analytical chemistry*, 3rd ed. Ellis Horwood,
 950 New York.

951 Millet, M.A., Dauphas, N., Greber, N.D., Burton, K.W., Dale, C.W., Debret, B., Macpherson,
 952 C.G., Nowell, G.M., Williams, H.M., 2016. Titanium stable isotope investigation of
 953 magmatic processes on the Earth and Moon. *Earth and Planetary Science Letters*, 449:
 954 197-205.

955 Molini-Velsko, C., Mayeda, T.K. and Clayton, R.N. (1986) isotopic composition of silicon in
 956 meteorites. *Geochim. Cosmochim. Acta* 50, 2719-2726.

957 Moynier, F., Agranier, A., Hezel, D.C. and Bouvier, A. (2010) Sr stable isotope composition
 958 of Earth, the Moon, Mars, Vesta and meteorites. *Earth Planet. Sci. Lett.* 300, 359-366.

959 Moynier, F., Albarede, F. and Herzog, G.F. (2006) Isotopic composition of zinc, copper, and
 960 iron in lunar samples. *Geochim. Cosmochim. Acta* 70, 6103-6117.

961 Nakamura, Y. (1983) Seismic velocity structure of the lunar mantle. *J. Geophys. Res.* 88.

962 Neal, C.R. (2001) Interior of the Moon: The presence of garnet in the primitive deep lunar
 963 mantle. *J. Geophys. Res.-Planets* 106, 27865-27885.

964 Neal, C.R. and Taylor, L.A. (1992) Petrogenesis of Mare Basalts - a Record of Lunar
 965 Volcanism. *Geochim. Cosmochim. Acta* 56, 2177-2211.

966 Oeser, M., Dohmen, R., Horn, I., Schuth, S. and Weyer, S. (2015) Processes and time scales
 967 of magmatic evolution as revealed by Fe-Mg chemical and isotopic zoning in natural
 968 olivines. *Geochim. Cosmochim. Acta* 154, 130-150.

969 Pahlevan, K. and Stevenson, D.J. (2007) Equilibration in the aftermath of the lunar-forming
 970 giant impact. *Earth Planet. Sci. Lett.* 262, 438-449.

971 Papike, J.J., Ryder, G. and Shearer, C.K. (1998) Lunar samples, in: Papike, J.J. (Ed.),
 972 Planetary Materials. Mineralogical Society of America, Washington, pp. 5-1:5-234.

973 Paniello, R.C., Day, J.M.D. and Moynier, F. (2012) Zinc isotopic evidence for the origin of
 974 the Moon. *Nature* 490, 376-380.

975 Penniston-Dorland, S., Liu, X.M. and Rudnick, R.L. (2017) Lithium Isotope Geochemistry,
 976 in: Teng, F.Z., Watkins, J., Dauphas, N. (Eds.), Non-Traditional Stable Isotopes.
 977 Mineralogical Soc Amer & Geochemical Soc, Chantilly, pp. 165-217.

978 Platzner, I.T. (1997) Modern isotope ratio mass spectrometry. John Wiley & sons, Chichester,
 979 514 p.

980 Poitrasson, F. (2006) On the iron isotope homogeneity level of the continental crust. *Chem.*
 981 *Geol.* 235, 195-200.

982 Poitrasson, F. (2007) Does planetary differentiation really fractionate iron isotopes? *Earth*
 983 *Planet. Sci. Lett.* 256, 484-492.

984 Poitrasson, F. (2009) Probes of the ancient and the inaccessible. *Science* 323, 882-883.

985 Poitrasson, F. (2017) Silicon isotope geochemistry, in: Teng, F.Z., Watkins, J.M., Dauphas,
 986 N. (Ed.), Non-traditional stable isotopes. Mineralogical Society of America, Washington,
 987 D.C., pp. 289-344.

988 Poitrasson, F., Delpech, G. and Grégoire, M. (2013) On the iron isotope heterogeneity of
 989 lithospheric mantle xenoliths: implications for mantle metasomatism, the origin of basalts
 990 and the iron isotope composition of the Earth. *Contrib. Mineral. Petrol.* 165, 1243-1258.

991 Poitrasson, F. and Freydier, R. (2005) Heavy iron isotope composition of granites determined
 992 by high resolution MC-ICP-MS. *Chem. Geol.* 222, 132-147.

993 Poitrasson, F., Halliday, A.N., Lee, D.C., Levasseur, S. and Teutsch, N. (2004) Iron isotope
994 differences between Earth, Moon, Mars and Vesta as possible records of contrasted
995 accretion mechanisms. *Earth Planet. Sci. Lett.* 223, 253-266.

996 Poitrasson, F., Levasseur, S. and Teutsch, N. (2005) Significance of iron isotope mineral
997 fractionation in pallasites and iron meteorites for the core-mantle differentiation of
998 terrestrial planets. *Earth Planet. Sci. Lett.* 234, 151-164.

999 Poitrasson, F., Roskosz, M. and Corgne, A. (2009) No iron isotope fractionation between
1000 molten alloys and silicate melt to 2000°C and 7.7 GPa: Experimental evidence and
1001 implications for planetary differentiation and accretion. *Earth Planet. Sci. Lett.* 278, 376-
1002 385.

1003 Poitrasson, F. and Zambardi, T. (2015) An Earth-Moon silicon isotope model to track silicic
1004 magma origins. *Geochim. Cosmochim. Acta* 167, 301-312.

1005 Polyakov, V.B. (2009) Equilibrium iron isotope fractionation at core-mantle boundary
1006 conditions. *Science* 323, 912-914.

1007 Polyakov, V.B. and Mineev, S.D. (2000) The use of Mössbauer spectroscopy in stable isotope
1008 geochemistry. *Geochim. Cosmochim. Acta* 64, 849-865.

1009 Polyakov, V.B., Osadchii, E.G., Chareev, D.A., Chumakov, A.I. and Sergeev, I.A. (2013) Fe
1010 beta-factors for sulfides from NRIXS synchrotron experiments, 2013 Goldschmidt
1011 Conference. *Mineralogical Magazine*, Prague, p. 1985.

1012 Potts, P.J. (1987) *A handbook of silicate rock analysis*. Blackie Academic and Professional,
1013 London, 622 p.

1014 Pringle, E.A., Moynier, F. (2017). Rubidium isotopic composition of the Earth, meteorites,
1015 and the Moon: Evidence for the origin of volatile loss during planetary accretion. *Earth and*
1016 *Planetary Science Letters*, 473: 62-70.

1017 Prissel, K.B., Krawczynski, M.J., Nie, N.X., Dauphas, N., Couvy, H., Hu, M.Y., Alp, E.E.
 1018 and Roskosz, M. (2018) Experimentally determined effects of olivine crystallization and
 1019 melt titanium content on iron isotopic fractionation in planetary basalts. *Geochim.*
 1020 *Cosmochim. Acta* 238, 580-598.

1021 Reufer, A., Meier, M.M.M., Benz, W., Wieler, R. (2012). A hit-and-run giant impact
 1022 scenario. *Icarus*, 221(1): 296-299.

1023 Richter, F.M., Chaussidon, M., Mendybaev, R.A. and Taylor, L.A. (2016) Magnesium
 1024 isotopic zoning of an olivine grain from lunar microgabbro 15555: Constraints on
 1025 crystallization and cooling, LPSC XXXXVII, Houston, p. 1146.pdf.

1026 Roskosz, M., Luais, B., Watson, H.C., Toplis, M.J., Alexander, C.M.O. and Mysen, B.O.
 1027 (2006) Experimental quantification of the fractionation of Fe isotopes during metal
 1028 segregation from a silicate melt. *Earth Planet. Sci. Lett.* 248, 851-867.

1029 Rustad, J.R. and Yin, Q.Z. (2009) Iron isotope fractionation in the Earth's lower mantle. *Nat.*
 1030 *Geosci.* 2, 514-518.

1031 Ryder, G. (1982) Lunar anorthosite 60025, the petrogenesis of lunar anorthosites, and the
 1032 composition of the Moon. *Geochim. Cosmochim. Acta* 46, 1591-1601.

1033 Ryder, G. (1985) Catalog of Apollo 15 rocks. Part 2. 15306-15468. NASA, Houston, pp. 339-
 1034 777.

1035 Ryder, G. and Schuraytz, B.C. (2001) Chemical variation of the large Apollo 15 olivine-
 1036 normative mare basalt rock samples. *J. Geophys. Res.-Planets* 106, 1435-1451.

1037 Ryder, G. and Steele, A. (1988) Chemical dispersion among Apollo 15 olivine-normative
 1038 mare basalts, Lunar and Planetary Science Conference, 18th. LPI, Houston, pp. 273-282.

1039 Sato, M. (1976) Oxygen fugacity and other thermochemical parameters of Apollo 17 high-Ti
 1040 basalts and their implications on the reduction mechanism. *Proceedings of the Seventh*
 1041 *Lunar and Planetary Science Conference* 7, 1323-1344.

1042 Sato, M., Hickling, N.L. and McLane, J.E. (1973) Oxygen fugacity values of Apollo 12, 14,
 1043 and 15 lunar samples and reduced state of lunar magmas. *Proceedings of the Fourth Lunar*
 1044 *and Planetary Science Conference* 4, 1061-1079.
 1045 Saunier, G., Poitrasson, F., Moine, B.N., Grégoire, M. and Seddiki, A. (2010) Effect of hot
 1046 desert weathering on the bulk-iron isotope composition of L6 and H5 ordinary chondrites.
 1047 *Meteorit. Planet. Sci.* 45, 195-209.
 1048 Savage, P.S., Armytage, R.M.G., Georg, R.B., Halliday, A.N., 2014. High temperature silicon
 1049 isotope geochemistry. *Lithos*, 190-191: 500-519.
 1050 Savage, P.S., Georg, R.B., Armytage, R.M.G., Williams, H.M. and Halliday, A.N. (2010)
 1051 Silicon isotope homogeneity in the mantle. *Earth Planet. Sci. Lett.* 295, 139-146.
 1052 Schauble, E.A. (2004) Applying stable isotope fractionation theory to new systems,
 1053 *Geochemistry of Non-Traditional Stable Isotopes*, pp. 65-111.
 1054 Schoenberg, R. and von Blanckenburg, F. (2006) Modes of planetary-scale Fe isotope
 1055 fractionation. *Earth Planet. Sci. Lett.* 252, 342-359.
 1056 Schoenberg, R., Merdian, A., Holmden, C., Kleinhanns, I.C., Hassler, K., Wille, M. and
 1057 Reitter, E. (2016) The stable Cr isotopic compositions of chondrites and silicate planetary
 1058 reservoirs. *Geochim. Cosmochim. Acta* 183, 14-30.
 1059 Schuessler, J., Schoenberg, R. and Sigmarsson, O. (2009) Iron and lithium isotope
 1060 systematics of the Hekla volcano, Iceland – Evidence for Fe isotope fractionation during
 1061 magma differentiation. *Chem. Geol.* 258, 78-91.
 1062 Sedaghatpour, F. and Jacobsen, S.B. (2019) Magnesium stable isotopes support the lunar
 1063 magma ocean cumulate remelting model for mare basalts. *Proc. Natl. Acad. Sci. U. S. A.*
 1064 116, 73-78.
 1065 Sedaghatpour, F., Teng, F.Z., Liu, Y., Sears, D.W.G., Taylor, L.A., 2013. Magnesium
 1066 isotopic composition of the Moon. *Geochimica et Cosmochimica Acta*, 120: 1-16.

1067 Seitz, H.M., Brey, G.P., Weyer, S., Durali, S., Ott, U., Munker, C., Mezger, K., 2006. Lithium
1068 isotope compositions of Martian and lunar reservoirs. *Earth and Planetary Science Letters*,
1069 245(1-2): 6-18.

1070 Shahar, A., Elardo, S.M. and Macris, C.A. (2017) Equilibrium Fractionation of Non-
1071 traditional Stable Isotopes: an Experimental Perspective, in: Teng, F.Z., Watkins, J.,
1072 Dauphas, N. (Eds.), *Non-Traditional Stable Isotopes*. Mineralogical Soc Amer &
1073 Geochemical Soc, Chantilly, vol. 82, pp. 65-83.

1074 Shahar, A., Hillgren, V.J., Horan, M.F., Mesa-Garcia, J., Kaufman, L.A. and Mock, T.D.
1075 (2015) Sulfur-controlled iron isotope fractionation experiments of core formation in
1076 planetary bodies. *Geochim. Cosmochim. Acta* 150, 253-264.

1077 Shearer, C.K., Hess, P.C., Wieczorek, M.A., Pritchard, M.E., Parmentier, E.M., Borg, L.E.,
1078 Longhi, J., Elkins-Tanton, L.T., Neal, C.R., Antonenko, I., Canup, R.M., Halliday, A.N.,
1079 Grove, T.L., Hager, B.H., Lee, D.C. and Wiechert, U. (2006) Thermal and magmatic
1080 evolution of the Moon, *New Views of the Moon*. MSA, pp. 365-518.

1081 Simon, J.I. and DePaolo, D.J. (2010) Stable calcium isotopic composition of meteorites and
1082 rocky planets. *Earth Planet. Sci. Lett.* 289, 457-466.

1083 Sio, C.K.I., Dauphas, N., Teng, F.Z., Chaussidon, M., Helz, R.T. and Roskosz, M. (2013)
1084 Discerning crystal growth from diffusion profiles in zoned olivine by in situ Mg-Fe
1085 isotopic analyses. *Geochim. Cosmochim. Acta* 123, 302-321.

1086 Snyder, G.A., Taylor, L.A. and Neal, C.R. (1992) A Chemical-Model for Generating the
1087 Sources of Mare Basalts - Combined Equilibrium and Fractional Crystallization of the
1088 Lunar Magmasphere. *Geochim. Cosmochim. Acta* 56, 3809-3823.

1089 Sossi, P.A. and Moynier, F. (2017) Chemical and isotopic kinship of iron in the Earth and
1090 Moon deduced from the lunar Mg-Suite. *Earth Planet. Sci. Lett.* 471, 125-135.

1091 Sossi, P.A., Nebel, O., Anand, M. and Poitrasson, F. (2016) On the iron isotope composition
 1092 of Mars and volatile depletion in the terrestrial planets. *Earth Planet. Sci. Lett.* 449, 360-
 1093 371.

1094 Spicuzza, M.J., Day, J.M.D., Taylor, L.A. and Valley, J.W. (2007) Oxygen isotope constraints
 1095 on the origin and differentiation of the Moon. *Earth Planet. Sci. Lett.* 253, 254-265.

1096 Taylor, L.A., Patchen, A., Mayne, R.G. and Taylor, D.H. (2004) The most reduced rock from
 1097 the moon, Apollo 14 basalt 14053: Its unique features and their origin. *Am. Mineral.* 89,
 1098 1617-1624.

1099 Telus, M., Dauphas, N., Moynier, F., Tissot, F.L.H., Teng, F.Z., Nabelek, P.I., Craddock, P.R.
 1100 and Groat, L.A. (2012) Iron, zinc, magnesium and uranium isotopic fractionation during
 1101 continental crust differentiation: The tale from migmatites, granitoids, and pegmatites.
 1102 *Geochim. Cosmochim. Acta* 97, 247-265.

1103 Teng, F.Z., 2017. Magnesium Isotope Geochemistry. In: Teng, F.Z., Watkins, J., Dauphas, N.
 1104 (Eds.), *Non-Traditional Stable Isotopes. Reviews in Mineralogy & Geochemistry.*
 1105 *Mineralogical Soc Amer & Geochemical Soc, Chantilly*, pp. 219-287.

1106 Teng, F.Z., Dauphas, N. and Helz, R.T. (2008) Iron isotope fractionation during magmatic
 1107 differentiation in Kilauea Iki Lava Lake. *Science* 320, 1620-1622.

1108 Teng, F.Z., Li, W.Y., Ke, S., Marty, B., Dauphas, N., Huang, S.C., Wu, F.Y. and Pourmand,
 1109 A. (2010) Magnesium isotopic composition of the Earth and chondrites. *Geochim.*
 1110 *Cosmochim. Acta* 74, 4150-4166.

1111 Tomascak, P.B., Langmuir, C.H., le Roux, P.J. and Shirey, S.B. (2008) Lithium isotopes in
 1112 global mid-ocean ridge basalts. *Geochim. Cosmochim. Acta* 72, 1626-1637.

1113 Tomascak, P.B., Magna, T. and Dohmen, R. (2016) *Advances in Lithium isotope*
 1114 *geochemistry.* Springer Verlag, 195 p.

1115 Torcivia, M.A. and Neal, C.R. (2017) Ferroan Anorthosite 60025: Magma ocean product -
 1116 and more!, LPSC XXXXVIII, Houston, p. 1480.pdf.

1117 Torcivia, M.A. and Neal, C.R. (2018) Ferroan Anorthosite 60025: A lunar breccia, LPSC
 1118 XXXXIX, Houston, p. 1331.pdf.

1119 Touboul, M., Puchtel, I.S. and Walker, R.J. (2012) W-182 Evidence for Long-Term
 1120 Preservation of Early Mantle Differentiation Products. *Science* 335, 1065-1069.

1121 Wang, K. and Jacobsen, S.B. (2016) Potassium isotopic evidence for a high-energy giant
 1122 impact origin of the Moon. *Nature* 538, 487.

1123 Wang, K., Jacobsen, S.B., Sedaghatpour, F., Chen, H. and Korotev, R.L. (2015) The earliest
 1124 Lunar Magma Ocean differentiation recorded in Fe isotopes. *Earth Planet. Sci. Lett.* 430,
 1125 202-208.

1126 Wang, K., Moynier, F., Dauphas, N., Barrat, J.A., Craddock, P. and Sio, C.K. (2012) Iron
 1127 isotope fractionation in planetary crusts. *Geochim. Cosmochim. Acta* 89, 31-45.

1128 Warren, P.H. (1993) A concise compilation of petrologic information on possibly nonmare
 1129 Moon rocks. *Am. Mineral.* 78, 360-376.

1130 Weyer, S., Anbar, A.D., Brey, G.P., Munker, C., Mezger, K. and Woodland, A.B. (2005) Iron
 1131 isotope fractionation during planetary differentiation. *Earth Planet. Sci. Lett.* 240, 251-264.

1132 Weyer, S., Anbar, A.D., Brey, G.P., Munker, C., Mezger, K. and Woodland, A.B. (2007) Fe-
 1133 isotope fractionation during partial melting on Earth and the current view on the Fe-isotope
 1134 budgets of the planets - (reply to the comment of F. Poitrasson and to the comment of B.L.
 1135 Beard and C.M. Johnson on "Iron isotope fractionation during planetary differentiation" by
 1136 S. Weyer, A.D. Anbar, G.P. Brey, C. Munker, K. Mezger and A.B. Woodland). *Earth*
 1137 *Planet. Sci. Lett.* 256, 638-646.

1138 Weyer, S. and Ionov, D.A. (2007) Partial melting and melt percolation in the mantle: The
 1139 message from Fe isotopes. *Earth Planet. Sci. Lett.* 259, 119-133.

1140 Wiechert, U. and Halliday, A.N. (2007) Non-chondritic magnesium and the origins of the
 1141 inner terrestrial planets. *Earth Planet. Sci. Lett.* 256, 360-371.

1142 Wieczorek, M.A., Jolliff, B.L., Khan, A., Pritchard, M.E., Weiss, B.P., Williams, J.G., Hood,
 1143 L.L., Richter, K., Neal, C.R., Shearer, C.K., McCallum, I.S., Tompkins, S., Hawke, B.R.,
 1144 Peterson, C., Gillis, J.J. and Bussey, B. (2006) The constitution and structure of the lunar
 1145 interior, *New Views of the Moon*. MSA, pp. 221- 364.

1146 Wiesli, R.A., Beard, B.L., Taylor, L.A. and Johnson, C.M. (2003) Space weathering processes
 1147 on airless bodies: Fe isotope fractionation in the lunar regolith. *Earth Planet. Sci. Lett.* 216,
 1148 457-465.

1149 Williams, H.M., Peslier, A.H., McCammon, C., Halliday, A.N., Levasseur, S., Teutsch, N.
 1150 and Burg, J.P. (2005) Systematic iron isotope fractionation variations in mantle rocks and
 1151 minerals: The effect of partial melting and oxygen fugacity. *Earth Planet. Sci. Lett.* 235,
 1152 435-452.

1153 Williams, H.M., Nielsen, S.G., Renac, C., Griffin, W.L., O'Reilly, S.Y., McCammon, C.A.,
 1154 Pearson, N., Viljoen, F., Alt, J.C. and Halliday, A.N. (2009) Fractionation of oxygen and
 1155 iron isotopes by partial melting processes: Implications for the interpretation of stable
 1156 isotope signatures in mafic rocks. *Earth Planet. Sci. Lett.* 283, 156-166.

1157 Williams, H.M. and Bizimis, M. (2014) Iron isotope tracing of mantle heterogeneity within
 1158 the source regions of oceanic basalts. *Earth Planet. Sci. Lett.* 404, 396-407.

1159 Yang, H., Lin, J.F., Hu, M.Y., Roskosz, M., Bi, W.L., Zhao, J.Y., Alp, E.E., Liu, J., Liu, J.C.,
 1160 Wentzowitch, R.M., Okuchi, T. and Dauphas, N. (2019) Iron isotopic fractionation in
 1161 mineral phases from Earth's lower mantle: Did terrestrial magma ocean crystallization
 1162 fractionate iron isotopes? *Earth Planet. Sci. Lett.* 506, 113-122.

1163 Zambardi, T., Poitrasson, F., Corgne, A., Meheut, M., Quitte, G. and Anand, M. (2013)
 1164 Silicon isotope variations in the inner solar system: Implications for planetary formation,
 1165 differentiation and composition. *Geochim. Cosmochim. Acta* 121, 67-83.

1166 Zhao, X.M., Zhang, H.F., Zhu, X.K., Tang, S.H. and Tang, Y.J. (2010) Iron isotope variations
 1167 in spinel peridotite xenoliths from North China Craton: implications for mantle
 1168 metasomatism. *Contrib. Mineral. Petrol.* 160, 1-14.

1169 Ziegler, K., Young, E.D., Schauble, E.A. and Wasson, J.T. (2010) Metal-silicate silicon
 1170 isotope fractionation in enstatite meteorites and constraints on Earth's core formation. *Earth*
 1171 *Planet. Sci. Lett.* 295, 487-496.

1172

1173 **Figure captions:**

1174

1175 Fig. 1: Initial planetary iron isotope composition ($\delta^{57}\text{Fe}$) estimates for the Earth and Moon
 1176 relative to IRMM-14 compared to chondrite and achondrite meteorite groups (Poitrasson,
 1177 2007). The arrows depict some more recent revisions towards putatively lighter compositions
 1178 for the Earth and Moon. See text for references and discussion.

1179

1180 Fig. 2: Iron isotope composition of bulk lunar rocks expressed in $\delta^{57}\text{Fe}$ relative to IRMM-14.
 1181 The group means and 2 standard error uncertainties are reported as continuous and dashed
 1182 lines/gray zones, respectively. The Earth reference value ($\delta^{57}\text{Fe} = 0.10 \pm 0.03\text{‰}$; from
 1183 Poitrasson et al., 2004; 2013) is shown for comparison. Note that highland rocks have been
 1184 split into anorthosites and Mg-suite rocks. Samples having extreme values or variable Fe
 1185 isotope compositions on different NASA CAPTEM aliquots are labeled, besides KREEP
 1186 basalt 15386. Data are from Table 1 and Supplementary Table. Iron isotope composition
 1187 estimates for the Lunar Upper Mantle ($0.142 \pm 0.026\text{‰}$) and Bulk Moon ($0.094 \pm 0.035\text{‰}$) from

1188 this work are reported at the bottom. Note that previous estimates of the Bulk Moon $\delta^{57}\text{Fe}$
1189 values ranged from -0.1 to +0.2%, depending on the authors (see text and Fig. 1).

1190

1191 Fig. 3: Iron isotope composition of bulk highland lunar rocks as a function of their pyroxene
1192 modal content. At low pyroxene modal contents the Fe isotope values are scattered, possibly
1193 due to the predominance of olivine affected by diffusive effects leading to low Fe isotope
1194 values or increasing plagioclase abundance having a heavy isotope composition. At higher
1195 pyroxene modal content, though, a homogeneous Fe isotope composition emerges at
1196 $0.094 \pm 0.035\%$ (2SE) for these highland rocks. Data are from Table 1, Supplementary Table
1197 and the Lunar Sample Compendium (Meyer, 2011).

1198

1199 Fig. 4: Relations between iron isotope composition estimates of planetary bodies against their
1200 Fe/Mn (a), Rb/Sr (b) and Mn/Na (c) estimates. APB stands for Angrite Parent Body; Note that
1201 the elemental ratio estimates are for CI chondrites. See Supplementary Table for data sources.

1202

Table 1: New iron isotope composition of bulk lunar samples from this study, along with previously published data on the same samples

	Rock type	Rock mass powdered (g)	$\delta^{57}\text{Fe}$ (‰)	2SE ^k	$\delta^{56}\text{Fe}$ (‰)	2SE ^k	n
Low-Ti basalts							
12009,130	Olivine vitrophyre basalt	0.856	0.101	0.031	0.045	0.020	6
12011,29	Pigeonite basalt	1.003	0.097	0.081	0.066	0.056	6
12031,37	Pigeonite basalt	0.824	0.154	0.071	0.098	0.066	6
12038,244	Feldspathic basalt	0.874	0.131	0.056	0.079	0.027	6
12045,30 ^a	Ilmenite basalt	1.032	0.136	0.094	0.082	0.062	6
12045,13 ^b		~1	0.138	0.078	0.092	0.052	3
12054,6	Ilmenite basalt	0.740	0.135	0.078	0.081	0.054	6
14053,237	Group C Al-rich basalt	0.634	0.070	0.024	0.038	0.023	6
14053,263 ^a		1.003	0.043	0.023	0.039	0.020	6
14053 ^c		~0.05	0.130	0.030	0.100	0.030	4
15555,988 ^a	Olivine-normative basalt	1.097	0.086	0.064	0.056	0.041	6
15555,115 ^b		~1	0.208	0.058	0.139	0.038	9
15555 ^d		~0.1	0.044	0.026	0.029	0.017	5
15555,955 ^e		~0.05-0.1	0.104	0.050	0.073	0.029	1
15555,139 ^f		0.0053	0.10	0.02	0.05	0.04	7
15555 ^c		~0.05	0.130	0.050	0.090	0.030	2
15556,198	Olivine-normative basalt	0.601	0.117	0.071	0.061	0.046	6
Mean low-Ti basalts (this study)ⁱ			0.113	0.023			9
Mean low-Ti basalts^j			0.127	0.012			27
High-Ti basalts							
10003,178	Low-K Group B2 basalt	0.856	0.320	0.071	0.213	0.047	6
10003 ^c		~0.05	0.200	0.040	0.130	0.040	2
10044,638	Low-K Group B1 basalt	0.827	0.354	0.118	0.235	0.072	6
10049,101 ^a	High-K Group A basalt	1.124	0.167	0.118	0.114	0.073	6
10057,268	High-K Group A basalt	0.599	0.189	0.028	0.158	0.038	6
10057 ^c		~0.05	0.210	0.060	0.140	0.050	2
10058,250	Low-K Group B1 basalt	0.852	0.290	0.040	0.198	0.030	9
10092,14	Low-K Group B3 basalt	0.699	0.327	0.039	0.212	0.031	6
70135,93	Type A basalt	1.048	0.238	0.061	0.146	0.045	9
70135 ^c		~0.05	0.220	0.060	0.150	0.050	3
70315,30	Type B1 basalt	0.991	0.325	0.050	0.196	0.034	6
74235,62	Type B2 basalt	0.595	0.253	0.086	0.166	0.053	6
74245,34	Type C basalt	0.604	0.213	0.079	0.107	0.059	6
74275,48	Type C basalt	0.820	0.273	0.081	0.177	0.040	6
74275,240 ^g		0.251	0.270	0.025	0.220	0.025	3
78598,6	Type A basalt	0.622	0.324	0.039	0.235	0.037	6
Mean high-Ti basalts (this study)ⁱ			0.273	0.039			12
Mean high-Ti basalts^j			0.274	0.020			25
KREEP							
15386	KREEP basalt	0.544	0.183	0.067	0.112	0.039	6
15386 ^b		~1	0.230	0.033	0.154	0.022	3
15386 ^d		~0.1	0.210	0.080	0.140	0.053	3
Highland rocks							
15415,190 ^a	Ferroan anorthosite	0.998	0.496	0.057	0.347	0.113	6
15455,306 ^a	Mg-suite Norite (CAN clast)	1.061	0.070	0.017	0.024	0.025	6
15455 ^c		~0.05	0.050	0.010	0.030	0.010	5

60015,785	Cataclastic ferroan anorthosite	0.160	0.054	0.038			6
60025,874 ^h	Ferroan anorthosite	0.167	-0.393	0.050	-0.276	0.037	12
60025 ^b		~1	0.225	0.063	0.150	0.042	3
62255,191	Ferroan anorthosite	0.139	0.265	0.119	0.181	0.088	3
62255,134 ^b		~1	0.176	0.062	0.118	0.042	3
62275,21	Ferroan anorthosite	0.144	0.272	0.026	0.064	0.042	3
67955,94	Mg-suite noritic anorthosite	0.166	0.156	0.049	0.060	0.114	6
77215,258 ^a	Mg-suite noritic	1.043	0.045	0.051	0.019	0.029	6
77215 ^b		~1	0.128	0.024	0.085	0.016	5
72415,82 ^h	Mg-suite dunite	0.127	-0.730	0.062	-0.446	0.085	12
72415 ^f		0.0011	-0.50	0.03	-0.35	0.02	11
72415 ^c		~0.05	-0.600	0.050	-0.400	0.040	2
76335,63 ^a	Mg-suite troctolite	1.012	-0.007	0.062	-0.050	0.122	4
Mean highland rocks (this study)ⁱ			0.023	0.235			10
Mean highland rocks^j			0.078	0.124			15
Impact melts and breccia							
14310	KREEP Basaltic impact melt	2.027	0.125	0.130	0.086	0.069	6
14310 ^d		~0.1	0.153	0.039	0.102	0.026	3
Mean impact melts and breccia^j			0.093	0.055			5
Mean Moon			0.166	0.032			73

^a New dissolution from a powder prepared in Toulouse. ^b Poitrasson et al. (2004), ^c Sossi and Moynier (2017), ^d Weyer et al. (2005), ^e Liu et al. (2010), ^f Wang et al. (2015) and ^g Wiesli et al. (2003). ^h The results presented were obtained from two separate sample aliquot dissolution and Fe purification to duplicate completely the measurements. ⁱ When several powder aliquots have been analyzed, the mean value of the sample was used for the calculation. ^j See the supplementary table for the extensive compilation used to compute this mean and uncertainty. When several values are available for a given rock, the mean of all these determination was used in the calculation to avoid overweighting samples with multiple determinations. ^k The iron isotope composition and two standard error (2SE) uncertainties quoted are calculated from the number of analyses indicated (n) and using the Student's t-correcting factors (Platzner, 1997).

Figure 1

Fig. 1

Figure 2

Fig. 2

Figure 3

Fig. 3

Figure 4

Fig. 4

Electronic Annex
[Click here to download Electronic Annex: Supplementary Table.xlsx](#)