Investigating the use of fallout and geogenic radionuclides as potential tracing properties to quantify the sources of suspended sediment in a mining catchment in New Caledonia, South Pacific

Virginie Sellier¹ • Oldrich Navratil² • J. Patrick Laceby³ • Michel Allenbach⁴ • Irène Lefèvre¹ • Olivier Evrard¹

¹Laboratoire des Sciences du Climat et de l'Environnement (LSCE), UMR 8212 (CEA/CNRS/UVSQ-IPSL),
Université Paris-Saclay, Gif-sur-Yvette, France

²Laboratoire Environnement-Ville-Société (EVS), UMR 5600/IRG, Université Lumière Lyon 2, Lyon (France)

³Environmental Monitoring and Science Division (EMSD), Alberta Environment and Parks (AEP Calgary,
Alberta, Canada

⁴LIVE-EA 4243, Université de Nouvelle-Calédonie, Nouméa, (Nouvelle-Calédonie, France) & LABEX Corail,
France

✉ Virginie Sellier

virginie.sellier@lsce.ipsl.fr

Abstract

Purpose. New Caledonia, a French island located in the south-west Pacific Ocean, has vast nickel resources. Open-cast mining has strongly increased soil erosion and the subsequent downstream transfer of sediments in river systems resulting in fundamental morphological changes (e.g. hyper-sedimentation, over-burden). Understanding the sediment source contributions from mining activities is therefore important to guide the implementation of effective management measures.

Materials and methods. A pilot sediment tracing study was conducted in the Thio River (400 km²) catchment draining the island's first mine. Sediment deposited during the February 25, 2015 and April 10, 2017 flood events was collected with a tributary tracing design, including main stem (n=19) and tributary (n=24) samples. The tributaries were classified into two source types: sub-catchments draining mining sites and sub-catchments devoid of mining activities. Three sets of potential tracing parameters (i.e. fallout radionuclides, geogenic radionuclides, and elemental geochemistry) were examined for their potential to model the contributions of sediment from mining versus non-mining tributaries to sediment collected on the Thio River.

Results and discussion. The very low fallout radionuclide activities (¹³⁷Cs and ²¹⁰Pb_{xs}) found in the source and sediment samples demonstrate that most material transiting the river network is derived from subsoil sources. Geogenic radionuclides and elemental geochemistry were therefore utilized in the tributary tracing approach. Accordingly, U and K were selected as the optimal tracers of the two main lithological regions supplying sediment to the main stem of the Thio River. Model results demonstrated that tributaries with mining activity dominated the sediment supply to the Thio River with mean sediment contributions of 68 % (SD 28 %) for the 2015 flood and 86 % (SD 7 %) for the 2017 event.

Conclusions and perspectives. Geogenic radionuclides and elemental geochemistry were more effective at discriminating between tributaries with and without mines than fallout radionuclides. The similarity of the model results from tracing with geogenic radionuclides and elemental geochemistry illustrates their potential to investigate sediment source contributions in similar catchments across the South Pacific Islands affected by mining activities.

Keywords Caesium-137 • Nickel mining • Sediment source fingerprinting • Soil erosion • X-ray fluorescence

1. Introduction

Soil erosion and the associated sediment transfer to river systems are exacerbated in tropical regions exposed to heavy rainfall, particularly during cyclones (Mertes 1994; Aalto and Nitttrouer 2012). In the South Pacific region, including the French Island of New Caledonia, these tropical depressions frequently generate overbank peak flows that may damage human settlements, public infrastructure and agricultural land (Terry et al. 2008). The flood events generated by cyclones and tropical depressions supply the major part of the sediment flux transiting rivers in this region. For example, during significant rainfall events, riverine suspended sediment concentrations (SSC) have exceeded concentrations of 350 g L^{-1} resulting in high sedimentation rates ($> 3 \text{ cm yr}^{-1}$) on floodplains in Fiji (Terry et al. 2002).

In New Caledonia, a significant proportion of the sediment load is potentially generated by mining activities, which increase the terrigenous inputs to downstream coastal lagoons (Bird et al. 1984). Although the mining sector has contributed to the economic growth of New Caledonia since the 1880s (5 % of GDP (IEOM 2016)), it is also considered to be the one of the main drivers of environmental degradation on the island (Sabinot and Lacombe 2015).

Suspended sediment is known to transport large quantities of nutrients and contaminants in river systems (Owens et al. 2001; Owens and Walling 2002; Walling et al. 2003). In New Caledonia, high concentrations of trace metals, including Ni and Cr, were found in pelagic and freshwater organisms (Hedouin et al. 2007; Hedouin et al. 2008). In addition, the suspended sediment itself degrades coral reef systems (Rogers 1990). Coral reefs are particularly sensitive to increases in turbidity as they often become stressed by the reduction in light penetration (Rogers 1979; Heintz et al. 2015; Restrepo et al. 2016). The degradation of 4500 km^2 of Caledonian coral reefs, listed as a UNESCO World Heritage site, threatens a variety of fundamental ecosystem services (e.g. fisheries, coastal protection and tourism) estimated at a value of \$340 million USD per year (Pascal 2010).

To protect these natural resources, the effective management of mining resources is required (Pascal et al. 2008; David et al. 2010). Open-cast mining is widespread in New Caledonia. The island's soils, developed on ultrabasic rocks (e.g. peridotites), are enriched in Fe and transition metals such as Mn, Ni, Cr and Co (Schwertmann and Latham 1986). The weathering of the peridotite results in the formation of Ni- and Fe-rich smectite, serpentine, goethite and hematite. The migration of these transition metals through the soil profile results in higher Ni concentrations in the saprolite (Ni: 0.5-3.5 %) and laterite layers (Ni: 0.9-1.8 %) (Trescases 1973). Both saprolite and laterite are mechanically mined, generating large quantities of overburden material

(e.g. mining tailings). This mining waste was historically (i.e. since 1880) stored on the foothills of the mines. In 1975, the inhabitants of New Caledonia, whose livelihood was already impacted by heavy rainfall events, became increasingly aware of the deleterious consequences of mining activities due to cyclone Alison, which generated extensive flooding and mudflows across the island.

Although the management of overburden material was improved through the prohibition of mining waste storage on foothills and the creation of storage ponds on the mining sites after 1975, a significant quantity of legacy material has accumulated on the hillsides where it is frequently exposed to erosion processes. In addition to these mining-related inputs of both fine sediment and coarser materials, other sources of sediment may supply a significant quantity of particulate material to New Caledonian river systems. Bushfires or land-clearing fires occur frequently in the region (Dumas et al. 2010). These fires can lead to shallow landslides and/or extensive soil erosion of surface material on hillslopes (Blake et al. 2009; Nyman et al. 2011; Smith et al. 2012). Moreover, the deer population significantly increased during the last 40 years. The associated trampling may have further accelerated surface soil degradation (Shellberg et al. 2010).

To help improve the management of deleterious sediment loads in mining-affected regions in New Caledonia, it is important to improve our understanding of the main sediment sources. Sediment fingerprinting techniques have been widely applied to investigate the relative contribution of multiple sources (Walling et al. 1979; Collins et al. 1996; Walling 2005). These techniques can provide information on the main erosion processes involved in sediment mobilization (e.g. mass movement, sheet, rill, gully and channel erosion) and the spatial origin of sources (e.g. land-use, lithology) that cannot be provided by traditional monitoring techniques. They consist of the measurement of multiple conservative properties, or fingerprints, in both sediment sources and river sediment in order to quantify the respective source contributions to material transiting river systems (Davis and Fox 2009; Koiter et al. 2013; Walling 2013). The commonly used properties can be grouped into four main categories; geochemical (e.g. elemental geochemical contents, isotopic ratios), biogeochemical (e.g. organic compounds), physical (e.g. mineral, magnetic parameters) and nuclear properties (e.g. radioactive elements) (Koiter et al. 2013). In recent years, these latter have diversified with the emergence of more alternative sediment fingerprinting techniques based on sediment color (Martínez-Carreras et al. 2010; Legout et al. 2013), pollen plant (Brown et al. 2008) or more recently environmental DNA (Ficetola et al. 2018; Evrard et al. 2019). This diversity in the use of fingerprints has been achieved in conjunction with the improvement of mixing models (e.g. reduction of associated uncertainties). Following this progress, Laceby et al. (2019) recommend to further develop the combination of multiple sets of fingerprints to enrich the knowledge base of

these parameters and the associated potential biases (e.g. source variability, particle size selectivity). These authors outlined in particular the need to improve our understanding of processes that establish the source fingerprint signals and their conservative behavior in order to be able to provide a better analysis of key processes driving the mobilization, generation, and deposition of sediment.

In New Caledonia, sediment originates from two main sources. In areas affected by bushfires and grassland degradation, sediment may be derived from a mixture of surface material (e.g. hillslope erosion) and subsurface soils (e.g. shallow landslides and channel bank erosion). In contrast, sediment associated with mining activities (e.g. overburden, bare mining soils, mining prospection and access roads) will likely consist predominantly of subsurface material. One potential approach to discriminate between surface and subsurface sediment sources is the comparison of their fallout radionuclide activities (^{137}Cs , $^{210}\text{Pb}_{\text{xs}}$). In particular, ^{137}Cs ($t_{1/2} = 30$ years), a by-product of thermonuclear bomb testing, which mainly occurred between 1952–1974 in the Southern Hemisphere, was demonstrated to be an effective tracer of surface and subsurface sources in numerous studies conducted in nearby Australia (Olley et al. 2013; Wilkinson et al. 2013; Hancock et al. 2014). Accordingly, ^{137}Cs could potentially discriminate between mining and non-mining sources, the latter of which is hypothesized to have higher ^{137}Cs concentrations owing to a greater proportion of surface material.

In addition to the investigation of fallout radionuclide analyses, the inclusion of other tracers including geochemical elements (Koiter et al. 2013; Laceby et al. 2015; Batista et al. 2018) holds potential to provide additional sediment source information such as whether non-mining or mining sources dominate the supply of sediment. Peridotite massifs where nickel ores are extracted cover one third of the island and are characterized by a depletion of Th, K and U. Contrarily, the volcano-sedimentary formation, that covers approximately the remaining two thirds of the island, are naturally enriched in Th, K and U (Sevin 2014). Accordingly, the Th, K and U contents may discriminate between sediment contributions from tributaries draining mines compared to watercourses flowing across areas devoid of mining activities.

The objective of this research is to quantify the contribution of mining and non-mining tributaries to the Thio River, one of the first catchments in New Caledonia to be mined for nickel. In particular, three sets of potentially discriminant tracing properties will be tested to quantify sediment source contributions: fallout radionuclides (i.e. ^{137}Cs and $^{210}\text{Pb}_{\text{xs}}$), geogenic radionuclides (i.e. Th, K and U) and elemental geochemistry. The model results will quantify the relative contribution of mining tributaries to sediment loads degrading the Thio River and its estuary. The results will also help to develop an approach to examine source contributions from

mining/non-mining catchments across New Caledonia and potentially across similar South Pacific and other tropical islands.

2. Materials and methods

2.1. Study area

The New Caledonian archipelago is located in the southwestern Pacific Ocean, in Melanesia. The Thio River catchment (397 km²) is located on the east coast of the main island, called La Grande-Terre (17,000 km²). The catchment has a variable topography, with a mean altitude of 416 m asl (above sea level); ranging from 0 to 1352 m asl (Figure 1-b). On the western part of the catchment, volcanoclastic sandstone, clay rock and magmatic rock formations are composed of cherts (22 %), sandstone (9 %), a mix of basalt, dolerite and gabbro (6 %), polymetamorphic rocks (6 %) and alluvia (4 %). Laterites (18 %), peridotites (17 %), serpentines (10 %) and hazburgites (1 %) constitute the peridotite massifs, which mainly cover the eastern part of the catchment (40 % of the catchment surface), which results in its steeper relief (Garcin et al. 2017) (Figure 1-a). The main land use in the catchment is permanent vegetation including forests (37 %), shrubland (35 %) and bushland (18 %) (Alric 2009). According to the mining registry, mining concessions cover 40% of the catchment area including 21% (Figure 1-c) corresponding to active and abandoned mining sites and exploration.

The climate is tropical and is characterized by the alternation of a hot wet season (November-April; mean temperature of 27 °C) and a cooler dry season (May-October; mean temperature of 20 °C). The mean annual rainfall in the Thio catchment is 1620 mm (1981–2008) (Alric 2009). About fifty cyclones have affected New Caledonia since the 1880s (i.e. a cyclone or a tropical depression occurs on average every 2.7 years) (Garcin 2010). According to local meteorological records, cyclones may supply more than 20 % of the annual rainfall in only one day (Météo France).

The catchment's river network is dense with 12 tributaries (Figure 1-b) flowing into the main stem of the Thio River (28 km long). Ninety-two percent of the river channel length is characterized with a slope lower than 5 %. The overall slope index I_g (34 ‰) reflects the torrential regime observed in the Thio River tributaries. The catchment is very reactive to heavy rainfall given the extensive bare soil surface associated with past mining activities (~10 sites), ongoing mining operations (e.g. 2 sites: *Thio Plateau*, *Camp des Sapins*) and the occurrence of 6 km² of mining roads, which all increase runoff production as they are well connected to the river network (Alric 2009). Extensive erosion processes are evident across the Thio catchment, including rills, gullies, landslides and channel bank erosion (Danloux and Laganier 1991).

2.2. Hydro-sedimentary monitoring

Rainfall is recorded at three stations (i.e. Thio Plateau, Thio village, Camps des Sapins) operated by Météo France and five stations by the island's local authorities (i.e. Kouaré, Bel-Air, Ningua, Kuenthio, Mont Do managed by the DAVAR – Direction des Affaires Vétérinaires Alimentaires et Rurales), with records available since 1952. Rainfall data at these eight stations was collected for two floods investigated in the current research, the tropical depression on February 25, 2015 and Cyclone Cook on April 10, 2017. Daily discharge has been monitored at a river gauging station located on the main stem of the Thio River section (at Saint-Michel) since 1981 by the DAVAR (Figure 1-b).

2.3. Source and river sediment sampling

A tributary tracing design was used to investigate sediment source contributions in the Thio River catchment (Olley and Caitcheon 2000; Evrard et al. 2015; Laceby et al. 2017). The tributary approach consists of sampling sediment in the different tributaries and using these samples as potential sources of sediment sampled further downstream on the main stem of the Thio River (Figure 1-c). Using lag deposits as potential source samples is a common technique in sediment fingerprinting studies (Bottrill et al. 1999). For those catchments that are not equipped with automatic sediment such as the Thio River catchment, lag deposit sampling is the most suitable alternative for suspended sediment sampling, particularly because it is well representative of the whole catchment area. Olley et al. (2013) showed that equivalent results (i.e fallout radionuclide activities) were observed for both lag deposits and suspended sediment samples (i.e. <10 µm fraction). However, this sampling technique also has disadvantages such as those outlined by Haddadchi et al. (2013), and including a lack of information on the hysteresis effect of flood events and the effect of discharge on sediment contribution from different sources.

Lag deposits were collected at exposed subaerial sites free of vegetation on channel bars. Forty-three samples were collected after two major floods (~10 yr return period): (1) the tropical depression of February 25, 2015 with lag deposits sampled between April 30 and May 5, 2015 ($n=31$) and (2) Cyclone Cook on April 10, 2017, sampled between May 16 and 17, 2017 ($n=12$). At each sampling site, five to ten subsamples of fine sediment were collected across a 10-m² surface with a plastic trowel. The subsamples were composited into one sample representative of the fine sediment deposited on the channel bars. Among these samples, 16 were

collected in tributaries with mine activities ('mining tributaries'), eight in tributaries draining areas devoid of mining activities ('non-mining tributaries') and 19 on the main stem of the Thio River (Figure 1-c). The samples were oven-dried at 40°C for ~48 hours and sieved to 2 mm and 63 µm.

2.4. Source and river sediment analysis

2.4.1. Particle size

Fine particles are preferentially mobilized and transported over longer distances. This process may lead to the accumulation of finer particles in the lower reaches of the main river channel and receiving waters. In the Thio catchment, this phenomenon has resulted in the hyper-sedimentation observed in the main stem and the lagoon. This particle size selectivity may affect biogeochemical properties (i.e. geochemical transformations, organic matter depletion/enrichment) and physical properties (i.e. particle size) of source and downstream sediment material (Lacey et al. 2017). Different techniques were implemented to address particle size effects in sediment fingerprinting studies (e.g. fractionation of source and sediment material to a narrow particle size range, concentration corrections) (Stout et al. 2013; Foucher et al. 2015; Le Gall et al. 2016). In the current research, fallout radionuclide activities, geogenic radionuclide and geochemical element contents measured in the <2 mm and <63 µm fractions of samples were compared. Gamma spectrometry and X-ray fluorescence analyses were carried out on a subset of samples (n=31) for which both fractions were analyzed. Linear correlations between the parameters measured in <2 mm and <63 µm fractions was assessed to select properties that were the less affected by a potential particle size effect.

2.4.2. Gamma spectrometry

Tributary and main stem samples (n=43) were packed in 15 mL polyethylene containers for radionuclide analysis with gamma spectrometry using coaxial N- and P-type HPGe detectors (Canberra/Ortec) at the Laboratoire des Sciences du Climat et de l'Environnement (LSCE, Gif-sur-Yvette, France). ¹³⁷Cs activities were measured at the 662 keV emission peak. ²¹⁰Pb_{xs} activities were calculated by subtracting the supported activity (determined by using two ²²⁶Ra daughters), ²¹⁴Pb (average count number at 295.2 and 351.9 keV) and ²¹⁴Bi (609.3 keV)) from the total ²¹⁰Pb activity measured at 46.5 keV (Le Gall et al. 2017). Assuming that ²³²Th and its daughter products were in secular equilibrium, ²²⁸Th activities were calculated using the average between the gamma rays of two of its daughter products, ²¹²Pb (239 keV) and ²⁰⁸Tl (583 keV) (Le Gall et al. 2016). Elemental concentrations in Th (mg kg⁻¹) were calculated from ²²⁸Th activities. In the same way, ²³⁸U was

estimated from the gamma rays of its daughter, ^{214}Pb (295 and 352 keV) and the U elemental content was expressed in mg kg^{-1} . ^{40}K was detected at 1460 keV and the K concentration was provided in mg kg^{-1} . Counting efficiencies and calibration were conducted using certified International Atomic Energy Agency (IAEA) standards (IAEA-444, 135, 375, RGU-1 and RGTh-1) prepared in the same containers as the samples.

2.4.3. *X-ray fluorescence*

X-ray fluorescence measurements were conducted on sediment samples stored in air double X-ray Mylar film small mass holder cells with a sample quantity between 0.2 and 0.5g. Major element contents (i.e. Mg, Al, Si, K, Ca, Ti, Cr, Mn, Fe, Ni, Cu, Co and Zn) were analyzed by energy dispersive X-ray fluorescence spectrometry (Epsilon 3, Malvern PANalytical) at LSCE, Gif-sur-Yvette, France. Samples were irradiated by a primary beam generated by an Rh anode X-ray tube emitting electromagnetic waves between 100eV and 1MeV with a maximum power, typical current and voltage fixed to 15 W, 3mA and 50 kV respectively. The associated Si-drift detector had a Be window thickness of 8 μm and recorded the sample spectrum in a 2D optical geometry configuration. Between 8 and 16 certified reference samples including Internal Atomic Energy Agency (IAEA) standards were used to derive the calibration curves for the geochemical elements of interest. X-ray intensities were converted into concentrations using the Epsilon 3 software program through the application of the fundamental parameters method. Correlations between the determined and the standard elemental contents were comprised between 0.90 And 0.99. To validate the calibration, three certified samples different from those used for the calibration were analysed, and the associated mean relative error was 17 % (SD 5 %). Due to imprecise calibration for Co (error: 264 %), this element was removed from further analysis.

2.5. Conservative behavior and source discrimination

A multi-parameter composite fingerprinting approach was developed by Collins et al. (1996) to identify the spatial sediment sources and quantify their contributions in the river network. This approach first selects an optimal suite of tracers that discriminate different sediment sources. Second, this approach estimates the relative source contributions to target sediments using end member mixing models.

The fingerprinting technique relies on the ability of tracers to discriminate between sediment sources while remaining conservative. The use of non-conservative tracers may lead to incorrect source group prediction resulting in inaccurate estimations of source contributions (Sherriff et al. 2015). Accordingly, it is generally considered important to restrict the tracing parameters to those that demonstrate conservative behavior.

Therefore, each set of potential tracing properties (i.e. fallout radionuclides, geogenic radionuclides and elemental geochemistry) was plotted to ensure that the tributary source properties plotted within the source range.

Thereafter, the Mann-Whitney U-test ($\alpha=0.05$) selected parameters that significantly discriminated between the two sediment sources (i.e. mining and non-mining tributaries) for each group of potential tracers. A stepwise discriminant function analysis (DFA; $\alpha=0.05$) was then used to select the optimal number of potential tracers through the selection of those that collectively maximized the discrimination between both sediment sources for each set of tracers. The optimal tracers for modelling were selected by minimizing the Wilks' lambda values in the backwards mode, with a $p<0.05$ used to select a tracer and $p<0.10$ used to remove a tracer.

2.6. Source contribution modelling

A distribution modelling approach was used to model the source contributions from the mining and non-mining tributaries to target sediment (Laceby et al. 2015; Laceby and Olley 2015). This approach incorporates distributions throughout the entire modelling framework, including the source contribution terms and the target sediment. For this two-source model, it is assumed that sediment samples constitute a discrete mixture of their sources, with the contribution from Source A being x , and the contribution of Source B being $(1-x)$ where:

$$Ax + B(1 - x) = C \quad \text{Equation 1}$$

C is the target Thio river sediment distribution, A and B are again the two source distributions (i.e. mining and non-mining tributaries), and x is modelled as a truncated normal distribution ($0 \leq x \leq 1$) with a mixture mean (μ_m) and standard deviation (σ_m) (Laceby and Olley 2015). The model is solved by minimizing the median difference between the distributions of both sides of Equation 1 (e.g. C and $Ax + B(1 - x)$) with the Optquest algorithm in Oracle's Crystal Ball software with a randomly generated mixture mean (μ_m) and standard deviation (σ_m). Source contributions (x) were determined by simulating all distributions in Equation 1 with 2500 Latin Hypercube samples from 500 bins and varying the mixture mean (μ_m) and standard deviation (σ_m). This simulation and solving procedure repeated 2500 times and the median proportional source contribution from these 2500 additional simulations being reported as the source contribution to the target sediment. Uncertainty was determined through summing three quantifiable model uncertainties: (1) the median absolute deviation of the individual source median contribution for the additional 2500 simulations; (2) the modelled standard deviation; and (3) the median absolute deviation of this modelled standard deviation for the 2500 model additional simulations (Laceby et al. 2015).

3. Results

3.1. Conservative behavior and source discrimination

3.1.1 Fallout radionuclides

Although the samples were analyzed in an ultra-low background laboratory, fallout radionuclide activities (i.e. ^{137}Cs , $^{210}\text{Pb}_{\text{xs}}$) were below detection limits for 10 samples for ^{137}Cs , and 5 samples for $^{210}\text{Pb}_{\text{xs}}$ for <2 mm fraction of samples (Table 1). When analyzing the <63 μm fraction, the results remained below detection limits for five sediment samples for ^{137}Cs and for two samples for $^{210}\text{Pb}_{\text{xs}}$. For the samples above the detection limit, mean activities for ^{137}Cs were 0.3 Bq kg^{-1} (SD 0.3 Bq kg^{-1}) for mining sources. Slightly higher mean ^{137}Cs activities were measured in non-mining sources (1 Bq kg^{-1} ; SD 0.5 Bq kg^{-1}). Similar values were found in the Thio River target sediment samples (i.e. on the main stem) in 2015 (0.7 Bq kg^{-1} ; SD 0.4 Bq kg^{-1}) (Table 1, <63 μm fraction). $^{210}\text{Pb}_{\text{xs}}$ activities were lower in mining sources (6 Bq kg^{-1} ; SD 7 Bq kg^{-1}) compared to non-mining sources (19 Bq kg^{-1} ; SD 15 Bq kg^{-1}). $^{210}\text{Pb}_{\text{xs}}$ activities in the Thio River target sediment samples were similar to non-mining sources in 2015 (15 Bq kg^{-1} ; SD 8 Bq kg^{-1}) (Figure 2-b, Table 1, <63 μm fraction).

Fallout radionuclide activities measured in both <63 μm and <2 mm fractions of sediment were correlated; with r^2 coefficients of 0.66 for ^{137}Cs and 0.73 for $^{210}\text{Pb}_{\text{xs}}$ (Figure 3-a, b). Owing to particle size effect affecting fallout radionuclides activities (Figures 2 and 3, Table 1) and a larger number of results above the detection limits in the <63 μm fraction, the sediment fingerprinting approach was carried out exclusively on the finest fraction of samples. ^{137}Cs and $^{210}\text{Pb}_{\text{xs}}$ activities found in river sediment samples plotted within the range of activities measured in the potential source samples, indicative of conservative behavior (Figure 2-b). However, the p-values associated with the Mann-Whitney U test ($\alpha=0.05$, $p<0.1$) for both ^{137}Cs (0.7720) and $^{210}\text{Pb}_{\text{xs}}$ (0.1109) indicated that these tracers were not able to significantly discriminate between sediment sources (i.e. mining and no-mining tributaries). Accordingly, fallout radionuclides were not used to model sediment source contributions in the Thio River catchment.

3.1.2 Geogenic radionuclides

There were 10-fold higher Th, U, K contents found in non-mining source samples compared to mining source samples (Figure 4, Table 1, <2 mm). Mean Th contents were 6 mg kg^{-1} (SD 2 mg kg^{-1}) in non-mining source samples versus 0.6 mg kg^{-1} (SD 0.7 mg kg^{-1}) for mining source samples. Mean K content was $12,097 \text{ mg kg}^{-1}$ (SD 3980 mg kg^{-1}) for non-mining source samples relative to 1386 mg kg^{-1} (SD 1848 mg kg^{-1}) in mining

source samples. Finally, mean U contents were 1.2 mg kg^{-1} (SD 0.3 mg kg^{-1}) in non-mining sources and 0.1 mg kg^{-1} (SD 0.1 mg kg^{-1}) in mining source samples (Table 1). Target river sediment samples were characterized by low geogenic radionuclide contents, similar to those found in mining source samples (Figure 4). These contents remained similar in sediment collected after both 2015 and 2017 flood events (Figure 4, Table 1). Mean Th contents in the Thio River sediment samples were 1 mg kg^{-1} (SD 0.6 mg kg^{-1}) in 2015 and 2 mg kg^{-1} (SD 1 mg kg^{-1}) in 2017. Mean K content was 4711 mg kg^{-1} (SD 2270 mg kg^{-1}) in 2015 relative to 2792 mg kg^{-1} (SD 880 mg kg^{-1}) in 2017. Finally, mean U contents were 0.3 mg kg^{-1} (SD 0.1 mg kg^{-1}) in 2015 and 0.4 mg kg^{-1} (SD 0.3 mg kg^{-1}) in 2017 (Table 1, $<2 \text{ mm}$).

Analyses conducted on the $<2 \text{ mm}$ and $<63 \mu\text{m}$ fractions ($n = 31$) demonstrated that Th, K and U contents were very well correlated in fine and coarse material ($r^2=0.96, 0.92, 0.90$ respectively) (Figure 5-a, b, c). Accordingly, these geogenic radionuclide properties were considered not to be significantly impacted by a potential particle size effect. As the elements were highly correlated between the two particle size fractions, the $<2 \text{ mm}$ was used in the modelling process.

According the Mann-Whitney U test results, all three geogenic radionuclides provided significant discrimination between the sediment sources (i.e. $p\text{-value} < 0.1$, Table 2). U was selected by the DFA to model sediment source contributions from mining and non-mining tributaries (Table 2) with a Wilk's lambda value was 0.1742 and 96 % of sources correctly classified.

3.1.3 Geochemistry

The concentrations of several geochemical elements were higher in mining source samples compared to non-mining sources, with 17-fold higher levels of Ni, 10-fold higher levels of Cr, 5-fold higher levels of Mg and two-fold higher concentrations of Fe and Mn (Figure 6). On the contrary, K, Ti and Al were enriched in non-mining source samples compared to mining sources, with 8-fold higher K concentrations, 3-fold higher Ti concentrations and two-fold Al concentrations. Zn contents were similar in both source types, with mean concentrations of 125 mg kg^{-1} (SD 4 mg kg^{-1}) in non-mining sources compared to 146 mg kg^{-1} (SD 47 mg kg^{-1}) in mining sources. Elemental concentrations measured in the river sediment samples generally plotted closer to the levels measured in mining sources compared to non-mining sources (Figure 6). They were similar in deposits collected in 2015 and 2017, except for the Mg contents that increased (32 %) and the Al and Ti contents that decreased (31 % and 44 %) between both flood events.

When comparing measurements on the <2 mm and <63 µm fractions, concentrations in most elements (i.e. Al, Ca, Cr, Fe, K, Mg, Mn, Ni, Si Ti and Zn) were not impacted by particle size (r^2 between 0.77 and 0.99) (Figure 7-a, b, c, e, f, g, h, i, j, k, l). In contrast, Cu contents were not well correlated in both fractions ($r^2=0.35$) (Figure 7-d). Accordingly, Cu was not included in as a potential tracer. As the elements were highly correlated between the two particle size fractions, the <2 mm was used in the modelling process.

All ten geochemical elements (i.e. Al, Ca, Cr, Fe, K, Mg, Mn, Ni, Si and Ti) that showed similar concentrations and correlations between the <2 mm and <63 µm fractions significantly discriminated between the potential sediment sources (i.e. Mann-Whitney U test ($\alpha=0.05$, $p<0.1$) (Table 2). The DFA selected K as the optimal tracer of mining and non-mining source sediments with a Wilk's lambda of 0.1691 and 96 % of the source samples were correctly classified (Table 2).

3.2. Modelling results

The geogenic model (U) estimated that 68 % (SD 31 %) of sediment transiting the Thio River during the 2015 flood event originated from mining tributaries. Nevertheless, the large standard deviation (31 %) (Table 3) reflects the change of dominant source after different confluence depending on the tributary characteristics. In uppermost areas of the catchment, the contribution of mining tributaries clearly dominated (97 %) (Figure 8, Table 3, [1]). After the confluence with the Kouaré River (Figure 8, [c]), the non-mining sources became the highest (95 %) (Table 3, [3]). Further downstream, the proportions of the mining sources increased again to reach 79 % (Thio Plateau input, Figure 8,[j, k, l], Table 3, [8]). Near the estuary, these mining sources remained largely dominant with sediment contributions between 80-86 % (Table 3, [9-11]).

During the 2017 flood event, the geogenic model also estimated that the mining tributaries dominated the supply of sediment throughout the main stem of the Thio River with a mean contribution of 87 % (SD 8 %) (Table 4). The only exception was the Kouaré River which likely supplied a significant contribution of non-mining sediment to the main stem of the Thio River (Figure 9, [c], Table 4, [4]). However, the mining tributary contribution remained dominant in the estuary with a contribution of 86 % (SD 0.1 %) (Figure 9, Table 4, [7-8]).

The elemental geochemistry model (K) demonstrated that mining tributaries supplied 68 % (SD 25 %) of the sediment material transiting the Thio River during the 2015 flood event. The results were similar with those obtained with the geogenic model (U): sediment material was mainly supplied by mining tributaries except after the confluence with the Kouaré River where the non-mining sources dominated with a mean contribution of 80 % (Table 3, [3]). In the estuary, the mining tributary contribution was estimated to vary between 63-89 % (Table

3, [9-11]). The largest difference between both model outputs was 29 % for the 2015 flood (Table 3, [5]). For the 2017 flood, results obtained with this model indicated that 84 % (SD 7 %) of sediment material originated from mining tributaries. Along the main stem of the Thio River, mining tributary contributions dominated the supply of sediment (Table 4 [1-8]). Again, the largest difference observed between the results of both models remained low (7 %).

4. Discussion

4.1. Implications for tracing the impact of mining activities on sediment supply in the South Pacific

One original assumption of the current research was that fallout radionuclides should provide discrimination between non-mining tributaries, with more surface soil erosion from shallow landslides and areas affected by bushfires and grassland degradation, compared to mining tributaries with higher subsoil erosion from mining related activities including the overburden, bare mining soils and access roads. Accordingly, higher ^{137}Cs and $^{210}\text{Pb}_{\text{xs}}$ activities were expected in potential source samples collected in non-mining tributaries whereas they were assumed to be depleted in mining tributaries where topsoil has been removed to provide access to the laterite and saprolite layers enriched in nickel. However, the results demonstrated low ^{137}Cs and $^{210}\text{Pb}_{\text{xs}}$ activities in both source tributaries. The fallout radionuclide activities in the non-mining tributary sediment were ~10 times lower than those measured in surface sources collected in nearby Australia. Wallbrink et al. (1998) showed that ^{137}Cs and $^{210}\text{Pb}_{\text{xs}}$ activities were comprised between 0.6-3 Bq kg⁻¹ for subsurface sources and 18-270 Bq kg⁻¹ for surface sources when analysing the <2 mm fraction. More recent studies focused on those fallout radionuclide activities contained in the finest fractions (i.e. <63µm and <10µm) of source samples, to which fallout radionuclides are preferentially bound (Smith et al. 2011; Olley et al. 2013; Wilkinson et al. 2013). In these studies, ^{137}Cs and $^{210}\text{Pb}_{\text{xs}}$ activities ranged from 4 to 10 Bq kg⁻¹ for subsurface sources and from 22 to 200 Bq kg⁻¹ for surface sources when analysing the <63 µm fraction (Smith et al. 2011). When restricting the analysis to the <10 µm fraction, fallout radionuclide activities were similar to those measured on the <2 mm and <63 µm fractions, varying between 0.15-15 Bq kg⁻¹ for subsurface sources and 10- 150 Bq kg⁻¹ for surface sources (Olley et al. 2013; Wilkinson et al. 2013). Consequently, the low ^{137}Cs and $^{210}\text{Pb}_{\text{xs}}$ activities found in both source tributaries in the current research should likely not to be attributed to the particle size fractions analyzed (<2 mm and <63 µm). On the contrary, Krause et al. (2003) showed that ^{137}Cs activity signatures equivalent to those measured in the Thio River catchment area were rather representative of gullied catchments. The results obtained in the current research indicate that both the mining and non-mining tributaries, and the target sediment collected

on the main stem of the Thio River, are derived primarily from subsurface sources. These results are consistent with earlier observations made by Danloux and Laganier (1991) who observed the dominance of subsoil erosion processes in the New Caledonian mining catchments with the widespread occurrence of rills, gullies, landslides and channel bank erosion. These authors also outlined the specific impacts of mining activities conducted during the last 150 years in New Caledonia (e.g. soil stripping over several meters) as responsible for the expansion of these subsoil erosion processes, resulting in the massive supply of sediment to river systems during extreme rainfall events occurring during the cyclone season (i.e. between January and April) (Terry et al. 2002; Terry et al. 2008).

Despite the occurrence of particle size effects during erosion and sediment transfer processes, the current research showed that a large number of tracers were not impacted by particle size. Geogenic radionuclides (i.e. Th, K, U) and elemental geochemistry (i.e. Al, Ca, Cr, Fe, K, Mg, Mn, Ni, Si, Ti, and Zn) were well correlated in both particle size fractions (i.e. < 2 mm and <63 μ m). The geogenic radionuclides are particularly promising as they provide a strong discrimination between both potential sediment sources with very minimal particle size differences. Furthermore, the source discrimination for the geogenic radionuclides is based on a well-established physico-chemical basis. Sevin (2014) reported that, in New Caledonia, the volcano-sedimentary rock formations naturally contained high U and K elemental contents, whereas peridotite massifs were depleted in these elements. As mining sites are concentrated in peridotite massifs, the application of these tracers for identifying the mining source contribution is straightforward. Although U and K trace the lithological regions supplying sediment to the rivers (contributions of peridotite massifs vs. volcano-sedimentary formations), they may also provide indirect information on the land-use contributions. About half of the peridotite massif surface area is occupied by mining exploration, active and abandoned mining sites, whereas these activities are virtually absent from the areas underlain by volcano-sedimentary formations.

4.2. Spatial and temporal variations of sediment source contributions

The geogenic and elemental geochemistry models were based on different fingerprints (i.e. U and K respectively) and particle size fraction analyzes (i.e. <2 mm and <63 μ m fractions). Nevertheless, both fingerprinting techniques provided similar estimations of source contributions to sediment even though local differences in modelling results were as high as 29 % (Tables 3 and 4).

Although the two investigated events corresponded to floods with a return period of 10 years ($3500 \text{ m}^3\text{s}^{-1}$), they were generated by storms that contributed very different proportions of the annual rainfall (7 % in 2015 vs. 25 % in 2017). Furthermore, the spatial pattern of rainfall across the catchment strongly differed. The first event (February 25, 2015) generated more rainfall on the western part of the catchment (230 mm at Kouaré rainfall station vs. 136 mm at Bel-Air rainfall station) (Figure A1-a). The Kouaré River sub-catchment received twice more rainfall than observed in the rest of the Thio River catchment. This heterogeneous spatial rainfall distribution may explain the relatively high contribution of the non-mining source to the sediment deposited in the Thio River after the confluence with the Kouaré River (Figure 8, [c] 80-95 %). On the contrary, rainfall preferentially affected the eastern part of the catchment (409 mm at Thio Plateau station, 493 mm at Camps des Sapins station vs. 60 mm at Kouaré station) during the second event (April 10, 2017) (Figure A1-b). The maximum rainfall was recorded at stations located nearby the mines currently in operation (*Thio Plateau mine*, *Camps des Sapins*) during the cyclone. During this event, the contribution of mining tributaries was larger than that of non-mining tributaries, reducing their relative contribution to 24-27 % (Figure 9, [c]). Although the mining source contributions largely dominated the sediment production across the entire catchment during the two investigated events, the results of the current research showed that the spatial distribution of rainfall may locally or sporadically modify the sediment supply to the river. Nevertheless, the dominant sediment source remained the mining tributaries with contributions varying between 63-82 % during the 2015 event and 84-86 % during the 2017 event near the estuary. These results demonstrate that nickel mining activities directly impact sediment dynamics in this New Caledonia catchment.

5. Conclusions

In New Caledonia, mining activities have been postulated to exacerbate soil erosion and increase sediment transfer to the rivers and the lagoons. In this current study, it was demonstrated that tributaries draining mining sources (e.g. overburden, bare mining soils, mining prospection and access roads) dominated the sediment supply to the main stem of the Thio River. During the 2015 event, 68 % (SD 28 %, combined results of both models) of sediment transiting the Thio River originated from mining tributaries. In the 2017 event, the mining tributaries provided 86 % (SD 7 %, combined results of both models) of the sediment input into the main stem of the Thio River. Although the spatial distribution of rainfall was shown to locally modify the respective contributions of non-mining and mining tributaries, the latter provided the main contribution to the sediment collected in the Thio River estuary after both 2015 and 2017 flood events (63-89 %). These results suggest that

catchment management should focus on mining tributaries, including those draining the Thio Plateau mine located near the estuary.

In this study, substantial efforts were made to address the potential effects of changes in particle size composition (i.e. <63 μm and <2 mm) on sediment properties in a mining catchment. Indeed, detachment, mobilization, transportation and deposition processes did generate a particle size effect illustrated by the differences in fallout radionuclide activities observed in the <63 μm and <2 mm fractions. These results remained consistent with those found in other studies compiled by Laceby et al. (2017). This study also provided additional information on the behaviour of geochemical tracers (i.e. geochemical properties and geogenic radionuclides) in a mining environment, in which very few sediment tracing studies have been conducted so far. Interestingly, geochemical properties and geogenic radionuclide contents were not or only slightly affected by the particle size effect. It would be useful to investigate whether such a conservative behaviour is observed in other mining catchments.

Future applications of the sediment tracing technique in mining catchments of New Caledonia and the South Pacific should rely on the measurement of geogenic radionuclide contents and geochemical properties in potential source and sediment material. Furthermore, the current research demonstrated that geogenic radionuclides had the highest discrimination potential between both potential sediment sources. In contrast, the very low fallout radionuclide levels found in sediment analyzed during the current research prevented their use for sediment tracing, although these results illustrated that subsoil erosion dominated in this mining catchment. In the future, additional tracers should be used (e.g. colour, organic matter properties) to provide more detailed source information, in relation with the diversity of land uses and soil types found in similar catchments of the South Pacific Islands in general, and New Caledonia in particular.

Acknowledgements

This work was supported by the National Technical Research Center (CNRT) « Nickel and its environment », Noumea, New Caledonia » (n°10PS2013-CNRT.UNC/IMMILA). Virginie Sellier received a PhD fellowship from the French Atomic Energy Commission (CEA, Commissariat à l'Energie Atomique et aux Energies Alternatives).

References

- Aalto R, Nittrouer CA (2012) 210Pb geochronology of flood events in large tropical river systems. *Philos Trans A Math Phys Eng Sci* 370:2040-2074. doi:10.1098/rsta.2011.0607
- Alric R (2009) Recueil des débits caractéristiques de la Nouvelle Calédonie. In: Direction des Affaires Vétérinaires Alimentaires et Rurales (DAVAR), Service de l'eau des statistiques et études rurales. Observatoire de la ressource en eau Nouméa,
- Batista PVG, Laceby JP, Silva MLN, Tassinari D, Bispo DFA, Curi N, Davies J, Quinton JN (2018) Using pedological knowledge to improve sediment source apportionment in tropical environments. *J Soils Sediments* doi:10.1007/s11368-018-2199-5
- Bird ECF, Dubois JP, Ittis JA (1984) The impacts of opencast mining on the rivers and coasts of New Caledonia. Tokyo, Japan, United Nations University.
- Blake WH, Wallbrink PJ, Wilkinson SN, Humphreys GS, Doerr SH, Shakesby RA, Tomkins KM (2009) Deriving hillslope sediment budgets in wildfire-affected forests using fallout radionuclide tracers. *Geomorphology* 104:105-116. doi:10.1016/j.geomorph.2008.08.004
- Bottrill L, Walling D, Leeks G (1999) Geochemical characteristics of overbank deposits and their potential for determining suspended sediment provenance; an example from the River Severn, UK. *Geological Society, London, Special Publications* 163:241-257.
- Brown A, Carpenter R, Walling D (2008) Monitoring the fluvial palynomorph load in a lowland temperate catchment and its relationship to suspended sediment and discharge. *Hydrobiologia* 607:27-40.
- Collins AL, Walling DE, Leeks GJL (1996) Composite fingerprinting of the spatial source of fluvial suspended sediment : a case study of the Exe and Severn river basins, United Kingdom. *Géomorphologie : relief, processus, environnement* 2:41-53. doi:10.3406/morfo.1996.877
- Danloux J, Laganier R (1991) Classification et quantification des phénomènes d'érosion, de transport et de sédimentation sur les bassins touchés par l'exploitation minière en Nouvelle Calédonie.
- David G, Leopold M, Dumas PS, Ferraris J, Herrenschmidt JB, Fontenelle G (2010) Integrated coastal zone management perspectives to ensure the sustainability of coral reefs in New Caledonia. *Mar Pollut Bull* 61:323-334. doi:10.1016/j.marpolbul.2010.06.020
- Davis CM, Fox JF (2009) Sediment fingerprinting: review of the method and future improvements for allocating nonpoint source pollution. *J Environ Eng* 135:490-504.
- Dumas P, Printemps J, Mangeas M, Luneau G (2010) Developing erosion models for integrated coastal zone management: a case study of The New Caledonia west coast. *Mar Pollut Bull* 61:519-529. doi:10.1016/j.marpolbul.2010.06.013
- Evrard O, Laceby JP, Ficetola GF, Gielly L, Huon S, Lefèvre I, Onda Y, Poulenard J (2019) Environmental DNA provides information on sediment sources: A study in catchments affected by Fukushima radioactive fallout. *Sci Total Environ* 665:873-881.
- Evrard O, Laceby JP, Huon S, Lefèvre I, Sengtaheuanghoung O, Ribolzi O (2015) Combining multiple fallout radionuclides (137Cs, 7Be, 210Pbxs) to investigate temporal sediment source dynamics in tropical, ephemeral riverine systems. *J Soils Sediments* 16:1130-1144. doi:10.1007/s11368-015-1316-y
- Ficetola GF, Poulenard J, Sabatier P, Messenger E, Gielly L, Leloup A, Etienne D, Bakke J, Malet E, Fanget B (2018) DNA from lake sediments reveals long-term ecosystem changes after a biological invasion. *Sci Adv* 4:eaar4292.
- Foucher A, Laceby PJ, Salvador-Blanes S, Evrard O, Le Gall M, Lefèvre I, Cerdan O, Rajkumar V, Desmet M (2015) Quantifying the dominant sources of sediment in a drained lowland agricultural catchment: The application of a thorium-based particle size correction in sediment fingerprinting. *Geomorphology* 250:271-281. doi:10.1016/j.geomorph.2015.09.007
- Garcin M (2010) Exploitation des granulats en lit vif des cours d'eau de la Grande-Terre, Nouvelle-Calédonie. BRGM/RP-58531-FR. 114 p., 90 fig., 3 tabl. Bureau des Recherches Géologiques et Minières
- Garcin M, Gastaldi Y, Lesimple S (2017) Quantification et évolution temporelle des apports miniers dans les rivières calédoniennes. BRGM/RP-66840-FR, 44 p., 23 fig., 5. Bureau des Recherches Géologiques et Minières

- Haddadchi A, Ryder DS, Evrard O, Olley J (2013) Sediment fingerprinting in fluvial systems: review of tracers, sediment sources and mixing models. *Int J Sediment Res* 28:560-578.
- Hancock GJ, Wilkinson SN, Hawdon AA, Keen RJ (2014) Use of fallout tracers ^7Be , ^{210}Pb and ^{137}Cs to distinguish the form of sub-surface soil erosion delivering sediment to rivers in large catchments. *Hydrol Process* 28:3855-3874.
- Hedouin L, Bustamante P, Fichez R, Warnau M (2008) The tropical brown alga *Lobophora variegata* as a bioindicator of mining contamination in the New Caledonia lagoon: a field transplantation study. *Mar Environ Res* 66:438-444. doi:10.1016/j.marenvres.2008.07.005
- Hedouin L, Pringault O, Metian M, Bustamante P, Warnau M (2007) Nickel bioaccumulation in bivalves from the New Caledonia lagoon: Seawater and food exposure. *Chemosphere* 66:1449-1457.
- Heintz T, Haapkyla J, Gilbert A (2015) Coral health on reefs near mining sites in New Caledonia. *Dis Aquat Organ* 115:165-173. doi:10.3354/dao02884
- IEOM (2016) Rapport d'activité 2016 de la Nouvelle-Calédonie. Institut d'Emission Outre-Mer
- Koiter A, Owens P, Petticrew E, Lobb D (2013) The behavioural characteristics of sediment properties and their implications for sediment fingerprinting as an approach for identifying sediment sources in river basins. *Earth-Sci Rev* 125:24-42.
- Krause A, Franks S, Kalma J, Loughran R, Rowan J (2003) Multi-parameter fingerprinting of sediment deposition in a small gullied catchment in SE Australia. *Catena* 53:327-348.
- Lacey JP, Evrard O, Smith HG, Blake WH, Olley JM, Minella JPG, Owens PN (2017) The challenges and opportunities of addressing particle size effects in sediment source fingerprinting: A review. *Earth-Sci Rev* 169:85-103. doi:10.1016/j.earscirev.2017.04.009
- Lacey JP, Gellis AC, Koiter AJ, Blake WH, Evrard O (2019) Preface evaluating the response of critical zone processes to human impacts with sediment source fingerprinting. *J Soils Sediments* doi:10.1007/s11368-019-02409-0
- Lacey JP, McMahon J, Evrard O, Olley J (2015) A comparison of geological and statistical approaches to element selection for sediment fingerprinting. *J Soils Sediments* 15:2117-2131. doi:10.1007/s11368-015-1111-9
- Lacey JP, Olley J (2015) An examination of geochemical modelling approaches to tracing sediment sources incorporating distribution mixing and elemental correlations. *Hydrol Process* 29:1669-1685. doi:10.1002/hyp.10287
- Le Gall M, Evrard O, Foucher A, Lacey JP, Salvador-Blanes S, Maniere L, Lefevre I, Cerdan O, Ayrault S (2017) Investigating the temporal dynamics of suspended sediment during flood events with ^7Be and $^{210}\text{Pb}_{\text{xs}}$ measurements in a drained lowland catchment. *Sci Rep* 7:42099. doi:10.1038/srep42099
- Le Gall M, Evrard O, Foucher A, Lacey JP, Salvador-Blanes S, Thil F, Dapoigny A, Lefevre I, Cerdan O, Ayrault S (2016) Quantifying sediment sources in a lowland agricultural catchment pond using (^{137}Cs) activities and radiogenic $(^{87}\text{Sr}/(^{86}\text{Sr}))$ ratios. *Sci Total Environ* 566-567:968-980. doi:10.1016/j.scitotenv.2016.05.093
- Legout C, Poulenard J, Nemery J, Navratil O, Grangeon T, Evrard O, Esteves M (2013) Quantifying suspended sediment sources during runoff events in headwater catchments using spectrophotometry. *J Soils Sediments* 13:1478-1492. doi:10.1007/s11368-013-0728-9
- Martínez-Carreras N, Udelhoven T, Krein A, Gallart F, Iffly JF, Ziebel J, Hoffmann L, Pfister L, Walling DE (2010) The use of sediment colour measured by diffuse reflectance spectrometry to determine sediment sources: application to the Attert River catchment (Luxembourg). *J Hydrol* 382:49-63.
- Mertes LA (1994) Rates of flood-plain sedimentation on the central Amazon River. *Geology* 22:171-174.
- Nyman P, Sheridan GJ, Smith HG, Lane PNJ (2011) Evidence of debris flow occurrence after wildfire in upland catchments of south-east Australia. *Geomorphology* 125:383-401. doi:10.1016/j.geomorph.2010.10.016

- Olley J, Brooks A, Spencer J, Pietsch T, Borombovits D (2013) Subsoil erosion dominates the supply of fine sediment to rivers draining into Princess Charlotte Bay, Australia. *J Environ Radioact* 124:121-129. doi:10.1016/j.jenvrad.2013.04.010
- Olley J, Caitcheon G (2000) Major element chemistry of sediments from the Darling–Barwon river and its tributaries: implications for sediment and phosphorus sources. *Hydrol Process* 14:1159-1175.
- Owens PN, Walling DE (2002) The phosphorus content of fluvial sediment in rural and industrialized river basins. *Water Res* 36:685-701.
- Owens PN, Walling DE, Carton J, Meharg AA, Wright J, Leeks GJL (2001) Downstream changes in the transport and storage of sediment-associated contaminants (P, Cr and PCBs) in agricultural and industrialized drainage basins. *Sci Total Environ* 266:177-186.
- Pascal M, De Forges BR, Le Guyader H, Simberloff D (2008) Mining and other threats to the New Caledonia biodiversity hotspot. *Conserv Biol* 22:498-499.
- Pascal N (2010) Ecosystèmes coralliens de Nouvelle-Calédonie Valeur économique des services écosystémiques Partie I: Valeur financière. IFRECOR Nouvelle-Calédonie, Nouméa 155
- Restrepo JD, Park E, Aquino S, Latrubesse EM (2016) Coral reefs chronically exposed to river sediment plumes in the southwestern Caribbean: Rosario Islands, Colombia. *Sci Total Environ* 553:316-329. doi:10.1016/j.scitotenv.2016.02.140
- Rogers CS (1979) The effect of shading on coral reef structure and function. *J Exp Mar Biol Ecol* 41:269-288.
- Rogers CS (1990) Responses of coral reefs and reef organisms to sedimentation. *Mar Ecol Prog Ser* 62:185-202.
- Sabinot C, Lacombe S (2015) La pêche en tribu face à l'industrie minière dans le sud-est de la Nouvelle-Calédonie. In: *Rev. Société Int. D'Ethnographie* 5, La mer et les Hommes. pp 120-137
- Schwertmann U, Latham M (1986) Properties of iron oxides in some New Caledonian oxisols. *Geoderma* 39:105-123.
- Sevin B (2014) Cartographie du régolithe sur formation ultrabasique de Nouvelle-Calédonie: Localisation dans l'espace et le temps des gisements nickélifères. Nouvelle Calédonie
- Shellberg J, Brooks A, Spencer J Land-use change from indigenous management to cattle grazing initiates the gullyng of alluvial soils in northern Australia. In: *19th World Congress of Soil Science: Soil Solutions for a Changing World*, 2010. pp 1-6
- Sherriff SC, Franks SW, Rowan JS, Fenton O, Ó'hUallacháin D (2015) Uncertainty-based assessment of tracer selection, tracer non-conservativeness and multiple solutions in sediment fingerprinting using synthetic and field data. *J Soils Sediments* 15:2101-2116. doi:10.1007/s11368-015-1123-5
- Smith HG, Sheridan GJ, Lane PNJ, Noske PJ, Heijnis H (2011) Changes to sediment sources following wildfire in a forested upland catchment, southeastern Australia. *Hydrol Process* 25:2878-2889. doi:10.1002/hyp.8050
- Smith HG, Sheridan GJ, Nyman P, Child DP, Lane PNJ, Hotchkis MAC, Jacobsen GE (2012) Quantifying sources of fine sediment supplied to post-fire debris flows using fallout radionuclide tracers. *Geomorphology* 139-140:403-415. doi:10.1016/j.geomorph.2011.11.005
- Stout JC, Belmont P, Schottler SP, Willenbring JK (2013) Identifying Sediment Sources and Sinks in the Root River, Southeastern Minnesota. *Ann Assoc Am Geogr* 104:20-39. doi:10.1080/00045608.2013.843434
- Terry JP, Garimella S, Kostaschuk RA (2002) Rates of floodplain accretion in a tropical island river system impacted by cyclones and large floods. *Geomorphology* 42:171– 182.
- Terry JP, Kostaschuk RA, Wotling G (2008) Features of tropical cyclone-induced flood peaks on Grande Terre, New Caledonia. *Water Environ J* 22:177-183. doi:10.1111/j.1747-6593.2007.00098.x
- Trescases JJ (1973) Weathering and geochemical behaviour of the elements of ultramafic rocks in New Caledonia. *BMRJ Aust Geol and Geophys* 141:149-161.

- Wallbrink P, Murray A, Olley J, Olive L (1998) Determining sources and transit times of suspended sediment in the Murrumbidgee River, New South Wales, Australia, using fallout ¹³⁷Cs and ²¹⁰Pb. *Water Resour Res* 34:879-887.
- Walling D, Peart M, Oldfield F, Thompson R (1979) Suspended sediment sources identified by magnetic measurements. *Nature* 281:110.
- Walling DE (2005) Tracing suspended sediment sources in catchments and river systems. *Sci Total Environ* 344:159-184. doi:10.1016/j.scitotenv.2005.02.011
- Walling DE (2013) The evolution of sediment source fingerprinting investigations in fluvial systems. *J Soils Sediments* 13:1658-1675. doi:10.1007/s11368-013-0767-2
- Walling DE, Owens PN, Carter J, Leeks GJL, Lewis S, Meharg AA, Wright J (2003) Storage of sediment-associated nutrients and contaminants in river channel and floodplain systems. *Applied Geochemistry* 18:195-220.
- Wilkinson SN, Hancock GJ, Bartley R, Hawdon AA, Keen RJ (2013) Using sediment tracing to assess processes and spatial patterns of erosion in grazed rangelands, Burdekin River basin, Australia. *Agric Ecosyst Environ* 180:90-102. doi:10.1016/j.agee.2012.02.002

667 **Tables**

668 **Table 1** Fallout radionuclide activities and geogenic radionuclide contents (mean, M; standard deviation, SD) in
669 the <2 mm and <63 μm fractions of sediment sources and Thio river sediment.

670 **Table 2** Results of Mann-Whitney U test and the DFA used to identify the optimum tracer combination to
671 differentiate sources supplying sediment to the Thio River for the <2 mm particle size fraction with the geogenic
672 approach and for the <63 μm particle size fraction with the geochemical approach.

673 **Table 3** Source contributions calculated by both approaches using gamma spectrometry or X-Ray Fluorescence
674 measurements for sediment deposited during the flood of February 25, 2015.

675 **Table 4** Source contributions calculated by both approaches using gamma spectrometry or X-Ray Fluorescence
676 measurements for sediment deposited during the flood of April 10, 2017.

677

678

679

Table 1

		Mining tributaries (<i>n</i> =16)				Non-mining tributaries (<i>n</i> =8)				Thio river sediment 2015 (<i>n</i> =11)				Thio river sediment 2017 (<i>n</i> =8)			
Fingerprinting property	Particle size	M	SD	Min	Max	M	SD	Min	Max	M	SD	Min	Max	M	SD	Min	Max
¹³⁷ Cs (<i>Bq kg⁻¹</i>)	<2 mm	0.3	0.1	0.1	0.6	0.6	0.3	0.3	1.2	0.7	0.1	0.2	1.3	0.5	0.4	0.1	0.5
	< 63 μm	0.3	0.3	0.01	0.9	1	0.5	0.5	1.4	0.7	0.4	0.2	1.3	ND	ND	ND	ND
²¹⁰ Pb _{xs} (<i>Bq kg⁻¹</i>)	<2 mm	3	3	1	12	10	10	2	34	9	5	4	18	8	5	2	15
	<63 μm	6	7	0.1	19	19	15	7	44	15	8	6	44	ND	ND	ND	ND
K (<i>mg kg⁻¹</i>)	<2 mm	1386	1848	57	5973	12097	3980	5181	17296	4711	2270	1593	9347	2792	880	1691	4292
	<63 μm	1140	1298	121	4742	12051	1394	10060	13697	5969	3030	1979	9240	ND	ND	ND	ND
Th (<i>mg kg⁻¹</i>)	<2 mm	0.6	0.7	0.08	2	6	2	2	7	1	0.6	0.6	6	2	1	0.6	2
	<63 μm	0.5	0.4	0.1	1	7	1	6	8	3	2	0.8	6	ND	ND	ND	ND
U (<i>mg kg⁻¹</i>)	<2 mm	0.1	0.1	0.04	0.5	1.2	0.3	0.3	1.5	0.3	0.1	0.2	1.2	0.4	0.3	0.4	0.4
	<63 μm	0.1	0.1	0.05	0.2	1.3	0.1	1.2	1.5	0.6	0.3	0.2	1	ND	ND	ND	ND

Table 2

	Mann-Whitney U test		DFA- backward mode			
Fingerprinting property	U value	p value	Wilk's Lambda	Variance explained by the variables (%)	Squared Mahalanobis distance	Correctly classified samples (%)
$^{137}\text{Cs} \text{ (Bq kg}^{-1}\text{)}$	38	0.7720	-	-	-	-
$^{210}\text{Pb}_{\text{xs}} \text{ (Bq kg}^{-1}\text{)}$	16	0.1109	-	-	-	-
K $\text{(mg kg}^{-1}\text{)}$	2	< 0.0001	-	-	-	-
Th $\text{(mg kg}^{-1}\text{)}$	2	< 0.0001	-	-	-	-
U $\text{(mg kg}^{-1}\text{)}$	2	< 0.0001	0.1742	82.6	19.6	96
Al $\text{(mg kg}^{-1}\text{)}$	8	0.000	-	-	-	-
Ca $\text{(mg kg}^{-1}\text{)}$	21	0.007	-	-	-	-
Cr $\text{(mg kg}^{-1}\text{)}$	124	< 0.0001	-	-	-	-
Cu $\text{(mg kg}^{-1}\text{)}$	23	0.011	-	-	-	-
Fe $\text{(mg kg}^{-1}\text{)}$	121	0.000	-	-	-	-
K $\text{(mg kg}^{-1}\text{)}$	2	< 0.0001	0.1691	83.1	20.3	96
Mg $\text{(mg kg}^{-1}\text{)}$	119	0.000	-	-	-	-
Mn $\text{(mg kg}^{-1}\text{)}$	108	0.006	-	-	-	-
Ni $\text{(mg kg}^{-1}\text{)}$	125	< 0.0001	-	-	-	-
Si $\text{(g kg}^{-1}\text{)}$	6	< 0.0001	-	-	-	-
Ti $\text{(mg kg}^{-1}\text{)}$	13	0.001	-	-	-	-
Zn $\text{(mg kg}^{-1}\text{)}$	68	0.834	-	-	-	-

Table 3

Sampling point	Mining tributary contribution (%)		Non mining tributary contributions (%)	
	U – Gamma spectrometry	K – XRF	U – Gamma spectrometry	K – XRF
1	97	95	3	5
2	96	95	4	5
3	5	20	95	80
4	64	69	36	31
5	75	46	25	54
6	76	79	24	21
7	11	33	89	67
8	79	86	21	14
9	86	89	14	11
10	80	69	20	31
11	82	63	18	37
M (%)	68	68	32	32
SD (%)	31	25	31	25
Minimum (%)	5	46	3	5
Maximum (%)	97	95	95	80

Table 4

Sampling point	Mining tributary contribution (%)		Non mining tributary contributions (%)	
	U – Gamma spectrometry	K – XRF	U – Gamma spectrometry	K – XRF
1	96	90	4	10
2	96	93	4	7
3	95	90	5	10
4	76	73	24	27
5	84	83	16	17
6	79	81	21	19
7	86	79	14	21
8	86	84	14	16
M (%)	87	84	13	16
SD (%)	8	7	8	7
Minimum (%)	76	73	4	7
Maximum (%)	96	93	24	27

Figure captions

Fig. 1 Main lithologies (a.), the location of rainfall and river monitoring stations (b.) and the sediment samples collected along with tributary source classifications and the location of the active and abandoned mining sites along with mining exploration (c.) conducted in the Thio catchment, New Caledonia

Fig. 2 Box-plots of fallout radionuclide activities (i.e. ^{137}Cs , $^{210}\text{Pb}_{\text{xs}}$) in the <2 mm (a) and the <63 μm (b) fractions of sediment collected on the mining tributaries (Mining), non-mining tributaries (Non mining) and the main Thio River (Thio river sediment (TRS) - flood events of 2015 and 2017). The box indicates the location of the first and third quartiles; the black line indicates the median value; the red line indicates the mean value

Fig. 3 Comparison of ^{137}Cs (a) and $^{210}\text{Pb}_{\text{xs}}$ (b) activities measured in the <63 μm and the <2 mm fractions of sediment samples collected in the Thio River catchment

Fig. 4 Box-plots of geogenic radionuclide contents (i.e. K, Th, U) in the <2 mm fraction of sediment collected on the mining tributaries (Mining), non-mining tributaries (Non mining) and the main Thio River (Thio river sediment (TRS) - flood events of 2015 and 2017). The box indicates the location of the first and third quartiles; the black line

Fig. 5 Comparison of K (a), Th (b) and U (c) contents measured in the <63 μm and the <2 mm fractions of sediment samples collected in the Thio River catchment

Fig. 6 Box-plots of geochemical element contents in the $<63\mu\text{m}$ fraction of sediment collected on the mining tributaries (Mining), non-mining tributaries (Non-mining) and the main Thio River (Thio river sediment (TRS) - flood events of 2015 and 2017). The box indicates the location of the first and third quartiles; the black line indicates the median value; the red line indicates the mean value

Fig. 7 Comparison of geochemical elemental contents measured in the <63 μm and the <2 mm fractions of sediment samples collected in the Thio River catchment

Fig. 8 Relative contributions of mining and non-mining tributaries to the sediment collected in the Thio River (2015 Flood Event) on the <2 mm fraction measurements using the geogenic model (U)

Fig. 9 Relative contributions of mining and non-mining tributaries to the sediment collected in the Thio River (2017 Flood event) based on the <2 mm fraction measurements using the geogenic model (U)

Figure 1

Fig. 1

b.

c.

Figure 2

Fig. 2

Fig. 3

Figure 4

Fig. 4

Fig.5

Figure 6

Fig. 6

Fig. 7

Fig. 8

2015 flood event

Figure 9

Fig. 9

2017 flood event

708 **Supplementary Material**

709 Fig. A1 Spatial rainfall distribution during flood events of February 25, 2015 (a) and April 10, 2017 (b) on the
710 Thio River catchment

711

Supplementary Material

Fig. A1

a.

b.