Cognitive task avoidance correlates with fatigue-induced performance decrement but not with subjective fatigue

Charles-Etienne Benoit ^{a,1}, Oleg Solopchuk ^{a,b,1}, Guillermo Borragán ^{a,c}, Alice Carbonnelle ^a, Sophie Van Durme ^a, Alexandre Zénon ^{a,b}

^a Institute of Neuroscience, Université catholique de Louvain (UCL), B-1200 Brussels, Belgium
^b INCIA, 33076, Bordeaux, France
^c Centre de Recherches en Cognition et Neurosciences and UNI – ULB Neurosciences Institute, Université Libre de Bruxelles (ULB), Belgium, Brussels, Belgium
¹ The first two authors contributed equally to this work

Corresponding author:
Alexandre Zénon, alexandre.zenon@u-bordeaux.fr

Abstract

Mentally demanding tasks feel effortful and are usually avoided. Furthermore, prolonged cognitive engagement leads to mental fatigue, consisting of subjective feeling of exhaustion and decline in performance. Despite the intuitive characterization of fatigue as an increase in subjective effort perception, the effect of fatigue on effort cost has never been tested experimentally. To this end, sixty participants in 2 separate experiments underwent a forced-choice working memory task following either a fatigue-inducing (i.e. cognitive task involving working memory, conflict and switch costs) or a control manipulation. We measured fatigue in terms of subjective feeling and performance decrement and assessed effort in terms of subjective perception and task avoidance. Surprisingly, we found that task avoidance did not systematically change following the fatigue manipulation. However, variations in task avoidance correlated with fatigue-induced performance decline, while the other measures of fatigue and effort were unrelated to each other. Our findings indicate that subjective fatigue develops independently of effort perception and suggest an “anticipatory regulation” model in which fatigue urges subjects to stop in anticipation of possible, future adverse consequences.

Keywords: Fatigue, Cognitive Effort, Neuroeconomics, N-Back task

1. Introduction

Demanding cognitive activities are typically aversive and lead to the percept of effort (Inzlicht et al., 2015; Westbrook and Braver, 2015). The subjective, introspective dimension of cognitive effort is assessed by means of self-report questionnaires, such as the NASA task-load index (Hart and Staveland, 1988), but cognitive effort can also be evaluated behaviorally by assessing task avoidance. This can be done by asking participants to choose between alternative options involving tasks of varied difficulty, associated with rewards of varied magnitude and by using neuroeconomic models to evaluate the effort cost of each task, i.e. how much performing the task discounts the associated reward (Hosking et al., 2014; Westbrook et al., 2013).

While the existence of a cost to cognition appears intuitively obvious, its origin remains unknown (Kurzban et al., 2013; Shenhav et al., 2017; Westbrook and Braver, 2015). What is clear, however, is that prolonged cognitive engagement leads to the subjective feeling of fatigue (Ackerman and Kanfer, 2009; Campagne et al., 2004; Deluca, 2005) and can also deteriorate performance (Holtzer et al., 2011; Schwid et al., 2003; Tanaka, 2015), sometimes referred to as objective fatigue (Bailey et al., 2007). The nature of these behavioral manifestations of cognitive

1
2
3
4 fatigue is diverse, including disruption of inhibition mechanisms (Kato et al., 2009), planning
5 (Lorist, 2008; Lorist et al., 2000; van der Linden et al., 2003), processing of new information
6 (Massar et al., 2010), attention (Boksem et al., 2006; Dorrian et al., 2007; Holtzer et al., 2011;
7 Lim et al., 2010), working memory (Gergelyfi et al., 2015) or task switching (Borragán et al.,
8 2017). Like cognitive effort, the origin of cognitive fatigue remains unclear. Some have proposed
9 that cognitive fatigue would result from metabolic depletion (Christie and Schrater, 2015;
10 Fairclough and Houston, 2004; Gailliot and Baumeister, 2007) but this view has been strongly
11 disputed (Carter et al., 2015; Vadillo et al., 2016). Other hypotheses include the spreading of
12 local sleep patterns in the brain (Krueger et al., 2008; Vyazovskiy et al., 2011), the buildup of a
13 decision signal aimed at limiting the opportunity cost of engaging cognitive resources in the
14 same task (Kurzban et al., 2013) or the extended use of high-consuming strategies aimed at
15 maintaining performance (i.e. Compensatory Control Model (CCM); Hockey, 2013, 1997).
16
17
18
19
20
21
22
23
24

25 One puzzling and pervasive observation in the fatigue literature is that the subjective dimension
26 typically fails to correlate with the behavioral consequences of fatigue (Deluca, 2005). According
27 to Hockey's influential CCM model (Hockey, 2013, 1997), this lack of relation between the
28 subjective and behavioral dimensions would be explained by the fact that subjective fatigue is
29 the manifestation of compensatory resource mobilization, needed to maintain performance
30 despite the progressive deterioration of capacity induced by fatigue. Therefore, initially,
31 subjective fatigue would predominate, while behavioral performance decrement would occur
32 later on in the building up of fatigue. Subjective feelings of fatigue would appear as an indication
33 that the resources needed to accomplish the ongoing demands are at risk or no longer
34 available. In support of this view, subjective task aversion has been shown to increase
35 concomitantly to time on task, even though it is unclear whether this increase is actually caused
36 by fatigue (Lorist et al., 2000). Along the same line, cognitive fatigue increases blood pressure
37 during cognitive task execution, but this effect has complex dependence on the likelihood and
38 importance of the outcome (Stewart et al., 2009; Wright et al., 2007). Finally, neurophysiological
39 activity suggestive of increased engagement during the compensatory phase of cognitive
40 fatigue has also been recently reported (Wang et al., 2016).
41
42
43
44
45
46
47
48
49
50
51
52

53 Phenomenologically, subjective fatigue is often reported as increased aversion for cognition and
54 increased percept of effort (Arai, 1912). In addition, the CCM assumes that subjective fatigue
55 should be accompanied with increased percept of cognitive effort, and presumably increased
56 task avoidance caused by the compensatory engagement of resources. To the extent of our
57 knowledge, the effect of fatigue on these behavioral correlates of effort has never been
58
59
60
61
62
63
64
65

1
2
3
4 experimentally assessed. We hypothesized that subjective fatigue would increase the cost of
5 effort: a fatigued person would feel more effort performing a task of equivalent difficulty and
6 would increase their preference for low-demand tasks. Sixty participants engaged in 2 separate
7 experiments in which subjective effort and task avoidance were measured following either a
8 fatigue-inducing or control manipulation. We evaluated fatigue both in its subjective dimension
9 and in its behavioral manifestation as performance decline, and controlled for other variables
10 such as sleepiness, arousal and motivation. We found that fatigue-related performance
11 decrement correlated with task avoidance, while subjective feelings of fatigue and effort were
12 unrelated to each other.
13
14
15
16
17
18
19
20
21

22 **2. Experiment 1**

23 **2.1. Materials and methods**

24 **2.1.1. Participants**

25
26
27 Thirty right-handed healthy participants took part in Experiment 1 (27 F, 3 M, Age: 22.8 +- 2.5,
28 mean, SD), receiving monetary compensation for their participation (70 - 100€). Each
29 experiment was performed over the span of three consecutive days. This included training on
30 day 1 and fatigue/control manipulation followed by the working memory task on the
31 experimental days 2 and 3, all occurring at the same time during the day, to control for circadian
32 effects between sessions. All participants reported a normal or corrected-to-normal vision, had
33 no auditory impairments or neurological disorders. Experiments were carried out according to
34 the Declaration of Helsinki and were approved by the Ethics Committee of the Université
35 catholique de Louvain. All participants provided a written informed consent.
36
37
38
39
40
41
42
43
44

45 **2.1.2. Subjective evaluation**

46
47 Participant's fatigue was assessed using the French version of the multidimensional fatigue
48 inventory (MFI) (Gentile et al., 2003), composed of fifteen statements to be rated on a 5-point
49 Likert scale (from "1 - Yes, that is true" to "5 - No, that is not true"). The answers to all
50 questions were summed to obtain a Fatigue Index, in which lower score reflected higher fatigue.
51 MFI was evaluated at the beginning and at the end of both experimental days.
52
53
54
55

56 In order to assess sleepiness, we used the 9-point Karolinska sleepiness scale (KSS)
57 (Åkerstedt et al., 2014; Åkerstedt and Gillberg, 1990). Participants had to select a number
58
59
60
61
62
63
64
65

corresponding to their level of sleepiness (from 1 = extremely alert to 9 = very sleepy, fighting against sleep). KSS was evaluated simultaneously with the MFI.

We also evaluated participants' intrinsic motivational state using intrinsic motivation inventory (IMI, <http://selfdeterminationtheory.org/intrinsic-motivation-inventory/>) (Ryan and Deci, 2000). The subjects provided the ratings on a 7-point Likert scale (from "Not at all true" to "Very true"), thus, higher scores represented higher task interest/enjoyment.

Subjective effort was evaluated using the National Aeronautics and Space Administration Task Load Index (NASA-TLX) rating scale (Hart and Staveland, 1988). It is divided in six subscales: mental demand, physical demand, temporal demand, performance, effort and frustration level. In the present study, we focused only on mental demand, performance and effort. Since the concepts of mental demand and effort were difficult to disentangle for participants, and since the scores strongly correlated with each other, we took the average of the mental demand and effort scores as an index of subjective effort cost. The participants had to score each of the items on a 20-point scale presented directly on screen. Participants completed the NASA-TLX after each block of the working memory task on both experimental days.

2.1.3. Fatigue/Control manipulation

On experimental days, participants underwent 2h of either fatigue or control manipulation (see Figure 1). Fatigue was induced by performing a conflict-switch-working memory (CSWm) task in which subjects had to make decision on the basis of conflicting sources of information (i.e. a variant of the Stroop task (MacLeod, 1991)), an approach that has been used widely as a way of promoting cognitive fatigue (Moeller et al., 2012; Pageaux et al., 2015; Wang et al., 2016). In addition, the CSWm task involved two different types of stimulus-response associations and participants had to switch regularly from one to the other, leading to switch costs, also known to promote fatigue (Lorist et al., 2000). Finally, the CSWm task required to maintain information in working memory, since the cue providing instruction for the task was given after the presentation of the stimulus. In the control session, participants watched emotionally neutral documentaries. The order of the sessions (fatigue/control) was counterbalanced between all participants. In the CSWm task, participants were instructed to be as accurate and fast as possible. The time between the cue onset and the initiation of the next trial was adapted to the performance of each participant, according to the following rule:

$$\mathcal{M} = \sum_{t=1}^T \mathcal{M} \times 0.75 + accuracy(t) \times 0.25$$

$$responseTime = \begin{cases} responseTime \times 1.1 & \text{if } \mathcal{M} \leq 0.8 \\ responseTime \times 0.9 & \text{if } \mathcal{M} > 0.8 \end{cases}$$

with T being the current trial number, *responseTime* being initialized to 1500 ms at the beginning of each block, $\mathcal{M}$ being initialized to zero, and *accuracy*(t) representing the accuracy of the response in trial t (one if correct, zero if incorrect). *responseTime* was 300 ms minimum and 3000 ms maximum. This procedure ensured that the task remained challenging for all subjects, independently of their capacity. Each block contained a total of 368 trials, distributed equally between 2 different tasks. In the number task, participants were presented with two numbers, on either side of the fixation point (left and right), and had subsequently to report the location of either the largest in value or the largest in size, depending on an instruction cue. In the arrow task, one arrow was presented either above or below the fixation point and participants were to report either its position or direction, depending on the instruction cue. Trials were equally divided into congruent (e.g. large digit with large value or upwards pointing arrow located on the upper half of the screen) and incongruent (e.g. large digit with small value or upwards pointing arrow located on the lower half of the screen) trials. The participants completed 8 blocks in 2 hours. Each trial proceeded as follows: the fixation cross appeared in the middle of the screen for 1000 ms, followed by 250 ms of stimuli presentation. After a delay, a blank screen appeared for 500-1000 ms (depending on the participant performance), and the instruction cue (“arrow” task: “direction” or “location”; “number” task: “value” or “size”) was presented for 550 ms. Participants had to answer within the cue presentation time, equal to the *responseTime* variable detailed above. The next trial began right after participants’ response. The probability of task switches (defined as a change in either congruency or task instruction) decreased geometrically with each successive trial, such that 45% of trials involved a task switch, 26% followed one trial with the same task, while 12%, 8%, 5% and 2% had 2, 3, 4 and 5 repetitions, respectively.

During the control session, participants watched animal documentaries on a computer for 2h. The documentaries were chosen according to their preference out of 7 different movies. Similar approaches have already been used in other experiments as a control condition (Badin et al., 2016; Van Cutsem et al., 2017; Zering et al., 2017).

2.1.4. Effort discounting evaluation

All tasks were implemented using version 3.0.9 of the Psychtoolbox (Brainard, 1997) in Matlab 7.5 (The MathWorks, Inc. Natick, Massachusetts, United States). Participants faced a 19-inch

CRT screen with a refresh rate of 100Hz. The distance between the screen and the chin support was 58 cm.

To evaluate effort and assess the consequences of fatigue, we used a variant of the auditory “N-back” working memory task, in which the subjects are required to judge whether the presented letter matches the one heard “N” letters before (Kirchner, 1958). Series of letters from the French alphabet, excluding F, J, M, P, W, Y, were presented aurally to participants. These letters were recorded from a young female voice and then normalized for consistency in pitch and volume. In each block, 25% of the letters were defined as targets, the others being considered as distractors. Task difficulty ‘N’ was calibrated individually during the training on the first day, and stayed constant during the whole experiment. The training included 15 blocks of increasing demand (N = 1 to 5), composed of 60 letters each. The letters were presented continuously while participants fixated the cross in the center of the screen. Response timeout was fixed to 1.5 seconds (indicated by change of fixation cross orientation), with an inter-stimulus interval of 2 to 3 seconds (uniformly distributed). A short pause was provided after every 5 blocks. The answers were collected via computer keyboard (distractor and target corresponded to button number ‘1’ and ‘2’ respectively). When the participant performance was greater than or equal to 70% (mean of the % of targets detected and the % of distractors correctly rejected) they progressed to the next level with a maximum difficulty level set to 5-back, otherwise they repeated the same level once more. The maximal difficulty reached by the participants in the last block was then used for all subsequent experimental acquisitions (see Figure 2C).

Following the fatigue or control manipulation on experimental days, participants performed the N-back tasks of different difficulty levels, presented in the context of a forced-choice paradigm, inspired from earlier studies (Westbrook et al., 2013). In order to avoid that participants’ preference for difficulty-reward combinations would be influenced by their auto-evaluation of performance, rather than by effort perception, we defined task difficulty as the ratio of 1-back and N-back tasks in the block of 80 letters (actual ‘N’ depending on participants’ performance during training, cfr. supra). These ratios are referred to as the block difficulty level (DL): % of N-back: DL-1 = 25%, DL-2 = 37.5%, DL-3 = 50%, DL-4 = 62.5%, DL-5 = 75%). The difficulty was represented visually using horizontal color bars (blue = 1-back; orange = n-back; see Figure 1). A fixed amount of 2€ was always associated with the DL-5 block while a variable amount, ranging between 0 and 2 € (with 0.20 € intervals) was offered for the easy option (i.e. DL-1 to DL-4). Participants had to make the choice by pressing the right or left mouse button depending

on whether they wanted to execute the task with the DL presented on the right or left part of the screen. The side of easy and difficult offers was pseudo-randomized for every participant. All possible difficulty pairings were evaluated for each possible reward level, which resulted in 44 forced-choices. Within these selections, participant had to perform the actual N-back task for 15 randomly selected choices. Any of the forced-choices could be selected for execution since the randomization did not take into account the proposed difficulty levels. Participants had to perform the selected block immediately following the choice. In each block, the accuracy (average percentage of target hits and distractor correct rejections) was averaged between both difficulty levels (1-back and N-back) to calculate the performance. This particular procedure ensured that, regardless of DL, performance in the 1-back and in the N-back tasks had the same impact on the amount of money earned at the end of the block. For instance, if participants had average performance of 90% in the easy and 60% in the difficult task, their reward would be 75% of the proposed amount, irrespective of the relative proportion of easy and difficult trials (DL). This provided incentives for participants to disregard their relative performance in the easy and difficult tasks, and to consider only cognitive effort associated with the DL when making their choice. The remuneration was calculated by multiplying the performance by the reward attributed to the selected block (i.e. 0 to 2 €). This information was provided on the screen at the end of each block. Prior to receiving performance feedback, participants had to answer the NASA-LTX questionnaire, presented on the computer screen.

2.1.5. Statistical analyses

We analyzed data with Matlab (The MathWorks, Inc., Natick, MA, USA) and performed statistical analyses with JASP (Version 0.8.2, JASP Team 2017). We performed both frequentist and Bayesian repeated-measure ANOVAs and Kendall correlations, to avoid undue influence of outliers. Results from the frequentist analyses are reported in terms of p-value (and F statistics for the ANOVA), whereas Bayesian analyses resulted in Bayes Factors (BF). BF represents the ratio of the likelihood of two different models with, most classically, one assuming the alternative hypothesis and the other assuming the null hypothesis. Bayesian analysis was used in the present study because it provides a better estimate of effect reliability (BF) than the p-value. In addition, it provides the ability to evaluate the probability that the null hypothesis is true. Finally, it mitigates considerably the issue of multiple comparison (Gelman and Tuerlinckx, 2000). The BF can be interpreted on the basis of the table proposed in Kass and Raftery (1995):

$\log BF_{10}$	BF_{10}	Evidence against H_0
0 to 1/2	1 to 3,2	Not worth more than a bare mention
1/2 to 1	3,2 to 10	Substantial
1 to 2	10 to 100	Strong
> 2	> 100	Decisive

For 1-way ANOVAs and correlation analyses, the BF_{10} is reported, which represents the comparison between the alternative and the null hypothesis. For 2-way ANOVAs, $BF_{inclusion}$ is reported instead. $BF_{inclusion}$ is obtained by comparing the likelihood of the model including, or excluding a given effect.

The behavior of the participant in the choice task was analyzed with a computational model of effort-based decision making inspired from earlier studies (Zénon et al., 2016)

$$Q(\alpha, \beta, R, D) = \alpha[(1 - \beta)\frac{R - 100}{10} + \beta(D - 1)]$$

$$\alpha = \frac{1}{e^{temperature}}$$

$$\beta = \frac{1}{1 + e^{-weight}}$$

$$P(easy|temperature, weight, R, D) = \frac{1}{1 + e^{-Q}}$$

with *temperature* and *weight* as free model parameters. R represented the amount of reward proposed for the difficult version of the task, and D accounted for the difficulty level (ranges of R and D in the model corresponded to their range in the actual experiment). Q represented an estimate of the net value of the proposed trial. $P(easy)$ indicated the probability of choosing the easy option, which was a function of the temperature and weight parameters and of the reward and difficulty conditions. The model was fit with multidimensional nonlinear minimization (Nelder-Mead, `fminsearch`) with ridge regularization.

The *weight* parameter can be interpreted as representing the relative impact of the difficulty level on the decisions of the participants, while the *temperature* parameter indicated how noisy their choices were. Increasing the weight parameter results in more effort-discounting (the

difficulty parameter D has more weight on the decision), while increasing the temperature parameter resulted in choices that were more stochastic (i.e. depended less on net value).

2.2. Results

CSWm task showed time-on-task increase in conflict and switch cost

Performance in the CSWm task was $92\pm3\%$, $93\pm2\%$, $45\pm2\%$ and $80\pm3\%$ (mean $\pm$ 95% confidence interval) for congruent/switch, congruent/no-switch, incongruent/switch and incongruent/no-switch trials, respectively, confirming that the task was challenging for the participants. We measured the difference in average performance block by block between congruent and incongruent conditions, either in switch or no-switch trials. We found that this incongruency effect increased over time in the switch (Bayesian t-test on Kendal correlation coefficients with block number; $BF_{10}=29637$), but decreased in no-switch trials ($BF_{10}=5.34$). Conversely, we found that the effect of switch on accuracy increased also over time in the incongruent ($BF_{10}=19.66$) but decreased in the congruent trials ($BF_{10}=28.80$). We considered the increase in congruency and switch effect over time as potential indicators of fatigue buildup and correlated the corresponding correlation coefficients with the other subjective and behavioral measures (NASA scores, N-back performance scores, model parameters, MFI, IMI and KSS). However, none of these correlations were significant ($BF_{10}<3$).

CSWm session increases MFI and subjective effort but not task avoidance or performance

This first analysis aimed at assessing the effect of the fatigue-inducing task on our different dependent variables. We performed a one-way repeated-measure ANOVA on the post-pre difference in MFI, KSS and IMI scores with SESSION as factor (control vs fatigue). There was no effect of SESSION on either KSS ($F(1,29)=2.188$, $p=0.15$, $BF_{10}=0.759$) or IMI ($F(1,29)=1.965$, $p=0.172$, $BF_{10}=0.672$) and a small influence of SESSION on MFI (see Figure 3A; $F(1,29)=5.075$, $p=0.032$, $BF_{10}=1.893$; i.e. significant according to frequentist statistics but anecdotal for Bayesian statistics). We then correlated these fatigue-related changes in subjective measures between each other. We found a very strong positive correlation between fatigue-control changes in MFI and KSS (Kendall's tau=0.508, $p<0.0001$, $BF_{10}=283.292$), suggesting that the subjects failed to distinguish clearly the subjective feeling of fatigue from that of sleepiness. Finally, neither KSS nor MFI correlated with IMI (all $p > 0.555$, all $BF_{10} < 2.33$), indicating that subjects' intrinsic motivation was not affected by fatigue manipulation, in accordance with previous studies (Gergelyfi et al., 2015).

We then looked at the effect of fatigue and difficulty level on the NASA questionnaires. We found that the NASA subjective effort and performance scores changed as a function of the DIFFICULTY level (2-way RM-ANOVA: NASA effort: $F(4,116)=9.564$, $p<0.001$, $BF_{inclusion}=4281$, NASA performance: $F(4,116)=4.767$, $p=0.001$, $BF_{inclusion}=17.9$). In addition, NASA effort, but not NASA performance changed with SESSION (see Figure 3C; NASA effort: $F(1,29)=4.581$, $p=0.041$, $BF_{inclusion}=361$, NASA performance: $F(1,29)=0.963$, $p=0.335$, $BF_{inclusion}=0.228$). There was no SESSION X DIFFICULTY interaction (NASA effort: $F(4,116)=0.180$, $p=0.949$, $BF_{inclusion}=0.025$, NASA performance: $F(4,116)=0.163$, $p=0.956$, $BF_{inclusion}=0.027$). There was no correlation between the fatigue-related changes in the two variables (Kendall's tau=-0.080, $p=0.547$, $BF_{10}=0.284$).

The temperature and weight parameters from the task avoidance models were also subjected to a one-way repeated-measure ANOVA with SESSION as factor (control vs fatigue). The weight parameter can be interpreted as an index of behavioral effort cost, while temperature indicates the level of randomness of participants' behavior. There was no change in either parameters between sessions (see Figure 2A, left panel, and Figure 3B; temperature: $F(1,29)=0.179$, $p=0.676$, $BF_{10}=0.285$, weight: $F(1,29)=0.677$, $p=0.418$, $BF_{10}=0.353$). There was a strong correlation between the fatigue-control changes in the two variables (Kendall's tau=0.526, $p<0.001$, $BF_{10}=704.0$). However, we noticed that many participants systematically picked the difficult task, irrespective of the reward value (13 subjects picked the difficult task in 100% of choices in at least one session). On average the percentage of choice of the difficult task was $86.4\pm2.7\%$ and $87.4\pm2.6\%$ for the fatigue and the control sessions, respectively (mean $\pm$ CI).

In terms of fatigue-related changes in task performance, a 2-way RM-ANOVA on the logit-transformed accuracy scores showed, besides the obvious difference between 1-back and N-back TASK (accuracy: $F(1,29)=165.363$, $p<0.001$, $BF_{10}>10^{29}$; RT: $F(1,29)=83.510$, $p<0.001$, $BF_{10}>10^{16}$), an absence of main SESSION effect (accuracy: $F(1,29)=2.475$, $p=0.127$, $BF_{10}=0.374$, RT: $F(1,29)=1.124$, $p=0.298$, $BF_{10}=0.354$) or interaction between the TASK and the SESSION effect (see Figure 2B; accuracy: $F(1,29)=2.651$, $p=0.114$, $BF_{10}=0.550$; RT: $F(1,29)=0.167$, $p=0.686$, $BF_{10}=0.266$).

Changes in performance correlate with changes in task avoidance

We investigated the link between the fatigue-related changes in task avoidance and the other effort and fatigue variables. We found strong evidence for a correlation between fatigue-induced changes in Model Weight and performance in the N-back task (see Figure 4A; Tau=-0.421, $p<0.001$, $BF_{10}=39.4$) but not with 1-back performance (see Figure 4A; Tau=-0.108, $p=0.415$,

BF₁₀=0.33). This correlation showed that participants exhibited larger effort-discounting when they had decreased performance in the fatigue session. In contrast, Model Weight failed to correlate with MFI (see Figure 4D; Tau=-0.005, p=0.972, BF₁₀=0.235) or NASA scores (see Figure 4B; effort: Tau=0.122, p=0.357, BF₁₀=0.362; performance: Tau=0.218, p=0.094, BF₁₀=0.938) changes.

Given the correlation between fatigue-related changes in task avoidance and N-back performance, we also looked at the correlation between changes in subjective effort and N-back performance. None of the correlations were significant (neither 1-back nor N-back task performance correlated with NASA performance or effort scores; see Figure 4C; $0.2 < BF_{10} < 0.3$).

Subjective fatigue correlates with changes in subjective effort and sleepiness

We also looked at the relation between changes in subjective fatigue and the effort variables. There was a moderate correlation between NASA Effort and MFI (Tau=0.306, p=0.018, BF₁₀=3.542) and between NASA Effort and KSS (Tau=0.341, p=0.011, BF₁₀=6.798). Finally, as classically reported in the literature, there was no correlation between the change in MFI and the change in performance (1-back: Tau=-0.133, p=0.323, BF₁₀=0.392; N-back: Tau=-0.070, p=0.592, BF₁₀=0.271).

2.3. Discussion

In experiment 1, we found only a moderate effect of the CSWm task on the subjective fatigue scores. However, the MFI questionnaire was provided at the end of the whole experiment and therefore, we suspected that the results were affected strongly by the performance of the intervening N-back task. In addition, MFI correlated strongly with KSS, suggesting that the participants failed to distinguish the concept of fatigue from that of sleepiness.

Subjective effort, as assessed through the NASA-TLX questionnaire, increased following the CSWm task, and its change correlated with that of MFI, apparently confirming our hypothesis that MF increases subjective effort cost. However, our other estimate of effort, task avoidance, correlated with behavioral (N-back performance), rather than subjective fatigue (MFI). One issue with the task avoidance estimate was that many participants always selected the difficult task, providing us with no information about their actual estimation of behavioral effort cost. This was presumably caused by their excessive motivation for the monetary reward.

To address these multiple methodological issues, we ran a second experiment to replicate the

findings in a more refined protocol. One supplementary MFI assessment was added right after the CSWm and documentary-watching sessions. The confound between sleepiness and fatigue was addressed by explaining in detail to each participant the difference between the 2 concepts (Borragán et al., 2017). In addition, in order to restore vigilance, we added an “arousal boost” after the CSWm and documentary-watching sessions and before the final evaluation (to equalize the conditions of questionnaire assessment). Also, we measured arousal by means of pupillometry. The assessment of task avoidance was improved by using virtual coins, rather than directly euros as rewards for the participants. These coins were translated into actual money at the end of the experiment but the participants were not aware of the conversion ratio. Moreover, instead of using all the combinations of difficulty and reward, we used an adaptive procedure to choose the conditions in each trial. Finally, the selection of N-back task difficulty during the training session was also less constrained, with no maximal value for the ‘N’ of the N-back.

3. Experiment 2

3.1. Materials and methods

Thirty right-handed healthy participants took part in Experiment 2 (19 F, 11 M, Age: 23.9±4.0). A larger proportion of male participants were sought after since males are known to consistently take more financial risks (Charness and Gneezy, 2012). MFI and KSS were assessed at the beginning and at the end of both experimental days, but also between the fatigue/control manipulation and behavioral effort-discounting task, following an arousal boost which consisted in running on the spot for 30 seconds. The arousal boost was introduced to minimize sleepiness. Since it was not expected to have any effect on fatigue, we expected this manipulation to make the MFI a purer measure of fatigue, dissociated from the confound of sleepiness. Each participant read the NASA-TLX instructions carefully prior to the experiment, in order to ensure their appropriate understanding of the different subscales.

The first phase of the training session in Experiment 2 was similar to that of Experiment 1 and comprised 10 blocks, now with 80 letters each. Participants were then engaged in 6 blocks of N-back task with mixed difficulties, similar to those experienced during the following experimental sessions. Different ratios of 1 back & N-back tasks were performed, while the difficulty level of the N-back (i.e. the value of the N) was adapted as a function of participants’ individual ability (see Figure 2C). When the participants’ performance was greater than or equal to 75% (mean of

the % of targets detected and the % of distractors correctly rejected) they progressed to the next level. If their performance was below 60%, they reverted back to the inferior N value. Otherwise they kept practicing the same level. We did not impose maximum difficulty level for the n-back. One hundred and twenty letters were presented in these blocks. Only 3 DL were now evaluated (% of N-back: DL-1 = 20%, DL-2 = 50%, DL-3 = 80%). The monetary incentive was switched from actual money to a point-based system, and the conversion rate was not disclosed in advance to the participants.

At the beginning of each experimental session, participants performed one warm-up block (DL-3). They then had to respond to 40 forced choices, out of which 10 randomly selected blocks were actually performed.

Instead of an exhaustive screening of all possible pairings between difficulty and reward, an adaptive procedure was implemented (Kontsevich and Tyler, 1999; Watson and Pelli, 1983). The idea behind this procedure is to evaluate the expected gain in information (i.e. decrease in entropy) for each possible condition. The algorithm then chooses the reward/difficulty condition that maximizes this expected information gain. Participants' choices were modeled with the same 2-parameter effort-based decision making model as described in Experiment 1. The prior probability distribution P_0 of the weight and temperature parameters were set to a normal distribution with mean zero and standard deviation equal to 5. In each trial, this probability distribution was updated as a function of participant's choice:

$$\begin{aligned} \log(P(\text{weight}, \text{temperature} | \text{choice}, R, D)) \\ &= \log(P(\text{choice} | \text{weight}, \text{temperature}, R, D)) + \log(P_0(\text{weight}, \text{temperature})) \\ &- \log\left(\sum_w \sum_t P(\text{choice} | w, t, R, E) \times P_0(w, t)\right) \end{aligned}$$

$$P(\text{choice} | \text{weight}, \text{temperature}, R, D) = \begin{cases} P(\text{easy} | \text{weight}, \text{temperature}, R, D), & \text{if easy option chosen} \\ 1 - P(\text{easy} | \text{weight}, \text{temperature}, R, D), & \text{if difficult option chosen} \end{cases}$$

Then the expected amount of information gained in the next trial was computed for each condition, each weight and temperature values and each possible choice, and allowed us to extract a probability of choosing a given reward and difficulty condition, $P(R, D)$, according to the following equations:

$$\begin{aligned} \log(P(\text{weight}, \text{temperature}, \text{choice} | R, D)) \\ &= \log(P(\text{choice} | \text{weight}, \text{temperature}, R, D)) + \log(P_0(\text{weight}, \text{temperature})) \\ P(\text{choice} | R, D) &= \sum_w \sum_t P(w, t, \text{choice} | R, D) \end{aligned}$$

$$\begin{aligned}
& \log(P(\text{weight}, \text{temperature} | \text{choice}, R, D)) \\
&= \log(P(\text{weight}, \text{temperature}, \text{choice} | R, D)) - \log\left(\sum_w \sum_t P(w, t, \text{choice} | R, D)\right) \\
& H(\text{choice}, R, D) = - \sum_w \sum_t P(w, t | \text{choice}, R, D) \times \log(P(w, t | \text{choice}, R, D)) \\
& H(R, D) = P(\text{easy} | R, D) \times H(\text{easy}, R, D) + P(\text{difficult} | R, D) \times H(\text{difficult}, R, D) \\
& P(R, D) = \frac{e^{-100 \times (H(R, D) - \min(H(R, D)))}}{\sum_r \sum_d e^{-100 \times (H(r, d) - \min(H(r, d)))}}
\end{aligned}$$

All other parameters and procedures were identical to Experiment 1.

In order to monitor participant's level of arousal, we recorded pupil size during performance of the N-back task. Pupil size was acquired using an Eyelink 1000+ eye tracker video-based system (SR Research Ltd., Kanata, Ontario, Canada), with a sampling frequency of 500 Hz. We also evaluated maximal pupil dilation by asking the subjects to apply physical force (maximal voluntary contraction, MVC) through a handgrip on a dynamometer Jamar® Hydraulic Hand Dynamometer). This step was performed twice at the beginning and at the end of both experimental days. No pupillometry data were acquired on the training day.

3.2. Results

CSWm task showed time-on-task increase in conflict and switch cost

In Experiment 2, performance in the CSWm task was 93±1%, 95±1%, 44±2% and 84±2% (mean±95% confidence interval) for congruent/switch, congruent/no-switch, incongruent/switch and incongruent/no-switch trials, respectively. Similarly to Experiment 1, the incongruency effect increased over time in the switch ($BF_{10} > 10^6$), but decreased in no-switch trials ($BF_{10} = 8.33$), while the effect of switch on accuracy increased over time in the incongruent ($BF_{10} > 60000$) but decreased in the congruent trials ($BF_{10} = 924$). Again, correlation of these correlation coefficients with the other subjective and behavioral measures (NASA scores, N-back performance scores, model parameters, MFI, IMI and KSS) resulted in no significant outcome ($BF_{10} < 3$).

CSWm session increases MFI, decreases performance but leaves task avoidance unchanged

We investigated the effect of the fatigue-inducing task on the dependent variables. A repeated-measure ANOVA on the post-pre difference in MFI and KSS scores showed a strong effect of SESSION on KSS (see Figure 3D; $F(1,29) = 10.69$, $p = 0.003$, $BF_{10} = 15.32$) and a decisive effect on the MFI ($F(1,29) = 23.06$, $p < 0.001$, $BF_{10} = 408.2$). There was a moderate correlation between the fatigue-control changes in the two variables (Kendall's tau=0.303, $p = 0.026$, $BF_{10} = 3.377$),

showing that the changes brought to Experiment 2 were only partly effective in separating fatigue and sleepiness (reduction of the correlation coefficient from 0.5 to 0.3). Similarly to Experiment 1, there was neither SESSION effect on IMI ($F(1,29)=1.965$, $p<0.172$, $BF_{10}=0.657$), nor correlation between IMI and MFI (Kendall's $\tau=0.016$, $p=0.900$, $BF_{10}=0.237$) or IMI and KSS (Kendall's $\tau=-0.015$, $p=0.914$, $BF_{10}=0.237$).

Regarding the NASA-TLX variables, we confirmed that the NASA effort and performance scores changed as a function of the DIFFICULTY level (NASA effort: $F(2,58)=26.654$, $p<0.001$, $BF_{inclusion}=2.880 \times 10^7$, NASA performance: $F(2,58)=14.884$, $p<0.001$, $BF_{inclusion}=10126$) but not SESSION (see Figure 3F; NASA effort: $F(1,29)=0.294$, $p=0.592$, $BF_{inclusion}=0.203$, NASA performance: $F(1,29)=0.124$, $p=0.728$, $BF_{inclusion}=0.171$) or SESSION X DIFFICULTY interaction (NASA effort: $F(2,58)=0.929$, $p=0.401$, $BF_{inclusion}=0.166$, NASA performance: $F(2,58)=0.257$, $p=0.775$, $BF_{inclusion}=0.116$). This time, there was a moderate, negative correlation between the fatigue-related changes in the two variables (Kendall's $\tau=-0.343$, $p=0.007$, $BF_{10}=7.037$).

We then looked at the task avoidance variables. The temperature and weight parameters from the task avoidance models were also subjected to a one-way Bayesian repeated-measure ANOVA with SESSION (control vs fatigue) as factor. There was no change in either parameters (see Figure 2A, right panel and Figure 3E; temperature: $F(1,29)=1.868$, $p=0.182$, $BF_{10}=0.594$, weight: $F(1,29)=0.001$, $p=0.981$, $BF_{10}=0.263$) between sessions. There was no correlation between the fatigue-control changes in the two variables (Kendall's $\tau=0.222$, $p=0.092$, $BF_{10}=0.978$). The percentage of acceptance of the difficult option in Experiment 2 was $61.0 \pm 4.2\%$ in the control and $58.4 \pm 5.4\%$ in the fatigue session (mean $\pm$ CI). None of the participants picked systematically the difficult option (all acceptance percentages were below 90%).

In Experiment 2, we found that fatigue had a significant impact on N-back task performance (see figure 2B). A RM-ANOVA on the logit-transformed accuracy scores showed, beside the difference between 1-back and N-back TASK (accuracy: $F(1,29)=299.4$, $p<0.001$, $BF_{10}>10^{37}$, RT: $F(1,29)=56.587$, $p<0.001$, $BF_{10}>10^{15}$), a moderate interaction between the TASK and the SESSION effect on accuracy ($F(1,29)=7.789$, $p=0.009$, $BF_{10}=4.979$). This interaction was absent from the RT data ($F(1,29)=0.105$, $p=0.748$, $BF_{10}=0.255$). The main SESSION effect was only anecdotal (accuracy: $F(1,29)=11.354$, $p=0.002$, $BF_{10}=2.593$, RT: $F(1,29)=8.024$, $p=0.008$, $BF_{10}=1.159$). Looking at the accuracy in each task individually, we found a strong decrease in performance with fatigue in the 1-back task ($F(1,29)=10.34$, $p=0.003$, $BF_{10}=12.97$) but not in the N-back task ($F(1,29)=0.132$, $p=0.719$, $BF_{10}=0.276$). There was no correlation between the fatigue-control

changes in the two variables (Kendall's tau=0.145, $p=0.272$, $BF_{10}=0.432$).

Decrease in performance correlates with changes in task avoidance and subjective effort

In agreement with Experiment 1, we found strong evidence for a correlation between fatigue-induced changes in Model Weight and performance (see Figure 4E; N-back: Tau=-0.374, $p=0.004$, $BF_{10}=13.376$; 1-back: Tau=-0.374, $p=0.004$, $BF_{10}=13.376$). In contrast, Model Weight changes failed to correlate with MFI (see Figure 4H; Tau=-0.082, $p=0.531$, $BF_{10}=0.286$), KSS (Tau=-0.223, $p=0.110$, $BF_{10}=0.999$) or NASA score changes (see Figure 4F; effort: Tau=0.109, $p=0.401$, $BF_{10}=0.332$; performance: Tau=-0.095, $p=0.464$, $BF_{10}=0.306$).

There was also a positive correlation between fatigue-induced changes in performance in the N-back task and NASA performance score (see Figure 4G, blue; N-back: Tau=0.393, $p=0.002$, $BF_{10}=20.615$; 1-back performance: Tau=0.264, $p=0.041$, $BF_{10}=1.784$), and a moderate correlation of the performance changes with the NASA effort score (see Figure 4G, red; N-back: Tau=-0.306, $p=0.018$, $BF_{10}=3.531$; 1-back: Tau=-0.149, $p=0.256$, $BF_{10}=0.450$).

No relation between fatigue and arousal

In Experiment 2, we measured pupil size in order to disentangle cognitive fatigue from unspecific time-on-task effects on arousal. Baseline arousal levels remained unaffected by the fatigue manipulation, as indexed by average pupil size during the MVC task ($F(1,29)=0.250$, $p=0.621$, $BF_{10}=0.281$). The change in pupil-indexed arousal correlated moderately with the change in KSS score but not with the MFI score change (KSS: Tau=0.302, $p=0.025$, $BF_{10}=3.294$; MFI: Tau=-0.005, $p=0.971$, $BF_{10}=0.235$).

A 2-way Bayesian RM-ANOVA on the peak pupil response during the task showed a moderate effect of session ($F(1,29)=1.912$, $p=0.177$, $BF_{inclusion}=3.157$), with larger pupil responses in the fatigue session. There was no DIFFICULTY ($F(2,58)=1.573$, $p=0.216$, $BF_{inclusion}=0.092$) or interaction ($F(2,58)=2.662$, $p=0.078$, $BF_{inclusion}=0.245$) effect. The between-session change in pupil size failed to correlate with any of the fatigue measures ($BF_{inclusion}$ between 0.2 and 0.3).

Subjective fatigue failed to correlate with the other fatigue or effort measures

Beside the previous correlations reported above, there was no correlation between NASA Effort and MFI (Tau=-0.012, $p=0.929$, $BF_{10}=0.236$) or KSS (Tau=-0.090, $p=0.504$, $BF_{10}=0.238$). Finally, there was no correlation between the change in MFI and the change in performance (1-back: Tau=-0.100, $p=0.442$, $BF_{10}=0.357$; N-back: Tau=0.063, $p=0.629$, $BF_{10}=0.264$).

3.3. Discussion

Important methodological limitations from Experiment 1 were addressed in the second experiment. After observing that behavioral effort cost was not affected by fatigue, and that many participants never selected the easy task, we hypothesized that the N-back task may not have been challenging enough and that the value of reward was too high relative to effort cost. Therefore, in Experiment 2 we did not restrict the difficulty of the working memory task, such that some participants were able to perform a 9-back version of the task. In order to decrease the incentive value of reward, we switched from using actual money to a point-based system. Finally, instead of testing exhaustively every possible combination of task difficulty and reward, we implemented a Bayesian search procedure, in which the proposed options were adapted based on the previous responses. Despite these improvements, one remaining limitation is noteworthy. We tested young university students, who are generally strongly motivated by reward and this may have masked partly the increases in behavioral effort costs associated with fatigue.

In experiment 2, we found a decisive effect of the CSWm task on the subjective fatigue scores, evaluated immediately following the fatigue-inducing manipulation, and on N-back performance. In contrast to Experiment 1, MFI correlated only moderately with KSS, confirming that better explanation of the fatigue and sleepiness constructs to the participants and/or the arousal boost manipulation decreased the confounding effect of sleepiness on fatigue. Average tonic pupil size, used as an independent measure of arousal, showed no systematic change across sessions, but correlated with the subjective sleepiness measure. Interestingly, however, we found that peak pupillary responses, thought to be indicative of the phasic variations in noradrenaline and/or acetylcholine (Aston-Jones and Cohen, 2005; Joshi et al., 2016; Reimer et al., 2016), increased following the fatigue manipulation, in agreement with earlier studies (Wright et al., 2007).

Another difference with respect to Experiment 1 was that the fatigue manipulation failed to have any impact on the subjective effort perception measures. We can only speculate on the reason for this difference, but one may suggest that subjective effort increased in Experiment 1 following the CSWm task because of increased sleepiness, rather than because of cognitive fatigue. In Experiment 2, since the impact of sleepiness was decreased, subjective effort did not change significantly.

Finally, the most important outcome of Experiment 2 is that we were able to replicate the two main findings from Experiment 1: task avoidance correlated with fatigue-induced performance

decrement but not with subjective fatigue.

4. General discussion

In the present paper, we hypothesized that cognitive fatigue would increase the cost of cognitive effort. We evaluated fatigue and effort in terms of subjective feeling, as well as in terms of behavioral manifestation such as task performance and reward discounting, respectively. We found that extensive involvement in a cognitively demanding task increased the feeling of fatigue and impaired task performance in the easy version of the task. Throughout the experiment, preference of the participants between task options reflected the attribution of a cost to cognitive effort (Westbrook et al., 2013). However, contrary to the hypothesis, effort cost did not increase with subjective fatigue but increased rather in proportion to fatigue-induced decrease in performance.

The cognitive mechanisms and neural substrates of effort and fatigue remain poorly understood. If subjective fatigue was a protective mechanism against forthcoming resource depletion (Gergelyfi et al., 2015; Muraven and Slessareva, 2003), or if it was the manifestation of increased engagement to maintain performance despite resource disruption (Hockey, 1997), it should also have increased task avoidance. The present findings go clearly against this hypothesis, by showing an absence of link between subjective fatigue and task avoidance. It is worth mentioning that we found a positive correlation between subjective fatigue and subjective effort in Experiment 1. However, this correlation was weak and disappeared in Experiment 2, in which we addressed multiple methodological limitations. In contrast, task avoidance correlated in both experiments with fatigue-related performance decrement. These findings emphasize the dissociation between the subjective and behavioral manifestations of fatigue and effort. Indeed, in both experiments, subjective and behavioral assessments of fatigue and effort failed to correlate with each other.

Parts of the present findings rest on the exclusion of the alternative hypothesis. Bayesian statistics, by considering that probabilities can be attributed to hypotheses (e.g. there is a 10% chance of raining tomorrow), allow us to evaluate the differences in likelihood between the null and alternative hypotheses in terms of Bayes Factor. Here, the correlation between MFI and model weight, reflecting task avoidance, resulted in both experiments in substantial evidence for the null hypothesis (Bayes Factor < 0.3). This replication of the null result provides, to our view, strong evidence that MFI, as a general measure of subjective fatigue, is not associated with increased cognitive task avoidance. However, it is important to note that this finding does not demonstrate that other phenomenological aspects of fatigue, that would fail to be captured by

the MFI, could not lead to increased task avoidance. Along the same line, it is possible that longer task performance, or more demanding tasks, could have led to increased task avoidance. Nevertheless, 2 hours of a task known to tax cognitive control resources, and in which difficulty was adjusted to lead to performance well below maximum, were not enough to induce increased task avoidance in the present experiment (BF below 0.3 in Exp2 and slightly above in Exp1).

We found that despite successful fatigue inducement, the participants maintained similar performance in the N-back task but showed significant decrease in 1-back performance. Our initial hypothesis was that fatigue was associated with disruption of vigilance, which typically leads to more mistakes in easy and monotonous tasks (Thiffault and Bergeron, 2003). We accounted for this possibility in Experiment 2, by adding an arousal boost manipulation to restore vigilance, by improving the dissociation between fatigue and sleepiness self-evaluation in participants, and by measuring participants' pupil size (Sara and Bouret, 2012; Zénon et al., 2014). We found neither change of arousal between the sessions nor any correlation with the measures of fatigue, sleepiness and performance. This suggests that the decrement in 1-back performance was not caused by decreased vigilance, even though measuring vigilance directly with psychomotor vigilance task would be necessary to confirm this claim more decisively. Therefore, understanding why fatigue disrupts specifically 1-back performance will require further investigation.

One potential issue with the neuroeconomical evaluation procedure is that the devaluation of reward value associated with task difficulty may be influenced by performance in the task, and not only by effort. This issue deserves a careful consideration in the present study, given that we found a correlation between fatigue-induced changes in task avoidance and performance. Previous attempts at solving this issue were based on persuading participants that their reward depended on task engagement, rather than on performance (Westbrook et al., 2013). Here, we relied on the design of the task-choice procedure to avoid any influence of performance on participants' decisions. Since block performance was computed by averaging accuracy in both the difficult and the easy tasks irrespective of their proportion in the block, a poor performance in the difficult task would have the same consequence on the reward, whatever choice the participants made. Therefore, there was no incentive for the participants to change their preference for the easy task ratio following fatigue. In fact, if participants task-choice depended on auto-evaluation of performance, decreased performance in N-back would have increased preference for low DL task conditions, whereas decreased performance in 1-

back would have increased preference for large DL tasks. However, we found that performance in both the 1-back and N-back tasks correlated with task avoidance. This indicates that task avoidance was affected by fatigue-induced performance decline rather than by the integration of performance as a cost in participants' decision. One question to address in future studies is whether the presence of the performance feedback at the end of each block is necessary to induce this relationship between performance decrements and task avoidance.

Another possible limitation of the present study is the reliance on the CSWm task to induce fatigue and N-back tasks to measure its effect. This choice was justified on the basis of several earlier studies of fatigue and effort discounting (Moeller et al., 2012; Pageaux et al., 2015; Wang et al., 2016; Westbrook et al., 2013). In addition, since we were interested in the link between the subjective feeling of fatigue - a task-independent subjective percept - and cognitive effort, there was no need to use the same task to induce and measure fatigue. Using different tasks has also the advantage of limiting the problem of boredom associated with the performance of the same monotonous task for several hours (Bench and Lench, 2013). Since little is known regarding the transferability of fatigue between different cognitive processes - available evidence point sometimes in the direction of global effect of fatigue, independent of the processes involved in fatigue induction (Arai, 1912; Wright et al., 2007), sometimes in the opposite direction (Persson et al., 2007) -, we decided to use a fatigue-inducing task involving a large range of cognitive processes, among which some should overlap with the ones involved in N-back performance. Along the same line, it could be argued that the N-back task was not demanding enough to lead to measurable effects on behavioral effort cost. We think it is unlikely because hit rate was only 80% on average in Experiment 2 and we found clear effect of task difficulty on reward discounting. Finally, the potential influence of the affective state of the participants (positive or negative) was not specifically evaluated following the completion of the fatigue / control manipulation (except for intrinsic motivation, which remained unaffected by fatigue induction). Affective state can indeed influence performance in cognitive tasks (Carver, C. S., & Scheier, 2011; Vohs et al., 2008) and should be considered as dependent variable in future studies.

It has been shown frequently that the feeling of fatigue occurs before any decrease of performance (Mullette-Gillman et al., 2015). Compensatory control hypothesis states that during the initial phase of cognitive fatigue, performance can be maintained at the cost of higher subjective effort (Hockey, 2013, 1997). However, the results of the present study failed to corroborate this hypothesis by showing a clear lack of relationship between subjective fatigue

1
2
3
4 and effort. We suggest that this lack of correlation of subjective fatigue with both performance
5 decrement and effort is compatible with an alternative “anticipatory regulation” view, according
6 to which cognitive fatigue would be an anticipatory protection mechanism, akin to anticipatory
7 regulatory mechanisms described in the physical fatigue literature (Crewe et al., 2008; Tucker,
8 2009; Tucker and Noakes, 2009). This protection mechanism would anticipate future adverse
9 consequences associated with task execution and would urge subjects to stop before any actual
10 consequence occurs (van der Linden, 2011). In other words, while the compensatory hypothesis
11 assumes that fatigue would affect cognitive resources early on and would lead to compensatory
12 increase in the recruitment of further resources in order to maintain performance (Hockey,
13 2011), the anticipatory regulation hypothesis assumes that subjective fatigue occurs before any
14 adverse consequence can be observed on the brain. According to this view, the engagement of
15 resources would remain constant during the initial buildup of fatigue, which would thus not lead
16 necessarily to increased subjective perception of effort or increased effort-discounting. Actual
17 alteration of cognitive resources would occur if one continues to engage in the task despite
18 subjective fatigue. At that point, performance would start to decrease and effortful control would
19 be engaged to compensate that decrease. This, in turn, might explain our findings that task
20 avoidance correlates with performance decrement.

21
22
23
24
25
26
27
28
29
30
31
32
33 This putative reliance of fatigue on such anticipatory mechanism points to several candidates in
34 terms of potential neural substrates. Anterior insula and cingulate cortex, in particular, have
35 been shown to encode anticipatory interoceptive signals (Barrett and Simmons, 2015).
36 Interestingly, these same regions are also thought to be involved in the phenomenology of
37 fatigue (Boksem and Tops, 2008; Dantzer et al., 2014; Pavese et al., 2010). Naturally, the
38 present findings provide only indirect support for this anticipatory regulation view and further
39 computational and neuroimaging studies are required to assess its validity in depth.

40 41 42 43 44 45 46 47 **Acknowledgements**

48
49
50 This work was supported by Fonds de la Recherche Scientifique (FNRS–FDP), Fondation
51 Médicale Reine Elisabeth (FMRE), the Fondation Louvain and IdEx Bordeaux. The authors
52 would like to acknowledge the contribution of Simon Van Hemelrijck for his support in data
53 acquisition.
54
55
56
57
58
59
60
61
62
63
64
65

References

- Ackerman, P.L., Kanfer, R., 2009. Test length and cognitive fatigue: An empirical examination of effects on performance and test-taker reactions. *J. Exp. Psychol. Appl.* 15, 163–181. doi:10.1037/a0015719
- Åkerstedt, T., Anund, A., Axelsson, J., Kecklund, G., 2014. Subjective sleepiness is a sensitive indicator of insufficient sleep and impaired waking function. *J. Sleep Res.* 23, 240–252. doi:10.1111/jsr.12158
- Åkerstedt, T., Gillberg, M., 1990. Subjective and Objective Sleepiness in the Active Individual. *Int. J. Neurosci.* 52, 29–37. doi:10.3109/00207459008994241
- Arai, T., 1912. Mental fatigue. Unpublished doctoral dissertation, Columbia University.
- Aston-Jones, G., Cohen, J.D., 2005. An integrative theory of locus coeruleus-norepinephrine function: adaptive gain and optimal performance. *Annu Rev Neurosci.*
- Badin, O.O., Smith, M.R., Conte, D., Coutts, A.J., 2016. Mental fatigue: Impairment of technical performance in small-sided soccer games. *Int. J. Sports Physiol. Perform.* 11, 1100–1105. doi:10.1123/ijsp.2015-0710
- Bailey, A., Channon, S., Beaumont, J.G., 2007. The relationship between subjective fatigue and cognitive fatigue in advanced multiple sclerosis. *Mult. Scler. J.* 13, 73–80. doi:10.1177/1352458506071162
- Barrett, L.F., Simmons, W.K., 2015. Interoceptive predictions in the brain. *Nat. Rev. Neurosci.* doi:10.1038/nrn3950
- Bench, S., Lench, H., 2013. On the Function of Boredom. *Behav. Sci. (Basel)*. 3, 459–472. doi:10.3390/bs3030459
- Boksem, M.A.S., Meijman, T.F., Lorist, M.M., 2006. Mental fatigue, motivation and action monitoring. *Biol. Psychol.* 72, 123–132.
- Boksem, M.A.S., Tops, M., 2008. Mental fatigue: Costs and benefits. *Brain Res. Rev.* doi:10.1016/j.brainresrev.2008.07.001
- Borragán, G., Slama, H., Bartolomei, M., Peigneux, P., 2017. Cognitive fatigue: A Time-based Resource-sharing account. *Cortex* 89, 71–84. doi:10.1016/j.cortex.2017.01.023
- Brainard, D.H., 1997. The Psychophysics Toolbox. *Spat. Vis.* 10, 433–436. doi:10.1163/156856897X00357

- Campagne, A., Pebayle, T., Muzet, A., 2004. Correlation between driving errors and vigilance level: Influence of the driver's age. *Physiol. Behav.* 80, 515–524.
doi:10.1016/j.physbeh.2003.10.004
- Carter, E.C., Kofler, L.M., Forster, D.E., McCullough, M.E., 2015. A series of meta-analytic tests of the depletion effect: Self-control does not seem to rely on a limited resource. *J. Exp. Psychol. Gen.* 144, 796–815. doi:10.1037/xge0000083
- Carver, C. S., & Scheier, M.F., 2011. Self-Regulation of Action and Affect, in: Baumeister, K.D.V. & R.F. (Ed.), *Handbook of Self Regulation: Research, Theory and Applications*. Guilford Press., New York, NY, USA, pp. 3–21.
- Charness, G., Gneezy, U., 2012. Strong Evidence for Gender Differences in Risk Taking. *J. Econ. Behav. Organ.* 83, 50–58. doi:10.1016/j.jebo.2011.06.007
- Christie, S.T., Schrater, P., 2015. Cognitive cost as dynamic allocation of energetic resources. *Front. Neurosci.* 9, 1–15. doi:10.3389/fnins.2015.00289
- Crewe, H., Tucker, R., Noakes, T.D., 2008. The rate of increase in rating of perceived exertion predicts the duration of exercise to fatigue at a fixed power output in different environmental conditions. *Eur. J. Appl. Physiol.* 103, 569–577. doi:10.1007/s00421-008-0741-7
- Dantzer, R., Heijnen, C.J., Kavelaars, A., Laye, S., Capuron, L., 2014. The neuroimmune basis of fatigue. *Trends Neurosci.* doi:10.1016/j.tins.2013.10.003
- Deluca, J., 2005. Fatigue, cognition and mental effort, in: *Fatigue as a Window to the Brain*. p. 17:52.
- Dorrian, J., Roach, G.D., Fletcher, A., Dawson, D., 2007. Simulated train driving: Fatigue, self-awareness and cognitive disengagement. *Appl. Ergon.* 38, 155–166.
doi:10.1016/j.apergo.2006.03.006
- Fairclough, S.H., Houston, K., 2004. A metabolic measure of mental effort. *Biol. Psychol.* 66, 177–190. doi:10.1016/j.biopsycho.2003.10.001
- Gailliot, M.T., Baumeister, R.F., 2007. The physiology of willpower: Linking blood glucose to self-control. *Personal. Soc. Psychol. Rev.* 11, 303–327. doi:10.1177/1088868307303030
- Gelman, A., Tuerlinckx, F., 2000. Type S error rates for classical and Bayesian single and multiple comparison procedures. *Comput. Stat.* 15, 373–390. doi:10.1007/s001800000040

- Gentile, S., Delarozière, J.C., Favre, F., Sambuc, R., San Marco, J.L., 2003. Validation of the French “multidimensional fatigue inventory” (MFI 20). *Eur. J. Cancer Care (Engl)*. 12, 58–64. doi:10.1046/j.1365-2354.2003.00295.x
- Gergelyfi, M., Jacob, B., Olivier, E., Zénon, A., 2015. Dissociation between mental fatigue and motivational state during prolonged mental activity. *Front. Behav. Neurosci*. 9. doi:10.3389/fnbeh.2015.00176
- Hart, S.G., Staveland, L.E., 1988. Development of NASA-TLX (Task Load Index): Results of Empirical and Theoretical Research. *Adv. Psychol*. 52, 139–183. doi:10.1016/S0166-4115(08)62386-9
- Hockey, G.R., 1997. Compensatory control in the regulation of human performance under stress and high workload; a cognitive-energetical framework. *Biol. Psychol*. 45, 73–93.
- Hockey, G.R.J., 2011. A Motivational Control Theory of Cognitive Fatigue. *Cogn. fatigue Multidiscip. Perspect. Curr. Res. Futur. Appl*. 167–187. doi:10.1037/12343-008
- Holtzer, R., Shuman, M., Mahoney, J.R., Lipton, R., Verghese, J., 2011. Cognitive Fatigue defined in the context of attention networks. *Aging, Neuropsychol. Cogn*. 18, 108–128. doi:10.1080/13825585.2010.517826
- Hosking, J.G., Cocker, P.J., Winstanley, C.A., 2014. Dissociable contributions of anterior cingulate cortex and basolateral amygdala on a rodent cost/benefit decision-making task of cognitive effort. *Neuropsychopharmacology* 39, 1558–67. doi:10.1038/npp.2014.27
- Inzlicht, M., Bartholow, B.D., Hirsh, J.B., 2015. Emotional foundations of cognitive control. *Trends Cogn. Sci*. doi:10.1016/j.tics.2015.01.004
- Joshi, S., Li, Y., Kalwani, R.M., Gold, J.I., 2016. Relationships between Pupil Diameter and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron* 89, 221–234. doi:10.1016/j.neuron.2015.11.028
- Kass, R.E., Raftery, A.E., 1995. Bayes Factor. *J. Am. Stat. Assoc*. doi:10.2307/2291091
- Kato, Y., Endo, H., Kizuka, T., 2009. Mental fatigue and impaired response processes: Event-related brain potentials in a Go/NoGo task. *Int. J. Psychophysiol*. 72, 204–211. doi:10.1016/j.ijpsycho.2008.12.008
- Kirchner, W.K., 1958. Age differences in short-term retention of rapidly changing information. *J. Exp. Psychol*. 55, 352–358. doi:10.1037/h0043688

- Kontsevich, L.L., Tyler, C.W., 1999. Bayesian adaptive estimation of psychometric slope and threshold. *Vision Res.* 39, 2729–2737. doi:10.1016/S0042-6989(98)00285-5
- Krueger, J.M., Rector, D.M., Roy, S., Van Dongen, H.P.A., Belenky, G., Panksepp, J., 2008. Sleep as a fundamental property of neuronal assemblies. *Nat. Rev. Neurosci.* 9, 910–919. doi:10.1038/nrn2521
- Kurzban, R., Duckworth, A., Kable, J.W., Myers, J., 2013. An opportunity cost model of subjective effort and task performance. *Behav. Brain Sci.* 36, 661–679. doi:10.1017/S0140525X12003196
- Lim, J., Wu, W. chau, Wang, J., Detre, J.A., Dinges, D.F., Rao, H., 2010. Imaging brain fatigue from sustained mental workload: An ASL perfusion study of the time-on-task effect. *Neuroimage* 49, 3426–3435. doi:10.1016/j.neuroimage.2009.11.020
- Lorist, M.M., 2008. Impact of top-down control during mental fatigue. *Brain Res.* 1232, 113–123. doi:10.1016/j.brainres.2008.07.053
- Lorist, M.M., Klein, M., Nieuwenhuis, S., De Jong, R., Mulder, G., Meijman, T.F., 2000. Mental fatigue and task control: planning and preparation. *Psychophysiology* 37, 614–625.
- MacLeod, C.M., 1991. Half a century of research on the Stroop effect: an integrative review. *Psychol. Bull.* 109, 163–203. doi:Doi 10.1037//0033-2909.109.2.163
- Massar, S.A.A., Wester, A.E., Volkerts, E.R., Kenemans, J.L., 2010. Manipulation specific effects of mental fatigue: Evidence from novelty processing and simulated driving. *Psychophysiology* 47, 1119–1126. doi:10.1111/j.1469-8986.2010.01028.x
- Moeller, S.J., Tomasi, D., Honorio, J., Volkow, N.D., Goldstein, R.Z., 2012. Dopaminergic involvement during mental fatigue in health and cocaine addiction. *Transl. Psychiatry* 2, e176.
- Mullette-Gillman, O.A., Leong, R.L.F., Kurnianingsih, Y.A., 2015. Cognitive Fatigue Destabilizes Economic Decision Making Preferences and Strategies. *PLoS One* 10, e0132022. doi:10.1371/journal.pone.0132022
- Muraven, M., Slessareva, E., 2003. Mechanisms of Self-Control Failure: Motivation and Limited Resources. *Personal. Soc. Psychol.*
- Pageaux, B., Marcora, S.M., Rozand, V., Lepers, R., 2015. Mental fatigue induced by prolonged self-regulation does not exacerbate central fatigue during subsequent whole-body endurance exercise. *Front. Hum. Neurosci.* 9. doi:10.3389/fnhum.2015.00067

- Paveze, N., Metta, V., Bose, S.K., Chaudhuri, K.R., Brooks, D.J., 2010. Fatigue in Parkinson ' s disease is linked to striatal and limbic serotonergic dysfunction 3434–3443. doi:10.1093/brain/awq268
- Persson, J., Welsh, K.M., Jonides, J., Reuter-Lorenz, P.A., 2007. Cognitive fatigue of executive processes: Interaction between interference resolution tasks. *Neuropsychologia* 45, 1571–1579. doi:10.1016/j.neuropsychologia.2006.12.007
- Reimer, J., Mccinley, M.J., Liu, Y., Rodenkirch, C., Wang, Q., Mccormick, D.A., Talias, A.S., 2016. Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. *Nat. Publ. Gr.* 7, 1–7. doi:10.1038/ncomms13289
- Ryan, R.M., Deci, E.L., 2000. Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *Am. Psychol.* 55, 68–78. doi:10.1037/0003-066X.55.1.68
- Sara, S.J., Bouret, S., 2012. Orienting and Reorienting: The Locus Coeruleus Mediates Cognition through Arousal. *Neuron* 76, 130–141. doi:10.1016/j.neuron.2012.09.011
- Schwid, S.R., Tyler, C.M., Scheid, E.A., Weinstein, A., Goodman, A.D., McDermott, M.P., 2003. Cognitive fatigue during a test requiring sustained attention: a pilot study. *Mult. Scler. J.* 9, 503–508. doi:10.1191/1352458503ms946oa
- Shenhav, A., Musslick, S., Lieder, F., Kool, W., Griffiths, T.L., Cohen, J.D., Botvinick, M.M., 2017. Toward a Rational and Mechanistic Account of Mental Effort. *Annu. Rev. Neurosci.* 40, 99–124. doi:10.1146/annurev-neuro-072116-031526
- Stewart, C.C., Wright, R.A., Azor Hui, S.K., Simmons, A., 2009. Outcome expectancy as a moderator of mental fatigue influence on cardiovascular response. *Psychophysiology* 46, 1141–1149. doi:10.1111/j.1469-8986.2009.00862.x
- Tanaka, M., 2015. Effects of Mental Fatigue on Brain Activity and Cognitive Performance: A Magnetoencephalography Study. *Anat. Physiol.* s4. doi:10.4172/2161-0940.S4-002
- Thiffault, P., Bergeron, J., 2003. Monotony of road environment and driver fatigue: A simulator study. *Accid. Anal. Prev.* 35, 381–391. doi:10.1016/S0001-4575(02)00014-3
- Tucker, R., 2009. The anticipatory regulation of performance: The physiological basis for pacing strategies and the development of a perception-based model for exercise performance. *Br. J. Sports Med.* doi:10.1136/bjsm.2008.050799
- Tucker, R., Noakes, T.D., 2009. The physiological regulation of pacing strategy during exercise:

- A critical review. *Br. J. Sports Med.* doi:10.1136/bjsm.2009.057562
- Vadillo, M.A., Gold, N., Osman, M., 2016. The Bitter Truth About Sugar and Willpower: The Limited Evidential Value of the Glucose Model of Ego Depletion. *Psychol. Sci.* 27, 1207–1214. doi:10.1177/0956797616654911
- Van Cutsem, J., De Pauw, K., Buyse, L., Marcora, S., Meeusen, R., Roelands, B., 2017. Effects of mental fatigue on endurance performance in the heat. *Med. Sci. Sports Exerc.* 49, 1677–1687. doi:10.1249/MSS.0000000000001263
- van der Linden, D., 2011. The urge to stop: The cognitive and biological nature of acute mental fatigue. *Cogn. Fatigue Multidiscip. Perspect. Curr. Res. Futur. Appl.* 149–164. doi:10.1037/12343-007
- van der Linden, D., Frese, M., Meijman, T.F., 2003. Mental fatigue and the control of cognitive processes: effects on perseveration and planning. *Acta Psychol. (Amst)*. 113, 45–65.
- Vohs, K.D., Baumeister, R.F., Schmeichel, B.J., Twenge, J.M., Nelson, N.M., Tice, D.M., 2008. Making Choices Impairs Subsequent Self-Control: A Limited-Resource Account of Decision Making, Self-Regulation, and Active Initiative. *J. Pers. Soc. Psychol.* 94, 883–898. doi:10.1037/0022-3514.94.5.883
- Vyazovskiy, V. V., Olcese, U., Hanlon, E.C., Nir, Y., Cirelli, C., Tononi, G., 2011. Local sleep in awake rats. *Nature* 472, 443–447. doi:10.1038/nature10009
- Wang, C., Trongnetrpunya, A., Samuel, I.B.H., Ding, M., Kluger, B.M., 2016. Compensatory Neural Activity in Response to Cognitive Fatigue. *J. Neurosci.* 36, 3919–24. doi:10.1523/JNEUROSCI.3652-15.2016
- Watson, A.B., Pelli, D.G., 1983. Quest: A Bayesian adaptive psychometric method. *Percept. Psychophys.* 33, 113–120. doi:10.3758/BF03202828
- Westbrook, A., Braver, T.S., 2015. Cognitive effort: A neuroeconomic approach. *Cogn. Affect. Behav. Neurosci.* 15, 395–415. doi:10.3758/s13415-015-0334-y
- Westbrook, A., Kester, D., Braver, T.S., 2013. What Is the Subjective Cost of Cognitive Effort? Load, Trait, and Aging Effects Revealed by Economic Preference. *PLoS One* 8, e68210. doi:10.1371/journal.pone.0068210
- Wright, R.A., Junious, T.R., Neal, C., Avello, A., Graham, C., Herrmann, L., Junious, S., Walton, N., 2007. Mental fatigue influence on effort-related cardiovascular response: Difficulty effects and extension across cognitive performance domains. *Motiv. Emot.* 31, 219–231.

doi:10.1007/s11031-007-9066-9

Zénon, A., Duclos, Y., Carron, R., Witjas, T., Baunez, C., Régis, J., Azulay, J.P., Brown, P., Eusebio, A., 2016. The human subthalamic nucleus encodes the subjective value of reward and the cost of effort during decision-making. *Brain* 139, 1830–1843. doi:10.1093/brain/aww075

Zénon, A., Sidibé, M., Olivier, E., 2014. Pupil size variations correlate with physical effort perception. *Front. Behav. Neurosci.* 8. doi:10.3389/fnbeh.2014.00286

Zering, J.C., Brown, D.M.Y., Graham, J.D., Bray, S.R., 2017. Cognitive control exertion leads to reductions in peak power output and as well as increased perceived exertion on a graded exercise test to exhaustion. *J. Sports Sci.* 35, 1799–1807. doi:10.1080/02640414.2016.1237777

Figure legends

Figure 1. Schematic description of the protocol. Participants were first evaluated with the multidimensional fatigue inventory (MFI) and Karolinska sleepiness scale (KSS). Following fatigue-inducing or control manipulation they again completed the questionnaires (only intrinsic motivation inventory (IMI) in Exp 1), and performed a forced-choice working memory task followed by the National Aeronautics and Space Administration Task Load Index (NASA-TLX). Experiment 2 also included arousal boosts consisting of 30-second run on the spot, prior to questionnaires completion.

Figure 2. A. Fitting curves from the behavioral model, averaged across participants for each difficulty level and session. Task preferences indicate a clear effect of difficulty level on probability of accepting more difficult task, with no clear difference between fatigue and control sessions in both experiments. B. Average performance in the 1-back (blue) and N-back tasks (red) in fatigue and control conditions. Error bars indicate the 95% confidence interval. C. Histogram of N-back difficulty levels used in both experiments.

Figure 3. Subjective evaluations and free model parameters in fatigue (gray bars) and control conditions (white bars) for Experiment 1 (A-C) and 2 (D-F). Error bars indicate the 95 % confidence interval. * $p < .05$. ** $p < .01$. *** $p < .001$.

Figure 4. Relationships between session changes in N-back performance, subjective evaluations and free model parameters for Experiment 1 (A-D) and Experiment 2 (E-H). * $p < .05$, ** $p < .01$, *** $p < .001$.

Figure 1
[Click here to download high resolution image](#)

Figure 2
[Click here to download high resolution image](#)

Exp. 1

Exp. 2

Figure 3
[Click here to download high resolution image](#)

Exp. 1

Exp. 2

Figure 4
[Click here to download high resolution image](#)

Exp. 1

Exp. 2