Asymmetric iodine catalysis-mediated enantioselective oxidative transformations

Aurélie Claraz, Géraldine Masson

► To cite this version:

Aurélie Claraz, Géraldine Masson. Asymmetric iodine catalysis-mediated enantioselective oxidative transformations. *Organic & Biomolecular Chemistry*, 2018, 16 (30), pp.5386-5402. 10.1039/C8OB01378K . hal-02410604

HAL Id: hal-02410604

<https://hal.science/hal-02410604>

Submitted on 13 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Asymmetric iodine catalysis-mediated enantioselective oxidative transformations

Aurélie Claraz* and Géraldine Masson*

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

The implementation of chiral iodine catalysis has tremendously been developed in the field of asymmetric synthesis during the last decade. It enables the stereoselective creation of C–O as well as C–C, C–N and C–X (X = halogen) bonds through oxidative transformations. Thanks to the low toxicity and ease of handling of iodine compounds, this strategy offers many advantages over classical metal-catalyzed oxidations with chiral ligands. The approaches rely on iodine (I/III) or (–I/+I) catalysis by using a chiral aryl iodine or ammonium iodide respectively in combination with a suitable terminal oxidant. As such, the design of iodine compounds with central, axial or even planar chirality has allowed to achieve high enantioselectivities. The goal of this review is to cover the different chiral iodine compounds-catalyzed oxidative transformations including α -functionalization of carbonyl compounds, dearomatization of phenol derivatives and difunctionalization of alkenes which should demonstrate that iodine catalysis has now found its place in the realm of asymmetric organocatalysis.

1. Introduction

Iodine catalysis has emerged as a versatile and eco-friendly strategy to perform oxidative transformations allowing for instance couplings with formation of a C–O, C–N, C–C or C–X (X = halogen) bond.¹ In order to describe mechanisms of such reactions, synthetic chemists frequently borrow the terms of oxidative additions, ligand exchanges and reductive eliminations to the field of metal-mediated processes. Nevertheless, iodine compounds are mild, stable, easy to handle and eco-compatible reagents, advantageously competing with heavy metals. As such, it is not surprisingly that iodine catalysis has attracted the interest of the synthetic community in the field of asymmetric synthesis with the aim to develop a greener and more environmentally benign chemistry.² Up to now, asymmetric iodine catalysis-mediated oxidative transformations rely on the catalytic use of either hypervalent organoiodine (III) reagents (also called λ^3 -iodanes) or organic hypoiodites (I) as chiral oxidants. In the first case, chiral organoiodanes (III)³ are generated *in situ* from aryl iodine precatalysts and a stoichiometric amount of achiral co-oxidant (Figure 1). The first enantioselective catalytic transformation was reported in 2007 by Wirth *et al.* for the α -oxytosylation of ketones by using an aryl iodine species bearing a chiral moiety *ortho* to the iodine atom in the aromatic ring.⁴ Since this original report, numerous chiral aryl iodide-mediated enantioselective oxidative transformations through iodine (I/III) catalysis have been investigated.⁵ Besides cost reductions and

atom economy, the development of catalytic processes allows to avoid the preliminary preparation and isolation of the chiral hypervalent organoiodine (III) reagent, since this reactive species is generated *in situ* from an aryl iodide. Obviously, this feature can be advantageous during the design of new chiral precatalysts, especially for unstable hypervalent organoiodine (III) derivatives. The currently known enantiopure precatalysts of hypervalent organoiodine (III) compounds exhibit either C₁- or C₂-symmetry with central, axial or even planar chirality (Figure 1).

Institut de Chimie des Substances Naturelles, CNRS UPR 2301
Université Paris-Sud, Université Paris-Saclay
1 av. de la Terrasse, 91198 Gif-sur-Yvette (France)
e-mail: aurelie.claraz@cnrs.fr; geraldine.masson@cnrs.fr
homepage: <http://www.icsn.cnrs-gif.fr/spip.php?article848&lang=en>

Figure 1 Concept of iodine (I/III)-enabled enantioselective oxidative transformations and general structure of chiral ayliodine precatalysts.

On the other hand, chiral hypiodites (I) are generated *in situ* from chiral organic iodides (-I) (typically chiral ammonium

iodides that are traditionally used in asymmetric phase transfer catalysis),⁶ and a stoichiometric amount of achiral co-oxidant (Figure 2).⁷

Figure 2 Concept of iodine (-I/I)-enabled enantioselective oxidative transformations and general structure of chiral ammonium iodide precatalysts.

In both catalytic cycles depicted above, (re)oxidation of the iodine atom of a precatalyst in low valent state to higher oxidation state is the key step.⁸ The choice of this stoichiometric achiral oxidant is critical for the success of the reaction. In an ideal fashion, it should not interact with the starting material or, at least, reoxidation of the precatalyst should be faster than the rates of undesired side reactions. In asymmetric processes, peroxides are the most commonly used oxidants that satisfy with more or less success these requirements.

To get cleaner reactions along with highest levels of enantiocontrol, a careful optimization of other classical reaction parameters such as solvent, temperature and the use of additives must be undertaken.

The focus of this review is to summarize the use of chiral iodine compounds in catalytic enantioselective oxidative processes with the aim of highlighting both catalysts design and synthetic applications. As such, we hope that this survey will help the reader in designing new chiral iodine precatalysts and applying those to new catalytic processes. For the sake of clarity, this review has been divided into different sections according to the kind of oxidative transformation considered:

- α -functionalization of carbonyl compounds,
- dearomatization of phenol derivatives,
- functionalization of alkenes.

2. α -Functionalization of carbonyl compounds

Hypervalent iodine compounds enable the introduction of a nucleophile at the α -position of a carbonyl compound. As such, numerous efforts have been made towards the development of enantioselective versions to take advantage of this inversion of polarity (*umpolung* reaction).

2.1. α -Oxytosylation

The enantioselective α -oxytosylation of ketones mediated by chiral hypervalent iodines gives access to synthetically useful enantioenriched carbonyl compounds. As mentioned above, the first catalytic version of this reaction was reported in 2007 by Wirth *et al.* Inspired by the works of Togo *et al.* who demonstrated a catalytic racemic version of this reaction,⁹ Wirth *et al.* employed organiodane **I** with central chirality as the chiral inducer, *meta*-chloroperoxybenzoic acid (mCPBA) as the stoichiometric oxidant and *p*-toluenesulfonic acid monohydrate (TsOH·H₂O) as a source of tosylate nucleophiles (Scheme 1).⁴ Contrarily to the process under stoichiometric conditions,¹⁰ the catalytic transformation needs to be carried out at room temperature, which suggests that oxidation of the iodoarene is the rate determining step in the catalytic cycle. Noteworthy, no racemization of the product in the reaction medium was observed. Nevertheless, only moderate enantioselectivities were obtained in this difficult transformation. From a mechanistic point of view, the authors proposed the *in situ* generation of a chiral Koser's-type reagent from **I**, mCPBA and TsOH·H₂O, furnishing either intermediate **3** or **4**. Subsequent nucleophilic addition of a tosylate anion on **3** (path A) or **4** (path B) via S_N2' and S_N2 type mechanisms would both provide enantioenriched ketone **2** along with the regeneration of aryl iodine **I**. Interestingly, the reaction is generally clean and the α -oxybenzoyl ketones which would result from the nucleophilic addition of the mCBA by-product were not formed.

Importantly, the authors pointed out the need of a chelating group at the benzylic position on the *ortho* substituent of the iodine atom in **I** to get the highest possible levels of enantioselectivity. Based on this observation, they synthesized compound **II**, having a chiral ester moiety as the chelating function at the benzylic position instead of the methoxy group, and resulting in an increase of enantioselectivity up to 39% during the α -oxytosylation of propiophenone (Scheme 2).¹¹ In 2010, the same group prepared derivative **III**, bearing a chiral ester at the *ortho* position of the iodine, and leading to lower enantioselectivity.¹²

In 2012, the group of Legault *et al.* developed chiral aryl iodine **IVa** bearing a chiral oxazoline as a coordinating motif and a methyl at the *ortho* and *ortho'*-positions of the iodine.¹³ Premixing catalyst, substrate, and TsOH, followed by slow addition of mCPBA slightly enhanced both yield and selectivity during the α -oxytosylation of propiophenone. According to computational analyses, this procedure would favor the formation of the key reactive iodane **5** by minimizing *N*-oxidation of the oxazoline moiety through *N*-protonation.¹⁴ Noteworthy, corresponding aryl iodine **IVb** without the *ortho'*-

methyl substituent is not able to promote the reaction as protonation of **IVb** is much more difficult. The best selectivity was finally obtained after the introduction of a chlorine atom *para* to the iodine (**IVc**).

Scheme 1 First catalytic enantioselective α -oxytosylation of ketones.

difficult.²⁰ The lack of control in the geometry of enol **3** might be also problematic. To avoid the passage through the *O*-bonded intermediate **3** and favor the *C*-bonded intermediate **4**, Legault *et al.* considered α -oxtosylation of enolacetate **6** (Scheme 5). With such a strategy, C_2 -symmetric 2-iodoresorcinol derivative **IX** offered the α -oxtosylated ketone **2** in good yield and with high enantioselectivity.²¹ It is worth pointing out that very poor enantioselectivity was obtained during the direct α -oxtosylation of ketone **2** with the same kind of catalyst. This observation supports two different mechanisms between the α -oxtosylation of ketone and enol acetate: S_N2' -type reductive elimination of a *O*-enolate intermediate in the former case providing low enantiocontrol in the former case and highly enantioselective S_N2 substitution of a *C*-enolate in the latter case.²⁰

Scheme 5 α -oxtosylation of enol acetates.

In 2016, Xiong, Coeffard *et al.* investigated the α -oxtosylation of α -substituted β -ketoesters **7** by using C_2 -symmetric iodoarene **IX** as chiral precatalyst in combination with mCPBA as stoichiometric oxidant at -20°C in chloroform (Scheme 6).²² Corresponding products **8** with a quaternary stereogenic center were obtained in poor yields albeit respectable enantiomeric excesses (up to 37% yield and 62% ee). Interestingly, 3 equivalents of trifluoroacetic acid (TFA) as additive were able to increase enantioselectivity.²³

Scheme 6 α -oxtosylation of α -substituted β -ketoesters.

2.2. Intramolecular α -oxygenation

The hypervalent iodine catalyzed α -oxygenation of carbonyl compounds has been extended by Moran *et al.* to an intramolecular version *via* lactonization of 5-oxo-5-phenylpentanoic acid **9** (Scheme 7).²⁴ Inspired by the works of Legault *et al.*,¹³ the authors synthesized ortho-tolyliodine **XI** bearing a pseudoephedrine-based chiral amide as chelating

Lewis agent. In the presence of substoichiometric amount of mCPBA and *p*TsOH, 5-benzoyldihydrofuran-2(3*H*)-one **10** was obtained with moderate yield and enantioselectivity.

Scheme 7 Oxidative cyclization of 5-oxo-5-phenylpentanoic acid.

In 2010, an elegant catalytic enantioselective oxidative cyclization of ketophenols **11** to the five-membered ring products 2-acyl-2,3-dihydrobenzofurans **12** has been developed by Ishihara *et al.* (Scheme 8a). Excellent yields and enantioselectivities at very low catalyst loading were obtained. Their approach relied on the use of a *in situ* generated chiral hypoiodite $[\text{IO}]^-$ catalyst from the corresponding quaternary ammonium iodide precatalyst **XII** with hydrogen peroxide or *tert*-butyl hydroperoxide (TBHP) as environmentally benign co-oxidants.²⁵ The unstable catalytic active species hypoiodite has been detected by a series of Raman spectra unveiling an unprecedented $\text{I}(-\text{I}/+\text{I})$ asymmetric catalysis.²⁶ This method has been extended to the less favorable six-membered ring oxidative cyclization to access 2-acyl-chromane derivatives.²⁶ In a similar fashion, highly enantioenriched 2-acyl tetrahydrofuran derivatives **14** have been obtained *via* cycloetherification of corresponding δ -hydroxyketones **13** using cumene hydroperoxide (CHP) as stoichiometric co-oxidant (Scheme 8b).²⁷

Scheme 8 *In situ* generated chiral hypoiodite-catalyzed oxidative cyclization.

2.3. α -Fluorination

A chiral iodoarene-catalyzed asymmetric nucleophilic α -fluorination of indanone-based β -ketoesters **15** has been initially reported by Kita, Shibata *et al.* using C_2 -symmetric axially chiral precatalyst **VI** in combination with mCPBA as a stoichiometric oxidant and n HF-pyridine as fluorine source (Scheme 9).²⁸ Moderate enantioselectivity was obtained with methyl ester **15a**, whereas sterically demanding adamantyl ester **15b** afforded a higher enantiomeric excess (56% *ee* vs. 25% *ee*). A breakthrough has been very recently accomplished by Rueping *et al.* by using chiral iodoarenes derived from 2-iodoresorcinol and lactate scaffolds with triethylamine pentahydrofluoride as the fluorine source.²⁹ More particularly, the authors developed the flexible C_2 -symmetrical menthol ester derivative **XIII** bearing a 3,5-bis(trifluoromethyl)-phenyl group *para* to the iodine providing fluorinated product **16** in good yields and with high enantioselectivities. Chiral iodoresorcinol derivative lacking C_2 -symmetry proved to be inferior. In parallel to Rueping's work, Lu, Zheng *et al.* have designed a novel planar chiral iodoarene **XIV** based on [2,2]-paracyclophane and bearing a benzoyl group as potentially coordinating function of the *ortho*-iodine atom. Triethylamine trihydrofluoride (TEA·3HF) was selected as fluorine source. It is worth mentioning that a positive effect on the enantioselectivity was observed with the introduction of methoxy-groups on the phenyl ring of the benzoyl.³⁰ In line with experimental observations, these three research groups assumed *in situ* generation of Ar^*-I-F_2 from Ar^*-I , mCPBA and HF along with activation of this key chiral reactive species by an excess of HF. Based on DFT calculations, Rueping *et al.* proposed a mechanism with formation of an *O*-bonded hypervalent iodine intermediate **17**.²⁹ Migration of the aryl iodonium from the *O*-ester enolate to the α -C atom would lead to intermediate **18** *via* transition state **17**→**18** and yield the α -fluorinated product through reductive elimination. The competitive α -hydroxylation of the carbonyl compounds is responsible of the moderate yield. Transition state **17**→**18** is anticipated to be the enantioselectivity-determining step. According to it, formation of (*R*)-**18** is favored as transition state (pro-*R*) **17**→**18** was found 1.4 kcal/mol lower in energy than transition state (pro-*S*) **17**→**18**. Reductive elimination would afford α -fluorinated (*R*)-product **16**, which is in agreement with experimentally observed stereochemistry. Unfortunately, the authors did not provide any calculations concerning the energy of a transition state corresponding to a pseudo-intramolecular S_N2' type reductive elimination of an *O*-iodonium keto-enolate, as it was proposed for the α -oxytosylation of carbonyl compounds by Legault *et al.* (scheme 1, intermediate **3**, L = F).²⁰

Scheme 9 α -fluorination of indanone derivatives.

2.4. Spirocyclization through $C(sp^2)$ - $C(sp^3)$ coupling

Although most hypervalent iodine (III) catalyzed-asymmetric oxidative coupling protocols enable the formation of C-X bonds (X = O, N, F), an elegant intramolecular $C(sp^2)$ - $C(sp^3)$ coupling reaction of N^1, N^3 -diphenylmalonamide **19** has been achieved by Gong *et al.* in 2014 (Scheme 10).³¹ This transformation provided access to biologically important spirobisoxindoles **20** with a quaternary carbon stereogenic center in good yields and with good enantiomeric excesses. The best reaction conditions involved the use of ethaneperoxoic acid ($MeCO_3H$) as the stoichiometric oxidant in nitromethane with C_2 -symmetric iodoresorcinol derivative **XV** as precatalyst. Notably, the tertiary proline amide on **XV** (at the end of the resorcinol arms) played key role in inducing the highest enantioselectivities compared to secondary amides, carboxylic acids or esters.

Moreover, the addition of 2 equivalents of trifluoroacetic acid enhanced the catalytic performances in terms of both yield and enantioselectivity. To shed light on the reaction mechanism and the origin of the stereoselectivities, computational studies have been undertaken by Sunoj *et al.*³² Based on DFT calculations, the authors proposed a mechanism during which the *in situ* generation of the key chiral hypervalent iodine intermediate $\text{Ar}^*\text{-I}(\text{TFA})_2$ was the initial step. Ligand exchange with the nitrogen atom of incoming dianilide substrate **19** followed by pseudo-intramolecular deprotonation of **21** by the leaving trifluoroacetate anion generated *O*-iodonium enolate **22**. The ensuing *O* to *C*-aryliodonium migration leads to *C*-iodonium enolate **23** with a lower energy. Intramolecular Friedel-Craft reaction assisted by the trifluoroacetate anion allowed the formation of the first oxindole ring **24** with concomitant regeneration of precatalyst **XV**. The second ring formation would be enabled by a similar reaction sequence yielding spirobisoxindole **19**. It is worth mentioning that direct ring closings from *O*-iodonium **22** and **26** have been calculated to be of higher energy, indicating that formation of *C*-iodonium enolates **23** and **27** might occur. As chiral information of intermediate **24** is lost during the second catalytic cycle, stereoselectivity of the reaction is solely governed during the second ring closure. More precisely, the 1,3-migration of the chiral aryl iodonium from *O* to *C*-enolate (**26**→**27**) is the stereocontrolling step of the reaction. The transition states leading to (*S*)-product **19** were found to be 2.1 kcal/mol lower than those leading to the (*R*)-product, which is in good agreement with the experimentally obtained enantioselectivity. Noteworthy, in the preferred conformer of key chiral hypervalent iodine intermediate $\text{Ar}^*\text{-I}(\text{TFA})_2$, resorcinol arms were found to organize in a C_2 -symmetric helical assembly with a right-handed P helicity in which the two trifluoroacetate ligands form a network of noncovalent interactions with the chiral arms.

Scheme 10 Spirocyclisation through two sequential C-C cyclizing couplings.

3. Dearomatization of phenol derivatives

The asymmetric dearomatization of phenol derivatives is of particular interest for the synthetic community as it can be a pivotal step in the total synthesis of natural products. Chiral hypervalent organoiodines reagents are organo-oxidants of choice to perform this transformation enantioselectively.³³ They can be generated *in situ* from a catalytic amount of chiral iodoarene and mCPBA as a stoichiometric oxidant. Ligand exchange between the phenol and *in situ* generated iodine (III) would furnish phenoxy- λ^3 -chiral aryliodarene **28** (Figure 3). Subsequently, two mechanisms are generally considered: direct attack of the nucleophile via a $\text{S}_{\text{N}}2'$ -type substitution (associative path) or formation of phenoxenium ion **29**, which is trapped by the nucleophile (dissociative path). The key feature to successively achieve asymmetric induction is to find reaction conditions (catalyst, solvent, additive...) capable of favorizing the associative mechanism.

Figure 3 Proposed mechanisms for organoiodane-mediated phenol dearomatization.

3.1. Intramolecular transformations

The intramolecular dearomatization of phenol derivatives provides a straightforward way to spirocyclic compounds. The asymmetric dearomatization of propanoic acid substituted 1-naphthol **30** to spiroactone **31** has been intensively studied as a model reaction for the design of new aryl iodine precatalysts. Indeed, precatalysts which exhibit either C_1 - or C_2 -symmetry and central or axial-chirality have been developed to achieve the highest enantioselectivity (Scheme 11).

In 2017, Nachtsheim and Pericàs *et al.* reported a new C_1 -symmetric triazole-based chiral iodoarenes (**XVI**). This catalyst was prepared thanks to an enzymatic kinetic resolution followed by a copper-catalyzed azide-alkyne cycloaddition and provided spiroactone **31a** with moderate yield and good enantioselectivity.

Ishihara *et al.* developed conformationally flexible C_2 -symmetric chiral iodoarenes **XVII** derived from lactate as a source of chirality affording spiroactone **31a** in good yield and with high enantioselectivity (55%, 92% ee).³⁴ More recently, an enhancement of the yield and the enantioselectivity was achieved in the same group by using catalyst **XVIII** derived from chiral amino-alcohol at low loadings.³⁵ X-ray and NMR analyses of *in situ* generated organoiodine(III) species from **XVIII** showed that intramolecular hydrogen-bonding interactions between acidic amido protons and ligands of iodine(III) allowed the construction of a suitable chiral environment around the iodine(III) center.³⁶

An original C_2 -symmetric iodoarene **XIX** with central chirality based on the rigid all-carbon anti-dimethanoanthracene framework displayed lower activity and selectivity.³⁷ However, a rigid C_2 -symmetric aryl iodine **Vb** based on the axially chiral spirobiindane scaffold has been disclosed by Kita *et al.* in 2008. Cyclized product **31a** has been obtained in good enantioselectivity with this catalyst.³⁸ An enhancement of the enantioselectivity was observed with **Vc** via introduction of ethyl-substituents at the *ortho*-positions of the iodine atoms, presumably thanks to an extension of surroundings around the reactive sites.³⁹

In 2016, Masson *et al.* explored axially chiral 1,1'-binaphthalene framework and derivatives for the development of novel chiral iodoarenes prepared from commercially

available (*R*)-1,1'-binaphthyl-2,2'-diamine (BINAM).⁴⁰ Spiroactone **31a** was obtained with little enantiomeric excess with H₈-BINI **XXa**. Introduction of 1-naphthyl substituents at the 3,3'-positions (**XXb**) exerted a slightly beneficial effect on enantioselectivity. Interestingly, a non C_2 -symmetric iodoarene precatalyst (**XXI**) harboring an amide group at the 2'-position afforded spiroactone **31a** in poor enantioselectivity but with an inversion of configuration. By analogy with Ishihara and Wirth's catalysts, additional non-covalent interactions between the amide group and the iodine (III) center or its ligands may be responsible of this reversal through a different transition state. Kita *et al.* designed a new type of chiral 1,1'-binaphthalene (**XXII**) substituted with the iodine atoms at the 8,8'-positions which dramatically improved enantioselectivity, but at the price of a high catalyst loading (0.55 equiv.).⁴¹ This precatalyst bearing the iodine atoms within the major groove of the naphthalene rings has been synthesized by a nickel-catalyzed reductive homocoupling, enantiomeric resolution of the obtained racemic diamine on preparative HPLC with chiral stationary phase column, followed by Sandmeyer-type iodination.

Scheme 11 Asymmetric dearomatization of propanoic acid substituted 1-naphthol **30a**.

In 2015, Ishihara *et al.* reported application of their strategy of *in situ* generated-chiral ammonium hypoiodite from the corresponding chiral ammonium iodide and hydrogen peroxide as stoichiometric oxidant to the catalytic spirocyclization of 1-naphthol derivatives with good yields and good enantioselectivities at room temperature (Scheme 12).⁴² The use of hydrogen-peroxide as stoichiometric oxidant offers the advantage of generating only water as a by-product instead of mCPBA, but limited the scope of the transformation to 3 or 4-substituted naphthols.

Scheme 12 Hypoiodite-catalyzed asymmetric dearomatization of 3 or 4 substituted 1-naphthol derivatives **30**.

In 2013, Ishihara *et al.* investigated oxidative cyclizative dearomatization of phenol derivatives **32** by using precatalyst **XVIII** (Scheme 13).³⁶ In the case of poorly reactive electron-deficient phenols, the addition of 50 equivalent of 1,1,1,3,3,3-hexafluoroisopropanol (HFIP) was required and dramatically enhanced efficiency of the process.⁴³ In contrast, addition of 25 equivalents of methanol was necessary in the case of highly reactive electron-rich phenols to presumably ensure a ligand exchange from acyloxyphenoxyiodine (III) **34**. This protocol would slow down the racemic dissociative pathway and would favor the enantioselective associative pathway as methoxy ligand is a less good leaving group than carboxylate ligand. Under these conditions, high yields and excellent enantioselectivities were obtained in both cases. Noteworthily, this protocol has proved to be also very efficient for the spirocyclization of poorly reactive 2-naphthol derivatives³⁵ and for dearomatization of *ortho*- and *para*-hydroquinone derivatives allowing the construction of *ortho*- and *para*-benzoquinones.⁴⁴

Proposed explanation for the role of MeOH if R = EDG:

Scheme 13 Spirocyclization of phenols.

In 2017, Ciufolini *et al.* described the hypervalent-iodine catalyzed enantioselective oxidative cycloetherification reaction of naphtholic alcohols **35** (Scheme 14).⁴⁵ The design of a new precatalyst was required as these substrates are less reactive than the corresponding carboxylic acids. In this context, a variant of the Ishihara's catalyst was examined. Considering that intramolecular H-bonding between the secondary amide NH group and the ligand of the iodine III is the key feature in these type of catalysts, the authors relocated the stereogenic center closer to the iodine atom. As a result, **XXIII** provided spiro compounds **36** in high yields and with excellent enantioselectivities. This procedure has been extended to oxidative cyclization of naphtholic sulfonamides with good enantioselectivity.

Scheme 14 A modification of Ishihara's type catalysts for the spirocyclization of naphtholic alcohols.

In their continuous efforts to exploit chiral hypervalent iodine catalysis for the creation of C-C bonds, Gong *et al.* have disclosed a highly enantioselective dearomatizative spirocyclization of 1-hydroxy-*N*-aryl-2-naphthamide derivatives **37**, providing access to spiroindoles derivatives **38** with good yields and enantioselectivity.⁴⁶ Precatalyst **XVII** was the aryl iodine of choice for this transformation in the presence of mCPBA as a stoichiometric oxidant. The reaction would proceed through the formation of phenoxy- λ^3 -iodine species **39**, which would undergo intramolecular S_N2' -like Friedel-Crafts substitution. The addition of a mixture of trifluoroethanol and water was critical for the improvement of both yields and enantioselectivity (Scheme 15).⁴³ As previously suggested by Ishihara *et al.*, these additives would facilitate ligand exchanges to favor the enantioselective associative pathway over the racemic dissociative pathway.

Scheme 15 Oxidative C-C coupling-enabled naphthols dearomatization.

3.2. Intermolecular transformations

Asymmetric intermolecular phenol dearomatization is a highly challenging task as this transformation can easily proceed through a racemic dissociative pathway.⁴⁷ Quideau *et al.* have pioneered catalytic enantioselective hydroxylative phenol dearomatization.⁴⁸ Axially chiral monoiodobinaphthyl **XXIV** was used in the presence of a stoichiometric amount of mCPBA to convert 2-methylnaphthol **40** into epoxy *ortho*-quinol **41** in good yield and with moderate enantioselectivity. Unfortunately, epoxidation of the *ortho*-quinol by the stoichiometric oxidant cannot be avoided. Secondary I...O interactions between the iodine (III) atom and the methoxy group and/or the carboxylic acid function were assumed, but the real nature (λ^3 or λ^5) of reacting hypervalent iodine involved in this transformation could not be determined with certainty.

Scheme 16 Intermolecular naphthol dearomatization.

Hypervalent iodine-catalyzed intermolecular dearomatization of phenol can also enable introduction of a nucleophile at the *para* position of the hydroxyl group. Designing an enantioselective version of such a reaction seems to be even more arduous, as the creation of the stereogenic center occurs quite far away from the chiral environment (namely the phenoxy- λ^3 -chiral aryl iodane). To address this challenge, Harned *et al.* designed C_1 -symmetric chiral iodine **XXV** derived from 8-iodotetralone and tartaric acid for the 4-

hydroxylation of phenols **42** in the presence of mCPBA as the stoichiometric oxidant (Scheme 17).⁴⁹ The resulting 2,5-cyclohexanedienediones **43** were obtained in moderate yields and with enantioselectivity up to 60% *ee*. In 2017, Muñiz *et al.* proposed the use of C₂-symmetric lactic amide **XXVI** (Ishihara's type precatalyst), which exhibited slightly lower stereoselectivities for the same transformation.⁵⁰ Very recently, Maruoka *et al.* have developed C₁-symmetric indanol-based chiral organoiodine **XXVII** allowing the access to 2,5-cyclohexadienediones **43** with the highest level of enantioselectivity reported to date.⁵¹

Scheme 17 4-hydroxylation of phenols.

isochromanones **45** were obtained in moderate yields and enantioselectivities up to 91% *ee*. The reaction would proceed through the diastereofacial attack of the chiral iodane on the double bond to form iodonium **46**. The latter would undergo nucleophilic substitutions by the internal hydroxy and carboxymethyl groups with inversion of configuration. The competitive direct oxidation of the double bond by mCPBA is responsible of the moderate yields.

Scheme 18 Oxidative double cyclization.

The background oxidation with mCPBA has been well identified by the same research group in the hypervalent-iodine catalyzed trifluoroacetoxylation of *ortho*-alk-1-enylbenzoates **47** (Scheme 19).⁵⁴ In this process, the catalytic oxidation mediated by chiral hypervalent iodine(III) species gives enantioenriched *syn*-products **48a** and **48b** via ring opening of iodonium **49**, while the direct oxidation with mCPBA delivers racemic *anti*-products **48c** and **48d** via opening of epoxide **50**. Good 6-*endo* regioselectivity and high enantioselectivity were obtained on *cis*-isochromanone **48a** with precatalyst **XXIX**. Unfortunately, the competitive background oxidation with mCPBA of the electron-rich internal double bond significantly affected the yield.

4. Functionalization of alkenes

Hypervalent iodine-mediated asymmetric difunctionalization of alkenes is a useful strategy for the preparation of polysubstituted molecules with the creation of one or two vicinal stereogenic centers from either two identical or different nucleophiles.⁵²

4.1. Intramolecular version

In its intramolecular version, such a reaction enables the construction of polysubstituted heterocycles in one step.

4.1.1. Lactonization. The first example of a catalytic difunctionalization of alkenes mediated by chiral hypervalent iodine has been reported by Fujita *et al.* In their original reports, the authors disclosed an oxidative double cyclization of 2-(4-hydroxybut-1-enyl)benzoates **44** (Scheme 18).⁵³ By using precatalyst **XXVIII**, mCPBA as the terminal oxidant and trifluoroacetic acid as an activator, the dihydrofuran-fused

Scheme 19 Oxy-lactonization of *ortho*-alk-1-enylbenzoates **47**.

In 2017, Masson *et al.* reported a tosyloxy- and phosphoryloxy-lactonization of flexible 4-pentenoic acid derivatives **51** (Scheme 20).⁵⁵ By using **XVII** and a stoichiometric amount of mCPBA in the presence of sulfonic or phosphoric acid, excellent *exo* selectivity was observed, providing access to valuable γ -butyrolactones **52** with good yields and good enantioselectivities. To minimize competitive direct oxidation of the terminal double bond by mCPBA, the reactions need to be carried out at 0 °C. Addition of 2,2,2-trifluoroethanol (TFE) as a cosolvent was required to improve the yield of the phosphoryloxycyclization.⁴³ From a mechanistic point of view, the authors proposed *in situ* generation of intermediate **53** from precatalyst **XVII**, mCPBA and the acid. Electrophilic attack of the double bond by this chiral λ^3 -aryliodonane would furnish chiral arylodonium **54**. Intramolecular nucleophilic attack of the carboxy group with *exo* selectivity followed by substitution of the iodonium with the sulfonyl or phosphoryl group would afford γ -lactone **52**.

Scheme 20 Tosyloxy- and phosphoryloxy-lactonization of flexible 4-pentenoic acid **51**.

The hypervalent iodine catalyzed fluorolactonization of *ortho*-alk-1-enylbenzoates **55** has been investigated by Jacobsen *et al.* (Scheme 21).⁵⁶ This method used HF·pyridine as a nucleophilic fluoride source with chiral precatalyst **XXX** and mCPBA, affording 3-alkyl-4-fluoroisochromanones **56** as a single *syn* diastereoisomer with moderate yields and high enantioselectivities. Noteworthy, the obtained regioselectivity (formation of the 6-membered ring) is complementary to that established in asymmetric fluorolactonization with an electrophilic fluoride source. This regioselectivity is explained by activation of the alkene by hypervalent iodine followed by addition of the fluoride anion onto the benzylic carbon, furnishing *anti*-vicinal fluoroiodonium intermediate **57**. The latter would undergo intramolecular nucleophilic substitution by the carboxymethyl group with inversion of configuration providing *syn*-diastereoisomer **56**.

Scheme 21 Hypervalent iodine catalysed asymmetric 6-endo-type-fluorolactonization.

4.1.2. Formation of aza-heterocycles. Kita, Shibata *et al.* have developed the chiral aryl iodine-catalyzed intramolecular aminofluorination of ω -amino-alkene **58** using mCPBA as the stoichiometric terminal oxidant and HF-pyridine as a nucleophilic fluoride source.²⁸ This transformation enabled the formation of 2-fluoropiperidine **59** with moderate yield and enantioselectivity. According to the observed 6-endo selectivity of the process, the aminofluorination would proceed through oxidation of the sulfonamide by *in situ* generated ArIF_2 , furnishing electrophilic species **60**. The latter would react with the double bond to release the precatalyst and concomitantly form intermediary aziridinium **61**. The latter would undergo nucleophilic attack of the fluoride on the tertiary carbon to deliver *endo*-type cyclized product **59**. The best aryl iodine precatalyst evaluated was C_2 -symmetric axially chiral BINI **VI**. Strikingly, while a stoichiometric amount of chiral aryl iodine difluoride **XXXIb** has been reported to give **59** with 81% ee,⁵⁷ no enantioselectivity was observed with parent aryl iodine **XXXIa** under catalytic conditions. This observation suggests different type of mechanisms between the catalytic and stoichiometric processes.

Scheme 22 Access to enantioenriched 3-aminopiperidine **59**.

Very recently, the highly enantioselective *syn*- β -fluoroaziridination of cinnamyltosylamide **62** has been disclosed by Jacobsen *et al.* (Scheme 23).⁵⁸ The optimal reaction conditions are similar to those previously reported by the group for fluorolactonization (see section 4.1.1) with **XXXII** as the precatalyst of choice. In contrast with the above fluoropiperidination reported by Kita, Shibata *et al.*, the reaction would be initiated by oxidation of the electron-rich internal alkene by the *in situ* generated ArIF_2 furnishing an electrophilic iodonium. Nucleophilic attack by the excess of fluoride would generate *anti*-vicinal fluoroiodonium **63**, which would be intramolecularly trapped by the sulfonamide to give fluoroaziridines **64**. This reaction sequence accounts for the total *syn*-selectivity. The process is limited to electron deficient cinnamyl tosylamides since substrates lacking electron withdrawing substituents gave rise to 1,1-difluoromethylated products **66** via phenonium ion **65** (Scheme 23, R = H, EDG). Carbonyl-based protected amines (carbamates, amides) led to the formation of 1,2-oxy-fluorinated products via intramolecular nucleophilic attack of the carbonyl oxygen onto the fluoroiodonium rather than the nitrogen.

Scheme 23 Synthesis of syn-β-fluoroaziridines and competitive rearrangement.

In 2014, Wirth *et al.* reported an elegant intramolecular diamination of homoallylic guanidine and diaminosulfone derivatives **67**.⁵⁹ In this transformation, the two attacking nucleophiles are on the substrate. Activation of the double bond by the hypervalent iodine triggered the first nucleophilic attack of one nitrogen to the more substituted carbon. S_N2 -type nucleophilic substitution of the resulting hypervalent moiety (intermediate **68**) delivered bicyclic molecules **69**. While poor enantioselectivity was obtained with the flexible C_2 -symmetric Ishihara's type catalyst, the design of a much more rigid catalyst **XXXIII** featuring a pyridine moiety attached to a chiral benzylic center afforded the diamines with up to 86% *ee*. The pyridine nitrogen would coordinate iodine (III). The authors did not rule out also a possible coordination of the oxygen atom belonging to the methoxy group to iodine (III). It is worth pointing out that sodium perborate in the presence of acetic acid as an activator was the stoichiometric oxidant of choice with this catalyst instead of the commonly employed mCPBA.

Scheme 24 Intramolecular diamination leading to dicyclic product.

4.2. Intermolecular version

4.2.1. Dioxygenation. In 2016, Ishihara and Muñiz *et al.* reported the first intermolecular asymmetric alkene dioxygenation under hypervalent iodine catalysis.⁶⁰ By using C_2 -symmetric bislactamide **XXXIV**, peracetic acid as the terminal oxidant and triflic acid as a

Brønsted acid co-catalyst in acetic acid, vicinal diacetoxylation of terminal styrenes **70** took place in good yields with up to 94% *ee* (Scheme 25). X-ray analyses and $^1\text{H-NMR}$ spectroscopy of the corresponding diacetate iodonium to **XXXIV** unveiled hydrogen bonding between the amide NH groups of the arms and ligand acetoxo, resulting in the creation of a supramolecular helical chirality around the central iodine (III) atom. From a mechanistic point of view, the catalytic cycle would start with the oxidation of **XXXIV** by peracetic acid in acetic acid to generate intermediary diacetoxylodinium **71**. Addition of triflic acid would allow the formation of a more reactive cationic catalyst state **72**⁶¹ which no longer exhibits C_2 -symmetry. Upon oxidation of the styrene and subsequent nucleophilic attack of acetate, intermediate **73** would be formed. Intramolecular nucleophilic addition of the acetate would regenerate catalyst **XXXIV** along with the formation of dioxolonium **74**. The latter would undergo ring opening by acetate or hydrolysis to provide vicinally dioxygenated product **75** upon treatment with acetic anhydride.

Scheme 25 Diacetoxylation of styrene.

Muñiz *et al.* have also demonstrated the use of bislactate **XXXV** to efficiently catalyze vicinal diacetoxylation of external and internal styrenes with up to 96% *ee* (Scheme 26).⁶² This protocol relies on the use of Selectfluor as the terminal oxidant

(to avoid epoxidation of the alkene observed with the commonly employed peracids) and a stoichiometric amount of trimethylsilyltriflate as an iodonium co-activator.

Scheme 26 Diacetoxylation of styrene with bis-lactate **XXXV** and selectfluor.

4.2.2. Difluorination. In 2016, Jacobsen *et al.* described asymmetric vicinal difluorination of trisubstituted cinnamide derivative **76** in 93% *ee* and with excellent *anti*-diastereoselectivity using resorcinol-based diester **XXXVI** (Scheme 27, left part).⁶³ In combination with *m*CPBA as stoichiometric oxidant and pyridine-9HF as nucleophilic fluorine source, the *in situ* generated reactive iodoarene difluoride would activate the internal alkene. Nucleophilic fluorination at the benzylic carbon would lead to *anti*-fluoroalkyl iodonium (III) intermediate **78**. Anchimeric nucleophilic assistance by the neighboring amide group (red arrows) would allow the second fluorination to take place via intermediate **79** with an overall total *anti*-diastereoselectivity to afford 1,2-difluorinated product **80**. In the case of disubstituted cinnamide derivatives **77** ($R^2 = H$), fluoroalkyl iodonium (III) intermediate **78** would preferentially undergo aryl migration (blue arrows) via phenonium ion **81** to afford gem-difluoro compounds **82** in good yields and with excellent enantioselectivities using precatalyst **XXXII** (Scheme 27, right part).⁶⁴ This elegant migratory geminal difluorination has been extended to secondary and tertiary cinnamate ester derivatives. It is worth mentioning that, in both transformations, higher enantioselectivities were obtained with aryl iodine bearing benzylic versus aliphatic substituents at the stereogenic centers probably due to attractive π -cation interactions in the transition states.

Scheme 27 Vicinal and geminal difluorination of alkenes.

Gilmour *et al.* reported vicinal difluorination of terminal aliphatic alkene **83** with modest yield and enantioselectivity using catalyst **XVIII**, Selectfluor as stoichiometric oxidant and amine-5HF (Scheme 28).⁶⁵

Scheme 28 Difluorination of terminal aliphatic alkene **83**.

4.2.3. Diamination

In 2017, Muñiz *et al.* reported an iodine (I/III)-catalyzed enantioselective vicinal diamination of internal and external styrenes using *m*CPBA as a terminal oxidant and bismesylimide as a nitrogen source (Scheme 29).⁶⁶ The undesired epoxidation background reaction was advantageously reduced when MTBE/HFIP or MTBE/TFE solvent combination were used at low temperature. Tertiary amide iodoresorcinol derivative **XXXVII** was the catalyst of choice to provide diamines **85** in good yields and with excellent enantioselectivity. Since the previously observed supramolecular arrangement of resorcinol derivatives with secondary amides through intramolecular hydrogen-bonding is not directly possible with this tertiary amide version, the intervention of water for the formation an analogous helical framework through intermolecular hydrogen-bonding has been proposed. From a mechanistic point of view, the authors

suggested the initial formation of key reactive catalyst diaminoiodonium **86** which would activate the alkene. Subsequent nucleophilic addition of the bismesylimide on the benzylic carbon would give *anti*-vicinal aminoalkyliodonium (III) **87**. Intramolecular displacement of the iodonium by the bisulfonimide with inversion of configuration would liberate the catalyst and generate cyclic intermediate **88**. Ring opening of this intermediate would allow the second C–N bond formation with second inversion of configuration, which would account for the *anti*-diastereoselectivity observed on diamines products **85**.

Scheme 29 Vicinal diaminations of alkenes.

Conclusions

The blossoming research field of asymmetric iodine catalysis has been summarized. Since the pioneering iodoarenes of Wirth *et al.* with a chiral coordinating group *ortho* to the iodine atom, we have seen that a handful of chiral arylidines has been developed and used efficiently as precatalyst of hypervalent iodine (III) to perform oxidative transformations with high enantiocontrol. Although the flexible and easily tunable C₂-symmetric resorcinol derivative seems to be a privileged structure in many transformations, other compounds with either central, axial or planar chirality have appeared to be also suitable precatalysts. Chiral quaternary ammonium iodides

have been also ingeniously employed as precursors of hypoiodites for iodine (-I/+I) catalysis. Both such catalytic systems have proven to be ideal methods for α -functionalization of carbonyl compounds, dearomatization of phenol derivatives and functionalization of alkenes with different type of nucleophiles (in an intra- or intermolecular way) in high enantioselectivities. Moreover, recent mechanistic investigations via X-Ray, NMR and computational analysis have allowed to shed on light on the structure of the reactive catalysts and the involved intermediaries in the catalytic cycles. These insights are expected to contribute in the near future to the design of new chiral iodine compounds and their applications in new asymmetric reactions.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The CNRS is gratefully acknowledged for financial supports.

Notes and references

- a) A. Yoshimura and V. V. Zhdankin, *Chem. Rev.*, 2016, **116**, 3328–3435; b) M. S. Yusubov and V. V. Zhdankin, *Resource-Efficient Technologies*, 2015, **1**, 49.
- F. Berthiol, *Synthesis*, 2015, **47**, 587.
- a) V. V. Zhdankin, *Arkivoc*, **2009**, 1; b) S. Ghosh, S. Pradhan and I. Chatterjee, *Beilstein J. Org. Chem.*, 2018, **14**, 1244.
- R. D. Richardson, T. K. Page, S. Altermann, S. M. Paradine, A. N. French, T. Wirth, *Synlett*, **2007**, 538.
- During the preparation of our manuscript, Muñiz *et al.* have published a review: A. Flores, E. Cots, J. Berges and K. Muñiz, *Adv. Synth. Catal.*, 2018, 10.1002/adsc.201800521.
- T. Hashimoto, K. Maruoka, *Chem. Rev.* 2007, **107**, 5656.
- M. Uyanik, K. Ishihara, *ChemCatChem*, 2012, **4**, 177.
- a) T. Dohi and Y. Kita, *Chem. Commun.* **2009**, 2073; b) F. V. Singh and T. Wirth, *Chem. Asian J.*, 2014, **9**, 950.
- Y. Yamamoto and H. Togo, *Synlett*, **2006**, 798.
- a) T. Wirth and U. H. Hirth, *Tetrahedron: Asym.*, 1997, **8**, 23; b) U. H. Hirt, B. Spingler and T. Wirth, *J. Org. Chem.*, 1998, **63**, 7674; c) U. H. Hirt, M. F. H. Schuster, A. N. French, O. G. Wiest and T. Wirth, *Eur. J. Org. Chem.*, **2001**, 1569.
- S. M. Altermann, R. D. Richardson, T. K. Page, R. K. Schmidt, E. Holland, U. Mohammed, S. M. Paradine, A. N. French, C. Richter, A. M. Bahar, B. Witulski and T. Wirth, *Eur. J. Org. Chem.*, **2008**, 5315.
- U. Farooq, S. Schäfer, A.-U.-H. A. Shah, D. M. Freudendahl and T. Wirth, *Synthesis*, **2010**, 1023.
- A.-A. Guilbault and C. Y. Legault, *ACS Catal.*, 2012, **2**, 219.
- A.-A. Guilbault, B. Basdevant, V. Wanie and Cl. Y. Legault, *J. Org. Chem.*, 2012, **77**, 11283.
- J. Yu, J. Cui, X.-S. Hou, S.-S. Liu, W.-C. Gao, S. Jiang, J. Tian and C. Zhang, *Tetrahedron Asym.*, 2011, **22**, 2039.
- S. Brenet, F. Berthiol and J. Einhorn, *Eur. J. Org. Chem.*, **2013**, 8094.
- S. Brenet, C. Minozzi, B. Clarens, L. Amiri and F. Berthiol, *Synthesis* 2015, **47**, 3859.
- (a) G. Levitre, A. Dumoulin, P. Retailleau, A. Panossian, F. R. Leroux and G. Masson, *J. Org. Chem.*, 2017, **82**, 11877. (b) J. Bortoluzzi, V. Jha, G. Levitre, M. J. Fer, J. Berreur, G. Masson,

- A. Panossian, F. R. Leroux, *Org. Chem.*, **2018**, doi: 10.1021/acs.joc.8b00648.
- 19 a) D. Augros, B. Yalcouye, A. Berthelot-Brehier, S. Choppin, M. Chesse, A. Panossian and F. R. Leroux, *Tetrahedron*, 2016, **72**, 5208; b) A. Berthelot-Brehier, A. Panossian, F. Colobert and F. R. Leroux, *Org. Chem. Front.*, 2015, **2**, 634; c) F. R. Leroux, A. Panossian and D. C. R. Augros, *Chim.* 2017, **20**, 682.
 - 20 S. Beaulieu and C. Y. Legault, *Chem. Eur. J.* 2015, **21**, 11206.
 - 21 a) B. Basdevant and C. Y. Legault, *Org. Lett.*, 2015, **17**, 4918. b) B. Basdevant, A.-A. Guilbault, S. Beaulieu, A. Jobin-Des Lauriers and C. Y. Legault, *Pure Appl. Chem.*, 2017, **89**, 781.
 - 22 Y. Feng, R. Huang, L. Hu, Y. Xiong and V. Coeffard, *Synthesis* 2016, **48**, 2637.
 - 23 T. Dohi, A. Maruyama, M. Yoshimura, K. Morimoto, H. Tohma and Y. Kita, *Angew. Chem. Int. Ed.*, 2004, **43**, 3595.
 - 24 A. Rodríguez and W. J. Moran, *Synthesis*, 2012, **44**, 1178.
 - 25 M. Uyanik, H. Okamoto, T. Yasui and K. Ishihara, *Science*, 2010, **328**, 1376.
 - 26 M. Uyanik, H. Hayashi and K. Ishihara, *Science*, 2014, **345**, 291.
 - 27 M. Uyanik, H. Hayashi, H. Iwata and K. Ishihara, *Chem. Lett.*, 2016, **45**, 353.
 - 28 S. Suzuki, T. Kamo, K. Fukushi, T. Hiramatsu, E. Tokunaga, T. Dohi, Y. Kita and N. Shibata, *Chem. Sci.*, 2014, **5**, 2754.
 - 29 R. Pluta, P. E. Krach, L. Cavallo, L. Falvienne and M. Rueping, *ACS Catal.*, 2018, **8**, 2582.
 - 30 Y. Wang, H. Yuan, H. Lu and W.-H. Zheng, *Org. Lett.*, 2018, **20**, 2555.
 - 31 H. Wu, Y.-P. He, L. Xu, D.-Y. Zhang and L.-Z. Gong, *Angew. Chem. Int. Ed.*, 2014, **53**, 3466.
 - 32 A. Sreenithya, C. Patel, C. M. Hadad and R. B. Sunoj, *ACS catal.*, 2017, **7**, 4189.
 - 33 a) S. Quideau, L. Pouységu, P. A. Peixoto and D. Deffieux, *Top. Curr. Chem.*, 2016, **373**, 25. b) A. M. Harned, *Tetrahedron Lett.*, 2014, **55**, 4681.
 - 34 a) M. Uyanik, T. Yasui, and Kazuaki Ishihara, *Angew. Chem. Int. Ed.*, 2010, **49**, 2175; b) M. Uyanik, T. Yasui and K. Ishihara, *Tetrahedron*, 2010, **66**, 5841.
 - 35 M. Uyanik, T. Yasui and K. Ishihara, *J. Org. Chem.*, 2017, **82**, 11946.
 - 36 M. Uyanik, T. Yasui and K. Ishihara, *Angew. Chem. Int. Ed.*, 2013, **52**, 9215.
 - 37 S. J. Murray and H. Ibrahim, *Chem. Commun.*, 2015, **51**, 2376.
 - 38 T. Dohi, A. Maruyama, N. Takenaga, K. Senami, Y. Minamitsuji, H. Fujioka, S. B. Caemmerer and Y. Kita, *Angew. Chem. Int. Ed.*, 2008, **47**, 3787.
 - 39 T. Dohi, N. Takenaga, T. Nakae, Y. Toyoda, M. Yamasaki, M. Shiro, H. Fujioka, A. Maruyama and Y. Kita, *J. Am. Chem. Soc.*, 2013, **135**, 4558.
 - 40 M. Bekkaye and G. Masson, *Synthesis*, 2016, **48**, 302.
 - 41 T. Dohi, H. Sasa, K. Miyazaki, M. Fujitake, N. Takenaga and Y. Kita, *J. Org. Chem.*, 2017, **82**, 11954.
 - 42 M. Uyanik, N. Sasakura, E. Kaneko, K. Ohori and K. Ishihara, *Chem. Lett.*, 2015, **44**, 179.
 - 43 Mechanistic studies on the activation of iodine (III) derivatives with fluorinated alcohols: I. Colomer, C. Batchelor-McAuley, B. Odell, T. J. Donohoe and R. G. Compton, *J. Am. Chem. Soc.*, 2016, **138**, 8855.
 - 44 M. Uyanik, N. Sasakura, M. Mizuno and K. Ishihara, *ACS Catal.*, 2017, **7**, 872.
 - 45 N. Jain, S. Xu, and M. A. Ciufolini, *Chem. Eur. J.*, 2017, **23**, 4542.
 - 46 D.-Y. Zhang, L. Xu, H. Wu and L.-Z. Gong, *Chem. Eur. J.*, 2015, **21**, 10314.
 - 47 L. Kürti, P. Herczegh, J. Visy, M. Simonyi, S. Antus and A. Pelter, *J. Chem. Soc., Perkin Trans. 1*, **1999**, 379.
 - 48 S. Quideau, G. Lyvinec, M. Marguerit, K. Bathany, A. Ozanne Beaudenon, T. Buffeteau, D. Cavagnat and A. Chénédé, *Angew. Chem. Int. Ed.*, 2009, **48**, 4605.
 - 49 K. A. Volp and A. M. Harned, *Chem. Commun.*, 2013, **49**, 3001.
 - 50 K. Muñiz and L. Fra, *Synthesis*, 2017, **49**, 2901.
 - 51 T. Hashimoto, Y. Shimazaki, Y. Omatsu and K. Maruoka, *Angew. Chem. Int. Ed.*, 2018, **57**, 7200.
 - 52 R. M. Romero, T. H. Wöste and K. Muñiz, *Chem. Asian J.*, 2014, **9**, 972; b) For a review on mechanistic aspects: M. Fujita, *Tetrahedron Lett.*, 2017, **58**, 4409.
 - 53 M. Fujita, K. Mori, M. Shimogaki and T. Sugimura, *Org. Lett.*, 2012, **14**, 1294.
 - 54 M. Shimogaki, M. Fujita and T. Sugimura, *Eur. J. Org. Chem.*, **2013**, 7128.
 - 55 C. Gelis, A. Dumoulin, M. Bekkaye, L. Neuville and G. Masson, *Org. Lett.*, 2017, **19**, 278.
 - 56 E. M. Woerly, S. M. Banik and E. N. Jacobsen, *J. Am. Chem. Soc.*, 2016, **138**, 13858.
 - 57 W. Kong, P. Feige, T. de Haro and C. Nevado, *Angew. Chem. Int. Ed.*, 2013, **52**, 2469.
 - 58 K. M. Mennie, S. M. Banik, E. C. Reichert and E. N. Jacobsen, *J. Am. Chem. Soc.*, 2018, **140**, 4797.
 - 59 P. Mizar, A. Laverny, M. El-Sherbini, U. Farid, M. Brown, F. Malmedy and T. Wirth, *Chem. Eur. J.*, 2014, **20**, 9910.
 - 60 S. Haubenreisser, T. H. Wöste, C. Martinez, K. Ishihara and K. Muñiz, *Angew. Chem. Int. Ed.*, 2016, **55**, 413.
 - 61 Y.-B. Kang and L. H. Gade, *J. Am. Chem. Soc.*, 2011, **133**, 3658.
 - 62 T. H. Wöste, K. Muñiz, *Synthesis* 2016, **48**, 816.
 - 63 S. M. Banik, J. W. Medley, E. N. Jacobsen, *J. Am. Chem. Soc.*, 2016, **138**, 5000.
 - 64 S. M. Banik, J. W. Medley and E. N. Jacobsen, *Science*, 2016, **353**, 51.
 - 65 I. G. Molnár, C. Thiehoff, M. C. Holland and R. Gilmour, *ACS Catal.*, 2016, **6**, 7167.
 - 66 K. Muñiz, L. Barreiro, R. M. Romero and C. Martinez, *J. Am. Chem. Soc.*, 2017, **139**, 4354.