A new link between the descent algebra of type B, domino tableaux and Chow's quasisymmetric functions

Ekaterina A. Vassilieva, Alina Mayorova

► To cite this version:

Ekaterina A. Vassilieva, Alina Mayorova. A new link between the descent algebra of type B, domino tableaux and Chow's quasisymmetric functions. *Discrete Mathematics*, 2019, 342 (6), pp.1658-1673. 10.1016/j.disc.2019.01.021 . hal-02410964

HAL Id: hal-02410964

<https://hal.science/hal-02410964>

Submitted on 14 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A new link between the descent algebra of type B, domino tableaux and Chow's quasisymmetric functions

Ekaterina A. Vassilieva^a, Alina R. Mayorova^{a,b}

^a*Laboratoire d'Informatique de l'Ecole Polytechnique, 91128 Palaiseau Cedex, France*

^b*Department of Higher Algebra, Moscow State University, Moscow, Russia*

Abstract

Introduced by Solomon in his 1976 paper, the descent algebra of a finite Coxeter group received significant attention over the past decades. As proved by Gessel, in the case of the symmetric group its structure constants give the comultiplication table for the fundamental basis of quasisymmetric functions. We show that this latter property actually implies several well known relations linked to the Robinson-Schensted-Knuth correspondence and some of its generalisations. This provides a new link between these results and the theory of quasisymmetric functions and allows to derive more advanced formulas involving Kronecker coefficients. Using the theory of type B quasisymmetric functions introduced by Chow, we extend this connection to the hyperoctahedral group and derive new formulas relating the structure constants of the descent algebra of type B, the numbers of domino tableaux of given descent set and the Kronecker coefficients of the hyperoctahedral group.

Keywords: descent algebra, type B quasisymmetric functions, Kronecker coefficients, RSK-correspondence, hyperoctahedral group, domino tableaux.

1. Introduction

For any positive integer n write $[n] = \{1, \dots, n\}$, S_n the symmetric group on $[n]$ and id_n the identity permutation of S_n . A **composition** $\alpha \models n$ is a sequence of positive integers $\alpha = (\alpha_1, \dots, \alpha_p)$ such that $\alpha_1 + \alpha_2 + \dots = n$. A composition λ of n with its parts sorted in decreasing order is called a **partition** and denoted $\lambda \vdash n$ or $|\lambda| = n$. A partition λ is usually represented as a Young diagram of $n = |\lambda|$ boxes arranged in $\ell(\lambda)$ left justified rows so that the i -th row from the top contains λ_i boxes. A **standard Young tableau** T is a Young diagram whose boxes are filled with the elements of $[n]$ such that the entries are strictly increasing along the rows and down the columns. The partition given by the number of boxes in each row is its **shape** and denoted $shape(T)$.

Email addresses: `ekaterina.vassilieva@lix.polytechnique.fr` (Ekaterina A. Vassilieva), `alina.r.m@yandex.ru` (Alina R. Mayorova)

Example 1.1. The following diagrams are standard Young tableaux of shape $\lambda = (6, 4, 2, 1, 1)$.

$$T_1 = \begin{array}{|c|c|c|c|c|c|} \hline 1 & 3 & 4 & 8 & 9 & 14 \\ \hline 2 & 6 & 7 & 10 & & \\ \hline 5 & 12 & & & & \\ \hline 11 & & & & & \\ \hline 13 & & & & & \\ \hline \end{array} \quad T_2 = \begin{array}{|c|c|c|c|c|c|} \hline 1 & 3 & 4 & 7 & 8 & 9 \\ \hline 2 & 6 & 12 & 14 & & \\ \hline 5 & 13 & & & & \\ \hline 10 & & & & & \\ \hline 11 & & & & & \\ \hline \end{array}$$

Compositions of n are in natural bijection with subsets of $[n-1]$. For $\alpha \models n$ and $I = \{i_1, i_2, \dots, i_m\}$ a subset of $[n-1]$ such that $i_1 < i_2 < \dots < i_m$, we denote $Des(\alpha) = \{\alpha_1, \alpha_1 + \alpha_2, \dots, \alpha_1 + \alpha_2 + \dots + \alpha_{p-1}\}$ and $comp(I) = (i_1, i_2 - i_1, \dots, i_m - i_{m-1}, n - i_m)$ this bijection and its inverse. As a result the number of compositions of n is 2^{n-1} .

One important feature of a permutation π of S_n is its **descent set** $Des(\pi) = \{1 \leq i \leq n-1 \mid \pi(i) > \pi(i+1)\}$. Similarly define the **descent set of a standard Young tableau** T as the subset of $[n-1]$ defined by $Des(T) = \{1 \leq i \leq n-1 \mid i \text{ is in a strictly higher row than } i+1\}$. For instance the descent set of the tableaux in Example 1.1 is $\{1, 4, 9, 10, 12\}$. We denote $d_{\lambda I}$ the number of standard Young tableaux of shape λ and descent set I .

Let D_I (resp. B_I) be the element of the algebra $\mathbb{C}S_n$ defined as the formal sum of all the permutations π such that $Des(\pi) = I$ (resp. $Des(\pi) \subseteq I$). In the more general context of finite Coxeter groups, Solomon showed in [19] that the D_I 's (resp. B_I 's) generate a subalgebra of $\mathbb{C}S_n$ of dimension 2^{n-1} usually referred to as the **Solomon's descent algebra**. More specifically, he showed that there exist (non negative integer valued) **structure constants** $(a_{IJ}^K)_{I,J,K \subseteq [n-1]}$ and $(b_{IJ}^K)_{I,J,K \subseteq [n-1]}$ that verify:

$$D_I D_J = \sum_{K \subseteq [n-1]} a_{IJ}^K D_K, \quad B_I B_J = \sum_{K \subseteq [n-1]} b_{IJ}^K B_K.$$

As a result the number of ways to write a fixed permutation π of D_K as the ordered product of two permutations $\pi = \sigma\tau$ such that $\sigma \in D_I$ and $\tau \in D_J$ depends only on $Des(\pi) = K$ and is equal to a_{IJ}^K . If we require instead σ and τ to be respectively in B_I and B_J then the number of such products is $\sum_{K' \supseteq K} b_{IJ}^{K'}$.

Remark 1.2. The two families of structure constants are linked through the formulas

$$\sum_{I' \subseteq I, J' \subseteq J} a_{I'J'}^K = \sum_{K' \supseteq K} b_{IJ}^{K'}. \quad (1)$$

Because of its important combinatorial and algebraic properties the descent algebra received significant attention afterwards. In particular, Garsia and Reutenauer in [11] provide a decomposition of its multiplicative structure and a new combinatorial interpretation of the coefficients b_{IJ}^K in terms of the

number of non-negative integer matrices with specified constraints. Bergeron and Bergeron give analogous results in [4] when the symmetric group is replaced by the hyperoctahedral group (Coxeter group of type B). As shown by Norton in [16], the descent algebra of a finite Coxeter group is also strongly related to its *0-Hecke algebra*. In particular, she proves that the dimension of each left principal indecomposable module of the 0-Hecke algebra is equal to the cardinality of the analogue of one the D_I 's. She further shows that the analogues of the $a_{IJ}^\emptyset$ are the entries of the *Cartan matrix* giving the number of times each irreducible module is a composition factor of each indecomposable module. In the specific case of the symmetric group S_n , Carter in [6] uses the celebrated *Robinson-Schensted (RS)* correspondence to explain the following relation obtained by computation of the Cartan matrix

$$a_{IJ}^\emptyset = \sum_{\lambda \vdash n} d_{\lambda I} d_{\lambda J}. \quad (2)$$

Finally, Solomon's descent algebra is dual to the Hopf algebra of *quasisymmetric functions* (see Section 2) introduced by Gessel in [12]. In particular, he shows that the comultiplication table for their fundamental basis is given by the a_{IJ}^K 's. In Section 2, we show that Equation (2) and its generalisations are a direct consequence of Gessel's result. Then we use the theory of quasisymmetric functions of type B introduced by Chow ([7]) in Section 3 to provide new analogous formulas in the case of the hyperoctahedral group. Section 4 provides additional results involving more general coefficients.

2. Descent algebra of the symmetric group

2.1. Quasisymmetric functions

Let $X = \{x_1, x_2, \dots\}$ be an alphabet of commutative indeterminates and $I \subseteq [n-1]$. A quasisymmetric function is a bounded degree formal power series in $\mathbb{C}[X]$ such that for any composition $(\alpha_1, \dots, \alpha_p)$ and any strictly increasing sequence of distinct indices $i_1 < i_2 < \dots < i_p$ the coefficient of $x_1^{\alpha_1} x_2^{\alpha_2} \dots x_p^{\alpha_p}$ is equal to the coefficient of $x_{i_1}^{\alpha_1} x_{i_2}^{\alpha_2} \dots x_{i_p}^{\alpha_p}$. Quasisymmetric functions admit the monomial and fundamental quasisymmetric functions as classical bases:

$$M_I(X) = \sum_{\substack{i_1 \leq \dots \leq i_n \\ k \in I \Leftrightarrow i_k < i_{k+1}}} x_{i_1} x_{i_2} \dots x_{i_n}, \quad F_I(X) = \sum_{\substack{i_1 \leq \dots \leq i_n \\ k \in I \Rightarrow i_k < i_{k+1}}} x_{i_1} x_{i_2} \dots x_{i_n}.$$

These two bases are related through

$$F_I(X) = \sum_{I \subseteq J \subseteq [n-1]} M_J(X). \quad (3)$$

Remark 2.1. In Gessel's paper [12], quasisymmetric functions are indexed by compositions and not subsets. However, the bijection between subsets and compositions given in Section 1 makes the definition equivalent.

For $\lambda \vdash n$ denote also $p_\lambda(X)$ and $s_\lambda(X)$ the power sum and Schur symmetric functions in the alphabet X indexed by λ . In what follows, we may remove the reference to X when there is no confusion.

A **semistandard Young tableau** is a Young diagram whose entries are strictly increasing down the column and non-decreasing along the rows. A tableau has weight $\mu = (\mu_1, \mu_2, \dots)$ if it has μ_i entries equal to i . Denote $K_{\lambda\mu}$ the number of *semistandard Young tableaux* of shape λ and weight μ . The decomposition of Schur symmetric functions in the fundamental (e.g. [20, 7.19.7]) and the monomial bases is

$$s_\lambda(X) = \sum_{I \subseteq [n-1]} d_{\lambda I} F_I(X) \quad (4)$$

$$= \sum_{I \subseteq [n-1]} K_{\lambda \text{ comp}(I)} M_I(X). \quad (5)$$

2.2. Gessel's relation and its consequences

For two commutative sets of variables $X = \{x_1, x_2, \dots, x_i, \dots\}$ and $Y = \{y_1, y_2, \dots, y_i, \dots\}$, we denote XY the set of indeterminates $\{x_i y_j; x_i \in X, y_j \in Y\}$ ordered by the lexicographical order. Gessel shows in [12, Thm. 11] that for any subset $K \subseteq [n-1]$

$$F_K(XY) = \sum_{I, J \subseteq [n-1]} a_{IJ}^K F_I(X) F_J(Y). \quad (6)$$

We can say that $F_K(XY)$ is the generating series for the coefficients a_{IJ}^K . As stated in introduction, Equation (2) and some of its generalisations are a direct consequence of Equation (6). More precisely, denote by χ^λ the irreducible character of S_n indexed by partition λ and for any $\lambda, \mu, \nu \vdash n$ define the **Kronecker coefficients** $g(\lambda, \mu, \nu) = 1/n! \sum_{\omega \in S_n} \chi^\lambda(\omega) \chi^\mu(\omega) \chi^\nu(\omega)$. We have the following theorem.

Theorem 2.2. *For $I, J, K \subset [n-1]$ let $a_{IJ}^\emptyset$ be the structure constant defined in introduction and let $a_{I,J,K}^\emptyset = [D_\emptyset] D_I D_J D_K$ be the number of triples of permutations $\sigma_1, \sigma_2, \sigma_3$ of S_n such that $\text{Des}(\sigma_1) = I, \text{Des}(\sigma_2) = J, \text{Des}(\sigma_3) = K$ and $\sigma_1 \sigma_2 \sigma_3 = \text{id}_n$. Equation (6) directly implies*

$$a_{IJ}^\emptyset = \sum_{\lambda \vdash n} d_{\lambda I} d_{\lambda J}, \quad (7)$$

$$a_{I,J,K}^\emptyset = \sum_{\lambda, \mu, \nu \vdash n} g(\lambda, \mu, \nu) d_{\lambda I} d_{\mu J} d_{\nu K}. \quad (8)$$

Proof. According to Equation (4) $F_\emptyset = s_{(n)}$. Then using the *Cauchy identity* for Schur functions

$$s_{(n)}(XY) = \sum_{\lambda \vdash n} s_\lambda(X) s_\lambda(Y) \quad (9)$$

and applying Equation (4) once again one gets:

$$\begin{aligned}
\sum_{I, J \subseteq [n-1]} a_{IJ}^\emptyset F_I(X) F_J(Y) &= F_\emptyset(XY) = s_{(n)}(XY) \\
&= \sum_{\lambda \vdash n} s_\lambda(X) s_\lambda(Y) \\
&= \sum_{\substack{\lambda \vdash n \\ I, J \subseteq [n-1]}} d_{\lambda I} d_{\lambda J} F_I(X) F_J(Y).
\end{aligned}$$

This can be generalised to prove Equation (8) by noticing that the structure constants verify $a_{I,J,K}^\emptyset = \sum_L a_{I,L}^\emptyset a_{J,K}^L$. As a result, Equation (6) extends to

$$F_\emptyset(XYZ) = \sum_{I, J, K} a_{I,J,K}^\emptyset F_I(X) F_J(Y) F_K(Z).$$

Finally use the generalised version of Equation (9)

$$s_\nu(XY) = \sum_{\lambda, \mu} g(\lambda, \mu, \nu) s_\lambda(X) s_\mu(Y) \quad (10)$$

to decompose $s_{(n)}(XYZ) = F_\emptyset(XYZ)$. $\square$

Remark 2.3. Denote λ' the partition corresponding to the transposed Young diagram of λ . For $I, J \subseteq [n-1]$, the numbers $a_{IJ}^{[n-1]}$ are given by

$$a_{IJ}^{[n-1]} = \sum_{\lambda \vdash n} d_{\lambda I} d_{\lambda' J}. \quad (11)$$

Proof. According to Equation (4) $F_{[n-1]} = s_{(1^n)}$. Then use the identity for Schur functions

$$s_{(1^n)}(XY) = \sum_{\lambda \vdash n} s_\lambda(X) s_{\lambda'}(Y)$$

and apply Equation (4) to s_λ and $s_{\lambda'}$. $\square$

2.3. Extension to the RSK-correspondence

Generalising the RS-correspondence to matrices with non-negative integral entries, Knuth ([14]) proved that for $r, c \models n$ the number $m_{r,c}$ of such matrices with row and column sums equal respectively to r and c is given by

$$m_{r,c} = \sum_{\lambda \vdash n} K_{\lambda r} K_{\lambda c}. \quad (12)$$

We have the following corollary to Theorem 2.2.

Corollary 2.4. Equation (12) is also a consequence of Equation (6). Furthermore let p, q and r be three compositions of n and denote $m_{p,q,r}$ the number of three-dimensional arrays $M = (M_{i,j,k})$ with non-negative integer entries such that $p_k = \sum_{i,j} M_{i,j,k}$, $q_j = \sum_{i,k} M_{i,j,k}$ and $r_i = \sum_{j,k} M_{i,j,k}$. Equation (6) implies

$$m_{p,q,r} = \sum_{\lambda, \mu, \nu \vdash n} g(\lambda, \mu, \nu) K_{\lambda p} K_{\mu q} K_{\nu r}. \quad (13)$$

Proof. We proceed with the proof of Equation (12).

Denote $A_{I,J}^K = \sum_{I' \subseteq I, J' \subseteq J} a_{I',J'}^K$. According to Equation (3)

$$\begin{aligned} \sum_{I', J' \subseteq [n-1]} a_{I', J'}^K F_{I'}(X) F_{J'}(Y) &= \sum_{I', J' \subseteq [n-1]} a_{I', J'}^K \sum_{I \supseteq I', J \supseteq J'} M_I(X) M_J(Y) \\ &= \sum_{I, J \subseteq [n-1]} M_I(X) M_J(Y) \sum_{I' \subseteq I, J' \subseteq J} a_{I', J'}^K \\ &= \sum_{I, J \subseteq [n-1]} A_{I,J}^K M_I(X) M_J(Y). \end{aligned}$$

As a result, $A_{I,J}^\emptyset$ is the coefficient in $M_I(X) M_J(Y)$ of $F_\emptyset(XY)$ which we know is equal to $\sum_{\lambda \vdash n} s_\lambda(X) s_\lambda(Y)$ (see above). Finally use Equation (5) to get

$$A_{I,J}^\emptyset = \sum_{\lambda \vdash n} K_{\lambda \text{ comp}(I)} K_{\lambda \text{ comp}(J)}. \quad (14)$$

It remains to prove that $A_{I,J}^\emptyset = m_{\text{comp}(I), \text{comp}(J)}$. The combinatorial interpretation of [11] states that $b_{I,J}^K$ is the number of non-negative integer matrices M with row and column sums equal to $\text{comp}(I)$ and $\text{comp}(J)$ respectively and with the word obtained by reading the entries of M row by row from top to bottom equal to K (zero entries being omitted). But according to Equation (1) $A_{I,J}^\emptyset = \sum_{K \subseteq [n-1]} b_{I,J}^K$. We get Equation (12).

In order to prove Equation (13), it suffices to notice that the interpretation of [11] generalises well to three dimensional arrays. $\square$

A bijective proof of Equation (13) is provided in [2]. The proof involves semi-standard Young tableaux and *Littlewood-Richardson tableaux*. Our approach allows us to recover these results immediately.

3. Descent algebra of the hyperoctahedral group

3.1. Signed permutations and domino tableaux

Let B_n be the **hyperoctahedral group** of order n . B_n is composed of all permutations π on the set $\{-n, \dots, -2, -1, 0, 1, 2, \dots, n\}$ such that for all i in $\{0\} \cup [n]$, $\pi(-i) = -\pi(i)$ (in particular $\pi(0) = 0$). As a result, such permutations usually referred to as **signed permutations** are entirely described by their

restriction to $[n]$. The **descent set** of π is the subset of $\{0\} \cup [n-1]$ defined by $Des(\pi) = \{0 \leq i \leq n-1 \mid \pi(i) > \pi(i+1)\}$. The main difference with respect to the case of the symmetric group is the possible descent in position 0 when $\pi(1)$ is a negative integer. We denote D_I^B (resp. B_I^B) the formal sum in $\mathbb{C}B_n$ of the signed permutations π with $Des(\pi) = I$ (resp. $Des(\pi) \subseteq I$). Let $c_{I_1, I_2, \dots, I_p}^K = [D_K^B] \prod_{j=1}^p D_{I_j}^B$ (resp. $e_{I_1, I_2, \dots, I_p}^K = [B_K^B] \prod_{j=1}^p B_{I_j}^B$) be the structure constants of Solomon's descent algebra of type B for $I_1, \dots, I_p, K \subseteq \{0\} \cup [n-1]$.

Remark 3.1. *These constants verify*

$$\sum_{I' \subseteq I, J' \subseteq J} c_{I'J'}^K = \sum_{K' \supseteq K} e_{IJ}^{K'}. \quad (15)$$

For $\lambda \vdash 2n$, a **standard domino tableau** T of shape λ is a Young diagram of shape λ tiled by **dominoes**, i.e. 2×1 or 1×2 rectangles filled with the elements of $[n]$ such that the entries are strictly increasing along the rows and down the columns. Denote $SDT(\lambda)$ the set of standard domino tableaux of shape λ . We denote $\mathcal{P}^0(n)$ the set of partitions $\lambda \vdash 2n$ whose associated Young diagram may be fully tiled by dominoes. Such partitions are usually called *empty 2-core partitions*. A standard domino tableau T has a descent in position $i > 0$ if $i+1$ lies strictly below i in T and has descent in position 0 if the domino filled with 1 is vertical. We denote $Des(T)$ the set of all its descents. For λ in $\mathcal{P}^0(n)$ and $I \subseteq \{0\} \cup [n-1]$, denote $d_{\lambda I}^B$ the number of standard domino tableaux of shape λ and descent set I .

Example 3.2. *The following standard domino tableaux have shape $(5, 5, 4, 1, 1)$ and descent set $\{0, 3, 5, 6\}$.*

1	2	3	5
			6
4		8	
7			

 $T_1 =$

1	2	3	5
		4	6
6		8	
7			

 $T_2 =$

There is a natural analogue of the RSK-correspondence for signed permutations involving **domino tableaux**. Barbash and Vogan ([3]) built a bijection between signed permutations of B_n and pairs of standard domino tableaux of equal shape in $\mathcal{P}^0(n)$. An independent development on the subject is due to Garfinkle in [8, 9, 10]. Van Leeuwen shows in [23] that the two approaches are actually equivalent. See also Stanton and White in [21] for a more general treatment of *rim hook tableaux*. Taşkın ([22, Prop. 26]) showed that the two standard domino tableaux associated to a signed permutation π by the algorithm of Barbash and Vogan have respective descent sets $Des(\pi^{-1})$ and $Des(\pi)$. As a result of this descent preserving property, we have the following analogue to Equation (2).

$$c_{IJ}^{\emptyset} = \sum_{\lambda \in \mathcal{P}^0(n)} d_{\lambda I}^B d_{\lambda J}^B. \quad (16)$$

3.2. Quasisymmetric functions of type B and main results

Chow defines in [7] an analogue of Gessel's algebra of quasisymmetric functions that is dual to the Solomon's descent algebra of type B.

Let $X = \{\cdots, x_{-i}, \cdots, x_{-1}, x_0, x_1, \cdots, x_i, \cdots\}$ be an alphabet of commutative indeterminates with $x_{-i} = x_i$ and I be a subset of $\{0\} \cup [n-1]$, he defines:

$$M_I^B(X) = \sum_{\substack{0 \leq i_1 \leq i_2 \leq \cdots \leq i_n \\ j \in I \Leftrightarrow i_j < i_{j+1}}} x_{i_1} x_{i_2} \cdots x_{i_n}, \quad F_I^B(X) = \sum_{\substack{0 \leq i_1 \leq i_2 \leq \cdots \leq i_n \\ j \in I \Rightarrow i_j < i_{j+1}}} x_{i_1} x_{i_2} \cdots x_{i_n}.$$

where $i_0 = 0$. Note the particular role played by variable x_0 .

Example 3.3. Let $n = 2$ and $X = \{x_{-2}, x_{-1}, x_0, x_1, x_2\}$ then $F_\emptyset = x_0^2 + x_1^2 + x_2^2 + x_0x_1 + x_0x_2 + x_1x_2$, $F_{\{1\}} = x_0x_1 + x_0x_2 + x_1x_2$, $F_{\{0\}} = x_1^2 + x_2^2 + x_1x_2$ and $F_{\{0,1\}} = x_1x_2$.

Remark 3.4. Prior to Chow, Poirier ([18]) introduced an alternative type B analogue of the quasisymmetric functions used for example in [1]. However Poirier's quasisymmetric functions are dual to the Mantaci-Reutenauer algebra and not Solomon's descent algebra of type B. See [17] for further details. The question of finding a similar development as in the present paper for Poirier's quasisymmetric functions and the Mantaci-Reutenauer algebra remains open.

Let Y be a second copy of X . Chow uses the theory of P -partitions of type B to show in [7, Thm. 2.3.4] that for $K \subseteq \{0\} \cup [n-1]$

$$F_K^B(XY) = \sum_{I, J \subseteq \{0\} \cup [n-1]} c_{I,J}^K F_I^B(X) F_J^B(Y). \quad (17)$$

Theorem 3.5. Equation (16) is a consequence of Equation (17).

The irreducible characters of B_n are naturally indexed by partitions of $\mathcal{P}^0(n)$ (see e.g. [15, I, Appendix B]). Denote ψ^λ the character indexed by $\lambda \in \mathcal{P}^0(n)$ and define the **Kronecker coefficients of B_n** , $g^B(\lambda, \mu, \nu) = 1/|B_n| \sum_{\omega \in B_n} \psi^\lambda(\omega) \psi^\mu(\omega) \psi^\nu(\omega)$. A more general form of Theorem 3.5 can be stated as follows.

Theorem 3.6. Let I, J and $K \subseteq \{0\} \cup [n-1]$. The following formula holds

$$c_{I,J,K}^\emptyset = \sum_{\lambda, \mu, \nu \in \mathcal{P}^0(n)} g^B(\lambda, \mu, \nu) d_{\lambda I}^B d_{\mu J}^B d_{\nu K}^B. \quad (18)$$

The proof of Theorems 3.5 and 3.6 uses Equation (17) and generating functions for domino tableaux. We detail it in the two following sections.

3.3. Domino functions

3.3.1. Definition

Generating functions for domino tableaux sometimes called *domino functions* are well studied objects (see e.g. [13]). We introduce a modified definition

to get an analogue of Schur functions verifying Equations (20) and (22) that we need to prove Theorem 3.6. Our development differs by the addition of "0" entries to the domino tableaux in some cases.

A **semistandard domino tableau** T of shape $\lambda \in \mathcal{P}^0(n)$ and weight $w(T) = \mu = (\mu_0, \mu_1, \mu_2, \dots)$ with $\mu_i \geq 0$ and $\sum_i \mu_i = n$ is a tiling of the Young diagram of shape λ with horizontal and vertical dominoes labelled with integers in $\{0, 1, 2, \dots\}$ such that labels are non decreasing along the rows, strictly increasing down the columns and exactly μ_i dominoes are labelled with i . If the top leftmost domino is vertical, it cannot be labelled 0 (we leave it to the reader to check that the only possible sub-pattern of dominos with label 0 in a semistandard domino tableau is a row composed of horizontal dominos). Denote $SSDT(\lambda)$ the set of semistandard domino tableaux of shape λ and $K_{\lambda\mu}^B$ the number of semistandard domino tableaux of shape λ and weight μ . The **standardisation** T^{st} of a semistandard domino tableau T of weight μ is the standard domino tableau obtained by relabelling the dominoes of T with $1, 2, \dots, n$ such that the dominoes labelled with $i_m = \min\{i \mid \mu_i > 0\}$ are relabelled with $1, 2, \dots, \mu_{i_m}$ from left to right and so on.

Example 3.7. Figure 5 shows a semistandard domino tableau of weight $\mu = (2, 0, 2, 0, 0, 4, 0, 1)$ and its standardisation.

Figure 1: A semistandard domino tableau and its standardisation.

Definition 3.8. Given alphabet X and a semistandard domino tableau T of weight μ , denote X^T the monomial $x_0^{\mu_0} x_1^{\mu_1} x_2^{\mu_2} \dots$. For $\lambda \in \mathcal{P}^0(n)$ we call the **domino function** indexed by λ the function defined in the alphabet X by

$$\mathcal{G}_\lambda(X) = \sum_{T \in SSDT(\lambda)} X^T. \quad (19)$$

3.3.2. Expansion into Chow's fundamental and monomial bases

We have the following proposition.

Proposition 3.9. Domino functions admit the following expansion into Chow's fundamental and monomial bases.

$$\mathcal{G}_\lambda = \sum_{I \subseteq \{0\} \cup [n-1]} d_{\lambda I}^B F_I^B = \sum_{I \subseteq \{0\} \cup [n-1]} K_{\lambda \text{comp}(I)}^B M_I^B. \quad (20)$$

In order to prove Proposition 3.9 we need the following lemma.

Lemma 3.10. *Let T_0 be a standard domino tableau. Then the fundamental quasisymmetric function $F_{Des(T_0)}^B$ is the generating function for semistandard domino tableaux T such that $T^{st} = T_0$, i.e.*

$$F_{Des(T_0)}^B = \sum_{T^{st}=T_0} X^T. \quad (21)$$

Proof. Given a fixed standard domino tableau T_0 , we reverse the standardisation operation to obtain the set of semistandard domino tableaux with standardisation T_0 . Since the standardisation operation is shape and domino tiling preserving, any such semistandard tableau T may be identified by the sequence of its entries $i_1, i_2, \dots, i_n$.

Figure 2: A standard domino tableau T_0 and the template for its semistandard preimages.

Relabel the entries of T_0 with the sequence $i_1, i_2, \dots, i_n$ such that k is mapped to i_k , as depicted on Figure 2. According to the standardisation procedure, we have $0 \leq i_1 \leq i_2 \leq \dots \leq i_n$ and $0 < i_1$ if the top leftmost domino is vertical. Furthermore such a sequence needs to verify additional conditions to give a valid semistandard domino tableau with standardisation equal to T_0 . Consider each constraint $i_k \leq i_{k+1}$ separately. If $i_k < i_{k+1}$ then locally the sequence gives a valid semistandard domino tableau (entries are increasing along the row and down the columns) and the standardisation process provides the expected result.

We look at the case $i_k = i_{k+1}$ and show two properties.

- First, if $i_k = i_{k+1}$ then $k \notin Des(T_0)$. Indeed assume $k \in Des(T_0)$. Domino k has to lay in a higher row than $k+1$ as in Figure 3. But domino i_{k+1} may not be outside the green area defined in Figure 3 as the entries in the tableau are strictly increasing down the columns. As a consequence, domino i_{k+1} has to be on the left of and below domino i_k , as in Figure 3, picture (c). This is a contradiction since according to the definition of the standardisation operation, we should relabel the domino i_{k+1} before the domino i_k with such a pattern and T_0 is not the standardisation of such a tableau. As a result $k \in Des(T_0)$ implies $i_k < i_{k+1}$.
- Secondly, when $k \notin Des(T_0)$ any sequence with $i_k = i_{k+1}$ locally gives a valid semistandard tableau whose standardisation is equal to T_0 . Indeed,

Figure 3: Layouts with a descent in position k . Green area corresponds to positions of domino i_{k+1} which do not violate the property of increasing entries along the rows and down the columns when $i_k = i_{i+1}$. In pictures (a) and (b) domino i_{k+1} lays outside the green area. Picture (c) shows a valid layout of dominoes i_k and i_{k+1} .

if $k \notin Des(T_0)$, domino k does not lay in a higher row than $k + 1$.

As a result, domino i_{k+1} cannot be outside the green area defined in Figure 4 as entries are increasing along the rows and down the columns, i.e. i_{k+1} is to the right of and above domino i_k . This makes sure that i_k is relabelled before i_{k+1} in the standardisation procedure.

Figure 4: Layouts with no descent in position k . Green area corresponds to positions of domino i_{k+1} which do not violate the property of increasing entries along the rows and down the columns when $i_k = i_{i+1}$. In pictures (a) and (b) domino i_{k+1} lays outside the green area. Picture (c) shows a valid layout of dominoes i_k and i_{k+1} .

As a consequence the preimages of T_0 can be identified as the sequences $0 \leq i_1 \leq i_2 \leq \dots \leq i_n$ such that $k \in Des(T_0) \Rightarrow i_k < i_{k+1}$ for $k \geq 0$ and $i_0 = 0$. Consider the constraints all together to ensure that such a sequence globally gives a valid semistandard tableau whose standardisation is equal to T_0 . $\square$

We are now ready to prove Proposition 3.9.

Proof of Proposition 3.9. Classify the set of semistandard domino tableaux according to their standardisation and add the monomials corresponding to the

tableaux in the same class. Using Lemma 3.10 we obtain the first part of the Proposition.

$$\begin{aligned}
\mathcal{G}_\lambda(X) &= \sum_{T \in SSDT(\lambda)} X^T \\
&= \sum_{T_0 \in SDT(\lambda)} \sum_{T \in SSDT(\lambda), T^{st}=T_0} X^T \\
&\stackrel{(21)}{=} \sum_{T_0 \in SDT(\lambda)} F_{Des(T_0)}^B(X) \\
&= \sum_{I \subseteq \{0\} \cup [n-1]} d_{\lambda I}^B F_I^B(X).
\end{aligned}$$

To get the second part of the proposition, relabel the entries of a semistandard tableau T with successive integers $0, 1, 2, \dots$ such that inequalities and equalities between entries in T are preserved after relabelling. If the top left-most domino is vertical skip 0. This operation removes the zeros in the weight of the tableau except, possibly, the first one. Denote $\tilde{T}$ the resulting tableau and for $\lambda \in \mathcal{P}^0(n)$, $\widetilde{SSDT}(\lambda)$ the set of semistandard domino tableaux of shape λ such that $\tilde{T} = T$. The second part of the statement is a consequence of the fact that the monomial quasisymmetric function $M_{set(w(T_0))}^B$ is precisely the generating function for all semistandard domino tableaux T mapped to the same $T_0 \in \widetilde{SSDT}(\lambda)$ by this operation.

$$\begin{aligned}
\mathcal{G}_\lambda(X) &= \sum_{T \in SSDT(\lambda)} X^T \\
&= \sum_{T_0 \in \widetilde{SSDT}(\lambda)} \sum_{T \in SSDT(\lambda), \tilde{T}=T_0} X^T \\
&= \sum_{T_0 \in \widetilde{SSDT}(\lambda)} M_{set(w(T_0))}^B \\
&= \sum_{I \subseteq \{0\} \cup [n-1]} K_{\lambda \text{comp}(I)}^B M_I^B(X).
\end{aligned}$$

Equation (20) is fully proven. $\square$

3.3.3. Cauchy identities

There is a well known (not descent preserving) bijection between semistandard domino tableaux of weight μ and **bi-tableaux**, i.e. pairs of semistandard Young tableaux of respective weights μ^- and μ^+ such that $\mu_i^+ + \mu_i^- = \mu_i$ for all i (see e.g. [5, Algorithm 6.1]). The respective shapes of the two Young tableaux depend only on the shape of the initial semistandard domino tableau. Denote (T^-, T^+) the bi-tableau associated to a semistandard domino tableau

T of shape λ and (λ^-, λ^+) their respective shapes. (T^-, T^+) (resp. (λ^-, λ^+)) is the **2-quotient** of T (resp. λ). Semistandard Young tableaux T^- and T^+ are built by filling each box of T (a domino is composed of two boxes) by a '-' or a '+' sign such that the top leftmost box is filled with '-' and two adjacent boxes have opposite signs. T^- (resp. T^+) is obtained from the sub tableau of T composed of the dominoes with top rightmost box filled with '-' (resp. '+').

Remark 3.11. *If the top leftmost domino of a semistandard domino tableau T is vertical (resp. horizontal), its label is used for T^- (resp. T^+). Therefore not labelling the top leftmost domino with 0 if it is vertical is sufficient to ensure that only T^+ may have entries equal to 0 and any bi-tableau with T^+ containing entries equal to 0 gives a valid semistandard domino tableau according to our constraints. As a result, in our setup, the bijection is actually between semistandard domino tableaux indexed by labels in $\{0, 1, 2, \dots\}$ and pairs of semistandard Young tableaux indexed respectively by $\{1, 2, \dots\}$ and $\{0, 1, 2, \dots\}$.*

Example 3.12. *Figure 5 shows a semistandard domino tableau of weight $\mu = (2, 0, 2, 0, 0, 4, 0, 1)$ and its 2-quotient.*

$$T = \begin{array}{|c|c|c|} \hline -0+ & -0+ & -5+ \\ \hline +2- & -2+ & -5+ \\ \hline - & +5- & + \\ \hline +5- & -7+ & \\ \hline \end{array} \longrightarrow \left(T^- = \begin{array}{|c|c|c|} \hline 5 & 5 & 5 \\ \hline \end{array}, T^+ = \begin{array}{|c|c|} \hline 0 & 0 \\ \hline 2 & 2 \\ \hline 5 & 7 \\ \hline \end{array} \right)$$

Figure 5: A semistandard domino tableau and its 2-quotient.

Denote $X^- = \{x_{-i}\}_{i>0}$ and $X^+ = \{x_i\}_{i\geq 0}$ (note that $X^- = X^+ \setminus \{x_0\}$ as $x_{-i} = x_i$).

Proposition 3.13. *For $\lambda \in \mathcal{P}^0(n)$, the domino function $\mathcal{G}_\lambda$ and the Schur symmetric functions are related through*

$$\mathcal{G}_\lambda(X) = s_{\lambda^-}(X^-)s_{\lambda^+}(X^+). \quad (22)$$

Proof. According to the definition of the 2-quotient above and Remark 3.11, one can get

$$\mathcal{G}_\lambda(X) = \sum_{\substack{\text{shape}(T^-)=\lambda^- \\ \text{shape}(T^+)=\lambda^+}} X^{-T^-} X^{T^+},$$

where the sum is on all pairs of semistandard Young tableau T^- and T^+ of shape λ^- and λ^+ such that no entry of T^- is equal to 0. $\square$

Finally we provide analogues of the Cauchy identities for domino functions.

Proposition 3.14. *The domino functions $\mathcal{G}_\lambda(XY)$ verify the following identities*

$$\begin{aligned}\mathcal{G}_{(2n)}(XY) &= \sum_{\lambda \in \mathcal{P}^0(n)} \mathcal{G}_\lambda(X) \mathcal{G}_\lambda(Y), \\ \mathcal{G}_\lambda(XY) &= \sum_{\mu, \nu \in \mathcal{P}^0(n)} g^B(\lambda, \mu, \nu) \mathcal{G}_\mu(X) \mathcal{G}_\nu(Y).\end{aligned}$$

Proof. Note that

$$(XY)^+ = \{x_i y_j\}_{(i,j) \geq (0,0)} = \{x_0 y_j\}_{j \geq 0} \cup \{x_i y_j\}_{i > 0, j \in \mathbb{Z}} = X^+ Y^+ \cup X^- Y^-.$$

Using Proposition 3.13 we get the first identity.

$$\begin{aligned}\mathcal{G}_{(2n)}(XY) &= s_{(n)}((XY)^+) \\ &= s_{(n)}(X^- Y^- \cup X^+ Y^+) \\ &= \sum_{k=0}^n s_{(k)}(X^- Y^-) s_{(n-k)}(X^+ Y^+) \\ &= \sum_{k=0}^n \sum_{\lambda^- \vdash k} s_{\lambda^-}(X^-) s_{\lambda^-}(Y^-) \sum_{\lambda^+ \vdash n-k} s_{\lambda^+}(X^+) s_{\lambda^+}(Y^+) \\ &= \sum_{\lambda \in \mathcal{P}^0(n)} s_{\lambda^-}(X^-) s_{\lambda^+}(X^+) s_{\lambda^-}(Y^-) s_{\lambda^+}(Y^+) \\ &= \sum_{\lambda \in \mathcal{P}^0(n)} \mathcal{G}_\lambda(X) \mathcal{G}_\lambda(Y).\end{aligned}$$

Using Proposition 3.13 and the theory of symmetric functions on wreath products we prove the second identity. Define as in [1] for any set of indeterminates U and V and any partition λ of n :

$$p_\lambda^+(U, V) = \prod_i [p_{\lambda_i}(U) + p_{\lambda_i}(V)], \quad p_\lambda^-(U, V) = \prod_i [p_{\lambda_i}(U) - p_{\lambda_i}(V)].$$

Note that if $U \cap V = \emptyset$, $p_\lambda^+(U, V) = p_\lambda(U \cup V)$ and if $V \subseteq U$, $p_\lambda^-(U, V) = p_\lambda(U \setminus V)$. Then for partitions $\lambda \in \mathcal{P}^0(n)$ and $\mu \in \mathcal{P}^0(n)$ of 2-quotient (μ^-, μ^+) , denote ψ_μ^λ the common value of character ψ^λ on all signed permutations of *cycle type* (μ^-, μ^+) . According to [15, I, Appendix B],

$$p_{\mu^-}^-(U, V) p_{\mu^+}^+(U, V) = \sum_{\lambda \in \mathcal{P}^0(n)} \psi_\mu^\lambda s_{\lambda^-}(V) s_{\lambda^+}(U).$$

Set $U = X^+$ and $V = X^-$ and use Proposition 3.13 to get for any $\mu \in \mathcal{P}^0(n)$

$$p_{\mu^+}(X) (x_0)^{n-|\mu^+|} = \sum_{\lambda \in \mathcal{P}^0(n)} \psi_\mu^\lambda \mathcal{G}_\lambda(X).$$

But $p_{\mu^+}(XY)(x_0y_0)^{n-|\mu^+|} = p_{\mu^+}(X)(x_0)^{n-|\mu^+|}p_{\mu^+}(Y)(y_0)^{n-|\mu^+|}$. We get

$$\sum_{\lambda \in \mathcal{P}^0(n)} \psi^\lambda \mathcal{G}_\lambda(XY) = \sum_{\mu, \nu \in \mathcal{P}^0(n)} \psi^\mu \psi^\nu \mathcal{G}_\mu(X) \mathcal{G}_\nu(Y).$$

That yields the second identity. $\square$

3.4. Proof of Theorems 3.5 and 3.6

Both Theorem 3.5 and Theorem 3.6 are consequences of Equation (17) and Proposition 3.14.

Proof of Theorem 3.5. Decompose the domino functions $\mathcal{G}_{(2n)}(XY)$ in two ways.

$$\begin{aligned} \mathcal{G}_{(2n)}(XY) &= F_\emptyset^B(XY) = \sum_{I,J} c_{I,J}^\emptyset F_I(X) F_J(Y) \\ \mathcal{G}_{(2n)}(XY) &= \sum_{\lambda \in \mathcal{P}^0(n)} \mathcal{G}_\lambda(X) \mathcal{G}_\lambda(Y) = \sum_{I,J} \sum_{\lambda \vdash n} d_{\lambda I}^B d_{\lambda J}^B F_I^B(X) F_J^B(Y). \quad \square \end{aligned}$$

Proof of Theorem 3.6. Similarly,

$$\begin{aligned} \mathcal{G}_{(2n)}(XYZ) &= F_\emptyset^B(XYZ) = \sum_{I,J,K} c_{I,J,K}^\emptyset F_I^B(X) F_J^B(Y) F_K^B(Z) \\ \mathcal{G}_{(2n)}(XYZ) &= \sum_{\nu \in \mathcal{P}^0(n)} \mathcal{G}_\nu(XY) \mathcal{G}_\nu(Z) \\ &= \sum_{\lambda, \mu, \nu \in \mathcal{P}^0(n)} g^B(\lambda, \mu, \nu) \mathcal{G}_\lambda(X) \mathcal{G}_\mu(Y) \mathcal{G}_\nu(Z) \\ &= \sum_{I,J,K} \sum_{\lambda, \mu, \nu \in \mathcal{P}^0(n)} g^B(\lambda, \mu, \nu) d_{\lambda I}^B d_{\mu J}^B d_{\nu K}^B F_I^B(X) F_J^B(Y) F_K^B(Z). \quad \square \end{aligned}$$

Remark 3.15. Theorem 3.5 is a special case of Theorem 3.6. However, we felt that an independent proof that does not use Kronecker coefficients but only the elementary first identity of Proposition 3.14 was of interest.

3.5. A corollary to Theorem 3.6

For $I \subseteq \{0\} \cup [n-1]$, denote $|I|$ the number of elements in I . The structure constants e_{IJ}^K also count templates M with non-negative integer entries of the following type.

$$M = (m_{i,j}) = \begin{pmatrix} a_{0,0} & a_{0,1} & \dots & a_{0,|I|} \\ & b_{1,1} & \dots & b_{1,|I|} \\ a_{1,0} & a_{1,1} & \dots & a_{1,|I|} \\ \vdots & & & \vdots \\ & b_{|J|,1} & \dots & b_{|J|,|I|} \\ a_{|J|,0} & a_{|J|,1} & \dots & a_{|J|,|I|} \end{pmatrix}$$

Given $q_s = a_{s,0} + \sum_k (a_{s,k} + b_{s,k})$ and $q_0 = a_{0,0} + \sum_k a_{0,k}$ we denote $q(M) = (q_0, q_1, \dots, q_{|I|})$ the **row sums**. The **column sums** $p(M) = (p_0, p_1, \dots, p_{|J|})$ verify $p_j = a_{0,j} + \sum_k (a_{k,s} + b_{k,s})$ and $p_0 = a_{0,0} + \sum_k a_{k,0}$. Successive reading of a -lines from left to right, and b -lines from right to left yields the **reading word**. Denote $n_{I,J}^K$ the number of such templates with row sums I , column sums J and reading word K and $n_{I,J} = \sum_K n_{I,J}^K$. Bergeron and Bergeron in [4] prove that $n_{I,J}^K = e_{I,J}^K$. We get as a corollary to Theorem 3.6:

Corollary 3.16. *Let I, J and $K \subseteq \{0\} \cup [n-1]$. The numbers $n_{I,J}$ and $n_{I,J}^K$ verify*

$$\begin{aligned} n_{I,J} &= \sum_{\lambda} K_{\lambda \text{ comp}(I)}^B K_{\lambda \text{ comp}(J)}^B \\ \sum_{R, D \subseteq \{0\} \cup [n-1]} n_{I,J}^D n_{D,K}^R &= \sum_{\lambda, \mu, \nu \vdash n} g^B(\lambda, \mu, \nu) K_{\lambda \text{ comp}(I)}^B K_{\mu \text{ comp}(J)}^B K_{\nu \text{ comp}(K)}^B. \end{aligned}$$

Proof. Considering relations (16) and (18) in the monomial basis one gets

$$\begin{aligned} \sum_{I,J} \sum_{I' \subseteq I, J' \subseteq J} c_{I',J'}^{\emptyset} M_I^B(X) M_J^B(Y) &= \\ \sum_{I,J} \sum_{\lambda \vdash n} K_{\lambda \text{ comp}(I)}^B K_{\lambda \text{ comp}(J)}^B M_I^B(X) M_J^B(Y), \end{aligned}$$

and

$$\begin{aligned} \sum_{I,J,K} \sum_{I' \subseteq I, J' \subseteq J, K' \subseteq K} c_{I',J',K'}^{\emptyset} M_I^B(X) M_J^B(Y) M_K^B(Z) &= \\ \sum_{I,J,K} \sum_{\lambda, \mu, \nu \vdash n} g^B(\lambda, \mu, \nu) K_{\lambda \text{ comp}(I)}^B K_{\mu \text{ comp}(J)}^B K_{\nu \text{ comp}(K)}^B M_I^B(X) M_J^B(Y) M_K^B(Z). \end{aligned}$$

The first equation of the statement is a consequence of the result of Bergeron and Bergeron and Equation (15).

$$n_{I,J} = \sum_K e_{I,J}^K = \sum_{I' \subseteq I, J' \subseteq J} c_{I',J'}^{\emptyset} = \sum_{\lambda \vdash n} K_{\lambda \text{ comp}(I)}^B K_{\lambda \text{ comp}(J)}^B.$$

To get the second identity compute

$$\begin{aligned} \sum_{R,D} n_{I,J}^D n_{D,K}^R &= \sum_{R,D} e_{I,J}^D e_{D,K}^R \\ &= \sum_R e_{I,J,K}^R \\ &= \sum_{I' \subseteq I, J' \subseteq J, K' \subseteq K} c_{I',J',K'}^{\emptyset} \\ &= \sum_{\lambda, \mu, \nu \vdash n} g^B(\lambda, \mu, \nu) K_{\lambda \text{ comp}(I)}^B K_{\mu \text{ comp}(J)}^B K_{\nu \text{ comp}(K)}^B. \quad \square \end{aligned}$$

4. Skew shape partitions

4.1. Type A

Let n be a non-negative integer and λ and μ , two integer partitions such that the diagram of μ is included in the diagram of λ and $|\lambda| - |\mu| = n$. A **standard (resp. semistandard) skew Young tableau** of shape λ/μ is a Young diagram of skew shape λ/μ whose boxes are filled with the elements of $[n]$ (resp. with positive integers) such that the entries are strictly increasing down the columns and along the rows (resp. strictly increasing down the columns and non decreasing along the rows). The descent set of a standard Young tableau is defined similarly as in the case of classical shapes. The **skew Schur function** $s_{\lambda/\mu}$ is the generating function for semistandard skew Young tableaux of shape λ/μ

$$s_{\lambda/\mu}(X) = \sum_{\text{shape}(T)=\lambda/\mu} X^T,$$

where the sum is on all semistandard skew Young tableaux of shape λ/μ . Given a skew shape λ/μ denote $k_{\mu\nu}^\lambda$ the **Littlewood-Richardson coefficients** i.e. the structure constants of the algebra of symmetric functions in the Schur basis. They also appear in the decomposition of skew Schur functions in terms of non-skew Schur functions

$$s_\lambda s_\mu = \sum_\nu k_{\lambda\mu}^\nu s_\nu, \quad s_{\lambda/\mu} = \sum_\nu k_{\mu\nu}^\lambda s_\nu. \quad (23)$$

Let I and J be two subsets of $[n-1]$. Gessel focuses in [12, Thm. 14] on the number $a_{IJ}^{\lambda/\mu}$ of couples of permutations α, β such that α has descent set I , β has descent set J and $\alpha\beta$ is *compatible* with the skew shape λ/μ (i.e. $\alpha\beta$ is a correct reading word for the skew shape λ/μ). We need some more notions introduced in [12] to define this properly.

Given any partial order P on $[n]$, denote $L(P)$ the set of permutations $\pi \in S_n$, such that the corresponding linear order $\pi(1) < \pi(2) < \dots < \pi(n)$ is a linearisation of P . To obtain a partial order $P_{\lambda/\mu}$ from the skew shape $|\lambda/\mu| = n$ we fill the corresponding skew Young diagram by integers $[n]$ from bottom to top and from left to right and then consider relations $i > k$ if k lay directly below or to the left of i . These relations generate the corresponding partial order $P_{\lambda/\mu}$.

Example 4.1. Let $\lambda/\mu = (4, 4, 3)/(2, 1)$. Figure 6 shows the corresponding partial order. Firstly fill the corresponding skew Young diagram by integers $[7]$ from bottom to top and from left to right and then rotate it by 135 degrees counterclockwise and take its vertical mirror image to get the plane presentation of the ordering $P_{(4,4,3)/(2,1)}$.

Due to Gessel [12, p. 295], $L(P_{\lambda/\mu})$ may be identified with Young tableaux of shape λ/μ . Given $\pi \in L(P_{\lambda/\mu})$ construct the corresponding skew Young tableau by filling the skew shape λ/μ with the letters of the word π^{-1} from bottom to top and from left to right, as in Example 4.2. The permutations from $L(P_{\lambda/\mu})$ are called **compatible** with the skew shape λ/μ .

Figure 6: A partial order $P_{(4,4,3)/(2,1)}$ obtained from the skew shape $(4,4,3)/(2,1)$.

Example 4.2.

Using the bijection above, one can notice that

$$a_{IJ}^{\lambda/\mu} = \sum_{K \subseteq [|\lambda| - |\mu| - 1]} d_{\lambda/\mu K} a_{IJ}^K$$

where $d_{\lambda/\mu K}$ is the number of standard Young tableaux of skew shape λ/μ and descent set K .

Proposition 4.3. *Let λ and μ two integer partitions such that λ/μ is a skew shape and $I, J \subseteq [|\lambda| - |\mu| - 1]$. The coefficients $a_{IJ}^{\lambda/\mu}$ verify the following equality involving the Littlewood-Richardson coefficients, the Kronecker coefficients of the symmetric group and the numbers of standard Young tableaux of given shape and descent*

$$a_{IJ}^{\lambda/\mu} = \sum_{\nu, \rho, \varepsilon} k_{\mu\nu}^{\lambda} g(\nu, \rho, \varepsilon) d_{\rho I} d_{\varepsilon J}.$$

Proof. Gessel showed in [12, Thm. 14] that

$$s_{\lambda/\mu}(XY) = \sum_{I, J} a_{IJ}^{\lambda/\mu} F_I(X) F_J(Y). \quad (24)$$

Use Equation (23) to decompose skew Schur function $s_{\lambda/\mu}(XY)$ using Littlewood-Richardson coefficients. Then use the generalised Cauchy identity (10) and decomposition (4) to prove the proposition.

$$\begin{aligned} s_{\lambda/\mu}(XY) &= \sum_{\nu} k_{\mu\nu}^{\lambda} s_{\nu}(XY) \\ &= \sum_{\nu, \rho, \varepsilon} k_{\mu\nu}^{\lambda} g(\nu, \rho, \varepsilon) s_{\rho}(X) s_{\varepsilon}(Y) \\ &= \sum_{I, J} \sum_{\nu, \rho, \varepsilon} k_{\mu\nu}^{\lambda} g(\nu, \rho, \varepsilon) d_{\rho I} d_{\varepsilon J} F_I(X) F_J(Y). \quad \square \end{aligned}$$

In the following section, we prove that there is a similar formula for the hyperoctahedral group provided some constraints on the skew-shapes.

4.2. Type B

4.2.1. Skew domino tableaux

Let λ and μ be two integer partitions such that λ/μ is a skew shape, $|\lambda| - |\mu| = 2n$ and such that the Young diagrams of shape μ and λ/μ (and by extension λ) may be tiled by horizontal and vertical dominoes. By abuse of notation, we also write $\lambda/\mu \in \mathcal{P}^0(n)$ when all these conditions are fulfilled. A **standard skew domino tableau** is a tiling of the skew Young diagram of shape λ/μ with horizontal and vertical dominoes labelled with integers of $[n]$ such that labels are strictly increasing along the rows and down the columns.

The bijection described in Carre and Leclerc [5] extends to the case of skew shapes. As a result, there is a bijection between standard skew domino tableaux of shape λ/μ and standard skew bi-tableaux of shape $(\lambda^-/\mu^-, \lambda^+/\mu^+)$. As noticed in Section 3.3.3, μ^- and μ^+ depends only on μ . One may apply the algorithm of Carre and Leclerc with an arbitrary tiling of the deleted part of shape μ . Note that the bijection is not descent preserving as in the case of non-skew shapes.

Example 4.4. Decompose the following standard skew domino tableau T .

$$T = \begin{array}{|c|c|c|c|c|c|} \hline - & + & - & + & - & + \\ \hline + & - & + & - & 2 & + \\ \hline - & + & - & 4 & + & - \\ \hline + & - & 1 & + & 6 & - \\ \hline - & + & 1 & + & 6 & - \\ \hline + & 5 & - & & & \\ \hline \end{array} \longrightarrow \left(T^- = \begin{array}{|c|c|c|} \hline 1 & 7 & 9 \\ \hline 5 & & \\ \hline \end{array}, T^+ = \begin{array}{|c|c|c|} \hline 2 & 3 & 8 \\ \hline 4 & & \\ \hline 6 & & \\ \hline \end{array} \right)$$

Given a standard skew domino tableau T , we call in the sequel **negative** (resp. **positive**) **domino**, a domino that goes to T^- (resp. T^+) according to the bijection above. We have the following definition.

Definition 4.5 (Descent set of a standard skew domino tableau). *A standard skew domino tableau T has a descent in position $i > 0$ if $i+1$ lies strictly below i in T and has descent in position 0 if the domino filled with 1 is negative. Denote $\text{Des}(T)$ the set of all the descents of T . For $\lambda/\mu \in \mathcal{P}^0(n)$ and $I \subseteq \{0\} \cup [n-1]$ denote also $d_{\lambda/\mu}^B I$ the number of standard skew domino tableaux of shape λ/μ and descent set I .*

Example 4.6. In Example 4.4 we have $\text{Des}(T) = \{0, 3, 4, 8\}$.

A **semistandard skew domino tableau** is a tiling of the skew Young diagram of shape λ/μ with horizontal and vertical dominoes labelled with integers in $\{0, 1, 2, \dots\}$ such that labels are non decreasing along the rows, strictly increasing down the columns, and negative dominoes cannot be labelled 0. The bijection of Carre and Leclerc [5] also generalises to semistandard domino tableaux and pairs of semistandard Young tableaux such that dominoes labelled 0 may appear only in T^+ .

Example 4.7. Decompose the following semistandard skew domino tableau.

$$T = \begin{array}{|c|c|c|c|c|} \hline - & + & - & + & - \\ \hline + & - & + & - & 0 \\ \hline - & + & - & 3 & + \\ \hline + & 3 & 6 & 7 & 9 \\ \hline - & + & - & + & + \\ \hline + & 6 & - & & \\ \hline \end{array} \longrightarrow \left(T^- = \begin{array}{|c|c|c|} \hline 3 & 7 & 9 \\ \hline 6 & & \\ \hline \end{array}, T^+ = \begin{array}{|c|c|c|} \hline 0 & 0 & 1 \\ \hline 3 & & \\ \hline 6 & & \\ \hline \end{array} \right)$$

The standardisation of a semistandard skew domino tableau is defined in the same way as for non-skew shapes. We provide the following example in lieu of a formal definition.

Example 4.8. The following picture shows a semistandard skew domino tableau T and its standardisation T^{st} .

$$T = \begin{array}{|c|c|c|c|c|} \hline & & & 1 & 8 \\ \hline & & 1 & 1 & \\ \hline & 4 & & & \\ \hline 1 & 6 & 6 & 10 & \\ \hline 6 & & & & \\ \hline \end{array} \longrightarrow T^{st} = \begin{array}{|c|c|c|c|c|} \hline & & & 3 & 8 \\ \hline & & 2 & & \\ \hline & 4 & & & \\ \hline 1 & 6 & 7 & 9 & \\ \hline 5 & & & & \\ \hline \end{array}$$

4.2.2. Skew domino functions

For $\lambda/\mu \in \mathcal{P}^0(n)$ define the **skew domino function** $\mathcal{G}_{\lambda/\mu}$ as the generating function for the semistandard skew domino tableaux of shape λ/μ . Proposition 3.13 generalises well to the case of skew shapes.

Proposition 4.9. For $\lambda/\mu \in \mathcal{P}^0(n)$, the domino function $\mathcal{G}_{\lambda/\mu}$ and the Schur symmetric functions are related through

$$\mathcal{G}_{\lambda/\mu}(X) = s_{\lambda^-/\mu^-}(X^-)s_{\lambda^+/\mu^+}(X^+).$$

Proof. This identity is a direct consequence of the bijection between semistandard skew domino tableaux of shape λ/μ and semistandard skew bi-tableaux of shape $(\lambda^-/\mu^-, \lambda^+/\mu^+)$ and the fact that only positive dominoes may be labelled 0. $\square$

Furthermore, we have the following proposition involving some type B analogue of the Littlewood-Richardson coefficients.

Proposition 4.10. For $\lambda/\mu \in \mathcal{P}^0(n)$, $\mu \in \mathcal{P}^0(m)$ and $\nu \in \mathcal{P}^0(n)$, denote $l_{\mu\nu}^\lambda$ the **type B analogues of the Littlewood-Richardson coefficients** that we define as $l_{\mu\nu}^\lambda = k_{\mu^-\nu^-}^{\lambda^-} k_{\mu^+\nu^+}^{\lambda^+}$. Domino functions verify

$$\mathcal{G}_\mu \mathcal{G}_\nu = \sum_{\lambda \in \mathcal{P}^0(n+m)} l_{\mu\nu}^\lambda \mathcal{G}_\lambda, \quad \mathcal{G}_{\lambda/\mu} = \sum_{\nu \in \mathcal{P}^0(n)} l_{\mu\nu}^\lambda \mathcal{G}_\nu.$$

Proof. Use Propositions 3.13 and 4.9 to show

$$\begin{aligned}
\mathcal{G}_\mu(X)\mathcal{G}_\nu(X) &= s_{\mu^-}(X^-)s_{\mu^+}(X^+)s_{\nu^-}(X^-)s_{\nu^+}(X^+) \\
&= \sum_{\lambda^- \vdash |\mu^-|+|\nu^-|} k_{\mu^- \nu^-}^{\lambda^-} s_{\lambda^-}(X^-) \sum_{\lambda^+ \vdash |\mu^+|+|\nu^+|} k_{\mu^+ \nu^+}^{\lambda^+} s_{\lambda^+}(X^+) \\
&= \sum_{\lambda \in \mathcal{P}^0(n+m)} \left(k_{\mu^- \nu^-}^{\lambda^-} k_{\mu^+ \nu^+}^{\lambda^+} \right) \mathcal{G}_\lambda(X),
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{G}_{\lambda/\mu}(X) &= s_{\lambda^-/\mu^-}(X^-)s_{\lambda^+/\mu^+}(X^+) \\
&= \left(\sum_{\nu^- \vdash |\lambda^-|-|\mu^-|} k_{\mu^- \nu^-}^{\lambda^-} s_{\nu^-}(X^-) \right) \left(\sum_{\nu^+ \vdash |\lambda^+|-|\mu^+|} k_{\mu^+ \nu^+}^{\lambda^+} s_{\nu^+}(X^+) \right) \\
&= \sum_{\nu \in \mathcal{P}^0(n)} \left(k_{\mu^- \nu^-}^{\lambda^-} k_{\mu^+ \nu^+}^{\lambda^+} \right) \mathcal{G}_\nu(X). \quad \square
\end{aligned}$$

In order to get an analogue of Proposition 3.9, the skew shapes have to verify some additional constraints. As noticed in Remark 3.11 for non-skew shapes, the constraints of not labelling the vertical top leftmost domino of a semistandard domino tableau T with 0 and the constraint that no negative domino may be labelled with 0 are equivalent. With skew shapes there is no analogue to this equivalence in the general case and we need to restrict the set of considered shapes. We proceed with the following definitions.

Definition 4.11 (Top domino). *Given a (semi-)standard domino tableau T , we say that domino d is a **top domino** of T if*

- (i) *there is no adjacent domino on the top of d ,*
- (ii) *there is no non-top adjacent domino to the left of d .*

*We further say that a top domino d is **minimum positive** if d is the leftmost downmost positive top domino of T such that there is no adjacent (negative top) domino to the left of d .*

Example 4.12. *In Example 4.4, dominoes labelled with 1, 2, 3 and 8 are top dominoes and 2 is the minimum positive one.*

We look at the following subset of $\mathcal{P}^0(n)$.

Definition 4.13 (Admissible skew shapes). *An **admissible skew shape** is a skew shape $\lambda/\mu \in \mathcal{P}^0(n)$ that cannot be tiled with horizontal and vertical dominoes in a way that*

- (i) *there is a minimum positive domino,*
- (ii) *a negative top domino is placed to the right and above the minimum positive domino.*

Denote $\widetilde{\mathcal{P}}^0(n)$ the set of admissible skew shapes λ/μ such that $|\lambda| - |\mu| = n$.

Example 4.14. The shape of the standard skew domino tableau in Example 4.4 is admissible as well as the following shapes.

These ones are not admissible.

Proposition 4.15. For $\lambda/\mu \in \widetilde{\mathcal{P}}^0(n)$, the skew domino function $\mathcal{G}_{\lambda/\mu}$ verify

$$\mathcal{G}_{\lambda/\mu} = \sum_{I \subseteq \{0\} \cup [n-1]} d_{\lambda/\mu}^B I F_I^B.$$

Proof. The proof is adapted from the one of Proposition 3.9. The first step is to show the analogue of Lemma 3.10, i.e. that for any standard skew domino tableau T_0 of shape λ/μ ,

$$F_{Des(T_0)}^B = \sum_{T^{st}=T_0} X^T.$$

To this end one needs to show that any sequence of relabelling of the dominoes $1, 2, \dots, n$ in T_0 by $i_1 \leq i_2 \leq \dots \leq i_n$ such that $k \in Des(T_0) \Rightarrow i_k < i_{k+1}$ with $k \geq 0$ and $i_0 = 0$ gives a valid semistandard skew domino tableau T such that $T^{st} = T_0$. If the domino labelled 1 in T_0 is positive, i.e. $\min(Des(T_0)) = k > 0$, then any subsequence $0 = i_1 = i_2 = \dots = i_l$ ($l \leq k$) gives a valid semistandard skew domino tableau. Indeed, since λ/μ is admissible, none of the dominoes $2, 3, \dots, k$ are negative. As a result, no negative domino is labelled with 0 (a restriction in our definition of semistandard skew domino tableaux required to get Proposition 4.9). Other cases are similar to the proof of Lemma 3.10 and not detailed here. $\square$

Remark 4.16. The proof of Proposition 4.15 makes it clear that admissibility is required to get at the same time Proposition 4.15 and Proposition 4.9. Indeed the later requires that only positive dominoes of a semistandard skew domino tableau may be labelled with 0. The former assumes that any sequence of dominoes labelled $1, 2, \dots, \min(Des(T_0))$ in a standard skew domino tableau T may be relabelled with 0 to get a valid preimage of T by standardisation. Only admissible shapes fulfil both conditions.

Define for $I, J \subseteq 0 \cup [n-1]$ the coefficients $c_{IJ}^{\lambda/\mu} = \sum_{K \subseteq 0 \cup [n-1]} d_{\lambda/\mu K}^B c_{IJ}^K$. Formula (24) admits the following type B analogue.

Proposition 4.17. *For $\lambda/\mu \in \widetilde{\mathcal{P}}^0(n)$ and $I, J \subseteq 0 \cup [n-1]$, the coefficients $c_{IJ}^{\lambda/\mu}$ verify*

$$\mathcal{G}_{\lambda/\mu}(XY) = \sum_{I, J \subseteq 0 \cup [n-1]} c_{IJ}^{\lambda/\mu} F_I^B(X) F_J^B(Y). \quad (25)$$

Proof. Apply Proposition 4.15 and then decompose Chow's fundamental quasisymmetric function $F_K^B(XY)$ into $F_I^B(X)$ and $F_J^B(Y)$ to prove the statement.

$$\begin{aligned} \mathcal{G}_{\lambda/\mu}(XY) &= \sum_{K \subseteq \{0\} \cup [n-1]} d_{\lambda/\mu K}^B F_K^B(XY) \\ &= \sum_{I, J} \left(\sum_K d_{\lambda/\mu K}^B c_{IJ}^K \right) F_I^B(X) F_J^B(Y) \\ &= \sum_{I, J} c_{IJ}^{\lambda/\mu} F_I^B(X) F_J^B(Y). \quad \square \end{aligned}$$

4.2.3. A type B analogue of Proposition 4.3

We are now ready to state the main result of this section.

Theorem 4.18. *For $\lambda/\mu \in \widetilde{\mathcal{P}}^0(n)$ and $\nu \in \mathcal{P}^0(n)$ denote $l_{\mu\nu}^\lambda$ the type B analogue of the Littlewood-Richardson coefficients defined as in Proposition 4.10 by $l_{\mu\nu}^\lambda = k_{\mu^- \nu^-}^\lambda - k_{\mu^+ \nu^+}^\lambda$. Then, for $I, J \subseteq 0 \cup [n-1]$, the coefficient $c_{IJ}^{\lambda/\mu}$ is given by*

$$c_{IJ}^{\lambda/\mu} = \sum_{\nu, \rho, \varepsilon \in \mathcal{P}^0(n)} l_{\mu\nu}^\lambda g^B(\nu, \rho, \varepsilon) d_{\rho I}^B d_{\varepsilon J}^B. \quad (26)$$

Proof. On the one hand expand $\mathcal{G}_{\lambda/\mu}(XY)$ using Proposition 4.17. On the other hand expand the same skew domino function using Proposition 4.10, Proposition 3.14 and Proposition 3.9. Namely

$$\begin{aligned} \sum_{I, J \subseteq 0 \cup [n-1]} c_{IJ}^{\lambda/\mu} F_I^B(X) F_J^B(Y) &= \mathcal{G}_{\lambda/\mu}(XY) \\ &= \sum_{\nu \in \mathcal{P}^0(n)} l_{\mu\nu}^\lambda \mathcal{G}_\nu(XY) \\ &= \sum_{\nu \in \mathcal{P}^0(n)} l_{\mu\nu}^\lambda \sum_{\rho, \varepsilon \in \mathcal{P}^0(n)} g^B(\nu, \rho, \varepsilon) \mathcal{G}_\rho(X) \mathcal{G}_\varepsilon(Y) \\ &= \sum_{\substack{I, J \subseteq 0 \cup [n-1] \\ \nu, \rho, \varepsilon \in \mathcal{P}^0(n)}} l_{\mu\nu}^\lambda g^B(\nu, \rho, \varepsilon) d_{\rho I}^B d_{\varepsilon J}^B F_I^B(X) F_J^B(Y). \end{aligned}$$

Acknowledgements

This project has been partially supported by the Vernadski scholarship.

References

- [1] R. Adin, C. A. Athanasiadis, S. Elizalde, and Y. Roichman. Character formulas and descents for the hyperoctahedral group. *Advances in Applied Mathematics*, 87:128–169, 2017.
- [2] D. Avella-Alaminosa and E. Vallejo. Kronecker products and the RSK correspondence. *Discrete Mathematics*, 312:1476–1486, 2012.
- [3] D. Barbash and D. Vogan. Primitive ideals and orbital integrals in complex classical groups. *Math. Ann.*, 259:153–199, 1982.
- [4] F. Bergeron and N. Bergeron. A decomposition of the descent algebra of the hyperoctahedral group. *Journal of Algebra*, 148(1):86–97, 1992.
- [5] C. Carré and B. Leclerc. Splitting the square of a Schur function into its symmetric and antisymmetric parts. *Journal of Algebraic Combinatorics*, 4:201–231, 1995.
- [6] R. Carter. Representation theory of the 0-Hecke algebra. *Journal of Algebra*, 104:89–103, 1986.
- [7] C. Chow. *Non commutative symmetric functions of type B*. PhD thesis, MIT, 2001.
- [8] D. Garfinkle. On the classification of primitive ideals for complex classical Lie algebras, I. *Compositio Mathematica*, 75(2):135–169, 1990.
- [9] D. Garfinkle. On the classification of primitive ideals for complex classical Lie algebras, II. *Compositio Mathematica*, 81(3):307–336, 1992.
- [10] D. Garfinkle. On the classification of primitive ideals for complex classical Lie algebras, III. *Compositio Mathematica*, 88(2):187–234, 1993.
- [11] A. Garsia and C. Reutenauer. A decomposition of Solomon’s descent algebra. *Advances in mathematics*, 77:189–262, 1989.
- [12] I. Gessel. Multipartite P-partitions and inner products of skew Schur functions. *Contemporary Mathematics*, 34:289–317, 1984.
- [13] A. Kirillov, A. Lascoux, B. Leclerc, and J. Thibon. Séries génératrices pour les tableaux de dominos. *C.R. Acad. Sci. Paris*, 318(I):395–400, 1994.
- [14] D. Knuth. Permutations, matrices and generalized young tableaux. *Pacific J. Math.*, 34:709–727, 1970.

- [15] I. Macdonald. *Symmetric functions and Hall polynomials*. Oxford University Press, 1999.
- [16] P. Norton. 0-Hecke algebras. *J. Austral. Math. Soc. Ser A*, 27:337–357, 1979.
- [17] T. K. Petersen. A note on three types of quasisymmetric functions. *The Electronic Journal of Combinatorics*, 12:R61, 2005.
- [18] S. Poirier. Cycle type and descent set in wreath products. *Discrete Math.*, 180:315–343, 1998.
- [19] L. Solomon. A Mackey formula in the group ring of a Coxeter group. *Journal of Algebra*, 41:255–264, 1976.
- [20] R. Stanley. *Enumerative combinatorics*, volume 2. Cambridge University Press, 2001.
- [21] D. Stanton and D. White. A Schensted algorithm for rim hook tableaux. *Journal of Combinatorial Theory, Series A*, 40:211–247, 1985.
- [22] M. Taşkin. Plactic relations for r-domino tableaux. *The Electronic Journal of Combinatorics*, 19:P38, 2012.
- [23] M. A. A. van Leeuwen. The Robinson-Schensted and Schützenberger algorithms, an elementary approach. *The Electronic Journal of Combinatorics*, 3(2):R15, 1996.