Functional traits modulate plant community responses to alien plant invasion

Guillaume Fried^{1*}, Marta Carboni^{2,3}, Lucie Mahaut⁴, Cyrille Violle³

*corresponding author, guillaume.fried@anses.fr

¹ Anses, Laboratoire de la Santé des Végétaux, Unité Entomologie et Plantes Invasives, 34988 Montferrier-sur-Lez, France.

² Department of Biological Sciences, University of Toronto-Scarborough, 265 Military Trail Toronto, ON, M1C 1A4, Canada

³ Department of Sciences, Università Roma TRE, V.le Marconi 446, 00146 Roma, Italy

⁴ Centre d'Ecologie Fonctionnelle et Evolutive, UMR 7372 CNRS – Université de Montpellier – Université Paul Valéry – EPHE, 34293 Montpellier, France.

Declarations of interest: none

Abstract (399 words)

It is well known that alien plant invasion modifies the composition and diversity of resident plant communities, yet our ability to predict patterns of vegetation responses to invasion is hampered by a poor understanding of which functional traits make some resident plants more or less vulnerable to invader impacts. For example, resident species may be more likely to persist and coexist with the invader if they display different strategies of resource use (i.e. high niche differences) or if they share similar highly competitive traits (i.e. competitive hierarchy). Here, we used a trait-based approach to 1) specifically test whether species changes in abundance following plant invasion is random or depends on specific functional trait values, 2) identify which functional strategies allow resident species to coexist with the invader and 3) assess to what extent communities where the invader has been removed return to a reference state. To do so, we applied a semi-experimental approach with the annual vine *Humulus japonicus*, an invasive species in riparian habitats of Southern France. We monitored plant communities seven times over two years in non-invaded reference plots, invaded plots and removal plots in which the invader had been removed at the seedling stage. We quantified species richness and functional richness as well as mean community trait values of seven traits. The comparison of invaded and removal plots highlighted a strong impact of *Humulus japonicus* on both species and functional richness of riparian communities. Resident species that had a distinct flowering onset from the invader faced less pronounced declines. Specifically, species that flowered before the invader reached high cover, and species that had a short flowering duration had lower risk of declining. In addition, species coexisting with the invader when it reached its maximum cover displayed high stature and high seed mass. These results suggest that both *niche differentiation* (in terms of flowering strategy) and *competitive hierarchies* (in terms of height and regeneration strategy) play a role in explaining plant community responses to plant invasion. Finally, while species richness recovered rapidly in

removal plots, functional richness remained lower than in non-invaded plots, pointing to a state-shift that would otherwise go undetected. Overall, our study highlights that the analysis of the functional traits of both invaders and resident species, combined with regular monitoring over time of non-invaded reference plots, invaded and removal plots can greatly improve our understanding of the impact of plant invasion on resident communities.

Keywords: flowering phenology; functional traits; impact; invasive plants; seed mass; temporal niche

Introduction

Research in invasion ecology has gradually shifted from explaining the success of invasive species' to understanding their effects on invaded communities and ecosystems (Vilà et al., 2011). Studies are increasingly finding evidence of strong impacts of invasive plants on plant communities, including species loss and changes in taxonomical composition (Fried et al., 2014; Hejda et al., 2009). Further advances was rendered possible by using functional traits (Castro-Díez et al., 2016; Hejda and Bello, 2013; Jucker et al., 2013), showing that impacts of alien plants are not random and that certain characteristics make some resident species more vulnerable than others (Hejda, 2013; Hejda et al., 2017). However, these studies did not take into account simultaneously the traits of the invasive species and the traits of the resident species in light of the coexistence theory (MacDougall et al., 2009). Here, we contribute to answering one of the basic questions associated with the impacts of invasive species: is it more advantageous for native species to possess characteristics contrasting to the invader (and therefore escape the invader's competition) or to possess similar characteristics and compete with the invader?

Trait differences between species are thought to influence biotic interactions and ultimately determine which species can coexist locally (i.e. "biotic filtering") (Weiher et al., 2011). Specifically, according to modern coexistence theory, both *niche differences* and *fitness differences* influence competitive outcomes (Chesson, 2000; Kraft et al., 2015). On the one hand, under strong biotic filtering by a dominant invasive alien plant, resident species may persist and coexist with the invader if they possess different traits, reflecting different strategies of resource use (i.e. high niche differences) (Figure 1A, on top). In the same way, resident species with similar traits will be more directly impacted, and as a result of their competitive exclusion, the average trait values of resident species in invaded communities should shift away from the traits of the invader (Figure 1A, at the bottom). On the other hand, species may persist

together with the invader if they have similar fitness and equivalent competitive effects (Figure 1B, on top), i.e. if they share similar highly competitive traits for resource use (Gallien et al., 2015; Mayfield and Levine, 2010). In this case, it is the species with less competitive traits that will be excluded (e.g. shorter species will be excluded by a taller invader that out-shades them) (Violle et al., 2009). Consequently, if only similarly competitive species (e.g., with similar plant height) coexist with the invader, the average trait values of resident species in the invaded communities will become closer to those of the invader (Figure 1B, at the bottom). Thus, identifying which resident species have persisted and which have been displaced according to their trait values and their differences with those of the invader can unravel the mechanisms underlying coexistence with the invader and the impacts observed (Gallien and Carboni, 2017).

From what precedes, the functional structure of the community (i.e., the composition and distribution of functional traits in the community) is expected to be altered as a consequence of plant invasion (Castro-Díez et al., 2016). The impact of biological invasion on the functional structure can be assessed by measuring two major components, which are the range (diversity) and the mean of trait values. First, the functional richness (FRic) is the amount of functional space filled by the community, an equivalent to trait range but in a multidimensional space. Hence, changes in FRic between non-invaded and invaded communities can inform us on whether species loss is associated with a decrease in the range of trait values (Villéger et al., 2008). Second, the community weighted mean (CWM) of a trait is the mean value of that trait across co-occurring species weighted by their abundance (Garnier et al., 2004). The analysis of CWMs of invaded communities is informative in two complementary ways. On the one hand, comparing the CWMs of non-invaded communities with those of the persisting species in invaded communities (i.e. excluding the invader) can help identifying which are the viable strategies for coexistence with the invader. On the other hand, including the invader in the calculation of the CWMs of invaded communities (i.e. the overall final species assemblage) can

102 reveal potential changes in ecosystem functions provided by the community in the post-invasion
103 state (Garnier et al., 2016).

104 So far, most studies quantifying the impacts of invasive species have relied on
105 comparing paired plots of invaded and non-invaded communities (e.g. Castro-Díez et al.
106 (2016); Fried et al. (2014); Jucker et al. (2013)). A much smaller number of studies, have also
107 experimentally removed the invader from the surveyed plots (e.g. Hejda (2012); Hulme and
108 Bremner (2006)). First, removal experiments provide a more direct and less biased measure of
109 impact because removal and invaded paired plots only differ by the presence-absence of the
110 invader, provided the invader is removed at an early stage (Kumschick et al., 2014). Second,
111 when coupled with regular surveys in time, removal plots allow following how communities
112 change after invasion, once the invader is controlled, and whether or not they recover towards
113 a similar state to that of uninvaded reference communities.

114 Here we apply a trait-based approach in order to address the following questions: 1)
115 what is the impact of an invasive plant on the species richness and functional richness of
116 resident communities? 2) Are certain species with particular traits more affected by invasion
117 than others? Does this depend on trait distance to the invader or on trait hierarchies? 3) To what
118 extent communities where the invader has been removed return to a reference state in terms of
119 species and functional richness, and in terms of dominant traits (community weighted means of
120 traits)? To answer these questions, we used a semi-experimental approach with the invasive
121 annual vine *Humulus japonicus* as a model species. Three kinds of plots were compared: 1)
122 invaded plots with high density of the invader, 2) removal plots in invaded sites where the
123 annual vine seedlings were removed early in the season, and 3) still non-invaded plots which
124 are considered as representing the reference state of the habitat before invasion. Comparing
125 these treatments can inform about the impact of the invader (invaded *versus* removal plots) but
126 also about the resilience of restored communities (removal *versus* non-invaded plots).

127

128 **Material and methods**

129 *Study area and habitat*

130 Our experiment was set up along the Gardon River, a tributary of the Rhône, in the
131 Mediterranean region of France (between 44°02'00.16"N, 4°07'53.35"E 43°59'11.93"N,
132 4°12'20.03"E). The Gardon River has a typical Mediterranean-type flow regime characterized
133 by a low mean annual discharge (33 m³/s) with higher flows during winter, extreme seasonal
134 variations, and rapid fluctuations in response to intense rainfall with flood peaks around 100
135 times greater than the mean discharge, usually in autumn (September, October) but occasionally
136 in spring (April) (Dezileau et al., 2014). Riparian vegetation of Mediterranean rivers such as
137 the Gardon River consists in a succession of different communities from herbaceous pioneer
138 communities dominated by *Persicaria lapathifolia*, *Veronica anagallis-aquatica* and *Bidens*
139 *frondosa* (*Bidention* communities) to riparian forests dominated by *Populus alba*. A previous
140 study showed that *H. japonicus* has its optimum in the *Bidention* communities in eutrophic soils
141 and full sun exposure (Fried et al., 2018).

142

143 *Experimental design*

144 We used a hierarchical sampling design with 60 plots nested into four sites: Vézénobres-1
145 (44°02'00.16"N, 4°07'53.35"E), Vézénobres-2 (44°01'56.46"N, 4°08'32.20"E), Ners
146 (44°01'24.77"N, 4°10'02.86"E), Brignon (43°59'11.93"N, 4°12'20.03"E). Invaded plots were
147 chosen at the invasion front downstream of the river (Ners, Brignon) and non-invaded plots
148 were chosen beyond the invasion front upstream of the river (Vézénobres-1, Vézénobres-2).
149 Plots within sites were located 3–500 meters apart from each other. The four sites were located
150 1–10 km apart.

151 In early March 2014, 40 experimental plots were set up at the invasion front in two of
152 the above mentioned sites (Brignon, Ners) where *H. japonicus* was identified as being
153 dominant, based on dense stands of *H. japonicus* seedlings and observations of *H. japonicus*
154 dominance during previous years. In each site, we randomly selected 10 small areas where two
155 paired-1m² permanent subplots were established. The paired subplots were distant 0.5–1.5 m
156 apart from each other, while different pairs were separated by a distance of 2–25 m. Due to the
157 spatial proximity, site conditions are very similar within a pair. Within each pair, we randomly
158 allocated the two treatments: one of the two subplots was left unmanipulated, i.e. invaded by
159 *H. japonicus* (Invaded plots), while in the second one, we removed all seedlings of *H. japonicus*
160 (Removal plots). We assume that removal of small seedlings (~1 cm height in March 2014 and
161 March 2015) of an annual plants resulted in minimal disturbances in the Removal plots
162 (Kumschick et al., 2014). In order to avoid shade effect from adjacent *H. japonicus*, the removal
163 was also applied in a buffer zone of 0.25 m. There were only few later emergences of *H.*
164 *japonicus* (only in April 2014 and April 2015), but we repeatedly had to cut stems of *H.*
165 *japonicus* coming from adjacent areas due to the lateral stem growth of the vine on the ground.

166 Additionally, 20 plots were randomly chosen within two other sites upstream of the river
167 (Vézénobres-1, Vézénobres-2) beyond the invasion front where we presumed that *H. japonicus*
168 has never occurred yet (all presences of *H. japonicus* were downstream of these two sites).
169 Choosing non-invaded reference plots in the same sites where the invader was dominant was
170 not possible as only very few uninvaded areas were left. Therefore, reference sites were chosen
171 based on similar position to the river (distance, topography), similar soil type and similar
172 species pool (vegetation dominated by *Bidens* spp.) compared to the sites used for Invaded and
173 Removal plots. In all plots in all sites we visually estimated cover (to the nearest 5%) of all
174 species at repeated intervals, four times in 2014 (March, April, May, July) and three times in
175 2015 (April, June, August).

In order to characterize the invasive species and the resident communities, we used seven traits from different sources: Raunkiaer's life forms (annuals, biennials with rosettes, erected perennials, perennials with stolons, perennials with rhizoms or tubers (geophytes), tree seedlings, and vines), flowering onset (month), flowering duration (month) (Julve, 1998), maximum plant height (m) (Tison and De Foucault, 2014), seed mass (g), specific leaf area (SLA, $\text{mm}^2.\text{mg}^{-1}$) and leaf dry matter content (LDMC, mg.g^{-1}) (Kleyer et al., 2008). We chose this set of traits because they capture differences in regeneration and acquisition strategies as well as the temporal use and competitive ability for resources in riparian habitats (Garnier et al., 2016).

*Assessing impact of *H. japonicus* on species and functional richness of resident communities*

First, we quantified how *H. japonicus* affected species and functional richness of resident communities. Preliminarily, in order to estimate and compare total species richness (γ richness) in the three treatments (considering all the 20 plots of each treatment), species' accumulation curves were used with the second-order jack-knife method to avoid a negatively biased estimate of species richness (Colwell and Coddington, 1994). Then, at the plot scale, the impact of *H. japonicus* on resident communities' diversity was assessed based on species richness (S) and functional richness (FRic). FRic is the amount of functional space filled by the community, an analogue of trait range in a multidimensional space (Villéger et al., 2008). Species richness (S) and functional richness (FRic) across invasion treatments were analyzed using mixed effect models to account for the nested sampling design and the repeated sampling in time in the same plots. Subplot identity (nested within Plots, with Plots nested within Sites) was considered as a random factor on the intercept. Site ("Ners", "Brignon", "Vézénobres-1", "Vézénobres-2"), Treatment ("Invaded", "Removal" and "Non-invaded"), Year (2014 and 2015) and Season, i.e. sampling dates of the consecutive censuses throughout one growing season, and their

interactions were included as fixed factors. For species richness S , we used a GLMM with a log link function and we assumed a Poisson distribution error. For FRic, we used a LMM and we assumed a Gaussian distribution error. Plant height and seed mass were log-transformed before the analyses. Starting from the maximum model, we created simplified models and tested with χ^2 the growth of residual deviance associated to the omission of each term or interactions (deletion tests). Given that riverbeds in the study area are strongly colonized by a number of additional alien plants (Chytrý et al., 2008; Fried et al., 2018), models were fitted also using only native species richness (S_{native}) and native functional richness ($\text{FRic}_{\text{native}}$) to assess if the patterns observed for native species follow the same trend as those observed for the whole community. Results were comparable (Appendix A), and in the following, except where noted, we report only the trends based on the entire community. We also developed another series of complementary mixed effect models in order to assess separately the effect of Treatment on S and FRic at the beginning (March 2014) and at the end of the experiment (August 2015) (see Appendix B).

These analyses were carried out in the R statistical framework (R Development Core Team, 2008) using package lme4, and post-hoc tests were performed with function lsmeans of package emmeans. Residuals were visually inspected to detect trends that could bias estimates but all assumptions of (G)LMMs were met.

Assessing which species and characteristics are most vulnerable

Second, in order to identify the response of individual species to invasion by *H. japonicus*, and whether this response can be explained by trait distances (*niche differentiation*) or trait hierarchies, we computed species abundance (the average cover a species occupied in the plots, A) and species frequency (the number of plots in which a species occurred, Fr) of each species in the Invaded and in the Removal plots. Here we focused on the Invaded-Removal paired plots

as they only differ by the presence-absence of the invader and represent therefore the most straightforward and unbiased method to measure the effect of the invader (Kumschick et al., 2014).

Preliminarily, to assess the overall changes on resident species abundance and frequency due to the presence of the invader, we compared the average abundance and frequency across all species (i.e., species is the unit of repetition) in Invaded and Removal plots (considering all sampling dates together) using a pairwise Wilcoxon test. Then, for each species, we calculated the Relative Impact (RI) of *H. japonicus* on species frequency and on species abundance, with $RI_x = \frac{x_{Rem} - x_{Inv}}{x_{Rem} + x_{Inv}}$, where x is the variable of interest (frequency or abundance), *Inv* represents the Invaded plots and *Rem* represents the Removal plots (Vila et al., 2006). RI_x ranges between -1 (species only in Invaded plots) and 1 (species only in Removal plots). When RI_x is computed by combining all the sampling dates (early spring to late summer, along an increasing cover gradient of *H. japonicus*), this index can detect species that can avoid the maximum development of *H. japonicus* with an early life-cycle (i.e., species that would be equally present in the early censuses of Invaded and Removal plots but absent in late censuses of both types of plots). When computed for the last censuses of the experiment (July 2014 and August 2015, which correspond to the maximum development of *H. japonicus* cover), RI_x will highlight species that are able to coexist with dense stands of the invader.

For both temporal scales (all censuses and last censuses), we tested whether the relative impacts of *H. japonicus* on the frequency (RI_{Fr}) and on the abundance (RI_A) of each species were related to their functional distance to *H. japonicus* or to trait hierarchy. For this purpose, we compared two models. In the first, under the niche differentiation hypothesis (Fig. 1A), the explanatory variables consisted for each species in the *absolute* difference of their trait values with *H. japonicus* (i.e., the distance to *H. japonicus*, always positive). In the second model, under the competitive hierarchy hypothesis (Fig. 1B), the explanatory variables consisted for

each species in the *relative* difference of their trait value with *H. japonicus* (e.g., a species that is shorter than *H. japonicus* will have a negative value for height difference). We used linear regressions with RI_{Fr} and RI_A of each species as the dependant variables and Raunkiær's life forms and absolute or relative distance of trait values between *H. japonicus* and each species as explanatory variables. Only the species occurring in both Removal and Invaded plots were considered in these analyses (discarded species represented less than 1% of species occurrences).

Assessing the impact of H. japonicus on the functional structure of resident communities

The impact on the functional structure was assessed by comparing $FRic$ (see above) and CWMs in the three treatments. Because we were interested in the impact of *Humulus japonicus* on the resident plant communities, we first computed CWMs excluding *Humulus japonicus* to account for shifts of the community *per se* (Thomsen et al., 2016). Second, we computed CWMs including *H. japonicus* to account for potential overall shifts in ecosystem functioning (Thomsen et al., 2016). We compared the CWMs (excluding *H. japonicus*) across treatments at the end of the season when the invader reached its maximum cover (July 2014 and August 2015). For this purpose, LMM and deletion tests were used with CWMs as the response variable, Treatment, Site and their interaction as fixed factors and subplot identity as a random factor on the intercept. We expect that CWMs of certain traits in Invaded plots will differ from CWMs in Removal plots and Non-Invaded plots, with values of traits in Invaded plots reflecting viable strategies in invaded sites dominated by *H. japonicus*. Finally, in order to assess to what extent the dominance of *H. japonicus* shift the mean traits of the communities, we repeated the same analysis including *H. japonicus* in the calculation of the CWMs.

Results

Impact on species and functional richness

A total of 179 species were recorded during the two years of the study. Species accumulation curves showed the presence of an asymptote indicating that the sampling was sufficient to capture the majority of the species pool in the three treatments (Appendix C). Estimates of maximum species richness showed that Non-Invaded plots had the greatest richness (154-160), followed by Removal plots (146-154) and Invaded plots (87-94). The functional richness of the cumulated species pool showed less differences between Non-Invaded plots (0.156), Removal plots (0.155) and Invaded plots (0.151).

In March 2014, at the beginning of the experiment (before removal), species richness and functional richness of Invaded and Removal paired plots were strongly correlated ($r=0.641$, $P<0.001$, and $r=0.698$, $P<0.001$, respectively), suggesting that there was no bias due to a potential early effect of *H. japonicus* seedlings (see Appendix D). For all 6 subsequent sampling dates there were no more significant correlations between pairs.

For the full model (over all the censuses), the marginal r^2 of the GLMM for species richness was 0.773 (while the conditional r^2 was 0.783) indicating that most of the variation of species richness was explained by the fixed factors. Overall, the number of species per plot was significantly different across treatments, across sites, between the two sampling years and there was also a strong effect of season (Table 1). There was no significant interactions between treatment and sites indicating no differences in the effects of invasion in different sites (Table 1). There were, however, significant interactions between treatments and season, revealing a different trend across treatments throughout the growing season, and significant interactions between treatments and year reflecting different trends in the two years (Table 1). Similar trends were found with functional richness (Table 1).

Species richness first increased in plots of all treatments at the beginning of the season (from March 2014 to April 2014) before a maximum value of species richness was reached in

301 April for both years with a lower level reached in Invaded plots (Table 2, Fig. 2). Then a
302 continuous decline was observed in all plots until the end of the summer with lowest values
303 reached in August 2015 and a sharper decline in Invaded plots (Table 2, Fig. 2). In parallel, the
304 total cover of *H. japonicus* increased throughout the growing season reaching 100% cover in
305 all Invaded plots in August 2015 (Table 2).

306 At the beginning of the experiment (March 2014), species richness varied according to
307 sites ($\chi^2=33.05$ $P<0.001$) and to treatments ($\chi^2=8.55$, $P=0.014$). Species richness was higher in
308 Non-Invaded plots compared to Removal (post-hoc tests, $z=2.80$, $P=0.014$) and Invaded
309 ($z=2.35$, $P=0.049$) plots, while there were no differences between Removal and Invaded plots
310 ($z=0.46$, $P=0.890$, Appendix B). Functional richness differed according to sites (LMM,
311 $\chi^2=14.58$, $P<0.001$) but not according to treatments (LMM, $\chi^2=2.81$, $P=0.246$) (Appendix B).

312 At the end of the experiment (August 2015), species richness varied according to
313 treatments (GLMM, $\chi^2=13.49$, $P<0.001$) but not anymore according to sites (GLMM, $\chi^2=3.73$,
314 $P=0.154$). There were less species in Invaded plots compared to both Removal plots (Tukey
315 Post-hoc test, $z=-6.67$, $P<0.001$) and Non-invaded plots ($z=-7.18$, $P<0.001$) while there were
316 no more differences between Removal plots and Non-invaded plots ($z=2.04$, $P=0.104$,
317 Appendix B). Similarly, functional richness differed among treatments (LMM, $\chi^2=58.60$,
318 $P<0.001$) but not across sites anymore (LMM, $\chi^2=2.02$, $P=0.364$). Functional richness was
319 lower in Invaded plots compared to both Removal plots (Tukey Post-hoc test, $z=-7.45$,
320 $P<0.001$) and Non-invaded plots ($z=-9.96$, $P<0.001$) while there were no more differences
321 between Removal plots and Non-invaded plots ($z=2.56$, $P=0.102$, Appendix B). When
322 accounting only for native species in the resident communities, similar results were found for
323 species richness (S_{native}), however native functional richness ($\text{FRic}_{\text{native}}$) remained lower in
324 Removal plots compared to Non-Invaded plots (post hoc test, $z=3.67$, $P=0.028$).

325

Magnitude of species decrease related to trait distances with the invader

H. japonicus reduced average species frequency of occurrence from 8.06% to 3.66% (Wilcoxon paired sign rank test, $V = 9262.5$, $P < 0.001$) and species mean cover from 2.26% to 1.12% (Wilcoxon paired sign rank test, $V = 6677.5$, $P < 0.001$). However the relative ranking of species remained the same for species frequency (Fig. 3a, Pearson's product-moment correlation, $t=30.40$, $r=0.916$, $P < 0.001$) and abundance (Fig. 3b, $t=11.55$, $r=0.706$, $P < 0.001$). Among the native species occurring in both Removal and Invaded-Plots, *Persicaria lapathifolia* showed the strongest decline in frequency (-42%) while *Ficaria verna* or *Alliaria petiolata* were unaffected (species positioned on the 1:1 line on Fig. 3a). *Lythrum salicaria* (-5.4%), *Agrostis stolonifera* (-5.1%) and *Galium aparine* (-4.5%) were the native with the strongest decline in cover abundance while *Phalaris arundinacea* was not affected (species positioned on the 1:1 line on Fig. 3b).

When we considered the *absolute* distance of traits between each species and *H. japonicus* (*niche differentiation hypothesis*), linear regressions showed that relative impact of *H. japonicus* on frequency of resident species was negatively correlated with the functional distance of flowering onset and positively correlated with the functional distance of flowering duration (Table 3, Fig. 4a), i.e. resident species displaying the same flowering onset as *H. japonicus* were more impacted as well as resident species with different flowering duration. The functional distances of all other traits between resident species and *Humulus japonicus* were not significantly related to the degree of impact (Table 3, Appendix E). When we considered the *relative* distance of traits between each species and *H. japonicus* (*trait hierarchy hypothesis*), the results were the same with a positive correlation between the magnitude of impact of *H. japonicus* and the relative distance to flowering onset and flowering duration (Table 3, Fig. 4b) meaning that early flowering species and species with short flowering

duration were less impacted than late flowering species and species with long flowering duration.

At the end of the season, the magnitude of impact of *H. japonicus* on native species frequency was negatively correlated with their *relative* distance in seed mass (i.e. species with larger seeds than *H. japonicus* were less impacted, Fig. 4c) while no relationships was found with absolute distance of traits (Table 3, Appendix E). There were no significant relationships between the impact of *H. japonicus* on abundance of resident species and the absolute or relative functional distances (Appendix F).

Impact on functional structure

Beside the strong shrinkage in trait range revealed by the decrease in FRic, *Humulus japonicus* also shifted the CWMs of invaded communities. At the end of the season of each year (July 2014 and August 2015), the persisting species in invaded communities differed from both Removal and Non-Invaded communities, in particular by having a higher CWM for seed mass ($\chi^2=17.4$, $P<0.001$, Fig. 5). They also differed from Removal plots by a later flowering onset (post-hoc test, $t=2.52$, $P=0.041$), a shorter flowering duration (post-hoc test, $t=-3.36$, $P=0.006$), a higher proportion of perennials with rhizomes (post-hoc test, $t=2.68$, $P=0.029$), a higher proportion of vines (post-hoc test, $t=-2.80$, $P=0.021$) and a lower proportion of annuals (post-hoc test, $t=2.61$, $P=0.034$, Fig. 5).

In contrast, the overall community CWMs (i.e. including trait values of *H. japonicus*) in invaded plots differed from Removal plots by having higher values for plant height (post-hoc test, $t=5.23$, $P<0.001$), LDMC (post-hoc test, $t=2.96$, $P=0.016$), and lower values for SLA (post-hoc test, $t=-2.97$, $P=0.013$) and flowering duration (post-hoc test, $t=-6.34$, $P<0.001$, Fig 6). In addition, invaded communities displayed a lower proportion of perennials with rhizomes or tubers ($\chi^2=13.2$, $P=0.001$), annuals ($\chi^2=8.9$, $P=0.011$), and a higher proportion of annual

vines ($\chi^2=192.2$, $P<0.001$) compared to Non-Invaded and Removal plots. The proportion of biennials was also lower compared to Removal plots (post-hoc test, $t=-3.28$, $P=0.006$) while the proportion of stoloniferous perennials was lower than in Non-Invaded plots (post-hoc test, $t=-3.88$, $P=0.023$). Finally, Removal and Invaded plots had a lower proportion of tree seedlings compared to Non-Invaded plots ($\chi^2=11.0$, $P<0.004$, Fig. 6).

Discussion

The aim of this study was to improve our understanding of alien species impacts on resident plant communities through a functional lens. Using paired plots (removal versus invaded) as a basis of comparison, we showed a strong impact of *Humulus japonicus* on riparian vegetation communities both taxonomically and functionally. At the landscape scale (γ -diversity) and across all the seven censuses, species richness was reduced by 60% while functional richness only decreased by 3% meaning that species remaining in invaded plots covered most of the functional space of removal plot communities. However, at the plot scale (α -diversity), a stronger decrease, between 77% (July 2014) and 97% (August 2015), was observed for species richness, as well as for functional richness, between 69 % (July 2014) and 100% (August 2015). This is a drastic impact, which can be explained by the high competitive exclusion ability of *Humulus japonicus* (Balogh and Dancza, 2008).

The trait values of species able to persist in invaded sites differed from the trait values of species that decrease in presence of the invader. This is remarkable for the traits: flowering onset, flowering duration, seed mass and life form. Perennial species with underground storage organs (rhizoms, tubers) were impacted less than species with an annual life cycle. Using a contrasting life strategy to the annual target invader appears to be an advantage for persisting in invaded sites. A similar result was described by Hejda (2013) who observed that annuals were proportionally more abundant in vegetation invaded by clonal, perennial polycarpic alien

plants. Contrary to this, annual vine, i.e. the same life form than the invader, was the second most represented life form strategy in the invaded communities. In this case, it is possible that the resident species using the same strategy than the invader have similar or higher fitness and competitive effects. Besides life forms, the comparison of trait values between persistent and declining species provides arguments in favour of both our hypotheses: niche differentiation- and competitive hierarchies-hypotheses.

Niche differentiation

The decrease in species occurrences in invaded communities was not random, with a strong signal according to the timing of flowering onset of resident species and its overlap with flowering onset of *Humulus japonicus*. Species with a similar flowering onset as *Humulus japonicus* were much more impacted compared to species with an earlier life cycle. This suggests that some species can coexist with *Humulus japonicus* in invaded sites if they are able to use resources before the development of the invader, as is the case for early flowering species (e.g. *Alliaria petiolata*, *Cardamine hirsuta*, *Cerastium glomeratum*). These species ended their life cycle before *H. japonicus* reached its full cover. In contrast, species whose flowering onset is similar to that of *H. japonicus* and that share a similar life cycle (spring germination, summer flowering and autumn fruiting) were the most heavily impacted (e.g. *Chenopodium album*, *Persicaria lapathifolia*). This result is in line with the hypothesis that native species need to have a certain degree of niche differentiation to be able to coexist with a strong invader and persist in invaded communities (Shea and Chesson, 2002). It also stresses the importance of traits related to plant phenology: it seems that flowering onset captured well the timing of species occupation of space and the timing of resources use, thus defining species temporal niches (Craine et al., 2012). Apart from competition for space, another hypothesis would be that a similar flowering phenology to the invader exposes resident species to increased competition for pollinator visitation and may lead to reduced seed output and population size

(Wolkovich and Cleland, 2011). This hypothesis is however unlikely in the case of *H. japonicus* that is largely pollinated by wind.

Competitive hierarchy

Our study provided evidence that the magnitude of impact on resident species is also associated to trait hierarchies, with less competitive species being more vulnerable. In particular, species that likely had more competitive seedlings than *H. japonicus* because of heavier seed mass, such as *Sicyos angulatus* or *Sambucus ebulus*, were also less impacted by dense stands of *H. japonicus*. Heavier seeds may also allow for longer distance dispersal for some species that would be able to co-occur with the invader each year if seeds are replaced in the seed bank from nearby non-invaded populations. However, three other traits known to be related to competitiveness (SLA, growth form and plant height) were also associated with the ability to co-exist with the invader. Although non-significant, we found a similar trend for specific leaf area (SLA), showing that species with faster acquisition ability (higher SLA), such as *Artemisia verlotiorum*, *Galium aparine* or *Helianthus tuberosus*, were less impacted by *H. japonicus*. The second approach based on community weighted means showed results consistent with these patterns, with species able to coexist with high cover of *H. japonicus* at the end of the season being mostly tall perennial species, with heavier seeds and late and short flowering times and able to reproduce vegetatively through rhizomes (e.g., *Artemisia verlotiorum*) or tubers (e.g., *Helianthus tuberosus*). This combination of trait values represents a different strategy than that of *H. japonicus* but resulted in almost similar fitness. In addition, the fact that only species that were taller, capable of lateral spread or that completed their life cycle early in the season (all of which can be considered strategies to avoid competitive shading) were able to persist in invaded communities, supports the idea that competition for light is one of the main mechanisms through which *H. japonicus* reduces germination and establishment of other species beneath its canopy.

450 *Potential impact at the ecosystem level*

451 The non-random impact of *H. japonicus* based on resident species traits, combined with its
452 tendency to become dominant, resulted in a modification of the overall dominant traits of the
453 invaded communities. When including *H. japonicus* in community weighted means
454 calculations, we showed that invaded communities became taller, which suggests potential
455 repercussions on productivity (Lavorel and Garnier, 2002). Flowering duration was overall
456 shorter which can have repercussions on pollinators' communities (Blaauw and Isaacs, 2014).
457 Last but not least, the invasion by *H. japonicus* strongly modified the life-form spectrum of the
458 communities with an annual vine becoming dominant at the expense of a diversity of life forms
459 including annuals, biennials, and different types of perennial species as well as tree seedlings.
460 Displacement of life forms that maintain leaves during the period of floods (e.g., biennials,
461 stoloniferous perennials) can enhance river banks erosion (Tickner et al., 2001). And finally,
462 absence of tree seedlings can also indicate a potential impact on tree recruitment and
463 successional changes in riparian habitats (Flory and Clay, 2010).

464 *Resilience of restored communities*

465 By allowing recruitment from the seedbank or from adjacent sites, Removal plots recovered
466 rapidly the level of species richness observed in Non-Invaded plots. This is consistent with a
467 previous study conducted on riparian habitats in Scotland where, after the invasive *Mumulus*
468 *guttatus* was removed, vegetation recovered rapidly (Truscott et al., 2008). More generally,
469 recover of species richness has also been highlighted in a meta-analysis comparing removal and
470 non-invaded plots using data from restoration studies (Andreu and Vilà, 2011). However,
471 during the same period, removal plots did not reach the same level of native functional richness
472 as non-invaded plots. These results suggest higher functional redundancy (i.e. more species
473 sharing similar functions) in restored communities compared to Non-Invaded communities.
474 Moreover, based on the community weighted means, we observed no full convergence of

Removal plots towards Non-Invaded plots. The higher proportion of annual species and the lower LDMC observed in Removal compared to Non-invaded sites exemplifies that plots in Removal sites are dominated by fast growing weedy ruderal species including other invasive alien species (e.g., *Ambrosia artemisiifolia*, *Artemisia annua*). This management-mediated secondary invasion is expected when the removal of an invasive plant species leaves place to bare soils (Gooden et al., 2009) which are typically colonized by species with a ruderal strategy, at least during the first years. Using species number only when assessing the resilience of communities may lead to unreasonably optimistic conclusions while the comparison of trait distribution between removal and non-invaded communities as done here provide a more complete assessment (Hejda et al., 2017).

Conclusions

Some previous studies have already highlighted the interest of using a functional approach to understand better the impact of invasive alien plants. However, most of these analyses (e.g., Hejda, 2013, Hejda et al., 2017) relied solely on multi-site comparative studies that cannot disentangle the invader as a passenger or driver of change in trait distribution of resident communities. Our study is novel as it integrates a functional approach coupled with observational and experimental field approaches, thereby offering greater power in ascertaining the causes and effects of invasion and community change. We opt for contrasting the traits of species able to persist versus species declining in presence of the invader, based on the regular comparison of their frequency of occurrence in removal and invaded plots over time as a more informative approach for measuring the impacts of invasion. Moreover, using functional traits and measuring absolute and relative differences between the invader and resident species allows for testing different hypotheses related to coexistence theory. On the one hand, we found that species that are able to use resources at a different period (mostly before) than the dominant

invasive species, can coexist and persist in the same site. On the other hand, we also highlighted that species that display a competitive advantage (high seed mass, perennial with rhizomes or tubers) are also less impacted. Combining these results suggests that 1) whatever their other traits, species that use a different temporal niche can coexist with the invader (*temporal niche differentiation*) while 2) the persistence of species that have the same phenology as the invader is contingent on trait values that confer a high competitive ability. The approach developed in this case study can be used more generally to improve the predictions of impact of invasive species on invaded communities and to follow the trajectory of restored communities.

Acknowledgments: The authors thank the Smage des Gardons and Jean-Philippe Reygrobellet for sites access facilities and for funding LM MSc work in 2014. This work was supported by Smage des Gardons and Anses. CV was supported by the European Research Council (ERC) Starting Grant Project “Ecophysiological and biophysical constraints on domestication in crop plants” (Grant ERC-StG-2014-639706- CONSTRAINTS).

Authors’ contributions: GF and LM designed the study and performed the field work in 2014 (GF, LM) and 2015 (GF). GF analyzed the data and lead the writing of the manuscript with significant inputs and ideas from MC and CV. All the authors reviewed substantially the manuscript.

References

- Andreu, J., Vilà, M., 2011. Native plant community response to alien plant invasion and removal. *Management of Biological Invasions* 2, 81-94.
- Balogh, L., Dancza, I., 2008. *Humulus japonicus*, an emerging invader in Hungary, in: Tokarska-Guzi, B., Brock, J.H., Brundu, G., Child, C.C., Daehler, C., Pysek, P. (Eds.), *Plant Invasions: Human perception, ecological impacts and management*. Backhuys Publishers, Leiden, The Netherlands, pp. 73-91.
- Blaauw, B.R., Isaacs, R., 2014. Flower plantings increase wild bee abundance and the pollination services provided to a pollination-dependent crop. *Journal of Applied Ecology* 51, 890-898.
- Castro-Díez, P., Pauchard, A., Traveset, A., Vilà, M., 2016. Linking the impacts of plant invasion on community functional structure and ecosystem properties. *Journal of Vegetation Science* 27, 1233-1242.
- Chesson, P., 2000. Mechanisms of maintenance of species diversity. *Annual Review of Ecology and Systematics* 31, 343-366.
- Chytrý, M., Maskell, L.C., Pino, J., Pysek, P., Vila, M., Font, X., Smart, S.M., 2008. Habitat invasions by alien plants: a quantitative comparison among Mediterranean, subcontinental and oceanic regions of Europe. *Journal of Applied Ecology* 45, 448-458.
- Colwell, R.K., Coddington, J.A., 1994. Estimating terrestrial biodiversity through extrapolation. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 345, 101.
- Craine, J.M., Wolkovich, E.M., Towne, E.G., Kembel, S.W., 2012. Flowering phenology as a functional trait in a tallgrass prairie. *New Phytologist* 193, 673-682.
- Dezileau, L., Terrier, B., Berger, J.F., Blanchemanche, P., Latapie, A., Freydier, R., Paquier, A., Lang, M., Delgado, J.L., 2014. Reconstitution des crues extrêmes du Gardon à partir

545 d'une analyse paléohydrologique. La Houille Blanche - Revue internationale de l'eau, 44-
546 52.

547 Flory, S.L., Clay, K., 2010. Non-native grass invasion suppresses forest succession.
548 *Oecologia* 164, 1029-1038.

549 Fried, G., Laitung, B., Pierre, C., Chagué, N., Panetta, F.D., 2014. Impact of invasive plants in
550 Mediterranean habitats: disentangling the effects of characteristics of invaders and
551 recipient communities. *Biological Invasions* 16 1639-1658

552 Fried, G., Mahaut, L., Pinston, A., Carboni, M., 2018. Abiotic constraints and biotic
553 resistance control the establishment success and abundance of invasive *Humulus japonicus*
554 in riparian habitats. *Biological Invasions* 20, 315-331.

555 Gallien, L., Carboni, M., 2017. The community ecology of invasive species: where are we and
556 what's next? *Ecography* 40, 335-352.

557 Gallien, L., Mazel, F., Lavergne, S., Renaud, J., Douzet, R., Thuiller, W., 2015. Contrasting
558 the effects of environment, dispersal and biotic interactions to explain the distribution of
559 invasive plants in alpine communities. *Biological invasions* 17, 1407-1423.

560 Garnier, E., Cortez, J., Billès, G., Navas, M.-L., Roumet, C., Debussche, M., Laurent, G.,
561 Blanchard, A., Aubry, D., Bellmann, A., Neill, C., Toussaint, J.-P., 2004. Plant functional
562 markers capture ecosystem properties during secondary succession. *Ecology* 85, 2630-
563 2637.

564 Garnier, E., Navas, M.-L., Grigulis, K., 2016. Plant functional diversity: Organism traits,
565 community structure, and ecosystem properties. Oxford University Press.

566 Gooden, B., French, K., Turner, P.J., Downey, P.O., 2009. Impact threshold for an alien plant
567 invader, *Lantana camara* L., on native plant communities. *Biological Conservation* 142,
568 2631-2641.

569 Hejda, M., 2012. What is the impact of *impatiens parviflora* on diversity and composition of
570 herbal layer communities of temperate forests? *Plos One* 7.

571 Hejda, M., 2013. Do species of invaded communities differ in their vulnerability to being
572 eliminated by the dominant alien plants? *Biological Invasions* 15, 1989-1999.

573 Hejda, M., Bello, F., 2013. Impact of plant invasions on functional diversity in the vegetation
574 of Central Europe. *Journal of Vegetation Science* 24, 890-897.

575 Hejda, M., Hanzelka, J., Kadlec, T., Štrobl, M., Pyšek, P., Reif, J., 2017. Impacts of an
576 invasive tree across trophic levels: Species richness, community composition and resident
577 species' traits. *Diversity and Distributions* 23, 997-1007.

578 Hejda, M., Pyšek, P., Jarošík, V., 2009. Impact of invasive plants on the species richness,
579 diversity and composition of invaded communities. *Journal of Ecology* 97, 393-403.

580 Hulme, P.E., Bremner, E.T., 2006. Assessing the impact of *Impatiens glandulifera* on riparian
581 habitats: partitioning diversity components following species removal. *Journal of Applied*
582 *Ecology* 43, 43-50.

583 Jucker, T., Carboni, M., Acosta, A.T.R., 2013. Going beyond taxonomic diversity:
584 deconstructing biodiversity patterns reveals the true cost of iceplant invasion. *Diversity and*
585 *Distributions* 19, 1566-1577.

586 Julve, P., 1998. *Baseflor*. Index botanique, écologique et chorologique de la flore de France.

587 Kleyer, M., Bekker, R.M., Knevel, I.C., Bakker, J.P., Thompson, K., Sonnenschein, M.,
588 Poschlod, P., Van Groenendael, J.M., Klimes, L., Klimesová, J., Klotz, S., Rusch, G.M.,
589 Hermy, M., Adriaens, D., Boedeltje, G., Bossuyt, B., Dannemann, A., Endels, P.,
590 Götzenberger, L., Hodgson, J.G., Jackel, A.-K., Kühn, I., Kunzmann, D., Ozinga, W.A.,
591 Römermann, C., Stadler, M., Schlegelmilch, J., Steendam, H.J., Tackenberg, O., Wilmann,
592 B., Cornelissen, J.H.C., Eriksson, O., Garnier, E., Peco, B., 2008. The LEDA Traitbase: a
593 database of life-history traits of Northwest European flora. *J. Ecol.* 96, 1266–1274.

594 Kraft, N.J.B., Adler, P.B., Godoy, O., James, E.C., Fuller, S., Levine, J.M., 2015. Community
 595 assembly, coexistence and the environmental filtering metaphor. *Functional Ecology* 29,
 596 592-599.

597 Kumschick, S., Gaertner, M., Vilà, M., Essl, F., Jeschke, J.M., Pyšek, P., Ricciardi, A.,
 598 Bacher, S., Blackburn, T.M., Dick, J.T.A., Evans, T., Hulme, P.E., Kühn, I., Mrugała, A.,
 599 Pergl, J., Rabitsch, W., Richardson, D.M., Sendek, A., Winter, M., 2014. Ecological
 600 Impacts of Alien Species: Quantification, Scope, Caveats, and Recommendations.
 601 *Bioscience* 65, 55-63.

602 Lavorel, S., Garnier, E., 2002. Predicting changes in community composition and ecosystem
 603 functioning from plant traits: Revisiting the Holy Grail. *Functional Ecology* 16, 545-556.

604 MacDougall, A.S., Gilbert, B., Levine, J.M., 2009. Plant invasions and the niche. *Journal of*
 605 *Ecology* 97, 609-615.

606 Mayfield, M.M., Levine, J.M., 2010. Opposing effects of competitive exclusion on the
 607 phylogenetic structure of communities. *Ecology letters* 13, 1085-1093.

608 Shea, K., Chesson, P., 2002. Community ecology theory as a framework for biological
 609 invasions. *Trends in Ecology and Evolution* 17, 170-176.

610 Thomsen, M.S., Wernberg, T., South, P.M., Schiel, D.R., 2016. To include or not to include
 611 (the invader in community analyses)? That is the question. *Biological Invasions* 18, 1515-
 612 1521.

613 Tickner, D.P., Angold, P.G., Gurnell, A.M., Mountford, J.O., 2001. Riparian plant invasions:
 614 hydrogeomorphological control and ecological impacts. *Progress in Physical Geography*
 615 25, 22-52.

616 Tison, J.-M., De Foucault, B., 2014. *Flora Gallica, Flore de France*. Biotope Éditions, Mèze
 617 (France).

618 Truscott, A.M., Palmer, S.C., Soulsby, C., Westaway, S., Hulme, P.E., 2008. Consequences
619 of invasion by the alien plant *Mimulus guttatus* on the species composition and soil
620 properties of riparian plant communities in Scotland. *Perspectives in Plant Ecology*
621 *Evolution and Systematics* 10, 231-240.

622 Vilà, M., Espinar, J.L., Hejda, M., Hulme, P.E., Jarošík, V., Maron, J.L., Pergl, J., Schaffner,
623 U., Sun, Y., Pyšek, P., 2011. Ecological impacts of invasive alien plants: A meta-analysis
624 of their effects on species, communities and ecosystems. *Ecology Letters* 14, 702-708.

625 Vila, M., Tessier, M., Suehs, C.M., Brundu, G., Carta, L., Galanidis, A., Lambdon, P., Manca,
626 M., Medail, F., Moragues, E., Traveset, A., Troumbis, A.Y., Hulme, P.E., 2006. Local and
627 regional assessments of the impacts of plant invaders on vegetation structure and soil
628 properties of Mediterranean islands. *Journal of Biogeography* 33, 853-861.

629 Villéger, S., Mason, N.W.H., Mouillot, D., 2008. New multidimensional functional diversity
630 indices for a multifaceted framework in functional ecology. *Ecology* 89, 2290-2301.

631 Violle, C., Garnier, E., Lecoœur, J., Roumet, C., Pothier, C., Blanchard, A., Navas, M.L.,
632 2009. Competition, traits and resource depletion in plant communities. *Oecologia* 160,
633 747-755.

634 Weiher, E., Freund, D., Bunton, T., Stefanski, A., Lee, T., Bentivenga, S., 2011. Advances,
635 challenges and a developing synthesis of ecological community assembly theory.
636 *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 366,
637 2403-2413.

638 Wolkovich, E.M., Cleland, E.E., 2011. The phenology of plant invasions: a community
639 ecology perspective. *Frontiers in Ecology and the Environment* 9, 287-294.

640

Figure 1. Expected effect of a dominant invasive plant on resident community based on A) the niche differentiation hypothesis or based on B) the competitive hierarchy hypothesis.

Legend:

A) *Niche differentiation hypothesis*: under this hypothesis, resident species that display similar traits than the invader are more impacted, while species displaying different traits can coexist with the invader. Plant invader's establishment decreases species richness and functional richness, but the coexistence of different strategies may limit functional richness decrease. The traits of the resulting invaded community are expected to shift away from the traits of the invader (increased distance).

B) *Competitive hierarchy hypothesis*: the introduction of a tall invasive plant will exclude shorter species by outshading them but species with similar or higher fitness will coexist with the invader. Plant invader's establishment decreases species richness and functional richness. If only similar competitive species (i.e., with similar plant height) coexist with the invader, the average trait values in the invaded communities will become closer to those of the invader. However, similar fitness can be achieved through different combinations of traits. A species with smaller plant height can still coexist with the invader thanks for example to strong lateral spread capacity. In such case, it would be difficult to predict how the average trait values in the invaded communities will shift compare to those of the invader.

Figure 2. Trends in species richness throughout the seasons and years for Removed (Rem.: light grey), Invaded (Inv.: dark grey) and Non-Invaded plots (N.-Inv.: white). For each box plot, top bar is maximum observation, lower bar is minimum observation, top of box is third quartile, bottom of box is first quartile, middle bar is median value and circles are possible outliers.

Figure 3. Changes in a) frequency of occurrence and b) abundance (mean cover) between paired Invaded and Removal plots. The red full line represent the 1:1 line, the black dashed line represent the regression slope of $Fr_{Inv.} \sim Fr_{Rem.}$ and $A_{Inv.} \sim A_{Rem.}$. Species names are coded with the first three letters of the genus followed by the first three letters of the species. Only the most common ($Fr > 5\%$) are shown.

Figure 4. Relationships between the relative impact of *Humulus japonicus* on resident species frequency (RI_{Fr}) and the distance between trait values of the resident species and *Humulus japonicus*. Only significant variable responses are shown (see Table 3 for the detailed results of the linear regressions and Appendix D for all the plots). Each point corresponds to a resident species common to removed and invaded plots. Distances of flowering in month, distance of seed mass in g (log).

a) Absolute functional distance, all censuses

b) Relative functional distance, all censuses

c) Relative functional distance, last censuses (July 2014 & August 2015)

Figure 5. Community-weighted means (CWMs) and proportion of lifeforms at the end of the season without *H. japonicus*. Different letters indicated significant differences of CWMs between treatments according to post-hoc tests. ns: non-significant, * $P < 0.05$, **, $P < 0.01$, *** $P < 0.001$. Units: maximum plant height (m), seed mass (g), specific leaf area (SLA, $\text{mm}^2.\text{mg}^{-1}$), leaf dry matter content (LDMC, mg.g^{-1}), flowering onset and duration (month). Abbreviations for Raunkiær's life forms: Bis_ros: biennials with rosettes, Erect_Per: erected perennials, Per_Sto: perennials with stolons, Per_rhi_tub: perennials with rhizoms or tubers (geophytes), Tree_seedl: tree seedlings.

Figure 6. Community-weighted means (CWMs) and proportion of lifeforms at the end of the season with *H. japonicus*. Different letters indicated significant differences of CWMs between treatments according to post-hoc tests. Ns: non-significant, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Units and abbreviations for Raunkiaer's life forms: see Figure 6.

Table 1. Analysis of deviance table of generalized mixed-effect models on species richness and functional richness. For native species richness and native functional richness, see Appendix A.

Factors	Species richness (S)				Functional richness (FRic)			
	deviance	Chisq	df	P	deviance	Chisq	df	P
Treatment	76.5	76.523	2	<0.001	85.9	85.936	2	<0.001
Site	20.5	20.472	2	<0.001	21.0	21.046	2	<0.001
Year	185.9	185.9	1	<0.001	23.5	23.503	1	<0.001
Season	154.2	154.25	1	<0.001	90.8	90.780	1	<0.001
Treatment:Site	3.5	3.4537	1	0.063	0.0	0.0093	1	0.923
Treatment:Year	25.8	25.79	2	<0.001	11.2	11.177	2	0.004
Treatment:Season	207.5	207.5	2	<0.001	34.2	34.221	2	<0.001
Year:Season	56.9	56.899	1	<0.001	11.1	11.141	1	0.001
Treatment:Year:Season	22.8	22.825	2	<0.001	11.5	11.431	2	0.003

Table 2. Mean native and alien species richness and cover of *Humulus japonicus* per treatment and time.

	<i>H. japonicus cover</i>	Native species richness			Alien species richness		
		Invaded plots	Removal plots	Non-invaded plots	Invaded plots	Removal plots	Non-invaded plots
Second week of March 2014	26.65	6.15±2.60	5.55+/-2.72	8.4+/-3.32	3.15+/-1.31	3.25+/-1.89	3.95+/-1.57
Second week of April 2014	52.40	10.8±3.05	16.0+/-3.20	14.74+/-4.16	5.35+/-1.60	8.5+/-1.91	6.9+/-2.47
Second week of May 2014	93.89	4.44±1.89	15.28+/-3.37	13.8+/-5.43	2.65+/-1.73	4.85+/-2.01	5.35+/-2.64
First week of July 2014	94.70	0.6±0.94	7.75+/-3.68	6.85+/-2.54	2.7+/-1.22	6.6+/-2.06	6+/-2.43
Last week of April 2015	44.83	6.33±3.01	10.29+/-2.14	8+/-4.00	1.7+/-1.89	4.3+/-2.30	4.7+/-2.72
Third week of June 2015	89.40	1.95±1.82	6.95+/-2.14	6.26+/-2.35	0.75+/-1.07	5.7+/-2.05	4.2+/-2.59
Third week of August 2015	100.00	0.05±0.22	4.45+/-1.64	5.9+/-3.42	0.15+/-0.37	3.2+/-1.70	4.55+/-2.89

Table 3. Analysis of variance table of the effect of lifeform and functional distance between resident species and *Humulus japonicus* on species changes in relative frequency (RI_{Fr}) for all the censuses and for the last census. Bold values show significant variables.

Functional <i>absolute</i> distance	df	All censuses				Last census (August 2015)			
		Estimate	SS	F-value	P	Estimate	SS	F-value	P
Lifeforms	5	-	0.442	1.749	0.146	-	0.116	0.296	0.908
Plant height	1	-0.060	0.063	1.246	0.271	0.129	0.043	0.547	0.471
SLA	1	-0.001	0.001	0.012	0.913	-0.016	0.138	1.756	0.205
Seed mass	1	0.025	0.063	1.253	0.270	0.063	0.158	2.010	0.177
LDMC	1	-0.000	0.003	0.059	0.810	-0.002	0.079	1.001	0.333
Flowering onset	1	-0.061	0.319	6.313	0.016*	0.083	0.184	2.340	0.147
Flowering duration	1	0.046	0.231	4.574	0.039*	-0.030	0.057	0.725	0.408
Residuals	40		2.023				1.176		
Functional <i>relative</i> distance	df	Estimate	SS	F-value	P	Estimate	SS	F-value	P
Lifeforms	5	-	0.607	2.430	0.052 (.)	-	0.145	0.419	0.828
Plant height	1	0.042	0.026	0.514	0.477	-0.107	0.096	1.626	0.217
SLA	1	-0.002	0.012	0.238	0.628	-0.011	0.254	4.305	0.051 (.)
Seed mass	1	-0.005	0.003	0.059	0.810	-0.072	0.420	7.106	0.015*
LDMC	1	0.000	0.018	0.356	0.554	0.002	0.160	2.705	0.116
Flowering onset	1	0.043	0.213	4.265	0.045*	-0.068	0.125	2.118	0.161
Flowering duration	1	0.039	0.374	7.478	0.009**	-0.016	0.000	0.001	0.972
Residuals	40		1.998				1.037		