**Successive shifts of the India-Africa transform plate boundary
during the Late Cretaceous-Paleogene interval:
implications for ophiolite emplacement along transforms**

Mathieu Rodriguez¹, Philippe Huchon², Nicolas Chamot-Rooke¹, Marc Fournier², Matthias Delescluse¹, Jeroen Smit³, Alexis Plunder², Grme Calvs⁴, Dia Ninkabou^{1,2}, Manuel Pubellier¹, Thomas Franois⁵, Philippe Agard², Christian Gorini²

1 - Laboratoire de Gologie, Ecole normale suprieure, PSL research university, CNRS UMR 8538, 24 rue Lhomond, 75005 Paris, France

2 - Sorbonne Universit, CNRS-INSU, Institut des Sciences de la Terre de Paris, IStP UMR 7193, F-75005 Paris, France

3 - Department of Earth Sciences, Utrecht University, PO box 80.021, 3508 TA, Utrecht, The Netherlands

4 - Universit Toulouse 3, Paul Sabatier, OMP-GET, 14 Avenue Edouard Belin, 31400-F, Toulouse, France

5 - GEOPS, Univ. Paris-Sud, CNRS, Universit Paris-Saclay, 91405 Orsay, France

Abstract

The Arabian Sea in the NW Indian Ocean is a place where two major transform boundaries are currently active : the Owen Fracture Zone between India and Arabia and the Owen Transform between India and Somalia. These transform systems result from the fragmentation of the India-Africa Transform boundary, which initiated about 90 Myrs ago, when the India-Seychelles block separated from Madagascar to move towards Eurasia. Therefore, the geological record of the Arabian Sea makes it possible to investigate the sensitivity of a transform system to several major geodynamic changes.

Here we focus on the evolution of the India-Africa transform system during the ~47-90 Ma interval. We identify the Late Cretaceous (~90-65 Ma) transform plate boundary along Chain Ridge, in the North Somali Basin. From 65 to ~42-47 Ma, the India-Africa transform is identified at the Chain Fracture Zone, which crossed both the Owen Basin and the North East Oman margin. Finally, the transform system jumped to its present-day location in the vicinity of the Owen Ridge. These shifts of the India-Africa boundary with time provide a consistent paleogeographic framework for the emplacement of the Masirah Ophiolitic Belt, which constitutes a case of ophiolite emplaced along a transform boundary. The successive locations of the India-Africa boundary further highlight the origin of the Owen Basin lithosphere incoming into the Makran subduction zone.

1. Introduction

Transform plate boundaries experience episodes of structural reorganization and migration related to geodynamic changes (e.g., the San Andreas and Queen Charlotte Faults: DeMets and Merkouriev, 2016; the Levant Fault: Smit *et al.*, 2010; the North Anatolian Fault: Hubert-Ferrari *et al.*, 2009; LePichon *et al.*, 2016; the Saint Paul transform: Maia *et al.*, 2016). In the oceanic domain, several transform systems have been active over more than 100 Myrs (Bonatti and Crane, 1982; Ligi *et al.*, 2002; Maia *et al.*, 2016; Maia, 2018), which allows us to investigate their behavior in the wake of major geodynamic changes.

Transform motion between India and Africa started around 90 Ma, when rifting began between Madagascar and the India-Seychelles block (Bernard and Munsch, 2000; Shuhail *et al.*, 2018). Since then, Africa has been split into different plates, including Arabia and Somalia, which still have active transform boundaries with India at the Owen Fracture Zone and the Owen transform, respectively (Fig. 1; Fournier *et al.*, 2008a, b; 2011; Rodriguez *et al.*, 2011). The India-Africa transform plate boundary therefore constitutes a good case-study to investigate the response of an oceanic transform system to geodynamic changes on the 100-Myr scale. The sensitivity of the India-Africa transform system since the Late Paleogene (~24 Ma) is described in Rodriguez *et al.* (2011; 2013; 2014a,b; 2016; 2018). However, the geological description of this transform fault is largely incomplete for the Late Cretaceous-Paleogene, a period marked by two major global plate reorganization events at 63-73 Ma (Cande and Patriat, 2015) and 42-47 Ma (Matthews *et al.*, 2016).

The aim of the present study is to identify the successive locations and configurations of the India-Africa transform boundary during this time interval, based on geological and free-air gravity maps, as well as seismic profiles crossing the Owen and the North Somali Basins. The successive shifts of the India-Africa transform boundary shed light onto 1) the mode of emplacement of the Masirah Ophiolite, and of the Bela, Muslim Bagh, Zhob, and Waziristan-Khost ophiolites in Pakistan and Afghanistan (Fig. 1), and on 2) the origin of the oceanic lithosphere presently subducting under eastern Makran (Pakistan).

2- Geological background & paleogeographic constraints

2-1- Opening of the Indian Ocean

From the Jurassic to the Cenomanian, two subduction zones drove the fragmentation of Gondwanaland and the subsequent opening of the Indian Ocean (Norton and Selater, 1979; Besse and Courtillot, 1988; Seton *et al.*, 2012; Matthews *et al.*, 2012): the Northern Neotethys Subduction (nowadays recorded along the Indus-Yarlong-Tsangpo suture zone) and the East Gondwana Subduction (which was located to the East of Australia and Antarctica Plates). A global plate reorganization event at 100-105 Ma corresponds to the initiation of a new subduction in the Southern Neotethys Ocean, while the East Gondwana subduction deactivated (Matthews *et al.*, 2012; Jolivet *et al.*, 2016). A late consequence of the 100-105 Ma plate reorganization event is the onset of the northwards migration of India in the Late Cretaceous (84-87 Ma; Fig. 2; Bernard and Munschy, 2000; Calvès *et al.*, 2011; Matthews *et al.*, 2012; Gibbons *et al.*, 2013; Battacharya and Yateesh, 2015). India's northwards drift is disturbed in the Late Maastrichtian, resulting in major changes in configuration of Indian Ocean's spreading centers. This change is recorded by the bending of fracture zones since chron 33 (74 Ma) at the SW Indian Ridge (Cande and Patriat, 2015) and a series of ridge jumps from the Mascarenes spreading centers to the Gop basin and eventually, the Carlsberg Ridge since 63 Ma (Royer *et al.*, 2002).

Seafloor spreading rates at the Carlsberg Ridge peaked at 18 cm.yr^{-1} at 52 Ma, before slowing down due to the onset of a global plate reorganization event, partly related to collision between India and the Kohistan-Ladakh arc (Burg, 2011; Cande and Stegman, 2011; Bouilhol *et al.*, 2013; Jagoutz *et al.*, 2015; Cande and Patriat, 2015). However, the role of India-Eurasia collision over global plate dynamics is still highly debated due to uncertainties in its precise timing (see the various paleogeographic scenarios proposing ages of collision between 24 and 55 Ma in Patriat and Achache, 1984; Chaubey *et al.*, 2002; Ali *et al.*, 2008; Molnar and Stock, 2009; van Hinsbergen *et al.*, 2012; Gibbons *et al.*, 2015; Matthews *et al.*, 2015; Buckman *et al.*, 2018). Regardless of the precise timing of the India-Eurasia collision, the Indian Ocean's fabric documents a global plate reorganization event between 47 and 42 Ma, expressed by changes in spreading rates and trends, as well as a major ridge jump from the Wharton Basin to the South East Indian Ridge (Coffin *et al.*, 2002).

2-2. *The fragmentation of the India-Seychelles-Madagascar block*

The series of global plate reorganization events that shaped the Indian Ocean induced several episodes of breakup within the India-Madagascar block initially separated from Africa and Australia in the Late Jurassic (Gibbons *et al.*, 2013).

First, a mid-oceanic ridge active during Albian-Senonian isolated the Kabul Block continental sliver from India (Fig. 2; Tapponnier *et al.*, 1981; Gnos and Perrin, 1997; Gaina *et al.*, 2015).

Second, a Late Cretaceous break-up between Madagascar and the India-Laxmi-Seychelles continental blocks (Fig. 1) led to the opening of the Mascarene and the East Somali Basins (Fig. 2; Schlich, 1982; Bernard and Munsch, 2000; Calvès *et al.*, 2011; Gibbons *et al.*, 2013; Gibbons *et al.*, 2015; Battacharya and Yateesh, 2015; Shuhail *et al.*, 2018).

Third, the break-up between the Seychelles-Laxmi block and India formed the Gop Basin (between magnetic chrons 31 (68 Ma) and 30 (67 Ma; Minshull *et al.*, 2008; Yateesh *et al.*, 2009; Eagles and Hoang, 2014). Following the culmination of the Deccan trap volcanism at ~65.5 Ma (Courtilot and Renne, 2003; Hooper *et al.*, 2010), the onset of the Carlsberg Ridge at chron 28 (63 Ma) marks the full Laxmi-Seychelles break-up (Dyment, 1998; Royer *et al.*, 2002).

2-3- *Late Cretaceous-Early Paleocene ophiolites in the NW Indian Ocean*

Ophiolites are remnants of oceanic lithosphere emplaced over continental lithosphere that are mainly encountered in fossil convergence zones, but also in contexts such as transform boundaries (Dilek and Furnes, 2014). To avoid any confusion, we use the term of ‘*obduction*’ where the ophiolites can be related to a subduction zone, and the term of ‘*ophiolite emplacement*’ where the ophiolite cannot be unambiguously tied to a subduction zone.

The initiation of an intra-oceanic subduction ~100-105 Ma in the southern part of the Neotethys Ocean led to obduction north of Arabia and India during the Campanian (Hacker *et al.*, 1996; Agard *et al.*, 2007, 2011; Hébert *et al.*, 2012; Morris *et al.*, 2017; Nicolas and Boudier, 2017; Guilmette *et al.*, 2018). In the surroundings of the Arabian Sea, the related ophiolites are the Semail in Oman (Searle and Cox, 1999; Breton *et al.*, 2004) and Zhob-Waziristan-Khost ophiolites in NE Pakistan (Cassaigneau, 1979; Badshah *et al.*, 2000; Sarwar, 1992; Khan *et al.*, 2007).

In south east Pakistan, the Bela and Muslim Bagh ophiolites record a different timing of subduction initiation, ranging between 65-70 Ma from the dating of metamorphic soles (Allemann, 1979; Mahmood *et al.*, 1995; Gnos *et al.*, 1998; Khan *et al.*, 2007), and 80 Ma from dating of Supra-Subduction Zone lavas (Kakar *et al.*, 2014). These ophiolites are derived from the basin extending between India and the Kabul Block.

The Masirah Ophiolite running along eastern Oman (Fig. 1) has a different paleogeographic origin. The oceanic crust exposed in the Masirah Ophiolite was formed at a latitude of ~30-50°S (Gnos and Perrin, 1997) in Tithonian-Berriasian times (~140-145 Ma; Peters and Mercolli, 1998). These ophiolites are therefore remnants of the Indian Ocean formed during the early stages of Gondwanaland break-up. The oldest Arabia-derived sediments (detrital Fayah formation) date back to Coniacian at Masirah (Immenhauser, 1996), indicating that the Masirah ophiolite was a part of India's lithosphere prior to its obduction (Gnos *et al.*, 1997). A particularity of the Masirah Ophiolite is the lack of metamorphic sole and hence, possibly no relationship to subduction (Wakabayashi and Dilek, 2003; Agard *et al.*, 2016). The age of emplacement is defined from stratigraphic and structural constraints (Immenhauser, 1996; Schreurs and Immenhauser, 1999; Immenhauser *et al.*, 2000). Late Maastrichtian folds affecting deep-sea sediments in the Batain plain record the first step of emplacement, whereas unconformable, Priabonian shallow water carbonates seal the ophiolite.

2-4- The Oman abyssal plain and the Owen Basin

The oceanic lithosphere incoming into the present-day Makran Subduction Zone is 70 to 100 Myrs-old according to heat flow measurements (Hutchison *et al.*, 1981), similar to ophiolites exposed at Semail. As a result, the lithosphere of the Oman abyssal plain is generally considered as a remnant of the Neotethys Ocean (McCall, 1997; Ravaut *et al.*, 1997; 1998; Ellouz-Zimmerman *et al.*, 2007; Barrier and Vrielynck, 2008; Frizon de Lamotte *et al.*, 2011; Burg, 2018).

However, offshore eastern Oman, most of the Owen basin basement is of Paleogene age (Fig. 3; Mountain and Prell, 1990; Rodriguez *et al.*, 2016). Considering a Neotethys origin for the entire Oman abyssal plain would imply that the Indian Ocean Tithonian lithosphere facing Arabia during the Late Maastrichtian emplacement of the Masirah ophiolites (Peters and Mercolli, 1998) must have been

subducted between the Owen Basin and the Oman abyssal plain. Such a subduction zone is not documented, however, questioning the fate of the Tithonian lithosphere now represented by the Masirah ophiolite, and the origin of the Late Cretaceous lithosphere in the eastern part of the Oman abyssal plain (Rodriguez *et al.*, 2016).

In addition, several features of the Oman abyssal plain remain enigmatic, including the Sonne lineament and the Qalhat Seamount-Little Murray Ridge (Fig. 3). Although the NW-SE trending Sonne lineament imaged on the free-air gravity map (Fig. 3) has been first interpreted as a still active strike-slip fault (Kukowski *et al.*, 2001), no trace of activity has been observed on seismic lines acquired since (Mouchot, 2009). The Qalhat Seamount and Little Murray Ridge constitute a chain of submarine volcanoes of Late Cretaceous age (Edwards *et al.*, 2000; Mouchot, 2009) whose paleogeographic origin remains unclear.

3- Materials and Methods

3-1- Geological, free-air gravity maps and plate reconstructions

The free-air gravity maps (Figs. 3, 4) have been designed using the DTU 13 database filtered for short wavelengths (Andersen *et al.*, 2013). Offshore, a few multibeam tracks have been acquired in the area of the Chain Ridge (Fig. 5) during transits of the AOC and VARUNA-CARLMAG cruises onboard the BHO Beautemps-Beaupré operated by the French navy in 2006 and 2019. The structure of the strike-slip fault system crossing the Huqf desert in Oman (Fig. 6) is mapped after geological maps of Oman at 1: 250 000 (sheets of Duqm, Madreka, Khalouf by Platel *et al.*, 1992) and maps built from seismic data (Filbrandt *et al.*, 2006). The paleogeographic reconstructions of India and Africa continents are drawn after the Gplates files provided in Matthews *et al.* (2016). The successive locations of the India-Africa plate boundary identified hereafter complete these reconstructions.

3-2- Seismic reflection

3-2-1. Sources of the datasets

For the Owen Basin, we use the seismic dataset from the OWEN 2 survey (Figs. 7 to 10; Rodriguez *et al.*, 2016), acquired in 2012 onboard the R/V Beautemps-Beaupré using a high-speed (10 knots)

seismic device. The source consists in two GI air-guns (one 105/105 c.i. and one 45/45 c.i.) fired every 10 seconds at 160 bars in harmonic mode, resulting in frequencies ranging from 15 to 120 Hz. The receiver is a 24-channel, 300-m-long seismic streamer, allowing a common mid-point spacing of 6.25 m and a sub-surface penetration of about 2 s two-way travel time. The standard processing consisted of geometry setting, water-velocity normal move-out, stacking, water-velocity f-k domain post-stack time migration, bandpass filtering and automatic gain control.

New interpretations for seismic lines crossing the North Somali Basin (Fig. 4) and the Batain plain in NE Oman (Fig. 11) are based on the dataset published in Bunce and Molnar (1977) and Beauchamp et al. (1995), respectively.

3-2-2. Stratigraphy in the Owen Basin

Seismic profiles have been tied to drilling sites available in the Arabian Sea from DSDP and ODP legs (Shipboard Scientific Party, 1974a, 1974b, 1989) and stratigraphic details can be found in Rodriguez *et al.* (2016). Two key seismic horizons have been identified on the seismic lines (Figs. 7 to 10). The cross-section of the Oman margin provided in Figure 7 summarizes the stratigraphic framework of the area. A first key reflector corresponds to an angular unconformity recording the end of the uplift of marginal ridges along the Oman margin in the Late Eocene. This unconformity is only recognized in the western part of the Owen Basin, and becomes concordant to the east. A second Late Oligocene-Early Miocene unconformity is observed across the entire Owen Basin, and reflects the diachronous flooding of the Owen Basin by the Indus turbiditic system (Rodriguez *et al.*, 2016). Finally, the top of the Masirah Ophiolites, expressed by a chaotic and highly reflective body on seismic lines (Fig. 7, 11), is considered as Late Maastrichtian, in agreement with onland studies (Immenhauser *et al.*, 1996; Immenhauser *et al.*, 2000).

4- Configurations of the India-Africa plate boundary during the Late Cretaceous-Eocene

4-1- Configuration of the Late Cretaceous India-Africa plate boundary

206 In the northern Somali Basin, on the western edge of the NE-SW trending Chain Ridge, a large sub-
207 vertical fault (offset over ~2 s TWT) is imaged on the seismic profile (Fig. 4; Bunce and Molnar,
208 1977). The fault lineament is well-expressed on the free-air gravity field, and partly mapped on the
209 multibeam coverage of Chain Ridge (Fig. 5). This fault juxtaposes two different oceanic lithospheres.
210 On the eastern flank of Chain Ridge, gabbros dredged at DSDP Site 235 indicate a 90 Myrs-old
211 seafloor (Shipboard Scientific Party, 1974a), i.e., a piece of the East Somali Basin. For the seafloor
212 west of Chain Ridge, magnetic anomalies document a Late Jurassic to Early Cretaceous age (160 to
213 130 Ma; Cochran, 1988; Gaina *et al.*, 2015). The western part of the northern Somali Basin is
214 therefore a remnant of the first stage of opening of the Indian Ocean. The juxtaposition of oceanic
215 lithospheres of different ages is explained by considering the fault observed west of Chain Ridge as
216 the fossil Late Cretaceous India-Africa transform plate boundary (Chain Ridge Transform, Fig. 4).
217 Evidence for Late Cretaceous (Santonian-Campanian) left-lateral strike-slip tectonics is further
218 identified in Oman (Fig. 6). Three-dimensional seismic reflection data document a distributed system
219 of conjugate strike-slip faults, which display typical flower structures, previously mapped by Filbrandt
220 *et al.* (2006). Left-lateral faults are also widely observed in the Huqf desert (Oman), cutting through
221 outcrops of Neoproterozoic remnants of the Pan-African orogeny, at the Khufai, Buah, Shuram,
222 Mukhaibah and Haushi anticlines (Fig. 6; Shackleton and Ries, 1990; Allen, 2007). This set of
223 observations defines the NW-SE trending transtensive Haushi-Nafun-Maradi Fault System (HNMFs
224 hereafter), mapped in Figs. 3 and 6. The HNMFs was active only during the deposition of the Fiqra
225 formation in Early Maastrichtian (Filbrandt *et al.*, 2006) and also locally accompanied by volcanic
226 activity (Glennie *et al.*, 1974; Wyns *et al.*, 1992).
227 Despite a poor sedimentary record due to low sedimentation rates during Late Maastrichtian-Early
228 Paleogene, the seismic lines crossing the western part of the Owen Basin at various latitudes document
229 some sedimentary layers beneath the Late Maastrichtian horizon sealing the Masirah ophiolites (Figs.
230 8-10). These Late Maastrichtian-Early Paleocene layers are trapped in half-graben structures and
231 display a fanning configuration (Figs. 8-10), which reflects tectonic activity. The low density of
232 available seismic lines does not allow us to map accurately these grabens and their precise relationship

with the HNMFS, but they are likely coeval with the tectonic deformation observed along the HNMFS.

4-2- Configuration of the Paleocene-Eocene India-Africa plate boundary

In the North Somali Basin, the Paleocene-Eocene India Africa plate boundary is expressed as the Chain Fracture Zone (Figs. 1, 4), which bounds the seafloor accreted at the Carlsberg Ridge since 63 Ma (Royer et al., 2002; Chaubey et al., 2002). However, the trace of the India-Africa transform boundary has not been clearly identified offshore Arabia in the Owen Basin, due to the complex history of the area related to various episodes of margin reactivation (Rodriguez et al., 2014, 2016). In this section we explore the remnants of the Paleocene-Eocene India Africa plate boundary preserved along the East Oman Margin and the Owen Basin.

Eastern Oman margin -Masirah :

The free-air gravity map of the Oman margin (Fig. 3) documents a series of en-échelon marginal ridges within a ~90-km-wide, N30°E-trending corridor running from Sawqirah to Ras Al Hadd (over ~600 km). The seismic line displayed in Fig. 11 (from Beauchamp et al., 1995) crosses the Masirah Ophiolite onshore, in one of the few places where the initial structure of the ophiolite is still preserved. There, the Masirah Ophiolites display a double vergent thrust stack of numerous ophiolite sheets, according to our revised interpretation (Fig. 11; Beauchamp et al., 1995). The cross section and the profiles displayed in Fig 7 show that elsewhere along the en-échelon marginal ridge system, the offshore segments of the Masirah Ophiolites are scattered. Vertical fault offsets (Fig. 7) indicate that the initial Masirah Ophiolite has been highly dismembered during the formation (i.e., uplift and shearing) of the en-échelon marginal ridge system (Fig. 3).

The stratigraphy of the eastern Oman margin documents the period of formation of this system of en-échelon marginal ridges. Onland, the Masirah ophiolite overlies a Late Maastrichtian, deep-sea detrital formation (Fayah Fm; Immenhauser, 1996), indicating that the marginal ridges did not exist at that time. On the other hand, the Priabonian shallow-water carbonates of the Aydim formation and laterites

cover both the ophiolites and the marginal ridges (Immenhauser, 1996; Shipboard Scientific Party, 1989). The Aydim formation is coeval with a major angular unconformity along the edges of the marginal ridges in the Owen Basin, marking the end of the marginal ridge uplift (Figure 3, 7; Rodriguez *et al.*, 2016). The uplift of the en-échelon system of marginal ridges therefore lasted over more than 25 Myrs, from 65-70 Ma up to ~40 Ma. This period is coeval with numerous geological events identified along the east Oman margin, including alkaline volcanism in the Batain and Haushi-Huqf areas (Gnos and Peters, 2003), and a reorganization of detrital sedimentary systems in SE Oman consistent with surface uplift (Filbrandt *et al.*, 2006; Robinet *et al.*, 2013).

Owen Basin:

The seismic lines displayed in Figs. 8 to 10 document the structural expression of the Paleogene India-Africa plate boundary within the Owen Basin, at various latitudes. In all these lines, we map the area where the indicators of pre-Late Maastrichtian tectonics vanish within the Owen Basin (Figs. 8-10). The area bounding the Late Maastrichtian basins displays a V-shape typical of transform valleys, with flanks characterized by a ~30° slope, shaped by erosion due to bottom current and mass wasting (frequently observed on figs. 8-10). Due to the low sedimentation rates during the Paleogene, the preservation of this extinct transform structure in the geological record is poor.

The location of the Chain Fracture Zone according to reconstructions by Royer *et al.* (2002) coincides with the eastern boundary of the pre-Maastrichtian basins identified on the seismic lines (Figs. 8-10). In reconstructions, the Chain Fracture Zone follows the trend of the northeast Oman margin north of 20°N (Fig. 3), where its steepness is compatible with a transform margin. The eastern part of the Owen Basin is Paleogene in age, consistent with basement ages obtained at ODP drilling sites along the Owen Ridge (Shipboard Scientific Party, 1974b, 1989). The composite age of the basement may therefore be explained by a major transform boundary crossing the Owen Basin (Fig. 3, Rodriguez *et al.*, 2016).

5- Discussion

Identification or reappraisal of these structural relationships highlight a major change in the configuration of the India-Africa boundary around ~65-70 Ma (i.e., coeval with the global plate reorganization event; Cande and Patriat, 2015). Data indicate that the India-Africa plate boundary migrated from the Chain Ridge Transform to the Chain Fracture Zone (Fig. 12). Here we address critical points of the structural evolution of the India-Africa plate boundary raised by the reconstruction of figure 12, as well as the paleogeographic implications of the transform plate boundary migration.

5-1-Structural evolution of the India-Africa plate boundary during the Late Cretaceous-Eocene

For the Late Cretaceous India-Africa plate boundary, our reconstruction highlights a lack of direct connection between the Chain Ridge transform offshore Somalia and the HNMFS in Oman (Fig. 12). While the HNMFS could be a distributed strike-slip system related to the partitioning of India-Africa motion along the offshore Chain Ridge Transform (Filbrandt *et al.*, 2006), its trend may reflect older Neoproterozoic structures (Allen, 2007) reactivated in response to the complex Late Cretaceous stress field, influenced by both the obduction of the Semail to the north and transform tectonics to the east.

For the Paleocene-Middle Eocene India-Africa plate boundary, the development of the en-échelon system of marginal ridges constitutes the clearest record of transform fault activity. The en-échelon system of marginal ridges may be interpreted as the result of a sheared transform margin (Fig. 13), in a transpressive, partitioned left-lateral strike-slip system of deformation related to the Chain Fracture Zone between ~65-70 Ma and ~40-45 Ma. Strain partitioning may explain the development of the marginal ridges along the southern part of the Oman margin, which was located more than 100 km away from the Chain Fracture Zone. In this framework, the en-échelon system of marginal ridges results from the reactivation of Tithonian passive margin structures, and probably structures reactivated during the activity of the HNMFS (Fig. 13; see below).

Trehu *et al.* (2015) demonstrated for a similar marginal ridge at the Queen Charlotte Transform (Carlson *et al.*, 1988; Barrie *et al.*, 2013; Rhor 2015) that such a configuration is typical of a slight (<15°) component of oblique convergence inducing transpression.

5-2- Emplacement of the Masirah Ophiolite along the east Oman transform margin

These results allow to reappraise the emplacement of the Masirah ophiolite of Jurassic age (Fig. 13). The story begins during the Late Maastrichtian, coeval with the migration of the India-Africa plate boundary at ~65-70 Ma from Chain Ridge to the Chain Fracture Zone. Paleogeographic reconstructions show that the Chain Fracture Zone must have crossed the Owen basin and the northeastern Oman margin in the area of the Batain plain (Royer *et al.*, 2002; Fournier *et al.*, 2010; Rodriguez *et al.*, 2016). There, the Masirah Ophiolite was a double vergent structure (Fig. 11) formed along a transpressive segment of the Chain Fracture Zone, or a remnant of Chain Ridge passing by the Chain Fracture Zone. The Masirah Ophiolite is uplifted during the Paleogene, when the en-échelon marginal ridges of the Oman transform margin developed (Fig. 10). During the Paleocene-Eocene, distributed left-lateral shear and uplift of the marginal ridges promoted the dismembering of the ophiolite fragments initially located in the Batain plain (Fig. 13). Dismembering of the ophiolites within the shear zone would explain why the double vergent structure identified at the Batain, as well as the fold system in front of the ophiolites (Schreurs and Immenhauser, 1999), are no longer observed to the south at Sawqirah or Masirah.

The initial double-vergent structure proposed for the Masirah ophiolites along the Chain Fracture Zone is very similar to many transverse ridges observed worldwide, such as the Davie Ridge offshore Mozambique (Mahanjane, 2014), the St Paul transverse ridge in the Atlantic (Maia *et al.*, 2016) or the MacQuarie Ridge at the Australia-Pacific plate boundary (Meckel *et al.*, 2003). The fact that double vergent structures may be a favorable setting for the initiation of ophiolite emplacement has already been proposed for the Meratus ophiolite in SE Borneo (Pubellier *et al.*, 1999) along an oblique convergent plate boundary.

5-3-Paleogeographic implications

Figure 14 presents paleogeographic reconstructions, which emphasize the successive locations of the India-Africa boundary and the related transfer of fragments of oceanic lithosphere from one plate to another. The migration of the India-Africa plate boundary from the Chain Ridge to the Chain Fracture Zone around 65-70 Ma induced a transfer of a sliver of Late Cretaceous oceanic lithosphere from the

East Somali Basin to Africa, explaining the age contrast with the western part of the North Somali Basin (Fig. 14c). The Late Cretaceous lithosphere from the East Somali Basin was displaced northwards along the Chain Fracture Zone, while seafloor was formed at Carlsberg Ridge (Fig. 14d). The plate boundary then jumped to its present-day location along the Owen Ridge, during the Late Eocene-Oligocene (Rodriguez *et al.*, 2016), probably as a consequence of India-Eurasia collision and the global plate reorganization event recorded around 47 Ma (Müller *et al.*, 2016). This migration of the India-Africa plate boundary lead to another episode of transfer to Africa of a piece of the oceanic lithosphere accreted at the Carlsberg Ridge (Fig. 14e; Rodriguez *et al.*, 2016). In this reconstruction, the composite origin of the Owen Basin is explained by the juxtaposition of remnants of a Tithonian passive margin to the west, slices of a Tithonian proto-Indian Ocean lithosphere preserved along the Masirah ophiolite, and Paleogene lithosphere formed at the Carlsberg Ridge to the east (Fig. 14). Therefore, the Late Cretaceous lithosphere subducting in front of the Makran has two different paleogeographic origins (Fig. 14): the Neotethys west of the Chain Fracture Zone and the East Somali Basin to the east (east of ~61°E).

This scenario also provides an explanation for the paleogeography of the Tithonian lithosphere at the origin of the Masirah ophiolite, that does not require a subduction zone between the Owen Basin and the Oman abyssal plain (Fig. 14). The East Somali Basin spreading center formed within the Tithonian oceanic lithosphere at the origin of the ophiolites, or at the tip of the Lower Cretaceous basin located between India and the Kabul block (Fig. 14). During the Late Cretaceous, the Tithonian lithosphere was progressively displaced to the north along the Chain Ridge Transform, while the East Somali and Mascarenes Basins opened to the south (the spreading center being at the tip of the Chain Ridge Transform). In the Late Maastrichtian, when the Masirah ophiolite emplacement started (unrelated to any subduction), the Tithonian lithosphere had reached the latitude of Oman (Fig. 12,14). After the Masirah obduction, the remaining Tithonian lithosphere located east of the Chain Fracture Zone was subducted beneath eastern Makran (Figs. 14c,d). In detail, this reconstruction suggests that the Sonne lineament, Qalhat seamount and Little Murray Ridge (Fig. 3) are parts of the fossil ocean-ocean transition between the East Somali Basin and the Carlsberg seafloor, when the Carlsberg Ridge formed in the wake of the Deccan plume (Dyment, 1998).

With regards to the eastern Pakistan ophiolites, the India-Africa plate boundary was located west of the Kabul block in the Late Maastrichtian according to our reconstruction (Fig. 14), making it difficult to consider these ophiolite sequences as remnants of an India-Africa transform, unless a complex stepover is involved. Instead, during the Paleogene-Early Eocene, the trend of the Chain Fracture Zone allows a connection with the Sistan Ocean (Fig. 14; Treolar and Izatt, 1993; McCall, 1997). More constraints from the surroundings of the Kabul block are needed to further document this episode of transform tectonics.

5-4- Migration of the India-Africa plate boundary at ~74-63 Ma as part of a global plate reorganization event?

The reorganization of the India-Africa plate boundary identified around 74-63 Ma may result from a major geodynamic reorganization event affecting the entire Indian Ocean. The driver of this plate reorganization event is a matter of debate. The Indonesian Slab reached the Lower Mantle during this period, which affects the slab pull force (Faccenna *et al.*, 2013). The volcanic activity of the Deccan-Réunion Plume peaked at 65 Ma, but earliest traces of Deccan type volcanism are encountered around 110-120 Ma (Mahoney *et al.*, 2002). Both slab penetration into the lower mantle and plume-push may affect oceanic spreading patterns at mid-oceanic ridges and global plate dynamics (Cande *et al.*, 2011; van Hinsbergen *et al.*, 2011; Faccenna *et al.*, 2013), even if the relative contribution of each process to force balance remains a matter of debate (Anderson, 2001; Bercovici, 2003; Cande and Stegman, 2011). However, these processes do not explain the abrupt change in plate boundary configuration.

Two punctual geological events (duration <5 myrs) may have contributed to the plate boundary reorganization event by affecting subduction dynamics (boundary forces):

- deactivation of the subduction at the origin of the Bela and Muslim-Bagh ophiolites and the Masirah Slab in tomography (Gaina *et al.*, 2015);
- collision of the Woyla Arc (Wajzer *et al.*, 1991; Gibbons *et al.*, 2015) and Burma Block by the Late Maastrichtian (Socquet and Pubellier, 2005) or the Early Tertiary (Searle *et al.*, 2007) with southeastern Eurasia.

While collision of continental terranes and subduction deactivation events are common phenomenon in plate tectonics, their relationship to transform boundary migration events is yet unclear (Maia, 2018). Our reconstructions nevertheless provide first order constraints on the context in which the migration of the India-Africa transform boundary occurred and a way to test the sensitivity of transform boundary to geodynamic changes. At ~70 Ma, spreading and convergence rates were high ($>10 \text{ cm.yr}^{-1}$), the age contrast between the adjacent oceanic lithospheres at the Chain Ridge transform was $> 30 \text{ Myrs}$, and the Chain Ridge transform offset was at least of 500-km (Figs. 12, 14). We propose that the integrated strength of the Chain Ridge transform may have been too high at the time of this plate reorganization to further accommodate the relative motion of India-Africa, hence resulting in the relocation of the transform boundary in a weaker area, at the Chain Fracture Zone.

6- Conclusions

Our reconstructions show that during the Late Cretaceous-Late Eocene interval, the India-Africa transform boundary migrated in response to broader geodynamic events. The Late Maastrichtian migration from the Chain Ridge Transform to the Chain Fracture Zone records the plate reorganization event at 74-63 Ma (Cande and Stegman, 2011). The Late Eocene migration from the Chain Fracture to the present-day location along the Owen Ridge records the well-defined global plate reorganization event at ~42-47 Ma (Muller *et al.*, 2016). The successive episodes of migration of the India-Africa plate boundary since the Late Cretaceous contributed to the transfer of oceanic slivers between both plates. Our updated paleogeographic framework suggests that the lithosphere presently subducting beneath eastern Makran originates from the East Somali Basin in the Indian Ocean, instead of the Neotethys as previously proposed.

This study also highlights how ophiolites may be emplaced along a transform boundary as a consequence of a transform migration event. Although the proposed scenario fits with all available geological constraints, a denser grid of seismic reflection profiles along the east Oman margin is needed to confirm and describe more precisely the mode of development of the en-échelon marginal ridges, and the related uplift and dismembering of the Masirah Ophiolitic Belt.

422

423 **Acknowledgements**

424 *Seismic lines were processed using Geocluster software from CGG Veritas. This study was supported*
425 *by SHOM, Ifremer and INSU-CNRS. We warmly thank the editor D. Wyman, Edwin Gnos and an*
426 *anonymous reviewer for their very helpful and constructive comments.*

427

428

429 **Figure captions:**

430 **Figure 1:** Structural map of the north western Indian Ocean, showing the currently active plate
431 boundaries, the distribution of the oceanic basins, as well as the distribution of the main ophiolites.

432 **Figure 2:** General paleogeographic reconstructions of the Indian Ocean from Early Cretaceous to
433 Early Tertiary, proposed by Gaina *et al.* (2015), and main geological events during the history of the
434 Indian Ocean. KB: Kabul Block; Masc. : Mascarenes; Sey: Seychelles; GPRE : Global Plate
435 Reorganization Events.

436 **Figure 3:** a) Free air gravity map of the Owen Basin offshore Oman and the Arabian Sea (filtered for
437 short wavelengths from DTU 13 database; Andersen *et al.*, 2013) and b) its interpretation,
438 with an emphasis on the composite age of the basement of the Owen Basin. HNMFS: Haushi-
439 Nafun-Maradi Fault System.

440 **Figure 4:** a) Free air gravity map of the North Somali Basin (filtered for short wavelengths from DTU
441 13 database; Andersen *et al.*, 2013) and b) its interpretation, with an emphasis on the composite age of
442 the basement of the North Somali Basin. c) New interpretation of a vintage seismic line from Bunce
443 and Molnar (1977) highlighting a major fossil transform fault at the western edge of Chain Ridge,
444 interpreted as a remnant of the Late Cretaceous India-Africa plate boundary.

445 **Figure 5 :** Multibeam map of the northern termination of the Chain Ridge and Chain Fracture Zone

446 **Figure 6:** a) Map of the Haushi Nafun Maradi Fault system; b) Simplified geological sketchmap of the
447 Huqf desert highlighting strike-slip offsets related to the Haushi-Nafun fault, after Platel *et al.*, 1992;
448 c) line drawing of a seismic profile crossing the Maradi fault, highlighting a negative flower structure,
449 redrawn from Filbrandt *et al.*, 2006.

450 **Figure 7:** a) Seismic line from the OWEN 2 cruise (Rodriguez *et al.*, 2016) crossing the western side
451 of the Sawqirah Ridge, where the Masirah ophiolites are tilted westwards, in direction of the Oman
452 platform; b) Seismic line from the OWEN 2 cruise (Rodriguez *et al.*, 2016) crossing the eastern edge
453 of the Sawqirah Ridge, showing a piece of the Masirah Ophiolite lying in the Owen Basin, and the
454 unconformity corresponding to the end of the formation of the marginal ridges. c) Cross section of the

Oman margin at the latitude of the Sawqirah Ridge (modified from Rodriguez *et al.*, 2016). Three main angular unconformities can be tracked across the Owen Basin.

Figure 8: Seismic lines from the Owen-2 cruise (Rodriguez *et al.*, 2016) crossing the Owen Basin.

Figure 9 : Seismic lines from the Owen-2 cruise (Rodriguez *et al.*, 2016) crossing the Owen Basin.

Figure 10 : Seismic lines from the Owen-2 cruise (Rodriguez *et al.*, 2016) crossing the Owen Basin.

These seismic lines highlight the presence of pre-Maastrichtian fanning configurations lying above the basement of the western part of the Owen Basin, and their absence in its eastern part. The boundary of the pre-Maastrichtian fanning configurations roughly coincides with the expected location of the fossil Chain Fracture Zone (according to reconstructions by Royer *et al.*, 2002).

Figure 11: a) Seismic profile crossing the Masirah ophiolite in the Batain plain (from Beauchamp *et al.*, 1995) and b) its revised interpretation, highlighting the positive flower structure of the Masirah Ophiolites. See fig. 3 for location.

Figure 12: Reconstruction of the configuration of the India-Africa plate Boundary at 65-70 Ma and 60 Ma, highlighting the major migration of the transform system occurring at that time.

Figure 13: Simplified reconstruction of the mode of emplacement of the Masirah ophiolites along with the development of the en-échelon marginal ridge system offshore Oman. Note that the emplacement of the ophiolites last over more than 25 Myrs and does not imply any subduction zone.

Figure 14: Revised paleogeographic reconstructions of the northwestern Indian Ocean, from 90 Ma to present-day, modified after Matthews *et al.* (2016). These reconstructions highlight the successive locations of the India-Africa plate boundary: the Chain Ridge Transform from 90 to 65 Ma, the Chain Fracture Zone from 65 to 47 Ma, in the vicinity of the Owen Ridge since the Oligocene. The main paleogeographic implications are the East Somali Basin origin of the lithosphere subducting beneath the eastern Makran, and the disconnection between the India-Africa transform boundary and the eastern Pakistan ophiolites.

References

482 Agard, P., Jolivet, L., Vrielynck, B., Burov, E., Monié, P., 2007. Plate acceleration: the
 483 obduction trigger? *Earth and planetary science letters* 258, 428-441.

484 Agard, P., Omrani, J., Jolivet, L., Whitechurch, H., Vrielynck, B., Spakman, W., Monié, P.,
 485 Meyer, B., Wortel, R., 2011. Zagros orogeny: a subduction-dominated process, In:
 486 Lacombe, O., Grasemann, B., Simpson, G. (Eds.), *Geodynamic Evolution of the Zagros*.
 487 *Geological Magazine* 692–725.

488 Agard, P., Yamato, P., Soret, M., Prigent, C., Guillot, S., Plunder, A., Dubacq, B., Chauvet,
 489 A., Monié, P., 2016. Plate interface rheological switches during subduction infancy:
 490 control on slab penetration and metamorphic sole formation. *Earth and Planetary Science*
 491 *Letters*, 451, 208-220.

492 Allen, P.A., 2007. The Huqf supergroup of Oman: basin development and context for
 493 Neoproterozoic glaciation. *Earth Science Reviews* 84, 139-185.

494 Allemann, F., 1979. Time of emplacement of the Zhob Valley ophiolites and Bela ophiolites,
 495 Baluchistan (preliminary report) in: *Geodynamics of Pakistan*, Geol. Surv. Pakistan, Spec.
 496 Mem. 215-242.

497 Ali, J.R., Aitchison, J.C., 2008. Gondwana to Asia : Plate tectonics, paleogeography and the
 498 biological connectivity of the Indian sub-continent from the Middle Jurassic through latest
 499 Eocene (166-35 Ma). *Earth Science Reviews* 88, 145-166.

500 Andersen O., Knudsen, P., Kenyon, S., Factor, J., Holmes, S., 2013. The dtu13 global marine
 501 gravity field—first evaluation. Technical report, DTU Space - National Space Institute

502 Anderson, D.L., 2001. Top-down Tectonics? *Science* 293, 2016-2018.

503 Badshah, M. S., Gnos, E., Jan, M. Q., Afridi, M. I., 2000. Stratigraphic and tectonic evolution
 504 of the northwestern Indian plate and Kabul Block, in Khan, M. A., Treloar, P. J., Searle,
 505 M. P., and Jan, M., eds., *Tectonics of the Nanga Parbat Syntaxis and the Western*
 506 *Himalaya*, Geological Society of London Special Publication, 170, 467-475.

507 Barrie, J.V., Conway, K.W., Harris, P.T., 2013. The Queen Charlotte Fault, British Columbia:
 508 seafloor anatomy of a transform fault and its influence on sediment processes. *Geo-Marine*
 509 *Letters* 33, 311-318

510 Barrier, E., Vrielynck, B., 2008. Palaeotectonic maps of the Middle East. Tectono-
 511 sedimentary-palinspastic maps from Late Norian to Pliocene. 14 maps. Paris
 512 CGMW/CCGM

513 Bhattacharya, G.C., Yatheesh, V., 2015. Plate-tectonic evolution of the deep ocean basins
 514 adjoining the western continental margin of India - a proposed model for the early opening
 515 scenario, In: Mukherjee, S. (Ed.), *Petroleum Geoscience: Indian Contexts*. Springer
 516 International Publishing, Switzerland, 1-61.

517 Beauchamp, W. H., Ries, A. C., Coward, M. P., Miles, J.A., 1995. Masirah Graben, Oman : a
 518 hidden Cretaceous rift basin? *American Association of Petroleum Geologists' Bulletin* 79,
 519 864-79.

520 Beck, R.A., Burbank, D.W., Sercombe, W.J., Riley, G.W., Barndt, J.K., Berry, J.R., Afzal, J.,
 521 Khan, A.M., Jurgen, H., Metje, J., Cheema, A., Shafique, N.A., Lawrence, R.D., Khan, A.,
 522 1995. Stratigraphic evidence for an early collision between northwest India and Asia.
 523 *Nature* 373, 55-58

524 Bercovici, D., 2003. The generation of plate tectonics from mantle convection. *Earth*
 525 *Planetary Science Letters* 205, 107-121.

526 Bernard, A., Munsch, M., 2000. Le bassin des Mascareignes et le bassin de Laxmi (océan
 527 Indien occidental) se sont-ils formés à l'axe d'un même centre d'expansion ? *Comptes*
 528 *Rendus de l'Académie des Sciences Paris* 330, 777–783

529 Besse, J., Courtillot, V., 1988. Paleogeographic maps of the continents bordering the Indian
 530 Ocean since the Early Jurassic. *Journal of Geophysical Research* 93, 11791-808.

531 Bouilhol, P., Jagoutz, O., Hanchar, J.M., Dudas, F.O., 2013. Dating the India-Eurasia
 532 collision through arc magmatic records. *Earth and Planetary Science Letters* 366, 163-175.

533 Breton, J.-P., Béchennec, F., Le Métour, J., Moen-Maurel, L., Razin, P., 2004. Eoalpine (Cretaceous)
 534 evolution of the Oman Tethyan continental margin : insights from a structural field study in Jabal
 535 Akhdar (Oman Mountains). *Georabia* 9, 1-18

536 Buckman, S., Aitchinson, J.C., Nutman, A.P., Bennett, V.C., Saktura, W.M., Walsh, J.M.J.,
 537 Kachovich, S., Hidaka, H., 2018. The Spongtang Massif in Ladakh, NW Himalaya: an Early
 538 Cretaceous record of spontaneous, intra-oceanic subduction initiation in the Neotethys. *Gondwana*
 539 *Research* 63, 226-249.

540 Bunce, E.T., Molnar, P., 1977. Seismic reflection profiling and basement topography in the
 541 Somali Basin: possible fracture zones between Madagascar and Africa. *Journal of*
 542 *Geophysical Research* 82, 5305-5311

543 Burg J.P., 2011. The Asia-Kohistan-India collision : review and discussion. D. Brown and
 544 P.D. Ryan (eds), *Arc-Continent Collision, Frontiers in Earth Sciences*, 279-309, doi
 545 10.1007/978-3-540-88558-0_10

546 Burg, J.P., 2018. Geology of the onshore Makran accretionary wedge: synthesis and tectonic
 547 interpretation. *Earth Science Reviews* 185, 1210-1231.

548 Calvès, G., Schwab, A. M., Huuse, M., Clift, P. D., Gaina, C., Jolley, D., Tabrez, A. R., Inam,
 549 A., 2011. Seismic volcanostratigraphy of the western Indian rifted margin: The pre-Deccan
 550 igneous province, *Journal of Geophysical Research* 116, B01101, doi:
 551 10.1029/2010JB000862.

552 Cande S.C., Stegman, D.R., 2011. Indian and African plate motions driven by the push force
 553 of the Réunion plume head. *Nature* 475, 47-52

554 Cande, S.,C. Patriat, P., 2015. The anticorrelated velocities of Africa and India in the Late
 555 Cretaceous and early Cenozoic. *Geophysical Journal International* 200, 227-243.

556 Carlson, P.R., Bruns, T.R., Plafker, G., 1988. Late Cenozoic offsets on the offshore
 557 connection between the Fairweather and Queen Charlotte faults off southeast Alaska.
 558 Marine Geology 85, 89-97.

559 Cassaigneau, C., 1979, Contribution à l'étude des structures Inde-Eurasie. La zone de suture
 560 de Khost dans le Sud-Est de l'Afghanistan, l'obduction Paléocène et la tectonique Tertiaire,
 561 PhD-thesis, 124: Université des Sciences et Techniques du Languedoc, Montpellier

562 Chaubey, A.K., Dymant, J., Bhattacharya, G.C., Royer, J.Y., Srinivas, K., Yatheesh, V., 2002.
 563 Paleogene magnetic isochrons and palaeo-propagators in the Arabian and Eastern Somali
 564 basins, NW Indian Ocean, In: Clift, P.D., Croon, D., Gaedicke, C., Craig, J. (Eds.), The
 565 Tectonic and Climatic Evolution of the Arabian Sea Region. Geological Society, London,
 566 Special Publication 195, 71-85.

567 Cochran, J. R., 1988. Somali Basin, Chain Ridge, and origin of the northern Somali Basin
 568 gravity and geoid low. Journal of Geophysical Research 93, 11985-12008.

569 Coffin, M. F., Pringle, M.S., Duncan, R.A., Gladchenko, T.P., Storey, M., Müller, R.D.,
 570 Gahagan, L.A., 2002. Kerguelen Hotspot magma output since 130 Ma. Journal of
 571 Petrology 43, 1121-1139.

572 Courtillot, V.E., Renne, P.R., 2003. On the ages of flood basalt events. Comptes rendus
 573 Géoscience 335, 113-140.

574 Delescluse, M., Montési, L. G. J., Chamot-Rooke, N., 2008. Fault reactivation and selective
 575 abandonment in the oceanic lithosphere. Geophysical Research Letters 35, L16312,
 576 doi:10.1029/2008GL035066.

577 DeMets, C., Merkouriev, S., 2016. High-resolution estimates of Nubia-Somalia plate motion
 578 since 20 Ma from reconstructions of the Southwest Indian Ridge, Red Sea and Gulf of
 579 Aden. Geophys. J. Int., 207, 317-322, doi.org/10.1093/gji/ggw276

580 Dilek, Y., Furnes, H., 2014. Ophiolites and their origins. Elements 10, 93-100.

581 Dymment, J., 1998. Evolution of the Carlsberg Ridge between 60 and 45 Ma: Ridge
582 propagation, spreading asymmetry, and the Deccan-Reunion hotspot. *Journal of*
583 *Geophysical Research* 103, 24067–24084 doi: 10.1029/98JB01759.

584 Eagles, G., Hoang, H.H., 2014. Cretaceous to present kinematics of the Indian, African and
585 Seychelles plates. *Geophysical Journal International* 196, 1-14

586 Edwards, R.A., Minshull, T. A., White, R. S., 2000. Extension across the Indian–Arabian
587 plate boundary: the Murray Ridge. *Geophysical Journal International* 142, 461-477.

588 Ellouz Zimmermann, N. et al., 2007, Offshore frontal part of the Makran accretionary prism (Pakistan)
589 the Chamak Survey, in *Thrust Belts and Foreland Basins: From Fold Kinematics to Hydrocarbon*
590 *Systems*, edited by O. L. Lacombe et al., pp. 349–364, Springer, Berlin.

591 Filbrandt, J. B., Al-Dhahab, S., Al-Habsy, A., Harris, K., Keating, J., Al-Mahruqi, S., Ozkaya,
592 S. I., Richard, P. D., Robertson, T., 2006. Kinematic interpretation and structural evolution
593 of North Oman, Block 6, since the Late Cretaceous and implications for timing of
594 hydrocarbon migration into Cretaceous reservoirs. *GeoArabia* 11, 97-140.

595 Fournier, M., Petit C., Chamot-Rooke, N., Fabbri, O., Huchon, P., Maillot, B., Lévrier, C.,
596 2008a. Do ridge-ridge-fault triple junctions exist on Earth? Evidence from the Aden-
597 Owen-Carlsberg junction in the NW Indian Ocean. *Basin Research* 20, 575-590, doi:
598 10.1111/j.1365-2117.2008.00356.x

599 Fournier, M., Chamot-Rooke, N., Petit, C., Fabbri, O., Huchon, P., Maillot, B., Lévrier, C.,
600 2008b. In-situ evidence for dextral active motion at the Arabia-India plate boundary.
601 *Nature Geoscience* 1, 54-58, doi: 10.1038/ngeo.2007.24.

602 Fournier, M., Chamot-Rooke, N., Petit, C., Huchon, P., Al-Kathiri, A., Audin, L., Beslier, M.-
603 O., d'Acremont, E., Fabbri, O., Fleury, J.-M., Khanbari, K., Lévrier, C., Leroy, S.,
604 Maillot B., Merkouriev, S., 2010. Arabia-Somalia plate kinematics, evolution of the Aden-
605 Owen-Carlsberg triple junction, and opening of the Gulf of Aden. *Journal of Geophysical*
606 *Research* 115, B04102, doi: 10.1029/2008JB006257

607 Fournier, M., Chamot-Rooke N., Rodriguez, M., Huchon, P., Petit, C., Beslier, M.-O.,
 608 Zaragosi S., 2011. Owen Fracture Zone: the Arabia-India plate boundary unveiled. *Earth*
 609 *and Planetary Science Letters* 302, 247-252, doi:10.1016/j.epsl.2010.12.027.

610 Frizon de Lamotte, D., Raulin, C., Mouchot, N., Wrobel-Daveau, J.-C., Blanpied, C.,
 611 Ringenbach, J.-C., 2011. The southernmost margin of the Tethys realm during the
 612 Mesozoic and Cenozoic : initial geometry and timing of the inversion processes. *Tectonics*
 613 30, TC3002, doi:10.1029/2010TC002691

614 Gaina, C., van Hinsbergen, D.J.J., Spakman, W., 2015. Tectonic interactions between India
 615 and Arabia since the Jurassic reconstructed from marine geophysics, ophiolite geology,
 616 and seismic tomography. *Tectonics* 34, 875-906, doi: 10.1002/2014TC003780.

617 Gibbons, A. D., Whittaker, J. M., Müller, R. D., 2013. The breakup of East Gondwana:
 618 Assimilating constraints from Cretaceous ocean basins around India into a best-fit tectonic
 619 model. *Journal of Geophysical Research Solid Earth* 118, doi:10.1002/jgrb.50079.

620 Gibbons, A.D., Zahirovic, S., Müller, R.D., Whittaker, J.M., Yateesh, V., 2015. A tectonic
 621 model reconciling evidence for the collisions between India, Eurasia and intra-oceanic arcs
 622 of the central-eastern Tethys. *Gondwana Research* 28, 451-492.

623 Glennie, K. W., Boeuf, M. G. A., Hughes-Clark, M. W., Moody-Stuart, M., Pilar, W. F. H.,
 624 and Reinhardt, B. M., 1974, *Geology of the Oman Mountains*: Verh. K. Ned. Geol.
 625 *Mijnbouwk. Genoot. Ged.*31, 1-423.

626 Gnos, E., Immenhauser, A., Peters, Tj., 1997. Late Cretaceous/early tertiary convergence
 627 between the Indian and Arabian plates recorded in ophiolites and related sediments.
 628 *Tectonophysics* 271, 1-19.

629 Gnos, E. and Perrin, M., 1997. Formation and evolution of the Masirah ophiolite constrained
 630 by paleomagnetic study of volcanic rocks. *Tectonophysics* 253, 53-64.

631 Gnos, E., Khan, M., Mahmood, K., Khan, A. S., Shafique, N. A., and Villa, I. M., 1998, Bela
 632 oceanic lithosphere assemblage and its relation to the Réunion hotspot: *Terra Nova* 10, 90-95.
 633 Gnos, E., and Peters, T., 2003, Mantle xenolith-bearing Maastrichtian to Tertiary alkaline
 634 magmatism in Oman.: *Geochemistry, Geophysics, Geosystems* 4, 8620.
 635 Guilmette, C., Smit, M.A., van Hinsbergen, D.J.J., Gürer, D., Corfu, F., Charette, B.,
 636 Maffione, M., Rabeau, O., Savard, D., 2018. Forced subduction initiation recorded in the
 637 sole and crust of the Semail Ophiolite of Oman. *Nature geosciences* 11, 688-695.
 638 <https://doi.org/10.1038/s41561-018-0209-2>
 639 Hacker, B.R., Mosenfelder, J.L., Gnos, E., 1996. Rapid emplacement of the Oman ophiolite:
 640 thermal and geochronologic constraints. *Tectonics* 15, 1230-1247.
 641 Hébert, R., Bezard, R., Guilmette, C., Dostal, J., Wang, C.S., Liu, Z.F., 2012. The Indus-
 642 Yarlung Zangbo ophiolites from Nanga Parbat to Namche Barwa syntaxes, southern
 643 Tibet : first synthesis of petrology, geochemistry, and geochronology with evidences on
 644 geodynamic reconstructions of Neo-Tethys. *Gondwana Research* 22, 377-397
 645 Hooper, P., Widdowson, M., Kelley, S., 2010. Tectonic setting and timing of the final Deccan
 646 flood basalt eruptions. *Geology* 38, 839-842
 647 Hubert-Ferrari, A., King, G., van der Woerd, J., Villa, I., Altunel, E., Armijo, R., 2009. Long
 648 term evolution of the North Anatolian Fault : new constraints from its eastern termination.
 649 From: VAN HINSBERGEN, D. J. J., EDWARDS, M. A. & GOVERS, R. (eds) *Collision*
 650 *and Collapse at the Africa–Arabia–Eurasia Subduction Zone*. The Geological Society,
 651 London, Special Publications, 311, 133–154. DOI: 10.1144/SP311.5
 652 Hutchison, I., Loudon, K. E., White, R.S., von Herzen, R. P., 1981. Heat flow and age of the
 653 Gulf of Oman. *Earth and Planetary Science Letters* 56, 252-262.

654 Immenhauser A., 1996. Cretaceous sedimentary rocks on the Masirah Ophiolite (Sultanate of
655 Oman): evidence for an unusual bathymetric history. *Journal of the Geological Society of*
656 *London* 153, 539-551.

657 Immenhauser, A., Schreurs, G., Gnos, E., Oterdoom, H. W., Hartmann, B., 2000. Late
658 Palaeozoic to Neogene geodynamic evolution of the northeastern Oman margin.
659 *Geological Magazine* 137, 1-18.

660 Jagoutz, O., Royden, L., Holt, A., Becker, T., 2015. Anomalously fast convergence of India
661 and Eurasia caused by double subduction. *Nature Geoscience* 8, 475-478.

662 Jolivet, L., Faccenna, C., Agard, P., Frizon de Lamotte, D., Menant, A., Sternai, P., Guillocheau, F.,
663 2016. Neo-Tethys geodynamics and mantle convection : from extension to compression in Africa
664 and a conceptual model for obduction. *Canadian Journal of Earth Science* 53, 1190-1204.

665 Kakar, M. I., Kerr, A.C., Mahmood, K., Collins, A.S., Khan, M., McDonald, I., 2014. Supra-
666 subduction zone tectonic setting of the Muslim Bagh ophiolite, north-western Pakistan: Insights
667 from geochemistry and petrology. *Lithos* 202-203, 190-206.

668 Khan, S.D., Mahmood, K., Casey, J.F., 2007. Mapping of Muslim Bagh ophiolite complex
669 (Pakistan) using new remote sensing and field data. *Journal of Asian Earth Sciences* 30,
670 333-343

671 Kukowski, N., Schillhorn, T., Huhn, K., von Rad, U., Husen, S., Flueh, E.R., 2001.
672 Morphotectonics and mechanics of the central Makran accretionary wedge off Pakistan.
673 *Marine Geology* 173, 1-19.

674 Le Pichon, X., A. M. C. Sengor, J. Kende, C. Imren, P. Henry, C. Grall, H. Karabulut, 2016.
675 Propagation of a strike-slip plate boundary within an extensional environment: the
676 westward propagation of the North Anatolian Fault. *Canadian Journal of Earth Sciences*
677 53, 1416-1439, <https://doi.org/10.1139/cjes-2015-0129>

678 Mahanjane, E.S., 2014. The Davie Fracture Zone and adjacent basins in the offshore
679 Mozambique Margin-A new insights for the hydrocarbon potential. *Marine and Petroleum*
680 *Geology* 57, 561-571.

681 Mahmood, K., Boudier, F., Gnos, E., Monié, P., Nicolas, A., 1995. $^{40}\text{Ar}/^{39}\text{Ar}$ dating of the
682 emplacement of the Muslim Bagh ophiolite, Pakistan. *Tectonophysics* 250, 169-181

683 Maia, M., S. Sichel, A. Briais, D. Brunelli, M. Ligi, N. Ferreira, T. Campos, B. Mougél, I.
684 Brehme, C. Hémond, A. Motoki, D. Moura, C. Scalabrin, I. Pessanha, E. Alves, A. Ayes,
685 and P. Oliveira (2016). Extreme mantle uplift and exhumation along a transpressive
686 transform fault, *Nat. Geosci.*, DOI:10.1038/NGEO2759

687 Matthews, K. J., Seton, M., Müller, R., 2012. A global-scale plate reorganization event at 105-100 Ma.
688 *Earth and Planetary Science Letters* 355-356, 283-298.

689 Matthews, K.J., Müller, R.D., Sandwell, D.T., 2015. Oceanic microplate formation records
690 the onset of India-Eurasia collision. *Earth and Planetary Science Letters*, 433, 204-214,
691 doi:10.1016/j.epsl.2015.10.040

692 Matthews, K.J., Maloney, K.T., Zahirovic, S., Williams, S.E., Seton, M., Müller, D.R., 2016.
693 Global plate boundary evolution and kinematics since the late Paleozoic. *Global and*
694 *Planetary Change* 146, 226-250.

695 McCall, G.J.H., 1997. The geotectonic history of the Makran and adjacent areas of southern
696 Iran. *Journal of Asian Earth Sciences* 15, 517-531.

697 Meckel, T.A., Coffin, M.F., Mosher, S., Symonds, P., Bernadel, G., Mann, P., 2003.
698 Underthrusting at the Hjort Trench, Australian-Pacific plate boundary: incipient
699 subduction? *Geochemistry, Geophysics, Geosystems* 12, doi:10.1029/2002GC000498

700 Merkouriev, S., Patriat, P., Sochevanova, N., 1996. Evolution de la dorsale de Carlsberg:
701 Evidence pour une phase d'expansion très lente entre 40 et 25 Ma (A18 à
702 A7). *Oceanologica Acta* 19, 1-13.

703 Minshull, T.A., Lane, C., Collier, J. S., Whitmarsh, R., 2008. The relationship between rifting
704 and magmatism in the northeastern Arabian Sea. *Nature Geoscience* 1, 463-467,
705 doi:10.1038/ngeo228

706 Molnar, P., Stock, J., 2009. Slowing of India's convergence with Eurasia since 20 Ma and its
707 implications for Tibetan mantle dynamics. *Tectonics* 28, TC3001, doi:
708 10.1029/2008TC002271

709 **MORRIS 2017**

710 Mouchot, N. 2009. Tectonique et sédimentation sur le complexe de subduction du Makran
711 pakistanais. PhD thesis, Univ. Cergy Pontoise, 208 pp.

712 Mountain, G. S., Prell, W. L., 1990. A multiphase plate tectonic history of the southeast
713 continental margin of Oman, *The Geology and Tectonics of the Oman Region*, edited by
714 Robertson, A. H. F., Searle, M. P. and Ries, A. C. Geological Society Special Publication
715 49, 725-743.

716 Müller, R.D., Seton, M., Zahirovic, S., Williams, S.E., Matthews, K.J., Wright, N.M., Shepard, G.E.,
717 Maloney, K.T., Barnett-Moore, N., Hosseinpour, M., Bower, D.J., Cannon, J., 2016. Ocean Basin
718 evolution and global-scale plate reorganization events since Pangea Breakup. *Annual Review of*
719 *Earth and Planetary Sciences* 44, 107-138, <https://doi.org/10.1146/annurev-earth-060115-012211>

720 Nicolas, A. and Boudier, F., 2017. Emplacement of Semail-Emirates ophiolite at ridge-trench
721 collision. *Terra Nova* 29, 127-134

722 Norton, I.O., Sclater, J.G., 1979. A model for the evolution of the Indian Ocean and the break-
723 up of Gondwanaland. *Journal of Geophysical Research* 84, 6803-6830.

724 Patriat, P., Achache, J., 1984. India-Eurasia collision chronology has implications for crustal
725 shortening and driving mechanism of plate. *Nature* 311, 615-621.

726 Patriat, P., Sloan, H., Sauter, D., 2008. From slow to ultraslow: a previously undetected event
727 at the Southwest Indian Ridge at ca. 24 Ma. *Geology* 36, 207-210, doi: 10.1130/G24270A.1

728 Peters, Tj., Mercolli, I., 1998. Extremely thin oceanic crust in the Proto-Indian Ocean:
 729 Evidence from the Masirah Ophiolite, Sultanate of Oman. *Journal of Geophysical Research*
 730 103, 677-689.

731 Platel, J. P., A. Berthiaux, J. Le Métour, M. Beurrier, J. Roger (1992), Geological map of
 732 Duqm and Madraka, Sultanate of Oman, sheet NE 40-03/07, scale 1: 250000, Oman
 733 Minist. of Pet. and Miner., Dir. Gen. of Miner., Muscat.

734 Pubellier, M., Girardeau, J., Tjashuri, J., 1999. Accretion history of Borneo inferred from the
 735 polyphase structural features in the Meratus Mountains. In: Metcalfe (Ed.), *Gondwana*
 736 *Dispersion and Asian Accretion-Final results of IGCP 321*. AA Balkema Publishers,
 737 Rotterdam, 141-160. Ravaut, P., Bayer, R., Hassani, R., Rousset, P., and Al-Yahya-ey, A., 1997,
 738 Structure and evolution of the northern Oman margin; gravity and seismic constraints over the
 739 Zagros-Makran-Oman collision zone: *Tectonophysics* 279, 253-280.

740 Ravaut, P., Carbon, D., Ritz, J.F., Bayer, R., Philip, H., 1998. The Sohar Basin, Western Gulf of
 741 Oman : description and mechanisms of formation from seismic and gravity data. *Marine and*
 742 *Petroleum Geology* 15, 359-377.

743 Robinet, J., Razin, P., Serra-Kiel, J., Gallardo-Garcia, A., Leroy, S., Roger, J., Grelaud, C.,
 744 2013. The Paleogene pre-rift to syn-rift succession in the Dhofar margin (northeastern gulf
 745 of Aden): stratigraphy and depositional environments. *Tectonophysics* 607, 1-16.

746 Rodriguez, M., Fournier, M., Chamot-Rooke, N., Huchon, P., Bourget, J., Sorbier, M.,
 747 Zaragosi, S., Rabaute, A., 2011. Neotectonics of the Owen Fracture Zone (NW Indian
 748 Ocean): structural evolution of an oceanic strike-slip plate boundary. *Geochemistry*
 749 *Geophysics Geosystems* 12, doi:10.1029/2011GC003731.

750 Rodriguez, M., Chamot-Rooke, N., Fournier, M., Huchon, P., Delescluse, M., 2013. Mode of
 751 opening of an oceanic pull-apart: The 20 °N Basin along the Owen Fracture Zone (NW
 752 Indian Ocean). *Tectonics* 32, 1-15, doi:10.1002/tect.20083.

753 Rodriguez, M., Chamot-Rooke, N., Huchon, P., Fournier, M., Delescluse, M., 2014a. The

754 Owen Ridge uplift in the Arabian Sea: implications for the sedimentary record of Indian
 755 monsoon in Late Miocene. *Earth and Planetary Science Letters* 394, 1-12, doi:
 756 10.1016/j.epsl.2014.03.011

757 Rodriguez, M., Chamot-Rooke, N., Huchon, P., Fournier, M., Lallemand, S., Delescluse, M.,
 758 Zaragosi, S., Mouchot, N., 2014b. Tectonics of the Dalrymple Trough and uplift of the
 759 Murray Ridge (NW Indian Ocean). *Tectonophysics*, 636, 1-17, doi:
 760 10.1016/j.tecto.2014.08.001

761 Rodriguez, M., Huchon, P., Chamot-Rooke, N., Fournier, M., Delescluse, M., François, T.
 762 2016. Tracking the Paleogene India-Arabia plate boundary. *Marine and Petroleum*
 763 *Geology*, 72, 336-358, doi:10.1016/j.marpetgeo.2016.02.019

764 Rodriguez, M., Fournier, M., Chamot-Rooke, N., Huchon, P., Delescluse, M., 2018. The
 765 geological evolution of the Aden-Owen-Carlsberg triple junction (NW Indian Ocean) since
 766 the Late Miocene, *Tectonics*, 37, 1552-1575, <https://doi.org/10.1029/2017TC004687>

767 Rohr, K.M.M., 2015. Plate Boundary Adjustments of the Southernmost Queen Charlotte
 768 Fault. *Bulletin of the Seismological Society of America*, 105, doi:10.1785/0120140162

769 Rollinson, H., 2017. Masirah-The other Oman ophiolite: a better analogue for mid-ocean
 770 ridge processes? *Geoscience Frontiers* 8, 1253-1262

771 Royer, J. Y., Chaubey, A. K., Dymant, J., Bhattacharya, G. C., Srinivas, K., Yateesh, V.,
 772 Ramprasad, T., 2002. Paleogene plate tectonic evolution of the Arabian and Eastern
 773 Somali basins, in *The Tectonic and Climatic Evolution of the Arabian Sea Region*, edited
 774 by P. D. Clift et al., *Geological Society Special Publication* 195, 7-23.

775 Sarwar, G., 1992. Tectonic setting of the Bela Ophiolites, southern Pakistan. *Tectonophysics*
 776 207, 359-381.

777 Searle, M., Cox, J., 1999. Tectonic setting, origin, and obduction of the Oman ophiolite,
 778 *Geological Society of America bulletin* 111, 104-122.

779 Searle, M.P., Noble, S.R., Cottle, J.M., Waters, D.J., Mitchell, A.H.G., Tin Hlaing,
 780 Horstwood, M.S.A., 2007. Tectonic evolution of the Mogok metamorphic belt,
 781 Burma(Myanmar) constrained by U-Th-Pb dating of metamorphic and magmatic rocks.
 782 *Tectonics* 26, TC3014, doi:10.1029/2006TC002083

783 Seton, M., Müller, R.D., Zahirovic, S., Gaina, C., Torsvik, T., Shepard, G., Talsma, A.,
 784 Gurnis, M., Turner, M., Maus, S., Chandler, M., 2012. Global continental and ocean basin
 785 reconstructions since 200 Ma. *Earth Sciences Reviews* 113, 212-270,
 786 doi:10.1016/j.earscirev.2012.03.002.

787 Shackleton, R. M., Ries, A. C., 1990. Tectonics of the Masirah fault zone and eastern Oman.
 788 In *The Geology and Tectonics of the Oman Region*, eds A. H. F. Robertson, M. P. Searle,
 789 and A. C. Ries. Geological Society Special Publication 49, 715-24.

790 Schlich, R., 1982. The Indian Ocean: aseismic Ridges, spreading centres and basins, In:
 791 Nairn, A.E.M., Stehli, F.G. (Eds.), *The Ocean Basins and Margins*. Plenum Press, New
 792 York, 51-147.

793 Schreurs, G., Immenhauser, A., 1999. West-northwest-directed obduction of the Batain group
 794 on the eastern Oman continental margin at the Cretaceous-Tertiary boundary. *Tectonics*
 795 18, 148-60.

796 Shipboard Scientific Party, 1974a. Site 235, In: Fisher, R.L., Bunce, E.T. et al., DSDP
 797 Init.Repts, leg 24. doi:10.2973/dsdp.proc.24.1974

798 Shipboard Scientific Party 1974b, Site 222. In: Whitmarsh, R.B., Weser, O.E., Ross, D.A.
 799 (Eds.), DSDP Init. Repts, leg 23 <http://dx.doi.org/10.2973/dsdp.proc.23.106>.

800 Shipboard Scientific Party 1989, Site 731. In: Prell, W.L., Niitsuma, N., et al. (Eds.), *Proc.*
 801 *ODP, Init.Repts.*, 117. College Station, TX (Ocean Drilling Program).

802 Socquet, A., Pubellier, M., 2005. Cenozoic deformation in Western Yunnan (China-Myanmar
 803 border), *Journal of Asian Earth Sciences*, 24, 495-511

804 Smit, J., Brun, J.-P., Cloething, S., Ben Avraham, Z., 2010. The rift like structure and asymmetry of
805 the Dead Sea Fault. *Earth and Planetary Science Letters* 290, 74-82.

806 Tapponnier, P., Mattauer, M., Proust, F., Cassaigneau, C., 1981. Mesozoic ophiolites, sutures,
807 and large-scale tectonic movements in Afghanistan. *Earth and Planetary Science Letters*
808 52, 355-371.

809 Tréhu, A.M., Scheidhauer, M., Rohr, K.M.M., Tikoff, B., Walton, M.A., Gulick, S.P.S.,
810 Roland, E., 2015. An abrupt transition in the mechanical response of the upper crust to
811 transpression along the Queen Charlotte Fault, *Bulletin of the Seismological Society of*
812 *America*, 105, doi:10.1785/0120159.

813 Treolar, P.J., Izatt, C.N., 1993. Tectonics of the Himalayan collision between the Indian Plate
814 and the Afghan Block: a synthesis. *Geological Society Special Publication*, 74, 69-87

815 van Hinsbergen, D.J.J., Steinberger, B., Doubrovine, P.V., Gassmöller, R., 2011. Acceleration
816 and deceleration of India-Asia convergence since the Cretaceous: Roles of mantle plumes
817 and continental collision. *Journal of Geophysical Research*, 116, B06101,
818 doi:10.1029/2010JB0085051

819 van Hinsbergen, D. J. J., Lippert, P. C., Dupont-Nivet, G., McQuarrie, N., Doubrovine, P. V.,
820 Spakman W., Torsvik, T. H., 2012. Greater India Basin hypothesis and a two-stage
821 Cenozoic collision between India and Asia. *Proceedings of the National Academy of*
822 *Sciences* 109, 7659-7664.

823 Wajzer, M.R., Barber, A.J., Hidayat, S., Suharsono, S., 1991. Accretion, collision and strike-
824 slip faulting: the Woyla group as a key to the tectonic evolution of North Sumatra. *Journal*
825 *of Southeast Asian Earth Sciences*, 6, 447-461

826 Wakabayashi, J., Dilek, Y., 2003. What constitutes ‘emplacement’ of an ophiolite?
827 Mechanisms and relationship to subduction initiation and formation of metamorphic soles.
828 *Geological Society of London, Special publications*, 218, 427-447.
829 <https://doi.org/10.1144/GSL.SP.2003.218.01.22>

830 Wyns, R., Béchenec, F., Le Métour, J., Roger, J., and Chevrel, S., 1992, Explanatory notes,
831 Geological map of Sur, sheet NF 40-08, scale 1:250'000, Muscat, Sultanate of Oman,
832 Ministry of Petroleum and Minerals, 1-103

833 Yateesh, V., Bhattacharya, G. C., Dymment, J., 2009. Early oceanic opening off Western India-
834 Pakistan margin : the Gop Basin revisited. *Earth and Planetary Science Letters* 284, 399-
835 408.

836 Zhuang, G., Najman, Y., Guillot, S., Roddaz, M., Antoine, P.-O., Métais, G., Carter, A.,
837 Marivaux, L., Solangi, S.H., 2015. Constraints on the collision and the pre-collision
838 tectonic configuration between India and Asia from detrital geochronology,
839 thermochronology, and geochemistry studies in the lower Indus basin, Pakistan. *Earth and*
840 *Planetary Science Letters* 432, 363-373.

Figure

Figure 1 Modified from Gaina et al. (2015)

Figure

Figure

Figure

Figure

★ Boundary of pre-Late Maastrichtian grabens on seismic lines

Figure c

Figure

Profile from Beauchamp et al., 1995

Figure

Figure

Figure

Seafloor age