Subtle effects of radiation on embryo development of the 3-spined stickleback

Adelaide Lerebours, Samuel Robson, Colin Sharpe, Jim Q. Smith

► To cite this version:

Adelaide Lerebours, Samuel Robson, Colin Sharpe, Jim Q. Smith. Subtle effects of radiation on embryo development of the 3-spined stickleback. *Chemosphere*, In press, 10.1016/j.chemosphere.2020.126005 . hal-02458327

HAL Id: hal-02458327

<https://hal.science/hal-02458327>

Submitted on 28 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Subtle effects of radiation on embryo development of the 3-spined stickleback

*^{1,2}*Adélaïde Lerebours, ³Samuel Robson, ²Colin Sharpe and ¹Jim T. Smith*

¹ School of Earth and Environmental Sciences, University of Portsmouth, Portsmouth, PO1 3QL, United Kingdom

² School of Biological Sciences, University of Portsmouth, Portsmouth, PO1 2DY, United Kingdom

³ Centre for Enzyme Innovation, University of Portsmouth, Portsmouth, PO1 2DT, United Kingdom

Keywords: radiation, fish, embryo.

Corresponding Author

* Adélaïde Lerebours, phone number: +33 (0)5 46 50 76 23, e-mail: adelaide.lerebours@univ-lr.fr;

Abstract

The Chernobyl and Fukushima nuclear power plant (NPP) accidents that occurred in 1986 and 2011 respectively have led to many years of chronic radiation exposure of wildlife. However, controversies remain on the dose threshold above which an impact on animal health occurs. Fish have been highly exposed immediately after both accidents in freshwater systems around Chernobyl and in freshwater and marine systems around Fukushima. The dose levels decreased during the years after the accidents, however, little is known about the effects of environmental low doses of radiation on fish health. The present laboratory study assesses the effects of an environmentally relevant dose range of radiation (0.1, 1 and 10 mGy/day) on early life stages of the 3-spined stickleback, *Gasterosteus aculeatus*.

The cardiac physiology and developmental features (head width, diameter, area) of high exposed embryos (10 mGy/day) showed no significant change when compared to controls. Embryos exposed to the medium and high dose were slower to hatch than the controls (between 166 and 195 hours post-fertilization). After 10 days of exposure (at 240 hours post-fertilization), larvae exposed to the high dose displayed comparable growth to controls. High-throughput sequence analysis of transcriptional changes at this time point revealed no significant changes in gene regulation compared to controls regardless of exposure conditions. Our results suggest that exposure of fish embryos to environmental radiation elicits subtle delays in hatching times, but does not impair the overall growth and physiology, nor the gene expression patterns in the recently hatched larvae.

Introduction

Serious nuclear accidents at both the Chernobyl Nuclear Power Plant (NPP) in April 1986 and the Fukushima NPP in March 2011 led to high levels of radiation exposure to wildlife. After the Chernobyl accident, the dose to fish found in the cooling pond was estimated at 10 mGy/day¹ and then rapidly declined due to the decrease of short life radionuclides and sedimentation processes².

Three decades after the Chernobyl NPP accident, the main radionuclides of concern are ⁹⁰Sr (a β emitter) and ¹³⁷Cs (a β and γ emitter) due to their long radioactive half-life (28 and 30 years respectively). Transuranium radioelements are also of concern due to their long radioactive half-life and high energetic alpha particle emission. However, their contribution to the total dose to fish at Chernobyl is very low³. In a highly contaminated lake called Glubokoye, located near the Chernobyl NPP, the total dose rate to perch (*Perca fluviatilis*) and roach (*Rutilus rutilus*) 30 years after the accident was up to 16 and 14 μ Gy/h respectively (less than 0.4 mGy/day)³.

³²P is a radioisotope of 14.26 days half-life that emits β particles of high energy (1.7 MeV) whose track length is a few millimetres in water. A β -particle directly ionizes molecules by the removal of an electron, whereas α particles (and X-ray) ionize molecules by generating a series of fast electrons (effectively β particles) after first hitting molecules, therefore, the use of a β -emitter is relevant to assess the effect of environmental radiation. In order to assess the effects of ionising radiation exposure on fish embryos under laboratory conditions, ³²P was selected due to its short half-life minimising the radioactive waste and its high energy β particles.

³²P is used in research laboratories, medical procedures and industry that may lead to discharge into freshwater systems. The Krasnoyarsk Mining and Chemical Industrial Complex in Russia discharged significant amounts of ³²P in the Yenisei river that were found to have accumulated in

fish, with ^{32}P activities of 2.2 Bq/L in water and 2900 Bq/kg in fish found at 200 km from the industrial site⁴. In the Columbia River contaminated by the cooling water from the Hanford plutonium reactors, the estimated level of ^{32}P in water between 1950 and 1971 varied from 0.1 to 7.7 Bq/L⁵ and the activity of ^{32}P in fish was varied from 0.7×10^3 to 22×10^3 Bq/kg between 1962 and 1964⁶. Waterborne uptake of ^{32}P was found to be negligible as compared to uptake via the dietary route in fish^{7,8}.

The 3-spined stickleback (*Gasterosteus aculeatus*) is a vertebrate model used in laboratory settings to assess the mechanistic effects of pollutants and in field surveys using multi-biomarker approaches⁹⁻¹⁰. The early stages of fish embryonic development are sensitive to ionising radiation¹¹.

Numerous studies describe the effects of acute exposure to radiation on fish embryos. However, much less is known about the effects of low doses of radiation in the environmental range of 0.1 to 10 mGy/day. Importantly, results from the literature on the effects of low doses on organism health differ¹². For instance, whilst some studies did not find evidence of radiation effects on populations of aquatic invertebrates¹³, fish³ or mammals¹⁴ at Chernobyl, others found adverse effects of radiation on the abundance of invertebrates¹⁵, birds¹⁶ and mammals¹⁷ at Chernobyl and Fukushima. Thus, the dose at which significant damage to wildlife populations occurs remains an open question¹².

Several studies have previously examined the effects of high doses of radiation (approximately 1000-fold above the environmental range) on the morphology of fish embryos. Exposure to acute dose of radiation induced various morphological abnormalities in embryos of zebrafish (*Danio rerio*)^{18,19,20,21}, mangrove killifish (*Kryptolebias marmoratus*)²² and medaka (*Oryzias latipes*)²³.

Numerous authors studied the effects of exposure to high doses of radiation on the hatching success of embryos but the results differed according to the type and nature of exposure. No difference in the percentage of embryos that hatched was observed in medaka embryos exposed to 35.42 mGy/h of tritiated water²³ and in zebrafish embryos exposed to 0.01 and 0.05 Gy of γ radiation²¹. However, other studies found significant effects on the hatching success. The percentage of zebrafish embryos that hatched decreased significantly after exposure to a total dose range of 0.1 - 10 Gy^{21,24}, and to a dose rate range of 0.3 - 2 Gy/day of γ radiation¹⁸. The percentage of mangrove killifish embryos that hatched decreased significantly after exposure to a total dose range of 2.5 – 10 Gy²². The hatching time is significantly delayed as compared to controls in zebrafish embryos exposed to a dose range of 0.1 - 10 Gy²⁴ and a dose rate range of 0.3 - 2 Gy/day of γ radiation¹⁸.

However, whilst many studies have examined the effects of high doses of radiation on hatching, fewer studies have examined hatching processes after exposure to lower doses, and no clear patterns of such dose effects has been observed. An acceleration of hatching was observed in zebrafish embryos exposed to a γ radiation dose of 10 and 1000 mGy/day, but no change was recorded after exposure to 1 and 100 mGy/day¹⁸. Hatching process appears to depend on the embryonic stage of exposure with some authors finding an acceleration of hatching in embryos exposed from 3 hpf (blastula stage) but not from 24 hpf (segmentation stage) to 1-1000 mGy/day of radiation¹¹.

Several studies investigated the effects of high doses of radiation on the cardiac physiology of fish embryos. Studies reported no effect¹⁹ or a decreased heart rate²⁰ in zebrafish embryos. To our knowledge, no studies in the literature have yet described the heart physiology of embryos exposed to environmentally relevant doses of radiation.

Several recent studies explored the effects of short-term exposure to radiation (from a few seconds to 96 hrs) on the transcriptional response of embryos. Gene expression changes were described in zebrafish embryos exposed to a total dose of 10 mGy to 100 mGy during 11 to 110 secs respectively (dose rate: 79.2 Gy/day), and the number of the differentially expressed genes was positively correlated to the dose¹⁹. Transcriptional response of genes involved in apoptosis²⁰ and DNA damage repair mechanisms^{20,25} were changed in zebrafish embryos exposed to 0.5 - 4 Gy of ⁵⁶Fe ion irradiation (Si et al. 2017)²⁰ or 9.6 and 96 mGy/day of γ radiation²⁵.

The present study is one of the few to assess of the effects of very low doses of radiation on fish embryo development under laboratory conditions, combining both high throughput molecular and biometric analyses. The aim was to investigate whether exposure to environmental low doses of radiation induces developmental, physiological and transcriptional changes in stickleback embryos.

Methods

Fish maintenance

Adult sticklebacks were kept in artificial reproduction conditions in order to generate the embryos for *in vitro* fertilization. The artificial water composition and experimental conditions were selected according to the OECD guideline for the testing of chemicals (CaCl₂: 294, MgSO₄: 123, NaHCO₃: 65, KCl: 6 mg/L, pH = 7.5, T°C = 19°C, photoperiod: 16h light/8h dark)²⁶. Four distinct couples were used to generate the embryos. The fertilization procedure was performed according to the protocol described in the OECD guideline for the testing of chemicals (test number 236). In total, 54 embryos were used for monitoring growth and physiology (27 controls and 27 exposed to the high dose), 108 for recording the hatching success (27 for each of the four

exposure conditions), 104 for next generation sequencing (NGS) analyses (Table S3) and 36 to check ^{32}P uptake in the chorion, prolarvae and larvae (Table S2).

Three hours after fertilization, each embryo was placed in an individual glass tube and waterborne exposed to 3 mL of a radioactive solution of 0.1, 1 and 10 mGy/day (or 4, 40 and 400 $\mu\text{Gy/h}$) using a solution of adenosine triphosphate labelled on the gamma phosphate group with ^{32}P (PerkinElmer). The final concentration of ATP in control, low and medium experimental tubes was adjusted to $1.1 \times 10^{-5} \mu\text{M}$ by addition of stable ATP. This ATP concentration is negligible as compared to the concentration found in a typical cell of 5 mM^{27} . The doses encompass the chronic low (L) (0.1 mGy/day) and medium (M) (1 mGy/day) doses encountered in the environment at Chernobyl 30 years after the accident, and the initial high dose (H) to fish after the accident (10 mGy/day). These doses span the Environmental Agency (EA) guidance level of 40 $\mu\text{Gy/h}$ (0.96 mGy/day) described in the radiological impacts on non- human species report of the EA in 2011²⁸. Control embryos were kept in clean artificial water. 2.5 mL of the water was renewed every 3 days. The embryos were exposed for 10 days and euthanized according to schedule 1 of the Home Office Licence (ASPA, 1986)²⁹ using tricaine methanesulfonate (Sigma).

Dose calculation and monitoring

Dose at the centre of the hemisphere was calculated from data in Berger (1971)³⁰ (Figure S1). Activity of ^{32}P was measured using a HIDEX 300SL liquid scintillation counter and associated MikroWin 2000 software (Version 4.43).

Developmental and physiological measurements

The morphological parameters were measured using a Zeiss axiozoom microscope and the Zen Pro software. The physiological parameter was measured using an optical microscope. The

growth of embryos was recorded at 4 dpf (days post-fertilization) by measuring the diameter (mm), area (mm²) and eye distance (mm). The hatching rate was calculated as the proportion of fish that hatched to the total number of fish from the same condition for each observation time. The cardiac physiology was assessed at 6 dpf by counting the heart beat rate (beats/min). The growth of larvae was recorded at 10 dpf by measuring the length (mm) and head width (mm) of the larvae through the glass tubes.

Next generation sequencing (NGS)

Differential expression analysis was conducted using NGS. For each of the 4 conditions (control, low dose, medium dose and high dose), 3 biological replicates of pools of 4 to 10 embryos aged of 10 days were used for NGS analyses (Table S3). Total RNAs were extracted using the High Pure RNA Tissue kit (Roche Diagnostics Ltd, West Sussex, U.K.) according to the manufacturer's instructions. RNA quality and integrity were evaluated using a bioanalyzer (Agilent, Santa Clara, USA). RNA integrity numbers ranged from 7.4 to 9.5 and showed low RNA degradation rates. NGS libraries for each pool were generated using the Illumina TruSeq mRNA library kit following the manufacturer's instructions. Libraries were sequenced using the Illumina HiSeq 2500 analyser, generating 125 base, paired-end sequences from libraries yielding an average of 33.4 ± 5.2 M paired reads per sample (Table S4). Quality control of raw fastq reads was conducted using fastQC (bioinformatics.babraham.ac.uk/projects/fastqc/). Reads were trimmed using the Trim Galore script to remove adapter sequences and low quality sequence tails (bioinformatics.babraham.ac.uk/projects/trim_galore/). Trimmed reads were mapped against the BROAD S1 *Gasterosteus aculeatus* 3-spined stickleback genome from Ensembl³⁰ using the STAR universal RNA seq aligner³¹ with parameters '--outSAMmultNmax 300'. Reads that

mapped uniquely to the genome in a proper pair with mapping quality score greater than 20 were used in downstream analyses.

Differential expression analyses

Differential expression analyses were conducted using the DESeq2 package³³ in R. Gene models were taken from Ensembl version 82³¹, and read counts over unique genes were quantified using the “summarizeOverlaps()” function in the GenomicAlignments package³⁴ using mode “Union”. Raw read counts were normalised using the regularised log transformation in DESeq2 for visualisation (Figure S2). P values were adjusted for multiple testing by using the Benjamini and Hochberg correction³⁵. To account for potential confounding effects of lineage due to a systematic difference in the pooling for replicate 3 compared to replicates 1 and 2 (Table S3), lineage was included as a covariate in the analysis. Significant differentially expressed genes were identified based on a fold-change of 2-fold or greater (up- or down-regulated) and an adjusted p-value less than 0.05. To avoid over-representing differential expression in low-abundance genes, significant genes were further filtered to remove those whose normalised expression was less than 1 for both the exposed and control groups. Gene ontology analysis was conducted using the clusterProfiler package³⁶.

Statistical analyses

Statistical analyses were performed using R version 3.1.2. After satisfying the assumptions of the normal distribution of the residuals, linear models were used to assess the potential differences. If the normality of the residuals wasn't respected a Kruskal-Wallis rank test was applied. A Fisher exact test was applied to assess any difference between hatching success at different times post-fertilization. When significant, post-hoc tests were performed and a Bonferroni correction of the α error was applied.

Results

Dose monitoring

The mean activity of ^{32}P solutions measured during the 10 days exposure experiment were 0.7 ± 0.2 , 27 ± 3 , 253 ± 24 and 2588 ± 199 Bq/mL ($n = 12$) (Table S1) and in accordance with the targeted activity of 2500, 250, 25 and 0 Bq/mL respectively, in each glass tube. After 3 days of exposure, the activities measured in the chorion were 1 ± 0 Bq for controls and low exposure conditions, and 3 ± 2 and 42 ± 11 Bq for the medium and high exposure conditions (mean $\pm$ SD, $n = 3$). The activities measured in dechorionated embryos were 1 ± 0 Bq for control, low and medium exposure conditions and 3 ± 2 Bq for the high exposure condition (mean $\pm$ SD, $n = 3$). After 9 days of exposure, the activities measured in the larvae were 1 ± 0 for control, low and medium exposure conditions and 2 ± 0 Bq for the high exposure condition (mean $\pm$ SD, $n = 3$) (Table S2).

Developmental and physiological parameters

Radiation exposure to 10 mGy/day did not significantly affect the growth of embryos. After 4 days, exposed embryos were equivalent in size. The diameter, surface and head width of exposed embryos were 1.46 ± 0.01 mm, 1.68 ± 0.02 mm² and 0.85 ± 0.03 mm respectively. These measures did not significantly differ from controls (diameter: 1.46 ± 0.01 mm, $p = 0.65$, area: 1.67 ± 0.02 mm², $p = 0.64$ and eye distance: 0.85 ± 0.03 mm, $p = 0.17$) (Table 1). After 6 days, the cardiac physiology of exposed embryos was not disturbed by exposure to radiation. The mean heart beats rate of exposed embryos were equal to 163.2 ± 12 beats/min and did not significantly differ from controls (150.0 ± 3.1 beats/min, $p = 0.22$) (Table 1). After 10 days, the development between exposed and control larvae remained similar. The length of exposed larvae

was 5.98 ± 0.06 mm and their head width 0.82 ± 0.01 mm. These values did not significantly differ from controls (length: 5.99 ± 0.05 mm, $p = 0.88$ and head width: 0.81 ± 0.01 mm, $p = 0.48$) (Table 1).

During the hatching process, embryos exposed to 1 (M) and 10 (H) mGy/day were slower to hatch at 170 hpf (hours post-fertilization), with 22% and 24% of the embryos respectively displaying a delay in hatching ($p = 0.014$ and $p = 0.010$ respectively) (Figure 1). At 174 hpf, the embryos exposed to the H condition reached 90% of hatching (HT₉₀) [169-176], 6 hours later than the controls that hatched at 168 hpf [164-169] (based on 95% confidence interval overlap) (Figure 2). There was no delay observed for the embryos exposed to L (HT₉₀: 171 hpf [168-173]) and M (HT₉₀: 172 hpf [166-175]) conditions as compared to the controls (Figure 2). Eventually, at 195 hpf, no significant difference in hatching success was observed between conditions ($p = 0.058$) (Figure 1) and the hatching percentage reached 100% [88-100] for the control embryos and 92%; 96% and 93% for the embryos exposed to 0.1 (L), 1 (M) and 10 (H) mGy/day conditions respectively (Figure 1).

Differential expression analysis

Analysis of the NGS data quality identified these data as showing excellent base calling qualities and post-filtering mapping rates of approximately 90% were seen throughout (Table S4). Following gene abundance identification, principal component analysis identified little difference between the dosage treatments in these data (Figure 3). A batch effect, resulting from the parentage of individuals pooled in the different replicates was found (Table S3). This batch effect was incorporated into the model for differential expression analysis.

Differential expression analysis was performed to identify genes whose expression was significantly deregulated following radiation exposure by comparing each of the three dosed

treatments against a control treatment as described. No significant change in gene expression was observed, with only a single gene showing significant differential expression in the low dosage after filtering (Figure S2).

Data availability

The RNA sequencing data can be obtained from ArrayExpress (<https://www.ebi.ac.uk/arrayexpress/>) with the accession number E-MTAB-7872.

Discussion

In the present study, embryo mortality was below 12% in each condition, in the range of what is considered as normal in studies using zebrafish embryos. In the literature, zebrafish embryos survival was reduced after exposures to higher doses such as 100 mGy¹⁹ and after 2 and 4 Gy of exposure to ⁵⁶Fe²⁰.

During the hatching process, embryos exposed to 1 mGy/day (M) and 10 mGy/day (H) conditions displayed reduced hatching successes, by 22 and 24% respectively. No change was observed in embryos exposed to the lowest dose 0.1 mGy/day (L). At the end of the hatching process, no significant difference on the hatching success of embryos was noticed between controls and exposure conditions with percentages reaching 92% (L), 96% (M) and 93% (H).

Exposure to different dose and nature of radiation can accelerate hatching. For instance, zebrafish embryos exposed to 9.6 mGy/h for 65h³⁶, 0.3 to 2 Gy/day¹⁸, 1-1000 mGy/day¹¹ and to an X-ray dose of 25 mGy³⁷ hatched earlier than control embryos. The embryonic stage at which radiation exposure occurs, appears to have consequences on the hatching sensitivity of embryos. For instance, mangrove killifish embryos displayed a higher sensitivity when exposed at an early

stage to 2.5, 5, 7.5, and 10 Gy of γ radiation. In addition, the hatching success was significantly decreased in embryos that were exposed early (10.5 hpf) for all doses²².

Other experiments showed that exposure to radiation induced a delay in hatching. Exposure to an X-ray dose above 25 mGy delayed hatching zebrafish embryos³⁸. Waterborne exposure to 20 and 100 $\mu\text{g/L}$ of ^{233}U induced a 12h delay in hatching time (HT_{50} : 59[54-66] and HT_{50} : 59[53-68] respectively), as compared to controls (HT_{50} : 47[45-48])³⁹. In our study, exposure to waterborne ^{32}P induced a 6h delay between embryos exposed to the highest dose, 10 mGy/day (H), (HT_{90} : 174 hpf [169-176]) as compared to controls (HT_{90} : 168 hpf [164-169]). A study performed on zebrafish embryos did not observe any modification of the hatching success of embryos exposed to a radiation dose range of 1, 2, 5 and 10 Gy⁴⁰.

During hatching, biochemical and behavioural process are synchronised to destroy the chorion⁴¹. Proteolytic enzymes and embryos movement contribute to the chorion disruption to allow hatching. Hatching may be delayed because of potential changes induced by radiation to those enzymes, as evidenced by studies on the effects of copper on rainbow trout eggs⁴¹. The delay could also reflect a protective response to stress where the chorion would protect the embryos from external hazard. A similar delay has been reported in zebrafish embryos exposed to metals^{42,43}. The present data suggest that a delay in hatching is a good indicator of exposure to environmental low dose of radiation in laboratory settings, which is in agreement with the study by Bourrachot et al (2008)³⁹ that uses waterborne ^{233}U . A developmental delay of maturing fish eggs has also been observed in organism exposed in their natural environment. A delay of oocyte growth has been evidenced in perch exposed to a total dose rate of 10-16 $\mu\text{Gy/h}$ (0.2-0.4 mGy/day) in exposed lakes at Chernobyl, 30 years after the accident, and was correlated to the radiation dose³. The precise mechanism by which radiation induces this delay is unknown. At the

molecular level, a recent study found that transcriptional response of genes involved in the circadian clock was modulated in zebrafish larvae (at 96 hpf) exposed to 0.4 and 4 mGy/h of tritiated water²⁵.

In the present study, radiation exposure to 10 mGy/day did not significantly affect the growth of embryos (no difference of diameter, area and eye distance at 4 dpf) and larvae (no difference of head width and length at 10 dpf) respectively. Similarly, no deformity (short tail, spinal curve, absence of pigment, failed hatching) and no length difference as compared to controls was observed in embryos exposed to 9.6 mGy/h³⁷. A recent environmental study found that the length and Fulton condition index of perch and roach were similar between lakes, in addition, no malformation of gonads and oocytes was recorded revealing that fish were in good health in general³. Other studies found a tail detachment in zebrafish exposed to 20 µg/L of ²³³U but not in embryos exposed to 100 µg/L of ²³³U and a decrease of the body length for both exposures³⁹. A reduction of body length was also found in zebrafish embryos exposed to high dose of radiation (1 - 10 Gy)^{40,24}. Exposure to a γ radiation dose of 0.3 to 2 Gy/day induced morphological abnormalities (tail atrophy and trunk axis malformations) in zebrafish embryos¹⁸. Using an acute dose of radiation, deformities (including spinal curvature, pericardial cyst enlargement and thoracic cavity variation) were noticed from 0.1 Gy, and hatching was reduced from 0.05 Gy¹⁹. Malformations such as tail deformity, pericardial edema and spinal curve were found to increase in zebrafish embryos exposed to 2 and 4 Gy of ⁵⁶Fe ion irradiation²⁰ and 0.01 to 1 Gy of γ radiation²¹. Pericardial and yolk sac edema, curved notochord and thin caudal fin were observed in the hermaphroditic fish embryos exposed to a total dose range of 2.5 - 10 Gy of γ radiation²². Vertebral malformations were reported in medaka embryos exposed to 35.42 mGy/h of tritiated water²³. Only a few studies have looked at the head development. Freeman et al. (2014)⁴⁰ found

that exposure to high dose of radiation (10 Gy) reduced eye diameter and head length. However, this dose is higher than the environmental dose range used in the present study.

The physiology of 6 dpf embryos exposed to 10 mGy/day was not changed as compared to controls based on the heart beats count. This is in agreement with a few previous studies that did not observed any change in zebrafish embryos exposed to a total dose of 10 to 100 mGy for 10.9 s to 109 s respectively (dose rate of 79.2 Gy/day)¹⁹ or after exposure to 2 Gy of ⁵⁶Fe ion irradiation (dose rate: 0.5 Gy/min)²⁰. The present work represents one of the rare studies that have assessed this physiological criterion at environmental low doses.

Interestingly, no significant transcriptional changes were observed in 10 dpf larvae after exposure to all dose levels. This may be a result of genes involved in protective mechanisms already being activated before the larval stage to compensate for the negative effects of radiation exposure, but returning to normal levels in later stages of development. In a recent transcriptomic study assessing the effects of tritiated water on zebrafish embryos, it was suggested that the onset of an early protective mechanism against oxidative stress may not be observed at the larval stage of development (96 hpf)²⁵. Indeed, antioxidant defence mechanisms that are activated in embryos may lead to a decrease in lipid peroxidation³⁷ and a reduction in DNA damage⁴⁴.

Another hypothesis is that the environmental doses used in the present study may be too low for eliciting a differential gene expression change. The basal gene transcriptional levels may be sufficient for the larvae to account for the effects of radiation. Results from the literature indicate a change in gene expression for higher exposure. Kumar et al (2017)²⁴ found that exposure to 5 Gy of radiation induced transcriptional changes of *sox* genes involved in development. The expression of genes seemed dependent on the embryonic stage of development and the dose level. Transcriptional changes of genes involved in antioxidant defence were found in early stage

zebrafish embryos exposed to low doses, but not to high doses. These gene transcriptional levels were unchanged at a later stage for both exposure conditions²⁵. The environmental doses used in the present study and currently existing at Chernobyl may be too low to induce a significant oxidative stress⁴⁵ and subsequent DNA damage. No genotoxic effect was evidenced as measured by micronuclei in erythrocytes of perch, roach³ and catfish⁴⁶ exposed to environmental radiation at Chernobyl.

These data suggest that low levels of radiation exposure have a negligible effect on gene expression profiles and embryo growth and physiology but result in subtle delays to hatching times that does not affect the final numbers of fish that hatched. The current levels of environmental radiation at NPP sites are therefore unlikely to negatively impact embryonic development of future offspring. These results support the findings from a previous large-scale environmental study that found a delay in the maturation of perch eggs, and that fish were otherwise in good general health³. Moreover, these results corroborate other environmental studies led at Chernobyl on aquatic macro-invertebrate development and physiology^{45,46}. Finally, this laboratory study is important as it provides environmentally relevant data to refine the current thresholds for which an effect is observed.

Acknowledgments

Authors are grateful to Neil Fuller and Graham Malyon for their help with fish maintenance. This work was completed as part of the TREE (Transfer-Exposure-Effects) consortium under the RATE programme (Radioactivity and the Environment), funded by the Environment Agency and Radioactive Waste Management Ltd (NERC grant NE/L000393/1).

References

- (1) Kryshev, A. I. Modelling the Accidental Radioactive Contamination and Assessment of Doses to Biota in the Chernobyl NPP's Cooling Pond. *Proceedings of the topical meeting of International Union of Radioecologists* **1998**, No. mol 1, 32–38.
- (2) *Chernobyl - Catastrophe and Consequences*; Smith, J. T., Beresford, N. A., Eds.; Springer-Praxis books in environmental sciences; Springer [u.a.]: Berlin, 2005.
- (3) Lerebours, A.; Gudkov, D.; Nagorskaya, L.; Kaglyan, A.; Rizewski, V.; Leshchenko, A.; Bailey, E. H.; Bakir, A.; Ovsyanikova, S.; Laptev, G.; et al. Impact of Environmental Radiation on the Health and Reproductive Status of Fish from Chernobyl. *Environmental Science & Technology* **2018**, 52 (16), 9442–9450.
- (4) Vakulovskii, S. M.; Kryshev, A. I.; Tertyshnik, é. G.; Chumichev, V. B.; Shishlov, A. E.; Savitskii, Yu. V.; Kudinov, K. G. ³² P Accumulation in Fish in the Enisei River and Reconstruction of the Irradiation Dose to the Public. *Atomic Energy* **2004**, 97 (1), 502–509.

- 385 (5) Walters, W. H.; Richmond, M. C.; Gilmore, B. G. Reconstruction of Radioactive
386 Contamination in the Columbia River: *Health Physics* **1996**, *71* (4), 556–567.
387
- 388 (6) Honstead, J. F.; Brady, D. N. The Uptake and Retention of ^{32}P and ^{65}Zn from the
389 Consumption of Columbia River Fish. *Health Phys* **1967**, *13* (5), 455–463.
390
- 391 (7) Winpenny, K.; Knowles, J. F.; Smith, D. L. The Uptake of Radioactive Phosphorus by
392 Brown Trout (*Salmo Trutta* L.) from Water and Food. *Journal of Environmental*
393 *Radioactivity* **1998**, *38* (2), 211–221.
394
- 395 (8) Smith, J. T.; Bowes, M. J.; Cailes, C. R. A Review and Model Assessment of ^{32}P and ^{33}P
396 Uptake to Biota in Freshwater Systems. *Journal of Environmental Radioactivity* **2011**, *102*
397 (4), 317–325.
398
- 399 (9) Katsiadaki, I.; Scott, A. P.; Mayer, I. The Potential of the Three-Spined Stickleback
400 (*Gasterosteus Aculeatus* L.) as a Combined Biomarker for Oestrogens and Androgens in
401 European Waters. *Marine Environmental Research* **2002**, *54* (3–5), 725–728.
402
- 403 (10) Sanchez, W.; Katsiadaki, I.; Piccini, B.; Ditché, J.-M.; Porcher, J.-M. Biomarker
404 Responses in Wild Three-Spined Stickleback (*Gasterosteus Aculeatus* L.) as a Useful Tool
405 for Freshwater Biomonitoring: A Multiparametric Approach. *Environment International*
406 **2008**, *34* (4), 490–498.

407

408 (11) Simon, O.; Massarin, S.; Coppin, F.; Hinton, T. G.; Gilbin, R. Investigating the
409 Embryo/Larval Toxic and Genotoxic Effects of γ Irradiation on Zebrafish Eggs. *Journal of*
410 *Environmental Radioactivity* **2011**, *102* (11), 1039–1044.

411

412 (12) Smith, J. Field Evidence of Significant Effects of Radiation on Wildlife at Chronic Low
413 Dose Rates Is Weak and Often Misleading. A Comment on “Is Non-Human Species
414 Radiosensitivity in the Lab a Good Indicator of That in the Field? Making the Comparison
415 More Robust” by Beaugelin-Seiller et Al. *Journal of Environmental Radioactivity* **2019**.

416

417 (13) Murphy, J. F.; Nagorskaya, L. L.; Smith, J. T. Abundance and Diversity of Aquatic
418 Macroinvertebrate Communities in Lakes Exposed to Chernobyl-Derived Ionising
419 Radiation. *Journal of Environmental Radioactivity* **2011**, *102* (7), 688–694.

420

421 (14) Deryabina, T. G.; Kuchmel, S. V.; Nagorskaya, L. L.; Hinton, T. G.; Beasley, J. C.;
422 Lerebours, A.; Smith, J. T. Long-Term Census Data Reveal Abundant Wildlife
423 Populations at Chernobyl. *Current Biology* **2015**, *25* (19), R824–R826.

424

425 (15) Moller, A. P.; Mousseau, T. A. Reduced Abundance of Insects and Spiders Linked to
426 Radiation at Chernobyl 20 Years after the Accident. *Biology Letters* **2009**, *5* (3), 356–359.

427

- (16) Møller, A. P.; Mousseau, T. A. Species Richness and Abundance of Forest Birds in Relation to Radiation at Chernobyl. *Biology Letters* **2007**, *3* (5), 483–486.
- (17) Møller, A. P.; Mousseau, T. A. Assessing Effects of Radiation on Abundance of Mammals and Predator–Prey Interactions in Chernobyl Using Tracks in the Snow. *Ecological Indicators* **2013**, *26*, 112–116.
- (18) Pereira, S.; Bourrachot, S.; Cavalie, I.; Plaire, D.; Dutilleul, M.; Gilbin, R.; Adam-Guillermin, C. Genotoxicity of Acute and Chronic Gamma-Irradiation on Zebrafish Cells and Consequences for Embryo Development. *Environmental Toxicology and Chemistry* **2011**, *30* (12), 2831–2837.
- (19) Zhao, W.; Hu, N.; Ding, D.; Long, D.; Li, S.; Li, G.; Zhang, H. Developmental Toxicity and Apoptosis in Zebrafish Embryos Induced by Low-Dose γ -Ray Irradiation. *Environmental Science and Pollution Research* **2019**, *26* (4), 3869–3881.
- (20) Si, J.; Zhou, R.; Song, J.; Gan, L.; Zhou, X.; Di, C.; Liu, Y.; Mao, A.; Zhao, Q.; Wang, Y.; et al. Toxic Effects of ^{56}Fe Ion Radiation on the Zebrafish (*Danio Rerio*) Embryonic Development. *Aquatic Toxicology* **2017**, *186*, 87–95.

- (21) Hu, M.; Hu, N.; Ding, D.; Zhao, W.; Feng, Y.; Zhang, H.; Li, G.; Wang, Y. Developmental Toxicity and Oxidative Stress Induced by Gamma Irradiation in Zebrafish Embryos. *Radiation and Environmental Biophysics* **2016**, *55* (4), 441–450.
- (22) Rhee, J.-S.; Kim, B.-M.; Kang, C.-M.; Lee, Y.-M.; Lee, J.-S. Gamma Irradiation-Induced Oxidative Stress and Developmental Impairment in the Hermaphroditic Fish, *Kryptolebias marmoratus* Embryo. *Environmental Toxicology and Chemistry* **2012**, *31* (8), 1745–1753.
- (23) Hyodo-Taguchi, Y.; Etoh, H. Vertebral Malformations in Medaka (Teleost Fish) after Exposure to Tritiated Water in the Embryonic Stage. *Radiation Research* **1993**, *135* (3), 400.
- (24) Praveen Kumar, M. K.; Shyama, S. K.; Kashif, S.; Dubey, S. K.; Avelyno, D.; Sonaye, B. H.; Kadam Samit, B.; Chaubey, R. C. Effects of Gamma Radiation on the Early Developmental Stages of Zebrafish (*Danio Rerio*). *Ecotoxicology and Environmental Safety* **2017**, *142*, 95–101.
- (25) Arcanjo, C.; Armant, O.; Floriani, M.; Cavalie, I.; Camilleri, V.; Simon, O.; Orjollet, D.; Adam-Guillermin, C.; Gagnaire, B. Tritiated Water Exposure Disrupts Myofibril Structure and Induces Mis-Regulation of Eye Opacity and DNA Repair Genes in Zebrafish Early Life Stages. *Aquatic Toxicology* **2018**, *200*, 114–126.

- (26) Lehninger, A. L.; Nelson, D. L.; Cox, M. M. *Lehninger Principles of Biochemistry*, 5th ed.; W.H. Freeman: New York, 2008.
- (27) ISO. *Water Quality — Determination of the Acute Toxicity of Waste Water to Zebrafish Eggs (Danio Rerio)*. International Standardization Organization, Geneva., 2007; Vol. ISO 15088.
- (28) Final assessment report. Radiological impacts on non- human species. UK EPRTM nuclear power plant design AREVA NP SAS and Electricité de France SA. Environment Agency, 2011.
- (29) *Guidance on the Operation of the Animals (Scientific Procedures) Act 1986*. 2014.
- (30) Berger, M. Distribution of Absorbed Doses around Point Sources of Electrons and Beta Particles in Water and Other Media. *Journal of Nuclear Medicine* **1971**, 12 (Supplement 5), 5–23.
- (31) Kersey, P. J.; Allen, J. E.; Allot, A.; Barba, M.; Boddu, S.; Bolt, B. J.; Carvalho-Silva, D.; Christensen, M.; Davis, P.; Grabmueller, C.; et al. Ensembl Genomes 2018: An Integrated Omics Infrastructure for Non-Vertebrate Species. *Nucleic Acids Research* **2018**, 46 (D1), D802–D808.

- 491 (32) Dobin, A.; Davis, C. A.; Schlesinger, F.; Drenkow, J.; Zaleski, C.; Jha, S.; Batut, P.;
492 Chaisson, M.; Gingeras, T. R. STAR: Ultrafast Universal RNA-Seq Aligner.
493 *Bioinformatics* **2013**, 29 (1), 15–21.
494
- 495 (33) Love, M. I.; Huber, W.; Anders, S. Moderated Estimation of Fold Change and Dispersion
496 for RNA-Seq Data with DESeq2. *Genome Biology* **2014**, 15 (12).
497
- 498 (34) Lawrence, M.; Huber, W.; Pagès, H.; Aboyoun, P.; Carlson, M.; Gentleman, R.; Morgan,
499 M. T.; Carey, V. J. Software for Computing and Annotating Genomic Ranges. *PLoS*
500 *Computational Biology* **2013**, 9 (8), e1003118.
501
- 502 (35) Benjamini, Y.; Hochberg, Y. Controlling the False Discovery Rate: A Practical and
503 Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*
504 *(Methodological)* **1995**, 57 (1), 289–300.
505
- 506 (36) Yu, G.; Wang, L.-G.; Han, Y.; He, Q.-Y. ClusterProfiler: An R Package for Comparing
507 Biological Themes Among Gene Clusters. *OMICS: A Journal of Integrative Biology* **2012**,
508 16 (5), 284–287.
509
- 510 (37) Hurem, S.; Gomes, T.; Brede, D. A.; Lindbo Hansen, E.; Mutoloki, S.; Fernandez, C.;
511 Mothersill, C.; Salbu, B.; Kassaye, Y. A.; Olsen, A.-K.; et al. Parental Gamma Irradiation

Induces Reprotoxic Effects Accompanied by Genomic Instability in Zebrafish (*Danio Rerio*) Embryos. *Environmental Research* **2017**, 159, 564–578.

- (38) Miyachi, Y.; Kanao, T.; Okamoto, T. Marked Depression of Time Interval between Fertilization Period and Hatching Period Following Exposure to Low-Dose X-Rays in Zebrafish. *Environmental Research* **2003**, 93 (2), 216–219.

- (39) Bourrachot, S.; Simon, O.; Gilbin, R. The Effects of Waterborne Uranium on the Hatching Success, Development, and Survival of Early Life Stages of Zebrafish (*Danio Rerio*). *Aquatic Toxicology* **2008**, 90 (1), 29–36.

- (40) Freeman, J. L.; Weber, G. J.; Peterson, S. M.; Nie, L. H. Embryonic Ionizing Radiation Exposure Results in Expression Alterations of Genes Associated with Cardiovascular and Neurological Development, Function, and Disease and Modified Cardiovascular Function in Zebrafish. *Frontiers in Genetics* **2014**, 5.

- (41) Westernhagen, H.; Dethlefsen, V.; Cameron, P.; Berg, J.; Fürstenberg, G. Developmental Defects in Pelagic Fish Embryos from the Western Baltic. *Helgoländer Meeresuntersuchungen* **1988**, 42 (1), 13–36.

- (42) Johnson, A.; Carew, E.; Sloman, K. The Effects of Copper on the Morphological and Functional Development of Zebrafish Embryos. *Aquatic Toxicology* **2007**, *84* (4), 431–438.
- (43) Fraysse, B.; Mons, R.; Garric, J. Development of a Zebrafish 4-Day Embryo-Larval Bioassay to Assess Toxicity of Chemicals. *Ecotoxicology and Environmental Safety* **2006**, *63* (2), 253–267.
- (44) Gagnaire, B.; Cavalié, I.; Pereira, S.; Floriani, M.; Dubourg, N.; Camilleri, V.; Adam-Guillermin, C. External Gamma Irradiation-Induced Effects in Early-Life Stages of Zebrafish, *Danio Rerio*. *Aquatic Toxicology* **2015**, *169*, 69–78.
- (45) Fuller, N.; Smith, J. T.; Nagorskaya, L. L.; Gudkov, D. I.; Ford, A. T. Does Chernobyl-Derived Radiation Impact the Developmental Stability of *Asellus Aquaticus* 30 Years On? *Science of The Total Environment* **2017**, *576*, 242–250.
- (46) Fuller, N.; Ford, A. T.; Nagorskaya, L. L.; Gudkov, D. I.; Smith, J. T. Reproduction in the Freshwater Crustacean *Asellus Aquaticus* along a Gradient of Radionuclide Contamination at Chernobyl. *Science of The Total Environment* **2018**, *628–629*, 11–17.

Figure and Table Legends.

Table 1. Morphological and physiological parameters recorded on 4 and 6 dpf embryos and 10 dpf larvae (mean $\pm$ SEM, $n = 27$).

Figure 1. Hatching success of embryos exposed to control, Low (0.1 mGy/day), Medium (1 mGy/day) and High (10 mGy/day) conditions, recorded at 166, 168, 172 and 195 dpf (% , IC95, $n = 27$).

Figure 2. Time necessary to reach 90% of hatching for embryos exposed to control, Low (0.1 mGy/day), Medium (1 mGy/day) and High (10 mGy/day) conditions ($n = 27$, IC95).

Figure 3. Principal component analysis showing the distribution of the 12 samples according to their gene expression profile over the top 500 genes based on their variance.

Table 1.

Morphological and physiological parameters	Exposure condition	
	Control	10 mGy/day
Embryos 4 dpf		
Diameter (mm)	1.46 ± 0.01	1.46 ± 0.01
Area (mm ²)	1.67 ± 0.02	1.68 ± 0.02
Eye distance (mm)	0.91 ± 0.02	0.85 ± 0.03
Embryos 6 dpf		
Heart beats (beats/min)	150.0 ± 3.1	163.2 ± 12
Larvae 10 dpf		
Head width (mm)	0.81 ± 0.01	0.82 ± 0.01
Length (mm)	5.99 ± 0.05	5.98 ± 0.06

Figure 1.

Figure 2.

626

627

628

629

630

631

632

633

634

635 **Figure 3.**

636

637

638

Supplemental Information.

Table S1. Table showing the ^{32}P activity measured for each exposure solutions (corresponding to the control, low: 0.1 mGy/day or 25 Bq/mL; medium: 1 mGy/day or 250 Bq/mL and high dose:

10 mGy/day or 2500 Bq/mL see Figure S1 for the dose calculation) renewed every 3 days (mean $\pm$ SD, $n = 3$). Measurements were performed using a HIDEX liquid scintillation counter.

Day	Activity (Bq/mL)			
	Control	Low	Medium	High
0	0.6 ± 0.1	22.6 ± 0.5	229.0 ± 6.8	2714.3 ± 155.3
3	0.8 ± 0.1	28.3 ± 1.2	251.5 ± 5.4	2533.9 ± 31.3
6	0.6 ± 0.1	31.4 ± 1.2	290.0 ± 3.2	2781.9 ± 14.7
9	0.8 ± 0.4	26.4 ± 0.1	240.0 ± 1.4	2320.5 ± 19.8

Table S2. Table showing the ^{32}P activity measured in the chorion and the dechorionated embryos after 3 days of exposure to control, medium and high dose (mean $\pm$ SD, $n = 3$) and in

the larvae after 9 days of exposure to control, medium and high dose (mean $\pm$ SD, $n = 3$).
Measurements were performed using a HIDEX liquid scintillation counter.

Tissue	Activity (Bq)			
	Control	Low	Medium	High
Chorion	1 $\pm$ 0	1 $\pm$ 0	3 $\pm$ 2	42 $\pm$ 11
Embryo	1 $\pm$ 0	1 $\pm$ 0	1 $\pm$ 0	3 $\pm$ 2
Larvae	1 $\pm$ 0	1 $\pm$ 0	1 $\pm$ 0	2 $\pm$ 0

Table S3. Table showing the composition of the 12 different pools used for next generation sequencing analyses.

Pool Id.	Treatment	Replicate (pool)	Number of embryos from each couple used to form the pool			
			Couple A	Couple B	Couple C	Couple D
1	Control	C01	3	3	3	1
2		C02	3	3	3	1
3		C03	1	0	0	5
4	Low	L11	3	3	3	1
5		L12	3	3	3	1
6		L13	2	2	2	4
7	Medium	M21	3	3	3	1
8		M22	3	3	3	1
9		M23	1	0	0	3
10	High	H31	3	3	3	1
11		H32	3	3	3	1
12		H33	0	0	0	4

Table S4. Number of reads per condition throughout sample processing. Raw and trimmed paired read counts are shown, along with the number of reads mapping to a unique locus

703 (“unique”) or multiple distinct loci (“multihits”). For multihits, the total number of hits and the
704 number of distinct reads are shown in brackets. Reads were filtered to keep only those mapping
705 uniquely in a proper pair with quality greater than 20 for downstream analyses.

706

SampleName	NumReads	Trimmed	Unique	Multihits	Filtered	Mapped
Control_Rep1	2x 31,910,348	2x 31,801,110	57,429,028	10,218,142 (3,078,701)	57,422,444	90.30%
Control_Rep2	2x 38,110,879	2x 37,980,438	68,172,637	12,151,094 (4,032,549)	68,164,538	89.70%
Control_Rep3	2x 38,495,511	2x 38,344,466	67,359,423	19,103,766 (5,234,609)	67,350,294	87.80%
Low_Rep1	2x 29,136,639	2x 29,035,323	52,487,727	8,766,747 (2,746,623)	52,481,744	90.40%
Low_Rep2	2x 41,686,354	2x 41,527,406	74,369,507	13,157,459 (4,353,014)	74,359,872	89.50%
Low_Rep3	2x 31,456,558	2x 31,330,688	56,101,672	9,229,210 (3,025,834)	56,093,802	89.50%
Medium_Rep1	2x 37,321,589	2x 37,195,851	66,814,130	11518101 (3805997)	66,807,080	89.80%
Medium_Rep2	2x 36,890,705	2x 36,763,379	65,732,666	11977756 (4093513)	65,725,120	89.40%
Medium_Rep3	2x 31,668,750	2x 31,548,680	55,669,106	14570879 (4057295)	55,662,160	88.20%
High_Rep1	2x 22,580,848	2x 22,504,598	40,755,145	6789854 (2074281)	40,750,928	90.50%
High_Rep2	2x 32,552,649	2x 32,441,090	58,867,494	9306884 (2943724)	58,861,186	90.70%
High_Rep3	2x 29,308,498	2x 29,172,425	51,473,727	13017738 (3565734)	51,465,422	88.20%

707

708

709

710

711

712

713

714

715

716

717 **Figure S1. Dose to fish egg.**

718 The mean dose to the egg calculated using the ERICA model is: 155 $\mu\text{Gy/h}$ per MBq/L in water.

719 The dose at the centre of the spherical egg calculated integrating the dose contribution from

720 activity in water at different distances x (see Figure below) using data in Berger (1967) is 164

721 $\mu\text{Gy/h}$ per MBq/L.

722 C : Activity (Bq/L)

723 A : Area of hemisphere

724 R_β : Absorbed dose rate per Bq at a given distance x (Gy/s)

725

726

727

728

729

730

731

732

733

Figure S2. MvA plots for the DeSeq2-based differential expression analysis for a) low exposure,
b) medium exposure and c) high exposure samples.

a)

b)

c)

742

743

744

745 **Figure S3.** Screen shot from the UCSC genome browser for the mapped read data for gene

746 ENSGACT00000015428.1, a homolog of POSTN.