Phenanthrene degradation using Fe(III)-EDDS photoactivation under simulated solar light: A model for soil washing effluent treatment

Yufang Tao, Marcello Brigante, Hui Zhang, Gilles Mailhot

► To cite this version:

Yufang Tao, Marcello Brigante, Hui Zhang, Gilles Mailhot. Phenanthrene degradation using Fe(III)-EDDS photoactivation under simulated solar light: A model for soil washing effluent treatment. Chemosphere, 2019, 236, pp.124366. 10.1016/j.chemosphere.2019.124366 . hal-02470626

HAL Id: hal-02470626

<https://hal.science/hal-02470626>

Submitted on 7 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Phenanthrene degradation using Fe(III)-EDDS photoactivation
under simulated solar light: A model for soil washing effluent
treatment**

Yufang Tao^{a,b}, Marcello Brigante^a, Hui Zhang^{b*}, Gilles Mailhot^{a*}

^a Université Clermont Auvergne, CNRS, SIGMA Clermont, Institut de Chimie de
Clermont-Ferrand, 63000 Clermont–Ferrand, France

^b Department of Environmental Engineering, School of Resources and Environmental Science,
Wuhan University, 430079, P. R. China

*Corresponding author:

Pr. Gilles Mailhot :

Present/permanent address: Institut de Chimie de Clermont-Ferrand, Université Clermont
Auvergne, 24 avenue Blaise Pascal, 63178 Aubière Cedex, France.

Tel : 00 33 (0)6 61 63 99 20

E-mail: gilles.mailhot@uca.fr

**Phenanthrene degradation using Fe(III)-EDDS photoactivation
under simulated solar light: A model for soil washing effluent
treatment**

Yufang Tao^{a,b}, Marcello Brigante^a, Hui Zhang^{b*}, Gilles Mailhot^{a*}

^a Université Clermont Auvergne, CNRS, SIGMA Clermont, Institut de Chimie de
Clermont-Ferrand, 63000 Clermont–Ferrand, France

^b Department of Environmental Engineering, School of Resources and Environmental Science,
Wuhan University, 430079, P. R. China

Abstract

In this work, for the first time, the nonionic surfactant polyoxyethylene-(20)-sorbitan
monooleate (Tween 80, C₆₄H₁₂₄O₂₆) aided soil washing effluent was treated by enhanced
activation of persulfate (PS) using Fe(III)-EDDS (EDDS: ethylenediamine-N, N-disuccinic
acid) complexes under simulated solar light irradiation. The performance of this system was
followed via the production and reactivity of radical species (SO₄^{•-}, HO[•], Cl₂^{•-}) and
degradation of phenanthrene (PHE) used as a model pollutant in soils. Different
physico-chemical parameters such as the concentration of reactive species and pH were
investigated through the PHE degradation efficiency. The second-order rate constants of the
reactions for generated radicals with PHE and Tween 80 in solution were identified through
competitive reaction experiments under steady-state conditions and application of nanosecond
laser flash photolysis (LFP) as well. A kinetic approach was applied to assess the selectivity

and reactivity of photo-generated radicals in aqueous medium in order to explain the observed degradation trends. This work proposes an innovative technology of management of soil washing solutions using Fe(III)-EDDS complexes and solar light for the activation of persulfate.

Keywords: Advanced oxidation processes, sulfate radical, iron complexes, soil depollution, photochemistry.

1. Introduction

Soil contamination is known to be a severe and ubiquitous problem worldwide. Specially, soil pollution caused by hydrophobic organic compounds (HOCs) have attracted more attention and been taken seriously by governments and enterprises due to the characters of high-toxicity and long-refractory of this kind of pollutant. Polycyclic aromatic hydrocarbons (PAHs) are typical representatives of HOCs, which are notable and distinguished by their high health risks of mutability, carcinogenicity and teratogenicity through long-term bioaccumulation in the environment (Yap et al., 2011; Li et al., 2012; Trellu et al., 2016). The characteristics of low solubility in water and poor mobility in soil of PAHs impede the elimination of this type of compound from soils. Comprehensive researches of *in-situ* technologies such as bioremediation (Tiehm et al., 1997), phytoremediation (Huang et al., 2004) and electrokinetic remediation (Reddy et al., 2006) technologies are available in the literature (Gan et al., 2009a). In Ye's (Ye et al., 1995) study of biodegradation of PAHs by *Sphingomonas paucimobilis*, after 16 h of incubation of pyrene, benz[a]anthracene (B[a]A), benzo[a]pyrene (B[a]P), benzo[b]fluoranthene (B[b]F) for each with 10mg/L, the degradation percentage was 1.3 ± 0.4 , 1.8 ± 0.1 , 9.6 ± 1 , 5.0 ± 0.6 , respectively. Lee's group (Lee et al., 2008) compared four native Korean plant species for phytoremediation of phenanthrene (PHE) and pyrene, the maximum removal efficiency of phenanthrene was >99%, for pyrene was 94% in planted soil

64 after 80 days of experiments. Maturi's group (Maturi and Reddy, 2006) investigated modified
65 cyclodextrin enhanced electrokinetic remediation of PHE and Nickel contaminated soil, the
66 optimal 10% removal efficiency of PHE was achieved with 1% cyclodextrin addition after
67 around 166 days elapsed duration. However, all these in-situ techniques present at least one
68 non-negligible disadvantages such as time-consuming treatment requirements, low
69 remediation efficiency and high economic input. In this case, *ex-situ* techniques represented
70 by surfactant aided soil washing processes appeal to more and more researchers. During the
71 past several decades, *ex-situ* soil washing (SW) and *in-situ* soil flushing (SF) techniques,
72 improved by application of surfactants as the extracting agents, have exhibited high
73 performances for PAH elimination from soils (Gan et al., 2009b; Mousset et al., 2014a).
74 Nonionic surfactant Polyoxyethylene (20) sorbitan monooleate (Tween 80), Dodecyl
75 polyglycol ether(Brij 35), Polyoxyethylene (10) isooctylphenyl ether (Triton X-100),
76 anionic surfactant sodium dodecyl sulfate (SDS) and sodium dodecyl benzene sulphonate
77 (SDBS) are commonly used in the literature of surfactant enhanced soil remediation (Mao et
78 al., 2015). Among these surfactants, the application of Tween 80 (TW80) was more extensive
79 and prominent due to its lower toxicity and higher solubility capacity (Cheng et al., 2008).
80 Except for the common application for PAHs, TW80 was supposed to be an excellent
81 alternative for numerous hydrophobic contaminants such as PAHs anthracene, fluoranthene,
82 pyrene and phenanthrene (Alcántara et al., 2008; Peng et al., 2011), Petroleum Hydrocarbons
83 (Huguenot et al., 2015), p-cresol (Rosas et al., 2011), Decabromodiphenyl ether (Zhou et al.,
84 2007), 4,4'-dichlorobiphenyl (Chu and Kwan, 2003), p-Nitrochlorobenzene (Guo et al., 2009)
85 and pesticides like Diazinon (Hernández-Soriano et al., 2012), DDT (Zheng et al., 2012). The
86 application of TW80 in the remediation of PAHs contaminated soil is especially ubiquitous
87 owing to its better solubilization characteristic than other surfactants. It was reported that
88 TW80 was efficient for solubilizing a plenty of PAHs such as Pyrene, PHE, Fluorene,

Acenaphthene, Naphthalene and TW80 exhibited superior capacity than other nonionic surfactants such as Tween 20, Triton X-100, Triton X-305, Triton X-405 (Zhu and Zhou, 2008). In another work, TW80 was suggested as the highest solubilization capacity than Brij35, TX100 and SDS for PHE on the basis of molar solubilization ratios (MSR) (Zhao et al., 2005). Nevertheless, as a consequence of these soil washing/flushing processes, a complicated mixture of waste liquids containing great loads of extractants, PAHs and other accompanying organics is generated and should be further treated to reduce the potential environment concern of discharge the effluent into environment. Owing to the complexity of the mixed components of SW solutions, the management and disposal are a tremendous challenge for environmental engineering (Rosas et al., 2013; Trellu et al., 2016). Soil washing effluent could be treated with biological method, adsorption techniques as well as Advanced oxidation process (AOPs). However, biological treatments bear the drawbacks of long duration and low efficiency (Gharibzadeh et al., 2016), while adsorption technologies possess potential risks since contaminants could not be completely decomposed (Ahn et al., 2008; Zhou et al., 2013; Li et al., 2014). In this case, AOPs including electrochemical oxidation (Gómez et al., 2010; Mousset et al., 2014c, b; Trellu et al., 2017a), ozone oxidation (Liu, 2018), photocatalytic oxidation (Zhang et al., 2011; Bai et al., 2019) have been developed to the most popular and common applied technology for PAHs and various surfactant involved soil washing effluent treatments. As reviewed in Table S1, AOPs based on radical oxidation mechanism could effectively decompose PAHs into various products during the processes of soil washing effluent treatment with PAHs dosage from 1 mg L^{-1} to around 200 mg L^{-1} and optimal surfactants ranging from 1 g L^{-1} to 10 g L^{-1} . Among multiple AOPs, Sulfate radicals based advanced oxidation processes (SR-AOPs) have been widely applied as effective and practicable techniques for remediation of contaminated waters and soils (Yen et al., 2011; Wacławek et al., 2017). Persulfate ($\text{S}_2\text{O}_8^{2-}$, PS) is an ideal alternative oxidant ($E_0 = 2.01\text{ V}$),

which can be activated to produce stronger sulfate radicals ($\text{SO}_4^{\bullet-}$) ($E_0 = 2.6 \text{ V}$) by various chemical or physical methods (Usman et al., 2012). In addition to its strong oxidation capacity, sulfate radical has revealed several advantages over other oxidants. $\text{SO}_4^{\bullet-}$ is more stable than hydroxyl radical ($\text{HO}^{\bullet}$) and exhibits a longer half-life time as well as a wider pH range for the application (Anipsitakis and Dionysiou, 2004).

Iron, one of the chemical activators of PS, is the most abundant transition metal present in the earth's crust and exists in a variety of forms in water such as soluble, colloidal and particulate forms. In natural water, insoluble iron-containing oxides account for a large proportion (Faust and Zepp, 1993), while dissolved iron is only found in a small percentage and most of them exists in the form of complex with organic ligands (Zhou et al., 2004). Polycarboxylates and (amino-)polycarboxylates like citrate, malonate, oxalate, EDTA, etc. can react with ferric ions (Fe^{3+}) to form stable and strong complexes and improve its solubility and stability in natural water (Chen et al., 2007). Moreover, such (amino-)polycarboxylate complexes are reported to have strong and fast photochemical reactivity under sunlight irradiation giving rise to oxidative species (Zhang et al., 2009). Those iron complexes such as Fe(III)-EDTA, Fe(III)-EDDS, Fe(III)-oxalate or Fe(III)-citrate have exhibited high efficiency as photo-catalysts and as persulfate and hydrogen peroxide activators for the elimination of a variety of contaminants (Manenti et al., 2015; Miralles-Cuevas et al., 2018; Soriano-Molina et al., 2019). Ethylenediamine-N,N-disuccinic acid (EDDS) is a structural isomer of EDTA, and [S,S]-EDDS, [R,R]-EDDS and [R,S/S,R]-EDDS are three stereo isomers of EDDS. Among those isomers, [S,S]-EDDS is recognized and employed as a environmentally safe and friendly substitution for EDTA for environmental restoration since it is easier biodegradable than EDTA (Nagaraju et al., 2007).

In this work, Fe(III)-EDDS complex was used to improve the PS photo-activation for the phenanthrene (PHE) degradation in soil washing solution. Main physico-chemical parameters

such as concentration of chemical species and pH were investigated using simulated solar radiation to generate the sulfate radical from PS. This work provides a novel proposal of application of Fe(III)-EDDS for soil washing effluent treatments.

2. Experimental Details

2.1 Chemicals

Tween 80 (TW80), phenanthrene (PHE) and ferric perchlorate ($\text{Fe}(\text{ClO}_4)_3 \cdot 9\text{H}_2\text{O}$) were obtained from Sigma-Aldrich. Ethylenediamine-N,N-disuccinic acid trisodium salt solution (35% in water) was purchased from Fluka. All the other used reagents and chemicals were of analytical grade and all the solutions employed in this work were prepared with purified water from Millipore Ultra-Pure System (18.2 M Ω cm). The stock solutions of Fe(III)-EDDS were a mixture of appropriate amount of newly prepared aqueous solutions of ferric perchlorate and EDDS at a stoichiometry ratio of 1:1.

2.2 Preparation of the synthetic SW solution

TW80 is one of the suggested and most applied enhancing-solubility nonionic surfactants that increases the solubility of HOCs due to its notable features of lower soil sorption capacity and critical micellar concentration (CMC), low toxic property as well as higher solubilization ability and higher cost-effectiveness (Mulligan et al., 2001). It is known that above CMC, the surface tension of HOCs varies with a much smaller slope and the solubility of HOCs is strongly improved (Trellu et al., 2016). The CMC of TW80 is equal to 0.016 g L⁻¹ (López-Vizcaíno et al., 2012) which is much lower compared to other surfactants. Due to the above advantages, TW80 was chosen as the aiding compound and its solubilization capacity of PHE was investigated.

2.3 Irradiation experiments and degradation kinetic

50 mL of samples were irradiated in a jacketed cylindrical Pyrex reactor connected to water

cooling system and placed in a rectangular wood box to prevent the interference from the external light sources. Four tubular lamps (Sylvania, F15W/350BL), used as irradiation sources, were fixed on the top of the wood box. The irradiation energy was determined employing an optical fiber with a charge coupled device (CCD) spectrophotometer (Ocean Optics USD 2000 + UV-vis) previously calibrated with a DH-2000-CAL reference lamp. The energy reaching the reactor determined between 300 and 400 nm ($1390 \mu\text{W cm}^{-2}$) was very closed to the typical solar emission in the same wavelength range (Bianco et al., 2016).

In Figure 1 emission spectrum of light sources and UV-vis absorption spectra of chemical species (carried out with a Cary 300 UV-visible spectrophotometer) were presented. An overlap between the emission spectrum of the lamps and the absorption spectrum Fe(III)-EDDS complex was highlighted. PHE absorbed in the ultraviolet region with two maxima at 252 nm and 295 nm. The main absorption region was located at $\lambda < 250$ nm for PS and $\lambda < 300$ nm for PHE in TW80 solution. Fe(III)-EDDS clearly exhibited a broad but gradual decrease absorption at the region of wavelengths up to approximately 400 nm and the overlapping absorption with the lamp emission spectrum from 300 nm and 400 nm as well.

All experiments were carried out at 293 ± 2 K and solutions were magnetically stirred with a magnetic bar to insure the homogenous mixture of the solution. At regular intervals, 0.5 mL of sample was withdrawn and analyzed by an UPLC-UV system (Alliance) equipped with a photodiode array detector. The eluent was a mixture of water and acetonitrile (30/70, v/v) with isocratic mode in a flow rate of 1 mL min^{-1} . The column was a Nucleodur 100-5 C18 of $150 \text{ mm} \times 4.6 \text{ mm}$ with particle size $5 \mu\text{m}$. The pseudo-first-order decay of PHE was determined using the following exponential equation:

$$\frac{[PHE]}{[PHE]_0} = \exp(-k't)$$

Where $[PHE]_0$ and $[PHE]$ are the initial and residual concentrations of PHE at time t , k' is the pseudo-first-order apparent rate constant (s^{-1}). The initial conversion rate of PHE (R_{PHE}) was calculated as $R_{PHE} (M s^{-1}) = k' \times [PHE]_0$. The retention time and the analysis of PHE were not modified by the presence of TW80. The pHs of the solutions were adjusted with 0.1 M of $HClO_4$ or NaOH solutions and no significant pH variation was observed during irradiation.

2.4 Laser flash photolysis

The determination of the second order rate constants were carried out with the nanosecond laser flash photolysis (LFP) apparatus from Applied Photophysics (LKS60). A 266 nm excitation was used to generate radicals from solution and apparatus description as well as competition kinetic approaches have been reported elsewhere and a brief description is given in the supplementary materials section (Huang et al., 2018).

3. Results and discussion

3.1 PHE degradation under various systems

The solubility of PHE was tested using different concentration of TW80 in water. The results reported in Figure SM1 indicate that PHE solubility increased with the concentration of TW80 and a maximum of PHE solubility was reached around 16 mg L^{-1} (corresponding to a concentration of $90 \text{ }\mu\text{M}$). This maximum of solubility of PHE was obtained from 0.5 g L^{-1} of TW80. This concentration of TW80 was then used to prepare PHE solution for all experiments.

The PHE degradation under simulated solar light at pH 3.5, using different mixture of Fe(III)–EDDS, Fe(III) and PS is reported in Figure 2. Control experiments showed that very

low degradation of PHE (less than 5%) occurred with the involvement of Fe(III)-EDDS complex and PS in the dark. Under irradiation, the direct photolysis (without PS and iron complexes) of PHE was negligible, less than 8% and addition of Fe(III)-EDDS (0.5 mM) not significantly changed the efficiency of PHE degradation after 150 min of irradiation. As a contrary, about 30% removal of PHE was being achieved in the presence of PS (5 mM) with UVA light due to the photogeneration of sulfate radicals that contributed to PHE degradation. Under UVA irradiation with addition of Fe(III) to PS solution, almost 35% of PHE removal was obtained under irradiation. So, although Fe(III) can be photo-reduced to Fe(II) (Brand et al., 1998) which is able to activate PS leading to the generation of sulfate radicals (Anipsitakis and Dionysiou, 2004; Liang et al., 2004), only 5% of PHE degradation was promoted compared to PS/UV system. In the system with replacement of Fe(III) by Fe(III)-EDDS (0.5 mM) to PS solution, the performance was much more pronounced and PHE removal was remarkably strengthened under irradiation, especially in the first 30 minutes. This difference can be explained by the lower photoreactivity of Fe(III) aquacomplexes in comparison with Fe(III)-EDDS at pH around 3.5 (Catastini et al., 2004; Huang, 2012) since Fe(II) formation from Fe(III) aquacomplexes is lower than from Fe(III)-EDDS at $\lambda > 300$ nm. Therefore, in the presence of PS, sulfate radical formation is more important with Fe(III)-EDDS (reactions 1 and 2):

Under sun-simulated irradiation in the presence of Fe(III)-EDDS, PHE kinetic exhibited two

different decay steps (Figure 2). In the first 30 mins of irradiation, Fe(III)-EDDS was almost completely degraded at pH 3.5 (Figure 3) to produce Fe(II) which represented the rate-limiting parameter for $\text{SO}_4^{\bullet-}$ formation. In the second step, after complete decomposition of Fe(III)-EDDS, iron remained in the form of Fe(III) aquacomplexes, and under the influence of UVA light, hydroxyl radical and Fe(II) could be generated as well (reaction 3) (Mailhot et al., 2002; Krýsová et al., 2003). In the presence of PS, Fe(II) led to Fe(III) again (reaction 2) and we were in the presence of the (photo)chemical cycle involving Fe(III) and Fe(II). In this system, the degradation of PHE was mainly attributed to the formation of $\text{SO}_4^{\bullet-}$ and $\text{HO}^{\bullet}$ through analogous reactions obtained with Fe(III) aquacomplexes in solution and the subsequent cycle of Fe(III)/Fe(II). In fact, the degradation kinetic after 30 minutes followed the same profile compared with those obtained where PS was mixed with Fe(III) aquacomplexes.

Same experiments conducted at pH 7.0 (Figure SM2) indicated that an enhancement of PHE degradation was still acquired with Fe(III)-EDDS, suggesting that the iron complex is photoactive also at higher pH (Wu et al., 2014). As a contrary, a strong inhibition of PHE degradation occurred with the dosing of Fe(III) aquacomplexes and PS. This effect could be ascribed to the precipitation of Fe(III) at $\text{pH} > 4.0$, resulting in absorption of light by iron particles and a potential screening effect for the photolysis of PS and also a very low photoreactivity of precipitated iron species.

What should be noted is on the contrary of many studies involving iron species, in our system UV/PS/Fe(III)-EDDS no significant effect of pH in the range 3.0 to 8.5 was obtained (Figure SM3). This observation was quite surprising, if we compare with former studies (Wu et al.,

2014; Wu et al., 2015) but could be due to the presence of TW80 which modified the reactivity of radical species (see following results).

3.2 Effect of Fe(III)-EDDS and PS concentrations

The effects of persulfate (PS) and Fe(III)-EDDS concentrations were investigated on the PHE degradation efficiency. PHE degradation involving different concentrations of Fe(III)-EDDS with 5 mM PS are reported in Figure 3A. An enhancement of PHE degradation, after 30 min of irradiation, more than 3 times was achieved with the Fe(III)-EDDS concentration increased from 0 to 0.5 mM (Figure 3B). However, the degradation efficiency of PHE decreased with the further increase of iron complex concentration. In fact, a scavenging effect of Fe(III)-EDDS and Fe(II) toward $\text{SO}_4^{\bullet-}$ was expected at higher concentrations (Peng et al., 2017; Yu et al., 2018). Considering that a second order rate constant of $9.9 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ has been determined for the reaction between sulfate radical and Fe(II) (E. et al., 1966), it was possible to argue that scavenging effect of ferrous ions, that were more stable in solution at pH 3.5, became significant at higher dosage of Fe(III)-EDDS. The maximum degradation of PHE (33%) after 30 mins was reached using 0.5 mM Fe(III)-EDDS, while after 150 minutes the highest degradation efficiency (55%) was obtained with 0.25 mM of Fe(III)-EDDS (Fig.3B). This shift of the maximum efficiency after 30 min was attributed to the higher load of Fe(II) and EDDS at higher concentration of Fe(III)-EDDS and so stronger scavenging effect of sulfate radicals.

To further explore the degradation performance of PHE in the system PS/Fe(III)-EDDS, the effect of persulfate concentration was investigated. As shown in Figure 4, PHE degradation efficiency was strongly promoted by increasing PS loading from 0 to 30 mM both for the photolysis of PS alone in solution and in the presence of Fe(III)-EDDS. The influence of Fe(III)-EDDS was much pronounced at higher concentration of persulfate. In the presence of Fe(III)-EDDS and after 150 min of irradiation, the enhancement of PHE degradation with 2

mM of PS was less than 2% while with 5 mM of PS the enhancement was around 18% and more than 50% with 15 and 30 mM of PS.

These results suggested that Fe(III)-EDDS had stronger impact at higher concentration of PS showing that the generation of sulfate radical through reaction between PS and Fe(II) was the main contribution of the efficiency of the system.

3.3. Effect of Fe(III)/EDDS ratio

According to previously reported works; the ratio of Fe(III) and EDDS concentrations represented a key parameter for the efficiency of the pollutants removal in water (Li et al., 2010). The concentration of PS was kept constant at 5 mM while various initial loads of Fe(III) were applied. As shown in Figure SM4, the initial degradation rate of PHE increased with a higher concentration of Fe(III). At 1 mM of Fe(III) the final elimination of PHE was higher than at 0.5 or 0.25 mM. This observation was resulted from the fact that at 1 mM of Fe(III), there were in solution 0.5 mM of Fe(III)-EDDS and 0.5 mM of Fe(III) aquacomplexes. The two iron species were photo-transformed into Fe(II) which could activate PS to generate sulfate radicals and moreover Fe(III) aquacomplexes generated photochemically also hydroxyl radicals. So in these conditions, a higher amount of sulfate and hydroxyl radicals were generated increasing the degradation of PHE. At concentrations of Fe(III) of 0.25 or 0.5 mM, the final removals of PHE were very close with a little higher efficiency at 0.25 mM. During the first step of reactions, Fe(III) was reduced into Fe(II) and EDDS was oxidized. However if there were some residual EDDS remaining in solution, Fe(II) species were easily re-oxidized to Fe(III) and combined with EDDS to generate Fe(III)-EDDS complex (Li et al., 2010). As a consequence, the formation of $\text{SO}_4^{\bullet-}$ and the degradation of PHE photoinduced by Fe(III)-EDDS complex could continue.

3.4 Effect of TW80 concentration.

To better explore the role of TW80 in this system, experiments with 0.5, 0.75 and 1 g L⁻¹ of

TW80 were carried out. The increase of the surfactant concentration from 0.5 to 1.0 g L⁻¹ inhibited significantly PHE degradation (Figure SM5A). In fact, when TW80 concentration increased to a factor of 1.5 and 2, the apparent rate constant of PHE degradation was reduced by a factor 1.8 and 3.3 (Figure SM5B). The negative effect acquired at higher TW80 concentrations could be ascribed to the competition between TW80 and PHE for the reactivity with photogenerated sulfate radicals. In fact, as reported in Table 1, a second order rate constant between SO₄^{•-} and TW80 ($k_{TW80,SO_4^{\bullet-}}$) of $4.6 \pm 0.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ has been determined (see SM for more details). At 0.5 g L⁻¹ of TW80 and considering the reactivity between SO₄^{•-} and Fe(III)-EDDS and PS (Neta et al., 1988; Bianco et al., 2017) reported in literature, we can argue that about 33 % of photogenerated SO₄^{•-} reacted with TW80 while only 7 % with PHE in solution. However, increasing the concentration of TW80 to 1 g L⁻¹, the values were respectively 49 and 5 %. Moreover, in previously works it has been suggested that PHE was trapped into the micelle core of the surfactant, leading to a lower availability toward oxidizing radicals (Trellu et al., 2016; Trellu et al., 2017b). The significant sulfate radical selectivity between TW80 and PHE clearly demonstrated that the concentration of TW80 was the major parameter determining the efficiency of oxidative treatment adopted for PHE removal.

3.5 Effect of chloride ions addition

Inorganic species such as chloride ions are widely present in soil, resulting in the presence of such ions into the effluent of the procedure of soil washing. However, the existence of chloride ions (Cl⁻) in wastewater has been reported to affect the degradation efficiency of contaminants during the advanced oxidation processes and more particularly in PS oxidation treatments (Huang et al., 2018; Zhang et al., 2018). Different roles of chloride ions (Cl⁻) are

323 reported in the literature on the efficiency of $\text{SO}_4^{\bullet-}$ and $\text{HO}^{\bullet}$ based oxidations. Generally, it is
 324 reported that Cl^- acts as a radical scavenger that substantially inhibits the efficiency of UV
 325 based activation of H_2O_2 and $\text{S}_2\text{O}_8^{2-}$ (Tsuneda et al., 2002). However, some researchers find
 326 that reactive species produced from chloride ions, chlorine and dichlorine radicals ($\text{Cl}^{\bullet}/\text{Cl}_2^{\bullet-}$),
 327 show high selectivity when react with organic pollutants and increase the efficiency of
 328 removal (Yang et al., 2014). Moreover, in other investigations it is suggested that Cl^- could
 329 react with $\text{SO}_4^{\bullet-}$ to generate $\text{HO}^{\bullet}$ in circumneutral and alkaline pH (Kiwi et al., 2000). In
 330 addition, the involvement of Cl^- is reported to modify the degradation mechanism using
 331 $\text{SO}_4^{\bullet-}$ based oxidation processes (Fang et al., 2012). As reported in Figure 5, complete
 332 degradation of PHE was achieved in 150 min of irradiation with 10 mM of Cl^- , while with 50
 333 mM or 100 mM of Cl^- , the same efficiency could be obtained within 60 min, demonstrating
 334 that the presence of higher dosages of Cl^- remarkably enhanced PHE degradation using
 335 Fe(III)-EDDS under UVA irradiation. This result was totally distinguished with other studies
 336 where Cl^- acted as an inhibitor or where its effect was negligible (Yang et al., 2016). To
 337 better understand the mechanism, for the first time, we determined the second order rate
 338 constants of $\text{Cl}_2^{\bullet-}$ with PHE ($k_{\text{PHE},\text{Cl}_2^{\bullet-}}$) and TW80 ($k_{\text{TW80},\text{Cl}_2^{\bullet-}}$) respectively at $4.6 \pm 0.3 \times 10^8$
 339 $\text{M}^{-1} \text{s}^{-1}$ and $7.1 \pm 0.1 \times 10^6 \text{M}^{-1} \text{s}^{-1}$ (table 1). This results suggested that second order rate constant
 340 between PHE and dichlorine radical anion ($k_{\text{PHE},\text{Cl}_2^{\bullet-}}$) was almost 65 times higher than the
 341 one between TW80 and the same radical ($k_{\text{TW80},\text{Cl}_2^{\bullet-}}$). Therefore, $\text{Cl}_2^{\bullet-}$ generated by Cl^-
 342 exhibited much more selectivity to PHE in our soil washing solution. Moreover, the form of
 343 kinetics demonstrated that reactivity between $\text{SO}_4^{\bullet-}$ and Cl^- played an important role to
 344 maintain PHE degradation, especially after 30 min of irradiation.

If we simplified our chemical system considering that generated radical species ($\text{HO}^\bullet$, $\text{SO}_4^{\bullet-}$, $\text{Cl}_2^{\bullet-}$) could react only with TW80 or PHE in solution, it was possible to explain, from a chemical kinetic point of view the impact of chloride on the system efficiency. In fact, in the absence of chloride ions mainly $\text{SO}_4^{\bullet-}$ was generated in solution through reaction (2) and photolysis of persulfate ions. Considering the concentration of chemical species (i.e. TW80 and PHE) and reactivity constant, about 82% of $\text{SO}_4^{\bullet-}$ were trapped by TW80. However, if we considered that all $\text{SO}_4^{\bullet-}$ was significantly converted into $\text{Cl}_2^{\bullet-}$ (about 90% using 100 mM of Cl^-), in this case, the stronger reactivity between $\text{Cl}_2^{\bullet-}$ and PHE compared to those determined for TW80 suggested that only 6% of $\text{Cl}_2^{\bullet-}$ react with the surfactant and 94% with PHE.

3.6 Radical species involvement

In order to clearly clarify the significance of the different radicals formed during the process and their roles in this system and so further elucidate the oxidation mechanism of PHE degradation, free radicals scavenging tests were carried out. Methanol (MeOH) was employed as scavenger for both $\text{HO}^\bullet$ ($k_{\text{HO}^\bullet, \text{MeOH}} = 9.7 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$) and $\text{SO}_4^{\bullet-}$ ($k_{\text{SO}_4^{\bullet-}, \text{MeOH}} = 1.0 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$), and tert-Butanol (t-But) has higher selectivity for reactivity with $\text{HO}^\bullet$ ($k_{\text{HO}^\bullet, \text{t-But}} = 3.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$) than with $\text{SO}_4^{\bullet-}$ ($k_{\text{SO}_4^{\bullet-}, \text{t-But}} = 8.4 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$) (Buxton et al., 1988; Neta et al., 1988). With the addition of 50 mM of tert-Butanol, PHE oxidation decreased (Figure.6), which revealed that $\text{HO}^\bullet$ was partly responsible for PHE oxidation. PHE removal was more attenuated with methanol addition indicating that $\text{SO}_4^{\bullet-}$ was also radical species involved in the transformation of PHE. The degradation efficiency of PHE reduced from 48% (without scavenger) to 40% and 23% with tert-Butanol and methanol, respectively.

With the different rate constants and concentrations of chemicals (TW80 and PHE) present in solution, we can deduce that, without scavengers about 18% of photogenerated $\text{SO}_4^{\bullet-}$ and 13% of $\text{HO}^\bullet$ reacted with PHE. However, i) In the presence of tert-Butanol (50 mM), $\text{HO}^\bullet$ radicals were almost completely trapped, only 0.3% of $\text{HO}^\bullet$ still reacted on PHE. There was no effect on the percentage of $\text{SO}_4^{\bullet-}$ reacting with PHE. The addition of tert-Butanol resulted in 40% PHE degradation instead of 48%. ii) With the addition of methanol (50 mM), about 1% of $\text{HO}^\bullet$ and 15% of $\text{SO}_4^{\bullet-}$ still reacted with PHE in solution. In this condition PHE concentration decreased by 25%.

As a consequence, in the presence of tert-Butanol almost no more $\text{HO}^\bullet$ reacted on PHE and the percentage of PHE degradation decreased by 8%, while with the dosing of methanol only 1% of $\text{HO}^\bullet$ and 15% of $\text{SO}_4^{\bullet-}$ reacted on PHE and the percentage of PHE degradation decreased by 25%. With these data, we can evaluate that 1% of $\text{SO}_4^{\bullet-}$ contributed to 6% of PHE degradation and 1% of $\text{HO}^\bullet$ contributed only to 0.6% of PHE degradation. In another word $\text{SO}_4^{\bullet-}$ radicals contributed 10 times more than the $\text{HO}^\bullet$ for the degradation of PHE in the presence of surfactant TW80. This result was not in agreement with the rate constant which was 1.4 times higher with $\text{HO}^\bullet$ than with $\text{SO}_4^{\bullet-}$ (table 1). Thus, this result highlighted the surrounding environment of the pollutant for the efficiency of the radical species. PHE in surfactant presented an opposite reactivity with $\text{HO}^\bullet$ and $\text{SO}_4^{\bullet-}$ if you compared with PHE alone in aqueous solution.

4 Conclusion

A novel application of Fe(III) complexes was studied to improve the photochemical activation of persulfate for treatments of soil washing effluent. Experimental parameters such as

Fe(III)-EDDS concentration, PS concentration, TW80 concentration exhibited great effects on the PHE photodegradation in this system. The PHE degradation included two stages and the first period of Fe(III)-EDDS decomposition was the rate-limiting step. Quenching experiments proved that $\text{SO}_4^{\bullet-}$ played a more important role than $\text{HO}^{\bullet}$ on PHE degradation more particularly in the presence of surfactant which completely modified the reactivity of radical species. Moreover, we provided the evidence that the presence of chloride ions could significantly improve the PHE removal in this system. The higher selectivity of $\text{Cl}_2^{\bullet-}$ between PHE and TW80 was responsible for the efficiency increase.

Acknowledgments

We gratefully acknowledge the Ministry of Education of the PR of China for providing financial support for Yufang TAO to stay at the Institute of Chemistry of Clermont-Ferrand and Clermont Auvergne University in France. Authors acknowledge financial support from the Region Council of Auvergne, from the “Fédération des Recherches en Environnement” through the CPER “Environment” founded by the Region Auvergne, the French government, FEDER from the European Community from PRC program CNRS/NSFC n°270437 and from CAP 20-25 I-site project.

References

- Ahn, C., Kim, Y., Woo, S., Park, J., 2008. Soil washing using various nonionic surfactants and their recovery by selective adsorption with activated carbon. *Journal of Hazardous Materials* 154, 153-160.
- Alcántara, M., Gómez, J., Pazos, M., Sanromán, M., 2008. Combined treatment of PAHs contaminated soils using the sequence extraction with surfactant–electrochemical degradation. *Chemosphere* 70, 1438-1444.
- Anipsitakis, G.P., Dionysiou, D.D., 2004. Radical generation by the interaction of transition metals with common oxidants. *Environ. Sci. Technol.* 38, 3705-3712.
- Bai, X., Wang, Y., Zheng, X., Zhu, K., Long, A., Wu, X., Zhang, H., 2019. Remediation of phenanthrene contaminated soil by coupling soil washing with Tween 80, oxidation using the UV/S₂O₈²⁻ process and recycling of the surfactant. *Chemical Engineering Journal*.
- Bianco, A., Passananti, M., Deguillaume, L., Mailhot, G., Brigante, M., 2016. Tryptophan and tryptophan-like substances in cloud water: Occurrence and photochemical fate. *Atmos. Environ.* 137, 53-61.
- Bianco, A., Polo-López, M.I., Fernández-Ibáñez, P., Brigante, M., Mailhot, G., 2017. Disinfection of water inoculated with *Enterococcus faecalis* using solar/Fe(III)EDDS-H₂O₂ or S₂O₈²⁻ process. *Water Res.* 118, 249-260.
- Brand, N., Mailhot, G., Bolte, M., 1998. Degradation photoinduced by Fe(III): Method of alkylphenol ethoxylates removal in water. *Environ. Sci. Technol.* 32, 2715-2720.
- Buxton, G.V., Greenstock, C.L., Helman, W.P., Ross, A.B., 1988. Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals ([•]OH/[•]O⁻) in aqueous solution. *J. Phys. Chem. Ref. Data* 17, 513-886.
- Catastini, C., Rafqah, S., Mailhot, G., Sarakha, M., 2004. Degradation of amitrole by excitation of iron(III) aquacomplexes in aqueous solutions. *J. Photochem. Photobiol., A* 162, 97-103.
- Chen, Y., Wu, F., Lin, Y., Deng, N., Bazhin, N., Glebov, E., 2007. Photodegradation of glyphosate in the ferrioxalate system. *J. Hazard. Mater.* 148, 360-365.
- Cheng, K., Lai, K., Wong, J., 2008. Effects of pig manure compost and nonionic-surfactant Tween 80 on phenanthrene and pyrene removal from soil vegetated with *Agropyron elongatum*. *Chemosphere* 73, 791-797.
- Chu, W., Kwan, C.Y., 2003. Remediation of contaminated soil by a solvent/surfactant system. *Chemosphere* 53, 9-15.
- E., H., A., H., G., B., 1966. Pulsradiolytische untersuchung des radikalanions SO₄⁻. *Ber. Bunsenges. Phys. Chem.* 70, 149-154.
- Fang, G.-D., Dionysiou, D.D., Wang, Y., Al-Abed, S.R., Zhou, D.-M., 2012. Sulfate radical-based degradation of polychlorinated biphenyls: Effects of chloride ion and reaction kinetics. *J. Hazard. Mater.* 227-228, 394-401.
- Faust, B.C., Zepp, R.G., 1993. Photochemistry of aqueous iron(III)-polycarboxylate complexes: roles in the chemistry of atmospheric and surface waters. *Environ. Sci. Technol.* 27, 2517-2522.
- Gan, S., Lau, E., Ng, H., 2009a. Remediation of soils contaminated with polycyclic aromatic hydrocarbons (PAHs). *Journal of hazardous materials* 172, 532-549.
- Gan, S., Lau, E.V., Ng, H.K., 2009b. Remediation of soils contaminated with polycyclic aromatic hydrocarbons (PAHs). *J. Hazard. Mater.* 172, 532-549.
- Gharibzadeh, F., Kalantary, R.R., Nasser, S., Esrafil, A., Azari, A., 2016. Reuse of polycyclic aromatic hydrocarbons (PAHs) contaminated soil washing effluent by bioaugmentation/biostimulation process. *Separation & Purification Technology* 168, 248-256.
- Gómez, J., Alcántara, M., Pazos, M., Sanromán, M., 2010. Remediation of polluted soil by a two-stage treatment system: desorption of phenanthrene in soil and electrochemical treatment to recover the extraction agent. *Journal of hazardous materials* 173, 794-798.

450 Guo, H., Liu, Z., Yang, S., Sun, C., 2009. The feasibility of enhanced soil washing of p-nitrochlorobenzene (pNCB)
 451 with SDBS/Tween80 mixed surfactants. *Journal of Hazardous Materials* 170, 1236-1241.

452 Hernández-Soriano, M.C., Mingorance, M.D., Peña, A., 2012. Desorption of two organophosphorous pesticides
 453 from soil with wastewater and surfactant solutions. *Journal of Environmental Management* 95, S223-S227.

454 Huang, W., 2012. Homogeneous and heterogeneous Fenton and photo-Fenton processes : impact of iron
 455 complexing agent ethylenediamine-N,N'-disuccinic acid (EDDS). *Université Blaise Pascal - Clermont-Ferrand II*.

456 Huang, W., Bianco, A., Brigante, M., Mailhot, G., 2018. UVA-UVB activation of hydrogen peroxide and persulfate
 457 for Advanced Oxidation Processes: Efficiency, mechanism and effect of various water constituents. *J. Hazard.*
 458 *Mater.* 347, 279-287.

459 Huang, X.-D., El-Alawi, Y., Penrose, D.M., Glick, B.R., Greenberg, B.M., 2004. A multi-process phytoremediation
 460 system for removal of polycyclic aromatic hydrocarbons from contaminated soils. *Environmental pollution* 130,
 461 465-476.

462 Huguenot, D., Mousset, E., Van Hullebusch, E.D., Oturan, M.A., 2015. Combination of surfactant enhanced soil
 463 washing and electro-Fenton process for the treatment of soils contaminated by petroleum hydrocarbons.
 464 *Journal of environmental management* 153, 40-47.

465 Kiwi, J., Lopez, A., Nadtochenko, V., 2000. Mechanism and kinetics of the OH-radical intervention during Fenton
 466 oxidation in the presence of a significant amount of radical scavenger (Cl⁻). *Environ. Sci. Technol.* 34, 2162-2168.

467 Krýsová, H., Jirkovský, J.r., Krýsa, J., Mailhot, G., Bolte, M., 2003. Comparative kinetic study of atrazine
 468 photodegradation in aqueous Fe(ClO₄)₃ solutions and TiO₂ suspensions. *Applied Catalysis B: Environmental* 40,
 469 1-12.

470 Lee, S.-H., Lee, W.-S., Lee, C.-H., Kim, J.-G., 2008. Degradation of phenanthrene and pyrene in rhizosphere of
 471 grasses and legumes. *Journal of Hazardous Materials* 153, 892-898.

472 Li, F., Guo, S., Hartog, N., 2012. Electrokinetics-enhanced biodegradation of heavy polycyclic aromatic
 473 hydrocarbons in soil around iron and steel industries. *Electrochim. Acta* 85, 228-234.

474 Li, H., Qu, R., Chao, L., Guo, W., Han, X., Fang, H., Ma, Y., Xing, B., 2014. Selective removal of polycyclic aromatic
 475 hydrocarbons (PAHs) from soil washing effluents using biochars produced at different pyrolytic temperatures.
 476 *Bioresour Technol* 163, 193-198.

477 Li, J., Mailhot, G., Wu, F., Deng, N., 2010. Photochemical efficiency of Fe(III)-EDDS complex: OH radical
 478 production and 17β-estradiol degradation. *J. Photochem. Photobiol., A* 212, 1-7.

479 Liang, C., Bruell, C.J., Marley, M.C., Sperry, K.L., 2004. Persulfate oxidation for in situ remediation of TCE. II.
 480 Activated by chelated ferrous ion. *Chemosphere* 55, 1225-1233.

481 Liu, J., 2018. Soil remediation using soil washing followed by ozone oxidation. *Journal of industrial and*
 482 *engineering chemistry* 65, 31-34.

483 López-Vizcaíno, R., Sáez, C., Cañizares, P., Rodrigo, M.A., 2012. The use of a combined process of
 484 surfactant-aided soil washing and coagulation for PAH-contaminated soils treatment. *Sep. Purif. Technol.* 88,
 485 46-51.

486 Mailhot, G., Sarakha, M., Lavedrine, B., Cáceres, J., Malato, S., 2002. Fe(III)-solar light induced degradation of
 487 diethyl phthalate (DEP) in aqueous solutions. *Chemosphere* 49, 525-532.

488 Manenti, D.R., Soares, P.A., Módenes, A.N., Espinoza-Quñones, F.R., Boaventura, R.A.R., Bergamasco, R., Vilar,
 489 V.J.P., 2015. Insights into solar photo-Fenton process using iron(III)-organic ligand complexes applied to real
 490 textile wastewater treatment. *Chem. Eng. J.* 266, 203-212.

491 Mao, X., Jiang, R., Xiao, W., Yu, J., 2015. Use of surfactants for the remediation of contaminated soils: A review.
 492 *Journal of Hazardous Materials* 285, 419-435.

493 Maturi, K., Reddy, K.R., 2006. Simultaneous removal of organic compounds and heavy metals from soils by

electrokinetic remediation with a modified cyclodextrin. *Chemosphere* 63, 1022-1031.

Miralles-Cuevas, S., Oller, I., Ruíz-Delgado, A., Cabrera-Reina, A., Cornejo-Ponce, L., Malato, S., 2018. EDDS as complexing agent for enhancing solar advanced oxidation processes in natural water: Effect of iron species and different oxidants. *Journal of Hazardous Materials*.

Mousset, E., Oturan, M.A., Van Hullebusch, E.D., Guibaud, G., Esposito, G., 2014a. Soil washing/flushing treatments of organic pollutants enhanced by cyclodextrins and integrated treatments: State of the art. *Crit. Rev. Env. Sci. Tec.* 44, 705-795.

Mousset, E., Oturan, N., Hullebusch, E.D.V., Guibaud, G., Esposito, G., Oturan, M.A., 2014b. Influence of solubilizing agents (cyclodextrin or surfactant) on phenanthrene degradation by electro-Fenton process – Study of soil washing recycling possibilities and environmental impact. *Water Research* 48, 306-316.

Mousset, E., Oturan, N., Hullebusch, E.D.V., Guibaud, G., Esposito, G., Oturan, M.A., 2014c. Treatment of synthetic soil washing solutions containing phenanthrene and cyclodextrin by electro-oxidation. Influence of anode materials on toxicity removal and biodegradability enhancement. *Applied Catalysis B Environmental* 160-161, 666-675.

Mulligan, C.N., Yong, R.N., Gibbs, B.F., 2001. Surfactant-enhanced remediation of contaminated soil: a review. *Eng. Geol.* 60, 371-380.

Nagaraju, V., Goje, T., Crouch, A.M., 2007. Determination of copper and iron using [S-S']-ethylenediaminedisuccinic acid as a chelating agent in wood pulp by capillary electrophoresis. *Anal. Sci.* 23, 493-496.

Neta, P., Huie, R.E., Ross, A.B., 1988. Rate constants for reactions of inorganic radicals in aqueous solution. *J. Phys. Chem. Ref. Data* 17, 1027-1284.

Peng, H., Zhang, W., Liu, L., Lin, K., 2017. Degradation performance and mechanism of decabromodiphenyl ether (BDE209) by ferrous-activated persulfate in spiked soil. *Chem. Eng. J.* 307, 750-755.

Peng, S., Wu, W., Chen, J., 2011. Removal of PAHs with surfactant-enhanced soil washing: influencing factors and removal effectiveness. *Chemosphere* 82, 1173-1177.

Reddy, K.R., Ala, P.R., Sharma, S., Kumar, S.N., 2006. Enhanced electrokinetic remediation of contaminated manufactured gas plant soil. *Engineering Geology* 85, 132-146.

Rosas, J., Vicente, F., Santos, A., Romero, A., 2011. Enhancing p-cresol extraction from soil. *Chemosphere* 84, 260-264.

Rosas, J.M., Vicente, F., Santos, A., Romero, A., 2013. Soil remediation using soil washing followed by Fenton oxidation. *Chem. Eng. J.* 220, 125-132.

Soriano-Molina, P., Plaza-Bolaños, P., Lorenzo, A., Agüera, A., García Sánchez, J.L., Malato, S., Sánchez Pérez, J.A., 2019. Assessment of solar raceway pond reactors for removal of contaminants of emerging concern by photo-Fenton at circumneutral pH from very different municipal wastewater effluents. *Chemical Engineering Journal* 366, 141-149.

Tiehm, A., Stieber, M., Werner, P., Frimmel, F.H., 1997. Surfactant-enhanced mobilization and biodegradation of polycyclic aromatic hydrocarbons in manufactured gas plant soil. *Environmental Science & Technology* 31, 2570-2576.

Trellu, C., Mousset, E., Pechaud, Y., Huguenot, D., van Hullebusch, E.D., Esposito, G., Oturan, M.A., 2016. Removal of hydrophobic organic pollutants from soil washing/flushing solutions: A critical review. *Journal of Hazardous Materials* 306, 149-174.

Trellu, C., Oturan, N., Pechaud, Y., van Hullebusch, E.D., Esposito, G., Oturan, M.A., 2017a. Anodic oxidation of surfactants and organic compounds entrapped in micelles - Selective degradation mechanisms and soil washing solution reuse. *Water Research* 118, 1.

538 Trellu, C., Oturan, N., Pechaud, Y., van Hullebusch, E.D., Esposito, G., Oturan, M.A., 2017b. Anodic oxidation of
 539 surfactants and organic compounds entrapped in micelles – Selective degradation mechanisms and soil washing
 540 solution reuse. *Water Res.* 118, 1-11.

541 Tsuneda, S., Ishihara, Y., Hamachi, M., Hirata, A., 2002. Inhibition effect of chlorine ion on hydroxyl radical
 542 generation in UV-H₂O₂ process. *Water Sci. Technol.* 46, 33-38.

543 Usman, M., Faure, P., Hanna, K., Abdelmoula, M., Ruby, C., 2012. Application of magnetite catalyzed chemical
 544 oxidation (Fenton-like and persulfate) for the remediation of oil hydrocarbon contamination. *Fuel* 96, 270-276.

545 Waclawek, S., Lutze, H.V., Grübel, K., Padil, V.V.T., Černík, M., Dionysiou, D.D., 2017. Chemistry of persulfates in
 546 water and wastewater treatment: A review. *Chem. Eng. J.* 330, 44-62.

547 Wu, Y., Bianco, A., Brigante, M., Dong, W., de Sainte-Claire, P., Hanna, K., Mailhot, G., 2015. Sulfate radical
 548 photogeneration using Fe-EDDS: influence of critical parameters and naturally occurring scavengers.
 549 *Environmental science & technology* 49, 14343-14349.

550 Wu, Y., Brigante, M., Dong, W., de Sainte-Claire, P., Mailhot, G., 2014. Toward a better understanding of
 551 Fe(III)–EDDS photochemistry: Theoretical stability calculation and experimental investigation of
 552 4-tert-butylphenol degradation. *J. Phys. Chem. A* 118, 396-403.

553 Yang, Y., Pignatello, J.J., Ma, J., Mitch, W.A., 2014. Comparison of halide impacts on the efficiency of
 554 contaminant degradation by sulfate and hydroxyl radical-based advanced oxidation processes (AOPs). *Environ.*
 555 *Sci. Technol.* 48, 2344-2351.

556 Yang, Y., Pignatello, J.J., Ma, J., Mitch, W.A., 2016. Effect of matrix components on UV/H₂O₂ and UV/S₂O₈²⁻
 557 advanced oxidation processes for trace organic degradation in reverse osmosis brines from municipal
 558 wastewater reuse facilities. *Water Res.* 89, 192-200.

559 Yap, C.L., Gan, S., Ng, H.K., 2011. Fenton based remediation of polycyclic aromatic hydrocarbons-contaminated
 560 soils. *Chemosphere* 83, 1414-1430.

561 Ye, D., Siddiqi, M.A., Maccubbin, A.E., Kumar, S., Sikka, H.C., 1995. Degradation of polynuclear aromatic
 562 hydrocarbons by *Sphingomonas paucimobilis*. *Environmental science & technology* 30, 136-142.

563 Yen, C.-H., Chen, K.-F., Kao, C.-M., Liang, S.-H., Chen, T.-Y., 2011. Application of persulfate to remediate
 564 petroleum hydrocarbon-contaminated soil: Feasibility and comparison with common oxidants. *J. Hazard. Mater.*
 565 186, 2097-2102.

566 Yu, S., Gu, X., Lu, S., Xue, Y., Zhang, X., Xu, M., Qiu, Z., Sui, Q., 2018. Degradation of phenanthrene in aqueous
 567 solution by a persulfate/percarbonate system activated with CA chelated-Fe(II). *Chem. Eng. J.* 333, 122-131.

568 Zhang, C., Wang, L., Pan, G., Wu, F., Deng, N., Mailhot, G., Mestankova, H., Bolte, M., 2009. Degradation of
 569 atrazine photoinduced by Fe(III)–pyruvate complexes in the aqueous solution. *J. Hazard. Mater.* 169, 772-779.

570 Zhang, W., Zhou, S., Sun, J., Meng, X., Luo, J., Zhou, D., Crittenden, J., 2018. Impact of chloride ions on UV/H₂O₂
 571 and UV/persulfate advanced oxidation processes. *Environ. Sci. Technol.* 52, 7380-7389.

572 Zhang, Y., Wong, J., Liu, P., Yuan, M., 2011. Heterogeneous photocatalytic degradation of phenanthrene in
 573 surfactant solution containing TiO₂ particles. *Journal of hazardous materials* 191, 136-143.

574 Zhao, B., Zhu, L., Li, W., Chen, B., 2005. Solubilization and biodegradation of phenanthrene in mixed
 575 anionic–nonionic surfactant solutions. *Chemosphere* 58, 33-40.

576 Zheng, G., Selvam, A., Wong, J.W.C., 2012. Enhanced Solubilization and Desorption of Organochlorine Pesticides
 577 (OCPs) from Soil by Oil-Swollen Micelles Formed with a Nonionic Surfactant. *Environmental Science &*
 578 *Technology* 46, 12062-12068.

579 Zhou, D., Wu, F., Deng, N., 2004. Fe(III)–oxalate complexes induced photooxidation of diethylstilbestrol in water.
 580 *Chemosphere* 57, 283-291.

581 Zhou, J., Jiang, W., Ding, J., Zhang, X., Gao, S., 2007. Effect of Tween 80 and β-cyclodextrin on degradation of

582 decabromodiphenyl ether (BDE-209) by white rot fungi. Chemosphere 70, 172-177.
583 Zhou, W., Wang, X., Chen, C., Zhu, L., 2013. Removal of polycyclic aromatic hydrocarbons from surfactant
584 solutions by selective sorption with organo-bentonite. Chemical Engineering Journal 233, 251-257.
585 Zhu, L., Zhou, W., 2008. Partitioning of polycyclic aromatic hydrocarbons to solid-sorbed nonionic surfactants.
586 Environmental pollution 152, 130-137.

587

588

589 Table 1: Second-order rate constants ($\text{M}^{-1} \text{s}^{-1}$) of TW80 and PHE with different radicals
 590 determined using LFP and competition kinetic method.

Compounds	$k_{HO^\bullet}$	$k_{SO_4^{\bullet-}}$	$k_{Cl_2^{\bullet-}}$
TW80	$9.9 \pm 0.1 \times 10^9$	$4.6 \pm 0.2 \times 10^9$	$7.1 \pm 0.1 \times 10^6$
PHE	$6.1 \pm 0.2 \times 10^9$	$4.3 \pm 0.4 \times 10^9$	$4.6 \pm 0.3 \times 10^8$

591

592

Figures Caption

1) Absorption spectra of Fe(III)-EDDS (0.5 mM), $\text{S}_2\text{O}_8^{2-}$ (5 mM) and 90 μM of PHE in 0.5 g L^{-1} of TW80. Emission spectrum of the adopted irradiation lamps with a maximum centred at 365 nm.

2) Degradation of PHE in TW80 0.5 g L^{-1} and degradation of Fe(III)-EDDS in different systems (Blue marked). Initial conditions are: $[\text{PS}] = 5 \text{ mM}$, $[\text{Fe(III)}] = 0.5 \text{ mM}$, $[\text{Fe(III)-EDDS}] = 0.5 \text{ mM}$, $[\text{TW80}] = 0.5 \text{ g L}^{-1}$, pH: 3.5 ± 0.2 .

3) **A)** Influence of Fe(III)-EDDS concentration on the photodegradation of PHE in aqueous solution. **B)** Degradation percentage of PHE after 30 and 150 minutes of irradiation as a function of different Fe(III)-EDDS concentrations. Initial conditions are: $[\text{PS}] = 5 \text{ mM}$, $[\text{TW80}] = 0.5 \text{ g L}^{-1}$ and pH: 3.5 ± 0.2 .

4) Influence of PS concentration on the photodegradation of PHE in aqueous solution. Initial conditions are: $[\text{Fe(III)-EDDS}] = 5 \text{ mM}$, $[\text{TW80}] = 0.5 \text{ g L}^{-1}$ and pH: 3.5 ± 0.2 .

5) Influence of concentration of Cl^- for photodegradation of PHE in aqueous solution. $[\text{PS}] = 5 \text{ mM}$, $[\text{Fe(III)-EDDS}] = 5 \text{ mM}$, $[\text{TW80}] = 0.5 \text{ g L}^{-1}$ and pH: 3.5 ± 0.2 .

6) Free radical scavenging tests photodegradation of PHE in aqueous solution. $[\text{PS}] = 5 \text{ mM}$, $[\text{Fe(III)-EDDS}] = 0.5 \text{ mM}$. $[\text{tert-Butanol}] = 50 \text{ mM}$, $[\text{Methanol}] = 50 \text{ mM}$, pH: 3.5 ± 0.2 .

Figure 1

Figure 2

641 **Figure 3**

Figure 4

Figure 5

668 **Figure 6**

669

670

671