The Southern Atlas Front in Tunisia and its foreland basin: structural style and regional-scale deformation

Amara Masrouhi ^{a,b,*}, Mohamed Gharbi ^b, Olivier Bellier ^c, Mohamed Ben Youssef ^b

^a King Abdulaziz University, Faculty of Earth Sciences, Geo-exploration Techniques Department, Jeddah, Saudi Arabia

^b Water Research and Technologies Center, Geo-Resources Laboratory, Borj-Cedria, Soliman, Tunisia

^c Aix Marseille Univ, CNRS, IRD, Coll France, CEREGE, Aix-en-Provence, France

* Corresponding author: amara.masrouhi@fsg.rnu.tn

Abstract:

Surface and subsurface data together with balanced cross-sections are used to emphasize the structural style and regional-scale deformation of the Southern Atlas Front in Tunisia and its foreland basin. This E–trending shaped belt reflects successive tectonic events. Mixed thick- and thin–skinned structural style characterizing this belt is defined by shallow thrusts rooted down into deep E–trending basement thrusts with major decollement level located within Triassic salt. A NW–trending system is also recognized in the Front and particularly responsible for the belt’s curved shape, which probably acts as transfer areas between the wide-ranging E–trending major structures. The intermediate areas are characterized by salt tectonics probably active during Jurassic and Lower Cretaceous, in response to active synsedimentary normal faults and differential sedimentation loading. The shortening amount varies and decreases from north to south and from west to east: ~7.45% in the West, ~3.3% in west-central part, ~2% in central area to insignificant (~1.4%) in the east. From north to south, the Northern Chotts Range narrow zone accommodates overall the Cenozoic shortening with ~6.95% and less than 1% in the Chotts region. This variation coincides with the pre-Cenozoic basin’s architecture acquired during Mesozoic Tethyan evolution, which generates the main Jurassic E–trending Chotts basin. Inside this basin, distributary sub-basins formed in relation with NW–trending faults active during Cretaceous period. The Chotts basin appears, therefore, as Mesozoic rift rollover structure well-preserved in the front of the belt. The compilation of data suggests that the Front of the southeasternmost end of the Atlassic domain is distinguished by the reactivation of inherited fault systems acting during alpine orogeny as major southward thrusts. The foreland shows a structural singularity to be formed by a preserved almost unmodified and well exposed large Mesozoic rollover structure.

Keywords: Southern Atlas Front; Tunisia; balanced cross-sections; tectonic inversion; structural inheritance

1. Introduction

The evolution of sedimentary basins typically involves extensional deformation of both basement and sedimentary cover. The initial rifting stage usually progress to continental passive margin, which is frequently characterized by an irregular sea floor dominated by repetitive depocenters associated to intra-basin active listric normal growth faulting. This is well observed in the present-day Atlantic-type passive margin and is well preserved in several inverted margins such as the Tethyan margins around the Mediterranean (present coordinates). These margins are characterized by associated sub-basins produced by major fault systems themselves made-up by regional tectonic extension (Allen and Allen, 1990). This system is in control of basin geometry (graben and half graben) strongly attested by important thickness variations of sedimentary sequences. The tectonic inversion is defined once the aforementioned basins are deformed by subsequent compressional tectonic (Letouzey, 1990). During the inversion of the margin, normal faults are partly or entirely inverted (Bally, 1983). The reactivation of normal faults is commonly well observed for those with an opposite strike to the subsequent compression direction. At the scale of structures, the reverse movement of the pre-existent extensional systems, is typically derived by preexistent normal faults that bound graben or half graben (Bally, 1983). In addition, the existence of salt layers in the sedimentary sequences characteristically made additional complication to the inversion process. The presence of salt is also involved in the mode of inversion, with classically occurrence of the preferential major decoupling along the salt layers (Roure et al., 1992). The Atlas Mountains of North Africa are recently documented by many authors as one of worldwide belt resulting from the basin inversion processes (e.g. Vially et al., 1994; Ziegler et al., 1995; Roure et al., 2012; Frizon de Lamotte et al., 2013; Lepretre et al., 2018a, b).

The Southern Atlas of Tunisia forms the front of the North African Alpine orogenic belt, between the deformed Atlassic domain to the North and the relatively undeformed Saharan platform area to the South (Fig. 1). This region also bounds the easternmost segment of the Cenozoic deformation front in the Maghreb region, which extends over 2000 km from High Atlas in Morocco to the West, the Algerian Saharan Atlas in the center and the Tunisian Atlas to the East. From West to East, the southern deformational front of the Maghrebides belt drastically changes with typical intra-continental chain in the High Atlas of Morocco and a fold-and-thrust belt in the Southern Atlas of Tunisia (Teixell et al., 2003; Benaouali-Mebarek et al., 2006; Missenard et al., 2007; Gharbi et al., 2015). The physiographic characteristic of the Southern Atlassic domain in Tunisia is a series of narrow elongated massifs separated by large basins (Figs. 2, 3). It shows a distinguished distribution of fold's geometry which led to

a varied structures and basin's interpretations by previous studies during the past 65 years. 1) 1950's to early 1990's: strike-slip faulting was proposed, which produced the fold distribution and the style of the Tunisian Southern Atlas (e.g. [Castany, 1954](#); [Zargouni and Ruhland, 1981](#); [Abdeljaouad and Zargouni 1981](#); [Abbes and Zargouni 1986](#); [Fakraoui 1990](#); [Ben Ayed, 1993](#); [Bedir et al., 2001](#)). Two major NW-trending right lateral strike-slip fault systems (Gafsa and Negrine-Tozeur, [Fig. 2](#)) were documented as the major ones, which are associated to left lateral NE-trending one recognized as less active faults systems. The structures are identified, in this model, as curved, right-stepping *en echelon* shaped folds related to the strike-slip fault systems. 2) 1990's late: few works acquaint with salt tectonics model ([Hlaiem et al., 1997](#); [Hlaiem, 1999](#)). These works introduce an evolution in which a significant diapirs took place during Early Jurassic period. This structural model locates the subsequent Tertiary folding as preferentially nucleated on the preexistent diapir apex. 3) Fault-related folding 2000's model ([Outtani et al., 1995](#); [Ahmadi et al., 2006](#)) was dominated by south verging thrust system on which folds are located. This model interprets the anticline structures in this zone as developed by the result of ramp-related folding, disturbing only the sedimentary cover (thin-skinned model) without basement involving. 4) 2010's: structural reconsideration of this belt was based on the role of the pre-existent faults systems in the subsequent tertiary deformation. This model highlights a structural evolution controlled by deep-seated basement faults ([Riley et al., 2011](#); [Said et al., 2011b](#); [Gharbi et al., 2015](#); [El Amari et al., 2016](#)).

This paper focuses on the structural style and regional-scale deformation of the Southern Atlas Front in Tunisia and its foreland basin as an inverted basin. To document these goals, surface data (geologic mapping, cross-sections, markers of tectonics deformation) are analyzed. New seismic reflection profile's interpretation, well data and long-transect sequences correlations are as well used to highlight the main structure and/or borders of different domains. Three long balanced cross-sections across the southern Atlassic belt are constructed to estimate the amount and the rate of shortening, as well as the variations in term of structural architecture and tectonic style using methods described in [Dalstrom, 1969](#); [Coward, 1996](#); [Decelles and Giles, 1996](#).

2. Geological setting and lithostratigraphic overview

The Southern part of the Atlas Mountains of Tunisia is located in the southeastern part of the Maghreb orogenic belt called the Maghrebides. The Maghrebides orogen is qualified as the North African Alpine chain, which is shaped during Tertiary–present-day compressional events related to Africa–Eurasia convergence setting. The timing of the deformation, as well as the main compressive events are now well established after sound arguments advanced during the 1990^s–2000^s (e.g. [Beauchamp et al., 1999](#);

Bosworth et al., 1999; Frizon de Lamotte et al., 2000; Guiraud et al., 2005; Masrouhi et al., 2007; Khomsi et al., 2009; Frizon de Lamotte et al., 2011; Gharbi et al., 2014). The convergence between African and Eurasian plates is expressed firstly in Tunisia by the positive inversion of major Tethyan basin and now it is mostly accepted as operated during Late Cretaceous (Guiraud and Bosworth, 1997; Benaouali-Mebarek et al., 2006). Even if in the southern Atlasic domain of Tunisia, the Late Cretaceous inversion markers are not yet well-identified; this issue is well highlighted in central and northern Tunisia (Masrouhi et al., 2008; Dhahri and Boukadi, 2010). The first significant compressional event causative of the Atlas building is the middle-late Eocene Atlasic compression (Vila et al., 1993; Frizon de Lamotte et al., 2000). This later is well underlined by local as well as regional unconformities on the overall Tunisian Atlas domain (Vila et al., 1975; Khomsi et al., 2006; Gharbi et al., 2015). In southern Tunisian domain, previous field data and regional geological mapping are consistent with this unconformity dating an Eocene compressional event (e.g. Slimane et al., 1991; Regaya et al., 1991; Riley et al., 2011). This compressional stage is followed by a period of tectonic quiescence during Oligocene and Lower Miocene times. The thick Numidian siliciclastic series deposited in the Tell-Rif sub-basins in addition to the general geodynamic setting (Miocene back-arc opening of the Algerian-Provencal basins and the rotation of Corsica and Sardinia) led to a model in which the northern edge of this chain is interpreted as an accretionary prism developed in the western Mediterranean as back-arc oceanic basin (Vergés and Sàbat, 1999; Frizon de Lamotte et al., 2009; Roure et al., 2012; Van Hinsbergen et al., 2014; Hamai et al., 2015). The Oligo-Miocene extensional setting is interpreted by several authors to be caused by the slab retreat (Jolivet and Faccenna, 2000; Deverchere et al., 2005). As summarized by several authors, events of widespread deformation in the Southern Atlasic Front of Tunisia are thought to have occurred during middle-Miocene to present-day times (Zargouni, 1985; Chihi, 1992; Swezey, 1996; Bouaziz et al., 2002; Gharbi et al., 2014). All authors, having worked in this region, agree that this event is the major compressional tectonic event building the southern Atlasic domain in Tunisia (e.g. Ben Ayed, 1980; Zargouni, 1985; Boukadi, 1989; Dlala, 1992; Chihi, 1992; Zouari et al., 1990; Gharbi et al., 2014; Bahrouni et al., 2014; Soumaya et al., 2015). This period is characterized in southern Atlasic domain by NW-trending main regional compression direction during Miocene-Pliocene and N-Trending tectonic during the late Pliocene-Quaternary (Dlala and Hfaiedh, 1993; Billi et al., 2011; Gharbi et al., 2014; Soumaya et al., 2018). The present structure of the Southern Atlasic Front in Tunisia comprises a series of elongated structural edifices widely accepted today as deriving from the inversion of Mesozoic depocenters recording the former passive margin basin and/or mini-basins along the south Tethyan margin (e.g. Chihi et al., 1992; Dhahri and Boukadi, 2010; Briki et al., 2018; Frifita et al., 2019).

The structural style of the southern Atlassic domain of Tunisia is dominated by E-W–trending structures. In major cases, these structures are shaped by deep E-W–trending fault systems. These systems are widely documented as inherited systems from the former south Tethyan margin (Gharbi et al., 2013; Gabtni et al., 2013; Tanfous-Amri et al., 2017). From a geodynamic point of view, this system is well correlated during Jurassic and Cretaceous periods with the Alpine Tethys evolution within the overall context of the relative left movement of Africa with respect to Eurasia (Boughdiri et al., 2007; Frizon de Lamotte et al., 2013). The breakup of Pangea has been expressed by successive extensional episodes that generate some major basins from Morocco to Tunisia (Moragas et al., 2016, 2017; Naji et al., 2018a; Jaillard et al., 2019). This period is started during the Late Permian and continued until Late Cretaceous. The Initial Mesozoic basin geometry was characterized by Triassic basin subject to prevalent evaporitic facies across overall Tunisia (Burolet, 1956; Kamoun et al., 2001). Triassic evaporitic sequences are indicative of an epicontinental environment which was recognized in entirely Maghreb sub-basins (e.g. Masrouhi and Koyi, 2012; Masrouhi et al., 2014a; Moragas et al., 2017). Triassic sequences are usually associated to subordinate basalts, locally preserved in Tunisia (Burolet, 1973; Laaridhi-Ouazâa, 1994) and commonly viewed as products of rift-related magmatism (Masrouhi et al., 2013; Masrouhi et al., 2014b). The Jurassic period is well documented as active rifting period overall the Maghreb area (Bracene et al., 2002; Boughdiri et al., 2007; Saura et al., 2014; Martin-Martin, 2017; Moragas et al., 2018). This stage is well preserved in south Tunisia by E-W–striking basins (such as the Chotts basin; Patriat et al., 2003) produced by major E-trending fault systems themselves made-up by regional N-trending tectonic extension. This system is locally in control of basin geometry (graben and half graben) strongly attested by important thickness variations of sedimentary sequences (Ben Ferjani et al., 1990; Bahrouni et al., 2016; Tanfous-Amri et al., 2017; this work). The N-S–extensional context prevailed during Jurassic become a NE-SW regional tectonic extensional setting during Early Cretaceous (Louhaichi and Tlig, 1993; Jaillard et al., 2017; Naji et al., 2018b). This second phase is mainly characterized by NW-trending faults systems and associated sub-basins. This translation from Jurassic E-W (present-coordinates) prevailing configuration to Early Cretaceous NW-SE shape is compatible with general E-W transform faults of Atlantic which change to NW-SE ones in the Alpine Tethys (Rosenbum et al., 2002; Frizon de Lamotte et al., 2011). In Southern Tunisia, the recent works establish a paleogeographic partition of the continental platform, which is scattered in some localities by related volcanic and halokinetic movement that are recognized overall Tunisian domain (Burolet, 1991; Mattoussi Kort et al., 2009; Jaillard et al., 2017; Dhahri and Boukadi, 2017). The main synsedimentary normal faults form a general tilted blocks basin’s geometry, for which the Aptian–Albian ages are documented to have the

maximum extensional related-structures period in Tunisia (Gharbi et al., 2013; Masrouhi et al., 2014a; Naji et al., 2018a). The rapid active extension characterizing Early Cretaceous period seems to decrease during late Cretaceous times. Basin shows a thick deposition of open marine shale and shallow carbonates sealing the above-mentioned tilted basin's geometry.

The compilation of lithostratigraphic sequences data in the Southern Atlasic mountains of Tunisia, presented in this work, is derived from petroleum wells and outcrops around the studied structures (Ben Ismail, 1982; Ben Youssef and Peybernes, 1986). The Triassic series are thick of about 400-800 m showing important thickness variations. In the Northern Chotts range, the Triassic series are mainly evaporitic showing carbonate and siliciclastic beds. The Jurassic strata consist of limestones of the Nara Formation (Soussi and Ben Ismail, 2000). Three members were recognized: lower, Middle and Upper Nara showing similarly important thickness variations (250 m to 2500 m). On the Jurassic limestones, Lower Cretaceous series show a siliciclastic, clay, dolomites and sandstones layers. The Lower Cretaceous shows in its bottom 300 m thick clay and dolomites of the Meloussi Formation. This later is overlain by the Boudinar Formation which is thick of about 200 m (Fig. 2).

The Bouhedma Formation consists of a repetitive mixed carbonate-clay-evaporite sequences and considered to be late Hauterivian age. This Formation shows also an important thickness variation (from 400 m to 2000 m). The Bouhedma Formation is overlain by Sidi Aïch Formation composed mainly of shaly sands and ferruginous sandstone. This Formation, thick of about 200 m, corresponds to fluvial environment and dated as Barremian age deduced from its stratigraphic position. In the northern Chotts range, the Sidi Aïch Formation is capped by the Orbata Formation dated as Aptian age (Ben Youssef et al., 1985). This later is composed of three members i.e. the lower Orbata corresponds to a dolomitic bar named in this range Barrani bar dated as Bedoulian age, the middle Orbata is a marly and sandy carbonates and the upper member consists of a sandy unit. In the Chotts basin, the upper Aptian to the lower and Middle Albian is characterized by hiatus. This discontinuity is due to non-deposition or to subsequent erosion. The Zebbag Formation was subdivided into four members, the upper Albian is formed by an alternation of marls and carbonates with *Knemiceras* named M'Razig member that is overlain by the massive dolomitic bar of the Rhadouan member. On this member rest the marls, carbonates and evaporites of Kerker member which is dated as Cenomanian age. The top of the Zebbag Formation is composed of massive bar of dolomite and dolomitic limestones named Guettar. This bar is dated as Turonian. The Aleg Formation, which is dated Turonian–Early Campanian, is characterized by marls and limestones layers. The Berda Formation, dated as Late Campanian–Early Maastrichtian, is

essentially formed by two carbonate members, which are separated by a middle member consisting of an alternation of marl and limestones (Fig. 2). Upper Cretaceous limestone sequences in the Chotts Basin are overlain by Lower Paleocene, lay the marine green shale of El Haria Formation that is overlain by the Eocene deposits which consist of the Metlaoui Formation limestones and the continental unit of Bouloufa Formation. This later is formed by conglomerates, red clay, and lacustrine limestones. In the southern Atlassic domain structures, the sequences are generally overlain unconformably by the Segui Formation, which is interpreted as syntectonic growth strata acknowledged to be the product of the erosion of the structures uplifted during Cenozoic compressional events.

3. Surface data

Recently the knowledge about the west Mediterranean geodynamic evolution is upgraded after several structural and geodynamics wide-ranging studies (e.g. Tricart et al., 1994; Faccenna et al., 2004; Lacombe and Jolivet, 2005; Vergés and Fernández, 2012, among others). The southern Tunisian Domain is rather understudied, compared to Algeria and Morocco, in term of regional-scale deformation. During the last decade, many studies carried out in the southern Atlassic front of Tunisia (e.g. Ahmadi et al., 2006; Riley et al., 2011; Said et al., 2011a; Gharbi et al., 2013, 2015). However, all these mentioned works were focused along the northern part of this belt (Metlaoui, Gafsa, Sehib and Orbata structures, Fig. 2). The southernmost segment (Northern Chotts range) and its relationships with the Saharan platform, together with taking into consideration the preexistent basin structure are yet to be appraised. The Northern Chotts Range (NCR) is a 165 Km-long and narrow 5 km-wide chain (Fig. 3). It constitutes the southernmost domain showing folded structures well highlighted by their physiographic characteristic as well as their remarkable and well-preserved closures. From West to East, this series of anticlines, with distinguished curved axes, are closely associated with E- to NW- to WNW-trending fault systems. The following sections address the structural analysis of the NCR, which is based on field observations across the eastern and the central parts of the Range as well as the northern isolated anticlines (Berda and Sehib), and the southern regions that is the Chotts basin together with the Southern Chotts Range (SCR). Data used are derived from this study, the preexisting 1/100000 geologic map published by the national official of Mines, Gharbi, (2013) and El Amari et al. (2016). This field work is controlled by systematic transects (with dips reading, faulting and folding analysis, entities relationships...), surface geologic cross-sections together with detailed mapping of the key-zones.

The Jebel Sehib structure, which extends in the South of Metlaoui Range, corresponds to an isolated anticline structure in the wide plain named the Segui plain (Figs. 3, 4UU'). The surface analysis shows a

dissymmetric folding (Ahmadi et al., 2006) for which the forelimb shows 70°–dipping strata and the backlimb displays gentle dipping strata of about 20° (Fig. 4TT'). This anticline illustrates a curved axis which changes from N120 in the West (near the NW-trending fault cutting the structure), to N80 in the middle and to N30 in the East (Fig. 3). This structure was for long time interpreted as the result of the master strike-slip movement and genetically related to strike-slip tectonics (Zargouni, 1985; Bedir et al., 2001; Zouari et al., 1991; Zouaghi et al., 2005, 2011). The elongated central part shows a constant and long (compared to East and West closure) geometry which mean that the formation of the fold was not complicated by the faults bordering the structure. The geometry of the fold shows a long gentle dipping backlimb and a short high dipping forelimb which seems to be not controlled by the adjacent strike-slip fault. The Sehib anticline structure is particularly well outlined by the Campanian limestones (Ahamdi et al., 2006). At the front, this anticline shows local-scale bend folding, which is probably the surface expression of deep thrust on which the fold is shaped (Fig. 4TT'). This anticline is characterized by south-verging bending style of folding showing a clear kink in the southern limb with Cretaceous and Eocene strata which suggests a north-dipping axial surface. The anticline is limited by two North and South flat domain plains filled by thick Mio-Plio-Quaternary Segui Formation displaying characteristics of syntectonic alluvial deposits exhibiting growth-strata pattern.

The Jebel Berda shows particularly unusual geometry showing an inverted northern back-limb and a long normal southern forelimb (Fig. 4UU'). The second singularity of this fold is the notable entangled form between an eastern part with an inverted northern limb which corresponds to the Ain El Beida, and a classically south verging anticline in the western part which corresponds to the Jebel Atra. This structure shows series extending from Cenomanian–Turonian sequences in the core to Mio-Plio-Quaternary deposits in the limbs. The interlacing between the two parts of the structure (Ain El Beida and Atra) is located within the Coniacian–Santonian marls sequences. The distinctive entangled form within Coniacian–Santonian marls is associated to thickness change, which vary from 140 m in the West to 50 m in the East. In the eastern part, the cross-section (Fig. 4UU') cuts this structure and illustrates the local Berda style with north-verging folding. This anticline is associated to E-trending fault, which corresponds to north-verging thrust. The hanging wall of Berda thrust is formed by north-verging anticline structure showing a north limb with 65–70°–north-dipping Cretaceous strata and a southern limb with 45°south-dipping strata. The footwall of the Berda thrust is formed by an overturned 75–80°–south-dipping layers. In the western part, i.e. Jebel Atra, the structure corresponds to N095/10°W anticline corresponding to a typical E–trending south–verging Atlassic structure, regarding the overall belt structure.

The central and the western part of the Northern Chotts Range is an E-trending narrow belt (Mahjoub and Fakraoui, 1990). This belt is generally characterized in the Jebel Askar, Sif El Laham and Teferna-Zitouna structures by two differentiated types of folding and/or style of structures between the northern and the southern limbs (Figs. 3, 4VV'). All these above-mentioned structures show high-dipping northern limbs with upper Cretaceous strata unconformably covered by Eocene and Miocene sequences. The southern limbs are formed by gentle-dipping to horizontal Lower Cretaceous sequences (Fig. 5). The southern side of all these structures are truncated by E-trending kilometric faults systems parallels to the axis of major anticlines (Fig. 3). These E-trending faults show usually inverse movement within Aptian–Albian sequences, in which the inverse throw rarely exceeds some tens of meters. This structural singularity of long faults with restricted throw is usually associated to hectometric folding well-arranged along faults, and which is observable over several kilometers (Fig. 4). In addition, along this E-trending fault systems a disharmonic folding is frequently observed particularly in Aptian marls and carbonates alternations. This fault systems (with restricted throw) vs folding is likely incompatible with the Tertiary compressions, which is more developed in the nearby structures. Regarding the folding style, with south gentle-dipping Early Cretaceous and North high-dipping Late Cretaceous, this south dipping fault system (with unimportant displacement, Figs. 4VV', 5) seems to be an inherited system deeply rooted and passively transported on north-dipping major thrust, which accommodate the Tertiary compressions.

Eastward, the Bir Oum Ali–Hachichina structure (BOAHS) comprises two major anticlines separated by the Bir Oum Ali–Hachichina fault systems (BOAHFS). This area exposes, in term of deformation, two different styles with a northern highly deformed anticline and a southern gentle deformed one (Fig. 4WW'). The BOAHS shows a curved-axis which changes from West to East (El Amari et al., 2016). As shown on the geologic map (Fig. 3), the BOAHS have a NE-trending axis in the Jebel Bir Oum Ali, a NW-trending one in the Jebel Hachichina and an ESE-trending axis in the Jebel Naimia. The field investigation (Figs. 5), along the BOAS anticline, highlight that the southern part is formed by the Early Cretaceous series showing a long southern limb and a short northern one characterized by ~10–15°S-dipping southern forelimb and ~45–65° N-dipping northern backlimb (El Amari et al., 2016). While the southern side is simple and shows a little deformed side of the BOAHS (Fig. 5), the northern part shows a complex pattern. This part is characterized by the anticline's succession with curved hinge line and showing a right stepping "*en echelon*" arrangement. These two different sides are separated by the BOAHFS and does not show a syncline in-between (Figs. 4UU'). The curved secondary (in respect to general structure) anticline succession is well observed along the central part of the NCR. Furthermore, from West to East geometric complexity increase with usually south overturned limbs (Figs. 4, 5). This area offers a well

traced ~15-20°-angular unconformity between the aforementioned Cretaceous series and Miocene–Pliocene sequences (Fig. 5). Eastward, this angular unconformity is localized between Paleocene–Eocene and Miocene–Pliocene series along 20 km northern limbs of this area (Figs. 3).

Eastward of the BOAHS, the Jebel Beidha structure, is the second remarkable fold in South Tunisia showing a northern inverted limb. The Jebel Beidha structure is an uncharacteristic sub-circular anticline cored by Lower Cretaceous sequences (Fig. 3). This anticline shows a 25°-south-dipping overturned northern backlimb and a 31°-south-dipping southern forelimb (Fig. 4XX'). The northern forelimb and Beidha anticline's core are abruptly affected by the 70°-south-dipping thrust, well observed in the Oum Aguel region. The Beidha structure exhibits along the core numerous ENE-trending normal faults, probably related to the Tethyan period regarding their tendency versus the fold deformation (Figs. 6, 7).

Eastward, the Beidha structure is bordered by rhomboidal area named the Bou Loufa basin typical of pull-apart basin shape (Zargouni, 1985; Abbes and Zargouni, 1986; Abbes et al., 1986; Fakraoui et al., 1991). This structure is shaped by northwest-trending dextral faults comprising two en echelon segments, which are the Jebel Stah – Chebket Bou Loufa and the Jebel El Haira segments (Fig. 3). These segments delimit right-stepover fault zone defining releasing stepover. The northwestern corner of this pull-apart basin developed at the end stage the formation of circular to elliptic salt intrusion; documented in previous literature as the Hadifa diapir (Abbes and Zargouni, 1986).

The eastern part of the NCR is an E-trending structure (Haidoudi structure, Fig. 3), which change at the easternmost segment (Zemlet el Beidha, Fig. 3) to NE-trending anticline. All this part is called the Zemlet el Beidha-Jebel Haidoudi structure (ZBHS). This structure, of about 24 km long, have a curve-shaped geometry with two main anticlines separated by the NW-trending Fejej fault (Fig. 3). The Zemlet el Beidha is a south-to-south-east verging asymmetric anticline with only an east closure and shows, from West to East, a curved axis that changes from E-strike to NE-strike (Fig. 3). This 16 Km long and 3-to -5 km wide structure, is located between the Menzel Habib plain to the North and the Chotts Fejej and Draa Oudhreh respectively Southwest and Southeast. The Zemlet el Beidha anticline is respectively cored by the Hauterivian–Barremian in the central part, the Albian in the Jebel Meida and the Coniacian–Santonian in the Jebel Romana and Gouada plain (Fig. 3). The north-eastern segment of the ZBHS is formed by gentle deformed Cretaceous sequences covered unconformably by Late Miocene–Pliocene to Quaternary series. The surface geometry of the ZBHS shows a ~20° N-dipping backlimb made of Coniacian–Santonian green marls alternated with bioclastic limestones unconformably overlain by Miocene continental sandstones (Fig. 4ZZ'). The forelimb is formed by the Aleg 45° S-dipping Formation,

itself unconformably overlain by the Segui late Plio-Quaternary Formation. Eastward, of the Jebel Tebaga Fatnassa a disturbance of dips is recorded in the limestones of the Aleg Formation possibly associated with N-dipping reverse faulting (Fig. 4XX'). Westward at the central part of the ZBHS, numerous N100–110°E apparent trending strike-slip faults, along Khanguet Aïcha and Khanguet Amor, regions affect the 45° S-dipping limb. Along the 9 km-long NNW–SSE surface cross section, across Jebel Es Smaïa at the North and the Jebel Haidoudi anticline at the South (Figs. 4YY'), the structure shows an asymmetric fold with a core occupied by Coniacian dolomites showing a ~15 to ~20° N-dipping backlimb and ~40° S-dipping southern forelimb. In this southern limb, the Late Cretaceous strata are unconformably overlain by Plio-Quaternary continental deposits. The Northern limb is affected by ENE-trending reverse fault that separates the Jebel Haidoudi anticline from the Oued Krenafess syncline in the south. This later syncline forms the hanging-wall of E-trending fault systems which likely correspond to the inversion of E-striking inherited system that delimits the Jebel Es Smaïa monocline at the north. In the southern limb of Jebel Haidoudi anticline, a main ~45 to 70°–NE-dipping normal faults are observed, most of them are a preserved syndimentary sealed normal faults. This normal faulting is usually sealed by the Santonian marls (Figs. 6, 7) attesting an ante-Santonian syndimentary activity. The Coniacian dolomitic sequence thickness increases toward the NE hanging-wall and are characteristic of syndimentary faults activity. In previous works, the numerical analysis of data (Gharbi et al., 2013), consisting of bedding rotation data to restore the horizontal bedding planes, highlights a ~N30 to 65°–E-trending tectonic extension that prevailed during the Coniacian–Santonian times.

The southern Chotts Range is an E-W striking physiographic chain, which extends over than 100 Km from Kebili in the West into Jebel Aziza in the El Hamma in the East (Ghanmi and Potfaj, 1987; Ghanmi et al., 1988). Two distinguished crests are formed by the lower Albian and Turonian dolomitic bars which are separated by marls intercalation. These two crests are well parted in the western part by the Cenomanian sequences. Both crests get closer in eastern part and becoming one on another corresponding to complete disappearance of the Cenomanian intercalations (Ghanmi and Potfaj, 1991). This chain was interpreted previously as the southern limb of a mega fold structure called the Chotts mega-anticline. Along this chain, the layer's dips collected during field work are usually between 5° and 10°, and never exceeds 15°, which cannot be taken necessarily as related to tectonic compressions.

In the end, the main fault systems controlling and associated to the folding, are of two types: 1) E-trending north-dipping faults act as major southward thrust systems and 2) E-trending south-dipping faults act as back-thrust likely connected down on the major north-dipping ones. These facts are

demonstrated in some structures by their preserved character as former normal faults deeply rooted (will be developed in the next seismic section) and passively transported on the north-dipping major ones.

4. Subsurface data

Eleven oil wells and nine seismic profiles are used in this work to describe and define the deep configurations of key-zones in the objective to connect and extend geological data in depth.

4.1. Correlation

Three correlations (C1, C2 and C3) between the sedimentary sequences of the study area and the nearby structures (Southern Central Tunisia in the North, and the Saharan Platform in the South) is used to shows lateral variations of thickness, facies and gaps (Figs. 8-10). The correlation is given here without major faults control with a primary objective to describe them in absence of structural adjustment, which will be integrated in the next section. For confidentiality reasons, in this work, the wells are numbered independently from their original names.

In the Saharan platform, the Triassic series are of typical Germanic facies composed of the three series (Soussi et al., 2017); (i) the basal shale-sandy unit (TAGI: Trias argilo-gréseux inférieur). The TAGI is interpreted as having been deposited in a fluvial deltaic environment, the TAGI shows habitually lateral injected volcanic sills (Ben Ismail, 1991), (ii) the intermediate marine carbonate unit is characterized by dolomites of the Messaoudi and Rehache Formations and (iii) the evaporitic unit of the Adjadj (Messaoudi and Mhira Evaporites) which show lagoonal deposits (Busson, 1970; Ben Ferjani et al., 1990; Bouaziz et al., 1999; Kammoun et al., 2001). The Triassic series thickness varies from 0 m (absent) in the P8 well to 500 m in the P8 and P9 wells. Towards the North, the subsurface data show a thick Triassic series, exceeding the 1500 m (Tanfous Amri et al., 2005, 2017). However, there are no wells that crossed the total Triassic sequences at the Southern Tunisian Atlas. At the P6 well (Fig. 8), the 1700 m-thick crossed sequences show a chaotic mixture facies of gypsum, clay, silt and anhydrite deposit with a few thin intercalations of dolomite (Buroillet, 1973; Ben Ferjani et al., 1990).

In the study area, many petroleum wells cross the totality of the Jurassic series. Its maximum thickness is found in the Chott Fejej where they reach a total thickness of 2500 m at the P2 well and passes even to 2650 m in nearby wells (Fig. 9). The thickness of Jurassic progressively decreases, and the facies becomes evaporitic and detrital, and reach 1100 m in the P12 and P10 wells in the far south and south-west part of the Chotts basin. Toward the North, the thickness of the Jurassic sediment does not exceed 500 m in

the P6 well and passes to 600 m at the P14 (Fig. 10) well where the Jurassic series represent the typical Nara Formation with three carbonate members indicating deposition in moderate deep marine environments of an external platform in southern and central Tunisian Atlas (Tlig, and Steinberg, 1980; Soussi, 2002).

In the Chott Fejej, the Melloussi, Boudinar, Bouhedma and Sidi Aich Formations, are thick of about 3000 m. Towards the North, the thickness decreases to reach 1100 m in the P6 and P14 wells (Figs. 9, 10). Laterally, in the Saharan Platform, the Tithonian-Barremian deposit passes to sandy facies with intercalations of clays, carbonates and evaporites named the "Wealdian facies" or Asfer Formation (Tlig, 2015). This Formation shows also important thickness variation ranging from 50 m in the P10 well to 600 m in the P11 well. From Aptian to Maastrichtian, the deposit is mainly made up of carbonates, clays and rare evaporitic series, outcropping widely in the two sides of the Fejej structure (Marie et al., 1984; Fakraoui, 1990; Fakraoui et al., 1991; Abdallah et al., 2000). The Aptian-Maastrichtian deposits typifies a transgressive cycle, showing several variations of thickness, facies and gaps. The Aptian–Late Turonian series are exposed along all the Southern Chotts range. The thickness changes from East to West showing a clear reduction towards the East (Ghanmi and Potfaj, 1991) and important thickness variations from South to North (Gharbi et al., 2015; Amri Tanfous et al., 2017; this work). These sequences have a 170 m–thickness in the eastern part of the SCR, in which the Turonian Guettar bar covers, via Cenomanian conglomeratic horizon, the Albian alternation. Towards the West, the maximum thickness of the Orbata and Zebbag Formations is recognized in the Fom El Argoub section (550 m). This E-W variation, is similarly observed along the NCR. In the eastern part (Zemlet el Beida), the Coniacian–Turonian Aleg Formation unconformably overlies the Aptian Barrani bar. Westward, at the Jebel Naimia, the thickness of the Aptian–Turonian series is about 95 m for the Orbata Formation and 215 m for the Zebbag Formation. Westward, At the khanguet Besbessa, the Orbata Formation shows a thickness of 120 m and the Zebbag Formation is about 390 m (Abdallah et al., 1995; Gharbi et al., 2013; El Amari et al., 2016). Toward the western part of the Range, the maximum thickness measured in the Khanguet El Asker is about 130 m for the Aptian–Orbata Formation and of 650 m for the Conomanian–Turonian Zebbag Formation. In subsurface, the P13 well crosses 1000 m of the Aptian-late Turonian series. Towards the South and North, the thickness of the Aptian-late Turonian series changes. It is about 550 m in the Saharan Platform (500 m in the P10, 550 m in P12 and 630 m in P3 wells) and about 870 m in the southern Central Tunisia (855 m in P6 and 885 m in the P14 wells).

In southern Atlassic domain of Tunisia, the Coniacian–Santonian Aleg Formation shows an important thickness and facies variations in both N-S and E-W striking. Along the SCR, this Formation is a 100 m–thick well exposed on the southern part of the Tebaga of Kebili mountains. Far South, the petroleum wells crossed 600 m of the Aleg formation. It is formed by two units, evaporitic with a lagoonal character at the base and carbonate at the top. Toward the North of the Chotts basin, the Aleg Formation is developed on the northern limbs of the NCR structures and even in two limbs of Jebel Berda. From East to West the variations are clear with 25 m at the Jebel Naimia, 130 m in Khanguet Besbessa (Abdallah, 1989; El Amari et al., 2016). Far west, the Aleg Formation have a thickness of 260 m at the Khanguet El Asker, 300 m at the Jebel Morra and reduced to 135 m at the Jebel Sidi Bou Hellel. At the outcrop, the thickness of the Aleg formation reached its maximum at Jebel Berda structure (450 m). Northward along the northern structures, the same variation is further accentuated with 450 m Jebel Berda and 1000 m in the P13 well (Fig. 10). The upper part of Cretaceous Formations shows the same thickness variations, although its rate of difference has been considerably reduced. The Campanian – Maastrichtian Berda Formation shows respectively 200 m in the Oued Batoum, 330 m in the Jebel Kherfane and 420 m at the P13 well.

4.2. Seismic data

Seismic interpretation is carried out in a structural goal. Nine seismic profiles are used and interpreted, for which some of them are composite ones. Seismic profile data are used to join the fault systems and/or tectonics block with the above-described lateral variations of thickness and facies of the sedimentary sequences. These Seismic profiles were correlated by many oil wells (Figs. 8-10 for more details) as well as the stratigraphic sequences established by field work. In the study area, three main parts are studied (northern, central and southern) based on the seismic data availability and structural style. We chose individual nine main stratigraphic limits (LS) that we considered to be important for the comprehension of the sedimentary history in addition to tectonics configuration. These three-selected areas well covered by the nine selected seismic profiles and well calibrated in north by P5, P6, P13 and P14; while in the central part by P1, P2, P4 and P7 and in the south P3, P8, P9, P10, P11 and P12 (Fig. 3). For confidentiality reasons, the seismic profiles are labelled as LSx independently from their original names.

The seismic section LS1 crossing the Chott Jerid area from SW to NE illustrates well the southwestern part of the Chotts basin (Fig. 11). This section proves a thickness increase of Jurassic and Lower Cretaceous sequences toward the middle part of the Chotts basin. Straight fault-plane reflections and

lithostratigraphic cutoffs constrains the location of high angle north-dipping normal fault systems. The hanging-wall of this faults is characterized by thick Jurassic (1000 m) and Lower Cretaceous (2000 m in P7 well) sequences. This thick block is probably deposited in deep area with high subsidence in response to the E-trending faults activity. In addition, this major E-trending fault seems to be sealed by the Cenomanian-Turonian Zebbag Formation. The northern side of this section shows also some normal faults that seem to be intra-basin growth normal faults. This section reproduces a good image of Lower Cretaceous beds highlighted by interference of irregular reflectors, which is probably the expression of growth strata pattern. A second section LS2, crossed by LS1 seismic profile, confirms this geometry of steep north-dipping normal faults associated to the growth of the Lower Cretaceous sequences (Fig. 12). These two seismic profiles (calibrated by nearby wells) argue for a basin showing in its southern side north-dipping normal faults responsible of the creation of deep zone during Lower Cretaceous in what is now the Chotts basin.

The lines LS3a and LS3b describe the eastern central part of basin (Figs. 13, 14). The central (north-south speaking) part of the Chotts basin, is constantly recording the initial Mesozoic extension. Along the LS3a section, the top Jurassic is highlighted by well-identified reflectors, which become discontinuous toward the southeastern part possibly related to the change of facies (Hlaïem, 1999). These reflectors are topped by irregular reflector of the Lower Cretaceous. This section shows an important normal faults system connecting the north dipping Chotts basin as revealed by the above-mentioned seismic profiles (Fig. 13). The key information given by this section is the increase to the Jurassic series thickness toward the East. This gives evidence that the Chotts basin, with major E-W striking governed by E-trending normal fault systems, shows also a lateral thickness variation from West to East parallel to the basin strike and confirm an eastward deepening.

This section (Fig. 13) shows that eastern part of NCR is marked by active salt tectonics. The northern side on this section exposes a remarkable thinning of Jurassic and Lower Cretaceous toward an area with internal confused reflection patterns. This configuration is characteristic of structure related salt rise as seen in the flanks of diapiric structures in many Atlantic domains (Hlaïem, 1999; Masrouhi et al., 2014a; Moragas et al., 2016, 2017). On this section, salt tectonics is probably controlled by Mesozoic normal faulting that is associated with rim synclines (salt mini-basins?) well developed in Jurassic thick sequences around the main salt body (Fig. 13).

Northward, the deep structure of the NCR is well illustrated by the NNW-SSE trending seismic reflection line LS3b (Fig. 14). Reflectors have been calibrated using projected data from P2 and P5 wells. To the

South, the Chott Fejej basin is formed above a normal movement of the fault system which delimited a deep graben beneath and associated with active subsidence during Jurassic and Early Cretaceous. To the North, a similar extensional structure created by listric normal faulting delimits a North-facing tilted blocks geometry beneath the Jebel Es Smaïa (Fig. 14).

Two long composite seismic profiles are derived along the Southern Atlas of Tunisia, i.e. the LS4a, approximately 100 km and the LS4b about 65 km-long (Figs. 15, 16). The seismic section LS4a crossing the SCR preserved subsurface structures of the Chott Fejej and the northern side of the Telemzane arch or the Telemzane High (Gabtni et al., 2012; Gabtni et al., 2013; Dhaoui et al., 2014). The southwestern part of the seismic section shows a considerable thinning of the Mesozoic series above the faulted Paleozoic basement? (Fig. 15). The faulted and probably folded Lower Paleozoic series form the anticline structure of Telemzane E-trending arch. These series are largely eroded and covered with an angular unconformity by the series of Permian-Carboniferous. The stratigraphic limit sited between the Telemzane high lower Paleozoic series and the Permian-Carboniferous series corresponds to the discontinuity known as Hercynian as suggested by the presence of Onlap figure in the seismic line LS4a. In the southwestern part of the study area, the P12 well shows that the upper part of the Paleozoic series (Carboniferous and Permian?) is absent due to the major uplift of Telemzane Arch during the Hercynian orogeny (Busson, 1967; Underdown and Redfern, 2007). The Permian-Carboniferous series showing chaotic seismic facies become increasingly thick towards the Chotts basin (Fig. 15). Eastward, the LS4b seismic section shows the remarkable geometry of the Hercynian unconformity between the Paleozoic and the Mesozoic sequences (Fig. 16). This unconformity correlates well with which was described far West in the Libyan domain (Boote et al., 1998). This section displays a remarkable angular unconformity configuration with successive toplap figures on the Paleozoic basement showing a northward slope varying from 2° to 4°. The toplap figure, the zones of truncated reflections and the correlation with nearby wells indicate successive unconformities. The succeeding unconformities are successively observed at the base of Triassic series, at the base of Jurassic and then at the base of Lower Cretaceous series filling all the basin (Fig. 16). The Early Triassic series are represented by continuous reflections and high amplitude can be interpreted as the continental sandstones of the Ouled Chebbi and Kirchaou Formations (Ben Ferjani et al., 1990). This Formation is overlain by chaotic seismic facies that corresponds to the Late Triassic evaporites. The Triassic series are affected by a set of an E-trending parallel normal faults dipping to the North defining a general system of half-graben containing smaller grabens. The major fault located in southern side of the Chott Fejej developing a structure of a tilted block.

The seismic reflection profile LS5, with Approximately E–W strike, is located in the Metlaoui plain between Sehib and Metlaoui mountains (Fig. 17). This line is presented here to discuss the NW–trending fault systems previously considered as the major faults controlled the formation and the growth of folds in South Tunisia (Zargouni and Tremoliere, 1981; Boukadi and Zargouni, 1991; Hlaïem, 1999; Bedir et al., 2001). On the basis of well continuous sub-horizontal reflectors on the both sides of the section and slightly deformed in the central, together with calibration by the P14 well, this section shows a series of NE-dipping steep normal faults characterizing different tectonics blocks with inconstant thick series. These faults are especially active during lower cretaceous and Coniacian–Santonian showing a clear variation of thickness. These data are consistent with an earlier hypothesis of inherited Mesozoic fault systems demonstrated by field data (see section 3). The reactivation of this fault system is clear in the fault located in the western part of this section, which show reverse movement. The LS6 seismic profile is also used to highlight at depth the Sehib structure. This profile shows in the forelimb a well identified reflector's truncated zone stating a thrust fault on which the Sehib fold is generated (Fig. 18). This section is well calibrated by the P13 well drilled in northern back-limb. This section is an individual one in the southern Tunisia that does not show any inherited closely controlling the subsequent compressional structure. The Sehib anticline seems to be a fault-related-fold developed during tertiary compressional events. However, the parallel near seismic line LS7 crossing the structure in the eastern part of the Sehib anticline shows that the northern part of the basin is formed above an inverted normal fault which delimits a deep graben beneath Sehib and associated with rapid subsidence during Early Cretaceous (Fig. 19). This unusual interconnection can be interpreted within two different scenarios: the eastern part of the Sehib anticline is separated from the main structure by NW–trending fault and forms in this case an isolated part like the Berda structures (Fig. 19). The second hypothesis is that the fault is formed above inherited systems, but in this case the LS6 section does not permit to highlight it. This last option can be sound if we know that the LS6 is not perpendicular to the structure and then to the deep inherited system if it exists.

As a summary of this section, the compilation of data coming from surface as well as the seismic and well data highlight a strong variation from North to South and from West to East, which coincide with the pre-Cenozoic shortening basin's architecture acquired during successive Mesozoic Tethyan extensional periods (Fig. 20). Data highlight a former Permian? to Early Cretaceous basin with multipart geometries that varies from East to West and from North to South. The Fig. 20 summarize a basin essentially controlled by earlier Mesozoic E- and NW–trending fault systems themselves related to the extensional period of the southern Tethyan margin. The E–trending faults are expressively active during Jurassic

controlling thick sequences especially in the eastern part of the basin (see [Fig. 20](#) for more details). The NW-trending faults are expressively active during Early Cretaceous showing a variation in the rate of sedimentation which become thicker in the central and western part of the basin. The E-trending fault systems, which generate a dominant E-W-striking basin (Chotts basin), which is bounded by two high Paleozoic zones: the Telemzame arch and Northern Chotts Range. Inside this main basin, distributary sub-basins established probably in conjunction with NW-trending faults. These results seem to be well integrated in the general Tethyan context ([Frizon de Lamotte et al., 2013](#)), which have demonstrate that the transform faults are E-W in the Atlantic and change to the NW-SE in the Alpine Tethys ([Rosenbaum, 2002](#)).

5. Balanced cross-sections and structural style

To illustrate the structural style and the regional-scale deformation as well as the evolution of the Southern Atlasic Front of Tunisia and its foreland, three long balanced cross-sections are constructed. Balanced cross-sections, further detailed below, are based on field extensive work, mapping of concerned areas, seismic reflection profiles across South Tunisia calibrated by detailed lithostratigraphic data from outcrops and petroleum wells. During processing, surface data are digitized from ASTER and DGEM, in which sections are built basically perpendicular to structures' axes to minimize the out of plane transport. Restoration and balancing of cross-sections are constructed referring the usual thrust tectonics concepts ([Suppe, 1983](#); [Suppe and Medwedeff, 1990](#); [Mitra, 2002](#); [Shaw et al., 2005](#); among others). Balancing and restoration were performed using the "Move 2017" structural modeling software (Midland Valley Inc.). For the cover, the thrust tectonics concepts fundamentally used in modeling are bed-length, thickness conservation and flexural slip algorithm ([Ramsay and Huber, 1987](#)). For the deep thrust tectonics' concepts and basement modeling, the restoration was based on the regional lithostratigraphic data ([Butler, 1989](#); [Butler et al., 2006](#)), well calibrated by petroleum wells.

As in the orogens worldwide, it was difficult to acknowledge the role of the basement tectonics control in the far Southern Atlas front of Tunisia in previous works. Currently, the outers part of orogens is recognized also to be controlled by thick-skinned deformation (e.g. [Mouthereau et al., 2007](#)). The basement-involved shortening as well as the tectonic style in the subsequent cross-sections, are constructed based on methods and/or natural examples recently published ([Mouthereau et al., 2012](#), [Lacombe and Bellahsen, 2016](#)).

The NNW-SSE constructed [S1] section was made from the Sehib structure in the North to the Saharan platform in the South ([Fig. 21](#)). This 110 km-long section is pinned in the Chebket el Mtarkeb plain in the

North and the southwestern Kebili plain in the South. This section highlights at least two different zones with northern deformed side and a southern part not as much of deformation compared to northern one. The northern portion shows tightly juxtaposed folds with two observable south-verging anticlines separated by syncline hidden under thin quaternary cover. The Jebel Sehib and the Jebel Taferna Zitouna anticlines connect down on large north-dipping faults themselves connected on the Triassic evaporites (transition basement-sedimentary cover?). Between these anticlines, the syncline is connected to depth on short-cut, possibly in sedimentary cover, itself connected to the north-dipping fault of the Jebel Sehib (Fig. 21). Considering all the above-presented data and taken in account the regional depth of dispositional accommodation zone and rate (Marshak and Wilkerson, 1992), the southern thrust -named in this work the Taferna thrust- involves possibly a high inherited basement structure beneath the Taferna–Zitouna. This thrust appears to be connected down on a major basement normal fault active during Mesozoic Tethyan rifting and controlled deep northern depositional area (regarding Taferna–Zitouna series) at least until lower Cretaceous times. Approximately the southern two-thirds of the section correspond to the Chott El Jerid basin and the northern part of the Saharan platform (Fig. 21). The Chott El Jerid basin, the Jebel Tebaga Kebili and the Saharan Platform structures are closely connected down on major north-dipping faults. From dense and abundant data (seismic reflection lines and petroleum wells), these structures are intended to connect three major normal faults controlling a thick Mesozoic sedimentary basin beneath this zone. The calculated data show that basement is at about a depth of 2 km (below sea level) in the Saharan platform, 5 km beneath the Chott El Jerid basin and about 3 km at Taferna–Zitouna (Fig. 21). Data are consistent with the hypothesis that the Taferna–Zitouna could be located on inherited basement high and forms the boundary between two different sedimentary depositional area. The present construction shows a model in which the major faults are preserved as normal and connected down onto a major sub-horizontal detachment at a depth of ~5 km beneath the southern part. The present model is also adequate when further combined with large geometrical sequences attitude. The hanging-wall, of the Chotts basins faults, reflects a pleasant reproduction of gently kinked rollover anticlines widely known in passive margin context (McClay, 1989; Brun and Nalpas, 1996). The hanging-wall rollover geometries were described in literature as closely controlled by the dips of the bounding faults (condition fulfilled in Chotts region) and amount of extension. In the southern Atlassic Front, a recent study of Gharbi et al. (2015), was estimated a rate of extension of about 6.5 km in Chotts region during rift stage from Permian? to Early Cretaceous. The conservation of layers length -based on the absence of observation of internal layers deformation- excepting the evaporitic Triassic series, reveals an estimation of shortening of about 7.6 km (~7.45%)

with 2.8 km (~2.7%) as deep shortening and 4.8 km (~4.7%) as shallow shortening. The shortening is widely dissimilar from a deformed northerly zone with 7.1 km shortening amount (~6.95%) and southerly almost undeformed zone with 0.5 km shortening amount (~0.5%). The Paleozoic high, defined in this study as delimited by major basement faults, had the role of an accommodation zone between the northern region with thick and thin-skinned tectonic style and the southern region showing preserved inherited extensional structures of the southern Tethyan margin (Fig. 21).

A second 107-km-long balanced cross-section [S2] is constructed from northwest to southeast crossing the following structures: Jebel Berda, Jebel Sif El Laham, Jebel Barrani, the Chotts basin, Jebel Tebaga Fatnassa and the Saharan Platform (Fig. 22). This NW-SE section is pinned in the Berda plain in the North and the southwestern El Hamma plain in the South. Similarly, this section shows two structurally distinct styles from North to South. The northern part, which we considered all the region located in the north of the NCR, corresponds to narrow anticline structures forming an E-W-striking major curved narrow range (Fig. 3). This part shows that shortening in the basement and cover is accommodated by different structural organization. Two distinguished anticlines, regarding the general style of NCR folds, bounded by south-dipping north-verging back-thrusts with moderate offset. The Jebel Berda as well as the Jebel Sif El Laham are plausibly up-lifted by these north-verging back-thrust systems. Back-thrust connects down on large thrust ones located between Basement and sedimentary cover and linked in the Triassic evaporitic layers at a depth of about 6 km (below sea level). This section records, from West to East, a net variation in depth of basement, which was computed in the last section at 3 km (below sea level). This geometry of shallow back-thrust connected down on major thrust systems shows a down-branches creating a basement short-cut beneath the Jebel Sif El Laham. We interpret this zone as Paleozoic high accentuated during Mesozoic extension which creating a horst structure beneath the NCR. During the subsequent Tertiary compression, the in-sequences forward propagation on the south-verging major deep thrusts attach on their footwall a north verging back-thrusts in favor of the formation of a deeper down triangle zone (Boyer and Elliot; 1982; Huyghe and Mugnier, 1992). These structures are very frequent at the front of many major thrust systems. They are described as the hinterland-dipping of normal faults that are inverted as foreland-directed thrusts, and early forelandward-dipping normal faults that are inverted as back-thrusts (McClay and Buchanan, 1992; Cooper et al., 1989; Bonini, et al., 2012). The restoration of this section suggests at least two phases of compressive deformation featured by the structures growth. The synclines, limited the two above-mentioned anticlines, shows a thin Oligocene-Miocene series unconformably resting onto the Eocene-Paleocene and Late Cretaceous sequences. Oligocene-Miocene series are themselves well deformed later such as the inverted ones

under the Berda back-thrust. This scenario reflects a first Middle to Late Eocene moderate tectonic compression and a post Lower Miocene important second event. The first one, is recognized as the Atlasic event and confirmed by recent studies in the southern Tunisia (Riley et al., 2011; Gharbi et al., 2015, El Amari et al., 2016, this work). This cross section suggests that during the Miocene–present-day successive compressions, the shortening within these structures is mainly accommodated by the inversion of north-dipping faults acting now as major thrusts deeply rooted in the basement with shallower thin-skinned south-dipping back-thrusts. Far South, the structure is approximately undeformed by Tertiary successive compressions and show a thick Jurassic and Lower Cretaceous sequences. The southern two-thirds of the section reproduce a large rollover anticline associated with the major north-dipping listric faults (Fig. 22). Mesozoic series filled a deep basin (basement is at about 7 km below sea level in the Chotts basin), broadly downthrown relative to northern and southern areas. This basin is also getting deeper from West to East compared to the western deep computed from the last section S1. This variation, from West to East, is perceived in term of deformation by decrease of shortening. The section reveals an estimation of shortening of about 3.65 km (~3.3%) with 1.5 km (~1.3%) as deep shortening and 2.15 km (~2%) as shallow shortening. The shallow shortening is widely different from the northern zone showing 3.4 km shortening amount (~3.1%) and southern zone practically undeformed with 0.25 km shortening amount (~0.2%). The Paleozoic High, beneath the NCR demonstrated by the present study which bounded by major basement faults, performs the role of an accommodation zone between north side region deformed by shallow thrust systems connected on deep basement ones and the southern region which is preserved as inherited extensional large rollover anticline formed during Mesozoic times (Fig. 22).

A short section [S3], compared to the western ones, is constructed eastward in the Chott Fejej zone based on well information and seismic reflection data (Fig. 23). This section is presented to confirm the same arrangement developed before and validates that the Cenozoic compressional deformation strongly decreases toward the East. This 43-km–long section illustrates the Chott Fejej basin as formed above a normal movement of the fault systems. These systems, which delimit a deep graben beneath the Chott Fejej, is associated with rapid subsidence during Jurassic and Early Cretaceous. To the North, a similar extensional structure created by listric normal faulting delimits a tilted blocks geometry with north-verging beneath the Jebel Es Smaïa. The constructed section shows a negligible shallow shortening of about 0.65 km (~1.4%).

6. Discussion

The Southern Atlas Front in Tunisia, which is the easternmost end of the Atlas Mountains across northwestern Africa, was presented for long time as developed in a strike-slip zone. The fold dispersal and the style of the Tunisian Southern Atlas were interpreted as linked to Gafsa and Negrine-Tozeur major NW-trending right lateral strike-slip fault systems (e.g. [Castany and Lapparent, 1952](#); [Abdeljaouad and Zargouni 1981](#); [Zargouni and Ruhland, 1981](#); [Zargouni et al., 1985](#); [Abbes and Zargouni 1986](#); [Fakraoui 1990](#)). The fold's distribution, in this model, was qualified of all of them as curved, right-stepping *en echelon* folds strictly related to the strike-slip fault systems. The second interpretation of structures is based on salt tectonics model, in which the Tertiary folding is preferentially nucleated on the preexistent diapir apex originally growing during Early Jurassic period ([Hlaiem et al., 1997](#), [Hlaiem, 1999](#); [Zouaghi et al., 2011](#)). The third thin-skinned interpretation with south verging thrust system ([Ahmadi et al. 2006](#); [Ahmadi et al., 2013](#); [Trigui et al., 2016](#)) interprets the anticline structures in this zone as developed and the result of ramp-related folding, deformed the sedimentary cover during a single tectonic phase (Miocene to recent). Recently new ideas have come to the fore, with introducing the role playing by structural heritage in the Cenozoic deformation of the belt ([Riley et al., 2011](#); [Gharbi et al., 2015](#); [El Amari et al., 2016](#); [Tanfous-Amri et al., 2017](#)). This study is integrated in the extensive present way for renewing the longstanding geological conceptions as it is the case in many West Mediterranean Alpine areas (e.g. [Vergés and Sàbat, 1999](#); [Faccenna et al., 2004](#); [Lacombe and Jolivet, 2005](#); [Deverchere et al., 2005](#); [Frizon de Lamotte et al., 2011](#); [Roure et al., 2012](#); [Bestani et al., 2015](#); [Lepretre et al., 2018a](#)).

The mixed thick- and thin-skinned structural style that typifies the southernmost deformed structures of the NCR in southern Tunisia is defined by shallower thrusts connected down on deep basement thrusts having a net variation throughout the NCR and the Chotts region. This study highlights a shortening amount, calculated from the above-exposed balanced sections, tending to decrease from North to South and from West to East. The total shortening (deep + shallow) amount varies from 7.6 km (this work, section S1) in the West to 3.65 km in the west-central part (this work, section S2), to 2 km ([Gharbi et al., 2015](#)) in central part and 0.65 km in the eastern part (this work, section S3). This variation coincides with variation of the top of basement which varies from 3.5 km below sea level in the S1 section, to 5.5 km in the S2 section and 6 km in S3 section. From North to South, the Cenozoic deformations are concentrated in the narrow zone of the NCR, which accommodate ~6.95 % of shortening and ~0.5% in the Chotts region. This variation coincides with net variation of total sequences thickness which is almost 4 km in the NCR and can reach 7.6 km of thickness in the Chotts. Additionally, this remarkable variation from North to South and from West to East, coincide with the pre-Cenozoic shortening basin architecture

acquired during successive Mesozoic Tethyan extensional periods. Surface and subsurface data analyzed before highlights a former Mesozoic basin with multipart geometries that varies from East to West and from North to South. To simplify the earlier Mesozoic basin's style, the general geometry can be described as follow: on a North-South strike, the central part consistent of the Chotts basin corresponds to a large rollover anticline structure developed during Jurassic and Early Cretaceous times. This large Mesozoic half-graben is developed between two high structures i.e. the NCR to the North and the Telemzane arch or the Telemzane High in the South (Gabtni et al., 2006; Gabtni et al., 2011; Dhaoui et al., 2014). The uplift of the Telemzane Arch occurred probably during the Hercynian orogeny (Busson, 1967; Underdown and Redfern, 2007) attested in southern Tunisia and Libya by remarkable large-scale unconformity of the Mesozoic series on an E-trending preexistent Paleozoic arch (Boote et al., 1998). On the northern side of the Chotts basin mega-rollover, an uplifted zone is defined by the present study as the horst structure in the basement along the NCR. This high zone is uplifted during Mesozoic times probable by south-dipping normal faults connecting down on the major listric large north-dipping fault systems. On an East-West strike evolution, this basin is deeper during Jurassic period in its eastern part showing Jurassic sequences with more than 2 km of thickness with southward dipping. The western side is not as much of deep and shows a north-dipping 1 km-thick Jurassic sequences. During Early Cretaceous an inversion of subsidence is started, and the western and central parts display deeper zone than the eastern part showing 2 km of sequences thickness near Kebili to the West and almost 1 km in the East near El Hamma region (see compilation on Fig. 20).

The central part of the NCR shows the outcrop of the only outcropping salt diapir called the Hadifa diapir, which is located in the western side of Hadifa graben (Fig. 3). Although the present work is not focused on this central part, the seismic section LS3a (Fig. 13) close to this area shows clearly the development of salt structures. The development of this diapir seems to be localized at the side of Mesozoic half graben. Triassic salt activity is probably initiated during Jurassic times and accentuated during Lower Cretaceous times. Salt structures in this central part look to be closely associated and triggered during continuous sedimentary loading. In addition, Hlaïem (1999), in the far west of the NCR (Bouhlef structure, Fig. 2), shows the emplacement of salt structure, in which salt migration is associated to synsedimentary normal faults during Jurassic and Lower Cretaceous times. Recently, Gharbi et al. (2015) were similarly highlight a salt activity beneath the Beidha structure (close to Hadifa salt diapir). Authors establish a scenario, in which northward salt migration is controlled by inherited normal faults responsible for the emplacement of the Beidha salt pillow.

Finally, the Southern Tunisian Fold-and-Thrust belt, which was interpreted for long time without incorporation of its structural inheritance, is comparable to many others famous Cenozoic belts governed by basin inversion and reactivation of inherited systems (Bonini et al., 2012; Espurt et al., 2014): e.g. the Provencal orogenic system (e.g. Roure and Colleta, 1996; Lacombe and Mouthereau, 2002; Espurt et al., 2012; Bestani et al., 2016), the Western Alps (e.g. Tricart and Lemoine, 1986; Bellahsen et al., 2012, 2014; Jourdon et al., 2014; Boutoux et al., 2014), the Apennines (e.g. Ghisetti and Vezzani, 1997; Scisciani et al., 2014) and the Pyrenees (e.g. Guimera, 1984; McClay et al., 2004; Mouthereau et al., 2014; Diaz et al., 2018).

The southern Atlasic domain of Tunisia appears to have strong influence of inherited former Mesozoic E-and NW-trending fault systems, as well as the sedimentation rate which reveals a dominant E-W-striking basin developed during successive rifting phases from Permian? to Early Cretaceous periods. The NCR's folded structures are interpreted to be formed above the E-trending inherited faults. These systems are established as inherited deep basement fault systems shaped an E-W high and deep zones during Mesozoic times. This E-trending systems is particularly active during Jurassic period. This period is characterized by the development of the Chotts basin limited by two major high Paleozoic zones (Telemzane arch in the South and the NCR high in the North). During Cenozoic compressions, the north-dipping major faults are inverted and subsequently acting as major deep thrust systems. The south-dipping ones are recognized as north-verging back-thrusts with moderate offset (i.e. Berda, Sif El Laham, Beidha) connecting down on large thrust ones with a strong coupling between Basement and sedimentary cover located in the Triassic evaporitic layers. During Cenozoic compressions, this system is responsible for the chain building and is expressly active during Miocene and Pliocene N-trending compression attested by thick Mio-Pliocene syntectonic strata. The NW-trending system is particularly active during Early Cretaceous time controlling a strong thickness variation of Early Cretaceous sequences. During Cenozoic compressions, this system is particularly responsible for the curved nature of this chain, which probably acts as transfer areas between the wide-ranging E-W strike major structures. The Chotts basin, which, until today, was considered as a large anticline related to the Cenozoic compressional events, is presumably a large rollover anticline preserved since the earlier rifting stage. The tertiary compressional front will be, therefore, repositioned in the front of NCR Front, and not in the SCR as it was previously accepted.

7. Conclusions

- The Southern Atlas Front in Tunisia is characterized by mixed thick- and thin-skinned structural style defined by shallow thrusts rooted down into deep basement thrusts with a preferential decollement level located within the Triassic evaporites. The E-trending N-dipping thrusts are recognized as major deep thrust systems. The S-dipping thrusts are recognized as N-verging back-thrusts with moderate offset connecting down on large north-dipping ones. A NW-trending system is also recognized, responsible for the curved nature of this chain, acting as transfer areas between the wide-ranging E-trending major structures. The intermediate areas are characterized by salt tectonics predominantly active during Jurassic and Lower Cretaceous, in response to active synsedimentary normal faults associated with differential sedimentation loading.
- Along the belt, the total shortening amount tends to decrease from North to South and from West to East. The calculated shortening varies from 7.6 km (~7.45%) in the West to 3.65 km in West-central part, about 2 km in central area to insignificant (0.65 km) in the eastern part. From north to south, deformations are concentrated in the narrow zone of the Northern Chotts Range, which accommodate almost 7% of deformation and less than 1% in the Chotts region.
- The strong variation from North to South and from West to East, coincide with the pre-Cenozoic shortening basin's architecture acquired during Mesozoic Tethyan evolution. This former evolution generates a main Jurassic E-trending basin (Chotts basin) bounded by two high Paleozoic zones. Inside the main basin, distributary sub-basins are recognized in relation to NW-trending faults mainly active during Cretaceous period. The Chotts basin appears, therefore, as well-preserved Mesozoic rift rollover structures taking place on the southern border of Atlasic orogen.
- The compilation of data, in this paper, suggests that the Front of the southeasternmost end of the Atlasic domain is distinguished by the reactivation of inherited fault systems acting during alpine orogeny as southward low-dip thrusts. The foreland shows a structural singularity to be formed by a preserved almost unmodified and well exposed large Mesozoic rollover structure.

Acknowledgments

The Authors gratefully acknowledge the ETAP (Tunisian National Oil Company) for granting permission to publish the data used in this work. Discussions with Nicolas Espurt, Hakim Gabtni, Riadh Ahmadi, Ferid Dhahri and Francois Roure contributed many different perspectives on this research. Midland Valley is acknowledged for providing academic license of Move for structural modeling. This work was financially supported by the Tunisian Ministry of Higher Education and Scientific Research (CERTE, Geo-resources

780 Laboratory funding) and the French Ministry of Foreign Affairs grant through French Embassy in Tunisia
781 (CMCU program). This work is a contribution to the program of the French National Research Agency
782 (ANR) through the A*MIDEX OT-Med project (nANR-11-LABX-0061) funded by the French Government
783 «Investissements d'Avenir» (n ANR-11-IDEX-0001-02). We are grateful to Olivier Lacombe, Charlotte
784 Fillon and two anonymous reviewers for providing constructive reviews that substantially improved the
785 structure, the scope and the focus of this paper.

786

787 **References**

- 788 Abbas, C., Abdeljaouad, S., Ben Oueddou, H., 1986. Geologic map of El Hamma 1:100000, sheet n°74,
789 National Geological survey, National Office of Mines, Tunisia.
- 790 Abbas, C., Zargouni, F., 1986. Anatomie d'un couloir de décrochements: le couloir de Hadifa (Chaîne
791 Nord des Chotts-Tunisie). *Rev. Sc. de la Terre*, vol. 4, Tunisie.
- 792 Abdallah, H., 1989. Les transgressions du Crétacé moyen entre les jeux tectoniques et les montées
793 eustatiques (Sud Tunisien). *Geobios*, 22, 1, 83–94.
- 794 Abdallah, H., Memmi, L., Damotte, R., Rat, P., Magniez, Jannin F., 1995. Le Crétacé supérieur de la chaîne
795 Nord des Chotts (Tunisie du centre-sud): Biostratigraphie et comparaison avec les régions voisines.
796 *Cretaceous Research*. 16, 5, 487–538.
- 797 Abdallah, H., Sassi, S., Meister, C., Souissi, R., 2000. Sequence stratigraphy and paleogeography at the
798 Cenomanian-Turonian boundary in the Gafsa-Chotts region, central Tunisia. *Cretaceous Research*, 21, 1,
799 35-106.
- 800 Abdeljaouad, S., Zargouni, F., 1981. Mise en évidence d'une tectonique intracretace dans le secteur de
801 Zemlet el Beidha (chaîne des Chotts). 1er Congr. Nat. Sci. Terre, Tunis.
- 802 Ahmadi, R., Ouali, J., Mercier, E., Mansy, J.-L., Van-Vliet Lanoë, B., Launeau, P., Rhekhiss, F., Rafini, S.,
803 2006. The geomorphologic responses to hinge migration in the fault-related folds in the Southern
804 Tunisian Atlas. *Journal of Structural Geology*, 28, 4, 721–728.
- 805 Ahmadi, R., Mercier, E., Ouali, J., 2013. Growth-strata geometry in fault-propagation folds: A case study
806 from the Gafsa basin, southern Tunisian Atlas. *Swiss Journal of Geosciences*, 106, 1, 91–107.
- 807 Allen, P. A., Allen, J. R., 1990. *Basins analysis: Principles and Applications*, 2d: New York, Wiley–Backwell,
808 560 p.
- 809 Bahrouni, N., Bouaziz, S., Soumaya, A., Ben Ayed, N., Attafi, K., Houla, Y., El Ghali, A., Rebai, N., 2014.
810 Neotectonic and seismotectonic investigation of seismically active regions in Tunisia: A multidisciplinary
811 approach. *Journal of Seismology*, 18, 2, 235–256.
- 812 Bahrouni, N., Houla, Y., Soussi, M., Boughdiri, M., Ali, W.B., Nasri, A., Bouaziz, S., 2016. Discovery of
813 Jurassic ammonite-bearing series in Jebel Bou Hedma (South-Central Tunisian Atlas): Implications for

814 stratigraphic correlations and paleogeographic reconstruction. *Journal of African Earth Sciences*, 113, 01,
815 101–113.

816 Bally, A.W., 1983. Seismic expression of structural styles. *AAPG studies in geology* 15, 3, 29 p.

817 Beauchamp, W., Allmendinger, R.W., Barazangi, M., Demnati, A., El, Alji M., Dahmani, M., 1999. Inversion
818 tectonics and the evolution of the High Atlas Mountains, Morocco, based on a geological-geophysical
819 transect. *Tectonics*, 18, 163–184.

820 Bedir, M., Boukadi, N., Tlig, S., Ben Timzal, F., Zitouni, L., Alouani, R., Slimane, F., Bobier, C., Zargouni, F.,
821 2001. Subsurface Mesozoic Basins in the Central Atlas of Tunisia, tectonics, sequence deposit distribution
822 and hydrocarbon potential. *A. A. P. G. Bulletin* 85, 885–907.

823 Bellahsen, N., Jolivet, L., Lacombe, O., Bellanger, M., Boutoux, A., Garcia, S., Mouthereau, F., Le Pourhiet,
824 L., Gumiaux, C., 2012. Mechanisms of margin inversion in the external Western Alps: Implications for
825 crustal rheology. *Tectonophysics*, 560-561, 62–83.

826 Bellahsen, N., Mouthereau, F., Boutoux, A., Bellanger, M., Lacombe, O., Jolivet, L., Rolland, Y., 2014.
827 Collision kinematics in the western external Alps, *Tectonics*, 33, 1055–1088.

828 Benaouali-Mebarek, N., D. Frizon de Lamotte, E. Roca, R. Bracène, J. L. Faure, W. Sassi, and F. Roure
829 (2006), Post-Cretaceous kinematics of the Atlas and Tell systems in central Algeria: Early foreland folding
830 and subduction-related deformation, *C. R. Geosci.*, 338, 115–125.

831 Ben Ayed, N., 1980. Le rôle des décrochements E-W dans l'évolution structurale de l'Atlas tunisien. *Bull.*
832 *Soc. Geol. France*, 1, 29–32.

833 Ben Ayed, N., 1993. Evolution tectonique de l'avant-pays de la chaîne alpine de Tunisie du début du
834 Mésozoïque à l'Actuel, *Ann. Mines Geol.*, Editions du Service géologique de Tunisie, no 32, 286 p.

835 Ben Ferjani, A., Burollet, P., Mejri, F., 1990. *Petroleum Geology of Tunisia*. Entreprise Tunisienne
836 d'activités pétrolières Ed., 194 p.

837 Ben Ismail, M. H., 1982. Le Trias et le Jurassique inférieur et moyen évaporitiques de l'extrême Sud
838 tunisien: étude de sondages profonds et de terrain, synthèses paléogéographiques. thesis 3ème cycle,
839 University Paris VI, 180 p.

840 Ben Ismail, M. H., 1991. Les bassins mésozoïques (Trias à Aptien) du sud de la Tunisie: stratigraphie
 841 intégrée, caractéristiques géophysiques et évolution géodynamique. Es-sciences thesis, University Tunis
 842 II, 446 p.

843 Ben Youssef, M., Biely, A., Memmi, L., 1985. La Formation Orbata (Aptien) en Tunisie méridionale.
 844 Précisions biostratigraphiques nouvelles. Notes du Service Geologique de Tunisie, 51, 105–120.

845 Ben Youssef, M., Peybernes, B., 1986. Données micropaléontologiques et biostratigraphiques nouvelles
 846 sur le Crétacé inférieur marin du Sud tunisien. Journal of African Earth Sciences, 5, 217–231.

847 Bestani, L., Espurt, N., Lamarche, J., Bellier, O., Hollender, F., 2016. Reconstruction of the Provence Chain
 848 evolution, southeastern France. Tectonics, 35, 6, 1506–1525.

849 Bestani, L., Espurt, N., Lamarche, J., Floquet, M., Philip, J., Bellier, O., Hollender, F., 2015. Structural style
 850 and evolution of the Pyrenean-Provence thrust belt, SE France. Bull. Soc. Geol. France, 186, 223–241.

851 Billi, A., Faccenna, C., Bellier, O., Minelli, L., Neri, G., Piromallo, C., Presti, D., Scrocca, D., Serpelloni, E.,
 852 2011. Recent tectonic reorganization of the Nubia–Eurasia convergent boundary heading for the closure
 853 of the western Mediterranean. Bull. Soc. Geol. Fr. 182, 279–303.

854 Bonini, M., Sani, F., Antonielli, B., 2012. Basin inversion and contractional reactivation of inherited
 855 normal faults: A review based on previous and new experimental models: Tectonophysics, 522–523, 55–
 856 88.

857 Boote, D. R. D., Clark-Lowes, D. D., Traut, M. W., 1998. Palaeozoic petroleum systems of North Africa, in
 858 Macgregor, D.S., Moody R.T.J., and Clark-Lowes, D.D., Ed. Petroleum geology of North Africa: Geological
 859 Society, London, Special Publication. 132, 7–68.

860 Bosworth, W., Guiraud, R., Kessler, L. G., 1999. Late Cretaceous (ca. 84 Ma) compressive deformation of
 861 the stable shelf of northeast Africa (Egypt); farfield stress effects of the “Santonian event” and origin of
 862 the Syrian arc. Geology, 27, 633–636.

863 Bouaziz, S., Barrier, E., Soussi, M. M., Turki, M.M., Zouari, H., 2002. Tectonic evolution of the northern
 864 African margin in Tunisia from paleostress data and sedimentary record. Tectonophysics, 357, 227–253.

865 Bouaziz, S., Barrier, E., Turki, M. M., Tricart, P., 1999. La tectonique permomésozoïque (anté-Vraconien)
 866 dans la marge sud téthysienne en Tunisie méridionale. Bull. Soc. Geol. France, 170, 1, 45–56.

867 Boughdiri, M., Cordey, F., Sallouhi, H., Maalaoui, K., Masrouhi, A., Soussi, M., 2007. Jurassic radiolarian-
868 bearing series of Tunisia: biostratigraphy and significance to western Tethys correlations. *Swiss J. Geosci.*
869 100, 431–441.

870 Boukadi, N., 1989. Disharmonic folds and dispersion of fold axes in strike-slip fault corridor: example of
871 Moulares fold, Tunisia, *Comptes Rendus - Academie des Sciences, Serie II*, 309, 20, 2105–2110.

872 Boukadi, N., Zargouni, F., 1991. Intersection of structural directions in the Tunisian Atlas: example of the
873 Mrhila and Labeid “tectonic knot”. *Comptes Rendus - Academie des Sciences, Serie II*, 312, 5, 529–534.

874 Boutoux, A., Bellahsen, N., Lacombe, O., Verlaguet, A., Mouthereau, F., 2014. Inversion of pre-orogenic
875 extensional basins in the external Western Alps: Structure, microstructures and restoration. *Journal of*
876 *Structural Geology*, 60, 13–29.

877 Boyer, S.E., Elliott, D., 1982. Thrust systems. *A.A.P.G. Bull.* 66, 1196–1230.

878 Bracène, R., Frizon de Lamotte, D., 2002. Origin of intraplate deformation in the Atlas system of western
879 and central Algeria: From Jurassic rifting to Cenozoic-Quaternary inversion, *Tectonophysics*, 357, 207–
880 226.

881 Briki, H., Ahmadi, R., Smida, R., Rekhiss, F., 2018. Structural evolution and tectonic style of the Tunisian
882 central Atlas; role of inherited faults in compressive tectonics (Ghoualguia anticline). *Tectonophysics*,
883 731-732, 48–63.

884 Brun, J. P., Nalpas, T., 1996. Graben inversion in nature and experiments. *Tectonics*, 15,3, 677–687.

885 Burollet, P.F., 1956. Contribution à l'étude stratigraphique de la Tunisie centrale. *Annales des Mines et*
886 *de la Géologie (Tunisie)*, vol. 18, 350 p.

887 Burollet, P. F., 1973, Importance des facteurs salifères dans la tectonique tunisienne, *In* Livre Jubilaire de
888 M. Solignac, *Ann. Mines Géol, Tunis*, no. 26, p. 111–120.

889 Burollet, P. F., 1991. Structures and tectonics of Tunisia. *Tectonophysics* 195, 359–369.

890 Busson, G., 1967, *Le Mésozoïque Saharien, 1ère partie: l'extrême Sud Tunisiens*, Paris: C.N.R.S., 194 p.

891 Busson, G., 1970. *Le Mésozoïque saharien. 2ème partie: Essai de synthèse des données des sondages*
892 *algéro-tunisiens: 2 tomes*, éd. C.N.R.S. Géol. N°. 11. Paris, 810p.

893 Butler, R. W. H., Tavarnelli, E., Grasso, M., 2006. Structural inheritance in mountain belts: An Alpine–
894 Apennine perspective. *Journal of structural geology*, 28, 11, 1893–1908.

895 Butler, R.W.H., 1989. The influence of pre-existing basin structure on thrust system evolution in the
896 western Alps. In: Cooper, M.A., Williams, G.D. (Eds.), *Inversion Tectonics*. Geological Society of London,
897 Special Publication, 44, pp. 105–122.

898 Castany, G., 1954. L'accident sud-tunisien, son age et ses relations avec l'accident sud-atlasique algerien.
899 C. R. Acad. Sci. Paris, 238, 916.

900 Castany, G., Lapparent, A. F., 1952. Discordances et phases de plissement dans l'extreme sud tunisien et
901 en Tripolitaine. *Bull. Soc. Geol. France*, 6, 1, 253–259.

902 Chihi, L., 1992. Seismotectonic study in Central and Southern Tunisia. *Tectonophysics*, 209, 1–4, 175–178.

903 Chihi, L., Ben Haj Ali, M., Ben Ayed, N., 1992. Structural significance and mechanism of folding in the
904 Chott chain, southern Tunisia. Analogy with folds associated with the Sbiba strike-slip fault, central
905 Tunisia, *Comptes Rendus - Academie des Sciences, Serie II*, 315, 10, 1245–1252.

906 Cooper, M. A., Williams, G. D., De Graciansky, P. C., Murphy, R.W., Needham, T., De Paor, D., Stoneley,
907 R., Todd, S. P., Turner J. P., Ziegler, P. A., 1989. Inversion tectonics – a discussion. In: Cooper, M.A.,
908 Williams, G. D. (eds.), *Inversion Tectonics*. Geological Society of London, Special Publication, 44, 335–347.

909 Coward, M. P., 1996. Balancing sections through inverted basins. *In* Buchanan, P.G. and Nieuwland, D.A.
910 (eds.) *Modern Developments in Structural Interpretation Validation and Modelling*. Geological Society,
911 London, Special Publication, 99, 51–78.

912 Dahlstrom, C. D. A., 1969. Balanced cross sections, *Canadian Journal of Earth Sciences*, 6, 4, 743–757.

913 DeCelles, P.G., Giles, K.N., 1996. Foreland basin systems. *Basin Research*. 8, 105–123.

914 Deverchere, J., Yelles, K., Domzig, A., De Lepinay, M., B., Bouillin, J. P., Gaullier, V., Bracene, R., Calais, E.,
915 Savoye, B., Kherroubi, A., Le Roy, P., Pauc, H., Dan, G., 2005. Active thrust faulting offshore Boumerdes,
916 Algeria, and its relations to the 2003 Mw 6.9 earthquake. *Geophys. Res. Letters*, 32, L04311.

917 Dhaoui, M., Gabtni, H., Jallouli, C., Jleilia, A., Mickus, K. L., Turki, M. M., 2014. Gravity analysis of the
918 Precambrian basement topography associated with the northern boundary of Ghadames Basin (southern
919 Tunisia). *Journal of Applied Geophysics*, 111, 299–311.

920 Dhahri, F., Boukadi, N., 2010. The evolution of pre-existing structures during the tectonic inversion
921 process of the Atlas chain of Tunisia. *Journal of African Earth Sciences*, 56, 139–149.

922 Dhahri, F., Boukadi, N., 2017. Triassic salt sheets of Mezzouna, central Tunisia: new comments on Late
923 Cretaceous halokinesis and geodynamic evolution of the Northern African margin. *Journal of African*
924 *Earth Sciences*, 129, 318–329.

925 Diaz, J., Vergés, J., Chevrot, S., Antonio-Vigil, A., Ruiz, M., Sylvander, M., Gallart, J., 2018. Mapping the
926 crustal structure beneath the eastern Pyrenees. *Tectonophysics*, 744, 296–309.

927 Dlala, M., 1992. Seismotectonic study in Northern Tunisia. *Tectonophysics*, 209, 1-4, 171–174.

928 Dlala, M., Hfaiedh, M., 1993. Le séisme du 7 novembre 1989 à Metlaoui (Tunisie méridionale): une
929 tectonique active en compression. *C. R. Acad. Sci. Paris Série II*, 317, 10, 1297–1302.

930 El Amari, E. A., Gharbi, M., Youssef, M. B., Masrouhi, A., 2016. The structural style of the southern
931 Atlasic foreland in northern Chotts range in Tunisia: field data from Bir Oum Ali structure. *Arabian*
932 *Journal of Geosciences*, 9:389.

933 Espurt, N., Hippolyte, J. C., Saillard, M., Bellier, O., 2012., Geometry and kinematic evolution of a long-
934 living foreland structure inferred from field data and cross section balancing, the Sainte-Victoire System,
935 Provence, France. *Tectonics*, 31, TC4021

936 Espurt, N., Hippolyte, J. C., Kaymakci, N., Sangu, E., 2014. Lithospheric structural control on inversion of
937 the southern margin of the black sea basin, central pontides, Turkey. *Lithosphere*, 6, 1, 26-34.

938 Faccenna, C., Piromallo, C., Crespo-Blanc, A., Jolivet, L., Rossetti, F., 2004. Lateral slab deformation and
939 the origin of the western Mediterranean arcs. *Tectonics*, 23, 1, TC1012, 1–21.

940 Fakraoui, M., 1990. Etude stratigraphique et structurale des chaînes de Chotts : Evolution géométrique et
941 cinématique liée à l'accident Sud-Atlasique. 3ème cycle Thesis, University, Tunis II, 231p.

942 Fakraoui, M., Rabiaa, C., Abbes, C., Zargouni, F., Bensalem, H., 1991. Geologic map of Bir Rakeb 1:100000,
943 sheet n°73, National Geological survey, National Office of Mines, Tunisia.

944 Frifita, N., Mickus, K., Zargouni, F., 2019. Gravity contribution to the geological study of grabens in the
945 northwest region of Tunisia including the Kef Basin region. *Journal of African Earth Sciences*, 150, 811–
946 822.

947 Frizon de Lamotte, D., Leturmy, P., Missenard, Y., Khomsi, S., Ruiz, G., Saddiqi, O., Guillocheau, F.,
 948 Michard, A., 2009. Mesozoic and Cenozoic vertical movements in the Atlas system (Algeria, Morocco,
 949 Tunisia): an overview. *Tectonophysics* 475, 9–28.

950 Frizon de Lamotte, D., Raulin, C., Mouchot, N., Wrobel-Daveau, J.-C., Blanpied, C., Ringenbach, J.-C.,
 951 2011. The southernmost margin of the Tethys realm during the Mesozoic and Cenozoic: Initial geometry
 952 and timing of the inversion processes. *Tectonics*, 30, TC3002.

953 Frizon de Lamotte, D., Tavakoli-Shirazi, S., Leturmy, P., Averbuch, O., Mouchot, N., Raulin, C.,
 954 Leparmentier, F., Blanpied, C., Ringenbach, J.-C., 2013. Evidence for Late Devonian vertical movements
 955 and extensional deformation in northern Africa and Arabia: Integration in the geodynamics of the
 956 Devonian world. *Tectonics*. 32, 1-16.

957 Frizon-de-Lamotte, D., Saint-Bezar, B., Bracene, R., Mercier, E., 2000. The two main steps of the Atlas
 958 building and geodynamics of the western Mediterranean. *Tectonics* 19, 740–761.

959 Gabtni, H., Jallouli, C., Mickus, K.L., Turki, M.M., 2013. Geodynamics of the Southern Tethyan Margin in
 960 Tunisia and Maghrebian domain: New constraints from integrated geophysical study. *Arabian Journal of*
 961 *Geosciences*, 6, 1, 271–286.

962 Gabtni, H., Jallouli, C., Mickus, K.L., Bédir, M., 2011. Geodynamics of the Southern Tethyan Margin in
 963 Tunisia and Maghrebian domain: new constraints from integrated geophysical study. *Arabian Journal of*
 964 *Geosciences*, 6, 271–286.

965 Gabtni, H., Jallouli, C., Mickus, K.L., Turki, M.M., Jaffal, M., Keating P., 2012. Basement structure of
 966 southern Tunisia as determined from the analysis of gravity data: Implications for petroleum exploration.
 967 *Petroleum Geoscience*, 18, 2, 143–152.

968 Gabtni, H., Jallouli, C., Mickus, K.L., Zouari, H., Turki, M.M., 2006. The location and nature of the
 969 Telemzan High-Ghadames basin boundary in southern Tunisia based on gravity and magnetic anomalies.
 970 *Journal of African Earth Sciences*, 44, 3, 303–3013.

971 Ghanmi, M., Potfaj, M., 1987. Geologic map of Kebili 1:100000, sheet n°81, National Geological survey,
 972 National Office of Mines, Tunisia.

973 Ghanmi, M., Potfaj, M., 1991. Données stratigraphiques sur la chaîne de Tebaga de Kebili Jebel Aziza
 974 (Tunisie méridionale): conséquences paléogéographiques et tectoniques, *Notes Serv. Geol. Tunis*. 58, 21–
 975 28.

976 Ghanmi, M., Potfaj, M., Mahjoub, K., 1988. Geologic map of Oglat Merteba 1:100000, sheet n°82,
977 National Geological survey, National Office of Mines, Tunisia.

978 Gharbi, M., Bellier, O., Masrouhi, A., Espurt, N., 2014. Recent spatial and temporal changes in the stress
979 regime along the southern Tunisian Atlas front and the Gulf of Gabes: New insights from fault kinematics
980 analysis and seismic profiles. *Tectonophysics*, 626, 120–136.

981 Gharbi, M., Masrouhi, A., Espurt, N., Bellier, O., Amari, E. A., Ben Youssef, M., Ghanmi, M., 2013. New
982 tectono-sedimentary evidences for Aptian to Santonian extension of the Cretaceous rifting in the
983 Northern Chotts range (Southern Tunisia). *Journal of African Earth Sciences*, 79, 58–73.

984 Gharbi, M., Espurt, N., Masrouhi, A., Bellier, O., Amari, E.A. 2015. Style of Atlasic tectonic deformation
985 and geodynamic evolution of the southern Tethyan margin, Tunisia. *Marine and Petroleum Geology*, 66,
986 801–816.

987 Ghisetti, F., Vezzani, L., 1997. Interfering paths of deformation and development of arcs in the fold-and-
988 thrust belt of the central Apennines (Italy). *Tectonics*, 16, 3, 523-536.

989 Guimerà, J. 1984. Palaeogene evolution of deformation in the northeastern Iberian Peninsula. *Geological*
990 *Magazine*, 121, 413–420.

991 Guiraud, R., Bosworth, W., 1997. Senonian basin inversion and rejuvenation of rifting in Africa and
992 Arabia: synthesis and implications to plate-scale tectonics. *Tectonophysics* 282, 39–82.

993 Guiraud, R., Bosworth, W., Thierry, J., Delplanque, A., 2005. Phanerozoic geological evolution of Northern
994 and Central Africa: an overview. *Journal of African Earth Sciences*, 43, 83–143.

995 Hamai, L., Petit, C. Abtout, A. Yelles Chaouche, A.; Déverchère, J. 2015. Flexural behaviour of the north
996 Algerian margin and tectonic implications. *Geophys. J. Int.*, 201, 1426–1436.

997 Hlaïem, A., Biju-Duval, B., Vially, R., Laatar, E., M'Rabet, A., 1997. Burial and thermal history modelling of
998 the Gafsa-Metlaoui intracontinental Basin (southern Tunisia): Implications for petroleum exploration.
999 *Journal of Petroleum Geology*, 20, 4, 403–426.

1000 Hlaïem, A., 1999. Halokinesis and structural evolution of the major features in eastern and southern
1001 Tunisian Atlas. *Tectonophysics* 306, 79–95.

1002 Huyghe. P., Mugnier, J. L. 1992. Short-cut geometry during structural inversion: competition between
1003 faulting and reactivation. *Bull. Soc. Géol. France*, 163, 691–700.

1004 Jaillard, E., Bouillin, J. P., Ouali, J., Dumont, T., Latil, J. L., Chihaoui, A., 2017. Albian salt-tectonics in
 1005 Central Tunisia: Evidences for an Atlantic-type passive margin. *Journal of African Earth Sciences*, 135,
 1006 220–234.

1007 Jaillard, E., Al Yacoubi, L., Reboulet, S., Robert, E., Masrour, M., Bouchaou, L., Giraud, F., El Hariri, K.,
 1008 2019. Late Barremian eustacy and tectonism in the western High Atlas (Essaouira-Agadir Basin),
 1009 Morocco. *Cretaceous Research*, 93, 225–244.

1010 Jolivet, L. Faccenna, C., 2000. Mediterranean extension and the Africa-Eurasia collision. *Tectonics*,
 1011 2000TC900018.

1012 Jourdon, A., Rolland, Y., Petit, C., Bellahsen, N., 2014. Style of Alpine tectonic deformation in the
 1013 Castellane fold-and-thrust belt (SW Alps, France): Insights from balanced cross-sections. *Tectonophysics*,
 1014 633, 1, 143–155.

1015 Kamoun, F., Peybernès, B., Ciszak, S., Calzada, S., 2001. Triassic palaeogeography of Tunisia.
 1016 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 172, 223-242.

1017 Khomsi, S., Bédir, M., Soussi, M., Ben Jemia, M.G., Ben Ismail-Lattrache, K., 2006. Mise en évidence en
 1018 subsurface d'événements compressifs Eocène moyen-supérieur en Tunisie orientale (Sahel): Généralité
 1019 de la phase atlasique en Afrique du Nord. *C. R. Géosci.*, 338, 1-2, 41–49.

1020 Khomsi, S., Ben Jemia, M.G., Frizon de Lamotte, D., Maherssi, C., Echihi, O., Mezni, R., 2009. An overview
 1021 of the Late Cretaceous-Eocene positive inversions and Oligo-Miocene subsidence events in the foreland
 1022 of the Tunisian Atlas: structural style and implications for the tectonic agenda of the Maghrebien Atlas
 1023 system. *Tectonophysics*, 475, 38–58.

1024 Laaridhi-Ouazaa, N., 1994. Etude minéralogique et géochimique des épisodes magmatiques mésozoïques
 1025 et miocènes de la Tunisie. Thèse Doc. Etat, University Tunis II, Tunisia, 466 p.

1026 Letouzey, J., 1990. Fault reactivation, inversion and fold-thrust-belt, in Letouzey, ed. *Petroleum and*
 1027 *tectonics in mobile belts*. Paris, Editions Technip, 101–128.

1028 Lacombe, O., Bellahsen, N., 2016. Thick-skinned tectonics and basement-involved fold-thrust belts:
 1029 Insights from selected Cenozoic orogens., *Geological Magazine*, 153, 5-6, 763–810.

1030 Lacombe, O., Mouthereau, F., 2002. Basement-involved shortening and deep detachment tectonics in
1031 forelands of orogens: Insights from recent collision belts (Taiwan, Western Alps, Pyrenees). *Tectonics*, 21,
1032 4, 10–30.

1033 Lacombe, O., Jolivet, L., 2005. Structural and kinematic relationships between Corsica and the Pyrenees-
1034 Provence domain at the time of the Pyrenean orogeny. *Tectonics*, 24, 1, TC1003, 1–20.

1035 Leprêtre, R., Missenard, Y., Barbarand, J., Gautheron, C., Juvie, I., Saddiqi, O., 2018b. Polyphased
1036 Inversions of an Intracontinental Rift: Case Study of the Marrakech High Atlas, Morocco. *Tectonics*, 37, 3,
1037 818–841.

1038 Leprêtre, R., De Lamotte, D.F, Combier, V., Gimeno-Vives, O., Mohn, G., Eschard, R., 2018a. The Tell-Rif
1039 orogenic system (Morocco, Algeria, Tunisia) and the structural heritage of the southern Tethys margin.
1040 *BSGF - Earth Sciences Bulletin*, 189, 2, Art. Nb. 2018009.

1041 Louhaichi, M. A., Tlig, S., 1993. Tectonique synsédimentaire des séries post-Barrémiennes au Nord-Est de
1042 la chaîne Nord des Chotts (Tunisie méridionale). *Géologie Méditerranéenne*, XX, 1, 53–74.

1043 Mahjoub, K., Fakraoui, M., 1990. Geologic map of Ksar El Askar 1:100000, sheet n°72, National Geological
1044 survey, National Office of Mines, Tunisia.

1045 Marie, J., Trouve, P., Desforges, G., Dufaure, P., 1984. Nouveaux éléments de paléogéographie du
1046 Crétacé de Tunisie. *Notes Mém. Comp. France Pétrol.*, 19.

1047 Marshak, S., and Wilkerson, M. S., 1992. Effect of overburden thickness on thrust belt geometry and
1048 development. *Tectonics*, 11, 560-566.

1049 Masrouhi, A., Bellier, O., Koyi, H., Vila, J. M., Ghanmi, M., 2013. The evolution of Lansarine-Baouala salt
1050 canopy in North African Cretaceous passive margin in Tunisia. *Geological Magazine*, 150 (5), 835–861.

1051 Masrouhi, A., Bellier, O., Koyi, H., 2014b. Geometry and structural evolution of Lorbeus diapir,
1052 northwestern Tunisia: polyphase diapirism of the North African inverted passive margin. *Int J Earth Sci*
1053 *(Geol Rundsch)*, 103, 881–900.

1054 Masrouhi, A., Bellier, O., Youssef, M.B., Koyi, H., 2014a. Submarine allochthonous salt sheets: gravity-
1055 driven deformation of North African Cretaceous passive margin in Tunisia e Bled Dogra case study and
1056 nearby salt structures. *Journal of African Earth Sciences*, 97, 125–142.

1057 Masrouhi, A., Ghanmi, M., Ben Slama, M.-M., Ben Youssef, M., Vila, J. M., Zargouni, F., 2008. New
 1058 tectono-sedimentary evidence constraining the timing of the positive tectonic inversion and the Eocene
 1059 Atlasic phase in northern Tunisia: Implication for the North African paleo-margin evolution. *C. R.*
 1060 *Geosciences*, 340, 771–778.

1061 Masrouhi, A., Ghanmi, M., Ben Youssef, M., Vila, J. M., Zargouni, F., 2007. Definition of a thrust nappe
 1062 with two Palaeogene units in the Lansarine plateau (northern Tunisia): a new structural element of the
 1063 Tunisian Atlas and revaluation of the Tertiary compression phases. *C. R. Geosciences*, 339, 441–448.

1064 Masrouhi, A., Koyi, H., 2012. Submarine “salt glacier” kinematics of Northern Tunisia, a case of Triassic
 1065 salt mobility in North African Cretaceous passive margin. In: Alsop, G.I., Archer, S.G., Hartley, A., Grant,
 1066 N.T., Hodgkinson, R. (Eds.), *J. Geol. R. Salt Tectonics, Sediments and Prospectivity*. Geological Society,
 1067 London, Special Publications, 363, 579–593.

1068 Martín-Martín, J.D., Vergés, J., Saura, E., Moragas, M., Messenger, G., Baqués, V., Razin, P., Grélaud, C.,
 1069 Malaval, M., Joussiaume, R., Casciello, E., Cruz-Orosa, I., Hunt, D.W., 2017. Diapiric growth within an
 1070 Early Jurassic rift basin: The Tazoult salt wall (central High Atlas, Morocco). *Tectonics*, 36, 1, 2–32.

1071 Mattoussi Kort, H., Gasquet, D., Ikenne, M., Laridhi Ouazaa, N., 2009. Cretaceous crustal thinning in
 1072 North Africa: Implications for magmatic and thermal events in the Eastern Tunisian margin and the
 1073 Pelagic Sea. *Journal of African Earth Sciences*, 55, 5, 257–264.

1074 McClay, K. R., Buchanan, P. G., 1992. Thrust faults in inverted extensional basins. In McClay, K. R. (ed.)
 1075 *Thrust Tectonics*, Chapman and Hall, London, 93–104.

1076 McClay, K.R., 1989. Analogue models of inversion tectonics. In: Cooper, M.A., Williams, G.D. (Eds.),
 1077 *Inversion tectonics*. Geological Society of London, Special Publication, 44, 41–62.

1078 McClay, K., Munoz, J. A., García-Senz, J., 2004. Extensional salt tectonics in a contractional orogen: A
 1079 newly identified tectonic event in the Spanish Pyrenees. *Geology*, 32, 9, 737–740.

1080 Missenard, Y., Taki, Z., Frizon de Lamotte, D., Benammi, M., Hafid, M., Leturmy, P., Sebrier, M., 2007.
 1081 Tectonic styles in the Marrakesh High Atlas (Morocco): the role of heritage and mechanical stratigraphy.
 1082 *Journal of African Earth Sciences*, 48, 247–266.

1083 Mitra, M., 2002. Fold-accommodation faults, *A.A.P.G. Bulletin*, 86, 671–693.

1084 Moragas, M., Vergés, J., Nalpas, T., Saura, E., Martín-Martín, J.D., Messenger, G., Hunt, D.W., 2017. The
 1085 impact of syn- and post-extension prograding sedimentation on the development of salt-related rift
 1086 basins and their inversion: clues from analogue modelling. *Marine and Petroleum Geology*, 88, 985–
 1087 1003.

1088 Moragas, M., Vergés, J., Saura, E., Martín-Martín, J.D., Messenger, G., Merino-Tomé, Ó., Suárez-Ruiz, I.,
 1089 Razin, P., Grélaud, C., Malaval, M., Joussiaume, R., Hunt, D.W. 2018. Jurassic rifting to post-rift
 1090 subsidence analysis in the central High Atlas and its relation to salt diapirism. *Basin Research*, 30, 336–
 1091 362.

1092 Mouthereau, F., Tensi, J., Bellahsen, N., Lacombe, O., De Boisgrollier, T., Kargar, S. 2007. Tertiary
 1093 sequence of deformation in a thin-skinned/thick-skinned collision belt: The Zagros Folded Belt (Fars,
 1094 Iran). *Tectonics*, 26, 5, art. no. TC5006.

1095 Mouthereau, F., Lacombe, O., Vergés, J., 2012. Building the Zagros collisional orogen: Timing, strain
 1096 distribution and the dynamics of Arabia/Eurasia plate convergence. *Tectonophysics*, 532-535, 27–60.

1097 Mouthereau, F., Filleaudeau, P. Y., Vacherat, A., Pik, R., Lacombe, O., Fellin, M. G., Castelltort, S.,
 1098 Christophoul, F., Masini, E., 2014. Placing limits to shortening evolution in the Pyrenees: Role of margin
 1099 architecture and implications for the Iberia/Europe convergence. *Tectonics*, 33, 12, 2283–2314.

1100 Naji, C., Gharbi, M., Amri, Z., Masrouhi, A., Bellier, O., 2018a. Temporal and spatial changes of the
 1101 submarine Cretaceous paleoslope in Northern Tunisia, inferred from slump folds analysis. *Proceedings of*
 1102 *the Geologists' Association* Volume 129, Issue 1, p. 40-56.

1103 Naji, C., Masrouhi, A., Amri, Z., Gharbi, M., Bellier, O., 2018b. Cretaceous paleomargin tilted blocks
 1104 geometry in northern Tunisia: stratigraphic consideration and fault kinematic analysis. *Arabian Journal of*
 1105 *Geosciences*, 11, 583.

1106 Outtani, F., Addoum, B., Mercier, E., de Frizon Lamotte, D., Andrieux, J., 1995. Geometry and kinematics
 1107 of the south Atlas front, Algeria and Tunisia. *Tectonophysics*, 249, 233–248.

1108 Patriat, M., Ellouz, N., Dey, Z., Gaulier, J.M., Ben, Kilani H., 2003. The Hammamet, Gabes and Chotts
 1109 basins (Tunisia) a review of the subsidence history. *Sediment. Geol.* 156, 241–262.

1110 Ramsay, J. G., Huber, M. I., 1987. *The Techniques of Modern Structural Geology*, vol. 2, *Folds and*
 1111 *Fractures*, Pergamon Press, London, 391 p.

1112 Regaya, K., Laatar, S., Chaouachi, A., Zargouni, F., 1991. Geologic map of Metlaoui 1:100000, sheet n°65,
1113 National Geological survey, National Office of Mines, Tunisia.

1114 Riley, P., Gordon, C., Simo, J. A., Tikoff, B., Soussi, M., 2011. Structure of the Alima and associated
1115 anticlines in the foreland basin of the southern Atlas Mountains, Tunisia. *Lithosphere* 3, 1, 76–91.

1116 Rosenbaum, G., Lister, G. S., Duboz, C., 2002. Relative motions of Africa, Iberia and Europe during Alpine
1117 orogeny. *Tectonophysics*, 359, 117–129.

1118 Roure, F., Brun, J. P., Colletta, B., Van Den Driessche, J., 1992. Geometry and kinematics of extensional
1119 structures in the alpine foreland basin of southeastern France. *Journal of Structural Geology*, 14, 503–
1120 519.

1121 Roure, F., Casero, P., Addoum, B., 2012. Alpine inversion of the North African margin and delamination of
1122 its continental lithosphere. *Tectonics*, 31, TC3006.

1123 Roure, F., Colletta B., 1996. Cenozoic inversion structures in the foreland of the Pyrenees and Alps, in
1124 *Peri-Tethys Memoir 2: Structure and Prospects of Alpine Basins and Forelands*, Mem. du Mus. Natl. Hist.
1125 Nat., vol. 170, edited by P. A. Ziegler, Mus. Natl. Hist. Nat., Paris, 173–209.

1126 Said, A., Baby, P., Dominique, C., Ouali, J., 2011b. Structure, paleogeographic inheritance, and
1127 deformation history of the southern Atlas foreland fold and thrust belt of Tunisia. *Tectonics*, 30, TC6004.

1128 Said, A., Dominique, C., Baby, P., Ouali, J., 2011a. Active oblique ramp faulting in the Southern Tunisian
1129 Atlas, *Tectonophysics*, 499, 178–189.

1130 Saura, E., Verges, J., Martin-Martin, J. D., Messenger, G., Moragas, M., Razin, P., Grélaud, C., Joussiaume,
1131 R., Malaval, M., Homke, S., Hunt, D.W., 2014. Syn- to post-rift diapirism and minibasins of the Central
1132 High Atlas (Morocco): The changing face of a mountain belt. *Journal of the Geological Society*. 171, 1, 97–
1133 105.

1134 Scisciani, V., Agostini, S., Calamita, F., Pace, P., Cilli, A., Giori, I., Paltrinieri, W., 2014. Positive inversion
1135 tectonics in foreland fold-and-thrust belts: A reappraisal of the Umbria-Marche Northern Apennines
1136 (Central Italy) by integrating geological and geophysical data. *Tectonophysics*, 637, 218–237.

1137 Shaw, J., Connors, C., Suppe, J., 2005. Seismic interpretation of contractional fault-related folds, AAPG
1138 seismic atlas, *Stud. Geol. Tulsa Okla.*, 53, 1–156.

1139 Slimane, F., Rabia, C., Zarai, N., Ben Youssef, M., Zargouni, F., 1991. Geologic map of Gafsa 1:100000,
1140 sheet n°66, National Geological survey, National Office of Mines, Tunisia.

1141 Soumaya, A., Ben Ayed, N., Delvaux, D., Ghanmi, M., 2015. Spatial variation of present-day stress field
1142 and tectonic regime in Tunisia and surroundings from formal inversion of focal mechanisms: Geodynamic
1143 implications for central Mediterranean. *Tectonics*, 34, 6, 1154–1180.

1144 Soumaya, A., Ben Ayed, N., Rajabi, M., Meghraoui, M., Delvaux, D., Kadri, A., Ziegler, M., Maouche, S.,
1145 Braham, A., 2018. Active Faulting Geometry and Stress Pattern Near Complex Strike-Slip Systems Along
1146 the Maghreb Region: Constraints on Active Convergence in the Western Mediterranean. *Tectonics*, 37, 9,
1147 3148–3173.

1148 Soussi, M., 2002. Le Jurassique de la Tunisie atlasique: Stratigraphie, dynamique sédimentaire,
1149 Paléogéographie et intérêt pétrolier, vol. 157. Documents des Laboratoires de Géologie de Lyon, 363 p.

1150 Soussi, M., Ben Ismail, H., 2000. Platform collapse and pelagic seamount facies: Jurassic development of
1151 Central Tunisia. *Sediment. Geol.* 133, 93–113.

1152 Soussi, M., Niedźwiedzki, G., Tałanda, M., Drózd, D., Sulej, T., Boukhalfa, K., Mermer, J., Błażejowski, B.,
1153 2017. Middle Triassic (Anisian-Ladinian) Tejra red beds and Late Triassic (Carnian) carbonate sedimentary
1154 records of southern Tunisia, Saharan Platform: Biostratigraphy, sedimentology and implication on
1155 regional stratigraphic correlations. *Marine and Petroleum Geology*, 79, 222–256.

1156 Suppe, J., 1983. Geometry and kinematics of fault-bend folding. *Am. J. Sci.*, 283, 684–721.

1157 Suppe, J., Medwedeff, D. A., 1990. Geometry and kinematics of fault propagation folding. *Eclogae Geol.*
1158 *Helv.*, 83, 3, 409–454.

1159 Swezey, C. S., 1996. Structural controls on Quaternary depocentres within the Chotts Trough region of
1160 southern Tunisia. *Journal of African Earth Sciences*, 22, 335–347.

1161 Tanfous Amri, D., Bédir, M., Soussi, M., Azaiez, H., Zitouni, L., Hédi Inoubli, M., Ben Boubaker, K., 2005.
1162 Early halokinesis associated to the Jurassic rift faulting in Central Tunisia (Majoura-El Hfay area). *Comptes*
1163 *Rendus – Geosciences*, 337, 7, 703–711.

1164 Tanfous Amri, D., Dhahri, F., Soussi, M., Gabtni, H., Bedir, M., 2017. The role of E-W basement faults in
1165 the Mesozoic geodynamic evolution of the Gafsa and Chotts basins, south-central Tunisia. *Journal of*
1166 *Earth System Science*, 126, 7, 104.

1167 Teixell, A., Arboleya, M. L., Julivert, M., Charroud, M., 2003. Tectonic shortening and topography in the
1168 central High Atlas (Morocco). *Tectonics*, 22, 5, 10–51.

1169 Tlig, S., 2015. The Upper Jurassic and Lower Cretaceous series of southern Tunisia and northwestern
1170 Libya revisited. *Journal of African Earth Sciences*, 110, 100–115.

1171 Tlig, S., Steinberg, M., 1980. Précisions sur le paléoenvironnement de la série carbonatée du Jurassique
1172 supérieur du Sud de la Tunisie. *Bull. Soc. Géol. France*, 7, XXII, n 2, 233–242.

1173 Tricart, P., Lemoine, M., 1986. From faulted blocks to megamullions and megaboudins: Tethyan heritage
1174 in the structure of the western Alps. *Tectonics*, 5, 1, 95–118.

1175 Tricart, P., Torelli, L., Argnani, A., Rekhiss, F., and Zitellini, N., 1994, Extensional collapse related to
1176 compressional uplift in the Alpine chain off northern Tunisia (Central Mediterranean): Tectonophysics,
1177 238, 317–329.

1178 Trigui, H., Ahmadi, R., Ouali, J., Khalfi, C., Mercier, E., 2016. Evidence of fault-propagation folds in
1179 foreland basin: the case of Chemsî and Belkhir anticlines of southern Tunisian Atlas. *Arabian Journal of*
1180 *Geosciences*, 9, 18, 706.

1181 Underdown, R., Redfern, J., 2008. Petroleum generation and migration in the Ghadames Basin, North
1182 Africa: A two-dimensional basin-modeling study. *AAPG Bulletin*, 92, 1, 53–76.

1183 Van Hinsbergen, D., Vissers, R., Spakman, W., 2014. Origin and consequences of western Mediterranean
1184 subduction, rollback, and slab segmentation. *Tectonics*, 33, 4, 393–419.

1185 Vergés, J., Fernández, M., 2012. Tethys–Atlantic interaction along the Iberia–Africa plate boundary: The
1186 Betic–Rif orogenic system. *Tectonophysics*, 579, 144–172.

1187 Vergés, J.; Sàbat, F. 1999. Constraints on the Neogene Mediterranean kinematic evolution along a 1000
1188 km transect from Iberia to Africa. *Geological Society, London, Special Publications*, 156, 63–80.

1189 Vially, R., Letouzey, J., Bernard, F., Haddadi, N., Desforges, G., Askri, H., Boudjema, A., 1994. Basin
1190 inversion along the northern African margin: the Saharan Atlas (Algeria). In: Roure, F. (Ed.), *Peri-Tethyan*
1191 *Platforms*. Edition Technip, Paris, 79–118.

1192 Vila, J. M., J. Sigal, P. E. Coiffait, J. C. Lahondère, and Y. Quinif, 1975. Sur l’allochtonie du Tell sétifien: La
1193 fenêtre de la Koudiat Touachra et la nappe de Djemila, *Bull. Soc. Geol. France*, 17, 1173–1175.

1194 Vila, J. M., Sigal, J., Lahondère, J. C., 1993. Position en fenêtre de la série néritique constantinoise sous la
1195 nappe de Djemila: Observations nouvelles du Djebel Mazela (massif du Djebel Oum Settas, Algérie du
1196 Nord-Est), C.R. Acad. Sci., Ser. II, 317, 395–401.

1197 Zargouni, F, Termolières J. C., 1981. Déformations tectoniques postérieures au dépôt de la série du Segui
1198 (Plio-Villafranchien) dans l’Atlas méridional tunisien. Actes 1er Cong. Nation. Sci. Terre, Tunis.

1199 Zargouni, F., 1985. Tectonique de l’Atlas méridional de Tunisie, évolution géométrique et cinématique
1200 des structures en zone de cisaillement. Thèse d’Etat, Univ. Louis Pasteur, Strasbourg, 304 p.

1201 Zargouni, F., Rabiaa, M. C., Abbès, C., 1985. Rôle des couloirs de cisaillement de Gafsa et de Négrine-
1202 Tozeur dans la structuration du faisceau des plis des Chotts, éléments de l’accident sud-atlasique. C.R.
1203 Acad Sci. Paris, 301, 11, 831–883.

1204 Zargouni, F., Ruhland, M., 1981. Style des déformations du quaternaire récent lié au coulissement de la
1205 faille de Gafsa et chronologie des phases tectoniques de l’Atlas méridional de Tunisie. C. R. Acad. Sc.
1206 Paris, 292, 913–915.

1207 Ziegler, P.A., Cloetingh, S., van Wees, J.D., 1995. Dynamics of intra-plate compressional deformation: the
1208 alpine foreland and other examples. Tectonophysics, 252, 7–59.

1209 Zouaghi, T., Bedir, M., Inoubli, M.H., 2005. 2 D Seismic interpretation of strike-slip faulting, salt tectonics,
1210 and Cretaceous unconformities, Atlas Mountains, central Tunisia. Journal of African Earth Sciences, 43,
1211 464–486.

1212 Zouaghi, T., Ferhi, I., Bedir, M., Ben Youssef, M., Gasmi, M., Inoubli, M.H., 2011. Analysis of Cretaceous
1213 (Aptian) strata in central Tunisia, using 2D seismic data and well logs. Journal of African Earth Sciences,
1214 61, 38–61.

1215 Zouari, H., Turki, M.M., Delteil, J., 1990. Nouvelles donnees sur l'evolution tectonique de la chaîne de
1216 Gafsa. Bull. Soc. Geol. France, 8, 621–628.

1217 Zouari, H., Ouled Ghrib, A., Ouezdou, Ben, Zargouni, F.H., Chouachi, A., Laatar, E., 1991. Geologic map of
1218 El Ayacha 1:100000, sheet n°67, National Geological survey, National Office of Mines, Tunisia.

1219

Figure captions

Figure 1. Simplified structural sketch of west Mediterranean region with location of the Southern Atlas Front of Tunisia with a base map is produced using ASTER and DGEM data.

Figure 2. (a) Simplified lithostratigraphic column of the Southern Atlassic area, derived from outcrop and petroleum wells data with thickness variation of Mesozoic sequences (minimum and maximum). (b) Simplified geologic map of the southern Tunisia showing the major E-trending structural ranges with location of the study area.

Figure 3. (a) Location of the studied seismic profiles (some of them are composite ones) of [Figs. 11-19](#) and the petroleum wells used in calibration, the base map is produced using ASTER and DGEM data. (b) Geologic map of the study area, data are derived from present work, [Abbes et al., 1986](#); [Ghanmi and Potfaj, 1987](#); [Ghanmi et al., 1988](#); [Mahjoub and Fakraoui, 1990](#); [Zouari et al., 1991](#); [Fakraoui et al., 1991](#); [Gharbi et al., 2015](#); [El Amari et al., 2016](#); revealing the detailed structures of southern Atlassic domain with location of surface geologic cross-sections of [Fig.4](#) and the long balanced cross-sections [S1-S2-S3] across the chain ([Figs. 21-23](#)).

Figure 4. Geologic cross sections prepared from surface data across the main structures of the study area, see [Fig. 3](#) for location.

Figure 5. (a) 3D Google earth WSW-looking view showing the central part of the Northern Chotts Range highlighting the interfering between E- and NW-trending fault systems with the location of the E-trending major thrust (Bir Oum Ali Structure) and Back-thrust (J. Sif Laham); the NW-trending acts as transfer areas between the wide-ranging E–W strike major structures. (b) West looking panoramic view of J. Sif Laham pointing the emplacement of the north-verging back-thrust between southern sub-horizontal Early Cretaceous sequences and northern folded Late Cretaceous sequences. West looking panoramic view of J. Bir Oum Ali showing the position of south-verging major thrust (incomplete reverse displacement from former normal throw). (c) Panoramic view, looking ENE, of reactivated normal fault (thickness variation from north to south) to thrust forming the fold above a triangular geometry established by north-verging back-thrust connected down on south-verging thrust. (d) East looking photo of the angular unconformity of Miocene–Pliocene series on Paleocene–Eocene sequences, which outcrops along 20 km of the Bir Oum Ali's northern limb.

Figure 6. Common preserved extensive deformation sealed within Cretaceous sequences in south Tunisia, (a) S-looking photo illustrating sealed normal faults within the Coniacian–Santonian sequences in

1250 the J. Haidoudi structure, (b) NW-looking photo of preserved normal faults within Hauterivian–Barremian
1251 sequences in the J. Beidha.

1252 **Figure 7.** E-looking panorama of an incompletely inversion of earlier normal fault as reverse observed
1253 within Late Cretaceous (Campanian) sequences in the northern side of the Bir Oum Ali structure.

1254 **Figure 8.** Lithostratigraphic correlation [C1] of regional extent showing the thickness variation from
1255 southwest to northeast. The correlation is given here without the major faults control with a primary
1256 objective to describe them in absence of structural adjustment (for this see [Fig. 20](#)).

1257 **Figure 9.** Lithostratigraphic correlation [C2] of regional extent showing the thickness variation across the
1258 central part of the study area. The correlation is given here without the major faults control with a
1259 primary objective to describe them in absence of structural adjustment (for this see [Fig. 20](#)).

1260 **Figure 10.** Lithostratigraphic correlation [C3] of regional extent showing the thickness variation across
1261 the eastern part of the study area. The correlation is given here without the major faults control with a
1262 primary objective to describe them in absence of structural adjustment (for this see [Fig. 20](#)).

1263 **Figure 11.** seismic profile LS1 (see [Fig. 3](#) for location), crossing the Chotts Jerid area from SW to NE,
1264 illustrates the southwestern part of the Chotts basin showing increase toward the middle part of the
1265 Chotts basin of Jurassic and Lower Cretaceous thickness associated to deep E–trending faults.

1266 **Figure 12.** seismic profile LS2, located in the central part of Chotts basin (see [Fig. 3](#) for location), shows a
1267 geometry of steep north-dipping normal faults associated to the growth strata pattern of the Lower
1268 Cretaceous sequences.

1269 **Figure 13.** seismic profile LS3a, located in the Eastern part of the Chotts basins (see [Fig. 3](#) for location),
1270 shows important normal fault systems relating the north dipping Chotts. The northern side exposes a
1271 remarkable thinning of Jurassic-Lower Cretaceous toward an area with internal confuse reflection
1272 patterns. This configuration is characteristically of structure related salt rise. salt tectonics is probably
1273 controlled by Mesozoic normal faulting that is associated with rim synclines (salt mini-basins?) well
1274 developed in Jurassic thick sequences around the salt body.

1275 **Figure 14.** seismic profile LS3b (see [Fig. 3](#) for location), Reflectors have been calibrated using projected
1276 data of P2 well onto the section. To the South, the Chott Fejej basin is formed above a normal movement
1277 of the fault systems which delimited a deep graben beneath Chott Fejej and associated with rapid
1278 subsidence during Jurassic and Early Cretaceous.

1279 **Figure 15.** 100 km-long composite seismic profile LS4a (see [Fig. 3](#) for location), from the Saharan
1280 platform to the Chotts basin crossing the Southern Chotts range, the southwestern part of the seismic
1281 section shows a considerable thinning of the Mesozoic series above the faulted Paleozoic series. the
1282 deep central part highlights intra-basin growth faulting associated to Jurassic and Lower Cretaceous
1283 growth strata pattern.

1284 **Figure 16.** 65 km-long composite seismic profile LS4b (see [Fig. 3](#) for location); southeastward, the seismic
1285 section shows a remarkable geometry indicating successive unconformities. The Hercynian unconformity
1286 between the Paleozoic and the Mesozoic sequences is well highlighted by successive toplap figures on
1287 the Paleozoic basement showing a northward slope varying from 2° to 4°.

1288 **Figure 17.** Seismic profile LS5 crossing the Metaloui plain (see [Fig. 3](#) for location), shows continuous sub-
1289 horizontal reflectors on the both side of the section and a little deformed area in the central part. This
1290 section shows a series of NE-dipping steep normal faults characterizing different tectonics blocks with
1291 inconstant thick series of Early Cretaceous and Coniacian–Santonian.

1292 **Figure 18.** Seismic profile LS6 (see [Fig. 3](#) for location), crossing the Sehib structure. The profile shows in
1293 the forelimb a well identified reflector's truncated zone stating a thrust fault above which the Sehib fold
1294 is formed.

1295 **Figure 19.** Seismic profile LS7 (see [Fig. 3](#) for location), crossing the structure in the eastern part of the
1296 Sehib anticline, shows that the north part basin is formed above inverted normal faults which bounds a
1297 deep graben beneath Sehib and associated with important subsidence during Early Cretaceous.

1298 **Figure 20.** schematic sketch collecting all data coming from seismic and wells together with surface data.
1299 Data highlight a dominant former Permian? to Early Cretaceous E-W–striking basin (Chotts basin)
1300 bounded by two high zones: the Telemzame arch and Northern Chotts Range, with multipart geometries
1301 that varies from east to west and from north to south (see text for more details).

1302 **Figure 21.** balanced cross-section [S1] across the western part of the southern Atlas Front of Tunisia, see
1303 [Fig. 3](#) for location.

1304 **Figure 22.** balanced cross-section [S2] across the central part of the southern Atlas Front of Tunisia, see
1305 [Fig. 3](#) for location.

1306 **Figure 23.** balanced cross-section [S3] across the eastern part of the southern Atlas Front of Tunisia, see
1307 [Fig. 3](#) for location.