Assignment[#] of 5-hydroxytryptamine receptor (HTR4) to human chromosome 5 bands q31-->q33 by in situ hybridization.

Sylvie Claeysen¹, Patrick Faye¹, Michèle Sebben¹, Stéphanie Lemaire², Joël Bockaert¹, Aline Dumuis¹ and Sylvie Taviaux^{2,*}

¹CNRS UPR 9023, Centre CNRS-INSERM de Pharmacologie-Endocrinologie (CCIPE), 141 rue de la Cardonille, 34094 Montpellier cedex 05, France

²CRBM-CNRS, 1919 route de Mende, BP 5051, 34033 Montpellier Cedex 1, France

* Author for correspondence

Phone: (33) 04 67-61-33-45 Fax: (33) 04 67-52-15-59

Email: taviaux@vega.crbm.cnrs-mop.fr

[#]To our knowledge this is the first time this gene has been mapped.

This work was supported by grants from the Centre National de la Recherche Scientifique, Laboratoires Fournier-Debat (Daix, France) and the Fondation pour la Recherche Médicale.

Rationale and significance

5-Hydroxytryptamine 4 receptors (5-HT₄R) were first characterized in 1988 (Dumuis *et al.*, 1988) in mouse colliculi neurons. Since then, it has been shown that 5-HT₄R mediates widespread effects in central and peripheral systems (Eglen *et al.*, 1995). These observations have stimulated the search and discovery of a very large series of selective and potent agonists and antagonists as well as the cloning of the receptor in rat and mouse brain (Eglen *et al.*, 1996; Gerald *et al.*, 1995; Claeyssen *et al.*, 1996). In human atria the concentration of 5-HT₄R and α_1 - and α_2 -adrenoceptors vary, depending on whether the patients have been treated with α -adrenergic-blocking agents or not (Kaumann *et al.*, 1994; Ouadid *et al.*, 1992). These observations suggest that a "cross-talk" may exist between the regulation of 5-HT₄R and α -adrenoceptors in human myocardium. To answer these questions, we cloned human 5-HT₄R (gene symbol: HTR4; Claeyssen *et al.*, 1997) from heart and mapped the chromosomal location by FISH.

Materials and methods

Human metaphase cells were prepared from phytohemagglutinin-stimulated lymphocytes. Probe was labeled with biotinyl-16-dUTP and FISH was performed as previously described (Taviaux and Demaille, 1993).

Probe name(s): h5-HT₄s

Probe type: cDNA

Insert size: 1.2kb

Proof of authenticity: sequencing

Gene reference: GDB:6381308; Claeyssen *et al.* (1997)

Results

Mapping data

Location: 5q31→q33

Number of cells examined: 52

Number of cells with specific signal: 1 (19), 2 (8), 3 (6), 4 (1) chromatids per cell

Most precise assignment: 5q32

Location of background signals (sites with >2 signals): none

References

- Claeysen S, Sebben M, Journot L, Bockaert J, Dumuis A: Cloning expression and pharmacology of the mouse 5-HT_{4L} receptor. *FEBS Letters* 398:19-25 (1996).
- Claeysen S, Faye P, Sebben M, Lemaire S, Bockaert J, Dumuis A: Cloning and expression of human 5-HT_{4S} receptors: effect of receptor density on their coupling to adenylyl cyclase. *Neuroreport* 8 issue 15 (1997)
- Dumuis A, Bouhelal R, Sebben M, Cory R, Bockaert J: A non classical 5-hydroxytryptamine receptor positively coupled with adenylate cyclase in the central nervous system. *Mol Pharmacol* 34:880-887 (1988).
- Eglen R, Wong EHF, Dumuis A, Bockaert J: Central 5-HT₄ receptors. *Trend Pharmacol* 16:391-398 (1995).
- Eglen RM, Hegde S: 5-Hydroxytryptamine (5-HT₄) receptors: physiology, pharmacology and therapeutic potential. *Exp Opin Invest Drugs* 5:373-388 (1996).
- Gerald C, Adham N, Kao HT, Olsen MA, Laz TM, Schechter LE, Bard JA, Vaysse PJ, Hartig PR, Branchek TA, Weinshank RL: The 5-HT₄ receptor: molecular cloning and pharmacological characterization of two splice variants. *EMBO J* 14:2806-2815 (1995).
- Kaumann AJ, Sanders L: 5-Hydroxytryptamine causes rate-dependent arrhythmias through 5-HT₄ receptors in human atrium-facilitation by chronic beta-adrenoceptor blockade. *Naunyn-Schmiedeberg's Arch Pharmacol* 349:331-337 (1994).
- Ouadid H, Seguin J, Dumuis A, Bockaert J, Nargeot J: Serotonin increases Ca⁺⁺ current in human atrial myocytes via the newly described 5-hydroxytryptamine₄ receptors. *Mol Pharmacol* 41:346-351 (1992).
- Taviaux AS, Demaille JG: Localization of human cell cycle regulatory genes CDC25C to 5q31 and WEE1 TO 11p15.3→p15.1 by fluorescence in situ hybridization. *Genomics* 15:194-196 (1993).

Figure legend

Figure 1

A) Hybridization of a human cDNA 5-HT₄ receptor probe to R-band human metaphases. Fluorescent spots are visualized at 5q31→q33. **B)** Ideogram of chromosome 5 homologue showing the fluorescent spot distribution to the 5q31→q33 band for 30 metaphases.