Phytochemical Screening and Polyphenolic Contents of *Dialium dinklagei* and *Diospyros monbuttensis*, Two Ivorian Medicinal Plants Used to Treat Malaria

Karim Tuo, Sylvain Béourou, André Offianan Touré, Karamoko Ouattara, Dieudonné Silué, Dominique Tano Konan, Marius Adagba, David Koffi, Stephane Yao, Joseph Djaman, et al.

► To cite this version:

Karim Tuo, Sylvain Béourou, André Offianan Touré, Karamoko Ouattara, Dieudonné Silué, et al.. Phytochemical Screening and Polyphenolic Contents of *Dialium dinklagei* and *Diospyros monbuttensis*, Two Ivorian Medicinal Plants Used to Treat Malaria. *Journal of Advances in Medical and Pharmaceutical Sciences*, 2015, 2 (4), pp.144-153. 10.9734/JAMPS/2015/13779 . hal-02509207

HAL Id: hal-02509207

<https://hal.science/hal-02509207>

Submitted on 17 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Phytochemical Screening and Polyphenolic Contents of *Dialium dinklagei* and *Diospyros monbuttensis*, Two Ivorian Medicinal Plants Used to Treat Malaria

**Tuo Karim^{1,2}, Béourou Sylvain^{2*}, Touré André Offianan², Ouattara Karamoko¹,
Silué Kigbafori Dieudonné³, Tano Konan Dominique², Adagba Marius⁴,
Koffi David^{1,2}, Yao Serge Stephane^{1,2}, Djaman Allico Joseph⁵
and Coulibaly Adama¹**

¹Pharmacodynamics Biochemical Laboratory, U.F.R. Biosciences, Felix Houphouet Boigny University,
Abidjan, 22 BP 582 Abidjan 22, Côte d'Ivoire.

²Department of Parasitology-Mycology, Malariology Unit, Institut Pasteur in Côte d'Ivoire, 01 Bp 490
Abidjan 01, Côte d'Ivoire.

³Department of Environment and Health, Swiss Center for Scientific Research in Côte d'Ivoire, 01 BP
1303 Abidjan 01, Côte d'Ivoire.

⁴Department Virus Epidemic, Institut Pasteur in Côte d'Ivoire, 01 Bp 490 Abidjan 01,
Côte d'Ivoire.

⁵Department of Medical Biochemistry and fundamental, Toxicology Unit, Institut Pasteur in Côte
d'Ivoire, 01 Bp 490 Abidjan 01, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. Author TK designed the study and wrote the protocol. Authors BS and TAO supervised the laboratory protocols and corrected the manuscript.

Authors OK, SKD and TKD corrected the manuscript. Authors YSS, AM and KD performed the statistical analysis and also wrote part of the manuscript. Authors DAJ and CA supervised the study.

All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMPS/2015/13779

Editor(s):

(1) Faiyaz Shakeel, Department of Pharmaceutics, King Saud University, Saudi Arabia.

(2) Dongdong Wang, Department of Pharmacognosy, West China College of Pharmacy, Sichuan University, China.

Reviewers:

(1) Alli Lukman Adewale, Medical Biochemistry, College of Health Sciences, University of Abuja, Nigeria.

(2) Norma Almaraz Abarca, Instituto Politécnico Nacional, México.

Complete Peer review History: <http://www.sciencedomain.org/review-history.php?id=888&id=36&aid=7803>

Original Research Article

**Received 3rd September 2014
Accepted 17th November 2014
Published 16th January 2015**

*Corresponding author: E-mail: be_sylvain@yahoo.fr, sylvainbeourou@pasteur.ci;

ABSTRACT

Introduction: Malaria is a deadly parasitic infection in its severe form. It is caused by an endoparasite protozoan transmitted by a mosquito vector. Despite considerable efforts to eradicate it, this endemic disease remains a public health problem because of its high frequency, severity and the consequences it generates. Furthermore, the existing treatments are increasingly ineffective; it is therefore urgent to find new anti-malarial drugs.

Objective: From the phytochemical screening and the determination of the polyphenolic content of *Dialium dinklagei* and *Diospyros monbuttensis* used in traditional medicine in Côte d'Ivoire to treat malaria, we identified families of chemical compounds that can guide us towards pharmacological studies. These studies allow us to identify new molecules which could be a possible alternative antimalarial reference used today because of their effectiveness in treating malaria.

Materials and Methods: From *Dialium dinklagei* and *Diospyros monbuttensis*, we obtained extracts decoction and hexane extracts, methanolic and aqueous extraction after three solvents of increasing polarity (n-hexane, methanol, water), respectively. The crude extracts obtained were tested for chemical identification of groups that may be the basis of the expected therapeutic effects. Moreover, the decoction and methanol extracts were used to determine the polyphenolic compounds.

Results: The results vary from one plant species to another and from one sample to another. Abundant presence of alkaloids in all samples and an absence of steroids in other extract except decoction extract of *Diospyros monbuttensis* were observed. Therefore, the other chemical groups are present in all plants in varying proportions but depending on the mode of extraction. A higher content of phenols and tannins in *Dialium dinklagei* over *Diospyros monbuttensis* was observed. However, the content of flavonoids of *Dialium dinklagei* is relatively lower compared to that observed in *Diospyros monbuttensis*.

Conclusion: The overall results of the phytochemical screening and polyphenolic content would partly explain the traditional healers' infatuation to these plants as antimalarials.

Keywords: Malaria; natural substances; chemical groups; *Dialium dinklagei*; *Diospyros monbuttensis*.

1. INTRODUCTION

The high population growth, the growing impoverishment and the lack of medical facilities are the major problems faced by the most of African populations [1]. Owing to this situation, they do not always have access to imported pharmaceuticals. According to Sandhya et al. [2] and Séguéna et al. [3], 80% of the world population relies on traditional medicine for their primary health care that requires the use of plant extract. Indeed, the use of traditional medicine is justified by the treatment of many infections against which the plants are used as effective as drugs imported by Africa [3-5]. Secondary metabolites present in these plants are not new to the medical properties that are attributed to them. The search for these secondary metabolites could be the first step towards the determination of active ingredients [6,7]. That is the reason why we have undertaken the phytochemical study of plants used in the Ivorian traditional medicine against malaria. In this context, *Diospyros monbuttensis* Gürk (Ebenaceae) and *Dialium dinklagei* (Caesalpinaceae) widely used by traditional

healers but little studied for their antimalarial activity have been selected.

Diospyros monbuttensis leaves have been reported to possess antibacterial and antimalarial properties [8-10]. The antimicrobial activities of the plant leaves extract was due to the presence of tannins [11,12]. Other constituents that have been reported include saponins, anthraquinones, cardiac glycosides and alkaloids [8]. *Dialium dinklagei* is a tree of Caesalpinaceae family with trunk cylindrical right. The sheets made up have an acid taste, the flowers are of color yellow and the bark is very thick [13]. Not previously reported from Ivory Coast. This species is distributed in West and Central Africa and has been reported from Angola, Cameroon, D.R. Congo, Gabon, Ghana, Guinea, Ivory Coast, Liberia, Nigeria and Sierra Leone [14].

The aim of this work was to investigate the correlation between the active ingredients present in these medicinal plants and their therapeutic effects, namely their use as antimalarial medicines.

2. MATERIALS AND METHODS

2.1 Plant Material

The plant material consisted of leaves of *Diospyros monbuttensis* and *Dialium dinklagei*. These are two plant species, which have been little studied for their antiparasitic activity, especially *Dialium dinklagei*. The leaves of *Diospyros monbuttensis* were collected in the month of January 2013 from Talahini-Sokoura in the Department of Sandegué (North-eastern Côte d'Ivoire). Those of *Dialium dinklagei* were collected in March 2013 in the District of Abidjan. Plants were adult and the harvest time was chosen the morning. Plants species were identified by Professor Ake-Assi from the "Centre National de Floristique", University of Felix-Houphouët-Boigny, Abidjan, Côte d'Ivoire. They were then dried in sunlight for one week at room temperature before being reduced to a fine powder by using mechanical grinder type IKAMAG. From the powder obtained after spraying the leaves, various crude extracts were prepared.

2.2 Preparation of Crude Extracts

First, from the fine powder obtained, decoctions of each plant were made to be as close as possible to the form of traditional use. Then we made three successive extractions with solvents of increasing polarity, according to the protocol of Békro et al. [15], Zirihé et al. [16] and Beourou et al. [17]. In the following order, we used three solvents of increasing polarity: hexane, methanol and water.

2.3 Decoction

The decoction was obtained by dissolving 20g of fine dry powder in 1L of distilled water. The whole was homogenized and boiled for 30minutes. The preparation was first wrung out in a square of white cloth, and then filtered successively twice on an absorbent cotton and once on a Whatman filter paper 3 mm. Filtrate 1 was thus obtained. The same operation was repeated with the residue by adding 1 liter of distilled water to obtain the filtrate 2. These two filtrates were gathered and evaporated in "Venticel" oven at 55°C. This series of operations led to the extract of decoction.

2.4 Extraction with Hexane

A sample (50 g) of fine dry powder were dissolved in 150 mL of hexane. The whole was vigorously agitated in a blender for 10minutes. The preparation was first wrung out in a square of white cloth, and then filtered successively with an absorbent cotton and filter paper. The first hexane filtrate was thus obtained (hexane filtrate 1). For filtrate 2, the same operation was repeated by addition of 150 mL of hexane on the residue found. This activity was repeated for filtrate 3. These three filtrates were then gathered and evaporated in the "Venticel" oven at 55°C. This operation led to a hexane extract.

2.5 Extraction with Methanol

After extraction with hexane, the residual powder was dried and then extracted with 150 mL of methanol. After 10 minutes of homogenization by stirring in a blender the extract was filtered through a piece of gauze for obtaining the methanolic filtrate 1. The same operation was repeated twice more to obtain the methanolic filtrates 2 and 3. All three filtrates were then pooled and evaporated in an oven "Venticel" at 55°C. This series of operations led to an extract called methanol extract.

2.6 Preparation of the Aqueous Extract

The same operations were repeated on the residual dry powder coming from the ethanolic extraction but using 150 mL of distilled water. We obtained the aqueous extract.

For the following tests, the aqueous and decoction extracts are dissolved in distilled water while the other extracts are dissolved in DMSO.

2.7 Phytochemical Screening

Phytochemical analysis was performed on the basis of tests for coloring characteristics to highlight the major chemical groups. It focused on decoction extracts, hexane, methanol and aqueous extracts of the studied plants. Various chemical groups were identified by reference to the methods described by Nemlin and Brunel [18], Békro et al. [15] and Bruneton [19].

The results have been classified according to: Highly positive: +++; Fairly positive: ++; weakly positive: +; Negative test: -

2.7.1 Saponins (Foam index)

Samples (0.1 g of dry extract) were dissolved in 10 mL of distilled water. The samples were shaken vigorously up and down for 30-45 seconds and then left for 15 minutes. The height of the foam was measured. Persistent foam for more than 1 cm high indicated the presence of saponins.

2.7.2 Steroids: Salkowski test

Five drops of concentrated H_2SO_4 were added to 1 mL of extract, the apparition of a red color indicated the presence of steroids.

2.7.3 Sterols and polyterpenes

Extracts (0.1 g) were dissolved in 1 mL of hot acetic anhydride in a capsule. The resulted solutions were poured and added with 0.5 mL H_2SO_4 . A violet coloration that turned in blue, and then in green revealed the presence of sterols and triterpenes.

2.7.4 Alkaloids

Two drops of Bouchard's reagent (reagent of iodine-iodide) were added to 1 mL of each extract. A red-brown precipitate indicated a positive reaction.

2.7.5 Polyphenols

A drop of alcoholic solution of 2% ferric chloride was added to 2 mL of extracts. A blue-blackish to green darkish coloration indicated a positive reaction.

2.7.6 Anthocyanins

Five milliliters of 10% H_2SO_4 was added to 5 mL of 5% extract, and then a base (five drops of 25% NH_4OH) was added. If the coloration is accentuated by acidification and then change into blue-purple in a basic environment, anthocyanins were present.

2.7.7 Leucoanthocyanins

One milliliter of extract was added to 5 mL of hydrochloric alcohol. The preparation was heated in the water-bath for 15 minutes. A cherry red or purplish coloration meant the presence of leucoanthocyanins.

2.7.8 Flavonoid

In a tube containing 3 mL of extract, a few drops of 10% NaOH were added. Appearance of yellow-orange color indicated the presence of flavonoids.

2.7.9 Quinones

An aliquot (0.1 g) of extract was dissolved in 5 mL of diluted HCl (1/5) and heated in a boiling water bath for 30 minutes, and then extracted with 20 mL of $CHCl_3$ after cooling. To the organic phase was added 0.5 mL of 50% NH_4OH diluted solution. The positivity of the reaction was indicated by a red to violet color.

2.7.10 Cardiotonic glycosides

This reaction is exothermic. It takes place in a glacial environment. To 2 mL of aqueous solution of extract were added 1 mL of glacial acetic acid and then 1 mL of concentrated sulfuric acid. Adding few drops of 2% $FeCl_3$ gave a green-bluish coloration, which is specific to cardiotonic glycosides.

2.7.11 Coumarins

To 5 mL of extract, 2 mL of hot water were added, and then the solution was shared between two test tubes. In the content, we added 0.5 mL of 25% NH_4OH . Under the UV light (366 nm), an intense fluorescence indicated the presence of coumarins.

2.7.12 Tannins

Two milliliters of water and few drops of 1% ferric chloride were added to 1 mL of extract. The appearance of a blue, blue-black or black coloration indicated the presence of gallic tannins, the green or dark green coloration showed the presence of catechic tannins.

2.8 Determination of Polyphenols

The polyphenol contents were determined for the foliar decoction and methanol extracts of *Diospyros monbuttensis* and *Dialium dinklagei*.

2.9 Determination of Total Phenols

The total phenolic content of plant extracts was determined by the Folin-Ciocalteu method [20,21]. To 0.5 mL of each plant extract of concentration 100 mg/mL were added 5 mL of

Folin-Ciocalteu reagent diluted 1/10 in distilled water and 4 mL of sodium carbonate (1M). The mixture was incubated at room temperature for 15 minutes. Then, the absorbance was registered at 765nm in a spectrophotometer against a blank. Gallic acid was used to construct a standard curve in order to calculate the phenol contents, which was prepared under the same conditions as above with a solvent mixture of methanol/water (50: 50, V/V) at concentrations ranging from 0 to 200 mg/mL. Experimentations were carried out in triplicate. The total phenolic contents of the extracts were expressed as gallic acid equivalents/g dry extract (mg GAE/g dry extract).

2.10 Determination of Total Flavonoids

The technique used to determine the contents of total flavonoids in the extracts was the colorimetric method of aluminum chloride [22,23]. And 0.5 mL (100 µg/mL) of each plant extract were taken, to which 1.5 mL of methanol, 0.1 mL of 10% aluminum chloride, 0.1 mL of potassium acetate (1 M) and 2.5 mL of distilled water were successively added. After incubation at room temperature for 30 minutes, the optical densities were measured in a spectrophotometer at 415 nm. A methanolic solution of quercetin in concentrations ranging from 0 to 100 µg/mL were used to construct a standard curve in order to calculate the flavonoid contents. Experimentations were carried out in triplicate. The contents of flavonoids were expressed as quercetin equivalents/g dry extract (mg QE/g dry extract).

2.11 Determination of Tannins

The tannin contents were calculated by using the modified vanillin method of Bainbridge et al. [24]. This method is based on the fact that the tannic acid (especially the flavan-3-ol) in the presence of the vanillin reagent (0.1 mg/mL) in an acid medium (sulfuric acid 70% (v/v) gives a red color whose maximum absorption is 500 nm. The standard solution consists of tannic acid (0.1 mg/mL) prepared in 98% methanol. A range of standard was prepared with a series of 6 tubes where the first represents a blank. In the other 5 tubes are introduced respectively 0.2; 0, 4; 0.6; 0.8; 1 ml of standard solution and complete the volume of each tube with 1 ml of distilled water. The test tube contains 1 ml of extract. To each of them are added 5 ml of vanillin reagent. After 30 minutes of incubation in the dark, the absorbance

was read at 500 nm against a blank (without extract). The tannic acid (1 mg/ml) was used to construct a standard curve in order to calculate the tannin contents. Experimentations were carried out in triplicate. The results were expressed as mg of tannic acid equivalents/g dry extract (mgTAE/g dry extract)

2.12 Statistical Analysis

Graphical representations of data were performed using the Graph Pad Prism 5.0 software (Microsoft, USA). The test results were expressed as mean $\pm$ SD. Multiple comparisons and determination of significance level are made by the univariate ANOVA followed by Tukey test. The differences were considered statistically significant at the 0.05 level.

3. RESULTS AND DISCUSSION

3.1 Plant Selection and the Output of the Extraction

The investigations were carried out with ethnobotanical approaches among stakeholders of traditional medicine, especially herbal traders operating on the markets of the District of Abidjan. These investigations resulted in the selection of *Diospyros monbuttensis* and *Dialium dinklagei*, widely used by Ivorian traditional pharmacopoeia and very little studied for their antimalarial activity. These species were mainly collected on the basis of indications by literature and information received from healers.

Decoctions produced the highest output (20%), and hexane was the smallest (1%). These results suggest the use of the method of extraction decoction as indicated for the preparation of extracts from these plants base (Figure 1). The very low output obtained with hexane shows that the studied plants contain very little non-polar compounds. This result is similar to the one obtained by Bekro et al. [15] (0, 75%) with *Caesalpinia benthamiana* (Baill.).

3.2 Phytochemical Screening

The results of the phytochemical screening carried out on the extracts of plants referred to malaria are mentioned in the Table 1. Chemical composition variations were found among the different extracts and between the two plant species analyzed.

3.3 *Diospyros monbuttensis*

The leaves contain sterols, polyterpenes, polyphenols, flavonoids, alkaloids, saponins, leucoanthocyanins, tannins, quinones and coumarins. Anthocyanins, cardiac glycosides and steroids are present at very small quantities. These results are different from those of Anie et al. [9], who reported the absence of tannins in roots and stem bark. They were similar to those of Bouquet and Debray [25] which showed the presence of quinones, tannins, sterols and saponins in the leaves and differ from ours by the absence of flavonoids and alkaloids. Previous studies [8] reported the presence of saponins, anthraquinones, cardiac glycosides and alkaloids. According Bouquet and Debray [25], this plant is considered to be a good remedy for stiffness, stomach ache, edema and leprosy by Baoule and Agni ethnies of Côte d'Ivoire. The antimalarial therapeutic indication may be related to the presence of alkaloids because we know that the majority of antimalarial drugs belong to the same chemical family [26]. Various studies have shown that coumarins have a high antioxidant capacity. This antioxidant activity is due to their capacity to scavenging free radicals and chelating metal ions [27]. Analgesics and anti-inflammatories activities are attributed to tannins. Tannins also contribute to wounds healing [28].

3.4 *Dialium dinklagei*

The decoction of the leaves contains sterols, polyterpenes, polyphenols, flavonoids, alkaloids,

coumarins and saponins in large quantities. Anthocyanins, leucoanthocyanins, tannins, quinones and cardiac glycosides are present in small quantities. There was also an absence of steroids. These results were comparable to those of Akoué [13], who noted a large presence of flavonoids, alkaloids, tannins and polyphenols in the ethanol extract of the leaves. Very little research has been made on this plant, but our results confirm those of Bouquet and Debray [25] about the presence of tannin in the leaves of *D. dinklagei*. Sterols, saponins, alkaloids and polyterpenes help explain its traditional therapeutic uses. In addition to their industrial applications as surfactants, traditionally saponins are widely used as detergents, pesticides and mollucides and foam, and can also have advantages on health [29].

3.5 Polyphenol Content of *Diospyros monbuttensis* and *Dialium dinklagei*

The contents of total phenols in the plant extracts, determined from a calibration curve of gallic acid ($Y=0.01810X + 0.07179$; $R^2=0.9911$) showed that *Dialium dinklagei* is richer in phenols than *Diospyros monbuttensis*, whatever the type of solvent used. The highest content was obtained in the methanolic extract of *Dialium dinklagei*. These levels ranged from 114.1 mgGAE/g and 205.1 mgGAE/g (Table 2). The difference between the contents of the phenol based on the plant and solvent used is highly significant ($p < 0.0001$).

Figure 1. Yields of extracts (%)

Table 1. Results of the foliar phytochemical tests

Natural substances		Chemical groups												
Plantspecies	Extracts	Steroids	Sterols and polyterpene s	Polyphenols	Flavonoids	Anthocyanins	Leucoanthocyanins	Gallic tannins	Catechiq tannins	Quinones	Alcaloids	Coumarins	Cardiotonic Glycosides	Saponins
<i>Diospyros monbuttensis</i>	Dec	+	+	+++	+	-	+	+++	-	+++	+	++	-	+++
	Hex	-	-	-	-	-	-	-	-	-	+++	+	-	-
	Met	-	+	+++	+	+	+++	+++	++	++	+	+	+	++
	Aq	-	+	+++	+	-	+	+	+	++	+	+	-	+++
<i>Dialium dinklagei</i>	Dec	-	+++	++	++	+	+	+	++	+	+++	++	+	+++
	Hex	-	++	-	-	-	-	-	+	-	+++	+	-	-
	Met	-	++	+	+++	++	+	+	-	-	+++	++	+	++
	Aq	-	-	+	+++	-	++	+	-	+	+++	+	-	+++

Strong positive: + + +; moderately positive: + +; Low positive: +; negative test:-
Dec: Decoction extract; Hex: Hexane extract; Met: Methanolic extract; Aq: Aqueous extract

Table 2. Total phenols, flavonoids and tannins contents of extracts

	Phenol content (mgGAE/g dry extract)		Flavonoid content (mgQE/g dry extract)		Tannin content (mgTAE/g dry extract)	
	Decoction extract	Methanolic extract	Decoction extract	Methanolic extract	Decoction extract	Methanolic extract
<i>Diospyros monbuttensis</i>	136,54±12,63	114,10±2,43	62,18±2,03	15,25±1,13	46,50±1,00	85,70±3,25
<i>Dialium dinklagei</i>	185,59±5,89	205,07±9,09	6,780±0,61	10,90±1,52	146,4±5,46	108,7±6,85

Values are expressed as mean ± SD (n = 3)

The results of the flavonoid contents of the different extracts of both plants species analyzed, determined from a calibration curve of quercetin ($Y = 0.02834X + 0.01368$; $R^2 = 0.9986$) showed that *Diospyros monbuttensis* is richer in flavonoids than *Dialium dinklagei*. The differences between the amounts of flavonoids according to the plant species and the solvent used were significant ($p < 0.0001$). These amounts vary between 6.78 mgQE/g and 62.18 mgQE/g (Table 2). The highest amount was obtained in the decoction extract of *Diospyros monbuttensis* (62.18 mgQE/g).

Tannins from different plant extracts were determined from a calibration curve of tannic acid ($Y = 0.2167X - 0.004417$; $R^2 = 0.9761$). Whatever the type of solvent used, the results showed that *Dialium dinklagei* is richer in tannins than *Diospyros monbuttensis*; the highest amount was obtained in the decoction extract of *Dialium dinklagei*. These contents ranged from 46.50 mgTAE/g and 146.4 mgTAE/g (Table 2). Statistically, the difference between the contents of the phenols concerning the plant species analyzed and the type of solvent used were highly significant ($p < 0.0001$).

The analysis of variance (ANOVA) and the post-ANOVA test (Tukey's multiple comparison test) have shown that the means of different contents of phenol, tannins and flavonoids of the two plant species were significantly different. When compared to those obtained by Akoue et al. [13] (4.42 ± 0.45 mgGAE/g for the aqueous extract and 2.27 ± 0.0038 mgGAE/g ethanolic extract), the contents of phenol *Dialium dinklagei* for methanolic and aqueous extracts were very high. This difference could be due to the extraction solvent and the method used.

In general, the decoction gave the highest contents of phenols, flavonoids and tannins. On the hand, the increase in temperature favors the diffusion and solubility of substances extracted. On the other hand, it destroys certain fragile substances [30]. This increase of the contents of flavonoids in the decocted extract of *Diospyros monbuttensis* can be explained by the destruction by the heat of polyphenol oxidases (PPO) that lowers the polyphenol content; thus the breaking of the connection between polyphenols and other substances leading to the accessibility of these active principles may explain this abundance [31]. The flavonoid content of *Dialium dinklagei*, relatively low in relation with that of *Diospyros monbuttensis*,

which is high in total phenol, could be explained by the presence of other phenolic compounds such as coumarins. The tannin content in *Dialium dinklagei* was higher than that of *Diospyros monbuttensis*. This difference may be related to the species of the studied plants, which were different. Tannins are usually defined as natural polyphenolic compounds with high molecular weight, capable of forming complexes with proteins, their effects on animals are various [32]. Thus, these compounds provide protection against cardiovascular diseases, immune/autoimmune diseases, brain disorders such as Parkinson's disease, and Alzheimer's disease [33]. Traditional medicinal use of these plants is well justified because it is now recognized that phenolic compounds have a significant therapeutic interest: fight against atherosclerosis, anticarcinogenic actions, a major source of natural antioxidants to fight against cellular aging [34,35]. Furthermore the phytochemical sorting revealed that in addition to phenols, flavonoids and tannins, these plants also contain polyterpenes, saponins and alkaloids which contribute to their various therapeutic activities.

4. CONCLUSION

The overall results of the phytochemical screening and polyphenolic content partly and rationally explain the interest of traditional healers for these plants as antimalarial plants. The therapeutic effects can be induced by various chemicals (alkaloid, flavonoids, polyphenols, polyterpenes, saponins, sterols and catechic tannins) which form the scientific basis of traditional therapeutic use of the studied plants. The *in vivo* and *in vitro* studies and the determination of the structure of active molecules will deepen the study of the antimalarial activity of these plants.

CONSENT

Not applicable.

ETHICAL APPROVAL

The study been performed in accordance with the ethical requirements in force in Ivory Coast.

ACKNOWLEDGEMENTS

We are indebted to Professor Aké-Assi Laurent (Laboratory of Botany, Unité de Formation et de Recherche-Biosciences, University of Felix

Houphouet-Boigny/Abidjan, Côte d'Ivoire) for botanical identification of the plants.

We also thank Dr. ETTIEN Yapo for the English translation of the manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Diaby B, Kroa E, Niare A, Coulibaly D, Traore Y, Giani S, et al. Connaissances, attitudes et pratiques des tradipraticiens de la sante de la ville d'Abidjan sur le paludisme. Mali Med. 2011;Tome XXVI n°3, 8-12.
- Sandhya B, Thomas S, Isabel W, Shenbagarathai R. Ethnomedicinal plants used by the valaiyan community of Piranmalai hills (Reserved forest), Tamil Nadu, India – A pilot study. Compl and Alter Med. 2006;3:101-114.
- Séguéna F, Soro K, Soro D, N'guessan K. Savoir-faire de populations locales des taxons du jardin Botanique de Bingerville, Côte d'Ivoire. J Appl Biosci. 2013; 68:5374-5393.
- Kerharo J, Bouquet A. Plante medecinale et toxique de Côte d'Ivoire et Haute-Volta, mission d'étude de la pharmacopée indigène en AOF. Ed Vigot frère, Paris. 1950;300.
- Pousset. Plantes médicinales Africaine. Utilisation pratique. Deuxième Tome, ACCT, Paris. 1989;156.
- Anthony JP, Fyfe I, Smith H. Plant active components—a resource for antiparasitic agents? Trends in Parasitol. 2005;21:462-468.
- Toyang NJ, Verpoorte R. A review of the medicinal potentials of plants of the genus Vernonia (Asteraceae). J of Ethnopharm. 2013;146:681–723.
- Kayode J, Omotoyinbo MA. Ethnobotanical utilization and conservation of chewing sticks plants species in Ekiti State, Nigeria. Res J. Botany. 2009;4:1-9.
- Anie CO, Matthew IA, Okeri HA. Antimicrobial activity of crude extracts of *Diospyros monbuttensis* (Fam: Ebenaceae) root and stem barks. Inter J of Biomedical Res. 2011;1:18-24.
- Olasehinde G, Olusola O, Obashola F, Singh R, Adesola A. *In-vitro* studies on the sensitivity pattern of Plasmodium falciparum to antimalarial drugs and local herbal extracts. Mal J. 2012;11(Suppl 1):P72.
- Odelola HA, Okorosobo VI. Preliminary investigation of *in vitro* Antimicrobial Activity of two Nigerian *Diospyros* sp. (Ebenaceae). Afri J Med Sci. 1988; 17(4):167-170.
- Mallavadhani UV, Panda AK, Rao YR. Pharmacology and chemotaxonomy of Diospyros. Phytochem. 1998;49:901-51.
- Akoué GN, Obame LC, Ondo JP, Brama I, Mbading-Mbading W, Otogo N'ngang ES, et al. Ethnotherapy study, phytochemical and antiradical activities of *Agelaea pentagyna* (Lam) Baill and *Dialium dinklagei* Harms. Medicinal plants from Gabon. Inter J of Adv Res. 2013;1:246-255.
- Lock JM. Legumes of Africa: a Check-List. Royal Botanic Gardens, Kew, Richmond, United Kingdom. 1989;619.
- Békro YA, Békro JAM, Boua BB, TRA BFH, Ehilé EE. Etude ethnobotanique et screening phytochimique de *Caesalpinia benthamiana* (Baill.) Herend. et Zarucchi (Caesalpiniaceae). Rev Sci Nat. 2007;4:217-225.
- Zirih GN, Kra AKM, Guede-Guina F. Évaluation de l'activité antifongique *Microglossa Pirifolia* (Lamarck) O. Kuntze (Asteraceae) « PYMI » sur la croissance *in vitro* de *Candida albicans*. Rev de Med et Ph Afr. 2003;17:11-18.
- Nemlin J, Brunel JE. Fascicule de Travaux Pratiques de Matière Médicale (3e année). Université Nationale de Côte-d'Ivoire. Faculté de Pharmacie. Département de Pharmacognosie. Laboratoire de Phytologie, 47 p ; 1995.
- Béourou S, Tuo K, Ouattara K, Touré AO, Meité S, Ako AAB, et al. Phytochemical screening of some medicinal plants used to treat malaria in Côte d'Ivoire (West Africa). Intern J of Chem and Pharm Sc. 2014;2:919-925.
- Bruneton J. Pharmacognosie, Phytochimie, Plantes médicinales. Paris, 4^{ème} Edition Lavoisier; 2009.
- Mcdonald S, Prenzler PD, Autolovich M, Robards K. Phenolic content and antioxidant activity of olive oil extracts. Food Chem. 2001;73:73-84.
- Li HB, Cheng KW, Wong CC, Fan KW, Chen F, Jiang Y. Evaluation of antioxidant capacity and total phenolic content of

- different fractions of selected microalgae. Food Chem. 2007;102:771-776.
22. Bahorun T, Gressier B, Trotin F, Brunete C, Dine T, Vasseur J, et al. Oxygen species scavenging activity of phenolic extracts from hawthorn fresh plant organs and pharmaceutical preparations. *Arzneimittelforschung*. 1996;46:1086-1089.
23. Chang C, Yang M, Wen H, Chern J. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J Food Drug Anal*. 2002;10:178-182.
24. Bainbridge Z, Tomlins K, Westby A. Analysis of condensed tannins using acidified vanillin. *J of F Sc and Agr*. 1996;29:77-79.
25. Bouquet A, Debray M. *Plantes médicinales de Côte-d'Ivoire, Imprimerie Louis Jean*, Paris (France). 1974;232.
26. Zirihi G N, N'guessan K, Etien DT, Seri KBP. Evaluation in vitro of antiparasitic activity of ethanolic extracts of *Funtumia elastica*, *Rauvolfia vomitoria* and *Zanthoxylum gillettii* on *Plasmodium falciparum* isolates from Côte d'Ivoire. *J of An & PI Sc*. 2009;5:406-413.
27. Tseng A. Chemoprevention of tumors in MTV-H ras transgenic mice with coumarins. *Proc Am Assoc Cancer Res*. 1991;32:2257.
28. Okwu DE, Josiah C. Evaluation of the chemical composition of two Nigerian medicinal plants. *Afr J Biotech*. 2006;5:357-361.
29. Shi J, Arunasalam K, Yeung DY, Kakuda Mittal G, Jiang Y. Saponins from edible legumes: Chemistry, processing and health benefits. *J Med Food*. 2004;7:67-78.
30. Jokić S, Velić D, Bilić M, Bucić-Kojić A, Planinić M, Tomas S. Modelling of the process of solid-liquid extraction of total polyphenols from soybeans. *J Food Sci*. 2010;28:206-212.
31. Lutz M, Henriques C, Escobar M. Chemical composition and antioxidant properties of mature and baby artichokes (*Cynara scolymus* L.), raw and cooked. *J of Food Compos Anal*. 2011;24:49-54.
32. Yang CMJ, Russell JB. Resistance of proline-containing peptides to ruminal degradation in vitro. *Appl Environ Microbiol*. 1992;58:3954-3958.
33. Han X, Shen T, Lou H. Dietary Polyphenols and their biological significance. *Inter J of Mol Sc*. 2007; 950-988.
34. Hagerman AE, Riedl KM, Jones GA, Sovik KN, Ritchard NT, Hartzfeld PW, Riechel TL. High molecular weight plant polyphenolics (tannins) as biological antioxidants. *J Agric Food Chem*. 1998;46:1887-1892.
35. Kumar GS, Nayaka H, Dharmesh SM; Salimath PV. Free and bound phenolic antioxidant in amla (*Emblia officinalis*) and turmeric (*Curcuma longa*). *J Food Compos Anal*. 2006;19:446-452.

© 2015 Tuo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?id=888&id=36&aid=7803>