



High order homogenization of the Poisson equation in a perforated periodic domain

Florian Feppon

► **To cite this version:**

Florian Feppon. High order homogenization of the Poisson equation in a perforated periodic domain. 2020. hal-02518528v4

HAL Id: hal-02518528

<https://hal.archives-ouvertes.fr/hal-02518528v4>

Preprint submitted on 24 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HIGH ORDER HOMOGENIZATION OF THE POISSON EQUATION IN A PERFORATED PERIODIC DOMAIN

F. FEPPON¹²

¹ *Centre de Mathématiques Appliquées, École Polytechnique, Palaiseau, France*
² *Safran Tech, Magny-les-Hameaux, France*

ABSTRACT. We derive high order homogenized models for the Poisson problem in a cubic domain periodically perforated with holes where Dirichlet boundary conditions are applied. These models have the potential to unify the three possible kinds of limit problems derived by the literature for various asymptotic regimes (namely the “unchanged” Poisson equation, the Poisson problem with a strange reaction term, and the zeroth order limit problem) of the ratio $\eta \equiv a_\varepsilon/\varepsilon$ between the size a_ε of the holes and the size ε of the periodic cell. The derivation relies on algebraic manipulations on formal two-scale power series in terms of ε and more particularly on the existence of a “criminal” ansatz, which allows to reconstruct the oscillating solution u_ε as a linear combination of the derivatives of its formal average u_ε^* weighted by suitable corrector tensors. The formal average is itself the solution of a formal, infinite order homogenized equation. Classically, truncating the infinite order homogenized equation yields in general an ill-posed model. Inspired by a variational method introduced in [52, 23], we derive, for any $K \in \mathbb{N}$, well-posed corrected homogenized equations of order $2K+2$ which yields approximations of the original solutions with an error of order $O(\varepsilon^{2K+4})$ in the L^2 norm. Finally, we find asymptotics of all homogenized tensors in the low volume fraction regime $\eta \rightarrow 0$ and in dimension $d \geq 3$. This allows us to show that our higher order effective equations converge coefficient-wise to either of the classical homogenized regimes of the literature which arise when η is respectively equivalent, or greater than the critical scaling $\eta_{\text{crit}} \sim \varepsilon^{2/(d-2)}$.

Keywords. Homogenization, higher order models, perforated Poisson problem, homogeneous Dirichlet boundary conditions, strange term.

AMS Subject classifications. 35B27, 76M50, 35330.

CONTENTS

| | |
|---|----|
| 1. Introduction | 2 |
| 2. Notation and summary of the derivation | 5 |
| 2.1. Notation conventions | 6 |
| 2.2. Summary of the derivation | 7 |
| 3. Derivation of the infinite order homogenized equation and of the criminal ansatz | 9 |
| 3.1. Traditional ansatz: definition of the cell tensors \mathcal{X}^k | 9 |
| 3.2. Properties of the tensors \mathcal{X}^k : odd order tensors \mathcal{X}^{2p+1*} are zero. | 10 |
| 3.3. Infinite order homogenized equation and criminal ansatz: tensors M^k and N^k | 11 |
| 3.4. Simplifications for the tensors \mathcal{X}^{k*} and M^k in case of symmetries | 15 |
| 4. Homogenized equations of order $2K+2$: tensors \mathbb{D}_K^{2k} | 17 |
| 4.1. Sufficient conditions under which an effective solution yields higher order approximations | 17 |
| 4.2. Construction of a well-posed higher order effective model by mean of a variational principle | 18 |
| 4.3. Asymptotic surprise: $\mathbb{D}_K^{2k} = M^{2k}$; error estimates for the homogenized model of order $2K+2$ | 22 |
| 5. Retrieving the classical regimes: low volume fraction limits when the size of the obstacles tends to zero | 24 |
| References | 30 |

Corresponding author. Email: florian.feppon@polytechnique.edu.

1. INTRODUCTION

One of the industrial perspectives offered today by the theory of homogenization lies in the development of more and more efficient topology optimization algorithms for the design of mechanical structures; there exists a variety of homogenization based techniques [6, 24, 17], including density based (or “SIMP”) methods [16]. Broadly speaking, the principle of these algorithms is to optimize one or several parameters of the microstructures which affect the coefficients of an effective model. The latter model accounts for the constitutive physics of the mixture of two materials (one representing the solid structure and one representing void) and is mathematically obtained by homogenization of the linear elasticity system. The knowledge of the dependence between the parameters of the microstructure and the coefficient of the effective model is the key ingredient of homogenization based techniques, because it allows to numerically—and automatically—interpret “gray designs” (i.e., for which the local density of solid is not a uniformly equal to 0 or 1) into complex composite shapes characterized by multi-scale patterns and geometrically modulated micro-structures [45, 10, 34, 36].

Several works have sought extensions of these methods for topology optimization of fluid systems, where the incompressible Navier–Stokes system is involved. In this context, an effective model is needed for describing the homogenized physics of a porous medium filled with either solid obstacles, fluid, or a mixture of both. However, classical literature [25, 3, 13, 49] identifies three possible kinds of homogenized models depending on how periodic obstacles of size a_ε scale within their periodic cell of size ε : depending on how the scaling $\eta \equiv a_\varepsilon/\varepsilon$ compares to the critical size $\sigma_\varepsilon := \varepsilon^{d/(d-2)}$ (in dimension $d \geq 3$), the fluid velocity converges as $\varepsilon \rightarrow 0$ to the solution of either a Darcy, a Brinkman or a Navier–Stokes equation. Unfortunately, there is currently no further result regarding an effective model that would be able to describe a medium featuring all possible sizes of locally periodic obstacles. The strategy that is the most commonly used in the density based topology optimization community consists in using either the Brinkman equation exclusively [20, 21, 28], or the Darcy model exclusively [54, 47]. These methodology have proved efficient in a number of works [46, 27], however, they remain inconsistent from a homogenization point of view, since these models are valid only for particular regimes of size of obstacles. In particular, this limitation makes impossible to interpret “gray” designs obtained with classical fluid topology optimization algorithms.

The main objective of this paper is to expose, accordingly, the derivation of a new class of—high order—homogenized models for perforated problems which have the potential to unify the different regimes of the literature. Our hope is that these models could permit, in future works, to develop new mathematically consistent and homogenization based topology optimization algorithms for fluid systems.




FIGURE 1. The perforated domain D_ε and the unit cell $Y = P \setminus (\eta T)$.

This article is a preliminary study towards such purpose: we propose to investigate here the case of the Poisson problem in a perforated periodic domain with Dirichlet boundary conditions on the holes,

$$\begin{cases} -\Delta u_\varepsilon = f & \text{in } D_\varepsilon \\ u_\varepsilon = 0 & \text{on } \partial\omega_\varepsilon \\ u_\varepsilon & \text{is } D\text{-periodic,} \end{cases} \quad (1.1)$$

which can be considered to be a simplified scalar and linear version of the full Navier–Stokes system. Let us mention that the analysis of the full Navier–Stokes system is much more challenging because of the

incompressibility constraint and the vectorial nature of the problem; the extension of the current work for the Stokes system—its linear counterpart—shall however be exposed in a future contribution [32, 33].

The setting considered is the classical context of periodic homogenization represented on Figure 1: $D := [0, L]^d$ is a d -dimensional box filled with periodic obstacles $\omega_\varepsilon := \varepsilon(\mathbb{Z}^d + \eta T) \cap D$. $P = (0, 1)^d$ is the unit cell, and $Y := P \setminus \eta T$ is the unit perforated cell. The parameter ε is the size of the periodic cell and is given by $\varepsilon := L/n$ where $n \in \mathbb{N}$ is an integer number assumed to be large. The parameter η is another rescaling of the obstacle T within the unit cell: the holes are therefore of size $a_\varepsilon := \eta\varepsilon$ which allow us to consider in section 5 so-called low volume fraction limits where η converges to zero. The boundary of the obstacle T is assumed to be smooth. $D_\varepsilon := D \setminus \omega_\varepsilon$ denotes the perforated domain and $f \in C^\infty(D)$ is a smooth D -periodic right-hand side. The periodicity assumption for u_ε and f is classical in homogenization and is used to avoid difficulties related to the arising of boundary layers (see [39, 19, 7]).

The literature accounts for several homogenized equations depending on how the size $a_\varepsilon = \eta\varepsilon$ of the holes compares to the critical size $\sigma_\varepsilon := \varepsilon^{d/(d-2)}$ in dimension $d \geq 3$ (or how $\log(a_\varepsilon)$ compares to $\log(\sigma_\varepsilon) := -1/\varepsilon^2$ in dimension $d = 2$) [43, 26, 3, 13, 49, 29, 37]:

- if $a_\varepsilon = o(\sigma_\varepsilon)$ (if $\log(a_\varepsilon) = o(\log(\sigma_\varepsilon))$ if $d = 2$), then the holes are “too small” and u_ε converges as $\varepsilon \rightarrow 0$ to the solution u of the Poisson equation in the homogeneous domain D (without holes):

$$\begin{cases} -\Delta u = f \text{ in } D \\ u \text{ is } D\text{-periodic.} \end{cases} \quad (1.2)$$

- if $a_\varepsilon = \sigma_\varepsilon$ (if $\log(a_\varepsilon) = -1/\varepsilon^2$ if $d = 2$), then u_ε converges as $\varepsilon \rightarrow 0$ to the solution u of the modified Poisson equation

$$\begin{cases} -\Delta u + Fu = f \text{ in } D \\ u \text{ is } D\text{-periodic,} \end{cases} \quad (1.3)$$

where the so-called *strange* reaction term Fu involves a positive constant $F > 0$ which can be computed by means of an exterior problem in $\mathbb{R}^d \setminus T$ when $d \geq 3$ (see (5.7) below), and which is equal to 2π if $d = 2$ (see [3, 25, 49, 37]).

- if $\sigma_\varepsilon = o(a_\varepsilon)$ (if $\log(\sigma_\varepsilon) = o(\log(a_\varepsilon))$ if $d = 2$) and $a_\varepsilon = \eta\varepsilon$ with $\eta \rightarrow 0$ as $\varepsilon \rightarrow 0$, then the holes are “large” and $a_\varepsilon^{d-2}\varepsilon^{-d}u_\varepsilon$ ($-\log(a_\varepsilon)\varepsilon^{-2}u_\varepsilon$ if $d = 2$) converges to the solution u of the zeroth order equation

$$\begin{cases} Fu = f \text{ in } D \\ u \text{ is } D\text{-periodic,} \end{cases} \quad (1.4)$$

where F is the same positive constant as in (1.3).

- if $a_\varepsilon = \eta\varepsilon$ with the ratio η fixed, then $\varepsilon^{-2}u_\varepsilon$ converges to the solution u of the zeroth order equation

$$\begin{cases} M^0 u = f \text{ in } D \\ u \text{ is } D\text{-periodic,} \end{cases} \quad (1.5)$$

where M^0 is another positive constant (which depends on η). Furthermore it can be shown that $M^0/|\log(\eta)| \rightarrow F$ if $d = 2$, and $M^0/\eta^{d-2} \rightarrow F$ (if $d \geq 3$) when $\eta \rightarrow 0$, so that there is a continuous transition from (1.5) to (1.4); see [4] and corollary 6 below.

The different regimes (1.2) to (1.5) occur because the heterogeneity of the problem comes from the zero Dirichlet boundary condition on the holes ω_ε in (1.1): this is the major difference with the setting commonly assumed in linear elasticity, where the heterogeneity induced by the mixture of two materials is instead inscribed in the coefficients of the physical state equation [53, 6]. (1.2) and (1.3) are the respectively the analogous of the Navier–Stokes and Brinkman regimes in the context of the homogenization of the Navier–Stokes equation, while the zeroth order equations (1.4) and (1.5) are analogous to Darcy models. As stressed above, the existence of these regimes raises practical difficulties in view of applying the homogenization method for shape optimization: the previous considerations show that one should use (1.2) and (1.3) in regions featuring none or very tiny obstacles, however one should use the zeroth order model (1.5) when the obstacles become large enough.

Our goal is to propose an enlarged vision of the homogenization of (1.1) through the construction, for any $K \in \mathbb{N}$, of a homogenized equation of order $2K + 2$,

$$\begin{cases} \sum_{k=0}^{K+1} \varepsilon^{2k-2} \mathbb{D}_K^{2k} \cdot \nabla^{2k} v_{\varepsilon, K}^* = f \text{ in } D, \\ v_{\varepsilon, K}^* \text{ is } D\text{-periodic,} \end{cases} \quad (1.6)$$

which yields an approximation of u_ε of order $O(\varepsilon^{2K+4})$ in the $L^2(D_\varepsilon)$ norm (for a fixed given scaling of the obstacles η). The function $v_{\varepsilon, K}^*$ denotes the higher order homogenized approximation of u_ε and $\mathbb{D}_K^{2k} \cdot \nabla^{2k}$ is a differential operator of order $2k$ with constant coefficients (the notation is defined in (2.3) below). Equation (1.6) is a ‘‘corrected’’ version of the zeroth order model (1.5) (for any K , it holds $\mathbb{D}_K^0 = M^0$), which yields a more accurate solution when ε is ‘‘not so small’’.

Our mathematical methodology is inspired from the works of Bakhvalov and Panasenko [15], Smyshlyaev and Cherednichenko [52], and Allaire et. al. [11]; it starts with the identification of a ‘‘classical’’ two-scale ansatz

$$u_\varepsilon(x) = \sum_{i=0}^{+\infty} \varepsilon^{i+2} u_i(x, x/\varepsilon), \quad x \in D_\varepsilon, \quad (1.7)$$

expressed in terms of Y -periodic functions $u_i : D \times Y \rightarrow \mathbb{R}$ which do not depend on ε . Our procedure involves then formal operations on related power series which give rise to several families of tensors and homogenized equations for approximating the formal, infinite order homogenized average u_ε^* :

$$u_\varepsilon^*(x) := \sum_{i=0}^{+\infty} \varepsilon^{i+2} \int_Y u_i(x, y) dy, \quad x \in D. \quad (1.8)$$

In proposition 5 below, we obtain that u_ε^* in (1.11) is the solution of a formal, ‘‘infinite order’’ homogenized equation,

$$\sum_{k=0}^{+\infty} \varepsilon^{2k-2} M^{2k} \cdot \nabla^{2k} u_\varepsilon^* = f, \quad (1.9)$$

where M^0 is the positive constant of (1.5) and $(M^k)_{k \geq 1}$ is a family of (constant) tensors of order k . From a computational point of view, one needs a well-posed finite order model. As it can be expected from other physical contexts [8, 11], the effective model obtained from a naive truncation of (1.9), say at order $2K$,

$$\sum_{k=0}^K \varepsilon^{2k-2} M^{2k} \cdot \nabla^{2k} v_{\varepsilon, K}^* = f \quad (1.10)$$

is in general not well-posed [8, 1]. Several techniques have been proposed in the literature to obtain well-posed homogenized models of finite order in the context of the conductivity or of the wave equation [12, 9, 1, 2, 11]. The derivation of the *well-posed* homogenized equation (1.6) relies on a minimization principle inspired from Smyshlyaev and Cherednichenko [52] and is marked by two surprising facts.

The first surprising result is the existence of a somewhat remarkable identity which expresses the oscillating solution u_ε in terms of its non-oscillating average u_ε^* :

$$u_\varepsilon(x) = \sum_{k=0}^{+\infty} \varepsilon^k N^k(x/\varepsilon) \cdot \nabla^k u_\varepsilon^*(x). \quad (1.11)$$

The functions N^k are P -periodic corrector tensors of order k (definition 2) depending only on the shape of the obstacles ηT and that vanish on $\partial(\eta T)$. Furthermore, N^0 is of average $\int_Y N^0(y) dy = 1$ and N^k is of average $\int_Y N^k(y) dy = 0$ for $k \geq 1$ (proposition 8): u_ε^* is consistently the average of u_ε with respect to the fast variable x/ε . While the derivation of (1.7) is very standard in periodic homogenization [50, 40, 18, 14], the existence of such a relation (1.11) (when compared to (1.7)) between the oscillating solution is less obvious; it has been noticed for the first time by Bakhvalov and Panasenko [15] for the conductivity equation, and then in further homogenization contexts in [51, 52, 23, 11, 2, 1]. Following the denomination of [11], we call the ansatz (1.11) ‘‘criminal’’ because the function u_ε^* has the structure of a formal power series in ε .

The second surprise lies in that our higher order homogenized equation (1.6) is obtained by adding to (1.10) a *single* term $\varepsilon^{2K} \mathbb{D}_K^{2K+2} \cdot \nabla^{2K+2}$; in other words $\mathbb{D}_K^{2k} = M^{2k}$ for any $0 \leq 2k \leq 2K$. This fact, which does not seem to have been noticed in previous works, is quite surprising because following [52, 23], the derivation of (1.6) is based on a minimization principle for the truncation of (1.11) at order K ,

$$W_{\varepsilon,K}(v_{\varepsilon,K}^*)(x) := \sum_{k=0}^K \varepsilon^k N^k(x/\varepsilon) \cdot \nabla^k v_{\varepsilon,K}^*(x), \quad x \in D_\varepsilon \quad (1.12)$$

which is expected to yield an approximation of order $O(\varepsilon^{K+3})$ only in the $L^2(D_\varepsilon)$ norm (u_ε^* and $v_{\varepsilon,K}^*$ are of order $O(\varepsilon^2)$). This is the order of accuracy stated in our previous work [32] and similarly obtained in the conductivity case by [52], or for the Maxwell equations by [23]; it is related to the observation that the first half of the coefficients of (1.6) and (1.9) coincide: $\mathbb{D}_K^{2k} = M^{2k}$ for any $0 \leq 2k \leq K$ (proposition 13). In fact, it turns out that *all coefficients* \mathbb{D}_K^{2k} and M^{2k} coincide except the one of the leading order; $\mathbb{D}_K^{2K+2} \neq M^{2K+2}$. As a result, we are able to show in the present paper that the reconstructed function obtained by adding more correctors,

$$W_{\varepsilon,2K+1}(v_{\varepsilon,K}^*)(x) := \sum_{k=0}^{2K+1} \varepsilon^k N^k(x/\varepsilon) \cdot \nabla^k v_{\varepsilon,K}^*(x), \quad x \in D_\varepsilon, \quad (1.13)$$

yields an approximation of u_ε of order $O(\varepsilon^{2K+4})$ (corollary 5 below):

$$\|u_\varepsilon - W_{\varepsilon,2K+1}(v_{\varepsilon,K}^*)\|_{L^2(D_\varepsilon)} + \varepsilon \|\nabla(u_\varepsilon - W_{\varepsilon,2K+1}(v_{\varepsilon,K}^*))\|_{L^2(D_\varepsilon)} \leq C_K(f) \varepsilon^{2K+4}. \quad (1.14)$$

Finally, we obtain in corollaries 6 and 7 (see also remark 8) that our homogenized models have the potential to “unify” the different regimes of the literature, in the sense that (1.6) and (1.9) converge formally (coefficient-wise) to either of the effective equations (1.3) and (1.4) when the scaling of the obstacle $\eta \rightarrow 0$ vanishes at rates respectively equivalent or greater than the critical size $\eta_{\text{crit}} \sim \eta^{2/(d-2)}$.

Unfortunately, we do not obtain in this work that this convergence holds for all possible rates, because the estimates of corollary 6 imply that higher order coefficients $\varepsilon^{2k-2} M^{2k}$ with $k > 2$ could blow up if the rate η vanishes faster than the critical size (i.e. when $\eta = o(\varepsilon^{2/(d-2)})$). However the coefficient-wise convergence holds for the homogenized equation (1.6) of order 2 (with $K = 0$). Although (i) the derivation of (1.6) has been performed by assuming η constant and (ii) all our error bounds feature constants $C_K(f)$ which depend *a priori* on η , these results seem to indicate that (1.6) has the potential to yield valid homogenized approximations of (1.1) in any regime of size of holes if $K = 0$ (which was our initial goal), and for any size $\eta \geq \eta_{\text{crit}}$ if $K \geq 1$.

The exposure of our work outlines as follows. Section 2 introduces the notation conventions and provides a brief summary of our derivations.

Section 3 then details the procedure which allows to construct the family of tensors M^k and $N^k(y)$ arising in the formal infinite order homogenized equation (1.9) and in the criminal ansatz (1.11). Additionally, we establish a number of algebraic properties satisfied by these tensors and we provide an account of the simplifications which occur in case of symmetries of the obstacle with respect to the unit cell axes.

Section 4 is devoted to the construction of the finite order homogenized equation (1.6) thanks to the method of Smyshlyaev and Cherednichenko. We prove the ellipticity of the model and we establish that $\mathbb{D}_K^{2k} = M^{2k}$ for any $0 \leq 2k \leq 2K$ (and not only for the first half coefficients with $0 \leq 2k \leq K$ as observed in [32, 52]). The high order homogenization process is then properly justified by establishing the error estimate (1.14).

Finally, section 5 examines the asymptotic properties of the tensors M^k in the low-volume fraction limit $\eta \rightarrow 0$ (in space dimension $d \geq 3$). This allows us to retrieve formally the classical regimes and the arising of the celebrated “strange term” (see [25]) at the critical scaling $\eta \sim \varepsilon^{2/(d-2)}$.

2. NOTATION AND SUMMARY OF THE DERIVATION

The full derivation of higher order homogenized equations involves the construction of a number of families of tensors such as $\mathcal{X}^k, M^k, N^k, \mathbb{D}_K^{2k}$. For the convenience of the reader, the notation conventions related to two-scale functions and tensor operations are summarized in section 2.1. We then provide a short synthesis of our main results and of the key steps of our derivations in section 2.2.

2.1. Notation conventions

Below and further on, we consider scalar functions such as

$$\begin{aligned} u : D \times P &\rightarrow \mathbb{R} \\ (x, y) &\mapsto u(x, y) \end{aligned} \tag{2.1}$$

which are both D and P -periodic with respect to respectively the first and the second variable, and which vanish on the hole $D \times (\eta T)$. The arguments x and y of $u(x, y)$ are respectively called the “slow” and the “fast” or “oscillating” variable. With a small abuse of notation, the partial derivative with respect to the variable y_j (respectively x_j) is simply written ∂_j instead of ∂_{y_i} (respectively ∂_{x_j}) where the context is clear, *i.e.* when the function to which it is applied depends only on y (respectively only on x).

The star-“*”-symbol is used to indicate that a quantity is “macroscopic” in the sense it does not depend on the fast variable x/ε ; e.g. $v_{\varepsilon, K}^*$ in (1.6), u_ε^* in (1.8) or J_K^* in (2.15) below. In the particular case where a two-variable quantity $u(x, y)$ is given such as (2.1), $u^*(x)$ always denotes the average of $y \mapsto u(x, y)$ with respect to the y variable:

$$u^*(x) := \int_P u(x, y) dy = \int_Y u(x, y) dy, \quad x \in D,$$

where the last equality is a consequence of u vanishing on $P \setminus Y = \overline{\eta T}$.

When a function $\mathcal{X} : P \rightarrow \mathbb{R}$ depends only on the y variable, we find sometimes more convenient (especially in section 5) to write its cell average with the usual angle bracket symbols:

$$\langle \mathcal{X} \rangle := \int_P \mathcal{X}(y) dy.$$

In all what follows and unless otherwise specified, the Einstein summation convention over repeated *subscript* indices is assumed (but never on *superscript* indices). Vectors $\mathbf{b} \in \mathbb{R}^d$ are written in bold face notation.

The notation conventions including those used related to tensor are summarized in the nomenclature below.

| | |
|---------------------------|--|
| \mathbf{b} | Vector of \mathbb{R}^d |
| $(b_j)_{1 \leq j \leq d}$ | Coordinates of the vector \mathbf{b} . |
| b^k | Tensor of order k ($b_{i_1 \dots i_k}^k \in \mathbb{R}$ for $1 \leq i_1, \dots, i_k \leq d$) |
| $b^p \otimes c^{k-p}$ | Tensor product of tensors of order p and $k-p$: |

$$(b^p \otimes c^{k-p})_{i_1 \dots i_k} := b_{i_1 \dots i_p}^p c_{i_{p+1} \dots i_k}^{k-p}. \tag{2.2}$$

| | |
|----------------------|---|
| $b^k \cdot \nabla^k$ | Differential operator of order k associated with a tensor b^k : |
|----------------------|---|

$$b^k \cdot \nabla^k := b_{i_1 \dots i_k}^k \partial_{i_1 \dots i_k}^k, \tag{2.3}$$

with implicit summation over the repeated indices $i_1 \dots i_k$.

| | |
|------------------------------------|---|
| $(\mathbf{e}_j)_{1 \leq j \leq d}$ | Vectors of the canonical basis of \mathbb{R}^d . |
| e_j | Tensor of order 1 whose entries are $(\delta_{i_1 j})_{1 \leq i_1 \leq d}$ (for any $1 \leq j \leq d$). Note: e_j and \mathbf{e}_j are the same mathematical object when identifying tensors of order 1 to vectors in \mathbb{R}^d . |
| δ_{ij} | Kronecker symbol: $\delta_{ij} = 1$ if $i = j$ and $\delta_{ij} = 0$ if $i \neq j$. |
| I | Identity tensor of order 2: |

$$I_{i_1 i_2} = \delta_{i_1 i_2}.$$

Note that the identity tensor is another notation for the Kronecker tensor and it holds $I = e_j \otimes e_j$.

| | |
|----------|----------------------------------|
| J^{2k} | Tensor of order $2k$ defined by: |
|----------|----------------------------------|

$$J^{2k} := \overbrace{I \otimes I \otimes \dots \otimes I}^{k \text{ times}}. \tag{2.4}$$

$\mathbb{B}_K^{k,l}$ Tensor of order $l + m$ associated with the quadratic form

$$\mathbb{B}_K^{l,m} \nabla^l v \nabla^m w := \mathbb{B}_{l,i_1 \dots i_k, j_1 \dots j_l}^{l,m} \partial_{i_1 \dots i_k}^l v \partial_{j_1 \dots j_l}^m w, \quad (2.5)$$

for any smooth scalar fields $v, w \in \mathcal{C}^\infty(D)$.

With a small abuse of notation, we consider zeroth order tensors b^0 to be constants (i.e. $b^0 \in \mathbb{R}$) and we still denote by $b^0 \otimes c^k := b^0 c^k$ the tensor product with a k -th order tensor c^k .

In all what follows, a k -th order tensor b^k truly makes sense when contracted with k partial derivatives, as in (2.3). Therefore all the tensors considered throughout this work are identified to their symmetrization:

$$b_{i_1 \dots i_k}^k \equiv \frac{1}{k!} \sum_{\sigma \in \mathfrak{S}_k} b_{i_{\sigma(1)} \dots i_{\sigma(k)}},$$

where \mathfrak{S}_k is the permutation group of order k . Consequently, the order in which the indices i_1, \dots, i_k are written in $b_{i_1 \dots i_k}^k$ does not matter and the tensor product \otimes is commutative under this identification:

$$b^k \otimes c^{k-p} = c^{k-p} \otimes b^k. \quad (2.6)$$

Finally, C, C_K or $C_K(f)$ denote universal constants that do not depend on ε but whose values may change from line to line (and which depend *a priori* on the shape of the hole ηT).

2.2. Summary of the derivation

One of the main results of this paper is the derivation of the higher order homogenized equation (1.6) for any desired order $K \in \mathbb{N}$, and the justification of the procedure by establishing the error estimate (1.14). Before summarizing the most essential steps of our analysis, let us recall that here and in all what follows, equalities involving infinite power series such as (1.11) are formal and without a precise meaning of convergence. Our derivation outlines as follows:

- (1) following classical literature [41, 40, 15, 19], we introduce a family of k -th order tensors $(\mathcal{X}^k(y))_{k \in \mathbb{N}}$ obtained as the solutions of cell problems (see definition 1 and proposition 1) which allows to identify the functions $u_i(x, y)$ arising in (1.11) and to rewrite the traditional ansatz more explicitly:

$$u_\varepsilon(x) = \sum_{i=0}^{+\infty} \varepsilon^{i+2} \mathcal{X}^i(x/\varepsilon) \cdot \nabla^i f(x), \quad x \in D_\varepsilon. \quad (2.7)$$

Introducing the averaged tensors of order i , $\mathcal{X}^{i*} := \int_Y \mathcal{X}^i(y) dy$, the formal average (1.8) reads

$$u_\varepsilon^*(x) = \sum_{i=0}^{+\infty} \varepsilon^{i+2} \mathcal{X}^{i*} \cdot \nabla^i f(x). \quad (2.8)$$

- (2) We construct (in proposition 5) constant tensors M^i by inversion of the formal equality

$$\left(\sum_{i=0}^{+\infty} \varepsilon^{i-2} M^i \cdot \nabla^i \right) \left(\sum_{i=0}^{+\infty} \varepsilon^{i+2} \mathcal{X}^{i*} \cdot \nabla^i \right) = I,$$

which yields, after left multiplication of (2.8) by $\sum_{i=0}^{+\infty} \varepsilon^{i-2} M^i \cdot \nabla^i$, the “infinite order homogenized equation” for $u_\varepsilon^*(x)$:

$$\sum_{i=0}^{+\infty} \varepsilon^{i-2} M^i \cdot \nabla^i u_\varepsilon^*(x) = f(x). \quad (2.9)$$

Note that (2.9) is exactly (1.9) because all tensors \mathcal{X}^{2k+1*} and M^{2k+1} of odd order vanish (proposition 3 and corollary 1).

- (3) We substitute the expression of $f(x)$ given by (2.9) into the ansatz (2.7) so as to recognize a formal series product:

$$u_\varepsilon(x) = \left(\sum_{i=0}^{+\infty} \varepsilon^i \mathcal{X}^i(x/\varepsilon) \cdot \nabla^i \right) \left(\sum_{i=0}^{+\infty} \varepsilon^i M^i \cdot \nabla^i \right) u_\varepsilon^*(x). \quad (2.10)$$

Introducing a new family of tensors $N^k(y)$ defined by the corresponding Cauchy product,

$$N^k(y) := \sum_{p=0}^k \mathcal{X}^p(y) \otimes M^{k-p}, \quad y \in Y,$$

we obtain the ‘‘criminal’’ ansatz (1.11):

$$u_\varepsilon(x) = \sum_{k=0}^{+\infty} \varepsilon^k N^k(x/\varepsilon) \cdot \nabla^k u_\varepsilon^*(x). \quad (2.11)$$

- (4) We now seek to construct well-posed effective models of finite order. Inspired by [52, 23], we consider truncated versions of functions of the form of (2.11): for any $v \in H^{K+1}(D)$, we define

$$W_{\varepsilon,K}(v)(x) := \sum_{k=0}^K \varepsilon^k N^k(x/\varepsilon) \cdot \nabla^k v(x), \quad x \in D. \quad (2.12)$$

where the tensors N^k are extended by zero in $P \setminus Y$ for this expression to make sense in $D \setminus D_\varepsilon$. Remembering that u_ε (identified with its extension by 0 in $D \setminus D_\varepsilon$) is the solution to the energy minimization problem

$$\begin{aligned} \min_{u \in H^1(D)} \quad & J(u, f) := \int_D \left(\frac{1}{2} |\nabla u|^2 - fu \right) dx \\ \text{s.t.} \quad & \begin{cases} u = 0 \text{ in } \omega_\varepsilon \\ u \text{ is } D\text{-periodic,} \end{cases} \end{aligned} \quad (2.13)$$

we formulate an analogous minimization problem for $v \in H^{K+1}(D)$ by restriction of (2.13) to the smaller space of functions $W_{\varepsilon,K}(v) \in H^1(D)$:

$$\begin{aligned} \min_{v \in H^{K+1}(D)} \quad & J(W_{\varepsilon,K}(v), f) = \int_D \left(\frac{1}{2} |\nabla W_{\varepsilon,K}(v)|^2 - f(x)W_{\varepsilon,K}(v)(x) \right) dx \\ \text{s.t.} \quad & v \text{ is } D\text{-periodic.} \end{aligned} \quad (2.14)$$

Averaging over the fast variable x/ε (by using lemma 3), we obtain a new minimization problem involving an approximate energy J_K^* (definition 4) which does not depend on x/ε ,

$$\begin{aligned} \min_{v \in H^{K+1}(D)} \quad & J_K^*(v, f, \varepsilon) \\ \text{s.t.} \quad & v \text{ is } D\text{-periodic.} \end{aligned} \quad (2.15)$$

Its Euler–Lagrange equation (see definition 5) defines finally our well-posed homogenized equation (1.6) and in particular the family of tensors $(\mathbb{D}_K^{2k})_{0 \leq 2k \leq 2K+22}$.

- (5) In view of (2.12), this procedure is expected to yield by construction an approximation $W_{\varepsilon,K}(v_{\varepsilon,K}^*)$ of u_ε with an error $O(\varepsilon^{K+3})$ (because $v_{\varepsilon,K}^*$ is of order $O(\varepsilon^2)$, see lemma 7). Surprisingly, we verify that all the tensors \mathbb{D}_K^{2k} and the tensors M^{2k} coincide for $0 \leq 2k \leq 2K$ (proposition 13). This allows to obtain in corollary 5 the error estimate (1.14), which states that $v_{\varepsilon,K}^*$ and the reconstruction (1.13) yield a *much better* approximation than expected, namely of order $O(\varepsilon^{2K+4})$ instead of $O(\varepsilon^{K+3})$.

The most essential point of our methodology is the derivation of the non-classical ansatz (2.11) of step (3). Let us stress that in the available works of the literature concerned with high order homogenization of scalar conductivity equations and its variants [15, 52, 11], the criminal ansatz (analogous to (2.11)) is readily obtained from the classical one (analogous to (1.11)) because the tensors N^k and \mathcal{X}^k coincide in these contexts (check for instance [15, 8]). Our case is very different because the heterogeneity comes from the Dirichlet boundary condition on the holes ω_ε .

3. DERIVATION OF THE INFINITE ORDER HOMOGENIZED EQUATION AND OF THE CRIMINAL ANSATZ

This section now presents the steps (1) to (3) of [section 2.2](#) in detail. We start in [section 3.1](#) by reviewing the definition of the family of cell tensors $(\mathcal{X}^k(y))_{k \in \mathbb{N}}$ which allows to identify the functions $u_i(x, y)$ involved in the “usual” two-scale ansatz [\(1.8\)](#) and to obtain an error estimate for its truncation in [proposition 2](#). This part is not new; it is a review of classical results available in a number of works, see e.g. [\[41, 40, 19\]](#). We then establish in [section 3.2](#) several properties of the tensor \mathcal{X}^k which are less found in the literature; the most important being that the averages of all odd order tensors vanish: $\mathcal{X}^{2p+1*} = 0$ for any $p \in \mathbb{N}$.

The next [section 3.3](#) focuses on the definition and on the properties of the family of tensors M^k and $N^k(y)$ which allow to we infer the infinite order homogenized equation [\(1.9\)](#) and the criminal ansatz [\(1.11\)](#).

Finally, we investigate in [section 3.4](#) how symmetries of the obstacle ηT with respect to the axes of the unit cell P reflect into a decrease of the number of independent components of the homogenized tensors \mathcal{X}^{k*} and M^k .

3.1. Traditional ansatz: definition of the cell tensors \mathcal{X}^k

Classically [\[50, 41, 40\]](#), the first step of our analysis is to insert formally the two-scale expansion [\(1.11\)](#) into the Poisson system [\(1.1\)](#). Because it will help highlight the occurrence of Cauchy products, we also assume (for the purpose of the derivation only) that the right-hand side $f \in \mathcal{C}^\infty(D)$ depends on ε and admits the following formal expansion:

$$f(x) = \sum_{i=0}^{+\infty} \varepsilon^i f_i(x), \quad x \in D.$$

Evaluating the Laplace operator against [\(1.11\)](#), we obtain formally

$$-\Delta u_\varepsilon = \sum_{i=-2}^{+\infty} \varepsilon^{i+2} (-\Delta_{yy} u_{i+2} - \Delta_{xy} u_{i+1} - \Delta_{xx} u_i),$$

where we use the convention $u_{-2}(x, y) = u_{-1}(x, y) = 0$, and where $-\Delta_{yy}$, $-\Delta_{xy}$, $-\Delta_{xx}$ are the operators

$$-\Delta_{xx} := -\operatorname{div}_x(\nabla_x \cdot), \quad -\Delta_{xy} := -\operatorname{div}_x(\nabla_y \cdot) - \operatorname{div}_y(\nabla_x \cdot), \quad -\Delta_{yy} := -\operatorname{div}_y(\nabla_y \cdot).$$

Identifying all powers in ε yields then the traditional cascade of equations (obtained e.g. in [\[40\]](#)):

$$\begin{cases} -\Delta_{yy} u_{i+2} = f_{i+2} + \Delta_{xy} u_{i+1} + \Delta_{xx} u_i & \text{for all } i \geq -2 \\ u_{-2}(x, y) = u_{-1}(x, y) = 0. \end{cases} \quad (3.1)$$

This system of equations is solved by introducing an appropriate family of cell tensors [\[41, 40\]](#).

Definition 1. We define a family of tensors $(\mathcal{X}^k(y))_{k \in \mathbb{N}}$ of order k by recurrence as follows:

$$\begin{cases} -\Delta_{yy} \mathcal{X}^0 = 1 \text{ in } Y \\ -\Delta_{yy} \mathcal{X}^1 = 2\partial_j \mathcal{X}^0 \otimes e_j \text{ in } Y \\ -\Delta_{yy} \mathcal{X}^{k+2} = 2\partial_j \mathcal{X}^{k+1} \otimes e_j + \mathcal{X}^k \otimes I \text{ in } Y, & \text{for all } k \geq 0 \\ \mathcal{X}^k = 0 \text{ on } \partial(\eta T) \\ \mathcal{X}^k \text{ is } P\text{-periodic.} \end{cases} \quad (3.2)$$

The tensors \mathcal{X}^k are extended by 0 inside the hole ηT in the whole unit cell P , namely $\mathcal{X}^k(y) = 0$ for $y \in \eta T$.

Remark 1. In view of [\(2.2\)](#), the third line of [\(3.2\)](#) is a short-hand notation for

$$-\Delta \mathcal{X}_{i_1 \dots i_{k+2}}^{k+2} = 2\partial_{i_{k+2}} \mathcal{X}_{i_1 \dots i_{k+1}}^{k+1} + \mathcal{X}_{i_1 \dots i_k}^k \delta_{i_{k+1} i_{k+2}}, \quad \text{for all } k \geq 0.$$

Proposition 1. The solutions $(u_i(x, y))_{i \geq 0}$ to the cascade of equations [\(3.1\)](#) are given by

$$\forall i \geq 0, u_i(x, y) = \sum_{k=0}^i \mathcal{X}^k(y) \cdot \nabla^k f_{i-k}(x), \quad x \in D, y \in Y. \quad (3.3)$$

Recognizing a Cauchy product, the ansatz (1.11) can be formally written as the following infinite power series product:

$$u_\varepsilon(x) = \sum_{i=0}^{+\infty} \varepsilon^{i+2} \mathcal{X}^i(x/\varepsilon) \cdot \nabla^i f(x) = \varepsilon^2 \left(\sum_{i=0}^{+\infty} \varepsilon^i \mathcal{X}^i(x/\varepsilon) \cdot \nabla^i \right) \left(\sum_{i=0}^{+\infty} \varepsilon^i f_i(x) \right). \quad (3.4)$$

Proof. See [41, 40, 32]. \square

We complete our review by stating a classical error estimate result which justifies in some sense the formal power series expansion (3.4).

Proposition 2. Denote $u_{\varepsilon,K}$ the truncated ansatz of (3.4) at order $K \in \mathbb{N}$:

$$u_{\varepsilon,K}(x) := \sum_{i=0}^K \varepsilon^{i+2} \mathcal{X}^i(x/\varepsilon) \cdot \nabla^i f(x), \quad x \in D_\varepsilon.$$

Then assuming $f \in C^\infty(D)$ is D -periodic, the following error bound holds:

$$\|u_\varepsilon - u_{\varepsilon,K}\|_{L^2(D_\varepsilon)} + \varepsilon \|\nabla(u_\varepsilon - u_{\varepsilon,K})\|_{L^2(D_\varepsilon)} \leq C_K \varepsilon^{K+3} \|f\|_{H^{K+2}(D)}$$

for a constant C_K independent of f and ε (but depending on K).

Proof. See [41, 40, 32]. \square

3.2. Properties of the tensors \mathcal{X}^k : odd order tensors \mathcal{X}^{2p+1*} are zero.

Following our conventions of section 2.1, the average of the functions u_i and \mathcal{X}^i with respect to the y variable are respectively denoted:

$$u_i^*(x) := \int_Y u_i(x, y) dy, \quad x \in D, \quad (3.5)$$

$$\mathcal{X}^{i*} := \int_Y \mathcal{X}^i(y) dy. \quad (3.6)$$

In the next proposition, we show that $\mathcal{X}^{2p+1*} = 0$ are of zero average for any $p \in \mathbb{N}$ and that \mathcal{X}^{2p*} depends only on the lower order tensors \mathcal{X}^p and \mathcal{X}^{p-1} . Similar formulas have been obtained for the wave equation in heterogeneous media, see e.g. Theorem 3.5 in [2], and also [1, 48].

Proposition 3. For any $0 \leq p \leq k$, the following identity holds for the tensor \mathcal{X}^{k*} :

$$\mathcal{X}^{k*} = \int_Y \mathcal{X}^k dy = (-1)^p \int_Y (\mathcal{X}^{k-p} \otimes (-\Delta_{yy} \mathcal{X}^p) - \mathcal{X}^{k-p-1} \otimes \mathcal{X}^{p-1} \otimes I) dy, \quad (3.7)$$

with the convention that $\mathcal{X}^{-1} = 0$. In particular, for any $p \in \mathbb{N}$:

- $\mathcal{X}^{2p+1*} = 0$
- \mathcal{X}^{2p*} depends only on the tensors \mathcal{X}^p and \mathcal{X}^{p-1} :

$$\mathcal{X}^{2p*} = (-1)^p \int_Y (\partial_j \mathcal{X}^p \otimes \partial_j \mathcal{X}^p - \mathcal{X}^{p-1} \otimes \mathcal{X}^{p-1} \otimes I) dy. \quad (3.8)$$

Proof. The result is proved by induction. Formula (3.7) holds true for $p = 0$ by using the convention $\mathcal{X}^{-1} = 0$ and $-\Delta_{yy} \mathcal{X}^0 = 1$. Assuming now the result to be true for $0 \leq p < k$, we perform the following integration by parts where we use the boundary conditions satisfied by the tensors \mathcal{X}^k and the commutativity property (2.6) of the tensor product:

$$\begin{aligned} \mathcal{X}^{k*} &= (-1)^p \int_Y (-\Delta_{yy} \mathcal{X}^{k-p} \otimes \mathcal{X}^p - \mathcal{X}^{k-p-1} \otimes \mathcal{X}^{p-1} \otimes I) dy \\ &= (-1)^p \int_Y ((2\partial_j \mathcal{X}^{k-p-1} \otimes e_j + \mathcal{X}^{k-p-2} \otimes I) \otimes \mathcal{X}^p - \mathcal{X}^{k-p-1} \otimes \mathcal{X}^{p-1} \otimes I) dy \\ &= (-1)^p \int_Y (-2\partial_j \mathcal{X}^p \otimes e_j - \mathcal{X}^{p-1} \otimes I) \otimes \mathcal{X}^{k-p-1} + \mathcal{X}^{k-p-2} \otimes \mathcal{X}^p \otimes I dy \\ &= (-1)^{p+1} \int_Y ((-\Delta_{yy} \mathcal{X}^{p+1}) \otimes \mathcal{X}^{k-p-1} - \mathcal{X}^{k-p-2} \otimes \mathcal{X}^p \otimes I) dy. \end{aligned}$$

Hence the formula is proved at order $p + 1$.
Now, the formula at order $p = k$ reads

$$\mathcal{X}^{k*} = (-1)^k \int_Y \mathcal{X}^0(-\Delta_{yy}\mathcal{X}^k)dy = (-1)^k \int_Y \mathcal{X}^k(-\Delta_{yy}\mathcal{X}^0)dy = (-1)^k \mathcal{X}^{k*},$$

which implies $\mathcal{X}^{k*} = 0$ if k is odd. Formula (3.8) follows easily from (3.7) with $k = 2p$. \square

For completeness, we provide a minor result which implies that there is no order k (even odd) such that $\mathcal{X}^k(y)$ is identically equal to zero. However, let us remark that some components $\mathcal{X}_{i_1 \dots i_k}^k(y)$ may vanish for some set of indices i_1, \dots, i_k , e.g. in case of invariances of the obstacle ηT along the cell axes.

Proposition 4. *The following identity holds:*

$$-\Delta_{yy}(\partial_{i_1 \dots i_k}^k \mathcal{X}_{i_1 \dots i_k}^k) = (-1)^k(k+1),$$

where we recall the implicit summation convention over the repeated indices $i_1 \dots i_k$.

Proof. The results clearly holds true for $k = 0$. For $k = 1$, it holds

$$-\Delta_{yy}\partial_i \mathcal{X}_i^1 = \partial_i(2\partial_i \mathcal{X}^0) = 2\Delta \mathcal{X}^0 = -2.$$

Assuming the result holds true till rank $k - 1$, the formula still holds at rank $k \geq 2$ because

$$\begin{aligned} -\Delta_{yy}\partial_{i_1 \dots i_k}^k \mathcal{X}_{i_1 \dots i_k}^k &= \partial_{i_1 \dots i_k}^k (2\partial_{i_k} \mathcal{X}_{i_1 \dots i_{k-1}}^{k-1} + \mathcal{X}_{i_1 \dots i_{k-2}}^{k-2} \delta_{i_{k-1} i_k}) \\ &= 2\Delta_{yy}(\partial_{i_1 \dots i_{k-1}}^{k-1} \mathcal{X}_{i_1 \dots i_{k-1}}^{k-1}) + \Delta_{yy}(\partial_{i_1 \dots i_{k-2}}^{k-2} \mathcal{X}_{i_1 \dots i_{k-2}}^{k-2}) \\ &= -2(-1)^{k-1}k - (-1)^{k-2}(k-1) \\ &= (-1)^k(k+1). \end{aligned}$$

\square

3.3. Infinite order homogenized equation and criminal ansatz: tensors M^k and N^k

This part outlines the steps (2) and (3) of the procedure outlined section 2.2. Let us recall that the first tensor \mathcal{X}^{0*} is a strictly positive number, since (3.8) implies $\mathcal{X}^{0*} = \int_Y |\nabla \mathcal{X}^0|^2 dy > 0$.

Proposition 5. *Let $(M^k)_{i \in \mathbb{N}}$ be the family of k -th order tensors defined by induction as follows:*

$$\begin{cases} M^0 = (\mathcal{X}^{0*})^{-1}, \\ M^k = -(\mathcal{X}^{0*})^{-1} \sum_{p=0}^{k-1} \mathcal{X}^{k-p*} \otimes M^p. \end{cases} \quad (3.9)$$

Then it holds, given the definitions (3.1) and (3.5) of u_i^* :

$$\forall i \in \mathbb{N}, f_i(x) = \sum_{k=0}^i M^k \cdot \nabla^k u_{i-k}^*(x). \quad (3.10)$$

Recognizing a Cauchy product, (3.10) can be rewritten formally in terms of the following ‘‘infinite order’’ homogenized equation for the ‘‘infinite order’’ homogenized average u_ε^* of (2.8):

$$\sum_{i=0}^{+\infty} \varepsilon^{i-2} M^i \cdot \nabla^i u_\varepsilon^* = f. \quad (3.11)$$

Proof. We proceed by induction. The case $i = 0$ results from the identity $u_0^*(x) = \mathcal{X}^{0*} f_0(x)$ which yields $f_0(x) = (\mathcal{X}^{0*})^{-1} u_0^*(x)$. Then, assuming (3.10) holds till rank $i - 1$ with $i \geq 1$, we average (3.3) with respect to the y variable to obtain

$$u_i^* = \sum_{p=0}^i \mathcal{X}^{p*} \cdot \nabla^p f_{i-p} = \mathcal{X}^{0*} f_i + \sum_{p=1}^i \mathcal{X}^{p*} \cdot \nabla^p f_{i-p}.$$

By using (3.10) at ranks $i - p$ with $1 \leq p \leq i$, we obtain the following expression for f_i :

$$\begin{aligned}
f_i &= (\mathcal{X}^{0*})^{-1} \left(u_i^* - \sum_{p=1}^i \sum_{q=0}^{i-p} (\mathcal{X}^{p*} \otimes M^q) \cdot \nabla^{p+q} u_{i-p-q}^* \right) \\
&= (\mathcal{X}^{0*})^{-1} \left(u_i^* - \sum_{p=1}^i \sum_{k=p}^i (\mathcal{X}^{p*} \otimes M^{k-p}) \cdot \nabla^k u_{i-k}^* \right) \quad (\text{by } k \leftarrow p + q) \\
&= (\mathcal{X}^{0*})^{-1} \left(u_i^* - \sum_{k=1}^i \sum_{p=1}^k (\mathcal{X}^{p*} \otimes M^{k-p}) \cdot \nabla^k u_{i-k}^* \right) \quad (\text{inversion of summation}) \\
&= (\mathcal{X}^{0*})^{-1} \left(u_i^* - \sum_{k=1}^i \left(\sum_{p=0}^{k-1} \mathcal{X}^{k-p*} \otimes M^p \right) \cdot \nabla^k u_{i-k}^* \right) \quad (\text{by } p \leftrightarrow k - p) \\
&= M^0 u_i^* + \sum_{k=1}^i M^k \cdot \nabla^k u_{i-k}^*,
\end{aligned}$$

which yields the result at rank i . □

Corollary 1. $M^k = 0$ for any odd value of k .

Proof. If k is odd, then $k - p$ and p have distinct parities in (3.9). Therefore, the result follows by induction and by using $\mathcal{X}^{k-p*} = 0$ for even values of p (proposition 3). □

It is possible to write a more explicit formula for the tensors M^k :

Proposition 6. The tensors M^k are explicitly given by $M^0 = (\mathcal{X}^{0*})^{-1}$ and:

$$\forall k \geq 1, M^k = \sum_{p=1}^k \frac{(-1)^p}{(\mathcal{X}^{0*})^{p+1}} \sum_{\substack{i_1 + \dots + i_p = k \\ 1 \leq i_1 \dots i_p \leq k}} \mathcal{X}^{i_1*} \otimes \dots \otimes \mathcal{X}^{i_p*}. \quad (3.12)$$

Proof. For $k = 1$, the result is true because

$$M^1 = -(\mathcal{X}^{0*})^{-1} M^0 \mathcal{X}^{1*} = -(\mathcal{X}^{0*})^{-2} \mathcal{X}^{1*}$$

which is exactly (3.12). Assuming (3.12) holds till rank $k \geq 1$, we compute

$$\begin{aligned}
M^{k+1} &= -(\mathcal{X}^{0*})^{-1} \sum_{p=0}^k \mathcal{X}^{k+1-p*} \otimes M^p \\
&= -(\mathcal{X}^{0*})^{-1} M^0 \mathcal{X}^{k+1*} \\
&\quad - (\mathcal{X}^{0*})^{-1} \sum_{p=1}^k \sum_{q=1}^p \frac{(-1)^q}{(\mathcal{X}^{0*})^{q+1}} \mathcal{X}^{k+1-p*} \otimes \sum_{\substack{i_1+\dots+i_q=p \\ 1 \leq i_1 \dots i_q \leq p}} \mathcal{X}^{i_1*} \otimes \dots \otimes \mathcal{X}^{i_q*} \\
&= -(\mathcal{X}^{0*})^{-2} \mathcal{X}^{k+1*} \\
&\quad - (\mathcal{X}^{0*})^{-1} \sum_{q=1}^k \frac{(-1)^q}{(\mathcal{X}^{0*})^{q+1}} \sum_{p=q}^k \sum_{\substack{i_1+\dots+i_q=p \\ 1 \leq i_1 \dots i_q \leq p}} \mathcal{X}^{k+1-p*} \otimes \mathcal{X}^{i_1*} \otimes \dots \otimes \mathcal{X}^{i_q*} \\
&= -(\mathcal{X}^{0*})^{-2} \mathcal{X}^{k+1*} \\
&\quad - (\mathcal{X}^{0*})^{-1} \sum_{q=1}^k \frac{(-1)^q}{(\mathcal{X}^{0*})^{q+1}} \sum_{\substack{i_1+\dots+i_{q+1}=k+1 \\ 1 \leq i_1 \dots i_{q+1} \leq k+1}} \mathcal{X}^{i_{q+1}*} \otimes \mathcal{X}^{i_1*} \otimes \dots \otimes \mathcal{X}^{i_q*} \\
&= -(\mathcal{X}^{0*})^{-2} \mathcal{X}^{k+1*} + \sum_{q=2}^{k+1} \frac{(-1)^q}{(\mathcal{X}^{0*})^{q+1}} \sum_{\substack{i_1+\dots+i_q=k+1 \\ 1 \leq i_1 \dots i_q \leq k+1}} \mathcal{X}^{i_1*} \otimes \dots \otimes \mathcal{X}^{i_q*},
\end{aligned}$$

from where the result follows. \square

Remark 2. This result essentially states that $\sum_{k=0}^{+\infty} \varepsilon^k M^k \cdot \nabla^k$ is the formal series expansion of

$$\left(\sum_{k=0}^{+\infty} \varepsilon^k \mathcal{X}^k \cdot \nabla^k \right)^{-1}.$$

Indeed, it is elementary to show the following identity for the inverse of a power series $\sum_{k=0}^{+\infty} a_k z^k$ with $(a_k) \in \mathbb{C}^{\mathbb{N}}$, $z \in \mathbb{C}$ and radius of convergence $R > 0$:

$$\left(\sum_{k=0}^{+\infty} a_k z^k \right)^{-1} = a_0^{-1} + \sum_{k=1}^{+\infty} \left(\sum_{p=1}^k \frac{(-1)^p}{a_0^{p+1}} \sum_{\substack{i_1+\dots+i_p=k \\ 1 \leq i_1 \dots i_p \leq k}} a_{i_1} a_{i_2} \dots a_{i_p} \right) z^k. \quad (3.13)$$

We now turn on the derivation of the ‘‘criminal’’ ansatz (2.11). Being guided by (2.10), this ansatz is obtained by writing the oscillatory part $u_i(x, y)$ in terms of the non oscillatory part $u_i^*(x)$:

Proposition 7. *Given the previous definitions (3.1) and (3.5) of respectively u_i and u_i^* , the following identity holds:*

$$\forall i \geq 0, u_i(x, y) = \sum_{k=0}^i \left(\sum_{p=0}^k M^p \otimes \mathcal{X}^{k-p}(y) \right) \cdot \nabla^k u_{i-k}^*(x). \quad (3.14)$$

Proof. We substitute (3.10) into (3.3), which yields

$$\begin{aligned}
u_i(x, y) &= \sum_{p=0}^i \sum_{q=0}^{i-p} (\mathcal{X}^p(y) \otimes M^q) \cdot \nabla^{p+q} u_{i-p-q}^*(x) \\
&= \sum_{p=0}^i \sum_{k=p}^i (\mathcal{X}^p(y) \otimes M^{p-k}) \cdot \nabla^k u_{i-k}^*(x) \quad (\text{change of indices } k = p + q) \\
&= \sum_{k=0}^i \sum_{p=0}^k (\mathcal{X}^p(y) \otimes M^{p-k}) \cdot \nabla^k u_{i-k}^*(x) \quad (\text{interversion of summation})
\end{aligned}$$

The result follows by performing a last change of indices $p \leftrightarrow k - p$. \square

This result motivates the definition of the tensors N^k of (2.11):

Definition 2. For any $k \geq 0$, we denote by $N^k(y)$ the k -th order tensor

$$N^k(y) := \sum_{p=0}^k M^p \otimes \mathcal{X}^{k-p}(y), \quad y \in Y. \quad (3.15)$$

Recognizing a Cauchy product, the identity (3.14) rewrites as expected as the ‘‘criminal’’ ansatz (2.11) which expresses the oscillating solution u_ε in terms of its formal homogenized averaged u_ε^* (defined in (2.8)):

$$u_\varepsilon(x) = \sum_{k=0}^{+\infty} \varepsilon^k N^k(x/\varepsilon) \cdot \nabla^k u_\varepsilon^*(x), \quad x \in D_\varepsilon.$$

The last proposition of this section gathers several important properties for the tensors N^k that are dual to those of the tensors \mathcal{X}^k stated in sections 3.1 and 3.2.

Proposition 8. *The tensor $N^k(y)$ satisfies:*

1. $\int_Y N^0(y) dy = 1$ and $\int_Y N^k(y) dy = 0$ for any $k \geq 1$.
2. For any $k \geq 0$ and assuming the convention $N^{-1} = N^{-2} = 0$:

$$-\Delta_{yy} N^{k+2} = 2\partial_j N^{k+1} \otimes e_j + N^k \otimes I + M^{k+2}. \quad (3.16)$$

3. For any $k \geq 0$,

$$-\Delta_{yy} (\partial_{i_1 \dots i_k}^k N_{i_1 \dots i_k}^k) = (-1)^k (k+1) M^0. \quad (3.17)$$

4. For any $k \geq 1$ and $1 \leq p \leq k-1$,

$$M^k = (-1)^{p+1} \int_Y (N^{k-p} \otimes (-\Delta_{yy} N^p) - N^{k-p-1} \otimes N^{p-1} \otimes I) dy, \quad (3.18)$$

In particular, M^{2p} depends only on the tensors N^p and N^{p-1} , which depend themselves only on the first $p+1$ tensors $\mathcal{X}^0 \dots \mathcal{X}^p$.

Proof. 1. For $k=0$, it holds $\int_Y N^0(y) dy = M^0 \mathcal{X}^{0*} = 1$. Furthermore, the definition (3.9) of the tensor M^k can be rewritten as

$$\forall k \geq 1, \int_Y N^k(y) dy = \sum_{p=0}^k \mathcal{X}^{k-p*} \otimes M^p = 0.$$

2. The cases $k=0$ and $k=1$ are easily verified. The case $k \geq 2$ is obtained by writing

$$\begin{aligned}
-\Delta_{yy} N^k &= \sum_{p=0}^k M^{k-p} \otimes (-\Delta_{yy} \mathcal{X}^p) \\
&= \sum_{p=2}^k M^{k-p} \otimes (2\partial_j \mathcal{X}^{p-1} \otimes e_j + \mathcal{X}^{p-2} \otimes I) + M^{k-1} \otimes (2\partial_j \mathcal{X}^0 \otimes e_j) + M^k \\
&= 2\partial_j \left(\sum_{p=1}^k M^{k-p} \otimes \mathcal{X}^{p-1} \right) \otimes e_j + \left(\sum_{p=0}^{k-2} M^p \otimes \mathcal{X}^{k-p-2} \right) \otimes I + M^k,
\end{aligned}$$

from where the result follows.

3. The proof of (3.17) is identical to that of [proposition 4](#).

4. We start by proving the result for $p = 1$ with $k > 1$: using the point 1., we write

$$\begin{aligned} M^k &= \int_Y N^0 \otimes M^k dy = \int_Y N^0 \otimes (-\Delta N^k - 2\partial_j N^{k-1} \otimes e_j - N^{k-2} \otimes I) dy \\ &= \int_Y M^0 \otimes N^k dy + \int_Y (2\partial_j N^0 \otimes e_j \otimes N^{k-1} - N^0 \otimes N^{k-2} \otimes I) dy \\ &= \int_Y ((-\Delta_{yy} N^1) \otimes N^{k-1} - N^0 \otimes N^{k-2} \otimes I) dy. \end{aligned}$$

Assuming now the result holds until rank p with $1 \leq p \leq k - 2$, we prove it at rank $p + 1$ thanks to analogous computations:

$$\begin{aligned} M^k &= (-1)^{p+1} \int_Y (M^{k-p} + 2\partial_j N^{k-p-1} \otimes e_j + N^{k-p-2} \otimes I) \otimes N^p \\ &\quad - N^{k-p-1} \otimes N^{p-1} \otimes I) dy \\ &= (-1)^{p+1} \int_Y ((-2\partial_j N^p \otimes e_j - N^{p-1} \otimes I) \otimes N^{k-p-1} \\ &\quad + N^{k-p-2} \otimes N^p \otimes I) dy \\ &= (-1)^{p+1} \int_Y ((\Delta_{yy} N^{p+1} + M^{p+1}) \otimes N^{k-p-1} + N^{k-p-2} \otimes N^p \otimes I) dy. \end{aligned}$$

□

3.4. Simplifications for the tensors \mathcal{X}^{k*} and M^k in case of symmetries

In this last part, we analyze how the homogenized tensors \mathcal{X}^{k*} and M^k reduce to small number of effective coefficients when the obstacle ηT is symmetric with respect to axes of the unit cell P . Such results are classical in the theory of homogenization; our methodology follows e.g. section 6 in [\[12\]](#).

In all what follows, we denote by $S := (S_{ij})_{1 \leq i, j \leq d}$ an arbitrary orthogonal symmetry (satisfying $S = S^T$ and $SS = I$). We shall specialize S in [corollary 3](#) below to either of the following cell symmetries:

- for $1 \leq l \leq d$, S^l denotes the symmetry with respect to the hyperplane orthogonal to e_l :

$$S^l := I - 2e_l e_l^T;$$

- for $1 \leq m \neq l \leq d$, S^{lm} denotes the symmetry with respect to the diagonal hyperplane that is orthogonal to $e_l - e_m$:

$$S^{lm} := I - e_l e_l^T - e_m e_m^T + e_l e_m^T + e_m e_l^T.$$

Recall the Laplace operator is invariant under such orthogonal symmetries S : for any smooth scalar field \mathcal{X} ,

$$-\Delta(\mathcal{X} \circ S) = -(\Delta \mathcal{X}) \circ S.$$

Proposition 9. *If the cell $Y = P \setminus \eta T$ is invariant with respect to a symmetry S , i.e. $S(Y) = Y$, then the following identity holds for the components of the solutions \mathcal{X}^k to the cell problem (3.2):*

$$\mathcal{X}_{i_1 \dots i_k}^k \circ S = S_{i_1 j_1} \dots S_{i_k j_k} \mathcal{X}_{j_1 \dots j_k}^k,$$

where we recall the implicit summation convention over the repeated indices $j_1 \dots j_k$.

Proof. The result is proved by induction on k . For $k = 0$, it holds

$$-\Delta_{yy}(\mathcal{X}^0 \circ S) = 1 \circ S = 1$$

and the symmetry of Y implies that $\mathcal{X}^0 \circ S$ also satisfies the boundary conditions of (3.2). This implies $\mathcal{X}^0 \circ S = \mathcal{X}^0$. For $k = 1$, we write

$$-\Delta_{yy}(\mathcal{X}_{i_1}^1 \circ S) = 2(\partial_{i_1} \mathcal{X}^0) \circ S = 2\partial_{j_1}(\mathcal{X}^0 \circ S) S_{i_1 j_1} = 2\partial_{j_1} \mathcal{X}^0 S_{i_1 j_1},$$

which implies similarly $\mathcal{X}_{i_1}^1 \circ S = S_{i_1 j_1} \mathcal{X}_{j_1}^1$. Finally, if the result holds till rank $k + 1$ with $k \geq 0$, then

$$\begin{aligned} -\Delta_{yy}(\mathcal{X}_{i_1 \dots i_{k+2}}^{k+2} \circ S) &= 2(\partial_{i_{k+2}} \mathcal{X}_{i_1 \dots i_{k+1}}^{k+1}) \circ S + \delta_{i_{k+1} i_{k+2}} \mathcal{X}_{i_1 \dots i_k}^k \circ S \\ &= 2S_{i_{k+2} j_{k+2}} \partial_{j_{k+2}} (\mathcal{X}_{i_1 \dots i_{k+1}}^{k+1} \circ S) + S_{i_{k+1} j_{k+1}} S_{i_{k+2} j_{k+2}} \delta_{j_{k+1} j_{k+2}} \mathcal{X}_{i_1 \dots i_k}^k \circ S \\ &= -S_{i_1 j_1} \dots S_{i_k j_{k+2}} \Delta_{yy} \mathcal{X}_{j_1 \dots j_{k+2}}^{k+2}, \end{aligned}$$

whence the result at rank $k + 2$. \square

Corollary 2. *If the cell $Y = P \setminus \eta T$ is invariant with respect to a symmetry S , then the components of the tensors \mathcal{X}^{k*} and M^k of respectively (3.6) and (3.9) satisfy:*

$$\mathcal{X}_{i_1 \dots i_k}^{k*} = S_{i_1 j_1} \dots S_{i_k j_k} \mathcal{X}_{j_1 \dots j_k}^{k*} \quad (3.19)$$

$$M_{i_1 \dots i_k}^k = S_{i_1 j_1} \dots S_{i_k j_k} M_{j_1 \dots j_k}^k \quad (3.20)$$

where we recall the implicit summation over the repeated indices $j_1 \dots j_k$.

Proof. Equality (3.19) results from the previous proposition and from the following change of variables:

$$\mathcal{X}_{i_1 \dots i_k}^{k*} = \int_Y \mathcal{X}_{i_1 \dots i_k}^k dy = \int_Y \mathcal{X}_{i_1 \dots i_k}^k \circ S dy.$$

Equality (3.20) can be obtained by using (3.19) in the formula (3.12). \square

Corollary 3. (1) *If the cell Y is symmetric with respect to all cell axes e_l , i.e. $S^l(Y) = Y$ for any $1 \leq l \leq d$, then*

$$\mathcal{X}_{i_1 \dots i_k}^{k*} = 0 \text{ and } M_{i_1 \dots i_k}^k = 0$$

whenever there exists a number r occurring with an odd multiplicity in the indices $i_1 \dots i_k$, i.e. whenever

$$\exists r \in \{1, \dots, d\}, \text{Card}\{j \in \{1, \dots, k\} \mid i_j = r\} \text{ is odd.}$$

(2) *If the cell Y is symmetric with respect to all diagonal axes $(e_l - e_m)$, i.e. $S^{l,m}(Y) = Y$ for any $1 \leq l < m \leq d$, then for any permutation $\sigma \in \mathfrak{S}_d$,*

$$\mathcal{X}_{\sigma(i_1) \dots \sigma(i_k)}^{k*} = \mathcal{X}_{i_1 \dots i_k}^{k*}.$$

$$M_{\sigma(i_1) \dots \sigma(i_k)}^k = M_{i_1 \dots i_k}^k.$$

Proof. (1) The symmetry S^l is a diagonal matrix satisfying $S^l e_l = -e_l$ and $S^l e_q = e_q$ for $q \neq l$. Hence, replacing S by S^l in (3.19), it holds

$$\mathcal{X}_{i_1 \dots i_k}^{k*} = (-1)^{\delta_{i_1 l} + \dots + \delta_{i_k l}} \mathcal{X}_{i_1 \dots i_k}^{k*},$$

which implies the result.

(2) Applying (3.19) to the symmetry $S^{l,m}$ yields the result for $\sigma = \tau$ where τ is the transposition exchanging l and m . Since this holds for any transposition $\tau \in \mathfrak{S}_d$, this implies the statement for any permutation $\sigma \in \mathfrak{S}_d$. \square

Let us illustrate how the previous corollary reads for the tensors M^2 and M^4 :

- if Y is symmetric with respect to the cell axes $(e_l)_{1 \leq l \leq d}$, then only the coefficients M_{ii}^2 , M_{ijj}^4 , M_{iii}^4 with $1 \leq i, j \leq d$ and $i \neq j$ are non zeros (in particular M^2 is diagonal).
- if in addition Y is symmetric with respect to the hyperplane orthogonal to $e_l - e_m$, then these coefficients do not depend on the values of the distinct indices i and j . As a result, M^2 is a multiple of the identity and M^4 reduces to two effective coefficients: there exists constants $\alpha, \beta, \nu \in \mathbb{R}$ such that

$$M^2 \cdot \nabla^2 = \nu \Delta \text{ and } M^4 \cdot \nabla^4 = \alpha \sum_{i=1}^d \partial_{iii}^4 + \beta \sum_{1 \leq i \neq j \leq d} \partial_{ijj}^4.$$

4. HOMOGENIZED EQUATIONS OF ORDER $2K + 2$: TENSORS \mathbb{D}_K^{2k}

This section details the steps (4) and (5) of [section 2.2](#) concerned with the process of truncating the infinite order homogenized equation (2.9) so as to obtain well-posed effective models of finite order. Recall that this process is needed because (1.10) is in general ill-posed, since the tensors M^k do not have any particular sign (in view of (3.12)).

The first [section 4.1](#) introduces the main technical results which allow to derive error estimates. More particularly we show in [section 4.1](#) that for any integer $K' \in \mathbb{N}$, any family of non-oscillating functions $(v_\varepsilon^*)_{\varepsilon > 0}$ yields an error estimate of order $O(\varepsilon^{K'+3})$ provided (i) v_ε^* is of order $O(\varepsilon^2)$ and (ii) v_ε^* solves the infinite order homogenized equation (1.9) up to a remainder of order $O(\varepsilon^{K'+1})$:

$$\sum_{k=0}^{K'} \varepsilon^{k-2} M^k \cdot \nabla^k v_\varepsilon^* = f + O(\varepsilon^{K'+1}). \quad (4.1)$$

In particular, this result reminds us that higher order models are generally not unique, they differ by the choice of extra differential operators of order greater than K' which turn (4.1) into a well-posed model.

Leaving momentarily these considerations aside, we propose in [section 4.2](#), a “variational” method inspired from [52, 23] which allows to construct a well-posed effective model (1.6) of order $2K + 2$ for any $K \in \mathbb{N}$. The procedure relies on an energy minimization principle based on the criminal ansatz (1.11); the coefficients \mathbb{D}_K^{2k} are inferred from an effective energy J_K^* ([definition 4](#)) and are *a priori* distinct from the tensors M^{2k} . These properties enable us to establish that the obtained model is *elliptic* (in particular, well-posed), hence amenable to numerical computations.

Finally, we obtain in [section 4.3](#) that surprisingly, it holds $\mathbb{D}_K^{2k} = M^{2k}$ for any $0 \leq 2k \leq K$, which implies that (4.1) is satisfied by the solution $v_\varepsilon^* \equiv v_{\varepsilon, K}^*$ with $K' = 2K + 1$ (recall $M^{2k+1} = 0$ for any $k \in \mathbb{N}$ from [corollary 1](#)). The error estimate (1.14) of order $O(\varepsilon^{2K+4})$ follows in [corollary 5](#).

4.1. Sufficient conditions under which an effective solution yields higher order approximations

The main result of this part is [proposition 10](#) where we provide sufficient conditions under which a sequence of sequence of macroscopic functions $v_\varepsilon^* \in \mathcal{C}^\infty(D)$ (depending on ε , K and f) yields a high order approximation of u_ε . The proof is based on the properties of the tensors N^k stated in [proposition 8](#) and the next three results.

Lemma 1 (see e.g. Lions (1981) [40]). *There exists a constant C independent of ε such that for any $\phi \in H^1(D_\varepsilon)$ satisfying $\phi = 0$ on the boundary $\partial\omega_\varepsilon$ of the holes, the following Poincaré inequality holds:*

$$\|\phi\|_{L^2(D_\varepsilon)} \leq C\varepsilon \|\nabla\phi\|_{L^2(D_\varepsilon, \mathbb{R}^d)}.$$

Corollary 4. *For any $h \in L^2(D_\varepsilon)$, let $r_\varepsilon \in H^1(D_\varepsilon)$ be the unique solution to the Poisson problem*

$$\begin{cases} -\Delta r_\varepsilon = h \text{ in } D_\varepsilon \\ r_\varepsilon = 0 \text{ on } \partial\omega_\varepsilon \\ r_\varepsilon \text{ is } D\text{-periodic.} \end{cases}$$

There exists a constant C independent of ε and h such that

$$\|r_\varepsilon\|_{L^2(D_\varepsilon)} + \varepsilon \|\nabla r_\varepsilon\|_{L^2(D_\varepsilon, \mathbb{R}^d)} \leq C \|h\|_{L^2(D_\varepsilon)} \varepsilon^2. \quad (4.2)$$

Proof. This result is classical; it is a consequence of [lemma 1](#) and of the energy estimate

$$\int_{D_\varepsilon} |\nabla r_\varepsilon|^2 dx = \int_{D_\varepsilon} h r_\varepsilon dx \leq \|h\|_{L^2(D_\varepsilon)} \|r_\varepsilon\|_{L^2(D_\varepsilon)} \leq C\varepsilon \|h\|_{L^2(D_\varepsilon)} \|\nabla r_\varepsilon\|_{L^2(D_\varepsilon, \mathbb{R}^d)}$$

which implies $\|\nabla r_\varepsilon\|_{L^2(D_\varepsilon, \mathbb{R}^d)} \leq C \|h\|_{L^2(D_\varepsilon)} \varepsilon$, then (4.2). □

Lemma 2. *Assume that the boundary of obstacle ηT is smooth. Then, for any $k \in \mathbb{N}$, the tensor \mathcal{X}^k is well-defined and is smooth, namely it holds $\mathcal{X}^k \in \mathcal{C}^\infty(\bar{Y})$. In particular, $\mathcal{X}^k \in L^\infty(Y) \cap H^1(Y)$.*

Proof. Since the constant function 1 is smooth, standard regularity theory for the Laplace operator $-\Delta_{yy}$ (see [35, 22, 31]) implies $\mathcal{X}^0 \in \mathcal{C}^\infty(\bar{Y})$. The result follows by induction by repeating this argument to \mathcal{X}^1 and \mathcal{X}^{k+2} for any $k \geq 0$. □

Proposition 10. Let $v_\varepsilon^* \in C^\infty(D)$ be a D -periodic function depending on ε (and possibly on K' and f) satisfying the following two hypotheses:

1. for any $m \in \mathbb{N}$, there exists a constant $C_{K',m}(f)$ depending only on m , K' and $f \in C^\infty(D)$ such that

$$\|v_\varepsilon^*\|_{H^m(D)} \leq C_{K',m}(f)\varepsilon^2 \quad (4.3)$$

2. v_ε^* solves the infinite order homogenized equation (1.9) up to a remainder of order $O(\varepsilon^{K'+1})$:

$$\left\| \sum_{k=0}^{K'} \varepsilon^{k-2} M^k \cdot \nabla^k v_\varepsilon^* - f \right\|_{L^2(D)} \leq C_{K'}(f)\varepsilon^{K'+1}. \quad (4.4)$$

Then the reconstructed function $W_{\varepsilon,K'}(v_\varepsilon^*)$ of (1.12) approximates the solution u_ε of the perforated Poisson problem (1.1) at order $O(\varepsilon^{K'+3})$, viz. there exists a constant $C_{K'}(f)$ independent of ε such that

$$\|u_\varepsilon - W_{\varepsilon,K'}(v_\varepsilon^*)\|_{L^2(D_\varepsilon)} + \varepsilon \|\nabla(u_\varepsilon - W_{\varepsilon,K'}(v_\varepsilon^*))\|_{L^2(D_\varepsilon, \mathbb{R}^d)} \leq C_{K'}(f)\varepsilon^{K'+3}.$$

Proof. Let us compute

$$\begin{aligned} -\Delta W_{\varepsilon,K'}(v_\varepsilon^*) &= \sum_{k=0}^{K'} \varepsilon^{k-2} (-\Delta N^k - 2\partial_l N^{k-1} \otimes e_l - N^{k-2} \otimes I)(\cdot/\varepsilon) \cdot \nabla^k v_\varepsilon^* \\ &\quad - \varepsilon^{K'-1} (2\partial_l N^{K'} \otimes e_l + N^{K'-1} \otimes I)(\cdot/\varepsilon) \cdot \nabla^{K'+1} v_\varepsilon^* \\ &\quad - \varepsilon^{K'} N^{K'}(\cdot/\varepsilon) \otimes I \cdot \nabla^{K'+2} v_\varepsilon^* \\ &= \sum_{k=0}^{K'} \varepsilon^{k-2} M^k \cdot \nabla^k v_\varepsilon^* - \varepsilon^{K'-1} (2\partial_l N^{K'} \otimes e_l + N^{K'-1} \otimes I)(\cdot/\varepsilon) \cdot \nabla^{K'+1} v_\varepsilon^* \\ &\quad - \varepsilon^{K'} N^{K'}(\cdot/\varepsilon) \otimes I \cdot \nabla^{K'+2} v_\varepsilon^*, \end{aligned}$$

where we have used (3.16) to obtain the second equality. Since the functions $(N^k)_{k \in \mathbb{N}}$ are smooth (lemma 2 and (3.15)), assumption (4.3) implies that the last two terms are lower than $\varepsilon^{K'+1}$:

$$\begin{aligned} \left\| \varepsilon^{K'-1} (2\partial_l N^{K'} \otimes e_l + N^{K'-1} \otimes I)(\cdot/\varepsilon) \cdot \nabla^{K'+1} v_\varepsilon^* \right\| &\leq C_{K'}(f)\varepsilon^{K'+1}, \\ \left\| \varepsilon^{K'} N^{K'}(\cdot/\varepsilon) \otimes I \cdot \nabla^{K'+2} v_\varepsilon^* \right\|_{L^2(D_\varepsilon)} &\leq C_{K'}(f)\varepsilon^{K'+1}. \end{aligned}$$

Using now assumption (4.4) and applying corollary 4 to $r_\varepsilon := u_\varepsilon - W_{\varepsilon,K'}(v_\varepsilon^*)$ yields the result. \square

4.2. Construction of a well-posed higher order effective model by mean of a variational principle

Leaving momentarily aside the result of proposition 10, we now detail the construction of our effective model (1.6) of finite order which is inspired from the works [15, 52, 23]. The construction of the coefficients \mathbb{D}_K^{2k} from an effective energy is exposed in section 4.2.1, and the well-posedness of the effective model is established in section 4.2.2.

4.2.1. The method of Smyshlyaev and Cherednichenko

According to the ideas outlined in step (4) of section 2.2, we consider truncations $W_{\varepsilon,K}(v)$ of the ‘‘criminal’’ ansatz (2.11) of the form:

$$W_{\varepsilon,K}(v)(x) := \sum_{k=0}^K \varepsilon^k N^k(x/\varepsilon) \cdot \nabla^k v(x), \quad x \in D, \quad (4.5)$$

where we seek a function $v \in H^{K+1}(D)$ which does not depend on the fast variable x/ε and which approximates the formal homogenized average u_ε^* of (2.8). The tensors $N^k(y)$ is extended by 0 in $P \setminus Y$ for (4.5) to make sense in $D \setminus D_\varepsilon$.

For any $u \in H^1(D)$ and $f \in L^2(D)$, we denote by $J(u, f)$ the energy

$$J(u, f) := \int_D \left(\frac{1}{2} |\nabla u|^2 - fu \right) dx.$$

Following the lines of the point (4) of [section 2.2](#), we consider the problem [\(2.14\)](#) of finding a minimizer of $v \mapsto J(W_{\varepsilon,K}(v), f)$. The key step of the strategy is to eliminate the fast variable x/ε in $J(W_{\varepsilon,K}(v), f)$ so as to obtain an effective energy $J_K^*(v, f, \varepsilon) \simeq J(W_{\varepsilon,K}(v), f)$ which does not involve oscillating functions. The main technical tool which allows us to perform this operation is the following classical lemma of two-scale convergence (see e.g. Appendix C. of [\[52\]](#) or [\[5\]](#)).

Lemma 3. *Let ϕ be a $P = [0, 1]^d$ -periodic function and $f \in \mathcal{C}^\infty(D)$ be a smooth D -periodic function. Then for any $k \in \mathbb{N}$ arbitrarily large, the following inequality holds:*

$$\left| \int_D f(x)\phi(x/\varepsilon)dx - \int_D \int_P f(x)\phi(y)dydx \right| \leq \frac{L^{d/2}}{|2\pi|^p} \left\| \phi - \int_P \phi dy \right\|_{L^2(P)} \|f\|_{H^p(D)} \varepsilon^p.$$

Proof. See Lemma 7.3 in [\[32\]](#) for a proof of this exact statement. \square

Before providing the definition of J_K^* based on the application of [lemma 3](#), we introduce several additional tensors that arise in the averaging process.

Definition 3 (Tensors $\mathbb{B}_K^{l,m}$). For any $K \in \mathbb{N}$, $1 \leq j \leq d$ and $0 \leq k \leq K + 1$, let $\tilde{N}_j^k(y)$ (with implicit dependence with respect to K) be the k -th order tensor defined by

$$\tilde{N}_j^k(y) = \begin{cases} \partial_j N^0(y) & \text{if } k = 0 \\ \partial_j N^k(y) + N^{k-1}(y) \otimes e_j & \text{if } 1 \leq k \leq K \\ N^K(y) \otimes e_j & \text{if } k = K + 1. \end{cases} \quad (4.6)$$

We define a family of constant bilinear tensors $\mathbb{B}_K^{l,m}$ of order $l + m$ by the formula

$$\mathbb{B}_K^{l,m} := \int_Y \tilde{N}_j^l(y) \otimes \tilde{N}_j^m(y) dy, \text{ for any } 0 \leq l, m \leq K + 1, \quad (4.7)$$

where the Einstein summation convention is still assumed over the repeated subscript index $1 \leq j \leq d$.

Definition 4 (Approximate energy J_K^*). For any $f \in L^2(D)$ and periodic function $v \in H^{K+1}(D)$, we define

$$J_K^*(v, f, \varepsilon) := \int_D \left(\frac{1}{2} \sum_{l,m=0}^{K+1} \varepsilon^{l+m-2} \mathbb{B}_K^{l,m} \nabla^l v \nabla^m v - f v \right) dx. \quad (4.8)$$

where we recall [\(2.5\)](#) for the definition of $\mathbb{B}_K^{l,m} \nabla^l v \nabla^m v$.

The definition of the energy $J_K^*(v, f, \varepsilon)$ is motivated by the following asymptotic—provided by [lemma 3](#)—which holds with any $p \geq 0$ arbitrarily large

$$J(W_{\varepsilon,K}(v), f) = J_K^*(v, f, \varepsilon) + o(\varepsilon^p).$$

More precisely, we have the following result:

Proposition 11. *Assume $f \in \mathcal{C}^\infty(D)$ is D -periodic. Let $v \in \mathcal{C}^\infty(D)$ be a smooth D -periodic function and $W_{\varepsilon,K}(v) \in \mathcal{C}^\infty(D_\varepsilon)$ be the truncated ansatz of the form of [\(4.5\)](#). The following estimate holds true with $p \geq 0$ arbitrarily large:*

$$|J(W_{\varepsilon,K}(v), f) - J_K^*(v, f, \varepsilon)| \leq C_{K,p} (\|v\|_{H^{p+2}(D)}^2 + \|f\|_{H^p(D)}^2) \varepsilon^p.$$

for a constant $C_{K,p}$ depending only on p and K (and η).

Proof. For any $1 \leq j \leq d$, the partial derivative $\partial_{x_j} W_{\varepsilon,K}(v)$ reads

$$\begin{aligned} \partial_{x_j} W_{\varepsilon,K}(v) &= \sum_{k=0}^K (\varepsilon^{k-1} \partial_{y_j} N^k(\cdot/\varepsilon) \cdot \nabla^k v + \varepsilon^k N^k(\cdot/\varepsilon) \otimes e_j \cdot \nabla^{k+1} v) \\ &= \sum_{i=0}^{K+1} \varepsilon^{i-1} \tilde{N}_j^i(\cdot/\varepsilon) \cdot \nabla^i v, \end{aligned}$$

by the definition (4.6) of the tensors \tilde{N}_j^k . The computation of the energy $J(W_{\varepsilon,K}(v), f)$ yields then

$$\begin{aligned} J(W_{\varepsilon,K}(v), f) &= \int_D \left(\frac{1}{2} \sum_{l,m=0}^{K+1} \varepsilon^{l+m-2} (\tilde{N}_j^l(\cdot/\varepsilon) \cdot \nabla^l v) (\tilde{N}_j^m(\cdot/\varepsilon) \cdot \nabla^m v) \right) dx \\ &\quad - \int_D \sum_{l=0}^K \varepsilon^l (N^l(\cdot/\varepsilon) \cdot \nabla^l v) f dx. \end{aligned}$$

The result follows by estimating both term after applying lemma 3: for any $0 \leq l, m \leq K+1$,

$$\begin{aligned} &\left| \int_D \varepsilon^{l+m-2} \left((\tilde{N}_j^l(\cdot/\varepsilon) \cdot \nabla^l v) (\tilde{N}_j^m(\cdot/\varepsilon) \cdot \nabla^m v) - \mathbb{B}_K^{l,m} \nabla^l v \nabla^m v \right) dx \right| \\ &\leq C_p \varepsilon^p \|\nabla^l v \otimes \nabla^m v\|_{H^{p-(l+m-2)}(D, \mathbb{R}^{d^{l+m}})} \\ &\leq C'_p \varepsilon^p (\|\nabla^l v\|_{H^{p-(l+m-2)}(D, \mathbb{R}^{d^l})}^2 + \|\nabla^m v\|_{H^{p-(l+m-2)}(D, \mathbb{R}^{d^m})}^2) \\ &\leq C''_p \varepsilon^p \|v\|_{H^{p+2}(D)}^2, \\ \forall 0 \leq l \leq K, &\left| \int_D \varepsilon^l (N^l(\cdot/\varepsilon) \cdot \nabla^l v f - f v) dx \right| \leq C_p \varepsilon^p \|f \nabla^l v\|_{H^{p-l}(D, \mathbb{R}^{d^l})} \\ &\leq C'_p \varepsilon^p (\|f\|_{H^{p-l}(D)}^2 + \|v\|_{H^p(D)}^2) \leq C'_p \varepsilon^p (\|f\|_{H^p(D)}^2 + \|v\|_{H^p(D)}^2), \end{aligned}$$

where we used that N^l is of average 1 if $l = 0$ and 0 otherwise (point 1. of proposition 8). \square

The approximate energy (4.8) is used (instead of $J(W_{\varepsilon,K}(v), f)$ in (2.14)) in order to construct our higher order homogenized model.

Definition 5. For any $K \in \mathbb{N}$, we call homogenized equation of order $2K+2$ the Euler–Lagrange equation associated with the minimization problem

$$\begin{aligned} &\min_{v \in H^{K+1}(D)} J_K^*(v, f, \varepsilon) \\ &\quad s.t. \quad v \text{ is } D\text{-periodic.} \end{aligned} \tag{4.9}$$

This equation reads explicitly in terms of a higher order homogenized solution $v_K^* \in H^{K+1}(D)$ as

$$\begin{cases} \sum_{k=0}^{K+1} \varepsilon^{2k-2} \mathbb{D}_K^{2k} \cdot \nabla^{2k} v_{\varepsilon,K}^* = f \\ v_{\varepsilon,K}^* \text{ is } D\text{-periodic,} \end{cases} \tag{4.10}$$

where the constant tensors \mathbb{D}_K^{2k} are defined for any $0 \leq 2k \leq 2K+2$ by:

$$\mathbb{D}_K^{2k} := \sum_{l=0}^{2k} (-1)^l \mathbb{B}_K^{l, 2k-l}, \tag{4.11}$$

assuming the convention $\mathbb{B}_K^{l,m} = 0$ for any $l > K+1$ or $m > K+1$.

Proof. Let us detail slightly the derivation of (4.10). The Euler–Lagrange equation of (4.9) reads, after an integration by parts:

$$\sum_{l,m=0}^{K+1} \varepsilon^{l+m-2} \frac{1}{2} ((-1)^m \mathbb{B}_K^{l,m} + (-1)^l \mathbb{B}_K^{m,l}) \nabla^{l+m} v_{\varepsilon,K}^* = f. \tag{4.12}$$

Since $\mathbb{B}_K^{l,m} = \mathbb{B}_K^{m,l}$ and $(-1)^l + (-1)^m$ vanishes when l and m are not of the same parity, only terms such that $l+m$ is even are possibly not zero in the above equation. Hence, (4.12) rewrites as (4.10) with

$$\mathbb{D}_K^{2k} = \sum_{l+m=2k} \frac{1}{2} ((-1)^l + (-1)^m) \mathbb{B}_K^{l,m} = \sum_{l=0}^{2k} \frac{1}{2} ((-1)^l + (-1)^{2k-l}) \mathbb{B}_K^{l, 2k-l}$$

for any $0 \leq 2k \leq 2K+2$, which eventually yields the desired expression (4.11). \square

Remark 3. As announced in the introduction, (4.10) turns out to be a simple correction of (1.10), see proposition 13 below.

4.2.2. Well-posedness of the homogenized model of order $2K + 2$

We now establish the well-posedness of the high order homogenized model (4.10). More precisely, we prove its ellipticity, which implies the existence and the uniqueness of the effective solution $v_{\varepsilon, K}^*$. Before stating the result, let us stress the following observation which is obvious, but a somewhat important consequence of the definition (4.7).

Lemma 4. *The dominant tensor $\mathbb{B}_K^{K+1, K+1}$ is symmetric and non-negative.*

Proposition 12. *Assume further that the dominant tensor $\mathbb{B}_K^{K+1, K+1}$ is non-degenerate, viz. there exists a constant $\nu > 0$ such that*

$$\forall \xi = \xi_{i_1 \dots i_{K+1}} \in \mathbb{R}^{d^{K+1}}, \mathbb{B}_K^{K+1, K+1} \cdot \xi \xi \geq \nu |\xi|^2. \quad (4.13)$$

Then (4.10) is elliptic and there exists a unique solution $v_K^ \in H^{K+1}(D)$ to the homogenized equation (4.10) of order $2K + 2$.*

Proof. Let us consider the space $V_K := \{v \in H^{K+1}(D) \mid v \text{ is } D\text{-periodic}\}$ and introduce $a : V_K \times V_K \rightarrow \mathbb{R}$ and $b : V_K \rightarrow \mathbb{R}$ the respective bilinear and linear forms defined for any $v \in V_K$ by

$$a(v, v) = \int_D \sum_{l, m=0}^{K+1} \varepsilon^{l+m-2} \mathbb{B}_K^{l, m} \nabla^l v \nabla^m v dx,$$

$$b(v) = \int_D f v dx.$$

The homogenized equation (4.10) reduces to the following variational problem:

$$\text{find } v_{\varepsilon, K}^* \in V_K \text{ such that } \forall v \in V_K, a(v_{\varepsilon, K}^*, v) = b(v). \quad (4.14)$$

From there, one could directly rely on the theory of Fredholm operators [42] to conclude to the existence of a solution $v_{\varepsilon, K}^*$. However we are going to show that a is coercive (meaning (4.10) is elliptic), which will allow us to apply Lax–Milgram theorem [30].

Under the non-degeneracy assumption (4.13), it is readily obtained that there exists a constant C_ε (depending on ε) such that for any $v \in V_K(D)$,

$$a(v, v) \geq (\varepsilon^{2K} \nu) \|\nabla^{K+1} v\|_{L^2(D, \mathbb{R}^d)}^2 + \varepsilon^{-2} M^0 \|v\|_{L^2(D)}^2 - C_\varepsilon \|v\|_{H^{K+1}(D)} \|v\|_{H^K(D)}.$$

Remembering $M^0 > 0$ and applying the following Young's inequality

$$\forall x, y \in \mathbb{R}, |xy| \leq \frac{x^2}{2\gamma} + \frac{\gamma y^2}{2}$$

for a sufficiently small $\gamma > 0$, we obtain the existence of two constants $\alpha_{\varepsilon, K} > 0$ and $\beta_{\varepsilon, K} > 0$ (that depend on ε and K) such that

$$\forall v \in V_K(D), a(v, v) \geq \alpha_{\varepsilon, K} \|v\|_{H^{K+1}(D)}^2 - \beta_{\varepsilon, K} \|v\|_{H^K(D)}^2. \quad (4.15)$$

Furthermore, (4.7) together with the proof of proposition 11 allow to rewrite $a(v, v)$ as

$$a(v, v) = \int_D \int_Y \left| (\varepsilon^{-1} \nabla_y + \nabla_x) \left(\sum_{k=0}^K \varepsilon^k N^k(y) \cdot \nabla^k v(x) \right) \right|^2 dx. \quad (4.16)$$

Then, $\int_D \int_Y u(x, y)^2 dy dx \geq \int_D |\int_Y u(x, y) dy|^2 dx$ and point 1. of proposition 8 imply the the following inequality:

$$\forall v \in V_K, a(v, v) \geq \|\nabla v\|_{L^2(D, \mathbb{R}^d)}^2. \quad (4.17)$$

We now prove that (4.15) and (4.17) together imply the coercivity of a on the space V_K , that is we claim there exists a constant $c_{\varepsilon, K} > 0$ such that

$$\forall v \in V_K, a(v, v) \geq c_{\varepsilon, K} \|v\|_{H^{K+1}(D)}^2. \quad (4.18)$$

Assume the contrary is true, then one can find a sequence (v_n) of functions satisfying $\|v_n\|_{H^{K+1}(D)} = 1$ and such that $a(v_n, v_n) \rightarrow 0$. Up to extracting a relevant subsequence, we may assume that $v_n \rightharpoonup v$ weakly in $H^{K+1}(D)$ and $v_n \rightarrow v$ strongly in $H^K(D)$. Then the polarization identity together with (4.15) and the positivity of a allow to show that (v_n) is a Cauchy sequence in V_K :

$$\begin{aligned} \forall p, q \in \mathbb{N}, \alpha_{\varepsilon, K} \|v_p - v_q\|_{H^{K+1}(D)}^2 &\leq a(v_p - v_q, v_p - v_q) + \beta_{\varepsilon, K} \|v_p - v_q\|_{H^K(D)}^2 \\ &= 2a(v_p, v_p) + 2a(v_q, v_q) - a(v_p + v_q, v_p + v_q) + \beta_{\varepsilon, K} \|v_p - v_q\|_{H^K(D)}^2 \\ &\leq 2a(v_p, v_p) + 2a(v_q, v_q) + \beta_{\varepsilon, K} \|v_p - v_q\|_{H^K(D)}^2 \xrightarrow{p, q \rightarrow \infty} 0. \end{aligned}$$

Therefore $v_n \rightarrow v$ strongly in V_K . Using the continuity of a , we infer then $a(v, v) = \lim_{n \rightarrow +\infty} a(v_n, v_n) = 0$. The property (4.17) yields then that v is a constant. Therefore, $0 = a(v, v) = \varepsilon^{-2} M^0 \|v\|_{L^2(D)}^2$, which implies $v = 0$. This is in contradiction with the fact that $\|v_n\|_{H^{K+1}(D)} = 1$ for any $n \geq 0$ and the strong convergence of (v_n) in $H^{K+1}(D)$.

Finally, the coercivity (4.18) and the continuity of a and b over V_K ensure that all the assumptions of the Lax–Milgram theorem are fulfilled, which yields existence and uniqueness to the problem (4.14). \square

Remark 4. It is always possible to add to \mathbb{D}_K^{2K+2} a small perturbation making (4.13) satisfied while keeping an “admissible” higher order homogenized equation. Indeed, since the other $2K + 1$ coefficients are kept unaffected, the error estimate provided by proposition 10 and corollary 5 below remain valid. Let us remark, however, that this non-degeneracy condition is automatically fulfilled for any shape of obstacle ηT when $K = 0$ because it is easily shown that $\mathbb{D}_0^2 = -(M^0)^2 \int_Y |\mathcal{X}^0(y)|^2 dy > 0$ (see (4.20)). In the general case $K \geq 1$, it could fail to be satisfied for particular obstacle shapes (e.g. in case of invariance of ηT along some of the cell axes).

4.3. Asymptotic surprise: $\mathbb{D}_K^{2k} = M^{2k}$; error estimates for the homogenized model of order $2K + 2$

We terminate this section by verifying the assumptions of proposition 10 which ensure the validity of the error estimate (1.14) claimed in the introduction. The next proposition establishes the point 1.

Lemma 5. *Assume the non-degeneracy condition (4.13). The solution $v_{\varepsilon, K}^*$ of (4.10) belongs to $\mathcal{C}^\infty(D)$ and for any $m \in \mathbb{N}$, there exists a constant $C_{m, K}$ that does not depend on ε such that*

$$\|v_{\varepsilon, K}^*\|_{H^m(D)} \leq C_{m, K} \|f\|_{H^m(D)} \varepsilon^2. \quad (4.19)$$

Proof. We solve (4.10) explicitly with Fourier expansions in the periodic domain $D = [0, L]^d$. Let $\widehat{f}(\xi)$ be the Fourier coefficients of f and $c(\xi, \varepsilon)$ the symbol of the differential operator of (4.10), namely:

$$\forall \xi \in \mathbb{Z}^d, \forall \varepsilon > 0, c(\xi, \varepsilon) = \sum_{k=0}^{K+1} (-1)^k |2\pi/L|^{2k} \varepsilon^{2k-2} \mathbb{D}_K^{2k} \cdot \xi^{2k},$$

where $\xi^0 = 1$ by convention and where we have used the short-hand notation $\mathbb{D}_K^{2k} \cdot \xi^{2k} := \mathbb{D}_{i_1 \dots i_{2k}}^{2k} \xi_{i_1 \dots i_{2k}}$. For $\xi \in \mathbb{Z}^d$, the Fourier coefficients $\widehat{v}_{\varepsilon, K}^*$ of v_K^* read:

$$\widehat{v}_{\varepsilon, K}^*(\xi) = \frac{\widehat{f}(\xi)}{c(\xi, \varepsilon)}.$$

Applying Parseval’s identity to the bilinear form (4.16), we obtain

$$\begin{aligned} c(\xi, \varepsilon) &= \int_Y \left| (\varepsilon^{-1} \nabla_y + (2i\pi/L)\xi) \left(\sum_{k=0}^K (2i\pi/L)^k \varepsilon^k N^k(y) \cdot \xi^k \right) \right|^2 dy \\ &= \varepsilon^{-2} \Phi(\varepsilon \xi). \end{aligned}$$

where Φ is the function defined by:

$$\begin{aligned} \Phi : \mathbb{R}^d &\longrightarrow \mathbb{R} \\ \lambda &\longmapsto \int_Y \left| (\nabla_y + (2i\pi/L)\lambda) \left(\sum_{k=0}^K (2i\pi/L)^k N^k(y) \cdot \lambda^k \right) \right|^2 dy. \end{aligned}$$

In order to obtain (4.19), it is enough to prove the existence of some positive constant $C_K > 0$ such that $\Phi(\lambda) \geq C_K$ for any $\lambda \in \mathbb{R}^d$. The function Φ is clearly continuous, nonnegative and satisfies $\Phi(\lambda) \rightarrow +\infty$ when $|\lambda| \rightarrow +\infty$ because of (4.13). Therefore it admits a minimum point $\lambda_0 \in (\mathbb{R}_+)^d$. Let us assume that $\Phi(\lambda_0) = 0$. Using the Cauchy-Schwartz inequality, we obtain

$$0 = \Phi(\lambda_0) \geq \left| \int_Y (\nabla_y + (2i\pi/L)\lambda_0) \left(\sum_{k=0}^K (2i\pi/L)^k N^k(y) \cdot \lambda_0^k \right) dy \right|^2 = \frac{4\pi^2}{L^2} |\lambda_0|^2.$$

Therefore it must hold $\lambda_0 = 0$, however this is not possible because

$$\Phi(0) = \int_Y |\nabla_y N^0(y)|^2 dy = M^0 > 0.$$

Consequently, $\forall \lambda \in \mathbb{R}^d$, $\Phi(\lambda) \geq \Phi(\lambda_0) > 0$ which concludes the proof. \square

The point 2 of proposition 10 is the object of the next result. More precisely, we prove that surprisingly, the coefficients \mathbb{D}_K^{2k} and M^{2k} coincide for $0 \leq 2k \leq 2K$. A similar fact was also observed for the (scalar) antiplane elasticity model considered in [52], but only for the first half $0 \leq 2k \leq K$ of the coefficients (and without providing a proof).

Proposition 13. *All the coefficients of the homogenized equation (1.6) of order $2K + 2$ (defined in (4.10)) coincide with those of the formal infinite order homogenized equation (1.9), except the leading order one:*

$$\begin{aligned} \mathbb{D}_K^{2k} &= M^{2k} \text{ for any } 0 \leq 2k \leq 2K, \\ \mathbb{D}_K^{2K+2} &= (-1)^{K+1} \int_Y N^K \otimes N^K \otimes Idy. \end{aligned} \quad (4.20)$$

Proof. First of all, (4.20) is only a rewriting of (4.11). We show the following, slightly more general, result:

$$\forall 0 \leq k \leq 2K, M^k = \sum_{l=0}^k (-1)^l \mathbb{B}_K^{l, k-l}, \quad (4.21)$$

which is enough for our purpose because of (4.11). We distinguish two cases.

1. *Case* $0 \leq k \leq K$. For $0 \leq k, l \leq K$, the coefficient $\mathbb{B}_K^{l, k-l}$ is given by (from (4.6))

$$\mathbb{B}_K^{l, k-l} = \int_Y (\partial_j N^l + N^{l-1} \otimes e_j) \otimes (\partial_j N^{k-l} + N^{k-l-1} \otimes e_j) dy, \quad (4.22)$$

where we use the convention $N^{-1} = N^{-2} = 0$. After an integration by parts, we rewrite $\mathbb{B}_K^{l, k-l}$ as follows:

$$\begin{aligned} \mathbb{B}_K^{l, k-l} &= \int_Y (-\Delta N^l - 2\partial_j N^{l-1} \otimes e_j - N^{l-2} \otimes I) \otimes N^{k-l} dy \\ &\quad + \int_Y (\partial_j N^l \otimes N^{k-l-1} \otimes e_j + N^{l-1} \otimes N^{k-l-1} \otimes I) dy \\ &\quad + \int_Y (\partial_j N^{l-1} \otimes N^{k-l} \otimes e_j + N^{l-2} \otimes N^{k-l} \otimes I) dy \\ &= \int_Y (M^l \otimes N^{k-l}) dy + B^{k, l} + B^{k, l-1} \end{aligned} \quad (4.23)$$

where $B^{k, l}$ is the k -th order tensor defined by

$$B^{k, l} := \int_Y (\partial_j N^l \otimes N^{k-l-1} \otimes e_j + N^{l-1} \otimes N^{k-l-1} \otimes I) dy.$$

Using now the point 1. of proposition 8 and recognizing a telescopic series, we obtain

$$\begin{aligned} \sum_{l=0}^k (-1)^l \mathbb{B}_K^{l, k-l} &= (-1)^k M^k + \sum_{l=0}^k ((-1)^l B^{k, l} - (-1)^{l-1} B^{k, l-1}) \\ &= (-1)^k M^k + (-1)^k B^{k, k} - (-1)^{-1} B^{k, -1}. \end{aligned}$$

This implies (4.21) by using the facts that $M^k = 0$ when k is odd and $B^{k,k} = B^{k,-1} = 0$ owing to our convention $N^{-1} = 0$.

2. *Case $K + 1 \leq k \leq 2K$.* The equality (4.22) is valid for any $K + 1 \leq k \leq 2K$ and $0 \leq l \leq k$ if we assume by convention (*in this part only*) that $N^m = 0$ whenever $m > K$, because of the definition (4.7). Then (4.23) remains true provided $0 \leq l \leq K$. Therefore, recognizing the same telescopic series, we are able to compute

$$\begin{aligned} \sum_{l=0}^K (-1)^l \mathbb{B}_K^{l,k-l} &= (-1)^K B^{k,K} \\ &= (-1)^K \int_Y (\partial_j N^K \otimes N^{k-K-1} \otimes e_j + N^{K-1} \otimes N^{k-K-1} \otimes I) dy. \end{aligned}$$

Remembering that $\mathbb{B}_K^{l,m} = 0$ whenever $l > K + 1$ or $m > K + 1$, we eventually obtain

$$\begin{aligned} \sum_{l=0}^k (-1)^l \mathbb{B}_K^{l,k-l} &= \sum_{l=0}^K (-1)^l \mathbb{B}_K^{l,k-l} + (-1)^{K+1} \mathbb{B}_K^{K+1,k-K-1} \\ &= (-1)^K \int_Y (\partial_j N^K \otimes N^{k-K-1} \otimes e_j + N^{K-1} \otimes N^{k-K-1} \otimes I) dy \\ &\quad + (-1)^{K+1} \int_Y (N^K \otimes e_j) \otimes (\partial_j N^{k-K-1} + N^{k-K-2} \otimes e_j) dy \\ &= (-1)^K \int_Y ((2\partial_j N^K \otimes e_j + N^{K-1} \otimes I) \otimes N^{k-K-1} - N^K \otimes N^{k-K-2} \otimes I) dy \\ &= (-1)^K \int_Y ((-\Delta_{yy} N^{K+1} - M^{K+1}) \otimes N^{k-K-1} - N^K \otimes N^{k-K-2} \otimes I) dy, \end{aligned} \tag{4.24}$$

where we have used (3.16) in the last equality. We now consider two cases:

- if $k = K + 1$, then the above expression reads

$$\begin{aligned} \sum_{l=0}^{K+1} (-1)^l \mathbb{B}_K^{l,K+1-l} &= (-1)^K \int_Y ((-\Delta_{yy} N^{K+1} - M^{K+1}) \otimes N^0) dy \\ &= (-1)^{K+1} M^{K+1} + (-1)^K \int_Y N^{K+1} \otimes M^0 dy \\ &= (-1)^{K+1} M^{K+1} = M^{K+1}, \end{aligned}$$

where the last equality is a consequence of [corollary 1](#).

- if $K + 2 \leq k \leq 2K$, then (4.24) coincides with M^k by using (3.18) (with $p = K + 1$).

□

Remark 5. As is highlighted by the proof [proposition 13](#), the “surprising” fact that $\mathbb{D}_K^{2k} = M^{2k}$ even for $K + 1 \leq 2k \leq 2K$ is partly due to the fact that for any $p \in \mathbb{N}$, the tensor M^{2p} can be computed from the first p homogenized tensors $(\mathcal{X}^i(y))_{0 \leq i \leq p}$ only (point 4. of [proposition 8](#)).

Since we have verified that all the requirements of [proposition 10](#) are satisfied with $v_\varepsilon^* \equiv v_{\varepsilon,K}^*$ and $L \equiv 2K + 1$ (remember $M^{2K+1} = 0$ from [corollary 1](#)), we are in position to state our main result.

Corollary 5. *The error estimate (1.14) holds for the reconstructed solution $W_{\varepsilon,2K+1}(v_{\varepsilon,K}^*)$ where $v_{\varepsilon,K}^*$ is the solution of (4.10).*

5. RETRIEVING THE CLASSICAL REGIMES: LOW VOLUME FRACTION LIMITS WHEN THE SIZE OF THE OBSTACLES TENDS TO ZERO

The goal of this section is to show that our higher order homogenized models (1.6) have the potential of being valid in any regime of size of holes. For this purpose, we obtain asymptotics for the tensors \mathcal{X}^{k*} and M^k in the low volume fraction limit when the scaling η of the obstacle vanishes to zero: $\eta \rightarrow 0$. Our main results are stated in [corollaries 6](#) and [7](#); they imply that both the infinite order homogenized equation (3.11) as well as the effective model (1.6) of order $2K + 2$ converge coefficient-wise to either of the three classical

regimes of the literature (namely, to the original Laplace equation (1.1), or to the analogue of the Brinkman or Darcy equation (1.3) and (1.4) if $K = 0$, and to either of (1.3) or (1.4) for $K \geq 1$ if η remains greater or comparable to the critical size $\eta_{\text{crit}} \sim \eta^{2/(d-2)}$).

In this whole subsection, it is assumed, for simplicity, that the space dimension is greater than 3:

$$d \geq 3.$$

We do not consider the case $d = 2$ which requires a specific treatment, although very similar results could be stated (see e.g. [4]).

In all what follows, the hole ηT is assumed to be strictly included in the unit cell for any $\eta \leq 1$ (it does not touch the boundary): $\eta T \subset\subset P$. Functions of the rescaled cell $\eta^{-1}P$ are indicated by a tilde $\tilde{\cdot}$ notation. For a given function $\tilde{v} \in L^2(\eta^{-1}P)$, we denote by $\langle \tilde{v} \rangle$ the average $\langle \tilde{v} \rangle := \eta^d \int_{\eta^{-1}P} \tilde{v}(y) dy$. With a small abuse of notation, when $\tilde{v} \in L^2(\eta^{-1}P \setminus T)$, we still denote by $\langle \tilde{v} \rangle$ this quantity where we implicitly extend \tilde{v} by 0 within T .

Let us recall that for any $v \in H^1(P \setminus (\eta T))$, if \tilde{v} is the rescaled function defined by $\tilde{v}(y) := v(\eta y)$ in the rescaled cell $\eta^{-1}P \setminus T$, then the L^2 norms of v and \tilde{v} and their gradients are related by the following identities:

$$\|v\|_{L^2(P \setminus (\eta T))} = \eta^{d/2} \|\tilde{v}\|_{L^2(\eta^{-1}P \setminus T)},$$

$$\|\nabla v\|_{L^2(P \setminus (\eta T), \mathbb{R}^d)} = \eta^{d/2-1} \|\nabla \tilde{v}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)}.$$

We follow the methodology of [4] (and also [37]), which relies extensively on the use of the next two lemmas:

Lemma 6. *Assume $d \geq 3$. There exists a constant $C > 0$ independent of $\eta > 0$ such that for any $\tilde{v} \in H^1(\eta^{-1}P \setminus T)$ which vanishes on the hole ∂T and which is $\eta^{-1}P$ periodic, the following inequalities hold:*

$$\|\tilde{v}\|_{L^2(\eta^{-1}P \setminus T)} \leq C \eta^{-d/2} \|\nabla \tilde{v}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)}, \quad (5.1)$$

$$|\langle \tilde{v} \rangle| \leq C \|\nabla \tilde{v}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)}, \quad (5.2)$$

$$\|\tilde{v} - \langle \tilde{v} \rangle\|_{L^2(\eta^{-1}P \setminus T)} \leq C \eta^{-1} \|\nabla \tilde{v}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)}, \quad (5.3)$$

$$\|\tilde{v} - \langle \tilde{v} \rangle\|_{L^{2d/(d-2)}(\eta^{-1}P \setminus T)} \leq C \|\nabla \tilde{v}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)}. \quad (5.4)$$

Proof. See [4, 38]. □

Lemma 7. *Consider $\tilde{h} \in L^2(\eta^{-1}P \setminus T)$ and let $\tilde{v} \in H^1(\eta^{-1}P \setminus T)$ be the unique solution to the Poisson problem:*

$$\begin{cases} -\Delta \tilde{v} = \tilde{h} & \text{in } \eta^{-1}P \setminus T \\ \tilde{v} = 0 & \text{on } \partial T \\ \tilde{v} & \text{is } \eta^{-1}P \setminus T\text{-periodic.} \end{cases} \quad (5.5)$$

There exists a constant $C > 0$ independent of η and \tilde{h} such that

$$\|\nabla \tilde{v}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} \leq C(\eta^{-1} \|\tilde{h} - \langle \tilde{h} \rangle\|_{L^2(\eta^{-1}P \setminus T)} + \eta^{-d} |\langle \tilde{h} \rangle|).$$

Proof. An integration by parts and the use of lemma 6 yields

$$\begin{aligned} \|\nabla \tilde{v}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} &= \int_{\eta^{-1}P \setminus T} \tilde{h} \tilde{v} dy \\ &= \int_{\eta^{-1}P \setminus T} (\tilde{h} - \langle \tilde{h} \rangle)(\tilde{v} - \langle \tilde{v} \rangle) dy + \int_{\eta^{-1}P \setminus T} \langle \tilde{h} \rangle \langle \tilde{v} \rangle dy \\ &\leq C(\|\tilde{h} - \langle \tilde{h} \rangle\|_{L^2(\eta^{-1}P \setminus T)} \|\tilde{v} - \langle \tilde{v} \rangle\|_{L^2(\eta^{-1}P \setminus T)} + \eta^{-d} |\langle \tilde{h} \rangle| |\langle \tilde{v} \rangle|) \\ &\leq C(\eta^{-1} \|\tilde{h} - \langle \tilde{h} \rangle\|_{L^2(\eta^{-1}P \setminus T)} + \eta^{-d} |\langle \tilde{h} \rangle|) \|\nabla \tilde{v}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)}. \end{aligned}$$

□

We also need to consider the so-called Deny-Lions (or Beppo-Levi) space $\mathcal{D}^{1,2}(\mathbb{R}^d \setminus T)$ whose definition is recalled below (the reader is referred to [4, 3, 13] and also [44], p.59. for more details).

Definition 6 (Deny-Lions space). The Deny-Lions space $\mathcal{D}^{1,2}(\mathbb{R}^d \setminus T)$ is the completion of the space of smooth functions by the L^2 norm of their gradients:

$$\mathcal{D}^{1,2}(\mathbb{R}^d \setminus T) := \overline{\mathcal{D}(\mathbb{R}^d \setminus T)}^{\|\nabla \cdot\|_{L^2(\mathbb{R}^d \setminus T, \mathbb{R}^d)}}.$$

When $d \geq 3$, it admits the following characterization:

$$\begin{aligned} \mathcal{D}^{1,2}(\mathbb{R}^d \setminus T) \\ = \{ \phi \text{ measurable} \mid \|\phi\|_{L^{2d/(d-2)}(\mathbb{R}^d \setminus T)} < +\infty \text{ and } \|\nabla \phi\|_{L^2(\mathbb{R}^d \setminus T, \mathbb{R}^d)} < +\infty \}. \end{aligned}$$

We introduce Ψ the unique solution to the exterior problem

$$\begin{cases} -\Delta \Psi = 0 \text{ in } \mathbb{R}^d - T \\ \Psi = 0 \text{ on } \partial T \\ \Psi \rightarrow 1 \text{ at } \infty, \end{cases} \quad (5.6)$$

and we denote by F the normal flux

$$F := \int_{\mathbb{R}^d \setminus T} |\nabla \Psi|^2 dx = - \int_{\partial T} \nabla \Psi \cdot \mathbf{n} ds, \quad (5.7)$$

where \mathbf{n} is the normal pointing *inward* T . The condition $\Psi \rightarrow 1$ at ∞ is to be understood in the sense that $\Psi - 1 \in \mathcal{D}^{1,2}(\mathbb{R}^d \setminus T)$.

The following result provides asymptotics for the tensors \mathcal{X}^k and their averages \mathcal{X}^{k*} . It extends Theorem 3.1 of [4] (see also [37]) where the special case $k = 0$ was obtained for the Stokes system.

Proposition 14. *Assume $d \geq 3$. For any $k \geq 0$, denote by $\tilde{\mathcal{X}}^{2k}$ and $\tilde{\mathcal{X}}^{2k+1}$ the rescaled tensors for any $x \in \eta^{-1}P \setminus T$ by:*

$$\tilde{\mathcal{X}}^{2k}(x) := \eta^{(d-2)(k+1)} \mathcal{X}^{2k}(\eta x) \text{ and } \tilde{\mathcal{X}}^{2k+1}(x) := \eta^{(d-2)(k+1)} \mathcal{X}^{2k+1}(\eta x).$$

Then:

(1) *there exists a constant $C > 0$ independent of $\eta > 0$ such that:*

$$\forall \eta > 0, \|\nabla \tilde{\mathcal{X}}^{2k}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} \leq C \text{ and } \|\nabla \tilde{\mathcal{X}}^{2k+1}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} \leq C; \quad (5.8)$$

(2) *the following convergences hold as $\eta \rightarrow 0$:*

$$\tilde{\mathcal{X}}^{2k} \rightharpoonup \frac{\Psi}{F^{k+1}} J^{2k}, \text{ weakly in } H_{loc}^1(\mathbb{R}^d \setminus T) \quad (5.9)$$

$$\tilde{\mathcal{X}}^{2k+1} \rightharpoonup 0 \text{ weakly in } H_{loc}^1(\mathbb{R}^d \setminus T) \quad (5.10)$$

$$\mathcal{X}^{2k*} \sim \frac{1}{\eta^{(d-2)(k+1)} F^{k+1}} J^{2k}, \quad (5.11)$$

where we recall $J^{2k} := \overbrace{I \otimes I \otimes \dots \otimes I}^{k \text{ times}}$ (definition (2.4)).

Remark 6. Let us recall that we already know $\mathcal{X}^{2k+1*} = 0$ for any $k \in \mathbb{N}$ (proposition 3).

Proof. The result is proved by induction.

(1) *Case $2k$ with $k = 0$.* The tensor $\tilde{\mathcal{X}}^0$ satisfies

$$-\Delta \tilde{\mathcal{X}}^0 = \eta^d \text{ in } \eta^{-1}P \setminus T \quad (5.12)$$

as well as the other boundary conditions of (5.5), hence lemma 7 yields

$$\|\nabla \tilde{\mathcal{X}}^0\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} \leq C \eta^{-d} \eta^d \langle 1 \rangle \leq C.$$

From (5.2), the average $\langle \tilde{\mathcal{X}}^0 \rangle$ is bounded. Therefore, there exists a constant $c^0 \in \mathbb{R}$ and a function $\hat{\Psi}^0$ such that, up to extracting a subsequence,

$$\langle \tilde{\mathcal{X}}^0 \rangle \rightarrow c^0 \text{ and } \tilde{\mathcal{X}}^0 \rightharpoonup \hat{\Psi}^0 \text{ weakly in } H_{loc}^1(\mathbb{R}^d \setminus T) \text{ when } \eta \rightarrow 0.$$

Furthermore, the lower semi-continuity of the $H_{loc}^1(\mathbb{R}^d \setminus T)$ norm and (5.4) imply that $\widehat{\Psi}^0 - c^0$ belongs to $\mathcal{D}^{1,2}(\mathbb{R}^d \setminus T)$ (see [4] for a detailed justification). Multiplying (5.12) by a compactly supported test function $\Phi \in C_c^\infty(\mathbb{R}^d \setminus \overline{T})$ and integrating by part yields

$$\int_{\eta^{-1}P \setminus T} \nabla \tilde{\mathcal{X}}^0 \cdot \nabla \Phi dy = \eta^d \int_{\eta^{-1}P \setminus T} \Phi dy.$$

Passing to the limit when $\eta \rightarrow 0$ entails that $\widehat{\Psi}^0$ is solution to the exterior problem

$$\begin{cases} -\Delta \widehat{\Psi}^0 = 0 \text{ in } \mathbb{R}^d \setminus T \\ \widehat{\Psi}^0 = 0 \text{ on } \partial T \\ \widehat{\Psi}^0 \rightarrow c^0 \text{ at } \infty. \end{cases} \quad (5.13)$$

By linearity, the unique solution to this problem is given by $\widehat{\Psi} = c^0 \Psi$ where Ψ is the solution to (5.6). Finally, the constant c^0 can be identified by integrating (5.12) against the constant test function $\Phi = 1$, which implies

$$c^0 F = - \int_{\partial T} \nabla(c^0 \Psi^0) \cdot \mathbf{n} ds = \lim_{\eta \rightarrow 0} - \int_{\partial T} \nabla \tilde{\mathcal{X}}^0 \cdot \mathbf{n} ds = 1.$$

Therefore, $c^0 = 1/F$ from where (5.9) follows for $k = 0$. Since the obtained limit is unique, the convergence holds for the whole sequence. Then (5.11) follows from a simple change of variable.

(2) *Case $2k + 1$ with $k = 0$.* A simple computation yields

$$-\Delta \tilde{\mathcal{X}}^1 = 2\eta \partial_j \tilde{\mathcal{X}}^0 \otimes e_j. \quad (5.14)$$

Applying lemma 7 and remarking that $\langle 2\eta \partial_j \tilde{\mathcal{X}}^0 \rangle = 0$, we obtain

$$\|\nabla \tilde{\mathcal{X}}^1\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} \leq C \eta^{-1} \eta \|\nabla \tilde{\mathcal{X}}^0\| \leq C'.$$

Integrating (5.14) against a compactly supported test function $\Phi \in C_c^\infty(\mathbb{R}^d \setminus T)$ and passing to the limit as $\eta \rightarrow 0$, we obtain with similar arguments the existence of a constant tensor c^1 (of order 1) such that, up to the extraction of a subsequence,

$$\tilde{\mathcal{X}}^1 \rightharpoonup c^1 \Psi \text{ weakly in } H_{loc}^1(\mathbb{R}^d \setminus T) \text{ and } \langle \tilde{\mathcal{X}}^1 \rangle \rightarrow c^1.$$

From proposition 3, it holds that $\langle \tilde{\mathcal{X}}^1 \rangle = 0$ which implies $c^1 = 0$, hence (5.10) for $k = 0$.

(3) *General case.* We now complete the proof by induction on k . Assuming the result holds till rank $k \geq 0$, we compute

$$\begin{aligned} -\Delta \tilde{\mathcal{X}}^{2k+2} &= 2\eta^{d-1} \partial_j \tilde{\mathcal{X}}^{2k+1} \otimes e_j + \eta^d \tilde{\mathcal{X}}^{2k} \otimes I, \\ -\Delta \tilde{\mathcal{X}}^{2k+3} &= 2\eta \partial_j \tilde{\mathcal{X}}^{2k+2} \otimes e_j + \eta^d \tilde{\mathcal{X}}^{2k+1} \otimes I. \end{aligned}$$

Applying lemmas 6 and 7 and (5.8) at rank k , we obtain

$$\begin{aligned} &\|\nabla \tilde{\mathcal{X}}^{2k+2}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} \\ &\leq C(\eta^{d-2} \|\nabla \tilde{\mathcal{X}}^{2k+1}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} + \eta^{d-1} \|\tilde{\mathcal{X}}^{2k} - \langle \tilde{\mathcal{X}}^{2k} \rangle\|_{L^2(\eta^{-1}P \setminus T)} \\ &\quad + \|\langle \tilde{\mathcal{X}}^{2k} \rangle\|_{L^2(\eta^{-1}P \setminus T)}) \\ &\leq C(\eta^{d-2} \|\nabla \tilde{\mathcal{X}}^{2k+1}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} + \eta^{d-2} \|\nabla \tilde{\mathcal{X}}^{2k}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} \\ &\quad + \|\nabla \tilde{\mathcal{X}}^{2k}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)}) \leq C', \end{aligned} \quad (5.15)$$

$$\begin{aligned} &\|\nabla \tilde{\mathcal{X}}^{2k+3}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)}^2 \\ &\leq C(\|\nabla \tilde{\mathcal{X}}^{2k+2}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} + \eta^{d-1} \|\tilde{\mathcal{X}}^{2k+1} - \langle \tilde{\mathcal{X}}^{2k+1} \rangle\|_{L^2(\eta^{-1}P \setminus T)} \\ &\quad + \|\langle \tilde{\mathcal{X}}^{2k+1} \rangle\|) \\ &\leq C(\|\nabla \tilde{\mathcal{X}}^{2k+2}\|_{L^2(\eta^{-1}P \setminus T, \mathbb{R}^d)} + \eta^{d-2} \|\nabla \tilde{\mathcal{X}}^{2k+1}\|) \leq C'. \end{aligned}$$

This implies (5.8) at rank $k+1$. Using similar arguments as previously, we infer the existence of constant tensors c^{2k+2} and c^{2k+3} such that

$$\begin{aligned}\tilde{\mathcal{X}}^{2k+2} &\rightharpoonup c^{2k+2}\Psi \text{ weakly in } H_{loc}^1(\mathbb{R}^d \setminus T) \text{ and } \langle \tilde{\mathcal{X}}^{2k+2} \rangle \rightarrow c^{2k+2} \\ \tilde{\mathcal{X}}^{2k+3} &\rightharpoonup c^{2k+3}\Psi \text{ weakly in } H_{loc}^1(\mathbb{R}^d \setminus T) \text{ and } \langle \tilde{\mathcal{X}}^{2k+3} \rangle \rightarrow c^{2k+3}.\end{aligned}$$

Since we know from proposition 3 that $\langle \tilde{\mathcal{X}}^{2k+3} \rangle = 0$, we obtain $c^{2k+3} = 0$ which yields (5.10) at rank $k+1$. Finally, we integrate (5.15) by parts against the test function $\Phi = 1$ in order to identify c^{2k+2} :

$$\begin{aligned}c^{2k+2}F &= - \int_{\partial T} \nabla(c^{2k+2}\Psi) \cdot \mathbf{n} ds = \lim_{\eta \rightarrow 0} - \int_{\partial T} \nabla \tilde{\mathcal{X}}^{2k+2} \cdot \mathbf{n} ds \\ &= \lim_{\eta \rightarrow 0} \langle \tilde{\mathcal{X}}^{2k} \rangle \otimes I = c^{2k} \otimes I.\end{aligned}$$

This implies $c^{2k+2} = J^{2k+2}/F^{k+2}$ and $c^{2k+3} = 0$, which concludes the proof. \square

Remark 7. The convergence (5.10) seems to indicate that we may have not found the optimal scaling for the odd order tensors \mathcal{X}^{2k+1} since we are able to identify only a zero weak limit in (5.10). A more elaborate analysis could be carried on by different techniques, see e.g. the use of layer potentials in [37].

We are now able to identify the asymptotic behavior of the constant tensors M^k . Recall already know that $M^{2k+1} = 0$ from corollary 1.

Corollary 6. *Assume $d \geq 3$. The following convergences hold for the tensors M^k as $\eta \rightarrow 0$:*

$$M^0 \sim \eta^{d-2}F, \quad (5.16)$$

$$M^2 \rightarrow -I, \quad (5.17)$$

$$\forall k > 1, M^{2k} = o\left(\frac{1}{\eta^{(d-2)(k-1)}}\right). \quad (5.18)$$

Proof. We replace the asymptotics of proposition 14 in the explicit formula (3.12) for the tensor M^k . (5.16) is a consequence of the definition $M^0 = (\mathcal{X}^{0*})^{-1}$. (5.17) is obtained by writing

$$M^2 = -((\mathcal{X}^{0*})^{-1})^2 \mathcal{X}^{2*} \sim -\frac{\eta^{2(d-2)}F^2}{\eta^{2(d-2)}F^2}I = -I.$$

Let us now prove (5.18). By eliminating terms of odd orders in (3.12), we may write for any $k \geq 1$,

$$\begin{aligned}M^{2k} &= \sum_{p=1}^{2k} \frac{(-1)^p}{(\mathcal{X}^{0*})^{p+1}} \sum_{\substack{i_1+\dots+i_p=k \\ 1 \leq i_1 \dots i_p \leq k}} \mathcal{X}^{2i_1*} \otimes \dots \otimes \mathcal{X}^{2i_p*} \\ &= \sum_{p=1}^{2k} (-1)^p \eta^{(p+1)(d-2)} F^{p+1} \sum_{\substack{i_1+\dots+i_p=k \\ 1 \leq i_1 \dots i_p \leq k}} \frac{J^{2i_1}}{\eta^{(d-2)(i_1+1)} F^{i_1+1}} \otimes \dots \otimes \frac{J^{2i_p}}{\eta^{(d-2)(i_p+1)} F^{i_p+1}} \\ &\quad + o\left(\frac{1}{\eta^{(k-1)(d-2)}}\right) \\ &= \frac{J^{2k}}{\eta^{(k-1)(d-2)} F^{k-1}} \left(\sum_{p=1}^{2k} (-1)^p \sum_{\substack{i_1+\dots+i_p=k \\ 1 \leq i_1 \dots i_p \leq k}} 1 \right) + o\left(\frac{1}{\eta^{(k-1)(d-2)}}\right).\end{aligned}$$

Then (5.18) results from the last summation being zero:

$$\forall k > 1, \sum_{p=1}^k (-1)^p \sum_{\substack{i_1+\dots+i_p=k \\ 1 \leq i_1 \dots i_p \leq k}} 1 = 0. \quad (5.19)$$

There are several ways to obtain the latter formula. A rather direct argument in the spirit of the proof of [proposition 6](#) is to apply the identity [\(3.13\)](#) to the power series $1/(1-z) = \sum_{k \in \mathbb{N}} z^k$ which yields

$$1 - z = 1 + \sum_{k=1}^{+\infty} \left(\sum_{p=1}^k (-1)^p \sum_{\substack{i_1 + \dots + i_p = k \\ 1 \leq i_1 \dots i_p \leq k}} 1 \right) z^k,$$

from where [\(5.19\)](#) follows by identifying the powers in z^k . \square

Remark 8. We retrieve formally in [corollary 6](#) the different classical asymptotic regimes [\(1.3\)](#) to [\(1.5\)](#) for the perforated problem [\(1.1\)](#) (at least for $K = 0$), because the coefficients of [\(1.9\)](#) read, as $\eta \rightarrow 0$:

$$\begin{aligned} \varepsilon^{-2} M^0 &\sim \frac{\eta^{d-2}}{\varepsilon^2} F, \\ \varepsilon^0 M^2 &\rightarrow -I, \\ \varepsilon^{2k-2} M^{2k} &= o\left(\left(\frac{\varepsilon^2}{\eta^{d-2}}\right)^{k-1}\right) \text{ for } k \geq 1. \end{aligned}$$

These asymptotics bring into play the ratio ε^2/η^{d-2} and so the critical scaling $\eta \sim \varepsilon^{2/(d-2)}$ corresponding to the ‘‘Brinkman’’ regime [\(1.3\)](#) (which implies $\varepsilon^{-2} M^0 \rightarrow F$ and $\varepsilon^{2k+2} M^{2k} = o(1)$). The Darcy regimes [\(1.4\)](#) and [\(1.5\)](#) correspond to the situation where $\eta^{d-2}/\varepsilon^2 \rightarrow +\infty$; in that case the zeroth order term $\varepsilon^{-2} M^0$ is dominant. The Stokes regime [\(1.2\)](#) is found for $K = 0$ and $\eta = o(\varepsilon^{2/(d-2)})$ (which implies $\varepsilon^{-2} M^0 \rightarrow 0$). Note that $\varepsilon^0 M^2 \rightarrow -I$ whatever the scaling $\eta \rightarrow 0$. Unfortunately, the inverse of the critical ratio ε^2/η^{d-2} could blow up as η vanishes at the rate $\eta = o(\varepsilon^{2/(d-2)})$: a more elaborate analysis shall be performed in future works to determine whether $\varepsilon^{2k-2} M^{2k}$ with $k > 1$ still converges to zero in this regime.

We conclude this paper with the statement that the regimes [\(1.4\)](#) and [\(1.5\)](#) are also captured in the low volume fraction limit $\eta \rightarrow 0$ by the homogenized equation [\(1.6\)](#) of finite order $2K + 2$. Because of [proposition 13](#), it is enough to establish that \mathbb{D}_K^{2K+2} satisfies the same asymptotics than M^{2K+2} (hence all the coefficients \mathbb{D}_K^{2k} for $0 \leq 2k \leq 2K + 2$ because of [proposition 13](#)). The proof of this result requires the following asymptotic bounds for the tensors N^k :

Proposition 15. *Assume $d \geq 3$. For any $k \geq 0$, denote \tilde{N}^{2k} and \tilde{N}^{2k+1} the rescaled tensors in $\eta^{-1}P \setminus T$ defined by:*

$$\forall y \in \eta^{-1}P \setminus T, \tilde{N}^{2k}(y) := \eta^{(d-2)k} N^{2k}(\eta y) \text{ and } \tilde{N}^{2k+1}(y) := \eta^{(d-2)k} N^{2k+1}(\eta y).$$

Then there exists a constant C independent of $\eta > 0$ such that

$$\forall \eta > 0, \|\nabla \tilde{N}^{2k}\|_{L^2(\eta^{-1}P \setminus T)} \leq C \text{ and } \|\nabla \tilde{N}^{2k+1}\|_{L^2(\eta^{-1}P \setminus T)} \leq C. \quad (5.20)$$

Moreover, the following convergences hold as $\eta \rightarrow 0$:

$$\tilde{N}^0 \rightharpoonup \Psi \text{ weakly in } H_{loc}^1(\mathbb{R}^d \setminus T) \quad (5.21)$$

$$\forall k \geq 1, \tilde{N}^k \rightharpoonup 0 \text{ weakly in } H_{loc}^1(\mathbb{R}^d \setminus T). \quad (5.22)$$

Proof. Using the definition [\(3.15\)](#) of the tensors N^k and eliminating odd order terms, the tensors \tilde{N}^{2k} and \tilde{N}^{2k+1} can be rewritten as:

$$\begin{aligned} \tilde{N}^0 &= \frac{M^0}{\eta^{d-2}} \tilde{\mathcal{X}}^0, \\ \forall k \geq 1, \tilde{N}^{2k} &= \frac{M^0}{\eta^{d-2}} \tilde{\mathcal{X}}^{2k} + M^2 \otimes \tilde{\mathcal{X}}^{2k-2} + \sum_{p=2}^k \eta^{(d-2)(p-1)} M^{2p} \otimes \tilde{\mathcal{X}}^{2(k-p)}, \\ \forall k \geq 0, \tilde{N}^{2k+1} &= \frac{M^0}{\eta^{d-2}} \tilde{\mathcal{X}}^{2k+1} + M^2 \otimes \tilde{\mathcal{X}}^{2k-1} + \sum_{p=2}^k \eta^{(d-2)(p-1)} M^{2p} \otimes \tilde{\mathcal{X}}^{2(k-p)+1}. \end{aligned}$$

By using the results of [proposition 14](#) and [corollary 6](#), we obtain

$$\begin{aligned}\tilde{N}^0 &= F\tilde{\mathcal{X}}^0 + o_{\mathcal{D}^{1,2}(\mathbb{R}^d \setminus T)}(1) \\ \forall k \geq 1, \tilde{N}^{2k} &= F\tilde{\mathcal{X}}^{2k} - \tilde{\mathcal{X}}^{2k-2} \otimes I + o_{\mathcal{D}^{1,2}(\mathbb{R}^d \setminus T)}(1) \\ \forall k \geq 1, \tilde{N}^{2k+1} &= F\tilde{\mathcal{X}}^{2k+1} - \tilde{\mathcal{X}}^{2k-1} \otimes I + o_{\mathcal{D}^{1,2}(\mathbb{R}^d \setminus T)}(1),\end{aligned}$$

where $o_{\mathcal{D}^{1,2}(\mathbb{R}^d \setminus T)} \rightarrow 0$ in $\mathcal{D}^{1,2}(\mathbb{R}^d \setminus T)$. This implies [\(5.20\)](#) to [\(5.22\)](#) by using once again [proposition 14](#). \square

Corollary 7. *It holds, as $\eta \rightarrow 0$:*

$$\begin{aligned}\mathbb{D}_0^2 &\rightarrow -I \\ \mathbb{D}_K^{2K+2} &= o\left(\frac{1}{\eta^{(d-2)K}}\right) = o\left(\frac{1}{\eta^{(d-2)((K+1)-1)}\right) \text{ for any } K \geq 1.\end{aligned}$$

Proof. According to [\(4.20\)](#), $\mathbb{D}_0^2 = -\int_Y N^0 \otimes N^0 \otimes Idy$. Then, [proposition 14](#) and [corollary 6](#) imply

$$\begin{aligned}-\int_Y N^0 \otimes N^0 \otimes Idy &= -(M^0)^2 \int_Y |\mathcal{X}^0|^2 dy I = -(M^0)^2 \eta^{-2(d-2)} \eta^d \int_Y |\tilde{\mathcal{X}}^0|^2 dy I \\ &\sim -F^2 \langle \tilde{\mathcal{X}}^0 \rangle^2 I \rightarrow -I.\end{aligned}$$

Finally, for $K \geq 1$ we estimate \mathbb{D}_K^{2K+2} as follows:

$$\begin{aligned}\mathbb{D}_K^{2K+2} &= (-1)^K \eta^d \eta^{-(d-2)2\lfloor K/2 \rfloor} \int_{\eta^{-1}P \setminus T} (\tilde{N}^K - \langle \tilde{N}^K \rangle) \otimes (\tilde{N}^K - \langle \tilde{N}^K \rangle) dy \\ &= O(\eta^{-(d-2)(2\lfloor K/2 \rfloor - 1)}) = o(\eta^{-(d-2)K}).\end{aligned}$$

\square

REFERENCES

- [1] Assyr Abdulle and Timothée Pouchon. Effective models for the multidimensional wave equation in heterogeneous media over long time and numerical homogenization. *Mathematical Models and Methods in Applied Sciences*, 26(14):2651–2684, 2016.
- [2] Assyr Abdulle and Timothée Pouchon. Effective models and numerical homogenization for wave propagation in heterogeneous media on arbitrary timescales. *arXiv preprint arXiv:1905.09062*, 2019.
- [3] Grégoire Allaire. *Homogénéisation des équations de Stokes et de Navier-Stokes*. PhD thesis, Université Paris 6, 1989.
- [4] Grégoire Allaire. Continuity of the Darcy’s law in the low-volume fraction limit. *Annali della Scuola Normale Superiore di Pisa. Classe di Scienze. Serie IV*, 18(4):475–499, 1991.
- [5] Grégoire Allaire. Homogenization and two-scale convergence. *SIAM Journal on Mathematical Analysis*, 23(6):1482–1518, 1992.
- [6] Grégoire Allaire. *Shape optimization by the homogenization method*, volume 146. Springer Science & Business Media, 2012.
- [7] Grégoire Allaire and Micol Amar. Boundary layer tails in periodic homogenization. *ESAIM: Control, Optimisation and Calculus of Variations*, 4:209–243, 1999.
- [8] Grégoire Allaire, Marc Briane, and Muthusamy Vanninathan. A comparison between two-scale asymptotic expansions and Bloch wave expansions for the homogenization of periodic structures. *SEMA journal*, 73(3):237–259, 2016.
- [9] Grégoire Allaire and Carlos Conca. Bloch wave homogenization and spectral asymptotic analysis. *Journal de mathématiques pures et appliquées*, 77(2):153–208, 1998.
- [10] Grégoire Allaire, Perle Geoffroy-Donders, and Olivier Pantz. Topology optimization of modulated and oriented periodic microstructures by the homogenization method. *Computers & Mathematics with Applications*, 2018.
- [11] Grégoire Allaire, Agnes Lamacz, and Jeffrey Rauch. Crime Pays; Homogenized Wave Equations for Long Times. *arXiv preprint arXiv:1803.09455*, 2018.
- [12] Grégoire Allaire and Takayuki Yamada. Optimization of dispersive coefficients in the homogenization of the wave equation in periodic structures. *Numerische Mathematik*, 140(2):265–326, 2018.
- [13] Grégoire Allaire. Homogenization of the Navier-Stokes equations with a slip boundary condition. *Communications on pure and applied mathematics*, 44(6):605–641, 1991.
- [14] Ivo Babuška. Homogenization and its application. Mathematical and computational problems. In *Numerical Solution of Partial Differential Equations—III*, pages 89–116. Elsevier, 1976.
- [15] N. Bakhvalov and G. Panasenko. *Homogenisation: averaging processes in periodic media*, volume 36 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1989.
- [16] M. P. Bendsoe and O. Sigmund. *Topology optimization*. Springer-Verlag, Berlin, 2003. Theory, methods and applications.
- [17] Martin Philip Bendsoe and Noboru Kikuchi. Generating optimal topologies in structural design using a homogenization method. *Computer methods in applied mechanics and engineering*, 71(2):197–224, 1988.

- [18] Alain Bensoussan, Jacques-Louis Lions, and George Papanicolaou. *Asymptotic analysis for periodic structures*, volume 374. American Mathematical Soc., 2011.
- [19] F Blanc and SA Nazarov. Asymptotics of solutions to the Poisson problem in a perforated domain with corners. *Journal de mathématiques pures et appliquées*, 76(10):893–911, 1997.
- [20] Thomas Borrvall and Joakim Petersson. Large-scale topology optimization in 3D using parallel computing. *Computer methods in applied mechanics and engineering*, 190(46-47):6201–6229, 2001.
- [21] Thomas Borrvall and Joakim Petersson. Topology optimization of fluids in Stokes flow. *International Journal for Numerical Methods in Fluids*, 41(1):77–107, 2003.
- [22] Haim Brezis. *Functional analysis, Sobolev spaces and partial differential equations*. Springer Science & Business Media, 2010.
- [23] Kirill D Cherednichenko and James A Evans. Full Two-Scale Asymptotic Expansion and Higher-Order Constitutive Laws in the Homogenization of the System of Quasi-static Maxwell Equations. *Multiscale Modeling & Simulation*, 14(4):1513–1539, 2016.
- [24] Andrej Cherkaev. *Variational methods for structural optimization*, volume 140. Springer Science & Business Media, 2012.
- [25] Doina Cioranescu and François Murat. A strange term coming from nowhere. In *Topics in the mathematical modelling of composite materials*, volume 31 of *Progr. Nonlinear Differential Equations Appl.*, pages 45–93. Birkhäuser Boston, Boston, MA, 1997.
- [26] Carlos Conca, François Murat, and Olivier Pironneau. The Stokes and Navier-Stokes equations with boundary conditions involving the pressure. *Japanese journal of mathematics. New series*, 20(2):279–318, 1994.
- [27] Ercan M Dede. Multiphysics topology optimization of heat transfer and fluid flow systems. In *proceedings of the COMSOL Users Conference*, 2009.
- [28] Sumer B. Dilgen, Cetin B. Dilgen, David R. Fuhrman, Ole Sigmund, and Boyan S. Lazarov. Density based topology optimization of turbulent flow heat transfer systems. *Structural and Multidisciplinary Optimization*, 57(5):1905–1918, May 2018.
- [29] Patrizia Donato and Jeannine Saint Jean Paulin. Homogenization of the Poisson equation in a porous medium with double periodicity. *Japan journal of industrial and applied mathematics*, 10(2):333, 1993.
- [30] Alexandre Ern and Jean-Luc Guermond. *Theory and practice of finite elements*, volume 159. Springer Science & Business Media, 2013.
- [31] Lawrence C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [32] Florian Feppon. *Shape and topology optimization of multiphysics systems*. PhD thesis, Thèse de doctorat de l’Université Paris Saclay préparée à l’École polytechnique, 2019.
- [33] Florian Feppon. High order homogenization of the Stokes system in a periodic porous medium. In preparation, 2020.
- [34] Perle Geoffroy Donders. *Homogenization method for topology optimization of structures built with lattice materials*. Theses, Université Paris-Saclay, December 2018.
- [35] Pierre Grisvard. *Elliptic problems in nonsmooth domains*, volume 69 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011.
- [36] Jeroen P Groen and Ole Sigmund. Homogenization-based topology optimization for high-resolution manufacturable microstructures. *International Journal for Numerical Methods in Engineering*, 113(8):1148–1163, 2018.
- [37] Wenjia Jing. A unified homogenization approach for the Dirichlet problem in Perforated Domains. *arXiv preprint arXiv:1901.08251*, 2019.
- [38] Olga A Ladyzhenskaya. *The mathematical theory of viscous incompressible flow*, volume 2. Gordon and Breach New York, 1969.
- [39] J.-L. Lions. *Perturbations singulières dans les problèmes aux limites et en contrôle optimal*. Lecture Notes in Mathematics, Vol. 323. Springer-Verlag, Berlin-New York, 1973.
- [40] J-L Lions. Some methods in the mathematical analysis of systems and their control. *Beijing, Science Press*, 1981.
- [41] Jacques-Louis Lions. Asymptotic expansions in perforated media with a periodic structure. *The Rocky Mountain Journal of Mathematics*, 10(1):125–140, 1980.
- [42] William Charles Hector McLean. *Strongly elliptic systems and boundary integral equations*. Cambridge university press, 2000.
- [43] François Murat and Jacques Simon. Sur le contrôle par un domaine géométrique. *Publication du Laboratoire d’Analyse Numérique de l’Université Pierre et Marie Curie*, 1976.
- [44] Jean-Claude Nédélec. *Acoustic and electromagnetic equations: integral representations for harmonic problems*. Springer Science & Business Media, 2001.
- [45] Olivier Pantz and Karim Trabelsi. A post-treatment of the homogenization method for shape optimization. *SIAM Journal on Control and Optimization*, 47(3):1380–1398, 2008.
- [46] M Pietropaoli, F Montomoli, and A Gaymann. Three-dimensional fluid topology optimization for heat transfer. *Structural and Multidisciplinary Optimization*, 59(3):801–812, 2019.
- [47] Nicolò Pollini, Ole Sigmund, Casper Schousboe Andreasen, and Joe Alexandersen. A “poor man’s” approach for high-resolution three-dimensional topology design for natural convection problems. *Advances in Engineering Software*, 140:102736, 2020.
- [48] Timothée Noé Pouchon. *Effective models and numerical homogenization methods for long time wave propagation in heterogeneous media*. PhD thesis, EPFL, 2017.

- [49] Jeffrey Rauch and Michael Taylor. Potential and scattering theory on wildly perturbed domains. *Journal of Functional Analysis*, 18:27–59, 1975.
- [50] Enrique Sanchez-Palencia. Fluid flow in porous media. *Non-homogeneous media and vibration theory*, pages 129–157, 1980.
- [51] Fadil Santosa and William W Symes. A dispersive effective medium for wave propagation in periodic composites. *SIAM Journal on Applied Mathematics*, 51(4):984–1005, 1991.
- [52] Valery P Smyshlyaev and KD Cherednichenko. On rigorous derivation of strain gradient effects in the overall behaviour of periodic heterogeneous media. *Journal of the Mechanics and Physics of Solids*, 48(6-7):1325–1357, 2000.
- [53] Luc Tartar. *The general theory of homogenization: a personalized introduction*, volume 7. Springer Science & Business Media, 2009.
- [54] X. Zhao, M. Zhou, O. Sigmund, and C. Andreasen. A ”poor man’s approach” to topology optimization of cooling channels based on a Darcy flow model. *International Journal of Heat and Mass Transfer*, 116:1108–1123, 2018.