

Reaction-diffusion equation on thin porous media

María Anguiano

► **To cite this version:**

María Anguiano. Reaction-diffusion equation on thin porous media. Bulletin of the Malaysian Mathematical Sciences Society, Springer Singapore, 2021. hal-02540887

HAL Id: hal-02540887

<https://hal.archives-ouvertes.fr/hal-02540887>

Submitted on 12 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reaction-diffusion equation on thin porous media

María ANGUIANO¹

Abstract

We consider a reaction-diffusion equation on a 3D thin porous media of thickness ε which is perforated by periodically distributed cylinders of size ε . On the boundary of the cylinders we prescribe a dynamical boundary condition of *pure-reactive* type. As $\varepsilon \rightarrow 0$, in the 2D limit the resulting reaction-diffusion equation has a source term coming from the dynamical-type boundary conditions imposed on boundaries of the original 3D domain.

AMS classification numbers: 35K57; 35B27

Keywords: Homogenization, energy method, dynamical boundary-value problems, porous media, thin films

1 Introduction and setting of the problem

We consider a parabolic problem in a thin porous media Ω_ε of thickness ε which is perforated by periodically distributed cylinders (obstacles) of size ε . On the boundary of the cylinders, we prescribe a dynamical boundary condition of *pure-reactive* type. The aim of this work is to prove the convergence of the homogenization process when ε goes to zero.

The asymptotic behavior of the solution of linear parabolic problems with dynamical boundary conditions of *pure-reactive* type in periodically perforated domains was analyzed by Timofte in [12]. The author obtained a new parabolic limiting problem defined on a unified domain with extra terms coming from the influence of the dynamical boundary conditions. More recently, in [1], we study the nonlinear case where the nonlinearity appears reflected in the limit equation. However, to our knowledge, there does not seem to be in the literature any study of the asymptotic behavior of the solution of parabolic models associated with dynamical boundary conditions of *pure-reactive* type in a thin porous media. Let us introduce the model we will be involved in this paper.

The geometrical setting. The periodic porous media is defined by a domain ω and an associated microstructure, or periodic cell $Y' = [-1/2, 1/2]^2$, which is made of two complementary parts: the obstacle part F' , and $Y'_f = Y' \setminus \overline{F'}$. More precisely, we assume that ω is a smooth, bounded, connected set in \mathbb{R}^2 , and that F' is an open connected subset of Y' with a smooth boundary $\partial F'$, such that $\overline{F'}$ is strictly included in Y' .

The microscale of a porous media is a small positive number ε . The domain ω is covered by a regular mesh of square of size ε : for $k' \in \mathbb{Z}^2$, each cell $Y'_{k',\varepsilon} = \varepsilon k' + \varepsilon Y'$ is divided in an obstacle part $F'_{k',\varepsilon}$ and $Y'_{f_{k',\varepsilon}}$, i.e. is similar to the unit cell Y' rescaled to size ε . We define $Y = Y' \times (0, 1) \subset \mathbb{R}^3$, which is divided in an obstacle part $F = F' \times (0, 1)$ and $Y_f = Y'_f \times (0, 1)$, and consequently $Y_{k',\varepsilon} = Y'_{k',\varepsilon} \times (0, 1) \subset \mathbb{R}^3$, which is also divided in an obstacle part $F_{k',\varepsilon}$ and $Y_{f_{k',\varepsilon}}$.

We denote by $\tau(\overline{F'_{k',\varepsilon}})$ the set of all translated images of $\overline{F'_{k',\varepsilon}}$. The set $\tau(\overline{F'_{k',\varepsilon}})$ represents the obstacles in \mathbb{R}^2 .

¹Departamento de Análisis Matemático. Facultad de Matemáticas. Universidad de Sevilla, 41012 Sevilla (Spain) anguiano@us.es

We define $\omega_\varepsilon = \omega \setminus \bigcup_{k' \in \mathcal{K}_\varepsilon} \overline{F}'_{k',\varepsilon} \subset \mathbb{R}^2$, where $\mathcal{K}_\varepsilon = \{k' \in \mathbb{Z}^2 : Y'_{k',\varepsilon} \cap \omega \neq \emptyset\}$, and we define the thin porous media $\Omega_\varepsilon \subset \mathbb{R}^3$ by

$$\Omega_\varepsilon = \{(x_1, x_2, x_3) \in \omega_\varepsilon \times \mathbb{R} : 0 < x_3 < \varepsilon\}. \quad (1)$$

We make the following assumption:

The obstacles $\tau(\overline{F}'_{k',\varepsilon})$ do not intersect the boundary $\partial\omega$.

We define $F'_{k',\varepsilon}^\varepsilon = F'_{k',\varepsilon} \times (0, \varepsilon)$. Denote by S_ε the set of the obstacles contained in Ω_ε . Then, S_ε is a finite union of obstacles, i.e.

$$S_\varepsilon = \bigcup_{k' \in \mathcal{K}_\varepsilon} \overline{F}'_{k',\varepsilon}^\varepsilon.$$

We define

$$\widetilde{\Omega}_\varepsilon = \omega_\varepsilon \times (0, 1), \quad \Omega = \omega \times (0, 1), \quad \Lambda_\varepsilon = \omega \times (0, \varepsilon). \quad (2)$$

We observe that $\widetilde{\Omega}_\varepsilon = \Omega \setminus \bigcup_{k' \in \mathcal{K}_\varepsilon} \overline{F}'_{k',\varepsilon}$, and we define $F_\varepsilon = \bigcup_{k' \in \mathcal{K}_\varepsilon} \overline{F}'_{k',\varepsilon}$ as the set of the obstacles contained in $\widetilde{\Omega}_\varepsilon$.

Position of the problem. We consider the following problem for a reaction-diffusion equation with dynamical boundary conditions of *pure-reactive* type on the surface of the obstacles and zero Dirichlet condition on the exterior boundary,

$$\left\{ \begin{array}{ll} \frac{\partial u_\varepsilon}{\partial t} - \Delta u_\varepsilon + \kappa u_\varepsilon = f(x', t) & \text{in } \Omega_\varepsilon \times (0, T), \\ \nabla u_\varepsilon \cdot \nu_\varepsilon + \varepsilon \frac{\partial u_\varepsilon}{\partial t} = \varepsilon g(x', t) & \text{on } \partial S_\varepsilon \times (0, T), \\ u_\varepsilon(x, 0) = u_\varepsilon^0(x), & \text{for } x \in \Omega_\varepsilon, \\ u_\varepsilon(x, 0) = \psi_\varepsilon^0(x), & \text{for } x \in \partial S_\varepsilon, \\ u_\varepsilon = 0, & \text{on } \partial \Lambda_\varepsilon \times (0, T), \end{array} \right. \quad (3)$$

where $u_\varepsilon = u_\varepsilon(x, t)$, $x = (x', x_3) \in \Omega_\varepsilon$, $t \in (0, T)$ and $T > 0$. The first equation states the law of standard diffusion in Ω_ε , $\Delta = \Delta_x$ denotes the Laplacian operator with respect to the space variable and $\kappa > 0$ is a given constant. The boundary equation (3)₂ is multiplied by ε to compensate the growth of the surface by shrinking ε , where the value of u_ε is assumed to be the trace of the function u_ε defined for $x \in \Omega_\varepsilon$, ν_ε is the outer normal to ∂S_ε . The term $\nabla u_\varepsilon \cdot \nu_\varepsilon$ represents the interaction domain-boundary. We assume that

$$u_\varepsilon^0 \in L^2(\Omega), \quad \psi_\varepsilon^0 \in L^2(\partial S_\varepsilon), \quad (4)$$

are given, and that there exists a positive constant C such that

$$\varepsilon^{-1} |u_\varepsilon^0|_{\Omega_\varepsilon}^2 + |\psi_\varepsilon^0|_{\partial S_\varepsilon}^2 \leq C, \quad (5)$$

where $|\cdot|_{\Omega_\varepsilon}$ and $|\cdot|_{\partial S_\varepsilon}$ denote the norm in $L^2(\Omega_\varepsilon)$ and $L^2(\partial S_\varepsilon)$, respectively.

Moreover, we assume that

$$f \in L^2(0, T; L^2(\omega)), \quad g \in L^2(0, T; H_0^1(\omega)), \quad (6)$$

are given. This choice of f and g is usual when we deal with thin domains. Since the thickness of the domain, ε , is small then the exterior force can be considered independent of the vertical variable.

In this paper, our main motivation is to study the asymptotic behavior of the solution u_ε of (3) as ε , the size of the cylinders and the corresponding thickness of the thin film, tend to zero. We use the related methods both in porous media and in thin film, respectively, in order to rigorously derive the homogenized model, which is the same as that in two dimensions.

For this purpose, we first perform a change of variables which consists in stretching in the x_3 -direction by a factor $1/\varepsilon$. As in [2], we use the dilatation in the variable x_3 , i.e.

$$y_3 = \frac{x_3}{\varepsilon}, \quad (7)$$

in order to have the functions defined in the open set with fixed height $\tilde{\Omega}_\varepsilon$ defined by (2). In this sense, we define \tilde{u}_ε by

$$\tilde{u}_\varepsilon(x', y_3, t) = u_\varepsilon(x', \varepsilon y_3, t), \quad a.e. (x', y_3, t) \in \tilde{\Omega}_\varepsilon \times (0, T), \quad (8)$$

and our goal then is to describe the asymptotic behavior of this new sequence \tilde{u}_ε when ε tends to zero.

Using the dilatation in the variable x_3 given by (7), we have that Δu_ε is transformed into

$$\Delta_{x'} \tilde{u}_\varepsilon + \frac{1}{\varepsilon^2} \partial_{y_3}^2 \tilde{u}_\varepsilon.$$

The factor $1/\varepsilon^2$ in front of the derivate in the y_3 direction means a very fast diffusion in the vertical direction. In some sense, we have substituted the thin porous media Ω_ε by a non-thin domain $\tilde{\Omega}_\varepsilon$, but with a very strong diffusion mechanism in the y_3 -direction. Because of the presence of this very strong diffusion mechanism, it is expected that solutions of problem which satisfies \tilde{u}_ε , defined in (20), become homogeneous in the y_3 -direction so that the limit solution will not have a dependence in this direction, and therefore, the limit problem will be two dimensional.

The method we follow in this paper is the so-called energy method of Tartar [13], which has been considered by many authors (see, for instance, Cioranescu and Donato [5], Conca and Donato [8], Conca *et al.* [9], Timofte [12] and [1]) and the technique introduced by Vanninathan [14] for the Steklov problems which transforms surface integrals into volume integrals. We get the following limit problem to (20), as ε goes to zero.

Theorem 1.1 (Main Theorem). *Under the assumptions (5)–(6), assume that $(u_\varepsilon^0, \psi_\varepsilon^0) \in H^1(\Omega_\varepsilon) \times H^{1/2}(\partial S_\varepsilon)$ and there exists a positive constant C such that*

$$\varepsilon^{-1/2} |\nabla u_\varepsilon^0|_{\Omega_\varepsilon} \leq C. \quad (9)$$

Let $(u_\varepsilon, \psi_\varepsilon)$ be the unique solution of the problem (3), where ψ_ε is the trace of the function u_ε defined for $x \in \Omega_\varepsilon$ and a.e. $t \in (0, T]$. Then, there exists an extension $\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon$ of \tilde{u}_ε , where \tilde{u}_ε is the dilation of u_ε given by (8), into all $\Omega \times (0, T)$, such that, as $\varepsilon \rightarrow 0$,

$$\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) \rightarrow u(t) \quad \text{strongly in } L^2(\Omega), \quad \forall t \in [0, T],$$

where $u \in L^2(0, T; H_0^1(\omega))$ is the unique solution of the following problem

$$\begin{cases} \left(\frac{|Y_f'|}{|Y'|} + \frac{|\partial F'|}{|Y'|} \right) \frac{\partial u}{\partial t} - \operatorname{div}_{x'} (Q \nabla_{x'} u) + \frac{|Y_f'|}{|Y'|} \kappa u = \frac{|Y_f'|}{|Y'|} f + \frac{|\partial F'|}{|Y'|} g, & \text{in } \omega \times (0, T), \\ u(x', 0) = u_0(x'), & \text{for } x' \in \omega, \\ u = 0, & \text{on } \partial\omega \times (0, T). \end{cases} \quad (10)$$

The homogenized matrix $Q = ((q_{i,j}))$, $1 \leq i, j \leq 2$, which is symmetric and positive-definite, is given by

$$q_{i,j} = \frac{1}{|Y_f'|} \int_{Y_f'} (e_i + \nabla_{y'} w_i) \cdot (e_j + \nabla_{y'} w_j) dy', \quad (11)$$

where $w_i \in \mathbb{H}_{\text{per}} \setminus \mathbb{R}$, $i = 1, 2$, is the unique solution of the cell problem

$$\begin{cases} -\operatorname{div}_{y'} (e_i + \nabla_{y'} w_i) = 0, & \text{in } Y_f', \\ (e_i + \nabla_{y'} w_i) \cdot \nu' = 0, & \text{on } \partial F', \\ w_i & \text{is } Y' \text{-periodic.} \end{cases} \quad (12)$$

Here, e_i is the i element of the canonical basis in \mathbb{R}^2 , ν' is the outer normal to $\partial F'$, and \mathbb{H}_{per} is the space of functions from $H^1(Y'_f)$ which are Y' -periodic.

The structure of the paper as follows. In Section 2, we introduce some notations which are used in the paper. In Section 3, we give a weak formulation of the problem, and establish the existence and uniqueness of solution. In Section 4, we establish the problem which satisfies \tilde{u}_ε given by (8). Some *a priori* estimates for \tilde{u}_ε are rigorously obtained in Section 5. A compactness result is addressed in Section 6. Finally, the main goal of proving the asymptotic behavior of the solution \tilde{u}_ε is established in Section 7.

2 Some notations

Along this paper, the points $x \in \mathbb{R}^3$ will be decomposed as $x = (x', x_3)$ with $x' \in \mathbb{R}^2$, $x_3 \in \mathbb{R}$. We also use the notation x' to denote a generic vector of \mathbb{R}^2 . We denote by $\chi_{\tilde{\Omega}_\varepsilon}$ the characteristic function of the domain $\tilde{\Omega}_\varepsilon$.

We denote by $(\cdot, \cdot)_{\Omega_\varepsilon}$ (respectively, $(\cdot, \cdot)_{\partial S_\varepsilon}$) the inner product in $L^2(\Omega_\varepsilon)$ (respectively, in $L^2(\partial S_\varepsilon)$), and by $|\cdot|_{\Omega_\varepsilon}$ (respectively, $|\cdot|_{\partial S_\varepsilon}$) the associated norm. We also denote $(\cdot, \cdot)_{\Omega_\varepsilon}$ the inner product in $(L^2(\Omega_\varepsilon))^3$.

We denote by $(\cdot, \cdot)_{\tilde{\Omega}_\varepsilon}$ (respectively, $(\cdot, \cdot)_{\partial F_\varepsilon}$) the inner product in $L^2(\tilde{\Omega}_\varepsilon)$ (respectively, in $L^2(\partial F_\varepsilon)$), and by $|\cdot|_{\tilde{\Omega}_\varepsilon}$ (respectively, $|\cdot|_{\partial F_\varepsilon}$) the associated norm. We also denote $(\cdot, \cdot)_{\tilde{\Omega}_\varepsilon}$ the inner product in $(L^2(\tilde{\Omega}_\varepsilon))^N$, with $N = 2, 3$.

We denote by $(\cdot, \cdot)_\Omega$ the inner product in $L^2(\Omega)$, and by $|\cdot|_\Omega$ the associated norm. We also denote $(\cdot, \cdot)_\Omega$ the inner product in $(L^2(\Omega))^3$. By $\|\cdot\|_\Omega$ we denote the norm in $H^1(\Omega)$.

We denote by $(\cdot, \cdot)_{\omega_\varepsilon}$ the inner product in $L^2(\omega_\varepsilon)$ and by $|\cdot|_{\omega_\varepsilon}$ the associated norm. We also denote $(\cdot, \cdot)_{\omega_\varepsilon}$ the inner product in $(L^2(\omega_\varepsilon))^2$. By $\|\cdot\|_{\omega_\varepsilon}$ we denote the norm in $H^1(\omega_\varepsilon)$, which is associated to the inner product

$$((\cdot, \cdot))_{\omega_\varepsilon} := (\nabla_{x'} \cdot, \nabla_{x'} \cdot)_{\omega_\varepsilon} + (\cdot, \cdot)_{\omega_\varepsilon}.$$

By $\|\cdot\|_{\tilde{\Omega}_\varepsilon}$ we denote the norm in $H^1(\tilde{\Omega}_\varepsilon)$, which is associated to the inner product

$$((\cdot, \cdot))_{\tilde{\Omega}_\varepsilon} := (\nabla_{x'} \cdot, \nabla_{x'} \cdot)_{\tilde{\Omega}_\varepsilon} + (\partial_{y_3} \cdot, \partial_{y_3} \cdot)_{\tilde{\Omega}_\varepsilon} + (\cdot, \cdot)_{\tilde{\Omega}_\varepsilon}.$$

By $\|\cdot\|_{\tilde{\Omega}_\varepsilon, T}$ we denote the norm in $L^2(0, T; H^1(\tilde{\Omega}_\varepsilon))$.

By $|\cdot|_{\tilde{\Omega}_\varepsilon, T}$ (respectively $|\cdot|_{\partial F_\varepsilon, T}$), we denote the norms in $L^2(0, T; L^2(\tilde{\Omega}_\varepsilon))$ and $L^2(0, T; (L^2(\tilde{\Omega}_\varepsilon))^3)$ (respectively $L^2(0, T; L^2(\partial F_\varepsilon))$). By $|\cdot|_{\Omega, T}$, we denote the norms in $L^2(0, T; L^2(\Omega))$ and $L^2(0, T; (L^2(\Omega))^3)$.

We denote by γ_0 the trace operator $u \mapsto u|_{\partial\Omega_\varepsilon}$. The trace operator belongs to $\mathcal{L}(H^1(\Omega_\varepsilon), H^{1/2}(\partial\Omega_\varepsilon))$, and we will use $\|\gamma_0\|$ to denote the norm of γ_0 in this space. Analogously, we denote by $\tilde{\gamma}_0$ the trace operator $\tilde{u} \mapsto \tilde{u}|_{\partial\tilde{\Omega}_\varepsilon}$.

We will use $\|\cdot\|_{\partial\tilde{\Omega}_\varepsilon}$ to denote the norm in $H^{1/2}(\partial\tilde{\Omega}_\varepsilon)$, which is given by $\|\tilde{\phi}\|_{\partial\tilde{\Omega}_\varepsilon} = \inf\{\|\tilde{v}\|_{\tilde{\Omega}_\varepsilon} : \tilde{\gamma}_0(\tilde{v}) = \tilde{\phi}\}$. We remember that with this norm, $H^{1/2}(\partial\tilde{\Omega}_\varepsilon)$ is a Hilbert space.

We denote by $H^r_{\partial\Lambda_\varepsilon}(\Omega_\varepsilon)$ and $H^r_{\partial\Lambda_\varepsilon}(\partial\Omega_\varepsilon)$, for $r \geq 0$, the standard Sobolev spaces which are closed subspaces of $H^r(\Omega_\varepsilon)$ and $H^r(\partial\Omega_\varepsilon)$, respectively, and the subscript $\partial\Lambda_\varepsilon$ means that, respectively, traces or functions in $\partial\Omega_\varepsilon$, vanish on this part of the boundary of Ω_ε , i.e.

$$H^r_{\partial\Lambda_\varepsilon}(\Omega_\varepsilon) = \{v \in H^r(\Omega_\varepsilon) : \gamma_0(v) = 0 \text{ on } \partial\Lambda_\varepsilon\},$$

and

$$H^r_{\partial\Lambda_\varepsilon}(\partial\Omega_\varepsilon) = \{v \in H^r(\partial\Omega_\varepsilon) : v = 0 \text{ on } \partial\Lambda_\varepsilon\}.$$

Let us notice that, in fact, we can consider an element of $H^{1/2}(\partial S_\varepsilon)$ as an element of $H^{1/2}_{\partial\Lambda_\varepsilon}(\partial\Omega_\varepsilon)$, and we can consider the given ψ_ε^0 as an element of $L^2_{\partial\Lambda_\varepsilon}(\partial\Omega_\varepsilon)$.

We denote by $H_{\partial\Omega}^r(\tilde{\Omega}_\varepsilon)$ and $H_{\partial\Omega}^r(\partial\tilde{\Omega}_\varepsilon)$, for $r \geq 0$, the standard Sobolev spaces which are closed subspaces of $H^r(\tilde{\Omega}_\varepsilon)$ and $H^r(\partial\tilde{\Omega}_\varepsilon)$, respectively, and the subscript $\partial\Omega$ means that, respectively, traces or functions in $\partial\tilde{\Omega}_\varepsilon$, vanish on this part of the boundary of $\tilde{\Omega}_\varepsilon$, i.e.

$$H_{\partial\Omega}^r(\tilde{\Omega}_\varepsilon) = \{\tilde{v} \in H^r(\tilde{\Omega}_\varepsilon) : \gamma_0(\tilde{v}) = 0 \text{ on } \partial\Omega\},$$

and

$$H_{\partial\Omega}^r(\partial\tilde{\Omega}_\varepsilon) = \{\tilde{v} \in H^r(\partial\tilde{\Omega}_\varepsilon) : \tilde{v} = 0 \text{ on } \partial\Omega\}.$$

Let us consider the space

$$H := L^2(\tilde{\Omega}_\varepsilon) \times L_{\partial\Omega}^2(\partial\tilde{\Omega}_\varepsilon),$$

with the natural inner product $((\tilde{v}, \tilde{\phi}), (\tilde{w}, \tilde{\varphi}))_H = (\tilde{v}, \tilde{w})_{\tilde{\Omega}_\varepsilon} + \varepsilon(\tilde{\phi}, \tilde{\varphi})_{\partial F_\varepsilon}$, which in particular induces the norm $\|(\cdot, \cdot)\|_H$ given by

$$\|(\tilde{v}, \tilde{\phi})\|_H^2 = |\tilde{v}|_{\tilde{\Omega}_\varepsilon}^2 + \varepsilon|\tilde{\phi}|_{\partial F_\varepsilon}^2, \quad (\tilde{v}, \tilde{\phi}) \in H.$$

Let us also consider the space

$$V_1 := \left\{ (\tilde{v}, \gamma_0(\tilde{v})) : \tilde{v} \in H_{\partial\Omega}^1(\tilde{\Omega}_\varepsilon) \right\}.$$

We note that V_1 is a closed vector subspace of $H_{\partial\Omega}^1(\tilde{\Omega}_\varepsilon) \times H_{\partial\Omega}^{1/2}(\partial\tilde{\Omega}_\varepsilon)$, and therefore, with the norm $\|(\cdot, \cdot)\|_{V_1}$ given by

$$\|(\tilde{v}, \gamma_0(\tilde{v}))\|_{V_1}^2 = \|\tilde{v}\|_{\tilde{\Omega}_\varepsilon}^2 + \|\gamma_0(\tilde{v})\|_{\partial F_\varepsilon}^2, \quad (\tilde{v}, \gamma_0(\tilde{v})) \in V_1,$$

V_1 is a Hilbert space.

For a vectorial function $v = (v', v_3)$ and a scalar function w , we introduce the operators: ∇_ε and div_ε , by

$$\nabla_\varepsilon w = (\nabla_{x'} w, \frac{1}{\varepsilon} \partial_{y_3} w)^t, \quad \text{div}_\varepsilon v = \text{div}_{x'} v' + \frac{1}{\varepsilon} \partial_{y_3} v_3. \quad (13)$$

We denote by $|\mathcal{O}|$ the Lebesgue measure of $|\mathcal{O}|$ (3-dimensional if \mathcal{O} is a 3-dimensional open set, 2-dimensional if \mathcal{O} is a curve).

Finally, we denote by C a generic positive constant, independent of ε , which can change from line to line.

3 Existence and uniqueness of solution of the problem (3)

We state in this section a result on the existence and uniqueness of solution of problem (3).

Definition 3.1. *A weak solution of (3) is a pair of functions $(u_\varepsilon, \psi_\varepsilon)$, satisfying*

$$u_\varepsilon \in C([0, T]; L^2(\Omega_\varepsilon)), \quad \psi_\varepsilon \in C([0, T]; L_{\partial\Lambda_\varepsilon}^2(\partial\Omega_\varepsilon)), \quad \text{for all } T > 0, \quad (14)$$

$$u_\varepsilon \in L^2(0, T; H_{\partial\Lambda_\varepsilon}^1(\Omega_\varepsilon)), \quad \text{for all } T > 0, \quad (15)$$

$$\psi_\varepsilon \in L^2(0, T; H_{\partial\Lambda_\varepsilon}^{1/2}(\partial\Omega_\varepsilon)), \quad \text{for all } T > 0, \quad (16)$$

$$\gamma_0(u_\varepsilon(t)) = \psi_\varepsilon(t), \quad \text{a.e. } t \in (0, T], \quad (17)$$

$$\begin{cases} \frac{d}{dt}(u_\varepsilon(t), v)_{\Omega_\varepsilon} + \varepsilon \frac{d}{dt}(\psi_\varepsilon(t), \gamma_0(v))_{\partial S_\varepsilon} + (\nabla u_\varepsilon(t), \nabla v)_{\Omega_\varepsilon} + \kappa(u_\varepsilon(t), v)_{\Omega_\varepsilon} \\ = (f(x', t), v)_{\Omega_\varepsilon} + \varepsilon (g(x', t), \gamma_0(v))_{\partial S_\varepsilon} \\ \text{in } \mathcal{D}'(0, T), \text{ for all } v \in H_{\partial\Lambda_\varepsilon}^1(\Omega_\varepsilon), \end{cases} \quad (18)$$

$$u_\varepsilon(0) = u_\varepsilon^0, \quad \text{and} \quad \psi_\varepsilon(0) = \psi_\varepsilon^0. \quad (19)$$

Thanks to [1, Theorem 3.3], we have the following result.

Theorem 3.2. *Under the assumptions (4)–(6), there exists a unique solution $(u_\varepsilon, \psi_\varepsilon)$ of the problem (3). Moreover, this solution satisfies the energy equality*

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} (|u_\varepsilon(t)|_{\tilde{\Omega}_\varepsilon}^2 + \varepsilon |\psi_\varepsilon(t)|_{\partial S_\varepsilon}^2) + |\nabla u_\varepsilon(t)|_{\tilde{\Omega}_\varepsilon}^2 + \kappa |u_\varepsilon(t)|_{\tilde{\Omega}_\varepsilon}^2 \\ &= (f(x', t), u_\varepsilon(t))_{\tilde{\Omega}_\varepsilon} + \varepsilon (g(x', t), \psi_\varepsilon(t))_{\partial S_\varepsilon}, \quad \text{a.e. } t \in (0, T). \end{aligned}$$

4 Dilatation in the variable x_3 : definition of \tilde{u}_ε

In this section, we use the dilatation in the variable x_3 , i.e. we use (7) in order to have the functions defined in the open set with fixed height $\tilde{\Omega}_\varepsilon$ defined by (2). In this sense, we define $\tilde{u}_\varepsilon \in L^2(0, T; H_{\partial\Omega}^1(\tilde{\Omega}_\varepsilon))$, for all $T > 0$, by (8), and using the transformation (7), the system (3) can be rewritten as

$$\left\{ \begin{array}{ll} \frac{\partial \tilde{u}_\varepsilon}{\partial t} - \operatorname{div}_\varepsilon(\nabla_\varepsilon \tilde{u}_\varepsilon) + \kappa \tilde{u}_\varepsilon = f(x', t) & \text{in } \tilde{\Omega}_\varepsilon \times (0, T), \\ \nabla_\varepsilon \tilde{u}_\varepsilon \cdot \tilde{\nu}_\varepsilon + \varepsilon \frac{\partial \tilde{u}_\varepsilon}{\partial t} = \varepsilon g(x', t) & \text{on } \partial F_\varepsilon \times (0, T), \\ \tilde{u}_\varepsilon(x', y_3, 0) = \tilde{u}_\varepsilon^0(x', y_3), & \text{for } (x', y_3) \in \tilde{\Omega}_\varepsilon, \\ \tilde{u}_\varepsilon(x', y_3, 0) = \tilde{\psi}_\varepsilon^0(x', y_3), & \text{for } (x', y_3) \in \partial F_\varepsilon, \\ \tilde{u}_\varepsilon = 0, & \text{on } \partial\Omega \times (0, T), \end{array} \right. \quad (20)$$

where $\operatorname{div}_\varepsilon$ and ∇_ε are given by (13), and $\tilde{\nu}_\varepsilon$ is the outer normal to ∂F_ε .

Taking in (3) as test function $\tilde{v}(x', x_3/\varepsilon)$ with $\tilde{v} \in H_{\partial\Omega}^1(\tilde{\Omega}_\varepsilon)$, applying the change of variables (7) and taking into account that $dx = \varepsilon dx' dy_3$ and $d\sigma(x) = \varepsilon d\sigma(x') dy_3$, the variational formulation of problem (20) is then the following one

$$\left\{ \begin{array}{l} \frac{d}{dt} \left(\int_{\tilde{\Omega}_\varepsilon} \tilde{u}_\varepsilon(t) \tilde{v} dx' dy_3 \right) + \varepsilon \frac{d}{dt} \left(\int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon(t)) \gamma_0(\tilde{v}) d\sigma(x') dy_3 \right) + \int_{\tilde{\Omega}_\varepsilon} \nabla_\varepsilon \tilde{u}_\varepsilon(t) \cdot \nabla_\varepsilon \tilde{v} dx' dy_3 \\ + \kappa \int_{\tilde{\Omega}_\varepsilon} \tilde{u}_\varepsilon(t) \tilde{v} dx' dy_3 = \int_{\tilde{\Omega}_\varepsilon} f(x', t) \tilde{v} dx' dy_3 + \varepsilon \int_{\partial F_\varepsilon} g(x', t) \gamma_0(\tilde{v}) d\sigma(x') dy_3 \\ \text{in } \mathcal{D}'(0, T), \text{ for all } \tilde{v} \in H_{\partial\Omega}^1(\tilde{\Omega}_\varepsilon). \end{array} \right. \quad (21)$$

Moreover, \tilde{u}_ε satisfies the energy equality

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} (|(\tilde{u}_\varepsilon(t), \gamma_0(\tilde{u}_\varepsilon(t)))|_H^2) + |\nabla_\varepsilon \tilde{u}_\varepsilon(t)|_{\tilde{\Omega}_\varepsilon}^2 + \kappa |\tilde{u}_\varepsilon(t)|_{\tilde{\Omega}_\varepsilon}^2 \\ &= (f(x', t), \tilde{u}_\varepsilon(t))_{\tilde{\Omega}_\varepsilon} + \varepsilon (g(x', t), \gamma_0(\tilde{u}_\varepsilon(t)))_{\partial F_\varepsilon}, \quad \text{a.e. } t \in (0, T). \end{aligned} \quad (22)$$

Considering in (5) the change of variables given in (7), and taking into account that $dx = \varepsilon dx' dy_3$ and $d\sigma(x) = \varepsilon d\sigma(x') dy_3$, we can deduce that

$$|\tilde{u}_\varepsilon^0|_{\tilde{\Omega}_\varepsilon}^2 + \varepsilon |\tilde{\psi}_\varepsilon^0|_{\partial F_\varepsilon}^2 \leq C. \quad (23)$$

5 *A priori* estimates for \tilde{u}_ε

Let us obtain some *a priori* estimates for \tilde{u}_ε .

Lemma 5.1. *Under the assumptions (4)–(6), there exists a positive constant C independent of ε , such that the solution \tilde{u}_ε of the problem (20) satisfies*

$$|\nabla_\varepsilon \tilde{u}_\varepsilon|_{\tilde{\Omega}_\varepsilon, T} \leq C, \quad |\tilde{u}_\varepsilon|_{\tilde{\Omega}_\varepsilon, T} \leq C, \quad (24)$$

$$|\tilde{u}_\varepsilon(t)|_{\tilde{\Omega}_\varepsilon} \leq C, \quad \sqrt{\varepsilon}|\gamma_0(\tilde{u}_\varepsilon(t))|_{\partial F_\varepsilon} \leq C, \quad (25)$$

for all $t \in (0, T)$.

Proof. By (22), we have

$$\frac{d}{dt} (|(\tilde{u}_\varepsilon(t), \gamma_0(\tilde{u}_\varepsilon(t)))|_H^2) + 2|\nabla_\varepsilon \tilde{u}_\varepsilon(t)|_{\tilde{\Omega}_\varepsilon}^2 + 2\kappa |\tilde{u}_\varepsilon(t)|_{\tilde{\Omega}_\varepsilon}^2 \leq |(\tilde{u}_\varepsilon(t), \gamma_0(\tilde{u}_\varepsilon(t)))|_H^2 + |f(t)|_{\omega_\varepsilon}^2 + \varepsilon |g(t)|_{\partial F'_\varepsilon}^2.$$

Integrating between 0 and t , we obtain

$$\begin{aligned} & |(\tilde{u}_\varepsilon(t), \gamma_0(\tilde{u}_\varepsilon(t)))|_H^2 + 2\min\{1, \kappa\} \int_0^t (|\nabla_\varepsilon \tilde{u}_\varepsilon(s)|_{\tilde{\Omega}_\varepsilon}^2 + |\tilde{u}_\varepsilon(s)|_{\tilde{\Omega}_\varepsilon}^2) ds \\ & \leq |(\tilde{u}_\varepsilon^0, \gamma_0(\tilde{u}_\varepsilon^0))|_H^2 + \int_0^t |(\tilde{u}_\varepsilon(s), \gamma_0(\tilde{u}_\varepsilon(s)))|_H^2 ds + \int_0^t (|f(s)|_{\omega_\varepsilon}^2 + \varepsilon |g(s)|_{\partial F'_\varepsilon}^2) ds. \end{aligned} \quad (26)$$

By [1, Lemma 4.1] with $p = 2$, we can deduce

$$\varepsilon |g(s)|_{\partial F'_\varepsilon}^2 \leq C (|g(s)|_{\omega_\varepsilon}^2 + \varepsilon^2 |\nabla g(s)|_{\omega_\varepsilon}^2) \leq C |g(s)|_{\omega_\varepsilon}^2,$$

which together with (6) gives

$$\int_0^T (|f(s)|_{\omega_\varepsilon}^2 + \varepsilon |g(s)|_{\partial F'_\varepsilon}^2) ds \leq C. \quad (27)$$

Taking into account (23) and (27) in (26) and applying Gronwall Lemma, in particular we obtain (25). Finally, taking into account (23), (25) and (27) in (26), we get (24). \square

Now, if we want to take the inner product in (20) with \tilde{u}'_ε , we need that $u'_\varepsilon \in L^2(0, T; H^1_{\partial\Omega}(\tilde{\Omega}_\varepsilon))$. However, we do not have it for our weak solution. Therefore, we use the Galerkin method in order to prove, rigorously, new *a priori* estimates for \tilde{u}_ε .

On the space V_1 we define a continuous symmetric linear operator $A_1 : V_1 \rightarrow V'_1$, given by

$$\langle A_1((\tilde{v}, \gamma_0(\tilde{v})), (\tilde{w}, \gamma_0(\tilde{w}))) \rangle = (\nabla_\varepsilon \tilde{v}, \nabla_\varepsilon \tilde{w})_{\tilde{\Omega}_\varepsilon} + \kappa (\tilde{v}, \tilde{w})_{\tilde{\Omega}_\varepsilon}, \quad (28)$$

for all $\tilde{v}, \tilde{w} \in H^1_{\partial\Omega}(\tilde{\Omega}_\varepsilon)$.

We observe that A_1 is coercive. In fact, taking into account that $\varepsilon < 1$, we have

$$\begin{aligned} \langle A_1((\tilde{v}, \gamma_0(\tilde{v})), (\tilde{v}, \gamma_0(\tilde{v}))) \rangle &= |\nabla_{x'} \tilde{v}|_{\tilde{\Omega}_\varepsilon}^2 + \frac{1}{\varepsilon^2} |\partial_{y_3} \tilde{v}|_{\tilde{\Omega}_\varepsilon}^2 + \kappa |\tilde{v}|_{\tilde{\Omega}_\varepsilon}^2 \\ &> |\nabla_{x'} \tilde{v}|_{\tilde{\Omega}_\varepsilon}^2 + |\partial_{y_3} \tilde{v}|_{\tilde{\Omega}_\varepsilon}^2 + \kappa |\tilde{v}|_{\tilde{\Omega}_\varepsilon}^2 \\ &\geq \min\{1, \kappa\} \|\tilde{v}\|_{\tilde{\Omega}_\varepsilon}^2 \\ &= \frac{1}{1 + \|\gamma_0\|^2} \min\{1, \kappa\} \|\tilde{v}\|_{\tilde{\Omega}_\varepsilon}^2 \\ &\quad + \frac{\|\gamma_0\|^2}{1 + \|\gamma_0\|^2} \min\{1, \kappa\} \|\tilde{v}\|_{\tilde{\Omega}_\varepsilon}^2 \\ &\geq \frac{1}{1 + \|\gamma_0\|^2} \min\{1, \kappa\} \|(\tilde{v}, \gamma_0(\tilde{v}))\|_{V_1}^2, \end{aligned} \quad (29)$$

for all $\tilde{v} \in H^1_{\partial\Omega}(\tilde{\Omega}_\varepsilon)$.

Let us observe that the space $H^1_{\partial\Omega}(\tilde{\Omega}_\varepsilon) \times H^{1/2}_{\partial\Omega}(\partial\tilde{\Omega}_\varepsilon)$ is compactly imbedded in H , and therefore, for the symmetric and coercive linear continuous operator $A_1 : V_1 \rightarrow V'_1$, where A_1 is given by (28), there exists a non-decreasing sequence $0 < \lambda_1 \leq \lambda_2 \leq \dots$ of eigenvalues associated to the operator A_1 with $\lim_{j \rightarrow \infty} \lambda_j = \infty$,

and there exists a Hilbert basis of H , $\{(w_j, \gamma_0(w_j)) : j \geq 1\} \subset D(A_1)$, with $\text{span}\{(w_j, \gamma_0(w_j)) : j \geq 1\}$ densely embedded in V_1 , such that

$$A_1((w_j, \gamma_0(w_j))) = \lambda_j(w_j, \gamma_0(w_j)) \quad \forall j \geq 1.$$

Taking into account the above facts, we denote by

$$(\tilde{u}_{\varepsilon,m}(t), \gamma_0(\tilde{u}_{\varepsilon,m}(t))) = (\tilde{u}_{\varepsilon,m}(t; 0, \tilde{u}_{\varepsilon}^0, \tilde{\psi}_{\varepsilon}^0), \gamma_0(\tilde{u}_{\varepsilon,m}(t; 0, \tilde{u}_{\varepsilon}^0, \tilde{\psi}_{\varepsilon}^0)))$$

the Galerkin approximation of the solution $(\tilde{u}_{\varepsilon}(t; 0, \tilde{u}_{\varepsilon}^0, \tilde{\psi}_{\varepsilon}^0), \gamma_0(\tilde{u}_{\varepsilon}(t; 0, \tilde{u}_{\varepsilon}^0, \tilde{\psi}_{\varepsilon}^0)))$ to (20) for each integer $m \geq 1$, which is given by

$$(\tilde{u}_{\varepsilon,m}(t), \gamma_0(\tilde{u}_{\varepsilon,m}(t))) = \sum_{j=1}^m \delta_{\varepsilon m j}(t)(w_j, \gamma_0(w_j)), \quad (30)$$

and is the solution of

$$\begin{aligned} & \frac{d}{dt} ((\tilde{u}_{\varepsilon,m}(t), \gamma_0(\tilde{u}_{\varepsilon,m}(t))), (w_j, \gamma_0(w_j)))_H \\ & + \langle A_1((\tilde{u}_{\varepsilon,m}(t), \gamma_0(\tilde{u}_{\varepsilon,m}(t))), (w_j, \gamma_0(w_j))) \rangle \\ & = (f(t), w_j)_{\tilde{\Omega}_{\varepsilon}} + \varepsilon(g(t), \gamma_0(w_j))_{\partial F_{\varepsilon}}, \quad j = 1, \dots, m, \end{aligned} \quad (31)$$

with initial data

$$(\tilde{u}_{\varepsilon,m}(0), \gamma_0(\tilde{u}_{\varepsilon,m}(0))) = (\tilde{u}_{\varepsilon,m}^0, \gamma_0(\tilde{u}_{\varepsilon,m}^0)), \quad (32)$$

where

$$\delta_{\varepsilon m j}(t) = (\tilde{u}_{\varepsilon,m}(t), w_j)_{\tilde{\Omega}_{\varepsilon}} + (\gamma_0(\tilde{u}_{\varepsilon,m}(t)), \gamma_0(w_j))_{\partial F_{\varepsilon}},$$

and $(\tilde{u}_{\varepsilon,m}^0, \gamma_0(\tilde{u}_{\varepsilon,m}^0)) \in \text{span}\{(w_j, \gamma_0(w_j)) : j = 1, \dots, m\}$ converge (when $m \rightarrow \infty$) to $(\tilde{u}_{\varepsilon}^0, \tilde{\psi}_{\varepsilon}^0)$ in a suitable sense which will be specified below.

Lemma 5.2. *Suppose the assumptions (4)–(6). Then, for any initial condition $(u_{\varepsilon}^0, \psi_{\varepsilon}^0) \in H^1(\Omega_{\varepsilon}) \times H^{1/2}(\partial S_{\varepsilon})$ of the problem (3) such that satisfies (9), there exists a positive constant C independent of ε , such that the solution \tilde{u}_{ε} of the problem (20) satisfies*

$$\sup_{t \in [0, T]} \|\tilde{u}_{\varepsilon}(t)\|_{\tilde{\Omega}_{\varepsilon}} \leq C, \quad (33)$$

$$|\tilde{u}'_{\varepsilon}|_{\tilde{\Omega}_{\varepsilon}, T} \leq C, \quad \sqrt{\varepsilon} |\gamma_0(\tilde{u}'_{\varepsilon})|_{\partial F_{\varepsilon}, T} \leq C. \quad (34)$$

Proof. First, considering in (9) the change of variables given in (7), and taking into account that $dx = \varepsilon dx' dy_3$ and $\partial_{y_3} = \varepsilon \partial_{x_3}$, we can deduce that $(\tilde{u}_{\varepsilon}^0, \tilde{\psi}_{\varepsilon}^0) \in V_1$ such that

$$|\nabla_{\varepsilon} \tilde{u}_{\varepsilon}^0|_{\tilde{\Omega}_{\varepsilon}} \leq C. \quad (35)$$

For all $m \geq 1$, there exists $(\tilde{u}_{\varepsilon,m}^0, \gamma_0(\tilde{u}_{\varepsilon,m}^0)) \in \text{span}\{(w_j, \gamma_0(w_j)) : 1 \leq j \leq m\}$, such that the sequence $\{(\tilde{u}_{\varepsilon,m}^0, \gamma_0(\tilde{u}_{\varepsilon,m}^0))\}$ converges to $(\tilde{u}_{\varepsilon}^0, \tilde{\psi}_{\varepsilon}^0)$ in V_1 . Then, taking into account (23) and (35), we know that there exists a positive constant C such that

$$|\nabla_{\varepsilon} \tilde{u}_{\varepsilon,m}^0|_{\tilde{\Omega}_{\varepsilon}}^2 + |\tilde{u}_{\varepsilon,m}^0|_{\tilde{\Omega}_{\varepsilon}}^2 \leq C. \quad (36)$$

For each integer $m \geq 1$, we consider the sequence $\{(u_{\varepsilon,m}(t), \gamma_0(u_{\varepsilon,m}(t)))\}$ defined by (30)–(32) with these initial data. Multiplying by the derivative $\delta'_{\varepsilon m j}$ in (31), and summing from $j = 1$ to m , we obtain

$$\begin{aligned} & |(\tilde{u}'_{\varepsilon,m}(t), \gamma_0(\tilde{u}'_{\varepsilon,m}(t)))|_H^2 \\ & + \frac{1}{2} \frac{d}{dt} (\langle A_1((\tilde{u}_{\varepsilon,m}(t), \gamma_0(\tilde{u}_{\varepsilon,m}(t))), (\tilde{u}_{\varepsilon,m}(t), \gamma_0(\tilde{u}_{\varepsilon,m}(t)))) \rangle) \\ & = (f(t), \tilde{u}'_{\varepsilon,m}(t))_{\tilde{\Omega}_{\varepsilon}} + \varepsilon(g(t), \gamma_0(\tilde{u}'_{\varepsilon,m}(t)))_{\partial F_{\varepsilon}}. \end{aligned}$$

Then, we deduce

$$\begin{aligned}
 & |(\tilde{u}'_{\varepsilon,m}(t), \gamma_0(\tilde{u}'_{\varepsilon,m}(t)))|_H^2 \\
 & + \frac{1}{2} \frac{d}{dt} (\langle A_1((\tilde{u}_{\varepsilon,m}(t), \gamma_0(\tilde{u}_{\varepsilon,m}(t))), (\tilde{u}_{\varepsilon,m}(t), \gamma_0(\tilde{u}_{\varepsilon,m}(t)))) \rangle) \\
 & \leq \frac{1}{2} |f(t)|_{\omega_\varepsilon}^2 + \frac{1}{2} |\tilde{u}'_{\varepsilon,m}(t)|_{\tilde{\Omega}_\varepsilon}^2 + \varepsilon \frac{1}{2} |g(t)|_{\partial F'_\varepsilon}^2 + \varepsilon \frac{1}{2} |\gamma_0(\tilde{u}'_{\varepsilon,m}(t))|_{\partial F'_\varepsilon}^2.
 \end{aligned}$$

Integrating now between 0 and t , taking into account the definition of A_1 and (29), we obtain that

$$\begin{aligned}
 & \int_0^t |(\tilde{u}'_{\varepsilon,m}(s), \gamma_0(\tilde{u}'_{\varepsilon,m}(s)))|_H^2 ds + \frac{\min\{1, \kappa\}}{1 + \|\gamma_0\|^2} \|(\tilde{u}_{\varepsilon,m}(t), \gamma_0(\tilde{u}_{\varepsilon,m}(t)))\|_{V_1}^2 \\
 & \leq \max\{1, \kappa\} \left(|\nabla_\varepsilon \tilde{u}_{\varepsilon,m}^0|_{\tilde{\Omega}_\varepsilon}^2 + |\tilde{u}_{\varepsilon,m}^0|_{\tilde{\Omega}_\varepsilon}^2 \right) + \int_0^t (|f(s)|_{\omega_\varepsilon}^2 + \varepsilon |g(s)|_{\partial F'_\varepsilon}^2) ds,
 \end{aligned} \tag{37}$$

for all $t \in (0, T)$.

Taking into account (27) and (36) in (37), we have proved that the sequence $\{(\tilde{u}_{\varepsilon,m}, \gamma_0(\tilde{u}_{\varepsilon,m}))\}$ is bounded in $C([0, T]; V_1)$, and $\{(\tilde{u}'_{\varepsilon,m}, \gamma_0(\tilde{u}'_{\varepsilon,m}))\}$ is bounded in $L^2(0, T; H)$, for all $T > 0$.

If we work with the truncated Galerkin equations (30)-(32) instead of the full PDE, using (29), we note that the calculations of the proof of Lemma 5.1 can be following identically to show that $\{(\tilde{u}_{\varepsilon,m}, \gamma_0(\tilde{u}_{\varepsilon,m}))\}$ is bounded in $L^2(0, T; V_1)$, for all $T > 0$.

Moreover, taking into account the uniqueness of solution to (20) and using Aubin-Lions compactness lemma (e.g., cf. Lions [10]), it is not difficult to conclude that the sequence $\{(\tilde{u}_{\varepsilon,m}, \gamma_0(\tilde{u}_{\varepsilon,m}))\}$ converges weakly in $L^2(0, T; V_1)$ to the solution $(\tilde{u}_\varepsilon, \gamma_0(\tilde{u}_\varepsilon))$ to (20). Since the inclusion $H^1(\tilde{\Omega}_\varepsilon) \subset L^2(\tilde{\Omega}_\varepsilon)$ is compact and $\tilde{u}_\varepsilon \in C([0, T]; L^2(\tilde{\Omega}_\varepsilon))$, it follows using [11, Lemma 11.2] that the estimate (33) is proved.

On the other hand, the sequence $\{(\tilde{u}'_{\varepsilon,m}, \gamma_0(\tilde{u}'_{\varepsilon,m}))\}$ converges weakly in $L^2(0, T; H)$ to $(\tilde{u}'_\varepsilon, \gamma_0(\tilde{u}'_\varepsilon))$, for all $T > 0$, and using the lower-semicontinuity of the norm and (37), we get

$$\begin{aligned}
 |\tilde{u}'_\varepsilon|_{\tilde{\Omega}_\varepsilon, T}^2 + \varepsilon |\gamma_0(\tilde{u}'_\varepsilon)|_{\partial F_\varepsilon, T}^2 & \leq \liminf_{m \rightarrow \infty} \left(|\tilde{u}'_{\varepsilon,m}|_{\tilde{\Omega}_\varepsilon, T}^2 + \varepsilon |\gamma_0(\tilde{u}'_{\varepsilon,m})|_{\partial F_\varepsilon, T}^2 \right) \\
 & \leq C \liminf_{m \rightarrow \infty} \left(|\nabla_\varepsilon \tilde{u}_{\varepsilon,m}^0|_{\tilde{\Omega}_\varepsilon}^2 + |\tilde{u}_{\varepsilon,m}^0|_{\tilde{\Omega}_\varepsilon}^2 + 1 \right) \\
 & = C \left(|\nabla_\varepsilon \tilde{u}_\varepsilon^0|_{\tilde{\Omega}_\varepsilon}^2 + |\tilde{u}_\varepsilon^0|_{\tilde{\Omega}_\varepsilon}^2 + 1 \right),
 \end{aligned}$$

which, jointly with (23) and (35), implies (34). \square

5.1 The extension of \tilde{u}_ε to the whole $\Omega \times (0, T)$: definition of $\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon$

Since the solution \tilde{u}_ε of the problem (20) is defined only in $\tilde{\Omega}_\varepsilon \times (0, T)$, we need to extend it to the whole $\Omega \times (0, T)$. For finding a suitable extension into all $\Omega \times (0, T)$, we shall use the following result.

Corollary 5.3 (Corollary 3 in [2]). *There exists an extension operator $\tilde{\Pi}_\varepsilon \in \mathcal{L}(H^1_{\partial\Omega}(\tilde{\Omega}_\varepsilon); H^1_0(\Omega))$ and a positive constant C , independent of ε , such that*

$$\begin{aligned}
 \tilde{\Pi}_\varepsilon \tilde{\varphi}(x', y_3) & = \tilde{\varphi}(x', y_3), \quad \text{if } (x', y_3) \in \tilde{\Omega}_\varepsilon, \\
 |\nabla_\varepsilon \tilde{\Pi}_\varepsilon \tilde{\varphi}|_\Omega & \leq C |\nabla_\varepsilon \tilde{\varphi}|_{\tilde{\Omega}_\varepsilon}, \quad \forall \tilde{\varphi} \in H^1_{\partial\Omega}(\tilde{\Omega}_\varepsilon).
 \end{aligned}$$

Let us obtain some a priori estimates for the extension of \tilde{u}_ε to the whole $\Omega \times (0, T)$. Using Corollary 5.3 together with Lemmas 5.1-5.2, we obtain the following result.

Corollary 5.4. *Assume the assumptions in Lemma 5.2. Then, there exists an extension $\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon$ of the solution \tilde{u}_ε of the problem (20) into $\Omega \times (0, T)$, such that*

$$|\nabla_\varepsilon \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon|_{\Omega, T} \leq C, \quad |\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon|_{\Omega, T} \leq C, \quad (38)$$

$$\sup_{t \in [0, T]} \left\| \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) \right\|_{\Omega} \leq C, \quad (39)$$

$$|\tilde{\Pi}_\varepsilon \tilde{u}'_\varepsilon|_{\Omega, T} \leq C, \quad (40)$$

where the constant C does not depend on ε .

6 A compactness result

In this section, we obtain some compactness results about the behavior of the sequence $\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon$ satisfying the *a priori* estimates given in Corollary 5.4.

Due to the periodicity of the domain $\tilde{\Omega}_\varepsilon$, one has, for $\varepsilon \rightarrow 0$, that

$$\chi_{\tilde{\Omega}_\varepsilon} \xrightarrow{*} \frac{|Y'_f|}{|Y'|} \text{ weakly-star in } L^\infty(\Omega), \quad (41)$$

where the limit is the proportion of the material in the cell Y' .

Set

$$\xi_\varepsilon = \nabla_{x'} \tilde{u}_\varepsilon \quad \text{in } \tilde{\Omega}_\varepsilon \times (0, T),$$

which, by the first estimate in (24), satisfies

$$|\xi_\varepsilon|_{\tilde{\Omega}_\varepsilon, T} \leq C. \quad (42)$$

Let us denote by $\tilde{\xi}_\varepsilon$ its extension with zero to the whole of $\Omega \times (0, T)$, i.e.

$$\tilde{\xi}_\varepsilon = \begin{cases} \xi_\varepsilon & \text{in } \tilde{\Omega}_\varepsilon \times (0, T), \\ 0 & \text{in } (\Omega \setminus \tilde{\Omega}_\varepsilon) \times (0, T). \end{cases} \quad (43)$$

Proposition 6.1. *Under the assumptions in Lemma 5.2, there exists a function $u \in L^2(0, T; H_0^1(\omega))$, where u is independent of y_3 , (u will be the unique solution of the limit system (10)) and a function $\xi \in L^2(0, T; (L^2(\Omega))^3)$ such that for all $T > 0$,*

$$\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) \rightharpoonup u(t) \quad \text{weakly in } H_0^1(\Omega), \quad \forall t \in [0, T], \quad (44)$$

$$\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) \rightarrow u(t) \quad \text{strongly in } L^2(\Omega), \quad \forall t \in [0, T], \quad (45)$$

$$\tilde{\xi}_\varepsilon \rightharpoonup \xi \quad \text{weakly in } L^2(0, T; (L^2(\Omega))^3), \quad (46)$$

where $\tilde{\xi}_\varepsilon$ is given by (43).

Proof. The estimates (38) read

$$|\nabla_{x'} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon|_{\Omega, T} \leq C, \quad |\partial_{y_3} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon|_{\Omega, T} \leq C \varepsilon \quad |\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon|_{\Omega, T} \leq C. \quad (47)$$

From the first and the last estimates in (47), we can deduce that $\{\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon\}$ is bounded in $L^2(0, T; L^2(0, 1; H^1(\omega)))$, for all $T > 0$. Let us fix $T > 0$. Then, there exists a subsequence $\{\tilde{\Pi}_{\varepsilon'} \tilde{u}_{\varepsilon'}\} \subset \{\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon\}$ and function $u \in L^2(0, T; L^2(0, 1; H^1(\omega)))$ such that

$$\tilde{\Pi}_{\varepsilon'} \tilde{u}_{\varepsilon'} \rightharpoonup u \quad \text{weakly in } L^2(0, T; L^2(0, 1; H^1(\omega))), \quad (48)$$

which implies

$$\partial_{y_3} \tilde{\Pi}_\varepsilon \tilde{u}_{\varepsilon'} \rightharpoonup \partial_{y_3} u \quad \text{weakly in } L^2(0, T; H^{-1}(0, 1; H^1(\omega))). \quad (49)$$

By the second estimate in (47), we can deduce that $\partial_{y_3} \tilde{\Pi}_\varepsilon \tilde{u}_{\varepsilon'}$ tends to zero in $L^2(0, T; L^2(\Omega))$. From (49) and the uniqueness of the limit, we have that $\partial_{y_3} u = 0$, which implies that u does not depend on y_3 .

By the estimate (39), for each $t \in [0, T]$, we have that $\{\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t)\}$ is bounded in $H_0^1(\Omega)$, and since we have (48), we can deduce

$$\tilde{\Pi}_\varepsilon \tilde{u}_{\varepsilon'}(t) \rightharpoonup u(t) \quad \text{weakly in } H_0^1(\Omega), \quad \forall t \in [0, T].$$

By the estimate (40), we see that the sequence $\{\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon'\}$ is bounded in $L^2(0, T; L^2(\Omega))$, for all $T > 0$. Then, we have that $\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) : [0, T] \rightarrow L^2(\Omega)$ is an equicontinuous family of functions.

Since, for each $t \in [0, T]$, we have that $\{\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t)\}$ is bounded in $H_0^1(\Omega)$, the compact embedding $H_0^1(\Omega) \subset L^2(\Omega)$, implies that it is precompact in $L^2(\Omega)$. Then, applying the Ascoli-Arzelà Theorem, we deduce that $\{\tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t)\}$ is a precompact sequence in $C([0, T]; L^2(\Omega))$. Hence, since we have (48), we can deduce that

$$\tilde{\Pi}_\varepsilon \tilde{u}_{\varepsilon'} \rightarrow u \quad \text{strongly in } C([0, T]; L^2(\Omega)),$$

and, in particular, we have

$$\tilde{\Pi}_\varepsilon \tilde{u}_{\varepsilon'}(t) \rightarrow u(t) \quad \text{strongly in } L^2(\Omega), \quad \forall t \in [0, T].$$

Finally, from the estimate (42) and (43), we have $|\tilde{\xi}_\varepsilon|_{\Omega, T} \leq C$, and hence, up a sequence, there exists $\xi \in L^2(0, T, (L^2(\Omega))^3)$ such that $\tilde{\xi}_{\varepsilon'} \rightharpoonup \xi$ weakly in $L^2(0, T; (L^2(\Omega))^3)$.

By the uniqueness of solution of the limit problem (10), we deduce that the above convergences hold for the whole sequence and therefore, by the arbitrariness of $T > 0$, all the convergences are satisfied, as we wanted to prove. \square

7 Homogenized model: proof of Theorem 1.1

In this section, we identify the homogenized model.

Let $v \in \mathcal{D}(\Omega)$ be a test function in (21), with v independent of y_3 . Then one has

$$\begin{aligned} & \frac{d}{dt} \left(\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) v dx' dy_3 \right) + \varepsilon \frac{d}{dt} \left(\int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon(t)) v d\sigma(x') dy_3 \right) + \int_{\Omega} \tilde{\xi}_\varepsilon \cdot \nabla_{x'} v dx' dy_3 \\ & + \kappa \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) v dx' dy_3 = \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} f(t) v dx' dy_3 + \varepsilon \int_{\partial F_\varepsilon} g(t) v d\sigma(x') dy_3, \end{aligned}$$

in $\mathcal{D}'(0, T)$.

We consider $\varphi \in C_c^1([0, T])$ such that $\varphi(T) = 0$ and $\varphi(0) \neq 0$. Multiplying by φ and integrating between 0 and T , we have

$$\begin{aligned} & -\varphi(0) \left(\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(0) v dx' dy_3 \right) - \int_0^T \frac{d}{dt} \varphi(t) \left(\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) v dx' dy_3 \right) dt \\ & - \varepsilon \varphi(0) \left(\int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon(0)) v d\sigma(x') dy_3 \right) - \varepsilon \int_0^T \frac{d}{dt} \varphi(t) \left(\int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon(t)) v d\sigma(x') dy_3 \right) dt \\ & + \int_0^T \varphi(t) \int_{\Omega} \tilde{\xi}_\varepsilon \cdot \nabla_{x'} v dx' dy_3 dt + \kappa \int_0^T \varphi(t) \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) v dx' dy_3 dt \\ & = \int_0^T \varphi(t) \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} f(t) v dx' dy_3 dt + \varepsilon \int_0^T \varphi(t) \int_{\partial F_\varepsilon} g(t) v d\sigma(x') dy_3 dt. \end{aligned} \quad (50)$$

For the sake of clarity, we split the proof in five parts. Firstly, we analyze the integrals on Ω , where we only require to use Proposition 6.1 and the convergence (41). Secondly, for the integrals on the boundary of the cylinders we make use of a convergence result based on a technique introduced by Vanninathan [14]. In the third step, we pass to the limit, as $\varepsilon \rightarrow 0$ in order to get the limit equation satisfied by u . In the fourth step, we identify $\int_0^1 \xi dy_3$ making use of the solutions of the cell-problems (12) and, finally we prove that u is uniquely determined.

Step 1. In this step, we analyze all the integrals on Ω . Passing to the limit, as $\varepsilon \rightarrow 0$, in the integrals on Ω and taking into account that u is independent of y_3 :

From (41) and (45), we have, for $\varepsilon \rightarrow 0$,

$$\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) v dx' dy_3 \rightarrow \frac{|Y'_f|}{|Y'|} \int_{\omega} u(t) v dx',$$

which integrating in time and using Lebesgue's Dominated Convergence Theorem, gives

$$\int_0^T \frac{d}{dt} \varphi(t) \left(\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) v dx' dy_3 \right) dt \rightarrow \frac{|Y'_f|}{|Y'|} \int_0^T \frac{d}{dt} \varphi(t) \left(\int_{\omega} u(t) v dx' \right) dt,$$

and

$$\kappa \int_0^T \varphi(t) \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) v dx' dy_3 dt \rightarrow \kappa \frac{|Y'_f|}{|Y'|} \int_0^T \varphi(t) \int_{\omega} u(t) v dx' dt.$$

By (41) and (45), we have

$$\varphi(0) \left(\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(0) v dx' dy_3 \right) \rightarrow \varphi(0) \frac{|Y'_f|}{|Y'|} \int_{\omega} u(0) v dx'.$$

By the assumption (6), (41) and using Lebesgue's Dominated Convergence Theorem, we get

$$\int_0^T \varphi(t) \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} f(x', t) v dx' dy_3 dt \rightarrow \frac{|Y'_f|}{|Y'|} \int_0^T \varphi(t) \int_{\omega} f(x', t) v dx' dt.$$

On the other hand, using (46), we obtain, for $\varepsilon \rightarrow 0$

$$\int_0^T \varphi(t) \int_{\Omega} \tilde{\xi}_\varepsilon \cdot \nabla_{x'} v dx' dy_3 dt \rightarrow \int_0^T \varphi(t) \int_{\Omega} \xi \cdot \nabla_{x'} v dx' dy_3 dt.$$

Step 2. Passing to the limit, as $\varepsilon \rightarrow 0$, in the surface integrals on the boundary of the cylinders and taking into account that u is independent of y_3 :

We make use of the technique introduced by Vanninathan [14] for the Steklov problem which transforms surface integrals into volume integrals. This technique was already used as a main tool to homogenize the non homogeneous Neumann problem for the elliptic case by Cioranescu and Donato [5].

Following the technical lemmas contained in [5, Section 3], let us introduce, for any $h \in L^{s'}(\partial F)$, $1 \leq s' \leq \infty$, the linear form μ_h^ε on $W_0^{1,s}(\Omega)$ defined by

$$\langle \mu_h^\varepsilon, \varphi \rangle = \varepsilon \int_{\partial F_\varepsilon} h(x'/\varepsilon, y_3) \varphi d\sigma(x') dy_3, \quad \forall \varphi \in W_0^{1,s}(\Omega),$$

with $1/s + 1/s' = 1$, such that

$$\mu_h^\varepsilon \rightarrow \mu_h \quad \text{strongly in } (W_0^{1,s}(\Omega))', \quad (51)$$

where $\langle \mu_h, \varphi \rangle = \mu_h \int_{\Omega} \varphi dx' dy_3$, with

$$\mu_h = \frac{1}{|Y'|} \int_{\partial F} h(y) d\sigma(y) dy_3.$$

In the particular case in which $h \in L^\infty(\partial F)$ or even when h is constant, we have

$$\mu_h^\varepsilon \rightarrow \mu_h \quad \text{strongly in } W^{-1,\infty}(\Omega).$$

In what follows, we shall denote by μ_1^ε the above introduced measure in the particular case in which $h = 1$. Notice that in this case μ_h becomes $\mu_1 = |\partial F'|/|Y'|$.

Observe that using [6, Corollary 4.2] with (44), we can deduce, for $\varepsilon \rightarrow 0$,

$$\varepsilon \int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon(t)) v d\sigma(x') dy_3 = \langle \mu_1^\varepsilon, \tilde{\Pi}_\varepsilon \tilde{u}_{\varepsilon|\tilde{\Omega}_\varepsilon}(t)v \rangle \rightarrow \mu_1 \int_\omega u(t) v dx' = \frac{|\partial F'|}{|Y'|} \int_\omega u(t) v dx',$$

which integrating in time and using Lebesgue's Dominated Convergence Theorem, gives

$$\varepsilon \int_0^T \frac{d}{dt} \varphi(t) \left(\int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon(t)) v d\sigma(x') dy_3 \right) dt \rightarrow \frac{|\partial F'|}{|Y'|} \int_0^T \frac{d}{dt} \varphi(t) \left(\int_\omega u(t) v dx' \right) dt.$$

Moreover, using [6, Corollary 4.2] with (44), we can deduce, for $\varepsilon \rightarrow 0$,

$$\varepsilon \int_{\partial F_\varepsilon} \gamma_0(u_\varepsilon(0)) v d\sigma(x') dy_3 = \langle \mu_1^\varepsilon, \tilde{\Pi}_\varepsilon \tilde{u}_{\varepsilon|\tilde{\Omega}_\varepsilon}(0)v \rangle \rightarrow \mu_1 \int_\omega u(0) v dx' = \frac{|\partial F'|}{|Y'|} \int_\omega u(0) v dx'.$$

On the other hand, note that using (51) with $s = 2$, taking into account (6) and by Lebesgue's Dominated Convergence Theorem, we can deduce, for $\varepsilon \rightarrow 0$,

$$\varepsilon \int_0^T \varphi(t) \int_{\partial F_\varepsilon} g(x', t) v d\sigma(x') dy_3 dt = \int_0^T \varphi(t) \langle \mu_1^\varepsilon, g(x', t)v \rangle dt \rightarrow \frac{|\partial F'|}{|Y'|} \int_0^T \varphi(t) \int_\omega g(x', t) v dx' dt.$$

Step 3. Passing to the limit, as $\varepsilon \rightarrow 0$, in (50): all the terms in (50) pass to the limit, as $\varepsilon \rightarrow 0$, and therefore taking into account the previous steps, we get

$$\begin{aligned} & -\varphi(0) \left(\frac{|Y'_f|}{|Y'|} + \frac{|\partial F'|}{|Y'|} \right) \int_\omega u(0) v dx' - \left(\frac{|Y'_f|}{|Y'|} + \frac{|\partial F'|}{|Y'|} \right) \int_0^T \frac{d}{dt} \varphi(t) \int_\omega u(t) v dx' dt \\ & + \int_0^T \varphi(t) \int_\Omega \xi \cdot \nabla_{x'} v dx' dy_3 dt + \kappa \frac{|Y'_f|}{|Y'|} \int_0^T \varphi(t) \int_\omega u(t) v dx' dt \\ & = \frac{|Y'_f|}{|Y'|} \int_0^T \varphi(t) \int_\omega f(t) v dx' dt + \frac{|\partial F'|}{|Y'|} \int_0^T \varphi(t) \int_\omega g(t) v dx' dt. \end{aligned}$$

Hence, $\int_0^1 \xi dy_3$ verifies

$$\left(\frac{|Y'_f|}{|Y'|} + \frac{|\partial F'|}{|Y'|} \right) \frac{\partial u}{\partial t} - \operatorname{div}_{x'} \left(\int_0^1 \xi dy_3 \right) + \frac{|Y'_f|}{|Y'|} \kappa u = \frac{|Y'_f|}{|Y'|} f + \frac{|\partial F'|}{|Y'|} g, \quad (52)$$

in $\omega \times (0, T)$.

Step 4. It remains now to identify $\int_0^1 \xi dy_3$. We shall make use of the solutions of the cell problems (12). For any fixed $i = 1, 2$, let us define

$$\Psi_{i\varepsilon}(x') = \varepsilon \left(w_i \left(\frac{x'}{\varepsilon} \right) + y'_i \right) \quad \forall x' \in \omega_\varepsilon, \quad (53)$$

where $y' = x'/\varepsilon$.

By periodicity

$$P_\varepsilon \Psi_{i\varepsilon} \rightharpoonup x'_i \quad \text{weakly in } H^1(\omega),$$

where $P_\varepsilon \Psi_{i\varepsilon}$ denotes the extension to ω given in [7, Lemma 1]. Then, by Rellich-Kondrachov Theorem, we can deduce

$$P_\varepsilon \Psi_{i\varepsilon} \rightarrow x'_i \quad \text{strongly in } L^2(\omega). \quad (54)$$

Let $\nabla_{x'} \Psi_{i\varepsilon}$ be the gradient of $\Psi_{i\varepsilon}$ in ω_ε . Denote by $\widetilde{\nabla_{x'} \Psi_{i\varepsilon}}$ the extension by zero of $\nabla_{x'} \Psi_{i\varepsilon}$ inside the holes. From (53), we have

$$\widetilde{\nabla_{x'} \Psi_{i\varepsilon}} = \nabla_{y'} (\widetilde{w_i + y'_i}) = \widetilde{\nabla_{y'} w_i(y')} + e_i \chi_{Y'_f},$$

and taking into account [3, Corollary 2.10], we have

$$\widetilde{\nabla_{x'} \Psi_{i\varepsilon}} \rightharpoonup \frac{1}{|Y'_f|} \int_{Y'_f} (e_i + \nabla_{y'} w_i(y')) dy' \quad \text{weakly in } L^2(\omega). \quad (55)$$

On the other hand, it is not difficult to see that $\Psi_{i\varepsilon}$ satisfies

$$\begin{cases} -\operatorname{div}_{x'} (\nabla_{x'} \Psi_{i\varepsilon}) = 0, & \text{in } \omega_\varepsilon, \\ \nabla_{x'} \Psi_{i\varepsilon} \cdot \nu'_\varepsilon = 0, & \text{on } \partial F'_\varepsilon. \end{cases} \quad (56)$$

Let $v \in \mathcal{D}(\Omega)$ with v independent of y_3 . Multiplying the first equation in (56) by $v \tilde{u}_\varepsilon$ and integrating by parts over ω_ε , we get

$$\int_{\omega_\varepsilon} \nabla_{x'} \Psi_{i\varepsilon} \cdot \nabla_{x'} v \tilde{u}_\varepsilon dx' + \int_{\omega_\varepsilon} \nabla_{x'} \Psi_{i\varepsilon} \cdot \nabla_{x'} \tilde{u}_\varepsilon v dx' = 0. \quad (57)$$

On the other hand, we multiply system (20) by the test function $v \Psi_{i\varepsilon}$ and integrating by parts over $\tilde{\Omega}_\varepsilon$, we obtain

$$\begin{aligned} & \frac{d}{dt} \left(\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 \right) + \varepsilon \frac{d}{dt} \left(\int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon) v \gamma_0(\Psi_{i\varepsilon}) d\sigma(x') dy_3 \right) \\ & + \int_{\tilde{\Omega}_\varepsilon} \nabla_{x'} \tilde{u}_\varepsilon \cdot \nabla_{x'} v \Psi_{i\varepsilon} dx' dy_3 + \int_{\tilde{\Omega}_\varepsilon} \nabla_{x'} \tilde{u}_\varepsilon \cdot \nabla_{x'} \Psi_{i\varepsilon} v dx' dy_3 + \kappa \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 \\ & = \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} f(x', t) v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 + \varepsilon \int_{\partial F_\varepsilon} g(x', t) v \gamma_0(\Psi_{i\varepsilon}) d\sigma(x') dy_3, \end{aligned} \quad (58)$$

in $\mathcal{D}'(0, T)$.

Using (57) in (58), we have

$$\begin{aligned} & \frac{d}{dt} \left(\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 \right) + \varepsilon \frac{d}{dt} \left(\int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon) v \gamma_0(\Psi_{i\varepsilon}) d\sigma(x') dy_3 \right) \\ & + \int_{\Omega} \tilde{\xi}_\varepsilon \cdot \nabla_{x'} v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 - \int_{\Omega} \widetilde{\nabla_{x'} \Psi_{i\varepsilon}} \cdot \nabla_{x'} v \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon dx' dy_3 + \kappa \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 \\ & = \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} f(x', t) v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 + \varepsilon \int_{\partial F_\varepsilon} g(x', t) v \gamma_0(\Psi_{i\varepsilon}) d\sigma(x') dy_3, \end{aligned}$$

in $\mathcal{D}'(0, T)$.

We consider $\varphi \in C_c^1([0, T])$ such that $\varphi(T) = 0$ and $\varphi(0) \neq 0$. Multiplying by φ and integrating between 0

and T , we have

$$\begin{aligned}
 & -\varphi(0) \left(\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(0) v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 \right) - \int_0^T \frac{d}{dt} \varphi(t) \left(\int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 \right) dt \\
 & -\varepsilon \varphi(0) \left(\int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon(0)) v \gamma_0(\Psi_{i\varepsilon}) d\sigma(x') dy_3 \right) - \varepsilon \int_0^T \frac{d}{dt} \varphi(t) \left(\int_{\partial F_\varepsilon} \gamma_0(\tilde{u}_\varepsilon(t)) v \gamma_0(\Psi_{i\varepsilon}) d\sigma(x') dy_3 \right) dt \\
 & \quad + \int_0^T \varphi(t) \int_{\Omega} \tilde{\xi}_\varepsilon \cdot \nabla_{x'} v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 dt - \int_0^T \varphi(t) \int_{\Omega} \widetilde{\nabla_{x'} \Psi_{i\varepsilon}} \cdot \nabla_{x'} v \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon dx' dy_3 dt \\
 & \quad + \kappa \int_0^T \varphi(t) \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon(t) v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 dt \\
 & = \int_0^T \varphi(t) \int_{\Omega} \chi_{\tilde{\Omega}_\varepsilon} f(x', t) v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 dt + \varepsilon \int_0^T \varphi(t) \int_{\partial F_\varepsilon} g(x', t) v \gamma_0(\Psi_{i\varepsilon}) d\sigma(x') dy_3 dt.
 \end{aligned} \tag{59}$$

Now, we have to pass to the limit, as $\varepsilon \rightarrow 0$. We will focus on the terms which involve the gradient. Taking into account (54), we reason as in steps 1 and 2 for the others terms.

Firstly, using (46) and (54), we have

$$\int_0^T \varphi(t) \int_{\Omega} \tilde{\xi}_\varepsilon \cdot \nabla_{x'} v P_\varepsilon \Psi_{i\varepsilon} dx' dy_3 dt \rightarrow \int_0^T \varphi(t) \int_{\Omega} \xi \cdot \nabla_{x'} v x'_i dx' dy_3 dt,$$

and by (45), (55) and Lebesgue's Dominated Convergence Theorem, we obtain

$$\int_0^T \varphi(t) \int_{\Omega} \widetilde{\nabla_{x'} \Psi_{i\varepsilon}} \cdot \nabla_{x'} v \tilde{\Pi}_\varepsilon \tilde{u}_\varepsilon dx' dy_3 dt \rightarrow \frac{1}{|Y'|} \int_0^T \varphi(t) \int_{\omega} \left(\int_{Y'_f} (e_i + \nabla_{y'} w_i) dy' \right) \cdot \nabla_{x'} v u dx' dt.$$

Therefore, when we pass to the limit in (59), we obtain

$$\begin{aligned}
 & -\varphi(0) \left(\frac{|Y'_f|}{|Y'|} + \frac{|\partial F'|}{|Y'|} \right) \left(\int_{\omega} u(0) v x'_i dx' \right) - \left(\frac{|Y'_f|}{|Y'|} + \frac{|\partial F'|}{|Y'|} \right) \int_0^T \frac{d}{dt} \varphi(t) \int_{\omega} u(t) v x'_i dx' dt \\
 & + \int_0^T \varphi(t) \int_{\Omega} \xi \cdot \nabla_{x'} v x'_i dx' dy_3 dt - \frac{1}{|Y'|} \int_0^T \varphi(t) \int_{\omega} \left(\int_{Y'_f} (e_i + \nabla_{y'} w_i) dy' \right) \cdot \nabla_{x'} v u(t) dx' dt \\
 & \quad + \kappa \frac{|Y'_f|}{|Y'|} \int_0^T \varphi(t) \int_{\omega} u(t) v x'_i dx' dt \\
 & = \frac{|Y'_f|}{|Y'|} \int_0^T \varphi(t) \int_{\omega} f(t) v x'_i dx' dt + \frac{|\partial F'|}{|Y'|} \int_0^T \varphi(t) \int_{\omega} g(t) v x'_i dx' dt.
 \end{aligned}$$

Using Green's formula and equation (52), we have

$$- \int_0^T \varphi(t) \int_{\Omega} \xi \cdot \nabla_{x'} v x'_i dx' dy_3 dt + \frac{1}{|Y'|} \int_0^T \varphi(t) \int_{\omega} \left(\int_{Y'_f} (e_i + \nabla_{y'} w_i) dy' \right) \cdot \nabla_{x'} v u dx' dt = 0.$$

The above equality holds true for any $v \in \mathcal{D}(\Omega)$ independent of y_3 , and $\varphi \in C_c^1([0, T])$. This implies that

$$- \left(\int_0^1 \xi dy_3 \right) \cdot \nabla_{x'} v x'_i + \frac{1}{|Y'|} \left(\int_{Y'_f} (e_i + \nabla_{y'} w_i) dy' \right) \cdot \nabla_{x'} v u = 0, \quad \text{in } \omega \times (0, T).$$

We conclude that

$$\operatorname{div}_{x'} \left(\int_0^1 \xi dy_3 \right) = \operatorname{div}_{x'} (Q \nabla_{x'} u), \quad (60)$$

where $Q = ((q_{ij}))$, $1 \leq i, j \leq 2$, is given by

$$q_{ij} = \frac{1}{|Y_f'|} \int_{Y_f'} (e_i + \nabla_{y'} w_i) \cdot e_j dy'.$$

Observe that if we multiply system (12) by the test function w_j , integrating by parts over Y_f' , we obtain

$$\int_{Y_f'} (e_i + \nabla_{y'} w_i) \cdot \nabla_{y'} w_j dy' = 0,$$

then we conclude that q_{ij} is given by (11).

Step 5. Finally, thanks to (52) and (60), we observe that u satisfies the first equation in (10). A weak solution of (10) is any function u , satisfying

$$u \in C([0, T]; L^2(\omega)), \quad \text{for all } T > 0,$$

$$u \in L^2(0, T; H_0^1(\omega)), \quad \text{for all } T > 0,$$

$$\left(\frac{|Y_f'|}{|Y'|} + \frac{|\partial F'|}{|Y'|} \right) \frac{d}{dt} (u(t), v) + (Q \nabla_{x'} u(t), \nabla_{x'} v) + \frac{|Y_f'|}{|Y'|} \kappa(u(t), v) = \frac{|Y_f'|}{|Y'|} (f(t), v) + \frac{|\partial F'|}{|Y'|} (g(t), v), \quad \text{in } \mathcal{D}'(0, T),$$

for all $v \in H_0^1(\omega)$, and

$$u(0) = u_0.$$

Due to that the homogenized matrix Q is positive-definite (see [5, Theorem 4.7]), applying a slight modification of [10, Chapter 2, Theorem 1.4], we obtain that the problem (10) has a unique solution, and therefore Theorem 1.1 is proved.

References

- [1] M. Anguiano, Existence, uniqueness and homogenization of nonlinear parabolic problems with dynamical boundary conditions in perforated media, *Mediterr. J. Math.*, (2020) **17**:18.
- [2] M. Anguiano and F.J. Suárez-Grau, Newtonian fluid flow in a thin porous medium with non-homogeneous slip boundary conditions, *Networks and Heterogeneous Media*, **14**, No. 2 (2019), 289–316.
- [3] D. Cioranescu, A. Damlamian and G. Griso, The periodic unfolding method in homogenization, *SIAM J. Math. Anal.*, **40**, No. 4 (2008), 1585–1620.
- [4] D. Cioranescu, A. Damlamian and G. Griso, *The periodic unfolding method: theory and applications to partial differential problems*, Series in Contemporary Mathematics, **3**, Springer, Singapore, 2018.
- [5] D. Cioranescu, P. Donato, Homogénéisation du problème du Neumann non homogène dans des ouvert perforés, *Asymptotic Analysis* **1** (1988), 115-138.
- [6] D. Cioranescu, P. Donato, H. Ene, Homogenization of the Stokes problem with non homogeneous slip boundary conditions, *Mathematical Methods in Applied Sciences* **19** (1996), 857-881.
- [7] D. Cioranescu, J. Saint Jean Paulin, Homogenization in open sets with holes, *J. Math. Anal. Appl.* **71** (1979), 590-607.

- [8] C. Conca, J.I. Díaz, C. Timofte Effective chemical processes in porous media, *Math. Models Methods Appl. Sci.*, **13**(10) (2003), 1437-1462.
- [9] C. Conca, P. Donato Non homogeneous Neumann problems in domains with small holes, *RAIRO Modél. Math. Anal. Numér.*, **22** (1988), 561-607.
- [10] J. L. Lions, *Quelques Méthodes de Résolution des Problèmes aux Limites Non linéaires*. Dunod, 1969.
- [11] J.C. Robinson, *Infinite-dimensional dynamical systems*. Cambridge University Press, 2001.
- [12] C. Timofte, Parabolic problems with dynamical boundary conditions in perforated media, *Mathematical Modelling and Analysis* **8**(4) (2003), 337-350.
- [13] L. Tartar, *Problèmes d'homogénéisation dans les équations aux dérivées partielles*, In: Cours Peccot, Collège de France, 1977.
- [14] M. Vanninathan, Homogenization of eigenvalues problems in perforated domains, *Proc. Indian Acad. of Sciences* **90** (1981), 239-271.