How maintenance and restoration measures mediate the response of riparian plant functional composition to environmental gradients on channel margins: Insights from a highly degraded large river.

Philippe Janssen^{1,2*}, Hervé Piégay¹, Bernard Pont³, André Evette²

¹ University of Lyon, UMR 5600 Environnement Ville société, CNRS, Site of ENS Lyon, F-69342 Lyon Cedex 07, France

² University Grenoble-Alpes, National Research Institute of Science and Technology for Environment and Agriculture (IRSTEA), UR LESSEM, 2 rue de la Papeterie BP 76, F-38402 Saint-Martin-d'Hères, France

³ Réserves Naturelles de France, 6 bis rue de la Gouge, BP 100, 21803 Quétigny Cedex, France

* Corresponding author, e-mail philippe.janssen@irstea.fr, phone +33 476762879

Email addresses of other authors:

H. Piégay (herve.piegay@ens-lyon.fr), B. Pont (bernard.pont@cen-isere.org), A. Evette (andre.evette@irstea.fr)

Running head: Human activity shapes riparian plants

ABSTRACT

Riparian habitats are transitional zones where strong environmental gradients shape community. To prevent flood risks and channel migration on managed rivers, civil engineering techniques have been widely used. Recently, ecological restoration of rivers has become a major issue. However, given the alteration of natural disturbance regimes induced by human infrastructures, the real added-value of these restoration actions is questionable. Thus, a major challenge is to better understand whether changes in abiotic conditions induced by human activity influence the response of plant communities to environmental gradients. Studying a highly degraded large river, we evaluated the effect of the elevation and soil texture gradients on plant functional composition and assessed whether human-mediated environment gradients, achieved through maintenance and restoration measures, shape community structure. In the summer of 2017, we sampled 17 geomorphic surfaces, mostly gravel bars, along the Rhône River and its tributaries that were either repeatedly cleared (brush clearing vs plowing), newly reprofiled or naturally rejuvenated by high flows. The results show shifts in trait values with elevation and convergence in plant traits with increasing proportion of fine sediments. The co-occurrence of species with contrasting traits was higher in highly disturbed environments, revealing the importance of rejuvenation processes. However, the influence of both environmental gradients was mediated by human activity. For maintenance measures, plowing was better able to promote species diversity and limited biotic homogenization along environmental gradients. Among the three geomorphic surfaces, naturally rejuvenated bars were the most stressful environments, hosting distinct functional assemblages, while communities on newly reprofiled banks were in the same ecological trajectories as repeatedly cleared bars. To promote an effective ecological restoration of riparian zones, (i) a greater variability of the minimum flow is needed, (ii)

45 bedload transport restoration should be a priority and (iii) reprofiled banks should better
46 mimic the landforms of natural river margins.

47

48 **KEY WORDS**

49 Biodiversity conservation, community structure, ecological restoration, environmental
50 gradient, trait-based approach, riparian zone

1. INTRODUCTION

At the interface between terrestrial and freshwater habitats, riparian zones represent a narrow portion of the landscape with disproportionate ecological importance, extending largely beyond their area (González et al., 2017). Indeed, riparian zones are dynamic ecosystems where the joint action of the flow and sediment regimes lead to great variability in habitat geomorphology (Naiman et al., 2005; Poff et al., 1997), which induce rapid changes in environmental conditions along the elevation gradient (Naiman and Decamps, 1997). Thus, from the lower to the upper part of banks, variations in soil moisture and inundation regime drives plant species turnover, i.e., from anoxic-tolerant species to species that tolerate periodic soil desiccation, and shape functional composition, i.e., from ruderal life strategy to competitor life strategy (Bejarano et al., 2017; Dawson et al., 2017; McCoy-Sulentic et al., 2017). Beside variations in species and traits composition along the elevation gradient, it has been shown that variations in soil texture are also of particular importance in explaining the functional composition of riparian plant communities (Kyle and Leishman, 2009; Xiong et al., 2001). Indeed, at low elevation level, an increase in the proportion of fine sediments may facilitate the establishment success of competitive species, leading to different patterns of species distribution along the flood gradient. The joint action of both the elevation and the soil texture gradients thus appears to have great importance in shaping riparian biodiversity. Worldwide riparian zones have been impacted by human activity (Feld et al., 2011; Nilsson et al., 2005). Civil engineering techniques have been widely used to control flood risks and prevent channel migration, which has greatly contributed to altering the flow and sediment regimes and, consequently, the ecological integrity of riparian zones (Poff and Zimmerman, 2010). Moreover, in highly modified large rivers, vegetation encroachment is a significant adjustment process (Comiti et al., 2011; Liébault and Piégay, 2002), and it is sometimes prevented through vegetation maintenance and clearing to maintain wide low-roughness

76 channels and control flooding risks (Shields and Nunnally, 1984). All together, these human-
77 induced changes have greatly altered the composition of riparian plant communities, leading
78 to a loss in biodiversity (Bejarano et al., 2017; Biswas and Mallik, 2010; Poff et al., 2007).
79 Specifically, by limiting rejuvenation processes, i.e., bedload supply and bank erosion due
80 related to flood events as well as reduction of peak flow magnitude and frequency, these
81 regulation measures have promoted a greater uniformity of environmental conditions with the
82 risk of increased the taxonomic and functional similarities of riparian communities over times
83 (i.e., biotic homogenization, Olden and Rooney, 2006).

84 To improve habitat and water quality along riparian zones, restoration efforts have become a
85 major issue in European countries (Gumiero et al., 2013). Through bank reprofiling following
86 dike and riprap removal or set-back, river managers aimed to recreate young patches and
87 potentially stimulate alluviation/rejuvenation processes within the riverbed (Biron et al., 2014;
88 Rohde et al., 2005). These measures are expected to increase riparian habitat heterogeneity
89 (e.g., substrates, resources) and allows for a greater diversity of plant species. Moreover,
90 despite their primary hydraulic objectives, maintenance measures are often viewed by river
91 managers as a way to increase the ecological value of riparian zones, e.g., by favoring
92 ecological succession and maintaining pioneer communities. However, given the irreversible
93 alteration of the natural disturbance regimes of most rivers and modifications to biotic and
94 abiotic conditions induced by human infrastructures (Poff et al., 1997; Richardson et al.,
95 2007), the real ecological added value of these actions is questionable. Indeed, ecological
96 communities of degraded riparian zones may have already been oriented toward transient or
97 new states (Bunn and Arthington 2002, Richardson et al. 2007), making the return to the
98 reference state difficult or impossible (Hobbs et al. 2009). From this view, it is thus of great
99 concern to better understand whether maintenance and restoration measures influence the
100 response of riparian plant communities to environmental gradients.

Here, we aimed to study the effect of the elevation and soil texture gradients on riparian plant functional composition. The elevation gradient was considered a proxy for flood duration, which is hypothesized to be the main disturbance controlling plant distributions in riparian zones (e.g., Fraaije et al., 2015; McCoy-Sulentic et al., 2017). The soil texture gradient was considered a proxy for soil moisture and fertility, which is hypothesized to be of great importance in structuring the functional composition of riparian plant communities (e.g., Kyle and Leishman, 2009; Xiong et al., 2001). Using a highly degraded river system, the Rhône River, as a case study, we further aimed to understand how human-mediated environmental gradients shape community structure. This result was achieved by evaluating the added value of maintenance and restoration measures applied along the Rhône River for the conservation of riparian biodiversity.

To date, few studies have used a functional approach to study assemblage patterns of riparian plants (Kyle and Leishman, 2009), particularly in the field of riparian vegetation restoration (González et al., 2015). However, unlike classical measures of species diversity, species traits capture different aspects of species' resource use and habitat requirements and are thus useful measures in unraveling complex patterns linking environmental changes, assemblage structures and ecosystem processes (Lavorel et al., 2008). By moving beyond species identity, trait-based approaches also enable comparisons of communities across systems (Cadotte et al., 2011), which allows to better evaluate successes or failures in restoration projects and thus better generalize research results. Moreover, in riparian zones, most of the previous studies using a functional approach have focused on a single environmental gradient, the flood gradient (Bejarano et al., 2017; McCoy-Sulentic et al., 2017), with little to no attention to the effects of interactions between it and other environmental gradients and/or human-mediated environmental gradients (but see, Brummer et al., 2016; Dawson et al., 2017). To accurately

define guidelines for the management and restoration of degraded riparian zones, this information is essential.

Specifically, we tested whether the mean and dispersion of selected trait values were influenced by the elevation and soil texture gradients as well as by maintenance and restoration measures on a range of riparian geomorphic surfaces (i.e., from gravel bars to channel banks). Analytical strategies that combine measures of the mean and dispersion of traits within a given species assemblage have been shown to describe two complementary aspects of the relationship between community structure and ecosystem functioning (de Bello et al., 2013; Ricotta and Moretti, 2011): (i) shifts in trait values due to environmental selection and (ii) patterns of trait convergence or divergence due to niche differentiation. Taking advantage of the large trait databases available currently (Kattge et al., 2011; Kleyer et al., 2008), we focused on three sets of traits widely used in riparian plant community studies (e.g., Bejarano et al., 2017; Dawson et al., 2017; Kyle and Leishman, 2009) and describing crucial dimensions of plant growth requirements as well as major axes of plant functional strategies (Ellenberg et al., 1992; Westoby, 1998). We then assessed the response of plant functional composition to environmental gradients among three types of geomorphic surfaces – repeatedly cleared, i.e., relict bars along the Rhône River maintained by brush clearing or plowing, newly reprofiled, i.e., banks reprofiled following dike removal and naturally rejuvenated, i.e., reference river margins, – and addressed the following two questions: (i) how do elevation and soil texture gradients shape riparian plant functional composition along a highly degraded large River?; (ii) How does human activities, achieved through maintenance and restoration measures, influence the response of plant functional composition to environmental gradients?

2. MATERIALS & METHODS

2.1. Study area and experimental design

The study was carried out along the Rhône River (total length = 810 km, catchment area = 96,500 km², mean annual discharge = 1,700 m³/s) in its French middle reach (approximately 250 km long) and three tributary rivers (Ain, Drôme and Doux Rivers, Figure 1). This area is characterized by a temperate climate with mean annual temperatures and precipitations of 13.6°C and 755 mm in the southern part and 11.6°C and 815 mm in the northern part. Within the study area, the Rhône River has been highly modified due to two historical development phases. During the 19th century, a rectification phase characterized by the building of a vast system of longitudinal submersible and transversal dikes in the main channel was carried out to facilitate navigation. In the second half of the 20th century, a derivation phase characterized by the construction of a series of lateral canals, running parallel to the natural Rhône River channel, was carried out to produce hydropower and facilitate irrigation and navigation. The bypassed sections, corresponding nowadays to part of the historical Rhône River channel, have been significantly impacted by the two phases, including changes in hydrological regime, channel dewatering, incision and lateral stabilization. They are characterized by a minimum flow that is observed most of the year, until the hydropower plant capacity is reached. Above this threshold, flood peaks are conveyed in the bypassed sections. Due to the peak-flow decrease, bedload supply interruption and associated bedload coarsening following gravel winnowing and bed armouring, no bedload transport is observed along most of these channels. Taken together, these hydro-morphological modifications have profoundly impoverished aquatic and terrestrial habitats. Since 2000, bypassed section have been subject to an ambitious ecological restoration program that mainly involves cut-off secondary channel dredging and reconnection and minimal flow increases (Lamouroux et al., 2015). In summer 2017, we sampled 17 geomorphic surfaces (mostly gravel bars that were somewhat subject to fine-grained overbank alluviation) that we classified into three types:

relict bars that were repeatedly cleared (n = 9), newly reprofiled banks (n = 4) and naturally rejuvenated bars (n = 4) (Figure 1). Relict bars are usually fixed in bypassed sections of the Rhône riverbed, i.e., coarse sediments are no longer remobilized during flood events and are subject to fine-grained overbank sedimentation and colonization by trees. These large fixed gravel bars correspond to a part of the river bottom before derivation and represent the only terrestrial habitats available for species within the Rhône riverbed. To prevent encroachment by trees and shrubs of these surfaces and associated channel narrowing, maintenance measures are implemented by river managers to scrap them by either brush clearing (n = 4) or plowing (n = 5). Reprofiled banks are newly restored surfaces due to bank reprofiling following dike removal in bypassed sections of the Rhône riverbed. These engineering actions were performed to widen the channel for improving flow conveyance. These new surfaces are not paved, with soil texture and topographical features quite comparable to those of repeatedly cleared bars. They represent new available terrestrial habitats for species colonization within the Rhône riverbed. These actions have been conducted on four reaches (one in 2009, one in 2015 and two in 2017) belonging to two bypassed sections. They are intended to be progressively eroded to promote self-gravel augmentation, bedload transport and new natural rejuvenated bar creation downstream. Naturally rejuvenated bars are still actively mobile bars in terms of bedload supply and erosion and are thus subject to bedload alluviation/rejuvenation processes, i.e., coarse sediments are partially remobilized during flood events. In the study area, such surfaces are mostly restricted to tributary rivers (Ain, Drôme and Doux Rivers), although a few bars of the Rhône riverbed are currently naturally rejuvenated because of existing bedload transport (e.g., Miribel reach, upstream of Lyon). These bars are considered natural reference geomorphic surfaces and are features that could potentially be restored in some of the bypassed sections.

2.2. Vegetation data

On each geomorphic surfaces, vegetation was surveyed along three transects spaced at a distance of approximately 100 m from each other (Figure 1). Transects were positioned perpendicularly to the river, in areas with clear elevation gradients relative to the water surface (means $\pm$ SD = 1.17 $\pm$ 0.93 m; range = 0.04-6.20 m). The transect length varied from 30 m to 105 m, depending on the length of the local gradient. They started directly from the water line and ended before the alluvial floodplain, which was mostly occupied by riparian forest. Along each transect, four 5 x 5 m quadrats (25 m²) were positioned, for a total of 12 quadrats per geomorphic surface. Quadrats were positioned in visually homogenous topographical and pedological areas, while avoiding aquatic parts of the river margin and targeting similar plant communities. Within each quadrat (n = 204), all vascular plants were surveyed following the Braun-Blanquet abundance-dominance methodology. Because geomorphic surfaces along the Rhône River were repeatedly cleared, tree and shrub layers were systematically absent and vegetation surveys were conducted by considering the herbaceous layer only. To characterize the entire plant community and account for plant phenology, two complete surveys were conducted (June and July 2017) and to limit observer biases, all surveys were performed by the first author (PJ).

2.3. Trait data

We used three groups of plant traits. Three ecological traits – plant species preferences for light, nitrogen content and soil moisture – were extracted from the Baseflor database (Julve, 1998), and the few missing values (n species = 22) were filled by using the ecological information of closely related species, i.e., same genus. These ecological traits, i.e., Ellenberg's indicator values (Ellenberg et al., 1992), were used to characterize variations in species habitat niches along environmental gradients and to highlight whether maintenance and restoration measures modify abiotic conditions along river margins. Three morphological traits – specific leaf area (SLA), plant height at maturity and seed mass – were extracted from

the TRY database (Kattge et al. 2011), and we calculated a mean trait value per species from them after removing all values with an error risk > 3 (see also A.1). For missing values, data were completed using the LEDA database (Kleyer et al., 2008) and, in a few cases (n species = 6), by using the information available for closely related species i.e., same genus. Within Westoby's L-H-S scheme (Westoby, 1998), SLA is related to resource acquisition and conservation, height at maturity to competitive ability and seed mass to dispersal ability. Three groups of species – annual, hydrochorous and alien species – were identified from the LEDA (annual species) and Baseflor databases (hydrochorous and alien species). Annual species characterize pioneer communities which are targeted by restoration measures along the Rhône River; hydrochorous species are dispersal-specialized species of high importance in riparian plant communities which are sensitive to alteration of the flow regime (Nilsson et al., 2010); alien species, defined as introduced species originated from America, Africa and Asia (i.e., based on species chorology), represent potential threats to native riparian plant communities which pose challenges in restoration measures (Richardson et al., 2007). Overall, these three species groups were used to highlight how environmental gradients drive changes in riparian zones diversity patterns and to assess whether maintenance and restoration measures affect these patterns. For each trait value, we computed community-weighted means (CWM) and functional dispersion (FDis). CWM is defined as the mean of trait values weighted by the relative abundance of each species bearing each value (Lavorel et al., 2008). FDis is defined as the mean distance of individual species to the weighted centroid of all species in the assemblage (Laliberté and Legendre, 2010). For each species group, we computed the total species richness and cover.

2.4. Elevation and soil texture gradients

At the center of each quadrat, the elevation difference to the water line was characterized using topographical surveys. Topographical measurements were conducted using a laser rangefinder (TruPulse 200X) positioned on a tripod at a constant height of 1.20 m and a target at a height of 2 m. The horizontal distance between the tripod and the target was taken from the water line at the beginning of each transect and the center of the four successive quadrats. Soil properties were derived from two soil core samples (30-cm depth) collected in two representative areas within each quadrat and pooled together. Each composite soil sample was then dried, weighed and sieved at the laboratory to estimate the proportion of large (>0.2 -mm mesh), medium (>50 - μ m mesh) and fine (<50 - μ m mesh) sediments. Due to the high correlation between these three variables and to avoid multicollinearity issues, we focused on only the proportion of subsoil fine sediments in subsequent analysis. This variable was preferred over others because it was considered the most meaningful variable to represent the process of fine-grained sediment alluviation that affects several gravel bars along the Rhône River.

2.5. Statistical analysis

Analyses were performed with R version 3.4.2 (R Core Team, 2017). Continuous independent variables were the elevation difference (denoted “elevation” in tables and figures) and the proportion of fine sediments (denoted “fine_sediments”); independent factors were maintenance measures, i.e., brush clearing vs plowing (denoted “maintenance”), and geomorphic surface type, i.e., repeatedly cleared relict bars vs newly reprofiled banks vs naturally rejuvenated bars (denoted “type”). Because independent variables were measured on different scales and because we were interested in interpreting the main effects of continuous variables in the presence of interactions, input variables were centralized (Schielzeth, 2010). Dependent variables were the CWM and the FDis of ecological and morphological trait values as well as the pooled richness and the cover of species groups at each quadrat. Because

the seed mass of a few species was very large, this trait value was log-transformed before CWM and FDis calculations. We then used one-way ANOVA with a type III sum of squares to determine if the elevation difference and the proportion of fine sediments varied differently between maintenance measures and among geomorphic surface types (car package). To determine whether human-mediated environment gradients shape the functional composition of riparian plants we used an analytical strategy involving two steps. First, by using quadrats located on only repeatedly cleared relict bars of the Rhône River ($n = 108$), we sought to understand whether environmental gradients shape riparian plant communities along a highly degraded large River and whether maintenance measures influence the response of plant functional composition to environmental gradients. Second, by using the whole dataset ($n = 204$), we sought to understand whether the response of plant functional composition to environmental gradients vary among repeatedly cleared relict bars, newly reprofiled banks of the Rhône riverbed and naturally rejuvenated bars, mostly sampled on tributary rivers (natural reference).

2.5.1. How do environmental gradients and maintenance measures shape plant functional composition along the Rhône River?

We considered 13 a priori models testing the main and interaction effects between the elevation and soil texture gradients and between environmental gradients and maintenance measures plus the null model (A.2.1). Because we used a nested sampling design in which “quadrats” were not really replicated along “transects” and “banks”, we used linear mixed models (LMMs, lme4 package) in which “transects” and “banks” were included as random effects. For continuous dependent variables (i.e., trait values), we fitted normal LMMs or log-normal LMMs for skewed response variables (i.e., plant height). For count dependent variables (i.e., richness and cover), to avoid statistical issues linked to overdispersion, we fitted negative binomial distribution general linear mixed models (GLMMs, lme4 package). In

all candidate models, the variance inflation factor was below five, indicating a lack of collinearity issues (Dormann et al., 2013). The variance explained by LMMs and GLMMs was estimated using the marginal coefficient of determination for fixed effect parameters alone (Nakagawa and Schielzeth, 2013). To identify the most parsimonious regression model, we used Akaike's information criterion corrected for small sample sizes (Burnham and Anderson, 2002). To estimate parameters and associated unconditional standard errors, we used model averaging based on the subset of top ranking models for which $\Delta AICc < 2$ (MuMin package).

2.5.2. How does the response of plant functional composition to environmental gradients vary among geomorphic surface types?

We considered 13 a priori models testing the main and interaction effects between both environmental gradients and between environmental gradients and geomorphic surface types plus the null model (A.2.2). In LMMs and GLMMs, the relict bar factor was set as the reference level. In all candidate models, the variance inflation factor was below five. The variance explained by models, models ranking and estimate of parameters and of unconditional standard errors were done in the same manner as previously (2.5.1.).

3. RESULTS

Overall, 273 plant species ($\text{mean} \pm \text{SD} = 28 \pm 13$) were recorded on the 204 quadrats along the Rhône River and tributaries. Distinguishing between surface types: 203 species (28 ± 12) were recorded on repeatedly cleared relict bars ($n = 108$), with 142 species (32 ± 10) on surfaces maintained by brush clearing ($n = 48$) and 171 species (25 ± 13) on surfaces maintained by plowing ($n = 60$); 198 species (31 ± 13) were recorded on newly reprofiled banks ($n = 48$); and 175 species (24 ± 13) were recorded on naturally rejuvenated bars ($n = 48$). Because ecological successions are blocked on repeatedly cleared relict bars, i.e., gravel bars are

scraped to avoid colonization by tree and shrub species, regeneration and seedling of tree species ($n = 15$) were removed from the dataset to avoid bias in the estimates of trait values (especially plant height at maturity). Subsequent analyses were thus conducted on a dataset of 258 herbaceous plant species.

3.1. How environmental gradients vary between maintenance measures and among geomorphic surface types?

On repeatedly cleared relict bars, ANOVAs showed that the variation in mean elevation and in the mean proportion of fine sediments was not related to maintenance measures (Figure 2, A.3 & A.4). This result indicates that in our study design, both environmental gradients varied consistently among geomorphic surfaces maintained by either brush clearing or plowing. Between repeatedly cleared relict bars, newly reprofiled banks and naturally rejuvenated bars, ANOVAs showed no significant variations in the mean elevation. However, the proportion of fine sediments significantly differed between surface types (p -value < 0.001 , Figure 2), which was due to the lower proportion of fine sediments on naturally rejuvenated bars (mean $\pm$ SD, 3.99 ± 7.47) than on repeatedly cleared relict bars (15.90 ± 16.42) and newly reprofiled banks (14.20 ± 10.71).

3.2. How do environmental gradients and maintenance measures shape plant functional composition along the Rhône River?

Results showed that variations in plant species preferences for soil moisture (mean and dispersion), light (dispersion) and nitrogen content (dispersion), in plant height (mean), seed mass (mean) and SLA (dispersion) trait values and in hydrochorous (richness and cover), annual (richness and cover) and alien (cover) species were best predicted by one of the a priori models (Table 1). The goodness of fit of these models varied from 14.6 % (dispersion of nitrogen content) to 51.7 % (hydrochorous species richness). However, for light (mean), nitrogen content (mean), SLA (mean), plant height (dispersion), seed mass (dispersion) and

alien species (cover), since the null model was either the top-ranked or part of top-ranked models (i.e., $\Delta AICc < 2$) none of the estimated parameters is of any inferential value.

3.2.1. Influence of environmental gradients

Model averaging (Table 2) revealed that an increase in the elevation level induced decreases in mean soil moisture trait value, the dispersion of SLA trait values and hydrochorous species richness but an increase in mean seed mass trait value. An increase in the proportion of fine sediments induced decreases in the dispersion of soil moisture, nitrogen content and SLA trait values and in the richness and cover of annual species. Mean plant height, the dispersion of light trait values, the richness of alien species and the cover of hydrochorous species were influenced by the interaction between the elevation and soil texture gradients. Graphical interpretation (Figure 3) revealed that the mean plant height and cover of hydrochorous species were higher on gravel bars with high proportion of fine sediments and decreased less with elevation when the proportion of fine sediments was low. The dispersion of light trait values and richness of alien species decreased with elevation when the proportion of fine sediments was low but increased with elevation when the proportion of fine sediments was high.

3.2.2. Influence of maintenance measures

Model averaging (Table 2) revealed that maintenance measures mediated the response of nitrogen content (dispersion) and SLA (dispersion) trait values and of annual species (richness and cover) to environmental gradients. Graphical interpretation (Figure 4) revealed that the dispersion of SLA trait values was higher on bars maintained by plowing and decreased less with elevation on bars maintained by brush clearing. The dispersion of nitrogen content trait values and annual species cover decreased with an increase in the proportion of fine sediments on bars maintained by brush clearing but did not change on those maintained by plowing. The richness of annual species was higher on bars maintained by plowing and

tended to decrease more with increasing proportion of fine sediments on bars maintained by brush clearing. Finally, alien species richness was higher on bars maintained by plowing.

3.3. How does the response of plant functional composition to environmental gradients vary among geomorphic surface types?

Results showed that variations in plant species preferences for soil moisture (mean and dispersion), light (mean and dispersion) and nitrogen content (mean and dispersion), in SLA (mean and dispersion), plant height (mean) and seed mass (mean and dispersion) trait values and in hydrochorous (richness and cover), annual (richness) and alien (cover) species were best predicted by one of the a priori models (Table 3). The goodness of fit of these models varied from 3.7 % (dispersion of seed mass) to 56.0 % (mean soil moisture). However, for plant height (dispersion), alien (richness) and annual (cover) species, since the null model was either the top-ranked or part of top-ranked models (i.e., $\Delta AICc < 2$), none of the estimated parameters is of any inferential value.

3.3.1. Influence of environmental gradients

Model averaging (Table 4) revealed that an increase in the elevation induced a decrease in the dispersion of SLA trait values but an increase in mean light trait value and the dispersion of seed mass trait values, which was consistent among the three geomorphic surfaces. An increase in the proportion of fine sediments induced a decrease in the dispersion of soil moisture, nitrogen content and SLA trait values and in the richness of annual species but an increase in the cover of alien species, which was consistent among the three geomorphic surfaces. Mean plant height and dispersion for light trait values were influenced by the interaction between the elevation and soil texture gradients, i.e., mean plant height decreased with elevation when the proportion of fine sediments was high but increased when the proportion of fine sediments was low while the dispersion for light trait values decreased with

elevation when the proportion of fine sediments was low but increased when the proportion of fine sediments was high.

3.3.2. *Influence of restoration measures*

Comparing the response of plant traits to environmental gradients between repeatedly cleared relict bars and newly reprofiled banks, model averaging (Table 4) revealed that soil moisture (mean), plant height (mean) and seed mass (mean) trait values and that hydrochorous (richness and cover) and alien (cover) species varied differently between the two geomorphic surfaces. Graphical interpretation (Figure 5-A) revealed that mean soil moisture trait value, mean plant height trait value and hydrochorous richness and cover decreased more rapidly with elevation on relict bars than on reprofiled banks. Mean seed mass trait value increased more rapidly with elevation on relict bars than on reprofiled banks. Alien cover was higher on reprofiled banks than on relict bars and tended to decrease more rapidly with elevation on reprofiled banks. Finally, mean light trait value was higher on reprofiled banks than on relict bars.

Comparing the response of plant traits to environmental gradients between repeatedly cleared bars and naturally rejuvenated bars, model averaging (Table 4) revealed that nitrogen content (mean), soil moisture (dispersion), SLA (mean), plant height (mean) and seed mass (mean and dispersion) traits values and that alien (cover) species varied differently between the two geomorphic surfaces. Graphical interpretation (Figure 5-B) revealed that mean nitrogen content, SLA and seed mass trait values decreased with elevation on naturally rejuvenated bars but increased on repeatedly cleared bars. Mean plant height increased with elevation on naturally rejuvenated bars but decreased on repeatedly cleared bars. The dispersion of soil moisture trait values decreased more rapidly with elevation level on naturally rejuvenated bars than on repeatedly cleared bars. Seed mass dispersion decreased more rapidly with proportion of fine sediments on naturally rejuvenated bars than on repeatedly cleared bars. Alien cover

was higher on naturally rejuvenated bars than on repeatedly cleared bars and tended to decrease less rapidly with elevation on naturally rejuvenated bars. Finally, the mean light trait value was higher, but the mean soil moisture trait value and hydrochorous species richness and cover were lower on naturally rejuvenated bars than on repeatedly cleared bars.

4. DISCUSSION

Using a trait-based approach, which is rarely used in studies evaluating the success of restoration actions on riparian vegetation (González et al., 2015), we showed that (i) elevation and soil texture gradients have strong and sometimes interactive effects on plant functional composition and that (ii) the influence of both environmental gradients is mediated by human activity. The co-occurrence of species with contrasting resource requirements and morphological traits was higher in highly disturbed environments, i.e., on surfaces not subject to fine-grained overbank alluviation and hydrological disconnection. Rejuvenation processes, through natural or human-induced disturbances, thus appear essential to avoid biotic homogenization and promote habitat heterogeneity.

4.1. Both the elevation and the soil texture gradients shape riparian plant functional composition

In riparian zones of the Rhône River, shifts in ecological traits were highlighted in relation to both environmental gradients. The elevation gradient drives changes in the plant species habitat niche, emphasizing patterns of hydrological niche segregation (Silvertown et al., 2015). In line with previous studies (Auble et al., 1994; Fraaije et al., 2015; Garssen et al., 2017), this shift confirms the importance of the soil moisture gradient in shaping riparian plant communities. Conversely, the soil texture gradient was related to patterns of trait convergence, i.e., an increase in the proportion of fine sediments induced decreases in the dispersion of soil moisture and nitrogen content trait values. Also, low proportion of fine

448 sediments allowed for a better co-occurrence of species with contrasting light resource
449 requirements, revealing a combination of xeric, mesic and hydric habitat conditions on gravel
450 bars made of coarse sediments. This result suggests that continuous alluviation of fines leads
451 to patterns of biotic homogenization (Olden and Rooney, 2006), underlining thus the
452 importance of rejuvenation processes in increasing habitat heterogeneity (Steiger et al., 2005).
453 Changes in the plant species habitat niche induced environmental selection toward specific
454 plant morphological traits. In line with previous studies (Kyle and Leishman, 2009; McCoy-
455 Sulentic et al., 2017), a shift in the mean seed mass trait value was highlighted along the
456 elevation gradient, confirming that species with small seed masses have better potential to
457 colonize disturbed environments than species with large seed masses (Gurnell et al., 2008).
458 Moreover, the interaction between the elevation and soil texture gradients influenced mean
459 plant height. Indeed, tall herb species, such as *Phragmites australis* and *Phalaris*
460 *arundinacea*, were more frequently surveyed close to the water line on relict gravel bars
461 subject to fine sediment alluviation. This relation may reflect the adaptation of plant species to
462 fairly constant flow during the vegetative period due to the regulation of flood events in
463 bypassed sections but also to the long-term decrease in bedload discharge, limiting the
464 mortality of perennial plants by coarse bed scour and burial disturbances (e.g., Kui and Stella,
465 2016; Polzin and Rood, 2006). Besides shifts in traits, patterns of trait
466 convergence/divergence were highlighted. Along both the elevation and the soil texture
467 gradients, SLA values converged, i.e., species with contrasting resource acquisition and
468 conservation strategies co-occurred less at high elevation and on gravel bars subject to fine
469 sediment alluviation, thus confirming patterns of biotic homogenization.
470 Finally, diversity patterns of plant species groups were also influenced by both environmental
471 gradients. A higher diversity of hydrochorous species occurred at low elevation level (Fraaije
472 et al., 2017), while a higher diversity of annual species was related to a low proportion of fine

sediments (Corenblit et al., 2009). Additionally, our results showed that the elevation and soil texture gradients interacted to shape diversity patterns. Indeed, the richness of alien species was maximal at low elevation but tended to increase along the elevation gradient when the proportion of fine sediments was high (Brummer et al., 2016). This pattern shows that the effect of both environmental gradients on plant communities may not be independent and argues for a better consideration of the soil texture gradient in riparian vegetation studies (Kyle and Leishman, 2009). Since taxonomic and functional diversity peaked at low elevation, on geomorphic surfaces not subject to fine-grained sediment alluviation, i.e., in frequently flooded areas, our results show that greater disturbance of environments within the Rhône riverbed promotes a greater diversification of habitats that benefit a larger diversity of species with contrasting functional strategies. Rejuvenation processes thus appear essential to conserving or restoring the functionality of riparian communities in highly degraded rivers.

4.2. Patterns of biotic homogenization are mitigated on gravel bars maintained by plowing

Distinguishing between relict bars of the Rhône River maintained by either brush clearing or plowing, we found that shifts in ecological and morphological traits in relation to environmental gradients were consistent. However, along the elevation gradient, the dispersion of SLA trait values diverged more at low elevation on river margins maintained by plowing than on other geomorphic surfaces. Also, with increasing proportion of fine sediments, nitrogen content trait values converged on only bars maintained by brush clearing. Co-occurrence of species with similar ecological and morphological traits were thus more pronounced on bars maintained by brush clearing, indicating patterns of biotic homogenization. Reversely, deeply disturbing the soil surface increases habitat heterogeneity, by providing a larger diversity of environmental conditions (microsites, resources) and thus a greater diversity of habitat niches. Specifically, plowing permits the active remobilization of

fine sediments, favoring erosion processes at local scale and substrate heterogeneity, which better allows for the co-occurrence of species with contrasting trophic requirements and resource acquisition strategies. Plowing also mediated the response of annual species to the soil texture gradient by enabling the maintenance of a more cover on gravel bars subject to fine-grained overbank alluviation, and promoted larger species richness of annual but also of alien species. Consistently with previous studies (e.g., Catford et al., 2012; Nilsson et al., 1989), this pattern highlight that colonization opportunities for ruderal species, among which annual and alien herbaceous species are well represented, increase in frequently disturbed riparian zones, mostly because competition for limiting resources is minimal. Overall, those results underlined the importance of rejuvenation processes in riparian zones to support functional and species diversity (e.g., Biswas and Mallik, 2010). Specifically, our results indicated that plowing better promotes pioneer communities within the Rhône riverbed, but also alien species, and better limits biotic homogenization along environmental gradients. Given the current alteration of the natural disturbance regimes of the Rhône River (mainly bar scouring through bedload transport), it may be argued that plowing could be a potential compromise between hydraulic (i.e., maintain wide low-roughness channels) and ecological (i.e., promote pioneer communities) considerations. However, beyond the fact that plowing does not only promote native pioneer communities, the strong economic and environmental costs of this maintenance operation must also be considered. In this view, bedload transport restoration in bypassed sections should be promoted first as a sustainable management perspective (González et al., 2017).

4.3. The response of plant functional composition to environmental gradients

diverged between repeatedly cleared relict bars, newly reprofiled banks and naturally rejuvenated bars.

Among the three geomorphic surface types, shifts in soil moisture and light requirement trait values with elevation and trait convergence for soil moisture and nitrogen content trait values with increasing proportion of fine sediments were consistent. However, strong differences in plant species habitat niches were also highlighted. Soil moisture trait values were lower on naturally rejuvenated bars than on other geomorphic surfaces and decreased less with elevation on newly reprofiled banks. Moreover, the abundance of full light-demanding species was higher on naturally rejuvenated and newly reprofiled surfaces than on repeatedly cleared relict bars. Thus, the environmental filtering from the lower to the upper part of river margins appears to be more stressful on naturally rejuvenated bars. This pattern is undoubtedly related to the low proportion of fine sediments on these surfaces and thus to the low water-holding capacity (Naiman and Decamps, 1997). Soil properties may also explain the shift in nitrogen content trait value along the elevation gradient that we found on only naturally rejuvenated bars. Indeed, the decrease in soil fertility (i.e., inferred from plant species trait values) with elevation may be related to both the low proportions of fine-grained overbank deposits at medium elevation and the young age of naturally rejuvenated bars, which thus prevent the establishment of the nitrogen-demanding species that are abundant on older relict bars (e.g., *Artemisia vulgaris*, *Solanum nigrum*, *Sonchus oleraceus*). Finally, the convergence of soil moisture trait values highlighted along the elevation gradient on only naturally rejuvenated bars may result from higher environmental selection toward plant species with similar hydrological requirements, confirming that these natural reference bars encompass stressful environmental conditions.

As expected, differences in plant species habitat niches among the three surface types were related to shifts in plant morphological traits. Along the elevation gradient, mean plant height increased on naturally rejuvenated bars but decreased on repeatedly cleared relict bars and newly reprofiled banks. This underlines the importance of competition in bypassed sections

and confirms long periods of growth opportunity between disturbances. Additionally, our results indicate that the mean seed mass increased with elevation on relict bars and reprofiled banks but decreased with elevation on naturally rejuvenated bars. Considering the critical role played by flooding in shaping patterns of seed deposition along the elevation gradient (Fraaije et al., 2017), the greater occurrence of small-seed-mass species at middle elevation on naturally rejuvenated bars may be due to a greater range of flooding here than at other geomorphic surfaces, allowing the accumulation of small buoyant seeds near the average water line. Indeed, as compared to naturally rejuvenated bars, which are periodically subject to flood events, relict bars and reprofiled banks in bypassed sections are subject to flow regulation, which imply a fairly constant minimum flow most of the year. Additionally, areas that are flooded more frequently may be associated with greater environmental selection toward small seed mass. This possibility is evidenced by differences in mean seed mass values among naturally rejuvenated bars (mean $\pm$ SD = 3.48 ± 2.88 mg), relict bars (4.15 ± 6.28 mg) and reprofiled banks (6.42 ± 8.51 mg). Finally, mean SLA values decreased along the elevation gradient on only naturally rejuvenated bars. This result reveals a shift in plant growth rate strategy with elevation levels, from low to high resource conservation strategy, and highlights adaptation to drought stress (Chaves et al., 2003). Overall, our results confirm that naturally rejuvenated bars are more stressful environments than the two other geomorphic surfaces types, hosting distinct functional assemblages of plant species.

For diversity patterns, results showed that the richness and cover of hydrochorous species decreased less along the elevation gradient on newly reprofiled banks than on other geomorphic surfaces and were less abundant on naturally rejuvenated bars. Since clear links exist between seed abundance and fine sediment amount (Goodson et al., 2003), this result may show that hydrochorous species require a certain minimum proportion of fine sediments to successfully establish along river margins (e.g., Corenblit et al., 2016). Moreover, the cover

of alien species was higher on naturally rejuvenated bars and reprofiled banks than other geomorphic surfaces. For naturally rejuvenated bars, this result is mostly due to the overrepresentation of the ruderal species *Ambrosia artemisiifolia* (frequency of occurrence: 81%); for reprofiled banks, this result is mostly due to the rapid colonization of newly available terrestrial habitats by *Symphytotrichum x salignum* (frequency of occurrence: 94%). Overall, we showed that the response of riparian plants to environmental gradients is mostly inconsistent among the three geomorphic surface types. This result is undoubtedly related to differences in environmental conditions and flow regimes between the Rhône River and its tributaries. This may confirm the alteration of the natural disturbance regimes of the Rhône River in terms of flow regulation and bedload supply and highlights their ecological consequences to the functionality of riparian habitats (e.g., Bejarano et al., 2017). Moreover, we showed that new terrestrial habitats resulting from restoration measures are in the same ecological trajectories of repeatedly cleared relict bars. Whereas the hydraulic efficiency of dike removal and bank reprofiling is not proven here (see Thorel et al. 2017), these findings suggest that their ecological added value is also questionable (e.g., Weber et al., 2009). In view of an effective ecological restoration of highly degraded river ecosystems (Molles, et al., 1998; Roni et al., 2002), we argue that (i) the minimum flow in bypassed sections of the Rhône River should vary more (see e.g., Hayes et al., 2018), (ii) bedload transport restoration should be a priority to allow for gravel bars that are more dynamic and (iii) newly reprofiled banks should better mimic landforms of natural river margins, by favoring low elevation surfaces and a high proportion of coarse sediments.

5. CONCLUSION

The need to disentangle interactions among natural and human-mediated environmental changes is important, and improved knowledge of these interactions could increase the

efficiency of conservation and restoration strategies (e.g., Dawson et al., 2017). Here, we showed that focusing on a single environmental gradient, such as the flood gradient, may lead to an incomplete understanding of the ecological processes that drives community assembly in riparian zones and that accounting for the influence that current human activity has on riparian biodiversity is fundamental to unraveling how environmental filtering acts on assembly processes. Specifically, we showed that biotic homogenization on river margins was clearly related to fine-grained alluviation processes. Since biotic homogenization alters ecosystem functioning and productivity and leads to the deterioration of ecosystem goods and services (Clavel et al., 2011), restoring the natural disturbance regimes should be a priority. Hence, on highly degraded rivers systems such as the Rhône River, the efficiency of local-scale restoration measures (e.g., bank reprofiling following dike removal) would benefit from a more global strategy of reactivation of bedload transport and supply at the reach and catchment scales (e.g., Rohde et al., 2005).

6. ACKNOWLEDGEMENTS

We thank Vincent Breton, Paul Cavaillé, Gilles Favier, Delphine Jaymond and Sophie Labonne for help in the field. The study has been supported by the TRY initiative on plant traits (<http://www.try-db.org>). The TRY initiative and database is hosted, developed and maintained by J. Kattge and G. Boenisch (Max Planck Institute for Biogeochemistry, Jena, Germany). TRY is currently supported by Future Earth/bioDISCOVERY and the German Center for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig. This work was cofunded by the Labex DRIIHM, French program "Investissements d'Avenir" (ANR-11-LABX-0010), which is managed by the ANR, the OHM Vallée du Rhône and IRSTEA.

7. REFERENCES

622 Auble, G.T., Friedman, J.M., Scott, M.L., 1994. Relating riparian vegetation to present and
 623 future streamflows. *Ecol. Appl.* 4, 544–554. <https://doi.org/10.2307/1941956>
 624 Bejarano, M.D., Nilsson, C., Aguiar, F.C., 2017. Riparian plant guilds become simpler and
 625 most likely fewer following flow regulation. *J. Appl. Ecol.*
 626 <https://doi.org/10.1111/1365-2664.12949>
 627 Biron, P.M., Buffin-Bélanger, T., Larocque, M., Choné, G., Cloutier, C.-A., Ouellet, M.-A.,
 628 Demers, S., Olsen, T., Desjarlais, C., Eyquem, J., 2014. Freedom space for rivers: A
 629 sustainable management approach to enhance river resilience. *Environ. Manage.* 54,
 630 1056–1073. <https://doi.org/10.1007/s00267-014-0366-z>
 631 Biswas, S.R., Mallik, A.U., 2010. Disturbance effects on species diversity and functional
 632 diversity in riparian and upland plant communities. *Ecology* 91, 28–35.
 633 Brummer, T.J., Byrom, A.E., Sullivan, J.J., Hulme, P.E., 2016. Alien and native plant
 634 richness and abundance respond to different environmental drivers across multiple
 635 gravel floodplain ecosystems. *Divers. Distrib.* 22, 823–835.
 636 <https://doi.org/10.1111/ddi.12448>
 637 Bunn, S.E., Arthington, A.H., 2002. Basic principles and ecological consequences of altered
 638 flow regimes for aquatic biodiversity. *Environ. Manage.* 30, 492–507.
 639 <https://doi.org/10.1007/s00267-002-2737-0>
 640 Burnham, K.P., Anderson, D.R., 2002. Model selection and multi-model inference: a practical
 641 information-theoretic approach, 2nd ed. Springer-Verlag, New York.
 642 Cadotte, M.W., Carscadden, K., Mirotchnick, N., 2011. Beyond species: functional diversity
 643 and the maintenance of ecological processes and services: Functional diversity in
 644 ecology and conservation. *J. Appl. Ecol.* 48, 1079–1087.
 645 <https://doi.org/10.1111/j.1365-2664.2011.02048.x>

Catford, J.A., Daehler, C.C., Murphy, H.T., Sheppard, A.W., Hardesty, B.D., Westcott, D.A.,
 Rejmánek, M., Bellingham, P.J., Pergl, J., Horvitz, C.C., Hulme, P.E., 2012. The
 intermediate disturbance hypothesis and plant invasions: Implications for species
 richness and management. *Perspect. Plant Ecol. Evol. Syst.* 14, 231–241.
<https://doi.org/10.1016/j.ppees.2011.12.002>

Chaves, M.M., Maroco, J.P., Pereira, J.S., 2003. Understanding plant responses to drought -
 from genes to the whole plant. *Funct. Plant Biol.* 30, 239–264.
<https://doi.org/10.1071/FP02076>

Clavel, J., Julliard, R., Devictor, V., 2011. Worldwide decline of specialist species: toward a
 global functional homogenization? *Front. Ecol. Environ.* 9, 222–228.
<https://doi.org/10.1890/080216>

Comiti, F., Da Canal, M., Surian, N., Mao, L., Picco, L., Lenzi, M.A., 2011. Channel
 adjustments and vegetation cover dynamics in a large gravel bed river over the last
 200years. *Geomorphology* 125, 147–159.
<https://doi.org/10.1016/j.geomorph.2010.09.011>

Corenblit, D., Steiger, J., Gurnell, A.M., Tabacchi, E., Roques, L., 2009. Control of sediment
 dynamics by vegetation as a key function driving biogeomorphic succession within
 fluvial corridors. *Earth Surf. Process. Landf.* 34, 1790–1810.
<https://doi.org/10.1002/esp.1876>

Corenblit, D., Vidal, V., Cabanis, M., Steiger, J., Garófano-Gómez, V., Garreau, A.,
 Hortobágyi, B., Otto, T., Roussel, E., Voldoire, O., 2016. Seed retention by pioneer
 trees enhances plant diversity resilience on gravel bars: Observations from the river
 Allier, France. *Adv. Water Resour.* 93, 182–192.
<https://doi.org/10.1016/j.advwatres.2016.02.015>

670 Dawson, S.K., Warton, D.I., Kingsford, R.T., Berney, P., Keith, D.A., Catford, J.A., 2017.
671 Plant traits of propagule banks and standing vegetation reveal flooding alleviates
672 impacts of agriculture on wetland restoration. *J. Appl. Ecol.*
673 <https://doi.org/10.1111/1365-2664.12922>

674 de Bello, F., Lavorel, S., Lavergne, S., Albert, C.H., Boulangeat, I., Mazel, F., Thuiller, W.,
675 2013. Hierarchical effects of environmental filters on the functional structure of plant
676 communities: a case study in the French Alps. *Ecography* 36, 393–402.
677 <https://doi.org/10.1111/j.1600-0587.2012.07438.x>

678 Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G.,
679 Gruber, B., Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean, C., Osborne,
680 P.E., Reineking, B., Schröder, B., Skidmore, A.K., Zurell, D., Lautenbach, S., 2013.
681 Collinearity: a review of methods to deal with it and a simulation study evaluating
682 their performance. *Ecography* 36, 27–46. [https://doi.org/10.1111/j.1600-](https://doi.org/10.1111/j.1600-0587.2012.07348.x)
683 [0587.2012.07348.x](https://doi.org/10.1111/j.1600-0587.2012.07348.x)

684 Ellenberg, H., Weber, H.E., Düll, R., Wirth, V., Werner, W., Paulißen, D., 1992. Zeigerwerte
685 von Pflanzen in Mitteleuropa. *Scr. Geobot.* 18, 1–248.

686 Feld, C.K., Birk, S., Bradley, D.C., Hering, D., Kail, J., Marzin, A., Melcher, A., Nemitz, D.,
687 Pedersen, M.L., Pletterbauer, F., Pont, D., Verdonschot, P.F.M., Friberg, N., 2011.
688 From Natural to Degraded Rivers and Back Again, in: Woodward, G. (Ed.), *Advances*
689 *in Ecological Research*. Elsevier, pp. 119–209. [https://doi.org/10.1016/B978-0-12-](https://doi.org/10.1016/B978-0-12-374794-5.00003-1)
690 [374794-5.00003-1](https://doi.org/10.1016/B978-0-12-374794-5.00003-1)

691 Fraaije, R.G.A., Moinier, S., van Gogh, I., Timmers, R., van Deelen, J.J., Verhoeven, J.T.A.,
692 Soons, M.B., 2017. Spatial patterns of water-dispersed seed deposition along stream
693 riparian gradients. *PLOS ONE* 12, e0185247.
694 <https://doi.org/10.1371/journal.pone.0185247>

695 Fraaije, R.G.A., ter Braak, C.J.F., Verduyn, B., Breeman, L.B.S., Verhoeven, J.T.A., Soons,
 696 M.B., 2015. Early plant recruitment stages set the template for the development of
 697 vegetation patterns along a hydrological gradient. *Funct. Ecol.* 29, 971–980.
 698 <https://doi.org/10.1111/1365-2435.12441>

699 Garssen, A.G., Baattrup-Pedersen, A., Riis, T., Raven, B.M., Hoffman, C.C., Verhoeven,
 700 J.T.A., Soons, M.B., 2017. Effects of increased flooding on riparian vegetation: Field
 701 experiments simulating climate change along five European lowland streams. *Glob.*
 702 *Change Biol.* 23, 3052–3063. <https://doi.org/10.1111/gcb.13687>

703 González, E., Felipe-Lucia, M.R., Bourgeois, B., Boz, B., Nilsson, C., Palmer, G., Sher, A.A.,
 704 2017. Integrative conservation of riparian zones. *Biol. Conserv.* 211, 20–29.
 705 <https://doi.org/10.1016/j.biocon.2016.10.035>

706 González, E., Sher, A.A., Tabacchi, E., Masip, A., Poulin, M., 2015. Restoration of riparian
 707 vegetation: A global review of implementation and evaluation approaches in the
 708 international, peer-reviewed literature. *J. Environ. Manage.* 158, 85–94.
 709 <https://doi.org/10.1016/j.jenvman.2015.04.033>

710 Goodson, J.M., Gurnell, A.M., Angold, P.G., Morrissey, I.P., 2003. Evidence for hydrochory
 711 and the deposition of viable seeds within winter flow-deposited sediments: the River
 712 Dove, Derbyshire, UK. *River Res. Appl.* 19, 317–334. <https://doi.org/10.1002/rra.707>

713 Gumiero, B., Mant, J., Hein, T., Elso, J., Boz, B., 2013. Linking the restoration of rivers and
 714 riparian zones/wetlands in Europe: Sharing knowledge through case studies. *Ecol.*
 715 *Eng.* 56, 36–50. <https://doi.org/10.1016/j.ecoleng.2012.12.103>

716 Gurnell, A., Thompson, K., Goodson, J., Moggridge, H., 2008. Propagule deposition along
 717 river margins: linking hydrology and ecology. *J. Ecol.* 96, 553–565.
 718 <https://doi.org/10.1111/j.1365-2745.2008.01358.x>

719 Hayes, D.S., Brändle, J.M., Seliger, C., Zeiringer, B., Ferreira, T., Schmutz, S., 2018.
 720 Advancing towards functional environmental flows for temperate floodplain rivers.
 721 Sci. Total Environ. 633, 1089–1104. <https://doi.org/10.1016/j.scitotenv.2018.03.221>
 722 Hobbs, R.J., Higgs, E., Harris, J.A., 2009. Novel ecosystems: implications for conservation
 723 and restoration. Trends Ecol. Evol. 24, 599–605.
 724 Julve, P., 1998. Baseflor. Index botanique, écologique et chorologique de la flore de France.
 725 Version : 2017. <http://perso.wanadoo.fr/philippe.julve/catminat.htm>.
 726 Kattge, J., Díaz, S., Lavorel, S., Prentice, I.C., Leadley, P., BöNisch, G., Garnier, E.,
 727 Westoby, M., Reich, P.B., Wright, I.J., Cornelissen, J.H.C., Violle, C., Harrison, S.P.,
 728 Van BODEGOM, P.M., Reichstein, M., Enquist, B.J., Soudzilovskaia, N.A., Ackerly,
 729 D.D., Anand, M., Atkin, O., Bahn, M., Baker, T.R., Baldocchi, D., Bekker, R.,
 730 Blanco, C.C., Blonder, B., Bond, W.J., Bradstock, R., Bunker, D.E., Casanoves, F.,
 731 Cavender-Bares, J., Chambers, J.Q., Chapin Iii, F.S., Chave, J., Coomes, D.,
 732 Cornwell, W.K., Craine, J.M., Dobrin, B.H., Duarte, L., Durka, W., Elser, J., Esser,
 733 G., Estiarte, M., Fagan, W.F., Fang, J., FernáNdez-MéNdez, F., Fidelis, A., Finegan,
 734 B., Flores, O., Ford, H., Frank, D., Freschet, G.T., Fyllas, N.M., Gallagher, R.V.,
 735 Green, W.A., Gutierrez, A.G., Hickler, T., Higgins, S.I., Hodgson, J.G., Jalili, A.,
 736 Jansen, S., Joly, C.A., Kerkhoff, A.J., Kirkup, D., Kitajima, K., Kleyer, M., Klotz, S.,
 737 Knops, J.M.H., Kramer, K., Kühn, I., Kurokawa, H., Laughlin, D., Lee, T.D.,
 738 Leishman, M., Lens, F., Lenz, T., Lewis, S.L., Lloyd, J., Llusià, J., Louault, F., Ma, S.,
 739 Mahecha, M.D., Manning, P., Massad, T., Medlyn, B.E., Messier, J., Moles, A.T.,
 740 MüLler, S.C., Nadrowski, K., Naeem, S., Niinemets, U., NöLlert, S., NüSke, A.,
 741 Ogaya, R., Oleksyn, J., Onipchenko, V.G., Onoda, Y., OrdoñEz, J., Overbeck, G.,
 742 Ozinga, W.A., PatiñO, S., Paula, S., Pausas, J.G., PeñUelas, J., Phillips, O.L., Pillar,
 743 V., Poorter, H., Poorter, L., Poschlod, P., Prinzing, A., Proulx, R., Rammig, A.,

Reinsch, S., Reu, B., Sack, L., Salgado-Negret, B., Sardans, J., Shiodera, S., Shipley, B., Siefert, A., Sosinski, E., Soussana, J.-F., Swaine, E., Swenson, N., Thompson, K., Thornton, P., Waldram, M., Weiher, E., White, M., White, S., Wright, S.J., Yguel, B., Zaehle, S., Zanne, A.E., Wirth, C., 2011. TRY - a global database of plant traits. *Glob. Change Biol.* 17, 2905–2935. <https://doi.org/10.1111/j.1365-2486.2011.02451.x>

Kleyer, M., Bekker, R.M., Knevel, I.C., Bakker, J.P., Thompson, K., Sonnenschein, M., Poschlod, P., van Groenendael, J.M., Klimeš, L., Klimešová, J., Klotz, S., Rusch, G.M., Hermy, M., Adriaens, D., Boedeltje, G., Bossuyt, B., Dannemann, A., Endels, P., Götzenberger, L., Hodgson, J.G., Jackel, A.-K., Kühn, I., Kunzmann, D., Ozinga, W.A., Römermann, C., Stadler, M., Schlegelmilch, J., Steendam, H.J., Tackenberg, O., Wilmann, B., Cornelissen, J.H.C., Eriksson, O., Garnier, E., Peco, B., 2008. The LEDA Traitbase: a database of life-history traits of the Northwest European flora. *J. Ecol.* 96, 1266–1274. <https://doi.org/10.1111/j.1365-2745.2008.01430.x>

Kui, L., Stella, J.C., 2016. Fluvial sediment burial increases mortality of young riparian trees but induces compensatory growth response in survivors. *For. Ecol. Manag.* 366, 32–40. <https://doi.org/10.1016/j.foreco.2016.02.001>

Kyle, G., Leishman, M.R., 2009. Plant functional trait variation in relation to riparian geomorphology: The importance of disturbance. *Austral Ecol.* 34, 793–804. <https://doi.org/10.1111/j.1442-9993.2009.01988.x>

Laliberté, E., Legendre, P., 2010. A distance-based framework for measuring functional diversity from multiple traits. *Ecology* 91, 299–305.

Lamouroux, N., Gore, J.A., Lepori, F., Statzner, B., 2015. The ecological restoration of large rivers needs science-based, predictive tools meeting public expectations: an overview of the Rhône project. *Freshw. Biol.* 60, 1069–1084. <https://doi.org/10.1111/fw.12553>

768 Lavorel, S., Grigulis, K., McIntyre, S., Williams, N.S.G., Garden, D., Dorrough, J., Berman,
 769 S., Quétier, F., Thébault, A., Bonis, A., 2008. Assessing functional diversity in the
 770 field – methodology matters! *Funct. Ecol.* 22, 134–147.
 771 <https://doi.org/10.1111/j.1365-2435.2007.01339.x>

772 Liébault, F., Piégay, H., 2002. Causes of 20th century channel narrowing in mountain and
 773 piedmont rivers of southeastern France. *Earth Surf. Process. Landf.* 27, 425–444.
 774 <https://doi.org/10.1002/esp.328>

775 McCoy-Sulentic, M.E., Kolb, T.E., Merritt, D.M., Palmquist, E., Ralston, B.E., Sarr, D.A.,
 776 Shafroth, P.B., 2017. Changes in community-level riparian plant traits over inundation
 777 gradients, Colorado River, Grand Canyon. *Wetlands* 37, 635–646.
 778 <https://doi.org/10.1007/s13157-017-0895-3>

779 Molles, M.C., Crawford, C.S., Ellis, L.M., Valett, H.M., Dahm, C.N., 1998. Managed
 780 flooding for riparian ecosystem restoration. *BioScience* 48, 749–756.
 781 <https://doi.org/10.2307/1313337>

782 Naiman, R.J., Bechtold, J.S., Drake, D.C., Latterell, J.J., O’keefe, T.C., Balian, E.V., 2005.
 783 Origins, patterns, and importance of heterogeneity in riparian systems, in: In G.
 784 Lovett, C. G. Jones, M. G. Turner, K. C. Weathers, Editors. *Ecosystem Function in*
 785 *Heterogeneous Landscapes*. Springer, New-York, USA, pp. 279–309.

786 Naiman, R.J., Decamps, H., 1997. The ecology of interfaces: riparian zones. *Annu. Rev. Ecol.*
 787 *Syst.* 28, 621–658.

788 Nakagawa, S., Schielzeth, H., 2013. A general and simple method for obtaining R^2 from
 789 generalized linear mixed-effects models. *Methods Ecol. Evol.* 4, 133–142.
 790 <https://doi.org/10.1111/j.2041-210x.2012.00261.x>

791 Nilsson, C., Brown, R.L., Jansson, R., Merritt, D.M., 2010. The role of hydrochory in
 792 structuring riparian and wetland vegetation. *Biol. Rev.* 85, 837–858.
 793 <https://doi.org/10.1111/j.1469-185X.2010.00129.x>

794 Nilsson, C., Grelsson, G., Johansson, M., Sperens, U., 1989. Patterns of plant species richness
 795 along riverbanks. *Ecology* 70, 77–84. <https://doi.org/10.2307/1938414>

796 Nilsson, C., Reidy, C.A., Dynesius, M., Revenga, C., 2005. Fragmentation and flow
 797 regulation of the world’s large river systems. *Science* 308, 405–408.

798 Olden, J.D., Rooney, T.P., 2006. On defining and quantifying biotic homogenization. *Glob.*
 799 *Ecol. Biogeogr.* 15, 113–120. <https://doi.org/10.1111/j.1466-822X.2006.00214.x>

800 Poff, N.L., Allan, J.D., Bain, M.B., Karr, J.R., Prestegard, K.L., Richter, B.D., Sparks, R.E.,
 801 Stromberg, J.C., 1997. The natural flow regime. *BioScience* 47, 769–784.
 802 <https://doi.org/10.2307/1313099>

803 Poff, N.L., Olden, J.D., Merritt, D.M., Pepin, D.M., 2007. Homogenization of regional river
 804 dynamics by dams and global biodiversity implications. *Proc. Natl. Acad. Sci.* 104,
 805 5732–5737.

806 Poff, N.L., Zimmerman, J.K.H., 2010. Ecological responses to altered flow regimes: a
 807 literature review to inform the science and management of environmental flows.
 808 *Freshw. Biol.* 55, 194–205. <https://doi.org/10.1111/j.1365-2427.2009.02272.x>

809 Polzin, M.L., Rood, S.B., 2006. Effective disturbance: Seedling safe sites and patch
 810 recruitment of riparian cottonwoods after a major flood of a mountain river. *Wetlands*
 811 26, 965–980. [https://doi.org/10.1672/0277-5212\(2006\)26\[965:EDSSSA\]2.0.CO;2](https://doi.org/10.1672/0277-5212(2006)26[965:EDSSSA]2.0.CO;2)

812 R Core Team, 2017. R: A language and environment for statistical computing. R Foundation
 813 for Statistical Computing, Vienna, Austria.

814 Richardson, D.M., Holmes, P.M., Esler, K.J., Galatowitsch, S.M., Stromberg, J.C., Kirkman,
 815 S.P., Pyšek, P., Hobbs, R.J., 2007. Riparian vegetation: degradation, alien plant
 816 invasions, and restoration prospects. *Divers. Distrib.* 13, 126–139.

817 Ricotta, C., Moretti, M., 2011. CWM and Rao's quadratic diversity: a unified framework for
 818 functional ecology. *Oecologia* 167, 181–188. [https://doi.org/10.1007/s00442-011-](https://doi.org/10.1007/s00442-011-1965-5)
 819 1965-5

820 Rohde, S., Schütz, M., Kienast, F., Englmaier, P., 2005. River widening: an approach to
 821 restoring riparian habitats and plant species. *River Res. Appl.* 21, 1075–1094.
 822 <https://doi.org/10.1002/rra.870>

823 Roni, P., Beechie, T.J., Bilby, R.E., Leonetti, F.E., Pollock, M.M., Pess, G.R., 2002. A review
 824 of stream restoration techniques and a hierarchical strategy for prioritizing restoration
 825 in Pacific Northwest watersheds. *North Am. J. Fish. Manag.* 22, 1–20.

826 Schielzeth, H., 2010. Simple means to improve the interpretability of regression coefficients:
 827 Interpretation of regression coefficients. *Methods Ecol. Evol.* 1, 103–113.
 828 <https://doi.org/10.1111/j.2041-210X.2010.00012.x>

829 Shields, F.D., Nunnally, N.R., 1984. Environmental aspects of clearing and snagging. *J.*
 830 *Environ. Eng.* 110, 152–165. [https://doi.org/10.1061/\(ASCE\)0733-](https://doi.org/10.1061/(ASCE)0733-9372(1984)110:1(152))
 831 9372(1984)110:1(152)

832 Silvertown, J., Araya, Y., Gowing, D., 2015. Hydrological niches in terrestrial plant
 833 communities: a review. *J. Ecol.* 103, 93–108. [https://doi.org/10.1111/1365-](https://doi.org/10.1111/1365-2745.12332)
 834 2745.12332

835 Steiger, J., Tabacchi, E., Dufour, S., Corenblit, D., Peiry, J.-L., 2005. Hydrogeomorphic
 836 processes affecting riparian habitat within alluvial channel-floodplain river systems: a
 837 review for the temperate zone. *River Res. Appl.* 21, 719–737.
 838 <https://doi.org/10.1002/rra.879>

839 Weber, C., Schager, E., Peter, A., 2009. Habitat diversity and fish assemblage structure in
840 local river widenings: A case study on a swiss river. *River Res. Appl.* 25, 687–701.
841 <https://doi.org/10.1002/rra.1176>
842 Westoby, M., 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant Soil* 199,
843 213–227.
844 Xiong, S., Nilsson, C., Johansson, M.E., Jansson, R., 2001. Responses of riparian plants to
845 accumulation of silt and plant litter: the importance of plant traits. *J. Veg. Sci.* 12,
846 481–490.

847 Table 1. Top-ranked models predicting the mean and dispersion of ecological and
 848 morphological trait values and the richness and cover of species groups vs elevation and soil
 849 texture gradients as well as maintenance measures on repeatedly cleared relict bars along the
 850 Rhône River (France), as assessed with Akaike's information criterion corrected for small
 851 sample size (AICc). Number of estimated parameters including the intercept (k), AICc, AICc
 852 weight (W), marginal coefficient of determination for fixed effect (R²) and evidence ratio
 853 (ER), i.e., Akaike weight of the best model/Akaike weight of the second best model, are
 854 provided.

Measure	Variable	Top-ranked model	k	AICc	W	R ²	ER
Ecological traits							
CWM:	Soil moisture	elevation + fine_sediments	6	257.5	0.225	0.506	1.02
	Light	elevation	5	138.5	0.199	0.022	0.00
	Nitrogen content	Null	4	138.0	0.258	0.000	0.00
FDis:	Soil moisture	fine_sediments	5	-44.3	0.443	0.166	0.99
	Light	elevation * fine_sediments + maintenance	8	-46.7	0.415	0.146	1.32
	Nitrogen content	fine_sediments * maintenance	7	-108.9	0.745	0.330	0.98
Morphological traits							
CWM:	SLA	Null	4	524.4	0.359	0.000	0.00
	Plant height	elevation * fine_sediments	7	24.5	0.530	0.212	0.99
	Seed mass	elevation	5	43.1	0.438	0.202	0.99
FDis:	SLA	elevation * maintenance + fine_sediments	8	-102.4	0.545	0.395	1.18
	Plant height	Null	4	165.2	0.238	0.000	0.00
	Seed mass	elevation + fine_sediments	6	64.9	0.238	0.059	2.95
Species groups							

Richness:	Hydrochorous	elevation	5	403.3	0.279	0.517	0.94
	Annual	$\text{fine_sediments} * \text{maintenance}$	7	691.6	0.520	0.389	0.99
	Alien	$\text{elevation} * \text{fine_sediments} + \text{maintenance}$	8	478.8	0.354	0.303	1.02
Cover:	Hydrochorous	$\text{elevation} * \text{fine_sediments}$	7	768.0	0.433	0.313	0.94
	Annual	$\text{fine_sediments} * \text{maintenance}$	7	878.6	0.681	0.223	0.97
	Alien	$\text{elevation} * \text{maintenance}$	7	741.8	0.189	0.077	0.97

855

856 Table 2. Average coefficients (Estimate ($\pm$ SE)) and confidence intervals (95% CI) for fixed effects predicting the mean and dispersion of
857 ecological and morphological trait values and the richness and cover of species groups vs elevation and soil texture gradients as well as
858 maintenance measures on repeatedly cleared relict bars along the Rhône River (France). The 95% confidence intervals of coefficients in bold
859 excluded 0.

Measure	Variable	elevation		fine_sediments		elevation:fine_sediments	
		Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)
Ecological traits							
CWM:	Soil moisture	-1.611 (±0.170)	(-1.945; -1.277)	0.331 (±0.202)	(-0.065; 0.726)	NA	NA
	Light	0.138 (±0.089)	(-0.036; 0.312)	-0.068 (±0.107)	(-0.277; 0.141)	NA	NA
	Nitrogen content	NA	NA	0.133 (±0.099)	(-0.061; 0.328)	NA	NA
FDis:	Soil moisture	NA	NA	-0.167 (±0.036)	(-0.238; -0.096)	NA	NA
	Light	0.015 (±0.041)	(-0.065; 0.094)	-0.130 (±0.050)	(-0.227; -0.032)	0.212 (±0.067)	(0.080; 0.344)
	Nitrogen content	NA	NA	-0.097 (±0.028)	(-0.152; -0.042)	NA	NA
Morphological traits							
CWM:	SLA	NA	NA	-0.419 (±0.597)	(-1.590; 0.752)	NA	NA
	Height	-0.188 (±0.056)	(-0.298; -0.078)	0.269 (±0.064)	(0.144; 0.394)	-0.201 (±0.092)	(-0.381; -0.021)

	Seed	0.295 (± 0.054)	(0.189; 0.401)	NA	NA	NA	NA
FDis:	SLA	-0.152 (± 0.030)	(-0.211; -0.093)	-0.087 (± 0.035)	(-0.156; -0.018)	NA	NA
	Height	-0.143 (± 0.106)	(-0.352; 0.065)	0.113 (± 0.117)	(-0.116; 0.342)	NA	NA
	Seed	0.135 (± 0.072)	(-0.005; 0.276)	-0.142 (± 0.079)	(-0.298; 0.013)	NA	NA
Species groups							
Richness:	Hydrochorous	-1.686 (± 0.185)	(-2.048; -1.324)	-0.171 (± 0.221)	(-0.603; 0.261)	NA	NA
	Annual	NA	NA	-0.385 (± 0.138)	(-0.655; -0.115)	NA	NA
	Alien	-0.173 (± 0.123)	(-0.415; 0.068)	-0.293 (± 0.155)	(-0.596; 0.010)	0.524 (± 0.200)	(0.132; 0.915)
Cover:	Hydrochorous	-2.417 (± 0.542)	(-3.479; -1.355)	0.560 (± 0.542)	(-0.503; 1.622)	-2.731 (± 1.115)	(-4.917; -0.545)
	Annual	NA	NA	-0.443 (± 0.218)	(-0.870; -0.016)	NA	NA
	Alien	-0.388 (± 0.245)	(-0.868; 0.093)	0.323 (± 0.279)	(-0.223; 0.869)	NA	NA

Measure	Variable	maintenance		elevation:maintenance		fine_sediments:maintenance	
		Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)
Ecological traits							
CWM:	Soil moisture	-0.179 (±0.234)	(-0.638; 0.280)	-0.353 (±0.275)	(-0.892; 0.186)	NA	NA
	Light	0.111 (±0.109)	(-0.103; 0.324)	NA	NA	NA	NA
	Nitrogen content	NA	NA	NA	NA	NA	NA
FDis:	Soil moisture	NA	NA	NA	NA	NA	NA
	Light	-0.085 (±0.050)	(-0.184; 0.014)	NA	NA	NA	NA
	Nitrogen content	0.103 (±0.037)	(0.030; 0.176)	NA	NA	0.242 (±0.057)	(0.130; 0.354)
Morphological traits							
CWM:	SLA	NA	NA	NA	NA	NA	NA
	Height	NA	NA	NA	NA	NA	NA
	Seed	NA	NA	NA	NA	NA	NA
FDis:	SLA	0.050 (±0.043)	(-0.034; 0.134)	-0.138 (±0.055)	(-0.246; -0.030)	NA	NA
	Height	NA	NA	NA	NA	NA	NA
	Seed	0.069 (±0.109)	(-0.145; 0.283)	NA	NA	NA	NA

Species groups

Richness:	Hydrochorous	0.380 (± 0.273)	(-0.155; 0.915)	NA	NA	NA	NA
	Annual	0.901 (± 0.203)	(0.503; 1.299)	NA	NA	0.722 (± 0.289)	(0.156; 1.288)
	Alien	0.633 (± 0.176)	(0.289; 0.978)	NA	NA	0.516 (± 0.282)	(-0.037; 1.069)
Cover:	Hydrochorous	-0.710 (± 0.512)	(-1.714; 0.294)	NA	NA	NA	NA
	Annual	0.843 (± 0.257)	(0.339; 1.347)	NA	NA	1.257 (± 0.411)	(0.451; 2.063)
	Alien	0.529 (± 0.359)	(-0.175; 1.232)	-0.889 (± 0.459)	(-1.788; 0.010)	0.852 (± 0.496)	(-0.120; 1.825)

Table 3. Top-ranked models predicting the mean and dispersion of ecological and morphological trait values and the richness and cover of species groups vs elevation and soil texture gradients as well as the type of geomorphic surface along the Rhône River and tributaries (France), as assessed with Akaike's information criterion corrected for small sample size (AICc). Number of estimated parameters including the intercept (k), AICc, AICc weight (W), marginal coefficient of determination for fixed effect (R²) and evidence ratio (ER), i.e., Akaike weight of the best model/Akaike weight of the second best model, are provided.

Measure	Variable	Top-ranked model	k	AICc	W	R ²	ER
Ecological traits							
CWM:	Soil moisture	elevation * type	9	468.2	0.654	0.560	1.89
	Light	elevation + type	7	230.1	0.372	0.143	1.94
	Nitrogen content	elevation * type	9	276.9	0.740	0.171	2.90
FDis:	Soil moisture	elevation * type + fine_sediments	10	-62.8	0.788	0.197	9.69
	Light	elevation * fine_sediments	7	-68.0	0.565	0.047	4.06
	Nitrogen content	fine_sediments	5	-181.3	0.431	0.050	1.83
Morphological traits							
CWM:	SLA	elevation * type + fine_sediments	10	980.2	0.570	0.112	1.34
	Plant height	elevation * fine_sediments + type	9	-33.7	0.541	0.175	2.44
	Seed mass	elevation * type	9	128.7	0.478	0.129	3.02
FDis:	SLA	elevation + fine_sediments	6	-180.8	0.395	0.247	1.76
	Plant height	Null	4	-358.2	0.259	0.000	1.54
	Seed mass	elevation + fine_sediments	6	134.9	0.305	0.037	1.81
Species groups							

Richness:	Hydrochorous	elevation * type	9	748.7	0.608	0.544	1.55
	Annual	elevation * fine_sediments	7	1367.2	0.335	0.049	1.34
	Alien	Null	4	905.8	0.201	0.000	1.14
Cover:	Hydrochorous	elevation * type	9	1342.4	0.517	0.225	1.22
	Annual	elevation + fine_sediments	6	1801.3	0.180	0.026	1.51
	Alien	elevation * type	9	1542.2	0.293	0.126	1.21

870

871 Table 4. Average coefficients (Estimate ($\pm$ SE)) and confidence intervals (95% CI) for fixed effects predicting the mean and dispersion of
872 ecological and morphological trait values and the richness and cover of species groups vs elevation and soil texture gradients as well as
873 geomorphic surface types (Cleared = repeatedly cleared relict bars, Reprof = newly reprofiled banks, Natura = naturally rejuvenated bars) along
874 the Rhône River and tributaries (France). The 95% confidence interval of coefficients in bold excluded 0.

Measure	Variable	elevation		fine_sediments		Elevation:fine_sediments	
		Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)
Ecological traits							
CWM:	Soil moisture	-1.817 (±0.174)	(-2.158; -1.477)	0.130 (±0.133)	(-0.131; 0.390)	NA	NA
	Light	0.176 (±0.061)	(0.057; 0.295)	-0.069 (±0.072)	(-0.211; 0.073)	NA	NA
	Nitrogen content	0.016 (±0.102)	(-0.184; 0.216)	NA	NA	NA	NA
FDis:	Soil moisture	-0.039 (±0.048)	(-0.133; 0.055)	-0.098 (±0.035)	(-0.167; -0.029)	NA	NA
	Light	-0.002 (±0.030)	(-0.061; 0.057)	-0.067 (±0.035)	(-0.136; 0.002)	0.167 (±0.058)	(0.053; 0.281)
	Nitrogen content	0.022 (±0.023)	(-0.023; 0.067)	-0.075 (±0.025)	(-0.125; -0.026)	NA	NA
Morphological traits							
CWM:	SLA	0.184 (±0.630)	(-1.052; 1.419)	-0.805 (±0.478)	(-1.743; 0.132)	NA	NA
	Height	-0.101 (±0.065)	(-0.228; 0.027)	0.138 (±0.039)	(0.062; 0.213)	-0.214 (±0.063)	(-0.337; -0.091)

	Seed	0.349 (± 0.070)	(0.212; 0.486)	NA	NA	NA	NA
FDis:	SLA	-0.135 (± 0.023)	(-0.179; -0.090)	-0.082 (± 0.027)	(-0.135; -0.029)	-0.045 (± 0.044)	(-0.132; 0.042)
	Height	-0.460 (± 0.028)	(-0.515; -0.406)	0.102 (± 0.008)	(0.086; 0.119)	NA	NA
	Seed	0.110 (± 0.050)	(0.013; 0.208)	-0.112 (± 0.062)	(-0.233; 0.009)	0.056 (± 0.096)	(-0.132; 0.243)
Species groups							
Richness:	Hydrochorous	-1.948 (± 0.215)	(-2.370; -1.527)	-0.172 (± 0.150)	(-0.466; 0.122)	NA	NA
	Annual	0.101 (± 0.086)	(-0.069; 0.270)	-0.324 (± 0.126)	(-0.571; -0.077)	0.328 (± 0.174)	(-0.012; 0.669)
	Alien	-0.091 (± 0.072)	(-0.233; 0.050)	-0.100 (± 0.097)	(-0.289; 0.089)	NA	NA
Cover:	Hydrochorous	-3.037 (± 0.536)	(-4.088; -1.987)	0.460 (± 0.348)	(-0.223; 1.142)	NA	NA
	Annual	0.256 (± 0.151)	(-0.039; 0.551)	-0.330 (± 0.177)	(-0.676; 0.017)	0.150 (± 0.276)	(-0.391; 0.691)
	Alien	-0.460 (± 0.028)	(-0.515; -0.406)	0.102 (± 0.008)	(0.086; 0.119)	NA	NA

Measure	Variable	Cleared-Reprof		Cleared-Natura		Elevation:Cleared-Reprof	
		Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)
Ecological traits							
CWM:	Soil moisture	-0.187 (±0.208)	(-0.595; 0.221)	-1.249 (±0.212)	(-1.664; -0.834)	0.111 (±0.231)	(0.658; 1.564)
	Light	0.269 (±0.122)	(0.030; 0.509)	0.300 (±0.124)	(0.057; 0.542)	NA	NA
	Nitrogen content	0.195 (±0.140)	(-0.079; 0.469)	-0.285 (±0.140)	(-0.559; -0.011)	-0.016 (±0.140)	(-0.290; 0.258)
FDis:	Soil moisture	-0.084 (±0.049)	(-0.180; 0.012)	-0.089 (±0.051)	(-0.189; 0.011)	-0.054 (±0.063)	(-0.177; 0.069)
	Light	NA	NA	NA	NA	NA	NA
	Nitrogen content	NA	NA	NA	NA	NA	NA
Morphological traits							
CWM:	SLA	-0.290 (±1.348)	(-2.933; 2.352)	0.301 (±1.366)	(-2.376; 2.979)	0.078 (±0.806)	(-1.502; 1.658)
	Height	0.003 (±0.046)	(-0.086; 0.093)	-0.113 (±0.049)	(-0.209; -0.017)	0.144 (±0.068)	(0.010; 0.278)
	Seed	-0.036 (±0.121)	(-0.273; 0.201)	0.149 (±0.121)	(-0.088; 0.386)	-0.262 (±0.096)	(-0.450; -0.074)
FDis:	SLA	0.025 (±0.039)	(-0.051; 0.102)	-0.051 (±0.040)	(-0.130; 0.027)	NA	NA
	Height	0.929 (±0.017)	(0.895; 0.963)	0.713 (±0.024)	(0.665; 0.761)	0.462 (±0.021)	(0.420; 0.504)
	Seed	0.069 (±0.127)	(-0.180; 0.318)	-0.065 (±0.138)	(-0.335; 0.206)	NA	NA

Species groups

Richness:	Hydrochorous	0.543 (± 0.233)	(0.087; 1.000)	-0.859 (± 0.292)	(-1.432; -0.286)	0.154 (± 0.279)	(0.607; 1.700)
	Annual	NA	NA	NA	NA	NA	NA
	Alien	0.372 (± 0.229)	(-0.078; 0.822)	0.155 (± 0.230)	(-0.295; 0.605)	NA	NA
Cover:	Hydrochorous	0.081 (± 0.552)	(-1.001; 1.163)	-1.403 (± 0.578)	(-2.536; -0.270)	2.304 (± 0.631)	(1.069; 3.540)
	Annual	0.609 (± 0.372)	(-0.121; 1.338)	0.674 (± 0.378)	(-0.066; 1.414)	NA	NA
	Alien	0.929 (± 0.017)	(0.895; 0.963)	0.713 (± 0.024)	(0.665; 0.761)	0.462 (± 0.021)	(0.420; 0.504)

Measure	Variable	Elevation:Cleared-Natura		fine_sediments:Cleared-Reprof		fine_sediments:Cleared-Natura	
		Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)	Estimate (±SE)	(95% CI)
Ecological traits							
CWM:	Soil moisture	0.259 (±0.326)	(-0.381; 0.898)	NA	NA	NA	NA
	Light	NA	NA	NA	NA	NA	NA
	Nitrogen content	-0.808 (±0.201)	(-1.202; -0.414)	NA	NA	NA	NA
FDis:	Soil moisture	-0.312 (±0.087)	(-0.483; -0.141)	NA	NA	NA	NA
	Light	NA	NA	NA	NA	NA	NA
	Nitrogen content	NA	NA	NA	NA	NA	NA
Morphological traits							
CWM:	SLA	-5.287 (±1.130)	(-7.503; -3.072)	NA	NA	NA	NA
	Height	0.324 (±0.095)	(0.138; 0.509)	NA	NA	NA	NA
	Seed	-0.365 (±0.139)	(-0.637; -0.093)	NA	NA	NA	NA
FDis:	SLA	NA	NA	NA	NA	NA	NA
	Height	0.187 (±0.016)	(1.154; 1.219)	NA	NA	NA	NA
	Seed	NA	NA	0.172 (±0.144)	(-0.111; 0.454)	-0.420 (±0.190)	(-0.793; -0.047)

Species groups

Richness:	Hydrochorous	-0.370 (± 0.559)	(-1.466; 0.726)	NA	NA	NA	NA
	Annual	NA	NA	NA	NA	NA	NA
	Alien	NA	NA	NA	NA	NA	NA
Cover:	Hydrochorous	-0.031 (± 0.987)	(-1.966; 1.904)	NA	NA	NA	NA
	Annual	NA	NA	NA	NA	NA	NA
	Alien	1.187 (± 0.016)	(1.154; 1.219)	NA	NA	NA	NA

877 Figure 1. Location of the study area within the Rhône River watershed, distribution of
 878 sampled geomorphic surfaces along the section of the Rhône River and the tributary rivers
 879 studied (Ain, Drôme and Doux Rivers) and form of the sampling design used to survey plant
 880 communities.

881 Figure 2. Variations in elevation and in the proportion of fine sediments in relation to A)
 882 maintenance measures along repeatedly cleared relict bars of the Rhône riverbed (n quadrats
 883 = 108) and B) the type of geomorphic surface (n quadrats = 204).

884

885 Figure 3. Interaction effect between elevation and soil texture on ecological and
 886 morphological traits and species groups on repeatedly cleared relict bars along the Rhône
 887 River (France). To provide representation, the proportion of fine sediments (continuous
 888 variable), has been divided into two equally sized groups.

889 Figure 4. Interaction effect between environmental gradients and maintenance measures on
 890 ecological and morphological traits and species groups on repeatedly cleared relict bars along
 891 the Rhône River (France).

892 Figure 5. Interaction effect between environmental gradients and the type of geomorphic
893 surface, i.e., A) repeatedly cleared relict bars vs newly reprofiled banks and B) repeatedly
894 cleared relict bars vs naturally rejuvenated bars, on ecological and morphological traits and
895 species groups along the Rhône River and tributaries (France).