Right or left? Determining the hand holding the tool from use traces

Alice Rodriguez, Emmanuelle Pouydebat, M. Gema Chacón, Marie-Hélène Moncel, Raphael Cornette, Ameline Bardo, Laurence Chèze, Radu Iovita, Antony Borel

► To cite this version:

Alice Rodriguez, Emmanuelle Pouydebat, M. Gema Chacón, Marie-Hélène Moncel, Raphael Cornette, et al.. Right or left? Determining the hand holding the tool from use traces. *Journal of Archaeological Science: Reports*, 2020, 31, 27p. 10.1016/j.jasrep.2020.102316 . hal-02615493

HAL Id: hal-02615493

<https://hal.science/hal-02615493>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Right or left? Determining the hand holding the tool from use traces.

Alice Rodriguez^{ah}, Emmanuelle Pouydebat^b, M. Gema Chacón^{cda}, Marie-Hélène Moncel^a, Raphaël Cornette^e, Ameline Bardo^f, Laurence Chèze^g, Radu Iovita^{hi}, Antony Borel^{aj}

a) Histoire Naturelle de l'Homme Préhistorique (HNHP), Muséum national d'histoire naturelle, CNRS, UPVD, 1 Rue René Panhard, 75013 Paris, France.

b) Mécanismes adaptatifs et évolution (MECADEV), Muséum national d'Histoire naturelle, CNRS, 57 rue Cuvier, CP 55, 75005 Paris, France.

c) Institut Català de Paleoecologia Humana i Evolució Social (IPHES), Zona Educacional 4, Campus Sescelades URV (Edifici W3), 43007 Tarragona, Spain.

d) Universitat Rovira i Virgili (URV), Àrea de Prehistòria, Avinguda Catalunya 35, 43002, Tarragona, Spain.

e) Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, 57 rue Cuvier, CP 50, 75005 Paris, France.

f) Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Marlowe Building, Canterbury, CT2 7NR, United Kingdom.

g) Univ. Lyon, Université Claude Bernard Lyon 1, IFSTTAR, LBMC UMR_T9406, F69622, Lyon, France.

h) Anthrotopography Laboratory, Center for the Study of Human Origins, Department of Anthropology, New York University, New York City, USA.

i) Early Prehistory and Quaternary Ecology, University of Tübingen, Tübingen, Germany.

j) Institute of Archaeological Sciences, Eötvös Loránd University, Múzeum krt. 4/b, 1088 Budapest, Hungary.

Corresponding authors: Alice Rodriguez; Antony Borel

E-mail addresses: aar596@nyu.edu ; antony.borel@mnhn.fr

Abstract:

Currently, approximately 90% of the human population is right-handed. This handedness is due to the lateralization of the cerebral hemispheres and is controlled by brain areas involved in complex motor tasks such as making stone tools or in language. In addition to describing the evolution of laterality in humans, identifying hand preference in fossil hominids can improve our understanding of the emergence and development of complex cognitive faculties during evolution. Several fields of prehistory like palaeoanthropology or lithic analysis have already investigated handedness in fossils hominins but they face limitations due to either the incomplete or the composite state of the skeleton remains or to results replication or method application failure. Wear analysis could provide new complementary data about hand preference evolution and the development of certain complex

39 cognitive functions using indirect evidence (use traces, micro-scars in particular) of the hand
40 holding the stone tool during use. Controlled experiment has been carried out in order to establish
41 a reference collection of tools used with the left and tools used with the right hand. Wear analysis
42 was performed on this corpus using “classical” microscopic approach and geometric morphometric
43 analysis. A machine learning algorithm, the k-NN method, was applied to verify if use traces
44 (micro-scars) could help determine the hand holding the tool during use. The best model, based on
45 parameters referring to invasiveness of micro-scars, was able to correctly determine the hand
46 holding the tool with 75% accuracy.

47

48

49 1. Introduction:

50

51 Hand preference is defined by the hand with which an individual performs a specific task
52 (e.g. Annett, 1985; Uomini and Gowlett, 2013). Several studies have demonstrated that hand
53 preference is linked to the lateralization of the cerebral hemispheres and is related to areas of the
54 brain similar to those involved in the control of other essential functions such as language (e.g.
55 Corballis, 2003; Meguerditchian et al., 2013; Stout et al., 2000; Uomini and Meyer, 2013; Willems
56 and Hagoort, 2007). Thus, in addition to describing the evolution of laterality in humans, identifying
57 hand preference in fossil hominids can improve our understanding of the emergence and
58 development of complex cognitive faculties during evolution (e.g. Steele and Uomini, 2009; Stout
59 et al., 2008).

60

61 Currently, between 74% and 96% of the human population preferentially uses the right
62 hand (Annett, 1985; Llaurens et al., 2009; Porac and Coren, 1981). Lefthanders are a minority in
63 all human populations. The highest rate of left-handers recorded in a population is found among
64 the Eipo in Papua New Guinea and is estimated at 26.9% (Faurie et al., 2005).

65 The control of fine manual actions (called “complex” motor actions) is attributed to the left
66 hemisphere which would explain the preferential use of the right hand for this type of action (e.g.
67 Greenfield, 1991; Johnson-Frey et al., 2005; Lewis, 2006; Schluter et al., 2001). Some authors
68 have thus hypothesized the development of a strong manual preference in relation to the
69 manufacture and complexity of tools (Corballis, 1987; Stock et al., 2013; Uomini and Gowlett,
70 2013; Uomini, 2009). Other studies show a greater effect of the nature and demands of the task as
71 well as the type of object involved and the speed of the action (Forrester et al., 2013, 2012, 2011;
72 Pouydebat et al., 2014; Quaresmini et al., 2014; Rogers, 2009). The complexity of the task is
73 therefore probably not the only criterion explaining the emergence of a strong manual preference.
74 Establishing a direct link between strong manual preference and the complexity of (stone) tools is
75 difficult and, so far, concrete archaeological data do not support it. Moreover, few studies highlight

the fact that the right hemisphere is also involved and essential in activities such as tool use (Frey and Gerry, 2006; Hartmann et al., 2005; Stout and Chaminade, 2012)

In non-human primates, especially great apes, similarly strong lateralization as in humans is missing (Vallortigara and Rogers, 2005). This differs according to the species but also within the same species at an individual level or a population level (Cashmore et al., 2008; Chapelain et al., 2011; Hopkins, 2006; Marzke, 2013; McManus, 2009). Differences in hand preference among non-human primates appear to be due to complex relationships between various factors, such as the environment (life in the wild or captivity), or the complexity and nature of the task (type, speed) (e.g. Fagot and Vauclair, 1991; Hopkins et al., 2011; McGrew and Marchant, 1997; Mosquera et al., 2012; Pouydebat et al., 2014).

The origin of this peculiarity during human evolution is for the moment unknown. So far, data about fossil hominin hand preference are mainly coming from the skeleton, either directly inferred from forelimb asymmetries or indirectly from the skull. The examination of bone asymmetries on the upper limbs of fossil hominids could deliver direct indications of hand preference. However, as suggested or stated in several studies, observed differences in hand entheses (the surfaces where ligaments are attached on bones) among individuals with the same occupation could be due to numerous variables including inter-individual genetic variability, nutrition or hormone levels, and not just manual activities (Foster et al., 2014; Karakostis et al., 2018, 2017; Karakostis and Lorenzo, 2016).

On the other hand, other measurements of upper limb asymmetries have been used to identify human hand preference (Shaw, 2011; Trinkaus et al., 1994). Based on such observation, Sparacello et al. (2017) studied a sample of humeri coming from 107 Late Pleistocene individuals and obtained a distribution of hand preference similar to the distribution obtained with recent human samples (i.e. right: 74.8%, left: 15.0%, ambiguous: 10.3%). Others suggested for example that the articular constraints model could help in the identification of hand preference (Lazenby et al., 2008) or that the analysis of trabecular bone could contribute to the identification of right/left directional asymmetry (Stephens et al., 2016).

Moving on to the skull, analyses of cranial endocranial asymmetries analyses (e.g. Balzeau et al., 2012; Holloway, 1981; Holloway and De La Costelareymondie, 1982; Poza-Rey et al., 2017) suggested possible relationships between hand preference, endocranial shape, and functional capacities in hominins. Other studies, investigating striations on teeth surfaces, show that even for the Middle Palaeolithic and species such as *Homo erectus* in Asia, or *Homo heidelbergensis* and *Homo neanderthalensis* in Europe, the right hand was already the chosen hand for the majority of individuals, both adults or young individuals (e.g. Bermúdez de Castro et al., 2003, 1988; Condemi et al., 2017; Estalrich and Rosas, 2013; Fiore et al., 2015; Frayer et al., 2012, 2010; Lozano et al., 2009; Poza-Rey et al., 2017; Xing et al., 2017). This could be the case as early as 1.8 Ma as suggested by the study of labial striations on the OH-65 fossil's anterior teeth (Lozano et al., 2017).

114 For younger periods, analysis of wall-paintings and striations on bone tools may give clues about
115 the hand preference of their authors or users respectively (see Spenneman, 1984 for an example
116 concerning Neolithic).

117

118 These studies do not provide data supporting a ratio of left- and right-handers clearly
119 different from now in Prehistory (see Llaurens et al., 2009). However, the more we go back in time
120 the more the application of these methods to extinct hominin species have serious limitations.
121 Relatively bad conservation of skeletal remains often prevents any analysis of traces that are
122 relevant for determining hand preference. Also, apart from the limited number of individuals which
123 could be studied (see table 2.1 of Faurie et al., 2016 for a summary of fossil hominin data for
124 laterality and corresponding references) palaeoanthropology faces the problem of the partial
125 skeleton remains which implies sometimes to study composite anatomical elements composed of
126 bones coming from different individuals (see Cashmore, 2009).

127 On the contrary, stone tools are often abundant in Palaeolithic sites. The data they could
128 provide regarding hand preference are therefore likely to be statistically more representative if
129 whole sets of assemblages are analysed. Different studies have intended to set up a methodology
130 to determine hand preference based on the observation of manufacture features appearing on
131 stone tools (size and shape of flake scatter patterns: Newcomer and Sieveking, 1980; dorsal cortex
132 position: Toth, 1985; occurrence of cortex and relic margins: Bradley and Sampson, 1986;
133 orientation of the cone of percussion: Rugg and Mullane, 2001; resharpening techniques: Cornford,
134 1986; combination of technical features: Bargalló et al., 2018, 2017; Bargalló and Mosquera,
135 2014). These methods give variable results and a recent work from Ruck et al. (2015), applying
136 Toth's, Rugg and Mullane's and Bargalló and Mosquera's methods, showed that these methods
137 were not able to determine the hand holding the hammer for the studied flakes production better
138 than a random guess (see also Patterson and Sollberger, 1986; Pobiner, 1999; Uomini, 2011,
139 2001).

140 The identification of the knapper hand preference based on the spatial organization of
141 debitage remains have also been investigated giving successful results in experimental conditions
142 but the authors highlight the difficulties in applying such method in an archaeological context
143 (Bargalló et al., 2018).

144

145 Some of these methods should be investigated further in order to improve their rate of
146 correct determination and their applicability to archaeological samples. However, producing a
147 stone flake is a quick and punctual activity. Therefore, any variation during this short production
148 moment might modify the types of features appearing on the flake. The poor rates of success in
149 identifying hand preference from patterns in flake production may be due to the high variability of
150 factors influencing the production of any particular flake. On the other hand, a repeated activity or
151 movement is likely to leave behind a stronger "signature" of the hand preference. Therefore,

152 focusing on stone tool use rather than stone tool production could provide better results in
153 identifying hominids hand preference. A handful of studies have attempted to extract hand
154 preference from stone tool use (based on asymmetries of the tool: Brinton, 1896; based on
155 direction of rotation of boring artefacts: Cahen et al., 1979; Cahen and Keeley, 1980; Keeley, 1977;
156 see also Uomini, 2008). Others simply discuss why it is difficult to do so (D'Errico, 1992; Semenov,
157 1970; Weber, 1990). However, to our knowledge, most of them only briefly mention this question
158 and none is based on an experimental reference collection that was statistically tested to allow the
159 identification of specific criteria corresponding to the use of the tool in the right or left hand.

160 Here, we propose to examine if hand preference can be identified from use-wear on stone
161 tools through qualitative (traditional wear observation) and quantitative (measurements and
162 geometric morphometric) wear descriptors. A machine learning algorithm is used for the first time
163 on these data in order to test which of the descriptors is/are the most relevant to recognize the
164 hand holding the tool based on the traces of use.

165

166

167 2. Materials and methods

168 2.1. Experimental setup

169 *Tools:*

170 Numerous variables (morphology and raw material type of the tool, worked material,
171 working angle, direction of movement, etc.) can have an effect on the formation of traces on the
172 surface of a stone tool. In order to characterise traces of use related to hand preference but also
173 quantify their variability, it is first essential to limit the effect of these variables and particularly the
174 effect of the form of the tool. Therefore, the form of the tool was standardized using soda-lime
175 glass replicas of a Levallois point (Iovita et al., 2014; figure 1). The use of this glass has the
176 advantage to simulate the type of conchoidal fracture of many raw materials used in prehistory.
177 These artefacts measured 64.5 mm long, 36.5 mm at the maximal width and 6 mm at the maximal
178 thickness, with a weight of 15.8 g on average.

179

180 **Figure 1 (1 column)**

181

182 *Users:*

183 A total of 20 volunteers participated in the experiment, 10 right-handed subjects (5 women
184 and 5 men) and 10 left-handed subjects (5 women and 5 men). They were beginners in using
185 stone tools; none of them had used prehistoric tools before. The subjects were not aware of the
186 fact that the aim of the experiment was related to hand preference in order to limit modification of
187 their behaviour. Their hand preference was documented along with a number of other variables,
188 including name, age, sex, size, weight, hand size, circumference of the last phalanx of each finger

189 of the hand holding the tool, palm length, palm width, expertise, previous injuries in order to
190 prevent them from identifying the aim of the study.

191

192 *Worked material:*

193 Wooden (*Pinus*) sticks with no bark were provided to each subject. These sticks were of
194 standardized diameter (18 mm) and length (50 cm).

195

196 *Task:*

197 Each individual had to sit on a chair and to saw wooden sticks with three different tools.
198 Each tool was used for a total duration of 20 minutes. The number of strokes was not constrained
199 but number of strokes (forward + backward = 1) per minute was measured during the 5th, 10th and
200 15th minutes in order to approximate the speed of the sawing movement (in mvt/min.) (Whole
201 experiment: min = 98.33 mvt/min., max = 191.33 mvt/min., mean = 135.96 mvt/min., SD = 23.59
202 mvt/min.; for left hand users: min = 98.33 mvt/min., max = 191.33 mvt/min., mean = 141.74
203 mvt/min., SD = 25.37 mvt/min.; for right hand users: min = 100.3 mvt/min., max = 166 mvt/min.,
204 mean = 129.9 mvt/min., SD = 20.41 mvt/min.; see the dataset in supplementary material). Before
205 the experiment proper, a five-minute test was carried out without any instructions to observe how
206 individuals hold and use the tool spontaneously, and to evaluate the constraints of our controlled
207 experiment on the tool grasping mode. The first tool of the proper experiment was used with the
208 only instruction to use the predefined edge. For the second and third tools, the participants had to
209 use the predefined edge and keep the wooden stick horizontal on their knees. Each experiment
210 was recorded individually and documented by one observer using a Samsung NV24HD video
211 camera.

212

213 2.2. Analysis

214 *Description of micro-scars*

215 Each stone tool was observed with a stereomicroscope (Zeiss Discovery V8 with Plan S
216 1.0x objective; FWD = 81 mm; it is equipped with a ToupTek camera with a resolution of 5
217 MPixels) before use at 10x (objective FOV = 23 mm) and 80x (objective FOV = 2.9 mm)
218 magnifications. Then, the tools were cleaned (in a solution of DERQUIM LM02 phosphate free
219 neutral soap at 5% in ultrasonic-bath during 15 min), and moulded (using Heraeus PROVIL®novo
220 Light Regular silicone) and casted (using Axson F180 polyurethane resin) to keep a record of their
221 original morphology before use. The tools were then used for a total duration of 20 minutes each.
222 After use, they were cleaned again (same procedure as before use) and observed at 10x and 80x
223 magnifications (with the Zeiss Discovery V8).

224 Micro-scars were documented before and after use on both sides of the tool. Their amount,
225 position (ventral or dorsal face), type, orientation, and (position of) maximum extent on the surface
226 perpendicularly to the edge were recorded (Figure 2). The type of micro-fractures was described

227 following the typology of the Ho Ho Committee (1979): fracture, feathered and step. We added the
228 type “abrupt” to refer to scars which were not straight as a fracture but too steep to enter in the
229 category “feathered”. Three categories were used for the orientation of the scars: perpendicular to
230 the edge, oriented distally (towards the point) and proximally (towards the butt).

231 Statistical tests were performed in R (R Core Team, 2019). As the assumptions of normality
232 and equality of variance for t-test were not met, we used non-parametric Mann-Whitney U tests to
233 verify if the number of micro-scars and if their invasiveness were different depending on the hand
234 holding the tool during use. The Cochran’s rule being respected, we used chi-square test to
235 examine if the type of micro-scars, their orientation and the face where they appeared were
236 independent of the user, of the user’s sex and of the hand holding the tool during use.

237

238 **Figure 2 (1 column)**

239

240 *Edge modification visualization and quantification*

241 Photograph of each tool was taken before and after use. The same camera was not
242 available during the whole study. Pictures were taken with a Nikon D300 equipped with a Tamron
243 16-300mm F/3.5-6.3 Di II VC PZD objective before use (pictures were taken at a focal length of
244 200mm) and with a NIKON 7100 with an AF-D Micro NIKKOR 60mm 1:2.8G ED objective after
245 use.

246 As they were taken with different conditions, the two batches of photos (i.e. from before and
247 after use) were first scaled. They were then binarized with Adobe Photoshop. The Momocs
248 (Bonhomme et al., 2014) and geomorph (Adams et al., 2019) libraries for R were used for the
249 analysis of the shape of the tool edges. The outlines of the tools were automatically extracted from
250 the binarized images. The outlines were then aligned and cut to keep only the area of interest
251 displaying the modifications due to use: the used edge of each tool (Figure 3). The remaining
252 curves corresponding to the used edges of each tool were defined by two landmarks, one at each
253 extremity, and by 1000 equidistant "sliding-semilandmarks" distributed between these two points. A
254 high number of points was chosen because the sought shape differences are due to the micro-
255 scarring of the edges and can be very fine. The space between each "semilandmark" was then
256 optimized by allowing them to slip between the previous and the next point in order to minimize the
257 Procruste distance with the pseudo-homologous point of the Procrustes mean shape (see Gunz
258 and Mitteroecker, 2013). This optimization aims to limit the effect of the former arbitrary spacing of
259 the points (Gunz and Mitteroecker, 2013; Perez et al., 2006), here equidistant. This optimization
260 and the superposition of the curves were carried out by Generalized Procrustean Analysis (GPA).
261 The new coordinates obtained by GPA were used to calculate a Principal Component Analysis
262 (PCA) and to visualize, with transformation grids (Klingenberg, 2013), the axes on which the shape
263 variations were the most important according to the manual preference. A Procrustes multivariate
264 analysis of variance (Procrustes ANOVA using the function “procD.lm” of package geomorph in R)

265 was used to test if the shape differences visualized according to the recorded factors (e.g. hand
266 holding the tool during use) were statistically significant. In other words, it allowed us to check if
267 specific scar distributions and specific used edge morphologies can be linked to the hand holding
268 the tool during use (i.e. left or right hand). The relationship between individuals' sex and the
269 resulting used edge shape was also tested.

270

271

Figure 3 (2 columns)

272

273 *k-Nearest Neighbours model*

274 In order to verify if the qualitative and quantitative descriptors of scars could be used to
275 discriminate a tool used with the right hand from a tool used with the left hand, we used the k
276 Nearest Neighbours (k-NN) machine learning algorithm. It was computed using the “caret” library
277 (Kuhn et al., 2019) in R. This algorithm uses Euclidean distances to find the k-Nearest Neighbours
278 of the artefact to be classified. The class which is the most frequent among the k Nearest
279 Neighbours is given to the unknown artefact (see Domínguez-Rodrigo and Baquedano, 2018 for
280 an example of application on cut-marks on bones). Twenty-one combinations of different
281 descriptors were tested:

282 1) Number of scars on the ventral and dorsal faces of the tool

283 2) types of scars on both faces

284 3) types of scars on dorsal face

285 4) types of scars on ventral face

286 5) directions of scars on both faces

287 6) directions of scars on dorsal face

288 7) directions of scars on ventral face

289 8) types and directions of scars on both faces

290 9) number, types and directions of scars on both faces

291 10) the 59 first principal components of the PCA on geometric morphometric data

292 11) the 2 first principal components of the PCA on geometric morphometric data

293 12) the 59 first principal components of the PCA on geometric morphometric data and the
294 number of scars on both faces

295 13) position of the scar with the maximum extent on the surface perpendicularly to the edge
296 on ventral face

297 14) position of the scar with the maximum extent on the surface perpendicularly to the edge
298 on dorsal face

299 15) position of the scar with the maximum extent on the surface perpendicularly to the edge
300 on both faces

301 16) maximum scar extent on the surface perpendicularly to the edge on ventral face

302 17) maximum scar extent on the surface perpendicularly to the edge on dorsal face

18) maximum scar extent on the surface perpendicularly to the edge on both faces

19) maximum scar extent and position of the scar with the maximum extent on the surface perpendicularly to the edge on both faces

20) all descriptors from microscopic observation

21) all descriptors from microscopic observation and geometric morphometrics

We normalized the descriptors so their values were comprised between 0 and 1 to avoid emphasizing descriptors with larger ranges of values. Each configuration was tested with each odd number between 3 and 59 as k value (3 being the minimum odd number for such model and 59 the maximum in our case as we had 60 artefacts). The k value of the final model was chosen based on the obtained accuracy and kappa for each model. As our sample size ($N = 60$) is quite small, we used leave-on-out cross-validation method to evaluate the models. In this method, the whole sample set of analysed artefacts but one is used as a training set and the removed artefact is used as a testing set. This procedure is repeated as many times as the number of artefacts in the sample set. In addition, we used kappa statistics (value between -1 and 1 which measures the agreement between observed and expected accuracy; negative values indicate disagreement, and the degree of agreement increase from >0 to 1), sensitivity (a measure of the proportion of attributed classes - i.e. positives - which are correctly attributed – i.e. true positives) and specificity (a measure of the proportion of rejected classes – i.e. negatives - which are correctly rejected – i.e. true negatives) to evaluate the performance of the selected model.

3. Results

3.1. Scars description

The scars qualitative description showed that independently of the hand holding the tool during use: 1) tools have more extended scars on their dorsal face than on the ventral face, 2) tools have mostly continuous scars on the dorsal face and discontinuous scars on the ventral face, 3) scars can be distributed all along the used edge and 4) scars are mostly perpendicular to the edge. Therefore, no significant difference of position, type and orientation of scars between tools used by left-handers and right-handers was noticed (Figure 4 and Supplementary figures 1 to 7).

However, right-handed subjects produced significantly more micro-scars than left-handed subjects ($\bar{X} = 81.42$, $sd = 39.84$ for the tools used by right-handers versus $\bar{X} = 53.04$, $sd = 18.60$ for the left-handers; Mann-Whitney U one-tailed test: $W = 240$, $p\text{-value} = 0.0011$). Micro-scars were also significantly less invasive on tools used by left-handed subjects ($\bar{X} = 2.45$ mm, $sd = 1.13$ mm for the tools used by right-handers versus $\bar{X} = 1.63$ mm, $sd = 0.73$ mm for the left-handers; Mann-Whitney U one-tailed test: $W = 771$, $p\text{-value} < 0.0001$).

The hypothesis of independence between the hand holding the tool and the type of micro-scars appearing on the surface of the tool could be rejected ($X^2 = 43.371$, $df = 4$, $p\text{-value} < 0.0001$). However, independence could not be rejected concerning the hand holding the tool and

341 the orientation of micro-scars ($X^2 = 0.66197$, $df = 2$, $p\text{-value} = 0.7182$) and the face where micro-
342 scars were located ($X^2 = 2.1323$, $df = 1$, $p\text{-value} = 0.1442$).

343

344 Among other chi-squared tests performed on this dataset we can notice that the type of
345 micro-scars is not independent from the sex of the user ($X^2 = 89.084$, $df = 4$, $p\text{-value} < 0.0001$) and
346 that the type of micro-scars, the orientation and the face where appeared micro-scars are not
347 independent from the user (respectively $X^2 = 587.75$, $df = 76$, $p\text{-value} < 0.0001$; $X^2 = 240.43$, $df =$
348 38 , $p\text{-value} < 0.0001$ and $X^2 = 114.05$, $df = 19$, $p\text{-value} < 0.0001$).

349

350

Figure 4 (2 columns) and Supplementary figures 1 to 7

351

352 3.2. Edge modification visualization and quantification through geometric morphometric

353 3.2.1. Comparison of edges before and after use

354 Before computing the geometric morphometric analysis, the extraction of each edge of
355 interest before and after use allowed comparing them to visualize the location of edge
356 modifications and the amount of removed material during use (Figure 5 and Supplementary figures
357 8 to 18. Most of the modification seem to be on the middle and the rear of the edge. Few tools
358 (40%, $N = 24/60$) present modifications on the front part of the edge but these modifications are
359 generally not invasive. No difference or clear tendency is noticed between tools used with left
360 versus the right hand regarding the amount of removed material and the location of modifications.

361

362

Figure 5 (1 column) and Supplementary figures 8 to 18

363

364 3.2.2. Used edges shapes quantification

365 The first two axes of the PCA on shape data account respectively for 23.26% and 11.79%
366 of the total variance (Figure 6). Groups of tools held with the right hand almost fully overlap with
367 the group of tools held with the left hand. Procrustes ANOVA confirmed that no pattern of shape of
368 used edge can be specifically attributed to tools used with the right hand versus tools used with the
369 left hand. (Procrustes ANOVA: $df = 1$, $SS = 0.0002$, $MS = 0.0002$, $Rsq = 0.0247$, $F = 1.4708$, $Z =$
370 1.4315 , $p\text{-value} = 0.101$). On the contrary, we noticed that the user and the sex of the user may
371 have a larger effect (than the hand holding the tool) on the shape of the used edges (Procrustes
372 ANOVA on shapes and users: $df = 19$, $SS = 0.0038$, $MS = 0.0002$, $Rsq = 0.4301$; $F = 1.5889$; $Z =$
373 4.5104 ; $p\text{-value} = 0.001$; Procrustes ANOVA on shapes and sex of users: $df = 1$, $SS = 0.0002$, MS
374 $= 0.0002$, $Rsq = 0.0324$; $F = 1.9407$; $Z = 2.3377$; $p\text{-value} = 0.029$).

375

376

Figure 6 (2 columns)

377

378 3.3. k Nearest Neighbours model

379 Among the 21 tested models, only five gave an accuracy of at least 75%, all are based on
380 descriptors concerning the maximum scar extent on the surface (perpendicularly to the edge). The
381 best model includes the maximum scar extent and the position of the scar with the maximum
382 extent on the surface (perpendicularly to the edge) on both faces. With a k value set up at 5, this
383 model provided an accuracy of 75% (95% confidence interval: 62%, 85%) with an acceptable
384 kappa of 0.51, a sensitivity of 85% and a specificity of 67% (Table 1).
385

386 **Table 1 (2 columns)**

387
388 4. Discussion

389 4.1. Can hand preference be identified from micro-scars?

390 We could not document any significant difference in position, type, and orientation of scars
391 between tools used by left-handers and right-handers. The microscopic observation showed that
392 tools have more extended scars on their dorsal face than on the ventral face regardless of which
393 hand the user preferred. Knowing that tools were always held with the same orientation during the
394 sawing action (with the dorsal face on the right) this result is counter intuitive. Indeed, during the
395 sawing action we usually observe a tilt of the tool toward the right for right-handers and towards the
396 left for left-handers usually resulting in more scars on the face placed on the left for right-handers
397 and on the right for left-handers. This result shows that hand preference has no influence on the
398 distribution of scars on either of the tool's faces. Combined with the fact that these tools had a
399 centred cutting edge, this result may show that the edge morphology and surface structure may
400 have an effect on the distribution of scars (when the glass casts were made, imperfections on the
401 edges were removed by polishing the ventral side of the edge only; this might have affected the
402 resilience of this face).

403 On the other hand, the type of micro-scars seems to be related to the hand holding the tool
404 during use. This is also a counter-intuitive result. However, as the tool had to be held in the same
405 orientation by left-handers and right-handers, the forces applied by left-handers, theoretically
406 symmetrical to those applied by right-handers, are applied on the opposite surface of the tool. As
407 mentioned above, the surface might have different resilience to contact during use and, as a
408 consequence, different types of micro-scars might have been produced. This should be tested in
409 future experiments. The suggestion that one of the surfaces (i.e. the dorsal face) of the tools might
410 be weaker than the other is somewhat supported by the fact that irrespective of which hand was
411 holding the tool, the dorsal face exhibited more scars.

412
413 Comparison of the edges before and after use did not show significant difference between
414 the amount of removed material and the location of modifications on tools used by left-handers and
415 right-handers. Right-handed subjects produced significantly more invasive and more numerous

416 micro-scars than left-handers but this is very likely to be due to an individual factor as, apart from
417 hand preference, the sex and the user seem to have a negligible effect on the characteristics of the
418 micro-scars. This was confirmed by geometric morphometric analyses which show that the shape
419 of the used edges depends more on the user and user's sex than on the hand preference.

420

421 Based on the data gathered through microscopic observation of scars and quantification of
422 edge modification with geometric morphometric, the k Nearest Neighbours method highlighted the
423 importance of the maximum scar extent (on the ventral face) and of the position of the more
424 invasive scar in discriminating hand preference during stone tool use. The model integrating these
425 variables gave 75% of good identification of the hand preference of the user.

426

427 4.2. Importance of individual characteristics

428 We decided not to choose experts at this step of the project because expertize in using
429 stone flakes (here more exactly soda lime-glass) is a complicated notion. How can we define an
430 expert user of stone flakes? Beginners are sometimes performing very well in such tasks. Like for
431 nowadays tools, a beginner might be more likely to damage the tool quicker than an expert might
432 but can we really consider someone as an expert stone flake user in the current world? Few have
433 more experience than others do and finding enough experienced people to produce statistically
434 acceptable data is not straightforward. In addition, experts may not have been the users of all the
435 archaeological pieces; expertise is not an information that we have archaeologically and we
436 intended to find parameters, which do not depend on the expertise to identify hand preference.

437 Thus, all of the individuals involved in this experiment were beginners in prehistoric tool
438 use. This lack of skill may be one of the reason why several scar features seem to be more related
439 to individual characteristics than to hand preference. The variability of grip and movements, even if
440 somehow constrained, likely influence scars production which may follow a pattern linked to the
441 way each individual used the tool. It can be supposed that in prehistory, frequent use of stones as
442 tools implies a certain degree of know-how and the transmission of this knowledge among
443 members of a particular community. A certain regularity in the way stone tools were used might be
444 expected. If so, individual characteristics might not exert such influence on scars features in
445 archaeological assemblages. However, in any discipline, experts often develop idiosyncratic
446 techniques which can be used to identify them. If stone tool use follows such a model, individual
447 characteristics may be very important in the process of scars formation and, for a specific
448 previously identified task, (even if unlikely because of the numerous variables possibly affecting
449 tool surfaces) it might be possible to evaluate the minimum number of different users within an
450 assemblage. This has to be tested in the future, involving experimenters skilled in stone tool use.

451

452 4.3. Hand holding the tool versus hand preference

453 The hand preference of an individual may vary depending on the action carried out
454 (Provins, 1997). For instance, in the frame of this experiment, one individual declared himself as
455 left-handed but spontaneously used the right hand to saw the wood during the whole experiment.
456 Also, some of the individuals asked to change the hand holding the tool (which was not allowed
457 here) because of pain due to this unusual task. In an unconstrained context (as in Prehistory), it is
458 likely that individuals would change hand if they feel pain or discomfort. Such modification of the
459 hand used during the tasks would highly complicate the identification of left-handed and right-
460 handed people based on archaeological stone tools. However, we can think that, during Prehistory,
461 stone tool use was a more frequent activity and pain and discomfort may have been less common.
462 In any case, such behaviours (as well as the case of ambidextrous people) demonstrate that what
463 we might intent to determine in an archaeological context is the hand holding the tool rather than
464 the hand preference of the user. Only the study of large samples through multi-proxy examination
465 might allow approaching hand preference.

466

467 4.4. Archaeological application

468 The results of this experiment, the k-NN results in particular, are promising and show that it
469 might be possible to identify the hand holding the tool during use based on micro-scars extent and
470 location. However, even if 75% of correct identification is acceptable, it remains valid for a
471 constrained experimental context in wood sawing activity only. We can expect a drastic drop of
472 accuracy in unconstrained situations. Indeed, it is for instance essential to consider that one tool
473 could have been used several times, for various activities, and by different users. Furthermore, in
474 an archaeological assemblage, it is very unlikely to deal only with artefacts used in sawing wood
475 activities. It is now essential to extend the experimental reference corpus with unstandardized
476 pieces (used for different activities), to improve the model with other traces (residues, micro-
477 polishes, striations), and to validate the results with blind-tests before any application on
478 archaeological assemblages. Such an application could be considered when a strong enough (i.e.
479 more than 90% accuracy) experimental model will be ready. It requires a large experimental
480 collection on which each of the traces will be examined and documented. Our model involves
481 parameters that are not dependent on the population ratio of right and lefthanders and is only
482 based on repeatable measurements and description. Once improved, we will be able to propose a
483 new proxy which could provide different result concerning hand preference ratio in Prehistory or
484 reinforce the obtained results from the other disciplines.

485

486 5. Conclusion

487 Wear analysis, geometric morphometric and machine learning provided complementary
488 results in order to examine if the hand holding the tool during use could be identified based on
489 micro-scars. However, the k-NN method showed that the data provided by geometric
490 morphometric and numerous data coming from wear analysis do not help in improving the

accuracy of the model of classification. Nevertheless, counter-intuitive results may indicate a bias related to the structure of the tool used in this experiment and these non-significant variables from geometric morphometric and wear analysis should not be rejected definitively before further experiments. Moreover, the heterogeneity of raw materials used in prehistoric times may lead to similar results. The best model (75% accuracy) is obtained when based on the maximum scar extent and the position of the scar with the maximum extent on the surface of the tool perpendicularly to the edge on both faces. Other variables show overlap for tools used with the left hand and tools used with the right hand, and are likely to depend rather on individual characteristics.

This first experiment, with standardized tools dedicated to hand preference identification, yielded promising results and already highlighted variables that could be used as discriminants. Nevertheless, the experimental corpus needs to be extended and the model tested further to provide reliable results from archaeological assemblages.

504

6. Acknowledgments

We thank Simon Puaud and Florent Détroit for their help and advices during the preparation of the experiments and Myriam Méziou, Anne-Cécile Haussonne and Anthony Herrel for logistic and funding administration. We are grateful to all the volunteers who participated.

Data acquisition and analysis were partly done thanks to the 2D/3D imaging and the Material characterisation platforms of the MNHN. The surface scanning laboratory of the IPHES also provided assistance for data acquisition.

Part of A. Rodriguez research was possible thanks to a grant to the Museum National d'Histoire Naturelle of Paris financed by the project "Robes de pierre", part of the POLYRE ('L'Aventure Polychrome') Chair of Excellence, supported by Sorbonne Universités. The PreMHo project is supported by funding from the French National Research Agency through LabEx ANR-10-LABX-0003-BCDiv, in the program "Investissements d'avenir" ANR-11-IDEX-0004-02.

M.G. Chacón. research is funded by CERCA Programme/Generalitat de Catalunya, by Spanish Ministry of Science, Innovation and Universities (MICINN) of the Spanish Government, project nº HAR2016-76760-C3-1-P and the Catalanian Government, research group's nº 2017 SGR 836.

A. Bardo research was funded by the Fyssen foundation.

We are grateful to the two anonymous reviewers and editors who provided constructive comments, which contributed to improve this manuscript.

524

References

- Adams, D.C., Collyer, M.L., Kaliontzopoulou, A., 2019. Geomorph: Software for geometric morphometric analyses. R package version 3.1.0.
- Annett, M., 1985. Left, right, hand and brain: the right shift theory. Erlbaum, London.

529 Balzeau, A., Gilissen, E., Grimaud-Hervé, D., 2012. Shared Pattern of Endocranial Shape Asymmetries among
530 Great Apes, Anatomically Modern Humans, and Fossil Hominins. *PLoS ONE* 7, e29581.
531 <https://doi.org/10/fzdr7w>

532 Bargalló, A., Mosquera, M., 2014. Can hand laterality be identified through lithic technology? *Laterality: Asymmetries of Body, Brain and Cognition* 19, 37–63. <https://doi.org/10/gfvtk9>

533 Bargalló, A., Mosquera, M., Lorenzo, C., 2018. Identifying handedness at knapping; an analysis of the
534 scatter pattern of lithic remains. *Archaeological and Anthropological Sciences* 10, 587–598.
535 <https://doi.org/10/gfvwd9>

536 Bargalló, A., Mosquera, M., Lozano, S., 2017. In pursuit of our ancestors' hand laterality. *Journal of Human Evolution* 111, 18–32. <https://doi.org/10.1016/j.jhevol.2017.06.001>

537 Bermúdez de Castro, J.M., Bromage, T.G., Jalvo, Y.F., 1988. Buccal striations on fossil human anterior teeth:
538 evidence of handedness in the middle and early Upper Pleistocene. *Journal of Human Evolution* 17,
539 403–412. <https://doi.org/10/c6mx2j>

540 Bermúdez de Castro, J.M., Martinon-Torres, M., Sarmiento, S., Lozano, M., Arsuaga, J.L., Carbonell, E., 2003.
541 Rates of anterior tooth wear in Middle Pleistocene hominins from Sima de los Huesos (Sierra de
542 Atapuerca, Spain). *Proceedings of the National Academy of Sciences* 100, 11992–11996.
543 <https://doi.org/10.1073/pnas.2034879100>

544 Bonhomme, V., Picq, S., Gaucherel, C., Claude, J., 2014. Momocs: Outline Analysis Using R. *Journal of Statistical Software* 56, 1–24. <https://doi.org/10.18637/jss.v056.i13>

545 Bradley, B., Sampson, C.G., 1986. Analysis of replication of two Acheulian artefact assemblages (Chiltern
546 Hills), in: *Stone Age Prehistory: Studies in Memory of Charles McBurney*. G.N. Bailey and P. Callow,
547 Cambridge, UK, pp. 29–45.

548 Brinton, D.G., 1896. Left-Handedness in North American Aboriginal Art. *American Anthropologist* 9, 175–
549 181.

550 Cahen, D., Keeley, L.H., 1980. Not less than two, not more than three. *World Archaeology* 12, 166–180.
551 <https://doi.org/10/fwptdp>

552 Cahen, D., Keeley, L.H., Van Noten, F.L., Behm, J.A., Busby, C.I., Dunnell, R.C., Hayden, B., Johnson, L.L., Katz,
553 P., Mohapatra, G.C., Movius, H.L., Narr, K.J., Newcomer, M., Newell, R.R., Odell, G.H., Ohel, M.Y.,
554 Paddayya, K., Pittioni, R., Roe, D.A., Tomenchuk, J., Wynn, T.G., 1979. Stone Tools, Toolkits, and
555 Human Behavior in Prehistory [and Comments and Reply]. *Current Anthropology* 20, 661–683.
556 <https://doi.org/10.1086/202371>

557 Cashmore, L., 2009. Can hominin 'handedness' be accurately assessed? *Annals of Human Biology* 36, 624–
558 641. <https://doi.org/10/d4rmtq>

559 Cashmore, L., Uomini, N., Chapelain, A., 2008. The evolution of handedness in humans and great apes: a
560 review and current issues. *Journal of Anthropological Sciences* 86, 7–35.

561 Chapelain, A.S., Hogervorst, E., Mbonzo, P., Hopkins, W.D., 2011. Hand Preferences for Bimanual
562 Coordination in 77 Bonobos (*Pan paniscus*): Replication and Extension. *International Journal of Primatology* 32, 491–510. <https://doi.org/10/bt29md>

563 Condemi, S., Monge, J., Quertelet, S., Frayer, D.W., Combier, J., 2017. Vergisson 4: a left-handed
564 Neandertal: CONDEMI et al. *Am. J. Phys. Anthropol.* 162, 186–190.
565 <https://doi.org/10.1002/ajpa.23101>

566 Corballis, M.C., 2003. From mouth to hand: Gesture, speech, and the evolution of right-handedness. *Behavioral and Brain Sciences* 26, 199–260. <https://doi.org/10.1017/S0140525X03000062>

567 Corballis, M.C., 1987. Straw monkeys. *Behavioral and Brain Sciences* 10, 269–270.
568 <https://doi.org/10/dkzpkc>

569 Cornford, J.M., 1986. Specialized resharpening techniques and evidence of handedness, in: *La Cotte de St. Brelade 1961-1978: Excavations by C.B.M. McBurney*. P. Callow and J. M. Cornford, Norwich, UK, pp. 337–351 & 413–414.

570 D'Errico, F., 1992. Technology, Motion, and the Meaning of Epipaleolithic Art. *Current Anthropology* 33, 94–
571 109. <https://doi.org/10/cp4mmp>

572 Domínguez-Rodrigo, M., Baquedano, E., 2018. Distinguishing butchery cut marks from crocodile bite marks
573 through machine learning methods. *Scientific Reports* 8, 5786. <https://doi.org/10.1038/s41598-018-24071-1>

582 Estalrich, A., Rosas, A., 2013. Handedness in Neandertals from the El Sidrón (Asturias, Spain): Evidence
583 from Instrumental Striations with Ontogenetic Inferences. *PLoS ONE* 8, e62797.
584 <https://doi.org/10.1371/journal.pone.0062797>

585 Fagot, J., Vauclair, J., 1991. Manual Laterality in Nonhuman Primates: A Distinction Between Handedness
586 and Manual Specialization. *Psychological Bulletin* 109, 76–89. <https://doi.org/10.1037/0033-2909.109.1.76>

587 Faurie, C., Raymond, M., Uomini, N., 2016. Origins, Development, and Persistence of Laterality in Humans,
588 in: *Laterality in Sports*. Elsevier, pp. 11–30. <https://doi.org/10.1016/B978-0-12-801426-4.00002-X>

589 Faurie, C., Schiefenhövel, W., le Bomin, S., Billiard, S., Raymond, M., 2005. Variation in the Frequency of
590 Left-handedness in Traditional Societies. *Current Anthropology* 46, 142–147.
591 <https://doi.org/10.1086/463bt>

592 Fiore, I., Bondioli, L., Radovč, J., Frayer, D.W., 2015. Handedness in the Krapina Neandertals: A Re-
593 Evaluation. *PaleoAnthropology* 19–36. <https://doi.org/10.4207/PA.2015.ART93>

594 Forrester, G.S., Leavens, D.A., Quaresmini, C., Vallortigara, G., 2011. Target animacy influences gorilla
595 handedness. *Animal Cognition* 14, 903–907. <https://doi.org/10.1007/s10071-011-0536-4>

596 Forrester, G.S., Quaresmini, C., Leavens, D.A., Mareschal, D., Thomas, M.S.C., 2013. Human handedness: An
597 inherited evolutionary trait. *Behavioural Brain Research* 237, 200–206. <https://doi.org/10.1016/j.bbr.2012.10.015>

598 Forrester, G.S., Quaresmini, C., Leavens, D.A., Spiezio, C., Vallortigara, G., 2012. Target animacy influences
599 chimpanzee handedness. *Animal Cognition* 15, 1121–1127. <https://doi.org/10.1007/s10071-012-0536-4>

600

601 Foster, A., Buckley, H., Tayles, N., 2014. Using Enthesis Robusticity to Infer Activity in the Past: A Review.
602 *Journal of Archaeological Method and Theory* 21, 511–533. <https://doi.org/10.1007/s10816-012-9156-1>

603

604 Frayer, D.W., Fiore, I., Lalueza-Fox, C., Radovčić, J., Bondioli, L., 2010. Right handed Neandertals: Vindija and
605 beyond. *Journal of Anthropological Sciences* 88, 113–127.

606 Frayer, D.W., Lozano, M., Bermúdez de Castro, J.M., Carbonell, E., Arsuaga, J.L., Radovčić, J., Fiore, I.,
607 Bondioli, L., 2012. More than 500,000 years of right-handedness in Europe. *Laterality: Asymmetries*
608 *of Body, Brain and Cognition* 17, 51–69. <https://doi.org/10.1080/13602542.2012.678888>

609 Frey, S.H., Gerry, V.E., 2006. Modulation of Neural Activity during Observational Learning of Actions and
610 Their Sequential Orders. *J. Neurosci.* 26, 13194–13201. <https://doi.org/10.1523/JNEUROSCI.3914-06.2006>

611

612 Greenfield, P.M., 1991. Language, tools and brain: The ontogeny and phylogeny of hierarchically organized
613 sequential behavior. *Behavioral and Brain Sciences* 14, 531–551.
614 <https://doi.org/10.1017/S0140525X00071235>

615 Gunz, P., Mitteroecker, P., 2013. Semilandmarks: a method for quantifying curves and surfaces. *Hystrix, the*
616 *Italian Journal of Mammalogy* 24. <https://doi.org/10.4404/hystrix-24.1-6292>

617 Hartmann, K., Goldenberg, G., Daumüller, M., Hermsdörfer, J., 2005. It takes the whole brain to make a cup
618 of coffee: the neuropsychology of naturalistic actions involving technical devices. *Neuropsychologia*
619 43, 625–637. <https://doi.org/10.1016/j.neuropsychologia.2004.07.015>

620 Ho Ho Committee, 1979. The Ho Ho Classification and Nomenclature Committee Report, in: *Lithic Use-*
621 *Wear Analysis*. Academic Press, pp. 133–135.

622 Holloway, R.L., 1981. Volumetric and asymmetry determinations on recent hominid endocasts: Spy I and II,
623 Djebel Ihroud I, and the salèHomo erectus specimens, with some notes on neandertal brain size.
624 *American Journal of Physical Anthropology* 55, 385–393. <https://doi.org/10.1002/ajpa.1330550308>

625 Holloway, R.L., De La Costelareymondie, M.C., 1982. Brain endocast asymmetry in pongids and hominids:
626 Some preliminary findings on the paleontology of cerebral dominance. *American Journal of Physical*
627 *Anthropology* 58, 101–110. <https://doi.org/10.1002/ajpa.1330580205>

628 Hopkins, W.D., 2006. Comparative and familial analysis of handedness in great apes. *Psychological Bulletin*
629 132, 538–559. <https://doi.org/10.1037/0033-2909.132.4.538>

630 Hopkins, W.D., Phillips, K.A., Bania, A., Calcutt, S.E., Gardner, M., Russell, J., Schaeffer, J., Lonsdorf, E.V.,
631 Ross, S.R., Schapiro, S.J., 2011. Hand Preferences for Coordinated Bimanual Actions in 777 Great
632 Apes: Implications for the Evolution of Handedness in Hominins. *J Hum Evol* 60, 605–611.
633 <https://doi.org/10.1016/j.jhevol.2011.05.004>

Iovita, R., Schönekeß, H., Gaudzinski-Windheuser, S., Jäger, F., 2014. Projectile impact fractures and launching mechanisms: results of a controlled ballistic experiment using replica Levallois points. *Journal of Archaeological Science* 48, 73–83. <https://doi.org/10.1016/j.jas.2013.01.031>

Johnson-Frey, S.H., Newman-Norlund, R., Grafton, S.T., 2005. A Distributed Left Hemisphere Network Active During Planning of Everyday Tool Use Skills. *Cereb Cortex* 15, 681–695. <https://doi.org/10.1093/cercor/bhh169>

Karakostis, F.A., Hotz, G., Scherf, H., Wahl, J., Harvati, K., 2017. Occupational manual activity is reflected on the patterns among hand entheses. *American Journal of Physical Anthropology* 164, 30–40. <https://doi.org/10.1002/ajpa.23253>

Karakostis, F.A., Hotz, G., Tourloukis, V., Harvati, K., 2018. Evidence for precision grasping in Neandertal daily activities. *Science Advances* 4, eaat2369. <https://doi.org/10.1126/science.adc2369>

Karakostis, F.A., Lorenzo, C., 2016. Morphometric patterns among the 3D surface areas of human hand entheses. *American Journal of Physical Anthropology* 160, 694–707. <https://doi.org/10.1002/ajpa.22999>

Keeley, L.H., 1977. The Functions of Paleolithic Flint Tools. *Scientific American* 237, 108–126. <https://doi.org/10.1038/sciam.1977.108>

Klingenberg, C.P., 2013. Visualizations in geometric morphometrics: how to read and how to make graphs showing shape changes. *Hystrix It. J. Mamm.* 24, 15–24. <https://doi.org/10.4404/hystrix-24.1-7691>

Kuhn, A., Wing, J., Weston, S., Williams, A., Keefer, C., Engelhardt, A., Cooper, T., Mayer, Z., Kenkel, B., R Core Team, Benesty, M., Lescarbeau, R., Ziem, A., Scrucca, L., Tang, Y., Candan, C., Hunt, T., 2019. caret: Classification and Regression Training. R package version 6.0-84.

Lazenby, R.A., Cooper, D.M.L., Angus, S., Hallgrímsson, B., 2008. Articular constraint, handedness, and directional asymmetry in the human second metacarpal. *Journal of Human Evolution* 54, 875–885. <https://doi.org/10.1016/j.jhevol.2007.12.001>

Lewis, J.W., 2006. Cortical Networks Related to Human Use of Tools. *Neuroscientist* 12, 211–231. <https://doi.org/10.1177/1073858406288327>

Llaurens, V., Raymond, M., Faurie, C., 2009. Why are some people left-handed? An evolutionary perspective. *Philosophical Transactions of the Royal Society B: Biological Sciences* 364, 881–894. <https://doi.org/10.1098/rstb.2008.0235>

Lozano, M., Estalrich, A., Bondioli, L., Fiore, I., Castro, J.-M.B. de, Arsuaga, J.L., Carbonell, E., Rosas, A., Frayer, D.W., 2017. Right-handed fossil humans. *Evolutionary Anthropology: Issues, News, and Reviews* 26, 313–324. <https://doi.org/10.1002/evan.21554>

Lozano, M., Mosquera, M., de Castro, J.M.B., Arsuaga, J.L., Carbonell, E., 2009. Right handedness of Homo heidelbergensis from Sima de los Huesos (Atapuerca, Spain) 500,000 years ago. *Evolution and Human Behavior* 30, 369–376. <https://doi.org/10.1016/j.ehbeh.2009.03.003>

Marzke, M.W., 2013. Tool making, hand morphology and fossil hominins. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368, 20120414–20120414. <https://doi.org/10.1098/rstb.2012.0414>

McGrew, W.C., Marchant, L.F., 1997. On the other hand: Current issues in and meta-analysis of the behavioral laterality of hand function in nonhuman primates. *Yearbook of Physical Anthropology* 40, 201–232.

McManus, I.C., 2009. The history and geography of human handedness, in: Sommer, I.E.C., Kahn, R.S. (Eds.), *Language Lateralization and Psychosis*. Cambridge University Press, Cambridge, pp. 37–58. <https://doi.org/10.1017/CBO9780511576744.004>

Meguerditchian, A., Vauclair, J., Hopkins, W.D., 2013. On the origins of human handedness and language: A comparative review of hand preferences for bimanual coordinated actions and gestural communication in nonhuman primates. *Developmental Psychobiology* 55, 637–650. <https://doi.org/10.1002/dev.22400>

Mosquera, M., Geribàs, N., Bargalló, A., Llorente, M., Riba, D., 2012. Complex Tasks Force Hand Laterality and Technological Behaviour in Naturalistically Housed Chimpanzees: Inferences in Hominin Evolution. *The Scientific World Journal* 2012, 1–12. <https://doi.org/10.1155/2012/123456>

Newcomer, M.H., Sieveking, G. de G., 1980. Experimental Flake Scatter-Patterns: A New Interpretative Technique. *Journal of Field Archaeology* 7, 345–352.

687 Patterson, J.B., Sollberger, L.W., 1986. Comments on Toth's Right-Handedness Study. *Lithic Technology* 15,
688 109–111. <https://doi.org/10/gfvtpx>

689 Perez, S.I., Bernal, V., Gonzalez, P.N., 2006. Differences between sliding semi-landmark methods in
690 geometric morphometrics, with an application to human craniofacial and dental variation. *Journal*
691 *of Anatomy* 208, 769–784. <https://doi.org/10.1111/j.1469-7580.2006.00576.x>

692 Pobiner, B.L., 1999. The Use of Stone Tools to Determine Handedness in Hominids. *Current Anthropology*
693 40, 90–92. <https://doi.org/10/c9w3md>

694 Porac, C., Coren, S., 1981. *Lateral Preferences and Human Behavior*. Springer-Verlag, New York.

695 Pouydebat, E., Borel, A., Chotard, H., Fragaszy, D., 2014. Hand preference in fast-moving versus slow-
696 moving actions in capuchin, *Sapajus* spp., and squirrel monkeys, *Saimiri sciureus*. *Animal Behaviour*
697 97, 113–123. <https://doi.org/10/f6prdn>

698 Poza-Rey, E.M., Lozano, M., Arsuaga, J.L., 2017. Brain asymmetries and handedness in the specimens from
699 the Sima de los Huesos site (Atapuerca, Spain). *Quaternary International* 433, 32–44.
700 <https://doi.org/10.1016/j.quaint.2015.10.004>

701 Provins, K.A., 1997. The Specificity of Motor Skill and Manual Asymmetry: A Review of the Evidence and Its
702 Implications. *Journal of Motor Behavior* 29, 183–192. <https://doi.org/10.1080/00222899709600832>

703 Quaresmini, C., Forrester, G.S., Spiezio, C., Vallortigara, G., 2014. Social environment elicits lateralized
704 behaviors in gorillas (*Gorilla gorilla gorilla*) and chimpanzees (*Pan troglodytes*). *Journal of*
705 *Comparative Psychology* 128, 276–284. <https://doi.org/10/gfvwfs>

706 R Core Team, 2019. R: A language and environment for statistical computing. R Foundation for Statistical
707 Computing. Vienna, Austria.

708 Rogers, L.J., 2009. Hand and paw preferences in relation to the lateralized brain. *Philosophical Transactions*
709 *of the Royal Society B: Biological Sciences* 364, 943–954. <https://doi.org/10/d2hdtb>

710 Ruck, L., Broadfield, D.C., Brown, C.T., 2015. Determining hominid handedness in lithic debitage: a review of
711 current methodologies. *Lithic Technology* 40, 171–188. <https://doi.org/10/gfvwft>

712 Rugg, G., Mullane, M., 2001. Inferring handedness from lithic evidence. *Laterality: Asymmetries of Body,*
713 *Brain and Cognition* 6, 247–259. <https://doi.org/10.1080/713754411>

714 Schluter, N.D., Krams, M., Rushworth, M.F.S., Passingham, R.E., 2001. Cerebral dominance for action in the
715 human brain: the selection of actions. *Neuropsychologia* 39, 105–113.
716 [https://doi.org/10.1016/S0028-3932\(00\)00105-6](https://doi.org/10.1016/S0028-3932(00)00105-6)

717 Semenov, S.A., 1970. *Prehistoric technology: an experimental study of the oldest tools and artefacts from*
718 *traces of manufacture and wear*. Cory, Adams & Mackay, London.

719 Shaw, C.N., 2011. Is 'hand preference' coded in the hominin skeleton? An in-vivo study of bilateral
720 morphological variation. *Journal of Human Evolution* 61, 480–487. <https://doi.org/10/cnp7dg>

721 Sparacello, V.S., Villotte, S., Shackelford, L.L., Trinkaus, E., 2017. Patterns of humeral asymmetry among
722 Late Pleistocene humans. *Comptes Rendus Palevol, Hominin biomechanics, virtual anatomy and*
723 *inner structural morphology: From head to toe. A tribute to Laurent Puymerail* 16, 680–689.
724 <https://doi.org/10/gbxm8s>

725 Spenneman, D.R., 1984. Handedness data on the European neolithic. *Neuropsychologia* 22, 613–615.
726 <https://doi.org/10/bvddpn>

727 Steele, J., Uomini, N., 2009. Can the Archaeology of Manual Specialization Tell Us Anything About Language
728 Evolution? A Survey of the State of Play. *Cambridge Archaeological Journal* 19, 97.
729 <https://doi.org/10/dbjrc5>

730 Stephens, N.B., Kivell, T.L., Gross, T., Pahr, D.H., Lazenby, R.A., Hublin, J.-J., Hershkovitz, I., Skinner, M.M.,
731 2016. Trabecular architecture in the thumb of *Pan* and *Homo*: implications for investigating hand
732 use, loading, and hand preference in the fossil record. *American Journal of Physical Anthropology*
733 161, 603–619. <https://doi.org/10.1002/ajpa.23061>

734 Stock, J.T., Shirley, M.K., Sarringhaus, L.A., Davies, T.G., Shaw, C.N., 2013. Skeletal evidence for variable
735 patterns of handedness in chimpanzees, human hunter-gatherers, and recent British populations.
736 *Annals of the New York Academy of Sciences* 1288, 86–99. <https://doi.org/10.1111/nyas.12067>

737 Stout, D., Chaminade, T., 2012. Stone tools, language and the brain in human evolution. *Philos Trans R Soc*
738 *Lond B Biol Sci* 367, 75–87. <https://doi.org/10.1098/rstb.2011.0099>

739 Stout, D., Toth, N., Schick, K., Chaminade, T., 2008. Neural correlates of Early Stone Age toolmaking:
 740 technology, language and cognition in human evolution. *Philosophical Transactions of the Royal*
 741 *Society B: Biological Sciences* 363, 1939–1949. <https://doi.org/10/fk2djc>
 742 Stout, D., Toth, N., Schick, K., Stout, J., Hutchins, G., 2000. Stone Tool-Making and Brain Activation: Position
 743 Emission Tomography (PET) Studies. *Journal of Archaeological Science* 27, 1215–1223.
 744 <https://doi.org/10/ds4vjg>
 745 Toth, N., 1985. Archaeological evidence for preferential right-handedness in the lower and middle
 746 pleistocene, and its possible implications. *Journal of Human Evolution* 14, 607–614.
 747 <https://doi.org/10/fj9wb7>
 748 Trinkaus, E., Churchill, S.E., Ruff, C.B., 1994. Postcranial robusticity in Homo. II: Humeral bilateral
 749 asymmetry and bone plasticity. *American Journal of Physical Anthropology* 93, 1–34.
 750 <https://doi.org/10/b8q2ws>
 751 Uomini, N., Gowlett, J., 2013. La latéralité manuelle préhistorique, les outils et le langage. *Revue de*
 752 *primatologie* 1–19. <https://doi.org/10/gfvwfv>
 753 Uomini, N.T., 2011. Handedness in Neanderthals, in: Conard, N.J., Richter, J. (Eds.), *Neanderthal Lifeways,*
 754 *Subsistence and Technology*. Springer Netherlands, Dordrecht, pp. 139–154.
 755 https://doi.org/10.1007/978-94-007-0415-2_14
 756 Uomini, N.T., 2009. The prehistory of handedness: Archaeological data and comparative ethology. *Journal*
 757 *of Human Evolution* 57, 411–419. <https://doi.org/10.1016/j.jhevol.2009.02.012>
 758 Uomini, N.T., 2008. In the knapper's hands: identifying handedness from lithic production and use, in:
 759 *Prehistoric Technology. 40 Years Later: Functional Studies and the Russian Legacy*, BAR
 760 *International Series*. pp. 51–62.
 761 Uomini, N.T., 2001. Lithic indicators of handedness: assessment of methodologies and the evolution of
 762 laterality in hominids (M.Sc. Dissertation). University of Durham, Durham.
 763 Uomini, N.T., Meyer, G.F., 2013. Shared Brain Lateralization Patterns in Language and Acheulean Stone Tool
 764 Production: A Functional Transcranial Doppler Ultrasound Study. *PLoS ONE* 8, e72693.
 765 <https://doi.org/10.1371/journal.pone.0072693>
 766 Vallortigara, G., Rogers, L.J., 2005. survival with an asymmetrical brain: advantages and disadvantages of
 767 cerebral lateralization. *Behavioral and Brain Sciences* 28, 575–589.
 768 <https://doi.org/10.1017/S0140525X05000105>
 769 Weber, T., 1990. Analysen der archäologischen Funde und des Befunds, in: *Neumark-Gröbern: Beiträge Zur*
 770 *Jagd Des Mittelpaläolithischen Menschen*, Veröffentlichungen Des Landesmuseums Für
 771 *Vorgeschichte in Halle*. D. Mania, M. Thomae, T. Litt and T. Weber, Berlin, pp. 237–255.
 772 Willems, R.M., Hagoort, P., 2007. Neural evidence for the interplay between language, gesture, and action:
 773 A review. *Brain and Language, Gesture, Brain, and Language* 101, 278–289.
 774 <https://doi.org/10.1016/j.bandl.2007.03.004>
 775 Xing, S., O'Hara, M., Guatelli-Steinberg, D., Ge, J., Liu, W., 2017. Dental Scratches and Handedness in East
 776 Asian Early Pleistocene Hominins: Handedness of East Asian Hominin. *Int. J. Osteoarchaeol.* 27,
 777 937–946. <https://doi.org/10.1002/oa.2601>
 778

Fig.1: Soda-lime glass replica used in the experiment.

Fig.2: Documentation of the type, orientation and position of maximum extent of the micro-scars.

Fig.3: Used edges curves extraction for geometric morphometric analysis. A) Example of photograph taken after use. B) Binarization. C) Scaling of outlines. D) The line going through the extremities of each outline is placed horizontally, the outlines are then wedged on the rightmost point. E) The average coordinates of the uppermost points of the outlines is calculated and the uppermost point of each outline is aligned on it. F) The edge of interest is extracted by cutting the outline at the uppermost point and at the leftmost point.

Fig.4: Examples of position and orientation of micro-scars for left and right-handers.

Fig.5: Comparison of tool edges before and after use for left and right-handers.

Fig.6: Two first axes of the principal component analysis of the coordinates obtained by Generalized Procrustean Analysis (GPA) of the curves representing the used edges of the tools used by left-handers and right-handers. The used edge shape reconstructions are displayed for the minimum and the maximum values of each PC.

Table 1: Results of the k Nearest Neighbours models. Models providing the best accuracy are in bold.

1cm

A)

Break

Feathered

Step

Abrupt

After Vaughan (1985)

B)

A)

B)

C)

D)

E)

F)

Tool used by a left-hander

Ventral face

Dorsal face

Tool used by a right-hander

Ventral face

Dorsal face

: General orientation of the micro-scars

: Micro-scars extent

: Edge modification visible at a magnification higher than 10X

MAX PC2

MIN PC2

MIN PC1

MAX PC1

Model n°	Included variables	K-value giving best results	Accuracy	95%CI	Kappa	Sensitivity	Specificity
1	Number of scars on the ventral and dorsal faces of the tool	23	0.65	0.5160, 0.7687	0.3092	0.7407	0.5758
2	Types of scars on both faces	9 & 11	0.6833	0.5504, 0.7974	0.3537	0.5926	0.7576
3	Types of scars on dorsal face	5	0.7	0.5679, 0.8115	0.398	0.7037	0.6970
4	Types of scars on ventral face	31	0.6	0.4654, 0.7244	0.1864	0.5185	0.6667
5	Directions of scars on both faces	17 & 27	0.6	0.4654, 0.7244	0.2131	0.7037	0.5152
6	Directions of scars on dorsal face	41	0.6333	0.499, 0.7541	0.2834	0.7778	0.5152
7	Directions of scars on ventral face	21	0.6333	0.499, 0.7541	0.2691	0.6667	0.6061
8	Types and directions of scars on both faces	25	0.65	0.516, 0.7687	0.3	0.6667	0.6364
9	Number, types and directions of scars on both faces	27	0.6167	0.4821, 0.7393	0.2333	0.6296	0.6061
10	The 59 first principal components of the PCA on geometric morphometrics data	55 & 57 & 59	0.55	0.4161, 0.6788	0	0	1
11	The 2 first principal components of the PCA on geometric morphometrics data	31	0.5833	0.4488, 0.7093	0.1776	0.6667	0.5152
12	The 59 first principal components of the PCA on geometric morphometrics data and the number of scars on both faces	13 & 15	0.65	0.516, 0.7687	0.3	0.6667	0.6364
13	Position of the scar with the maximum extent on the surface perpendicularly to the edge on ventral face	17 & 47	0.7333	0.6034, 0.8393	0.4539	0.6296	0.8182
14	Position of the scar with the maximum extent on the surface perpendicularly to the edge on dorsal face	21 & 23	0.5833	0.4488, 0.7093	0.1611	0.5556	0.6061
15	Position of the scar with the maximum extent on the surface perpendicularly to the edge on both faces	11	0.7333	0.6034, 0.8393	0.4613	0.7037	0.7576
16	Maximum scar extent on the surface perpendicularly to the edge on ventral face	25 & 37	0.75	0.6214, 0.8528	0.4966	0.7407	0.7576
17	Maximum scar extent on the surface perpendicularly to the edge on dorsal face	21 & 23	0.7333	0.6034, 0.8393	0.4613	0.7037	0.7576
18	Maximum scar extent on the surface perpendicularly to the edge on both faces	23	0.75	0.6214, 0.8528	0.5	0.7778	0.7273
19	Maximum scar extent and position of the scar with the maximum extent on the surface perpendicularly to the edge on both faces	5	0.75	0.6214, 0.8528	0.5066	0.8519	0.6667
20	All descriptors from microscopic observation	5	0.75	0.6214, 0.8528	0.5	0.7778	0.7273
21	All descriptors from microscopic observation and geometric morphometrics	9	0.75	0.6214, 0.8528	0.5066	0.8519	0.6667