



Visible-Light-Induced Metal-Free Trifluoromethylselenolation of Electron-Rich Heteroarenes Using the Nucleophilic Me₄N][SeCF₃] Reagent

Arnaud de Zordo-Banliat, Lucas Barthélémy, Flavien Bourdreux, Béatrice N. Tuccio, Guillaume Dagousset, Bruce Pégot, Emmanuel Magnier

► To cite this version:

Arnaud de Zordo-Banliat, Lucas Barthélémy, Flavien Bourdreux, Béatrice N. Tuccio, Guillaume Dagousset, et al.. Visible-Light-Induced Metal-Free Trifluoromethylselenolation of Electron-Rich Heteroarenes Using the Nucleophilic Me₄N][SeCF₃] Reagent. *European Journal of Organic Chemistry*, 2020, 2020 (4), pp.506-509. 10.1002/ejoc.201901793 . hal-02861697

HAL Id: hal-02861697

<https://hal.science/hal-02861697>

Submitted on 22 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Visible-Light-Induced Metal-Free Trifluoromethylselenenolation of Electron-Rich Heteroarenes using the Nucleophilic [Me₄N][SeCF₃] Reagent

Arnaud De Zordo-Banliat,^[a] Lucas Barthélémy,^[b] Flavien Bourdreux,^[a] Béatrice Tuccio,^[b] Guillaume Dagousset,^{*[a]} Bruce Pégot,^{*[a]} and Emmanuel Magnier^{*[a]}

Abstract: A metal-free visible-light-promoted regioselective trifluoromethylselenenolation of electron-rich heteroarenes has been developed via C–H functionalization. This eco-friendly, atom-economical, and easy-to-operate protocol provides direct access to a wide range of functionalized SeCF₃-containing heteroarenes in high yields, and is amenable to continuous flow techniques. A radical mechanism was supported by EPR experiments.

Organofluorine molecules have attracted continuous interest due to the specific properties of this halogen. These fluorinated compounds have found many applications in various areas, from life sciences to materials science.^[1] More recently, it has been shown that the association of a CF₃ moiety with a heteroatom (such as oxygen, sulfur, or selenium) can deeply modulate these physicochemical and biological properties.^[2] Amongst these new emergent fluorinated moieties, the SeCF₃ group has probably been the least studied, but a clear resurgence of the appeal for this group recently appeared in the literature. Selenoproteins are indeed useful endogenous molecules, and some organoselenium compounds have interesting biological activities at small dose,^[3] or can find valuable applications in materials science.^[4] Another interest of this group is the lipophilicity induced in organic moieties. The recent determination of its Hansch parameter shows a value slightly lower than the parent SCF₃ group and similar to the one of the SF₅ group, another emerging group.^[5] Therefore, the development of new methods to incorporate the SeCF₃ moiety directly into organic molecules is highly desirable.

To date, most of the available strategies to directly introduce a SeCF₃ group are nucleophilic or electrophilic approaches (using either nucleophilic MSeCF₃ reagents, M = 1/2 Hg, Cu, Me₄N, or electrophilic XSeCF₃ reagents, X = SeCF₃, Cl, Tosyl, Scheme 1a,b).^[6,7] Very recently, Billard, Tlili and co-workers reported the first radical trifluoromethylselenenolation reaction under visible-light photoredox catalysis, using the electrophilic reagent **1**.^[8] This radical process was notably applied to the efficient synthesis of trifluoromethylselenenolated arenes from the corresponding diazonium salts (Scheme 1c).^[8a] However,



Scheme 1. The different strategies for the direct introduction of the SeCF₃ moiety.

considering the low commercial availability of diazonium salts, a greener approach would consist in the direct C–H trifluoromethylselenenolation of non-prefunctionalized arenes. Based on our previous works on the photoredox-catalysed synthesis of fluorinated compounds,^[9] we anticipated that the \cdot SeCF₃ radical could also be generated from a nucleophilic source of SeCF₃ via a visible-light-induced SET (Single Electron Transfer) oxidation. This fluorinated radical would then easily react with aromatic substrates (Scheme 1d). We wish to report herein a new visible-light-promoted aerobic C–H trifluoromethylselenenolation of heteroarenes, which proceeds under very mild, metal-free and eco-friendly reaction conditions.

We focused our initial investigations on the trifluoromethylselenenolation of indole (**1a**) with the easily accessible and air-stable Me₄NSeCF₃ reagent **2** (Table 1).^[6d] We were pleased to see that the reaction using 2 mol% of Ru(bpy)₃(PF₆)₂ (**3a**, bpy = 2,2'-bipyridine) as the photocatalyst with blue LEDs irradiation under air atmosphere furnished a high yield of the 3-trifluoromethylselenenolated indole **4a** (87%, entry 1). A survey of organic dyes in place of Ru(bpy)₃(PF₆)₂ revealed that Eosin Y (**3d**) was the optimal organophotocatalyst for this transformation, affording an almost quantitative yield of

[a] A. De Zordo-Banliat, F. Bourdreux, Dr. G. Dagousset, Dr. B. Pégot, Dr. E. Magnier
Institut Lavoisier de Versailles, UMR 8180, Université de Versailles-Saint-Quentin 78035 Versailles Cedex, France,
<http://www.ilv.uvsq.fr/>
Fax: (+) 33 1 39254452
[b] L. Barthélémy, Dr. B. Tuccio
Aix-Marseille Université-CNRS, Institut de Chimie Radicalaire, UMR 7273, F-13397 Marseille Cedex 20, France

Table 1. Survey of reaction conditions for the photocatalyzed trifluoromethylselenenolation of indole **1a**.


Entry	x	Photoredox catalyst 3	Yield [%] ^[a,b]
1 ^[c]	2	Ru(bpy) ₃ (PF ₆) ₂ 3a	87
2	2	Rhodamine B 3b	35
3	2	Rose Bengal 3c	34
4	2	Eosin Y 3d	99 (89) ^[d]
5 ^[e]	2	3d	48
6	1.5	3d	60
7 ^[f]	2	3d	0
8 ^[g]	2	3d	0
9	2	none	7

[a] General conditions: **1a** (0.20 mmol), **2** (x equiv), **3** (0.02 equiv) in MeCN (2 mL) irradiated with green LEDs at RT for 18 h. [b] Yields determined by ¹⁹F NMR spectroscopy using PhOCF₃ as an internal standard. [c] Reaction performed with blue LEDs irradiation. [d] Yields referred to chromatographically pure product into brackets. [e] Reaction performed with 1 mol% of **3d**. [f] Reaction performed in the dark. [g] Reaction performed under argon atmosphere.

4a (entries 2-4). Attempts to use reduced amount of SeCF₃ reagent or of the photocatalyst led to diminished yields (entries 5-6), and air, light and catalyst were all essential partners for the reaction to proceed (entries 7-9).


Having established the optimal reaction conditions, the scope of this novel photocatalysed trifluoromethylselenenolation was next investigated and the results are summarised in Scheme 2. Pleasingly, a wide range of indole derivatives **1a-1s** were smoothly trifluoromethylselenolated. The mild reaction conditions were compatible with a large array of electron-donating or –withdrawing functionalities at position 2, 4, 5 or 6 of the indole scaffold, including ethers, nitro, halogens, nitriles, esters, aldehydes, ketones and boronic esters. This process was fully regioselective in favour of the 3-position, and it is worth to note that no reaction took place in the case of 3-substituted indoles. Remarkably, this reaction was successfully broadened to other electron-rich heteroaromatic substrates: 7-azaindole (**1t**) and 4-azaindole (**1u**) were efficiently converted into the corresponding SeCF₃-adducts **4t-4u** in high yields, while functionalized pyrroles **1v-1z** selectively afforded the corresponding 2-trifluoromethylselenolated pyrroles **4v-4z**, albeit in moderate yields.

In addition, this novel protocol was also optimized with the use of continuous-flow techniques, which are perfectly well adapted to photochemical transformations: indeed, the narrow PFA tubing (1.3 mm, PFA = perfluoroalkoxy) of the flow device allows for an optimal and homogeneous irradiation of the reaction mixture, heat and mass transfer characteristics are enhanced,




Scheme 2. Scope of the visible-light-mediated trifluoromethylselenenolation of heteroarenes. General conditions: **1** (0.2 mmol), **2** (0.4 mmol), **3d** (2 mol%) in 2 mL of MeCN irradiated with green LEDs under air for 18 h at rt. Yields referred to chromatographically pure products, and yields into brackets were determined by ¹⁹F NMR spectroscopy using PhOCF₃ as internal standard. [a] 1 mmol scale. [b] 36 h of irradiation.

and scale-up experiments are facilitated under safer reaction conditions.^[10] In order to reach complete conversion of indole **1a**, slight modifications of the reaction conditions were made. The concentration of the reaction was decreased to 0.033M to achieve complete dissolution, and air was replaced by O₂ (see SI for more details). With these optimal conditions, the residence time could be decreased to only 50 min of irradiation, and the expected SeCF₃ product **4a** was obtained in 81% yield. Furthermore, the same experiment was performed on a larger scale without any significant loss of the yield (1 mmol, Scheme 3).



Scheme 3. Synthesis of 3-(trifluoromethylselanyl)indole (**4a**) using continuous flow.

We then decided to investigate the mechanism of this transformation. The fact that both Eosin Y and continuous irradiation were needed for the reaction to proceed (see Table 1 and the Supporting information for light on/off experiments) suggested that a radical pathway was involved. To verify this hypothesis, some spin trapping / electron paramagnetic resonance (ST/EPR) experiments were carried out with 2-methylnitrosopropane (MNP) as the spin trap. In a first step, irradiating the radical source Me₄NSeCF₃ **2** in presence of Eosin Y **3d** and MNP led to record a very intense three line EPR spectrum ($a_N = 15.7$ G, spectrum I, Scheme 4a). Since blank tests performed in the absence of either Me₄NSeCF₃ **2** or of Eosin Y **3d** never yielded any signal, spectrum I could be assigned to the spin adduct formed after trapping the radical $\cdot\text{SeCF}_3$, *i.e.* MNP-SeCF₃, the satellite lines being due to a hyperfine coupling with ¹³C in β -position (natural abundance, $a_{13C} = 4.2$ G). The generation of $\cdot\text{SeCF}_3$ was further confirmed by ST/EPR experiments performed with the nitron spin trap DMPO (see ESI). In a second step, additional spin trapping experiments were performed in the presence of excess of indole **1a**, added to the medium containing reagent **2**, MNP and photocatalyst **3d**. The EPR spectrum II given in Scheme 4a was thus recorded. Its analysis revealed the presence of two spin adducts, the minor one (10%, $a_N = 15.7$ G) corresponding to MNP-SeCF₃. The major compound (90%) shows a more complex signal due to hyperfine couplings with the nitroxide nitrogen ($a_N = 14.5$ G), but also with one hydrogen ($a_{H\beta} = 2.7$ G) and one nitrogen ($a_{N\beta} = 1.8$ G) nuclei, both in β position towards the nitroxide function. This is perfectly consistent with adduct MNP-B obtained after trapping of the carbon-centered radical **B** coming from the addition of $\cdot\text{SeCF}_3$ onto indole (Scheme 4b). It is worth noting that, to the best of our knowledge, such α -amino radical **B** has



Scheme 4. (a) EPR spectra obtained using MNP as the spin trap (red dotted lines correspond to the calculated spectra). For experimental details, see the SI. (b) Proposed mechanism.

never been characterized in other similar radical C–H functionalizations of indoles.

In addition, redox potentials of indole **1a** and Me₄NSeCF₃ **2** were measured by cyclic voltametry (see the Supporting Information). The respective values of oxidation potential ($E_{1a} = +1.46$ V and $E_2 = +0.72$ V vs. SCE) clearly showed that the oxidation of **2** by the excited state EY* of **3d** ($E_{EY^*/EY\cdot-} = +0.83$ V vs. SCE)^[11] is thermodynamically feasible, while oxidation of **1a** by EY* is a thermodynamically unfavourable process. Furthermore, luminescence quenching experiments were performed. No quench of the excited state EY* of **3d** was observed with indole **1a**, which also invalidated the direct oxidation of **1a** by EY*. We also monitored the UV-Vis absorbance of EY in the presence of **1a** or **2**. While no change was noted with **1a**, a small hypsochromic shift (from 538 nm to 527 nm) was observed when increasing concentrations of Me₄NSeCF₃ **2** were added. This observation might suggest the formation of a ground-state complex between EY and **2**, which was further corroborated by NMR experiments (see the Supporting Information). Moreover, the measurement of the quantum yield of this reaction (0.48, see the Supporting Information for more details) suggested that a radical chain mechanism is not the major operative pathway of this transformation.

On the basis of all these studies, a plausible mechanism is depicted in Scheme 4b. Eosin Y would first associate with reagent **2**, leading to a ground-state complex [EY·2]. Then, irradiation of this complex with green LEDs would give rise to the $\cdot\text{SeCF}_3$ radical **A**, which would add onto indole **1**, leading to radical intermediate **B**. Both radicals **A** and **B** were clearly

identified by ST/EPR experiments. Then radical **B** would be easily oxidized with concurrent loss of a proton, affording the desired 3-trifluoromethylselenolated indole **4**. It is also worth noting that no dimeric species $\text{CF}_3\text{SeSeCF}_3$ was observed during the reaction, even when using *in operando* ^{19}F NMR spectroscopy. However, a polar mechanism involving this dimer, which would account for the observed C-3 regioselectivity, cannot be completely ruled out.

In conclusion, we have developed a new visible-light mediated aerobic C–H trifluoromethylselenolation of heteroarenes using the air-stable and readily accessible nucleophilic reagent $\text{Me}_4\text{NSeCF}_3$. This atom-economical methodology operates under very mild and metal-free conditions, and has been successfully applied to a wide range of indoles, azaindoles, and pyrroles bearing various functional groups, either in batch or in flow. Extension of this work to other radical trifluoromethylselenolation reactions is currently underway in our laboratory and will be reported in due course.

Experimental Section

General procedure: In a test tube were added the indole **1** (0.2 mmol, 1 equiv), Eosin Y **3d** (0.004 mmol, 2 mol %) and $\text{Me}_4\text{NSeCF}_3$ **2** (0.4 mmol, 2 equiv.). MeCN (2 mL) was added and the solution was stirred at room temperature under green LED irradiation for 18 h opened to air atmosphere. The crude was then purified by preparative TLC to afford the corresponding pure product.


Acknowledgements

AZB thanks LabEx Charmmmat for a Master's student fellowship. BT acknowledges the CNRS research infrastructure RENARD for access to the EPR facilities and the NIEHS for providing free the WINSIM software used to simulate the EPR spectra. LabEx Charmmmat is gratefully acknowledged for the funding of the continuous flow device. Agence Nationale de la Recherche (ANR) is warmly acknowledged for funding (ANR 18-CE07-0039-01).

Keywords: selenium • fluorine • photoredox catalysis • trifluoromethylselenation • indoles

- [1] a) P. Kirsch in *Modern Fluoroorganic Chemistry: Synthesis, Reactivity, Applications*, 2nd, Completely Revised and Enlarged edition, Wiley: New-York, **2013**; b) K. Müller, C. Faeh, F. Diederich, *Science* **2007**, 317, 1881; c) *Bioorganic and Medicinal Chemistry of Fluorine* (Eds.: J.-P. Begue, D. Bonnet-Delpon), Wiley, Hoboken, **2008**; d) J. Wang, M. Sanchez-Rosello, J. L. Acena, C. del Pozo, A. E. Sorochinsky, S. Fustero, V. A. Soloshonok, H. Liu, *Chem. Rev.* **2014**, 114, 2432.
- [2] Selected reviews: a) A. Tlili, F. Toulgoat, T. Billard, *Angew. Chem.* **2016**, 128, 11900; *Angew. Chem. Int. Ed.* **2016**, 55, 11726; b) T. Besset, P. Jubault, X. Pannecoucke, T. Poisson, *Org. Chem. Front.* **2016**, 3, 1004; c) A. Tlili, E. Ismalaj, Q. Glenadel, C. Ghiazza, T. Billard *Chem. Eur. J.* **2018**, 24, 3659; d) C. Ghiazza, T. Billard, A. Tlili, *Chem. Eur. J.* **2019**, 25, 6482.
- [3] a) A. Krief, L. Hevesi in *Organoselenium Chemistry I. Functional Group Transformations*; Springer: Berlin, **1988**; b) H. E. Ganther, *Carcinogenesis* **1999**, 20, 1657; c) R. Naithani, *Mini-Rev. Med. Chem.* **2008**, 8, 657; d) L. V. Papp, A. Holmgren, K. K. Khanna, *Antioxid. Redox Signaling* **2010**, 12,793; e) M. C. Ledesma, B. Jung-Hynes, T. L. Schmit, R. Kumar, H. Mukhtar, N. Ahmad, *Mol. Med.* **2011**, 17, 134; f) R. Mousa, R. N. Dardashti, N. Metanis, *Angew. Chem.* **2017**, 129, 16027; *Angew. Chem. Int. Ed.* **2017**, 56, 15818.
- [4] L. V. Romashov, V. P. Ananikov, *Chem. Eur. J.* **2013**, 19, 17640.
- [5] Q. Glenadel, E. Ismalaj, T. Billard, *Eur. J. Org. Chem.* **2017**, 530.
- [6] Selected trifluoromethylselenolations with nucleophilic SeCF_3 reagents: a) N. V. Kondratenko, A. A. Kolomeyts, V. I. Popov, L. M. Yagupolskii, *Synthesis* **1985**, 667; b) C. Chen, L. Ouyang, Q. Lin, Y. Liu, C. Hou, Y. Yuan, Z. Weng, *Chem. Eur. J.* **2014**, 20, 657; c) Y. Wang, Y. You, Z. Weng, *Org. Chem. Front.* **2015**, 2, 574; d) Q. Lefebvre, R. Pluta, M. Rueping, *Chem. Commun.* **2015**, 51, 4394; e) M. Aufiero, T. Sperger, A. S.-K. Tsang, F. Schoenebeck, *Angew. Chem.* **2015**, 127, 10462; *Angew. Chem. Int. Ed.* **2015**, 54, 10322; f) C. Matheis, V. Wagner, L. J. Goossen, *Chem. Eur. J.* **2016**, 22, 7; g) C. Matheis, T. Krause, V. Bragoni, L. J. Goossen, *Chem. Eur. J.* **2016**, 22, 12270; h) J.-B. Han, T. Dong, D. A. Vicio, C.-P. Zhang, *Org. Lett.* **2017**, 19, 3919; i) T. Dong, J. He, Z.-H. Li, C.-P. Zhang, *ACS Sustain. Chem. Eng.* **2018**, 6, 1327; j) Q.-Y. Han, C.-L. Zhao, T. Dong, J. Shi, K.-L. Tan, C.-P. Zhang, *Org. Chem. Front.* **2019**, 10.1039/C9QO00631A.
- [7] Selected trifluoromethylselenolations with electrophilic SeCF_3 reagents: a) E. Magnier, C. Wakselman, *Collect. Czech. Chem. Commun.* **2002**, 67, 1262; b) Q. Glenadel, E. Ismalaj, T. Billard, *J. Org. Chem.* **2016**, 81, 8268; c) C. Ghiazza, Q. Glenadel, A. Tlili, T. Billard, *Eur. J. Org. Chem.* **2017**, 3812; d) C. Ghiazza, A. Tlili, T. Billard, *Eur. J. Org. Chem.* **2018**, 3680; e) S. Dix, M. Jakob, M. N. Hopkinson, *Chem. Eur. J.* **2019**, 25, 7635.
- [8] a) C. Ghiazza, V. Debrauwer, C. Monnereau, L. Khrouz, M. Médebielle, T. Billard, A. Tlili, *Angew. Chem.* **2018**, 130, 11955; *Angew. Chem. Int. Ed.* **2018**, 57, 11781; b) C. Ghiazza, L. Khrouz, C. Monnereau, T. Billard, A. Tlili, *Chem. Commun.* **2018**, 54, 9909; c) C. Ghiazza, C. Monnereau, L. Khrouz, M. Médebielle, T. Billard, A. Tlili, *Synlett* **2019**, 30, 777.
- [9] a) A. Carboni, G. Dagousset, E. Magnier, G. Masson, *Org. Lett.* **2014**, 16, 1240; b) A. Carboni, G. Dagousset, E. Magnier, G. Masson, *Synthesis* **2015**, 47, 2439; c) L. Jarrige, A. Carboni, G. Dagousset, G. Levitre, E. Magnier, G. Masson *Org. Lett.* **2016**, 18, 2906; d) G. Dagousset, C. Simon, E. Anselmi, B. Tuccio, T. Billard, E. Magnier, *Chem. Eur. J.* **2017**, 23, 4282; e) M. Daniel, G. Dagousset, P. Diter, P.-A. Klein, B. Tuccio, A.-M. Goncalves, G. Masson, E. Magnier, *Angew. Chem.* **2017**, 129, 4055; *Angew. Chem. Int. Ed.* **2017**, 56, 3997; f) G. Levitre, G. Dagousset, E. Anselmi, B. Tuccio, E. Magnier, G. Masson, *Org. Lett.* **2019**, 21, 6005.
- [10] Selected reviews: a) J. P. Knowles, L. D. Elliott, K. I. Booker-Milburn, *Beilstein J. Org. Chem.* **2012**, 8, 2025; b) Y. Su, N. J. W. Straathof, V. Hessel, T. Noël, *Chem. Eur. J.* **2014**, 20, 10562; c) Z. J. Garlets, J. D. Nguyen, C. R. J. Stephenson, *Isr. J. Chem.* **2014**, 54, 351; d) M. B. Plutschack, C. A. Correia, P. H. Seeberger, K. Gilmore, *Top. Organomet. Chem.* **2016**, 57, 43; e) D. Cambié, C. Bottecchia, N. J. W. Straathof, V. Hessel, T. Noël, *Chem. Rev.* **2016**, 116, 10276
- [11] D. Prasad Hari, B. König, *Chem. Commun.* **2014**, 50, 6688.

COMMUNICATION



Green trifluoromethylselenenolation : a radical pathway, demonstrated by physico-chemical studies, allows the regioselective introduction of the SeCF₃ group on indoles and other heterocycles. This metal free photoredox pathway occurs under irradiation with green led on batch and flow conditions.

Arnaud De Zordo-Banliat,^[a] Lucas Barthélémy,^[b] Flavien Bourdreux,^[a] Béatrice Tuccio,^[b] Guillaume Dagousset,^{*[a]} Bruce Pégot,^{*[a]} and Emmanuel Magnier^{*[a]}

Page No. – Page No.

Visible-Light-Induced Metal-Free Trifluoromethylselenenolation of Electron-Rich Heteroarenes using the Nucleophilic [Me₄N][SeCF₃] Reagent