



# Analysis of the Feshbach-Schur method for the Fourier Spectral discretizations of Schrödinger operators

Geneviève Dusson, Israel Michael Sigal, Benjamin Stamm

## ► To cite this version:

Geneviève Dusson, Israel Michael Sigal, Benjamin Stamm. Analysis of the Feshbach-Schur method for the Fourier Spectral discretizations of Schrödinger operators. *Mathematics of Computation*, 2023, 10.1090/mcom/3774 . hal-02919770v2

HAL Id: hal-02919770

<https://hal.science/hal-02919770v2>

Submitted on 3 Dec 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ANALYSIS OF THE FESHBACH–SCHUR METHOD FOR THE FOURIER SPECTRAL DISCRETIZATIONS OF SCHRÖDINGER OPERATORS

GENEVIEVE DUSSON, ISRAEL MICHAEL SIGAL, AND BENJAMIN STAMM

**ABSTRACT.** In this article, we propose a new numerical method and its analysis to solve eigenvalue problems for self-adjoint Schrödinger operators, by combining the Feshbach–Schur perturbation theory with the spectral Fourier discretization. In order to analyze the method, we establish an abstract framework of Feshbach–Schur perturbation theory with minimal regularity assumptions on the potential that is then applied to the setting of the new spectral Fourier discretization method. Finally, we present some numerical results that underline the theoretical findings.

## 1. INTRODUCTION

In this article, we address the problem of the computation of eigenvalues of self-adjoint Schrödinger operators (quantum Hamiltonians) of the form

$$\mathcal{H} = -\Delta + V.$$

Our main results are a priori error estimates for the approximation error of the eigenvalue and eigenfunction for a new method allowing irregular potentials beyond the regularity assumptions within the standard variational setting [2, 1, 12]. The main ingredient of this method and its analysis is a reduction of this infinite-dimensional problem to a finite-dimensional one in a fully controlled way with an effective estimate of the error terms, using the Feshbach–Schur map (FSM) method. This method originated in works of I. Schur on the Dirichlet problem in planar domains and H. Feshbach, on resonances in nuclear physics, and was then developed independently in numerical analysis, computational quantum chemistry and mathematical physics, see [16, 18] with the original techniques called variously the Feshbach projection and Schur complements methods.

We combine this approach with spectral Fourier discretizations which are widely used in numerical methods in electronic structure calculation, especially for condensed matter simulations and in materials science, and known in this community as planewave discretization. Electronic structure calculation is indeed one of the problems we have in our sight. And one particular very useful aspect of planewaves is that they are eigenfunctions of the Laplace operator, entering the Hamiltonian  $\mathcal{H} = -\Delta + V$  that needs to be diagonalized in order to determine the electronic structure of the system.

Our analysis is thus relying on non-variational perturbative techniques. Perturbation-based approaches have a long history in quantum mechanics. For example, different perturbation methods have been proposed, such as [4, 22], traditionally to introduce more physical details e.g. many-particle interactions in a given approximation. The mathematical justification has been provided by the seminal work of Kato [21]. Perturbation methods have also been used to study van der Waals interactions between two hydrogen atoms in the dissociation limit in a mathematically rigorous way in [7, 11]. More recently, a post-processing strategy has been proposed by some of the authors for planewave discretizations for non-linear eigenvalue problems [8, 9, 10, 15], which considers the exact solution as a perturbation of the discrete (using the planewave basis) approximation.

This is in spirit not so far from so-called two-grid methods, where a first problem is solved on a coarse basis, i.e. in a small discretization space, and a small problem is solved on a fine basis. In the case of eigenvalue problems using a same Hamiltonian  $\mathcal{H} = -\Delta + V$  as in this article, two-grid and three-grid methods have been proposed e.g. in [14, 27] within a variational approximation. A two-grid method has also been proposed for nonlinear eigenvalue problems of a Gross–Pitaevskii type equation in [5].

As emphasized above, we extend in this article the FSM-method to establish finite-dimensional approximations based on the spectral Fourier basis to solve the Schrödinger eigenvalue problem with controlled errors on the eigenvalues and eigenvectors. To be a little more concrete, we define a new problem in a coarse, finite-dimensional, subspace  $\mathbf{X}_M \subset \mathbf{X}$  spanned by Fourier modes yielding the exact eigenvalue one would obtain when computing it in the infinite-dimensional space  $\mathbf{X}$ . Indeed, our contribution follows a new Ansatz based on the question: Can we find a discrete Hamiltonian acting on the finite-dimensional space  $\mathbf{X}_M$  that has the exact eigenvalue  $\lambda_*$  of the original Hamiltonian (acting on  $\mathbf{X}$ ) as eigenvalue? It turns that the answer is yes, but that the discrete Hamiltonian depends itself on  $\lambda_*$ , through the Feshbach–Schur (FS) -map, leading to an eigenvalue problem in  $\mathbf{X}_M$  that is nonlinear in the spectral parameter. Not surprisingly, the map cannot be computed exactly but only be approximated through a fast decaying series, that is truncated based on a parameter  $K$ , and which requires computations in a larger space  $\mathbf{X}_N$  with  $\mathbf{X}_M \subset \mathbf{X}_N \subset \mathbf{X}$ . However, this defines a new numerical method that is defined by the three parameters  $\sigma = (N, M, K)$  and that can be rigorously analysed. In this work we quantify the error introduced due to the discretization parameters  $\sigma = (N, M, K)$ .

This article is organized as follows. In Section 2 we present the problem and numerical method that is used to find approximations thereof, as well as the main approximation result of the article and the error bounds on the eigenvalues. Section 3 provides the above-mentioned abstract framework of Feshbach–Schur perturbation theory based on the regularized version of form-boundedness whereas Section 4 contains some technical results needed to prove the main result which

follows in Section 5. Finally, we present in Section 6 some numerical results to illustrate the convergence as well as the error bounds, and we conclude with some perspectives in Section 7.

## 2. SET-UP AND RESULTS

**2.1. Problem statement.** In order to simplify the notation, we consider a cubic lattice  $\mathcal{R} = L\mathbb{Z}^d$  ( $L > 0$ ,  $d = 1, 2, 3$ ), but all our arguments straightforwardly apply to the general case of any Bravais lattice. In this paper we are interested in the spectral theory of the self-adjoint Schrödinger operators (quantum Hamiltonians)

$$\mathcal{H} := -\Delta + V,$$

with reasonably regular,  $\mathcal{R}$ -periodic potentials  $V$ , acting on the Hilbert space

$$\mathsf{L}_{\text{per}}^2 := \{u \in L_{\text{loc}}^2(\mathbb{R}^d) \mid u \text{ is } \mathcal{R}\text{-periodic}\},$$

endowed with the scalar product  $\langle u, v \rangle = \int_{\Omega} u(\mathbf{r}) v(\mathbf{r}) d\mathbf{r}$  and the induced norm  $\|\cdot\|$ , where  $\Omega = [0, L]^d$  is the chosen fundamental cell of the lattice  $\mathcal{R} = L\mathbb{Z}^d$ . Note that  $V$  acts as a multiplicative operator, whereas  $V$  is either a function (in the more regular case) or a distribution (in the less regular case), whose regularity will be discussed shortly.

Specifically, we would like to solve the eigenvalue problem

$$\mathcal{H}\varphi = \lambda\varphi, \tag{1}$$

in a space  $\mathsf{X} \subset \mathsf{H}_{\text{per}}^1$ . Here,  $\mathsf{H}_{\text{per}}^1$  is the Sobolev space of index 1 of periodic functions on  $\Omega$ , which is defined in precise terms later on and equation (1) is considered in the weak sense.

To this end we use the Feshbach–Schur method to reduce the problem to a finite dimensional one. To simplify the exposition, we will assume that the eigenvalue of interest  $\lambda_*$  is isolated, which is true for the smallest eigenvalue under fairly general assumptions of  $V$ , see Theorems XIII.46 - XIII.48 of the textbook [25]. We denote by  $\|\cdot\|$  the operator norm on  $\mathcal{L}(\mathsf{L}_{\text{per}}^2)$ , the space of bounded linear operators on  $\mathsf{L}_{\text{per}}^2$ . To formulate our condition on the potential  $V$ , we introduce the following norm measuring its regularity

$$\|V\|_r := \|(-\Delta + \text{Id})^{-1/2+r/2} V (-\Delta + \text{Id})^{-1/2+r/2}\|,$$

where the operator  $(-\Delta + \text{Id})^s$  is defined by the Fourier transform (cf. Appendix A). In what follows, we thus assume that the potential  $V$  satisfies the following condition.

**Assumption 1.** *The potential  $V$  is real,  $\mathcal{R}$ -periodic and satisfies*

$$\|V\|_r < \infty \quad \text{for some } r > 0.$$

Assumption 1 implies that  $V$  is  $\Delta$ -form bounded [13, 24], which corresponds to  $r = 0$ . The latter, weaker property implies that  $\mathcal{H}$  (a) is self-adjoint; (b) is bounded below and (c) has purely discrete spectrum (see e.g. [13, 24, 25, 20]).

Moreover, potentials  $V$  belonging to the Sobolev spaces,  $\mathsf{H}_{\text{per}}^s := (-\Delta + 1)^{s/2} \mathsf{L}_{\text{per}}^2$  satisfy this assumption as shown in Appendix A, Lemma 13 for  $r \leq s + 1$  and  $r < 1 + \frac{s}{2} - \frac{d}{4}$ . In terms of Sobolev spaces, Assumption 1 states that  $V$ , as an operator, maps  $\mathsf{H}_{\text{per}}^{1-r}$  into  $\mathsf{H}_{\text{per}}^{-1+r}$ .

As an example, we note that the Coulomb potential  $V(x) = \frac{Z}{|x|}$  in three dimensions ( $d = 3$ ) satisfies  $V \in \mathsf{H}_{\text{per}}^s$ ,  $s < 1/2$ . Thus, in view of the above discussion we conclude that Assumption 1 is satisfied for any  $r < 1/2$ . For potentials  $V \in \mathsf{L}_{\text{per}}^2$ , thus with  $s = 0$ , the above discussion yields that Assumption 1 is satisfied for any  $r < 1 - \frac{d}{4}$ .

**2.2. Approach.** In our approach, we reduce the exact infinite dimensional eigenvalue problem to a finite dimensional one in a controlled way for fairly irregular potentials. Of course, we have to pay a price for this, which is that at one point we solve a one-dimensional fixed point problem that can be equivalently seen as a non-linear eigenvalue problem. A key ingredient of our method is the finite dimensional space and the corresponding orthogonal projection onto which we map the original problem to obtain a reduced, finite-dimensional one.

Let  $\mathsf{X}_M$  denote the subspace of  $\mathsf{L}_{\text{per}}^2$  spanned by the eigenfunctions of  $-\Delta$  on  $\mathcal{R}$ , with eigenvalues smaller than  $\rho_M$ , as

$$\mathsf{X}_M = \left\{ \sum_{k \in \mathcal{R}^*, |k| \leq M} \hat{u}_k e_k(x) \mid \hat{u}_k^* = \hat{u}_{-k}, \hat{u}_k \in \mathbb{C} \right\}, \quad (2)$$

where  $e_k(x) = |\Omega|^{-\frac{1}{2}} e^{ik \cdot x}$  (Fourier modes, also called planewaves),  $\mathcal{R}^* = \frac{2\pi}{L} \mathbb{Z}^d$ , and

$$\rho_M := \left( \frac{2\pi M}{L} \right)^2.$$

Let  $\mathsf{P}_M$  be the  $\mathsf{L}_{\text{per}}^2$ -orthogonal projection onto  $\mathsf{X}_M$  and  $\mathsf{P}_M^\perp := \text{Id} - \mathsf{P}_M$ . We consider the Galerkin approximation of the linear Hamiltonian  $\mathcal{H} := -\Delta + V$ ,

$$\mathcal{H}_M := \mathsf{P}_M(-\Delta + V)\mathsf{P}_M.$$

Let  $\varphi$  denote an eigenfunction of (1), introduce the projections  $\varphi_M = \mathsf{P}_M \varphi$  and  $\varphi_M^\perp = \mathsf{P}_M^\perp \varphi$  and project the exact eigenvalue problem (1) onto the subspace  $\mathsf{X}_M$  and its complement  $\mathsf{X}_M^\perp$  to obtain

$$\mathsf{P}_M(\mathcal{H}_M - \lambda)\varphi_M = -\mathsf{P}_M V \varphi_M^\perp, \quad (3)$$

$$\mathsf{P}_M^\perp(\mathcal{H}_M^\perp - \lambda)\varphi_M^\perp = -\mathsf{P}_M^\perp V \varphi_M, \quad (4)$$

where  $\mathcal{H}_M^\perp := \mathsf{P}_M^\perp \mathcal{H} \mathsf{P}_M^\perp$ . Here and in the remainder of this article, we abuse notation and write  $\lambda$  instead of  $\lambda \text{Id}$  in order to denote the multiplicative operator. Next, in Appendix A, we prove the following

**Lemma 1.** *Let Assumption 1 hold and define  $\kappa_M := \rho_M - (\rho_M + 1) \rho_M^{-r} \|V\|_r$ . Then*

$$\mathcal{H}_M^\perp \geq \kappa_M \text{ on } \text{Ran } \mathsf{P}_M^\perp. \quad (5)$$

Here,  $\text{Ran}$  denotes the range (or image) of the following operator.

Thus for  $\lambda < \kappa_M$ , the operator  $\mathcal{H}_M^\perp - \lambda$  is invertible and we can solve (4) for  $\varphi_M^\perp$  and thus  $\varphi_M^\perp = -(\mathcal{H}_M^\perp - \lambda)^{-1} \mathsf{P}_M^\perp V \varphi_M$ . Substituting the result into (3), we obtain the non-linear eigenvalue problem

$$(\mathcal{H}_M + U_M(\lambda)) \varphi_M = \lambda \varphi_M, \quad (6)$$

where we introduced the *effective interaction*  $U_M(\lambda) : \mathsf{X}_M \rightarrow \mathsf{X}_M$ , or a Schur complement,

$$U_M(\lambda) := -\mathsf{P}_M V \mathsf{P}_M^\perp (\mathcal{H}_M^\perp - \lambda)^{-1} \mathsf{P}_M^\perp V \mathsf{P}_M. \quad (7)$$

We then have the following proposition, which is proved in Appendix A.

**Proposition 1.** *For each  $\lambda$  such that  $\lambda < \kappa_M$ ,  $U_M(\lambda)$  is a well-defined operator as a product of three maps:  $\mathsf{P}_M, V$  and  $\mathsf{P}_M^\perp (\mathcal{H}_M^\perp - \lambda)^{-1} \mathsf{P}_M^\perp$  between various but matching Sobolev spaces.*

Now, we construct a completely computable approximation of the eigenvalue problem (6), with the operators involved being sums of products of finite matrices. Namely, we expand the resolvent  $(\mathcal{H}_M^\perp - \lambda)^{-1}|_{\text{Ran } \mathsf{P}_M^\perp} = (-\Delta + V_M^\perp - \lambda)^{-1}|_{\text{Ran } \mathsf{P}_M^\perp}$  in (7) in the formal Neumann series in  $V_M^\perp := \mathsf{P}_M^\perp V \mathsf{P}_M^\perp$ , then truncate this series at  $K \in \mathbb{N}$  and replace the projections  $\mathsf{P}_M^\perp = \text{Id} - \mathsf{P}_M$  by  $\mathsf{P}_M^N := \mathsf{P}_N - \mathsf{P}_M$ , with  $N > M$ . Introducing the notation

$$\mathsf{G}_M^N(\lambda) := (-\Delta - \lambda)|_{\text{Ran } \mathsf{P}_M^N}^{-1} \quad (8)$$

and  $V_M^N := \mathsf{P}_M^N V \mathsf{P}_M^N$ , we obtain the following truncated effective interaction

$$U_\sigma(\lambda) := -\mathsf{P}_M V \mathsf{P}_M^N R_\sigma(\lambda) \mathsf{P}_M^N V \mathsf{P}_M, \quad (9)$$

where  $\sigma := (N, M, K)$  and  $R_\sigma(\lambda) := \sum_{k=0}^K (-1)^k [\mathsf{G}_M^N(\lambda) V_M^N]^k \mathsf{G}_M^N(\lambda)$ . Since all the operators involved in (9) are finite matrices, this family is well-defined and computable. Now, we define  $\mathcal{H}_\sigma(\lambda) := \mathcal{H}_M + U_\sigma(\lambda)$  on  $\mathsf{X}_M$  and consider the eigenvalue problem: find an eigenvalue  $\lambda_{\sigma i}$  and the corresponding eigenfunctions  $\varphi_{\sigma i} \in \mathsf{X}_M$  such that

$$\mathcal{H}_\sigma(\lambda_{\sigma i}) \varphi_{\sigma i} = \lambda_{\sigma i} \varphi_{\sigma i}. \quad (10)$$

Next, we define the approximate ‘lifting’ operator whose origin will be become clear in the next section:

$$Q_\sigma(\lambda) := \text{Id} - R_\sigma(\lambda) \mathsf{P}_M^N V \mathsf{P}_M. \quad (11)$$

Note that in the case of  $N = M$  there holds  $\mathsf{P}_M^N = 0$ ,  $U_\sigma(\lambda) = 0$ , and thus, equation (10) (and (6)) simply reduces to the variational approximation involving the Hamiltonian  $\mathcal{H}_M$ .

**2.3. Main results.** Within this manuscript, we denote by  $\lesssim$  upper bounds involving constants that do not depend on the parameters  $\sigma = (N, M, K), \alpha, r, \|V\|_r$ . Then, we have the following result, whose proof will be provided in Section 5.

**Theorem 1.** *Let Assumption 1 hold, let  $\lambda_\star$  be an isolated eigenvalue of  $\mathcal{H}$  of finite multiplicity  $m$ , with eigenfunctions  $\varphi_i$ , and let  $\gamma_0$  denote the gap between  $\lambda_\star$  and the rest of the spectrum of  $\mathcal{H}$ .*

*Then, there exists  $\alpha > 0$  and  $M_0 \in \mathbb{N}$  such that for  $N \geq M \geq M_0$ , problem (10) has  $m$  solutions  $(\varphi_{\sigma i}, \lambda_{\sigma i}) \in \mathsf{X}_M \times [\lambda_\star - \frac{\gamma_0}{2}, \lambda_\star + \frac{\gamma_0}{2}]$  approximating  $(\varphi_i, \lambda_\star)$  in the following sense:*

$$|\lambda_\star - \lambda_{\sigma i}| \lesssim (\lambda_\star + \alpha) \frac{\|V\|_r^2}{\alpha^r} \varepsilon(\sigma, r, V), \quad (12)$$

$$\|\varphi_i - Q_\sigma(\lambda_{\sigma i})\varphi_{\sigma i}\| \lesssim \|V\|_r \left[ 1 + \frac{\lambda_\circ}{\gamma_0} \frac{\|V\|_r}{\alpha^r} \right] \varepsilon(\sigma, r, V), \quad (13)$$

where  $\lambda_\circ = \lambda_\star + \gamma_0 + \alpha$  and

$$\varepsilon(\sigma, r, V) := \rho_N^{-r} + \rho_M^{-r} [4\rho_M^{-r} \|V\|_r]^{K+1}.$$

This Theorem is subject to several remarks.

**Remark 1.** Note that  $\varepsilon$  is equivalent to

$$\varepsilon(\sigma, r, V) \approx N^{-2r} + M^{-2r} \left[ 4 \left( \frac{L}{2\pi} \right)^{2r} M^{-2r} \|V\|_r \right]^{K+1},$$

where the equivalence constants do not depend on the parameters  $\sigma = (N, M, K), r, \alpha, V$ .

**Remark 2.** In some cases, for instance in multi-scale problems, one might be only interested in the coarse-scale solution, i.e. the best-approximation in the coarse space  $\mathsf{X}_M$  given by  $\mathsf{P}_M \varphi_i$ . In such cases, a useful byproduct of the proof of Theorem 1 is the following estimate

$$\|(-\Delta + \text{Id})^s (\mathsf{P}_M \varphi_i - \varphi_{\sigma i})\| \lesssim \frac{\lambda_\circ}{\gamma_0} \frac{\|V\|_r^2}{\alpha^r} \rho_M^s \varepsilon(\sigma, r, V), \quad (14)$$

for any  $s \geq 0$ , which thus compares the eigenfunctions in the space  $\mathsf{X}_M$ .

**Remark 3.** Note that convergence of the eigenvalues and the eigenfunctions can be achieved by taking the limit  $K, N \rightarrow \infty$  for fixed  $M \geq M_0$ . For practical purposes, the idea is to set  $N$  large enough so that the error is dominated by the error introduced in  $K < +\infty$ .

Further, note that the eigenvalue and eigenvector errors have the same rate of convergence with respect to  $K$ . However, the error in the eigenvector depends on the gap  $\gamma_0$  while the error in the eigenvalue does not.

The estimate with respect to  $N$  in Theorem 1 is not sharp in all cases, in particular for sufficiently regular potentials  $V$ . Nonetheless, our analysis has the merit of presenting the convergence result in one combined analysis based on perturbative techniques which also holds for low regularities of the potential where standard *a priori* convergence results of the variational approximation do not hold. In fact, still in the low regularity regime, an estimate of the variational problem can be obtained by setting  $K \rightarrow \infty$  or  $M = N$ .

Note that we can adapt the result whenever *a priori* approximation results are available by employing the triangle inequality. Indeed, if the potential  $V$  belongs to the Sobolev space  $V \in \mathsf{H}_{\text{per}}^s$ , with  $s > d/2$ , we resort to *a priori* results in a first place to obtain a sharp bound with respect to  $N$ , see e.g., [1, 6, 3], and also [23] for a certain class of discontinuous potentials in  $H^{1/2-\varepsilon}$  for all  $\varepsilon > 0$ , in two dimensions.

More precisely, we consider  $\mathcal{H}$  acting on  $\mathsf{X}_N$  directly, i.e. substituting  $\mathcal{H}$  by  $\mathcal{H}_N := \mathsf{P}_N \mathcal{H} \mathsf{P}_N$  and using  $\mathsf{X} = \mathsf{X}_N$  with variational solution  $(\varphi_N, \lambda_N)$ , assuming a simple eigenvalue for simplicity. It is important to note that problem (10) remains unchanged and thus, the result of Theorem 1 holds with

$$\tilde{\varepsilon}(\sigma, r, V) := \rho_M^{-r} [4\rho_M^{-r} \|V\|_r]^{K+1},$$

but where the exact solution is substituted by  $(\varphi_N, \lambda_N)$ . Proceeding then by the triangle inequality yields

$$\begin{aligned} |\lambda_\star - \lambda_{\sigma i}| &\leq |\lambda_\star - \lambda_N| + |\lambda_N - \lambda_{\sigma i}|, \\ \|\varphi_i - Q_\sigma(\lambda_{\sigma i})\varphi_{\sigma i}\| &\leq \|\varphi_i - \varphi_N\| + \|\varphi_N - Q_\sigma(\lambda_{\sigma i})\varphi_{\sigma i}\|. \end{aligned}$$

Combining then the aforementioned *a priori* estimates from [1, 6, 3] for the first terms of the right hand sides with Theorem 1 for the latter parts yields the following corollary.

**Corollary 2.** *Under the conditions of Theorem 1 and if  $V \in \mathsf{H}_{\text{per}}^s$ ,  $s > d/2$ , then*

$$\begin{aligned} |\lambda_\star - \lambda_{\sigma i}| &\leq C (N^{-(2s+2)} + \tilde{\varepsilon}(\sigma, r, V)), \\ \|\varphi_i - Q_\sigma(\lambda_{\sigma i})\varphi_{\sigma i}\| &\leq C (N^{-(s+2)} + \tilde{\varepsilon}(\sigma, r, V)), \end{aligned}$$

for some constant  $C > 0$  independent on  $\sigma = (N, M, K)$ .

**Remark 4.** *Instead of the space  $\mathsf{X}_M$  defined in (2) spanned by the eigenfunctions of  $-\Delta$  on  $\mathcal{R}$ , we could have taken another finite dimensional approximation of the space  $L^2(\mathbb{R})$ .*

**Remark 5.** *Given a sequence of larger and larger finite dimensional spaces, one could apply the FS maps consequently with larger and larger projections. In particular, this could lead to a version of a multigrid technique.*

**2.4. Theoretical background.** Let us shed light on the theoretical foundation on the eigenvalue formulation in form of (6), instead of (1). For a pair of projections  $P$  and  $P^\perp$  such that  $P + P^\perp = \text{Id}$ , we define the set  $D_P$  of operators  $H$  on a Hilbert space such that  $H^\perp := P^\perp H P^\perp$  is invertible on  $\text{Ran } P^\perp$  and the operators  $HP$  and  $PH$  are bounded. Furthermore, for an operator  $H \in D_P$ , we define the bounded operator

$$F_P(H) := P(H - HR^\perp H)P, \quad (15)$$

where  $R^\perp := P^\perp(H^\perp)^{-1}P^\perp$ , acting on the subspace  $\text{Ran } P$ . Our approach is based on the following results, originally presented in [18, Theorem 11.1].

**Theorem 3.** *Let  $P$  and  $P^\perp$  be a pair of projections such that  $P + P^\perp = \text{Id}$ . Then  $F_P: H \rightarrow F_P(H)$ , considered as a map from the subspace  $D_P$  into the space of bounded operators acting on the subspace  $\text{Ran } P$ , is isospectral in the following sense: if an operator  $H$  and a number  $\lambda \in \mathbb{R}$  are such that  $H - \lambda \in D_P$ , then*

- (a)  $\lambda \in \sigma(H) \iff 0 \in \sigma(F_P(H - \lambda))$ ;
- (b)  $H\psi = \lambda\psi \iff F_P(H - \lambda)\varphi = 0$ ;
- (c)  $\dim \text{Null}(H - \lambda) = \dim \text{Null}F_P(H - \lambda)$ .

Moreover,  $\psi$  and  $\varphi$  in (b) are related as  $\varphi = P\psi$  and  $\psi = Q_P(\lambda)\varphi$ , where

$$Q_P(\lambda) := P - P^\perp(H^\perp - \lambda)^{-1}P^\perp HP.$$

Finally, if  $H$  is self-adjoint, then so is  $F_P(H)$ .

Here,  $\text{Null}$  denotes the null space (or kernel) of the following operator. The map  $F_P$  on the space of operators, is called the *Feshbach–Schur map*. The relation  $\psi = Q_P(\lambda)\varphi$  allows us to reconstruct the full eigenfunction from the projected one. By statement (a), we have

**Corollary 4.** *Let  $\nu_i(\lambda)$  denote the  $i$ -th eigenvalue of the operator  $F_P(H - \lambda) + \lambda$  for each  $\lambda$  in an interval  $I \subset \mathbb{R}$ . Then, there exists a bijection between the eigenvalues of  $H$  in  $I$  and the solutions of the equation*

$$\nu_i(\lambda) = \lambda.$$

In the current setting of the spectral Fourier approximations,  $P = P_M$ ,  $P^\perp = P_M^\perp$ , Proposition 1 implies that the results of Theorem 3 apply for each choice of  $M \in \mathbb{N}$  and yield

$$F_{P_M}(\mathcal{H} - \lambda) = \mathcal{H}_M(\lambda) - \lambda P_M, \quad (16)$$

where we introduced the notation

$$\mathcal{H}_M(\lambda) := \mathcal{H}_M + U_M(\lambda). \quad (17)$$

Note that  $\mathcal{H}_M(\lambda)$  is exactly the operator entering (6). Thus, we have the following.

**Corollary 5.** *Let  $\lambda \in \mathbb{C}$  with  $\text{Re } \lambda < \kappa_M$ . Then*

- (a)  $\mathcal{H}\psi = \lambda\psi \iff (\mathcal{H}_M(\lambda) - \lambda)\varphi_M = 0$ ;
- (b)  $\dim \text{Null}(\mathcal{H} - \lambda) = \dim \text{Null}(\mathcal{H}_M(\lambda) - \lambda)$ .

(c)  $\psi$  and  $\varphi$  in (a) are related as  $\varphi_M = \mathbf{P}_M\psi$  and  $\psi = \mathbf{Q}_M(\lambda)\varphi_M$ , where

$$\mathbf{Q}_M(\lambda) = \text{Id} - (\mathcal{H}_M^\perp - \lambda)^{-1}\mathbf{P}_M^\perp V \mathbf{P}_M. \quad (18)$$

i.e. the corresponding eigenfunction can be reconstructed from  $\varphi_M$  by an explicit linear map.

This result shows that the original infinite-dimensional spectral problem (1) is equivalent to the finite dimensional spectral problem (6) which is nonlinear in the spectral parameter  $\lambda$ . We now state a few properties of the effective interaction  $U_M(\lambda)$ , in order to characterize the solutions of the fixed-point problems  $\nu_i(\lambda) = \lambda$ . First, we give a definition. A family of bounded, self-adjoint operators  $T(\lambda) : X \rightarrow X$ , with  $\lambda \in I \subset \mathbb{R}$ , is said to be monotonically decreasing if  $T(\lambda) < T(\lambda')$ , in the sense of quadratic forms (i.e.  $\langle u, T(\lambda)u \rangle < \langle u, T(\lambda')u \rangle$ ), for all  $\lambda > \lambda'$ . Similarly, one can define increasing, non-increasing, and non-decreasing families of operators. If  $T(\lambda)$  is a weakly differentiable family, then, by the fundamental theorem of calculus, if  $T'(\lambda) \leq 0$  and is not identically 0, then  $T(\lambda)$  is monotonically decreasing.

**Proposition 2.** For  $\lambda \in \mathbb{R}$  such that  $\lambda < \kappa_M$ ,  $U_M(\lambda)$  is (i) non-positive, (ii) monotonically decreasing with  $\lambda$ , (iii) vanishing as  $\lambda \rightarrow -\infty$ . For  $\lambda \in \mathbb{C}$  such that  $\text{Re } \lambda < \kappa_M$ ,  $U_M(\lambda)$  is (iv) complex analytic in  $\lambda$  and (v) symmetric.

*Proof.* Properties (i)-(iv) follow directly from definition (7) and Lemma 1 above. For the last one, we use that  $\mathcal{H}$  is self-adjoint.  $\square$

**Proposition 3.** Denote by  $\nu_{Mi}(\lambda)$  the  $i$ -th eigenvalue of  $\mathcal{H}_M(\lambda)$  and assume that the  $i$ -th eigenvalue of  $\mathcal{H}$  is less than  $\kappa_M$ . Then, the equation  $\nu_{Mi}(\lambda) = \lambda$  has a unique solution in the interval  $(-\infty, \kappa_M)$ .

*Proof.* Since, for  $\lambda < \kappa_M$ ,  $U_M(\lambda)$  is symmetric, the operator  $\mathcal{H}_M(\lambda)$  defined by (17) is (a) self-adjoint, (b) monotonically decreasing with  $\lambda$ , (c) converging to  $\mathcal{H}_M$  as  $\lambda \rightarrow -\infty$ , (d) is complex analytic in  $\lambda$  for  $\text{Re } \lambda < \kappa_M$ . We deduce from (b) that the functions  $\nu_{Mi}$  are decreasing on  $(-\infty, \kappa_M)$  and thus, if the  $i$ -th eigenvalue of  $\mathcal{H}$  is less than  $\kappa_M$ , the equation  $\nu_{Mi}(\lambda) = \lambda$  has a unique solution (that equals the  $i$ -th eigenvalue of  $\mathcal{H}$ ).  $\square$

Note also that  $\lim_{\lambda \rightarrow -\infty} \nu_{Mi}(\lambda)$  is the  $i$ -th eigenvalue of  $\mathcal{H}_M$  which is larger than the  $i$ -th eigenvalue of  $\mathcal{H}$  due to the variational principle.

These considerations motivate the numerical strategies to compute solutions to (10) in the following section.

**2.5. Numerical strategy.** In order to find solutions to the non-linear eigenvalue problem (10), we propose two strategies:

*Strategy 1:* For a fixed index  $i = 1, \dots, M$ , consider the sequence of iterates  $\lambda_\sigma^{(j)}$  obtained by

$$\lambda_\sigma^{(j)} : \text{ is the } i\text{-th eigenvalue of } \mathcal{H}_\sigma(\lambda_\sigma^{(j-1)}). \quad (19)$$

We thus introduce the notation  $\nu_{\sigma i}(\lambda)$  denoting the  $i$ -th eigenvalue (counting multiplicities) of the Hamiltonian  $\mathcal{H}_\sigma(\lambda)$  and thus have  $\lambda_\sigma^{(j)} = \nu_{\sigma i}(\lambda_\sigma^{(j-1)})$ . The limit value  $\lambda_\sigma := \lim_{j \rightarrow \infty} \lambda_\sigma^{(j)}$  then satisfies  $\lambda_\sigma = \nu_{\sigma i}(\lambda_\sigma)$  and thus (10).

*Strategy 2:* For a given target value  $\lambda_t \in \mathbb{R}$ , consider the sequence of iterates  $\lambda_\sigma^{(j)}$  obtained by

$$\lambda_\sigma^{(j)} : \text{ is the eigenvalue of } \mathcal{H}_\sigma(\lambda_\sigma^{(j-1)}) \text{ closest to } \lambda_t. \quad (20)$$

We thus introduce the notation  $\nu_{\sigma t}(\lambda)$  denoting the eigenvalue of the Hamiltonian  $\mathcal{H}_\sigma(\lambda)$  closest to  $\lambda_t$  and thus have  $\lambda_\sigma^{(j)} = \nu_{\sigma t}(\lambda_\sigma^{(j-1)})$ . The limit value  $\lambda_\sigma := \lim_{j \rightarrow \infty} \lambda_\sigma^{(j)}$  then satisfies  $\lambda_\sigma = \nu_{\sigma t}(\lambda_\sigma)$  and thus (10).

In both cases, as outlined in the upcoming Remark 10, convergence of the fixed-point maps (19) and (20) can be guaranteed under some conditions and for  $N, M$  large enough.

**Remark 6.** *This numerical strategy can be easily extended for eigenvalues with multiplicity higher than one. Indeed, using Proposition 3, provided that the sought eigenvalue is less than  $\kappa_M$ , one can iteratively compute a given number of the lowest eigenvalues of the operator  $\mathcal{H}_\sigma(\lambda_\sigma)$ , with  $\lambda_\sigma$  updated at each iteration, matching a close value of the multiple eigenvalue. At convergence, this eigenvalue and the multiplicity will match the exact one, up to an error given in Theorem 1. Regarding bands of eigenvalues, one could as well iteratively compute a given number of the lowest eigenvalues of the operators  $\mathcal{H}_\sigma(\lambda_{\sigma i})$ , with the  $\lambda_{\sigma i}$  updated at each iteration, but this multiplies the number of problems that have to be solved by the number of eigenvalues in the band. To do it in an efficient manner would require to consider a density-matrix formalism, in order to consider all interesting eigenvectors together, but this goes beyond the scope of the present paper.*

The complexity of the numerical method can be summarized as follows:

- (1) We assume that the resolution of the non-linearity with the iterative scheme (19) or (20) requires  $J$  iterations.
- (2) At each iteration  $j = 1, \dots, J$ , one requires the resolution of an eigenvalue problem with  $O(M^d)$  degrees of freedom and we assume that this problem is solved with an iterative solver using  $J_{\text{evp}}^M$  matrix-vector multiplications.
- (3) The computation of one matrix-vector product corresponding to the Hamiltonian  $\mathcal{H}_M + U_M(\lambda) = -P_M \Delta P_M + P_M V P_M + U_M(\lambda)$  contains three parts.
  - (i) the matrix corresponding to  $P_M \Delta P_M$  is diagonal in the Fourier basis;
  - (ii) the application of  $P_M V P_M$  can be effected using the Fast Fourier Transform (FFT) on the grid  $X_M$  and scales as  $M^d \log(M^d)$ ;
  - (iii) the matrix-vector product corresponding with the application of the

multiplicative potential  $U_\sigma(\lambda)$ . For convenience we recall here its definition:

$$U_\sigma(\lambda) = -\mathbf{P}_M V \mathbf{P}_M^N \left( \sum_{k=0}^K (-1)^k \left[ \mathbf{G}_M^N(\lambda) V_M^N \right]^k \mathbf{G}_M^N(\lambda) \right) \mathbf{P}_M^N V \mathbf{P}_M,$$

with  $\mathbf{G}_M^N(\lambda) = (-\Delta - \lambda)|_{\text{Ran } \mathbf{P}_M^N}^{-1}$ . The application of  $\mathbf{G}_M^N(\lambda)$  scales as  $N^d - M^d$  and the application of the potentials  $\mathbf{P}_M V \mathbf{P}_M^N$ ,  $V_M^N$  and  $\mathbf{P}_M^N V \mathbf{P}_M$  can be performed again using the FFT on the entire grid  $X_N$  and it scales as  $N^d \log(N^d)$ .

The overall complexity is thus proportional to

$$J \times J_{\text{evp}}^M \times \left( M^d + M^d \log(M^d) + (K+3)N^d \log(N^d) + (K+2)(N^d - M^d) \right)$$

In contrast, following the same notation, the variational problem requires a complexity of

$$J_{\text{evp}}^N \times N^d \log(N^d)$$

Therefore in order to compare the methods, it depends on  $J \times J_{\text{evp}}^M(K+3)$  versus  $J_{\text{evp}}^N$  at comparable accuracy.

Finally, we emphasize that the focus of the present paper is to present the numerical analysis of this new method, which covers cases where the analysis of the standard variational approximation does not hold and efficiency becomes a secondary factor.

### 3. PERTURBATION ESTIMATES

In this article, we often deal with the following eigenvalue perturbation problem: Given an operator  $H$  on a Hilbert space  $X$  of the form

$$H = H_0 + W, \tag{21}$$

where  $H_0$  is an operator with some isolated eigenvalues and  $W$  is small in an appropriate norm, show that  $H$  has eigenvalues near those of  $H_0$  and estimate these eigenvalues and the corresponding eigenvectors. We therefore start by presenting an abstract theory which will be applied to our concrete problem in the following sections.

Specifically, we assume that  $H$  and  $H_0$  are self-adjoint and bounded from below and that  $W$  is  $\alpha$ -form-bounded w.r.t. of  $H_0$ , in the sense that for  $\alpha \in \mathbb{R}$  such that  $H_0 + \alpha$  is a positive operator ( $H_0 + \alpha > 0$ ), we have

$$\|W\|_{H_0, \alpha} := \|(H_0 + \alpha)^{-1/2} W (H_0 + \alpha)^{-1/2}\| < \infty, \tag{22}$$

where  $(H_0 + \alpha)^{-s}$ ,  $s > 0$ , is defined either by the spectral theory or by the explicit formula

$$(H_0 + \alpha)^{-s} := c_\alpha \int_0^\infty (H_0 + \alpha + \omega)^{-1} d\omega / \omega^s,$$

where  $c_\alpha := [\int_0^\infty (\alpha + \omega)^{-1} d\omega / \omega^s]^{-1}$ . This notion is equivalent to that of the relative form-boundedness, but it gives an important quantification of the latter.

We also note here that, by a known result about relatively form-bounded operators (see e.g. [24, 20]), if  $H_0$  is a self-adjoint, bounded below operator on  $X$  and  $W$  is symmetric and  $\alpha$ -form-bounded w.r.t. of  $H_0$ , then  $H = H_0 + W$  is self-adjoint.

We start with a general result on the eigenvalue difference.

**Proposition 4.** *Let  $H_0$  be a self-adjoint bounded below operator on  $X$  and  $W$  symmetric and  $\alpha$ -form-bounded w.r.t. of  $H_0$ , and let  $H = H_0 + W$ . Let  $\alpha \in \mathbb{R}$  be such that  $H_0 + \alpha > 0$ . Then the eigenvalues of  $H$  and  $H_0$  satisfy the estimates*

$$|\nu_i(H) - \nu_i(H_0)| \leq (\nu_i(H_0) + \alpha) \|W\|_{H_0, \alpha}, \quad (23)$$

where  $\nu_i(A)$  denotes the  $i$ -th eigenvalue of the operator  $A$ .

*Proof.* Let  $u \in X$  be arbitrary and define  $v = (H_0 + \alpha)^{1/2}u$  noting that  $(H_0 + \alpha) > 0$ . Then,

$$\langle u, Hu \rangle = \langle u, H_0 u \rangle + \langle v, (H_0 + \alpha)^{-1/2}W(H_0 + \alpha)^{-1/2}v \rangle.$$

Note that

$$\langle v, (H_0 + \alpha)^{-1/2}W(H_0 + \alpha)^{-1/2}v \rangle \leq \|W\|_{H_0, \alpha} \langle v, v \rangle = \|W\|_{H_0, \alpha} \langle u, (H_0 + \alpha)u \rangle,$$

and therefore

$$\begin{aligned} \langle u, H_0 u \rangle (1 - \|W\|_{H_0, \alpha}) - \alpha \|u\|^2 \|W\|_{H_0, \alpha} &\leq \langle u, Hu \rangle \\ &\leq \langle u, H_0 u \rangle (1 + \|W\|_{H_0, \alpha}) + \alpha \|u\|^2 \|W\|_{H_0, \alpha}. \end{aligned}$$

Using the min-max principle (Courant–Fisher), there holds

$$\nu_i(H_0) (1 - \|W\|_{H_0, \alpha}) - \alpha \|W\|_{H_0, \alpha} \leq \nu_i(H) \leq \nu_i(H_0) (1 + \|W\|_{H_0, \alpha}) + \alpha \|W\|_{H_0, \alpha},$$

which leads to the result.  $\square$

Let us now assume that  $\lambda_0$  is an isolated eigenvalue of  $H_0$  of finite multiplicity  $m$  and let  $P_0$  be the orthogonal projection onto the span of the eigenfunctions of  $H_0$  corresponding to the eigenvalue  $\lambda_0$ , and let  $P_0^\perp := \text{Id} - P_0$ . We further introduce  $H_{0,-\lambda} := H_0^\perp - \lambda P_0^\perp$  and thus  $H_{0,\alpha} = H_0^\perp + \alpha P_0^\perp$ .

Let  $\gamma_0$  denote the gap of  $\lambda_0$  to its closest eigenvalue in the remaining spectrum of  $H_0$  and we introduce the spectral interval  $I_0 = [\lambda_0 - \frac{1}{2}\gamma_0, \lambda_0 + \frac{1}{2}\gamma_0]$ .

Our next result gives estimates on the difference of eigenvectors of  $H$  and  $H_0$ , as well as on the difference of their corresponding eigenvalues. For standard approaches to the spectral perturbation theory, see [26, 21, 25, 20].

**Theorem 6.** *Let  $H_0$  be a self-adjoint bounded below operator on  $X$ , with the eigenvalue  $\lambda_0$  as above, and  $W$  symmetric and  $\alpha$ -form-bounded w.r.t.  $H_0$ , and let  $H = H_0 + W$ . Let  $\alpha \in \mathbb{R}$  be such that  $H_0 + \alpha > 0$ . If  $\|W\|_{H_0, \alpha} \leq \frac{1}{2} \frac{\gamma_0}{\lambda_0 + \alpha}$ , then*

the self-adjoint operator  $H$  has exactly  $m$  eigenvalues (counting the multiplicities), denoted by  $\mu_i$ , in the interval  $I_0 = [\lambda_0 - \frac{1}{2}\gamma_0, \lambda_0 + \frac{1}{2}\gamma_0]$  which satisfy

$$|\mu_i - \lambda_0| \leq (\lambda_0 + \alpha) \|W\|_{H_0, \alpha} \leq \frac{1}{2}\gamma_0. \quad (24)$$

Further, if  $\|W\|_{H_0, \alpha} \leq \frac{1}{4}\frac{\gamma_0}{\lambda_0}$ , then any normalized eigenfunction,  $\psi_i$ , of  $H$  for the eigenvalue  $\mu_i$  satisfies the estimates

$$\|H_{0, \alpha}^{1/2}(\psi_{0i} - \psi_i)\| \leq 4\frac{\lambda_0}{\gamma_0}(\lambda_0 + \alpha)^{1/2} \|W\|_{H_0, \alpha}, \quad (25)$$

$$\|\psi_{0i} - \psi_i\| \leq 4\frac{\lambda_0}{\gamma_0} \|W\|_{H_0, \alpha}, \quad (26)$$

where  $\lambda_0 = \lambda_0 + \alpha + \gamma_0$  and  $\psi_{0i}$  is an appropriate eigenfunction of  $H_0$  corresponding to the eigenvalue  $\lambda_0$ , namely  $\psi_{0i} := P_0\psi_i$ .

**Remark 7.** We note that similar estimates can be obtained for normalized eigenfunctions  $\tilde{\psi}_{0i} := P_0\psi_i/\|P_0\psi_i\|$  with an additional factor 2 using the estimate

$$\|\tilde{\psi}_{0i} - \psi_i\| \leq |1 - \|\psi_{0i}\|| + \|\psi_{0i} - \psi_i\| \leq |\|\psi_i\| - \|\psi_{0i}\|| + \|\psi_{0i} - \psi_i\| \leq 2\|\psi_{0i} - \psi_i\|.$$

We first develop the following preliminary results.

**Lemma 2.** Let  $\alpha \in \mathbb{R}$  be such that  $H_0 + \alpha > 0$  and  $\lambda_0 := \lambda_0 + \gamma_0 + \alpha$ . Then, for all  $\lambda \in I_0^c := \{z \in \mathbb{C} : \operatorname{Re} z \in I_0\}$ , there holds  $|P_0^\perp - (\lambda + \alpha)H_{0, \alpha}^{-1}P_0^\perp| \geq \frac{\gamma_0}{2\lambda_0}$ .

*Proof.* The eigenvalues of  $|P_0^\perp - (\lambda + \alpha)H_{0, \alpha}^{-1}P_0^\perp|$  on  $\operatorname{Ran} P_0^\perp$  are

$$\left|1 - \frac{\lambda + \alpha}{\lambda_{0i} + \alpha}\right| = \left|\frac{\lambda_{0i} - \lambda}{\lambda_{0i} + \alpha}\right|,$$

where  $\lambda_{0i}$  denotes the eigenvalues of  $H_0$  and the index  $i$  runs over all eigenvalues except  $i$  such that  $\lambda_{0i} = \lambda_0$ . For  $\lambda \in I_0^c$ , we write  $\lambda = \lambda_r + i\lambda_i$ , with  $\lambda_r \in I_0$ ,  $\lambda_i \in \mathbb{R}$ . Since for any  $x \in \mathbb{R}$ ,  $|x - \lambda| \geq |x - \lambda_r|$  we have thus to study the function

$$f(x) = \left| \frac{x - \lambda}{x + \alpha} \right|, \quad x \in K_\alpha := [-\alpha, +\infty) \setminus (\lambda_0 - \gamma_0, \lambda_0 + \gamma_0),$$

for  $\lambda \in I_0$  in order to lower bound the eigenvalues. Since

$$f'(x) = \frac{x - \lambda}{|x - \lambda|} \cdot \frac{\alpha + \lambda}{(x + \alpha)^2},$$

if  $\alpha + \lambda \leq 0$ , there holds  $f'(x) < 0$  for  $x > -\alpha$  so that

$$\min_{x \in K_\alpha} f(x) \geq 1.$$

If  $\alpha + \lambda > 0$ , there holds

$$f'(x) < 0 \text{ for } x < \lambda, \quad f'(x) > 0 \text{ for } x > \lambda,$$

and thus, for  $\lambda \in I_0$ ,

$$\min_{x \in K_\alpha} f(x) = \min(f(\lambda_0 - \gamma_0), f(\lambda_0 + \gamma_0)) = \min\left(\frac{|\lambda_0 - \gamma_0 - \lambda|}{\lambda_0 - \gamma_0 + \alpha}, \frac{|\lambda_0 + \gamma_0 - \lambda|}{\lambda_0 + \gamma_0 + \alpha}\right) \geq \frac{1}{2} \frac{\gamma_0}{\lambda_\circ},$$

yielding the result.  $\square$

Denote  $H^\perp := P_0^\perp H P_0^\perp|_{\text{Ran } P_0^\perp}$  and  $R^\perp(\lambda) := P_0^\perp(H^\perp - \lambda)^{-1}P_0^\perp$ . We have

**Lemma 3.** *Let  $\alpha \in \mathbb{R}$  be such that  $H_0 + \alpha > 0$  and  $\lambda_\circ := \lambda_0 + \gamma_0 + \alpha$ . Let  $I_0^c := \{z \in \mathbb{C} : \text{Re } z \in I_0\}$  and assume  $\|W\|_{H_{0,\alpha}} \leq \frac{1}{4} \frac{\gamma_0}{\lambda_\circ}$ . Then, for  $\lambda \in I_0^c$ , the following statements hold*

- (a) *The operator  $H^\perp - \lambda$  is invertible on  $\text{Ran } P_0^\perp$ ;*
- (b) *The inverse  $R^\perp(\lambda) := P_0^\perp(H^\perp - \lambda)^{-1}P_0^\perp$  defines a bounded, analytic operator-family;*
- (c) *The expression*

$$U(\lambda) := -P_0 H R^\perp(\lambda) H P_0 \tag{27}$$

*defines a finite-rank, analytic operator-family and bounded as*

$$\|U(\lambda)\|_{H_{0,\alpha}} \leq 4 [\lambda_\circ / \gamma_0] \|P_0 W P_0^\perp\|_{H_{0,\alpha}}^2 \leq 4 [\lambda_\circ / \gamma_0] \|W\|_{H_{0,\alpha}}^2. \tag{28}$$

*Further,  $U(\lambda)$  is symmetric for any  $\lambda \in I_0$ .*

*Proof.* (a) Since  $H^\perp$  is self-adjoint, the operator  $H^\perp - \lambda$  is invertible for any  $\lambda \in \mathbb{C} \setminus \mathbb{R}$ . For  $\lambda \in I_0$ , we argue as follows. With the notation  $A^\perp := P_0^\perp A P_0^\perp|_{\text{Ran } P_0^\perp}$ , we write

$$H^\perp = H_0^\perp + W^\perp.$$

Now, we write

$$H^\perp - \lambda P_0^\perp = H_{0,\alpha}^{1/2} [P_0^\perp - (\lambda + \alpha) H_{0,\alpha}^{-1} + K_\lambda] H_{0,\alpha}^{1/2}, \tag{29}$$

with  $K_\lambda = H_{0,\alpha}^{-1/2} W^\perp H_{0,\alpha}^{-1/2}$ . Lemma 2 yields that  $|P_0^\perp - (\lambda + \alpha) H_{0,\alpha}^{-1} P_0^\perp| \geq \frac{\gamma_0}{2\lambda_\circ}$  and thus, the operator  $\text{Id} - (\lambda + \alpha) H_{0,\alpha}^{-1} + K_\lambda$  is invertible as soon as  $\|K_\lambda\| < \frac{\gamma_0}{2\lambda_\circ}$ , which is in particular the case if  $\|W\|_{H_{0,\alpha}} \leq \frac{1}{4} \frac{\gamma_0}{\lambda_\circ}$ . Then, we also have  $|\text{Id} - (\lambda + \alpha) H_{0,\alpha}^{-1} + K_\lambda| \geq \frac{1}{4} \frac{\gamma_0}{\lambda_\circ}$ . Hence the operator  $H^\perp - \lambda$  is a product of three invertible operators and therefore is invertible itself on  $\text{Ran } P_0^\perp$ .

For (b), since  $H^\perp - \lambda$  is invertible on  $\text{Ran } P_0^\perp$ , the interval  $I_0$  is contained in the resolvent set,  $\rho(H^\perp|_{\text{Ran } P_0^\perp})$ , of  $H^\perp|_{\text{Ran } P_0^\perp}$  and therefore, since  $H^\perp|_{\text{Ran } P_0^\perp}$  is self-adjoint,  $I_0^c \subset \rho(H^\perp|_{\text{Ran } P_0^\perp})$ . Since

$$R^\perp(\lambda) := P_0^\perp(H^\perp - \lambda)^{-1}P_0^\perp \tag{30}$$

is the resolvent of the operator  $H^\perp - \lambda$  restricted to  $\text{Ran } P_0^\perp$ , it is analytic on its resolvent set and in particular on  $I_0^c$ .

To prove statement (c), we note that the operators  $R^\perp(\lambda)$ ,  $H P_0$  and  $P_0 H = (H P_0)^*$  are bounded and  $R^\perp(\lambda)$  is symmetric for  $\lambda \in I_0$ . Hence so is  $U(\lambda)$ . The

analyticity of  $U(\lambda)$  follows from the analyticity of  $R^\perp(\lambda)$ . It is clear that  $U(\lambda)$  is of finite rank due to its definition.

Finally, to prove estimate (28), we first show that the operator  $R^\perp(\lambda) := P_0^\perp(H^\perp - \lambda)^{-1}P_0^\perp$ ,  $\lambda \in I_0^c$ , satisfies

$$\|H_{0,\alpha}^{1/2}R^\perp(\lambda)H_{0,\alpha}^{1/2}\| \leq 4 \frac{\lambda_\circ}{\gamma_0}. \quad (31)$$

To this end, we invert (29) on  $\text{Ran } P_0^\perp$  and use that  $|\text{Id} - (\lambda + \alpha)H_{0,\alpha}^{-1} + K_\lambda| \geq \frac{1}{4} \frac{\gamma_0}{\lambda_\circ}$  to obtain (31) for  $\lambda \in I_0^c$ .

Finally, we prove inequality (28). Since  $P_0 H_0 = H_0 P_0$  and  $P_0 P_0^\perp = 0$ , we have

$$P_0 H P_0^\perp = P_0 W P_0^\perp, \quad P_0^\perp H P_0 = P_0^\perp W P_0.$$

These relations and definition (27) yield

$$U(\lambda) = -P_0 W R^\perp(\lambda) W P_0. \quad (32)$$

Combining (31) and (32), we obtain (28).  $\square$

**Remark 8.** In addition, we have the estimate

$$\|U(\lambda)\| \leq 4 \frac{\lambda^2}{\gamma_0} \|P_0 W P_0^\perp\|_{H_{0,\alpha}}^2. \quad (33)$$

Indeed, since  $H_0 P_0 = \lambda_0 P_0$ , we have  $\|(H_0 + \alpha)^{1/2} P_0\|^2 = \lambda_0 + \alpha$ , which implies the estimate

$$\|P_0 A P_0\| = (\lambda_0 + \alpha) \|P_0 A P_0\|_{H_{0,\alpha}}, \quad (34)$$

which, together with estimate (28), yields (33).

Hence, under the conditions of Lemma 3 and for  $\lambda \in I_0$  the following Hamiltonian is well defined

$$H(\lambda) := P_0 H P_0 + U(\lambda). \quad (35)$$

Note that  $P_0 H P_0 = \lambda_0 P_0$ . Lemma 3 above implies

**Corollary 7.** The operator family  $H(\lambda)$  is (i) self-adjoint for  $\lambda \in I_0$  and (ii) complex analytic in  $\lambda \in I_0^c$ .

In what follows, we label the eigenvalue families  $\nu_i(\lambda)$ ,  $i = 1, \dots, m$ , of  $H(\lambda)$  in the order of their increase and so that

$$\nu_1(\lambda) \leq \dots \leq \nu_m(\lambda). \quad (36)$$

Note that the eigenvalue branches  $\nu_i(\lambda)$  can also be of higher multiplicity. On a subinterval  $I_i \subset I_0$ , we say that the branch  $\nu_i(\lambda)$  is isolated on  $I_i$  if each other branch  $\nu_j(\lambda)$ , with  $\lambda \in I_i$ , either i) coincides with  $\nu_i(\lambda)$  or ii) satisfies

$$\min_{\lambda \in I_i} |\nu_i(\lambda) - \nu_j(\lambda)| \geq \gamma_i > 0. \quad (37)$$




FIGURE 1. (Left) Schematic illustration of the eigenvalues  $\nu_i(\lambda)$  of  $H(\lambda)$  in the neighborhood of  $\lambda_0$  for the case of  $m = 3$ . (Right) Illustration of the spectrum of  $H_0$  consisting of five eigenvalues  $\lambda_{0,1}, \dots, \lambda_{0,5}$  of multiplicity  $m_{0,1} = 1, m_{0,2} = 2, m_{0,3} = 4, m_{0,4} = 2, m_{0,5} = 1$  and the corresponding situation when zooming in close to  $\lambda_0 = \lambda_{0,i}$ .

Further, we have the following result.

**Proposition 5.** *Let  $\alpha \in \mathbb{R}$  be such that  $H_0 + \alpha > 0$  and let  $I_i \subset I_0$  be such that the branch  $\nu_i(\lambda)$  is isolated on  $I_i$ . For  $\lambda \in I_i$ , (i) the eigenvalues  $\nu_i(\lambda)$  of  $H(\lambda)$  are continuously differentiable; (ii) the derivative  $\nu'_i(\lambda)$  is non-positive; (iii) the solutions to the equations  $\nu_i(\lambda) = \lambda$  are unique if  $\lambda \in I_i$ ; (iv) if  $\|W\|_{H_{0,\alpha}} \leq \frac{1}{4}\frac{\gamma_0}{\lambda_\circ}$ , the derivatives  $\nu'_i(\lambda)$ ,  $\lambda \in I'_0 := [\lambda_0 - \frac{1}{4}\gamma_0, \lambda_0 + \frac{1}{4}\gamma_0] \cap I_i$ , are bounded as*

$$|\nu'_i(\lambda)| \leq \frac{8}{\pi} \frac{(\lambda_0 + \alpha)\lambda_\circ}{\gamma_0^2} \|P_0 W P_0^\perp\|_{H_{0,\alpha}}^2.$$

where  $\lambda_\circ := \lambda_0 + \gamma_0 + \alpha$ .

*Proof.* Proof of (i) of a simple eigenvalue  $\lambda_0$ , i.e.,  $m = 1$ . In such a case,  $P_0$  is a rank-one projector on the space spanned by the eigenvector  $\varphi_0$  of  $H_0$  corresponding to the eigenvalue  $\lambda_0$  and therefore Eq. (35) implies that  $H(\lambda) = \nu_1(\lambda)P_0$ , with

$$\nu_1(\lambda) := \langle \varphi_0, H(\lambda)\varphi_0 \rangle. \quad (38)$$

This and Corollary 7 show that the eigenvalue  $\nu_1(\lambda)$  is analytic.

We now prove (i) in the general case. First, we claim the following well-known formula

$$\nu'_i(\lambda) = \langle \chi_i(\lambda), U'(\lambda)\chi_i(\lambda) \rangle, \quad (39)$$

for  $\lambda \in I_i$ , where  $\chi_i(\lambda)$  are well-chosen normalized eigenfunction of  $H(\lambda)$  corresponding to the eigenvalue  $\nu_i(\lambda)$ , namely that they are differentiable in  $\lambda$ . To this end, we observe that for each  $\mu \in I_i$ , we can find a local neighborhood  $I_\mu \subset I_i$  of  $\mu$  such that

$$\bigcup_{j \neq i} \{\nu_j(\lambda) \mid \lambda \in I_\mu\} \cap \{\nu_i(\lambda) \mid \lambda \in I_\mu\} = \emptyset, \quad (40)$$

due to the isolated branch property, i.e.,  $\gamma_i > 0$  in (37). Second, since  $H(\lambda)$  is self-adjoint for  $\lambda \in I_0$  and analytic (say, in the resolvent sense) in  $\lambda \in I_0^c$ , the Riesz projection, corresponding to the eigenvalue  $\nu_i(\lambda)$ :

$$P_i(\lambda) := \frac{1}{2\pi i} \oint_{\Gamma_i(\mu)} (H(\lambda) - z)^{-1} dz, \quad (41)$$

where  $\Gamma_i(\mu)$  is a closed curve in the resolvent set of  $H(\lambda)$  surrounding the eigenvalue branch  $\{\nu_i(\lambda) : \lambda \in I_\mu\}$ , is also self-adjoint and analytic in  $\lambda \in I_\mu^c$  and therefore in  $\lambda \in I_0^c$  (see [25, 20]), condition (40) guarantees that we can choose such a closed curve which contains no other points of  $\sigma(H(\mu))$  on  $I_\mu$ , and that, combining all neighborhoods of  $\mu$  for  $\mu \in I_i$ , there holds that  $P_i(\lambda)$  is analytic in  $I_i$ . From [25, Theorem XII.12], there exists an analytic family of unitary operators  $V_i(\lambda)$  such that  $P_i(\lambda) = V_i(\lambda)P_i(\lambda_0)[V_i(\lambda)]^{-1}$ ,  $\lambda_0$  being possibly replaced by some arbitrary  $\mu \in I_i$  if  $\lambda_0$  does not belong to  $I_i$ . We then define

$$\chi_i(\lambda) = V_i(\lambda)\psi_{0i},$$

where  $\psi_{0i}$  is an eigenvector of  $H_0$  corresponding to the eigenvalue  $\lambda_0$ . Since  $V_i(\lambda)$  is analytic,  $\chi_i(\lambda)$  is also analytic in  $I_i$ , so in particular differentiable, and one can easily check that  $\chi_i(\lambda)$  is of norm 1 and that  $P_i(\lambda)\chi_i(\lambda) = \chi_i(\lambda)$ , which guarantees that  $\chi_i(\lambda)$  is a normalized eigenfunction of  $H(\lambda)$ . Now, we use that

$$\begin{aligned} \langle \chi'_i(\lambda), H(\lambda)\chi_i(\lambda) \rangle + \langle \chi_i(\lambda), H(\lambda)\chi'_i(\lambda) \rangle &= \nu_i(\lambda)(\langle \chi'_i(\lambda), \chi_i(\lambda) \rangle + \langle \chi_i(\lambda), \chi'_i(\lambda) \rangle) \\ &= \langle \chi_i(\lambda), \chi_i(\lambda) \rangle' = 0 \end{aligned}$$

to obtain  $\nu'_i(\lambda) = \langle \chi_i(\lambda), H'(\lambda)\chi_i(\lambda) \rangle$ , which gives (39). The differentiability of  $\chi_i(\lambda)$  and the analyticity of  $H(\lambda)$  then implies the differentiability of  $\nu_i$  in each neighborhood of  $\lambda$ .

In order to prove (ii), note that  $U'(\lambda) \leq 0$ , as follows by the explicit formula

$$U'(\lambda) := -PWP_0^\perp(H^\perp - \lambda)^{-2}P_0^\perp WP \leq 0. \quad (42)$$

Hence,  $\nu'_i(\lambda) < 0$  by (39). The monotonicity of  $\nu_i(\lambda)$  also implies the well-posedness of the equations  $\nu_i(\lambda) = \lambda$  under the condition that  $\lambda \in I_i$ , thus statement (iii).

We now aim to prove (iv). Starting from (39), we estimate  $\nu'_i(\lambda)$  with

$$|\nu'_i(\lambda)| \leq \|(H_0 + \alpha)^{1/2}P_0\|^2 \|U'(\lambda)\|_{H_0, \alpha}. \quad (43)$$

The first factor on the right hand side is exactly known as

$$\|(H_0 + \alpha)^{1/2}P_0\|^2 = (\lambda_0 + \alpha). \quad (44)$$

To investigate the second factor on the r.h.s. of (43), we use the analyticity  $U(\lambda)$  and the estimate (28). Indeed, by the Cauchy integral formula, we have

$$\|U'(\lambda)\|_{H_0,\alpha} \leq \frac{1}{2\pi R} \sup_{\substack{\mu \in \mathbb{C}, \\ |\mu - \lambda| = R}} \|U(\mu)\|_{H_0,\alpha},$$

where  $R$  is such that  $\{\mu \in \mathbb{C} : |\mu - \lambda| \leq R\} \subset I_0^c$ . Taking  $R = \frac{1}{4}\gamma_0$  gives, under the conditions of Lemma 3, the estimate

$$\|U'(\lambda)\|_{H_0,\alpha} \leq \frac{8}{\pi} \frac{\lambda_\circ}{\gamma_0^2} \|P_0 W P_0^\perp\|_{H_0,\alpha}^2. \quad (45)$$

Combining equations (43), (44) and (45) shows (iv).  $\square$

**Corollary 8.** *Let  $\alpha \in \mathbb{R}$  be such that  $H_0 + \alpha > 0$  and let  $I_i \subset I_0$  be such that the branch  $\nu_i(\lambda)$  is isolated on  $I_i$ . Under the condition that*

$$\frac{8}{\pi} \frac{(\lambda_0 + \alpha)\lambda_\circ}{\gamma_0^2} \|P_0 W P_0^\perp\|_{H_0,\alpha}^2 < 1,$$

*and that the unique solution  $\lambda$  of  $\nu_i(\lambda) = \lambda$  satisfies  $\lambda \in I'_0$ , the fixed-point iteration  $\lambda^{(k+1)} = \nu_i(\lambda^{(k)})$  converges to  $\lambda$  for initial values in  $I'_0$ .*

Now, we proceed directly to the proof of Theorem 6.

*Proof of Theorem 6.* For the estimate on the eigenvalues, we first remark that applying Proposition 4 to the  $m$  eigenvalues corresponding to  $\lambda_0$  for  $H_0$  provides the first inequality in (24). The second one follows immediately from the condition  $\|W\|_{H_0,\alpha} \leq \frac{1}{2} \frac{\gamma_0}{\lambda_0 + \alpha}$  and thus  $\mu_i \in I_0$ .

The fact that the operator  $H$  has exactly  $m$  eigenvalues (counting the multiplicities) in  $I_0$  follows from Corollary 4 and Proposition 5(iii) and the fact that  $H(\lambda)$  is a  $m \times m$  symmetric matrix.

For the estimates on the eigenfunctions, recall from Theorem 3 that  $Q_0(\mu_i)\psi_{0i} = \psi_i$ , where  $\mu_i = \nu_i(H)$  and the operator  $Q_0(\lambda)$  is given by

$$Q_0(\lambda) := \text{Id} - R^\perp(\lambda)P_0^\perp W P_0, \quad (46)$$

with  $R^\perp(\lambda)$  defined in (30). This yields

$$\psi_{0i} - \psi_i = \psi_{0i} - Q_0(\mu_i)\psi_{0i} = R^\perp(\mu_i)P_0^\perp W P_0 \psi_{0i}. \quad (47)$$

Then, for  $\gamma \in \{0, 1/2\}$ ,

$$\begin{aligned} \|H_{0,\alpha}^\gamma(\psi_{0i} - \psi_i)\| &\leq \|H_{0,\alpha}^\gamma R^\perp(\mu_i)P_0^\perp W P_0\| \\ &\leq \|H_{0,\alpha}^\gamma H_{0,\alpha}^{-1/2} P_0^\perp\| \|H_{0,\alpha}^{1/2} R^\perp(\mu_i)H_{0,\alpha}^{1/2}\| \|P_0^\perp W P_0\|_{H_0,\alpha} \|P_0 H_{0,\alpha}^{1/2}\|. \end{aligned}$$

In the previous expression, we can use (31) to estimate  $\|H_{0,\alpha}^{1/2} R^\perp(\mu_i)H_{0,\alpha}^{1/2}\|$ . Then, we note that

$$\|P_0^\perp W P_0\|_{H_0,\alpha} \leq \|W\|_{H_0,\alpha},$$

as well as

$$\|P_0 H_{0,\alpha}^{1/2}\| = (\lambda_0 + \alpha)^{1/2}.$$

Finally, in the case  $\gamma = 0$

$$\|H_{0,\alpha}^{-1/2} P_0^\perp\| \leq (\lambda_0 + \alpha)^{-1/2},$$

and for  $\gamma = 1/2$ ,

$$\|H_{0,\alpha}^\gamma H_{0,\alpha}^{-1/2} P_0^\perp\| \leq 1.$$

Combining the four bounds leads to (25) and (26).  $\square$

**Remark 9.** Note that by Theorem 3, any solution  $\mu_i$  to the equation  $\nu_i(\mu_i) = \mu_i$ , for  $i = 1, \dots, m$  and where the  $\mu_i$  are in ascending order, is an eigenvalue of  $H$ . Under the condition

$$\|W\|_{H_0,\alpha} \leq \frac{1}{2} \frac{\gamma_0}{\lambda_0 + \alpha}, \quad (48)$$

Theorem 6 guarantees that the eigenvalues  $\mu_i$  satisfy  $|\mu_i - \lambda_0| \leq \frac{\gamma_0}{2}$  and thus  $\mu_i \in I_0$ .

On the contrary, the eigenvalues  $\mu_i$  are the only  $m$  eigenvalues of  $H$  belonging to  $I_0$  if a similar condition as (48), but for the next larger eigenvalue of  $H_0$  than  $\lambda_0$  holds whereas such a condition is automatically satisfied by (48) for the preceding eigenvalue of  $H_0$ .

#### 4. PRELIMINARY RESULTS

We now derive a few preliminary results that will be useful for proving Theorem 1. For the following proofs, we define the following quantities:  $h_\lambda := -\Delta - \lambda$ ,  $V_M^\perp = P_M^\perp V P_M^\perp$ ,  $V_M^N = P_M^N V P_M^N$ .

**Lemma 4.** For  $\lambda \in \mathbb{C}$  with  $\operatorname{Re} \lambda < \frac{1}{2}\rho_M$  and  $\rho_M \geq 1$ , the following bounds hold

$$\|h_\lambda^{-1/2} V_M^N h_\lambda^{-1/2}\| \leq \|h_\lambda^{-1/2} V_M^\perp h_\lambda^{-1/2}\| \leq 4\rho_M^{-r} \|V\|_r. \quad (49)$$

*Proof.* First, we note that

$$\|h_\lambda^{-1/2} V_M^N h_\lambda^{-1/2}\| = \|P_N h_\lambda^{-1/2} V_M^\perp h_\lambda^{-1/2} P_N\| \leq \|h_\lambda^{-1/2} V_M^\perp h_\lambda^{-1/2}\|.$$

Then, we estimate for any  $s \geq 0$ , using the assumption  $\operatorname{Re} \lambda \leq \frac{1}{2}\rho_M$

$$\|h_{-1}^s h_\lambda^{-s} P_M^\perp\| \leq \frac{|\rho_M + 1|^s}{|\rho_M - \lambda|^s} \leq 2^s (1 + \rho_M^{-1})^s \leq 4^s, \quad (50)$$

This implies in particular that  $\|h_\lambda^{-1/2} V_M^\perp h_\lambda^{-1/2}\| \leq 4 \|V_M^\perp\|_0$ . The result follows noting that  $\|V_M^\perp\|_0 \leq \rho_M^{-r} \|V\|_r$ .  $\square$

**Lemma 5.** *For  $\lambda \in \mathbb{C}$  with  $\operatorname{Re} \lambda \leq \min(\frac{1}{2}\rho_M, \kappa_M)$  and  $\rho_M \geq 1$ , the following bound holds*

$$\|U_\sigma(\lambda)\|_r \leq 4\rho_M^{-r} \|V\|_r^2 \sum_{k=0}^K [4\rho_M^{-r} \|V\|_r]^k. \quad (51)$$

Moreover, if  $4\rho_M^{-r} \|V\|_r < 1$ ,

$$\|U_\sigma(\lambda)\|_r \leq \rho_M^{-r} \frac{4 \|V\|_r^2}{1 - 4\rho_M^{-r} \|V\|_r}, \quad (52)$$

and in particular

$$\|U_M(\lambda)\|_r \leq \rho_M^{-r} \frac{4 \|V\|_r^2}{1 - 4\rho_M^{-r} \|V\|_r}. \quad (53)$$

*Proof.* By definition (9), we can write  $U_\sigma(\lambda)$  which is well-defined for  $\lambda < \kappa_M$  as

$$U_\sigma(\lambda) = - \sum_{k=0}^K \mathsf{P}_M V \mathsf{P}_M^N h_\lambda^{-1/2} \left[ -h_\lambda^{-1/2} V_M^N h_\lambda^{-1/2} \right]^k h_\lambda^{-1/2} \mathsf{P}_M^N V \mathsf{P}_M. \quad (54)$$

Using estimate (49), there holds

$$\|U_\sigma(\lambda)\|_r \leq \sum_{k=0}^K [4\rho_M^{-r} \|V\|_r]^k \|h_{-1}^{-1/2+r/2} \mathsf{P}_M V \mathsf{P}_M^N h_\lambda^{-1/2}\| \|h_\lambda^{-1/2} \mathsf{P}_M^N V \mathsf{P}_M h_{-1}^{-1/2+r/2}\|.$$

and by (50),  $\|h_\lambda^{-1/2} \mathsf{P}_M^N V \mathsf{P}_M h_{-1}^{-1/2+r/2}\| \leq 2\rho_M^{-r/2} \|V\|_r$ , so that we obtain (51). The bound (52) is easily obtained from (51) and taking  $K, N = \infty$  in (52), we arrive at (53).  $\square$

**Lemma 6.** *For  $\lambda < \frac{1}{2}\kappa_M$ ,  $\rho_M \geq 1$  and if  $4\rho_M^{-r} \|V\|_r < 1$ , the following bounds hold*

$$\|U'_M(\lambda)\|_r \leq \frac{1}{\pi(\kappa_M - 2\lambda)} \rho_M^{-r} \frac{4 \|V\|_r^2}{1 - 4\rho_M^{-r} \|V\|_r}. \quad (55)$$

*Proof.* Since from Proposition 2(iv),  $U_M(\lambda)$  is complex analytic in  $\lambda$  for  $\operatorname{Re} \lambda < \kappa_M$ , by the Cauchy integral formula, we have

$$\|U'_M(\lambda)\|_r \leq \frac{1}{2\pi R_M(\lambda)} \sup_{\substack{\mu \in \mathbb{C}, \\ |\mu - \lambda| = R_M(\lambda)}} \|U_M(\mu)\|_r,$$

with  $R_M(\lambda) = \frac{1}{2}\kappa_M - \lambda > 0$ . Using (53) and noting that  $\frac{1}{2}\kappa_M \leq \frac{1}{2}\rho_M$ , we obtain (55).  $\square$

**Lemma 7.** For  $\lambda < \min(\kappa_M, \frac{1}{2}\rho_M)$ ,  $\rho_N \geq \rho_M > 1$  and if  $4\rho_M^{-r}\|V\|_r < 1$  and  $4\rho_N^{-r}\|V\|_r + \frac{16\rho_M^{-2r}\|V\|_r^2}{1-4\rho_M^{-r}\|V\|_r} < 1$ , the following bound holds

$$\begin{aligned} & \|U_M(\lambda) - U_\sigma(\lambda)\|_r \\ & \leq \frac{4\rho_N^{-r}\|V\|_r^2}{1-4\rho_N^{-r}\|V\|_r - \frac{16\rho_M^{-2r}\|V\|_r^2}{1-4\rho_M^{-r}\|V\|_r}} \left[ 1 + \frac{4\rho_M^{-r}\|V\|_r}{1-4\rho_M^{-r}\|V\|_r} \right]^2 + \frac{4\rho_M^{-r}\|V\|_r^2}{1-4\rho_M^{-r}\|V\|_r} [4\rho_M^{-r}\|V\|_r]^{K+1}. \end{aligned} \quad (56)$$

*Proof.* We first write  $U_M(\lambda) - U_\sigma(\lambda)$  as

$$U_M(\lambda) - U_\sigma(\lambda) = (U_M(\lambda) - U_{MN}(\lambda)) + (U_{MN}(\lambda) - U_\sigma(\lambda)), \quad (57)$$

where, using the notation  $\mathcal{H}_M^N = \mathbf{P}_M^N \mathcal{H} \mathbf{P}_M^N$

$$\begin{aligned} U_{MN}(\lambda) & := -\mathbf{P}_M V \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V \mathbf{P}_M \\ & = -\mathbf{P}_M V \mathbf{P}_M^N (-\Delta + V_M^N - \lambda)^{-1} \mathbf{P}_M^N V \mathbf{P}_M. \end{aligned}$$

Since  $U_M(\lambda) = -\mathbf{P}_M V \mathbf{P}_M^\perp (\mathcal{H}_M^\perp - \lambda)^{-1} \mathbf{P}_M^\perp V \mathbf{P}_M$ , with  $\mathcal{H}_M^\perp = \mathbf{P}_M^\perp \mathcal{H} \mathbf{P}_M^\perp$ , the first term  $U_M(\lambda) - U_{MN}(\lambda)$  is estimated as follows.

Denoting by  $\mathcal{H}_N^\perp = \mathbf{P}_N^\perp \mathcal{H} \mathbf{P}_N^\perp$ , and the Schur complement

$$A = \mathbf{P}_N^\perp ((\mathcal{H}_N^\perp - \lambda \mathbf{P}_N^\perp) - \mathbf{P}_N^\perp V \mathbf{P}_M^N (\mathcal{H}_M^\perp - \lambda)^{-1} \mathbf{P}_M^N V \mathbf{P}_N^\perp)^{-1} \mathbf{P}_N^\perp,$$

there holds, using a block matrix inversion

$$\begin{aligned} \mathbf{P}_M^\perp (\mathcal{H}_M^\perp - \lambda)^{-1} \mathbf{P}_M^\perp & = \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N + \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V A V \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N \\ & \quad - \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V A - A V \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N + A. \end{aligned} \quad (58)$$

Therefore,  $U_M(\lambda) - U_{MN}(\lambda)$  can be decomposed into four terms as

$$\begin{aligned} U_M(\lambda) - U_{MN}(\lambda) & = -\mathbf{P}_M V \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V A V \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V \mathbf{P}_M, \\ & \quad + \mathbf{P}_M V \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V A V \mathbf{P}_M, \\ & \quad + \mathbf{P}_M V A V \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V \mathbf{P}_M, \\ & \quad - \mathbf{P}_M V A V \mathbf{P}_M. \end{aligned}$$

Then, the  $r$ -norm can be estimated as

$$\begin{aligned} \|U_M(\lambda) - U_{MN}(\lambda)\|_r & \leq \|V\|_r^2 \left[ \|h_{-1}^{1/2-r/2} \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V A V \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N h_{-1}^{1/2-r/2}\| \right. \\ & \quad + \|h_{-1}^{1/2-r/2} \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V A h_{-1}^{1/2-r/2}\| \\ & \quad + \|h_{-1}^{1/2-r/2} A V \mathbf{P}_M^N (\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N h_{-1}^{1/2-r/2}\| \\ & \quad \left. + \|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\| \right]. \end{aligned}$$

Introducing appropriate  $h_{-1}^{1/2-r/2}$  and  $h_{-1}^{-1/2+r/2}$  terms, we obtain

$$\begin{aligned} \|U_M(\lambda) - U_{MN}(\lambda)\|_r &\leq \|V\|_r^2 \left[ \|h_{-1}^{1/2-r/2} P_M^N (\mathcal{H}_M^N - \lambda)^{-1} P_M^N h_{-1}^{1/2-r/2}\|^2 \|V\|_r^2 \|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\| \right. \\ &\quad + 2 \|h_{-1}^{1/2-r/2} P_M^N (\mathcal{H}_M^N - \lambda)^{-1} P_M^N h_{-1}^{1/2-r/2}\| \|V\|_r \|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\| \\ &\quad \left. + \|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\| \right] \\ &\leq \|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\| \|V\|_r^2 \\ &\quad \times \left[ 1 + \|h_{-1}^{1/2-r/2} P_M^N (\mathcal{H}_M^N - \lambda)^{-1} P_M^N h_{-1}^{1/2-r/2}\| \|V\|_r \right]^2. \end{aligned}$$

We are therefore left with the estimation of  $\|h_{-1}^{1/2-r/2} P_M^N (\mathcal{H}_M^N - \lambda)^{-1} P_M^N h_{-1}^{1/2-r/2}\|$  and  $\|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\|$ . First, noting from (49) that

$$\|h_\lambda^{1/2} P_M^N (\mathcal{H}_M^\perp - \lambda)^{-1} P_M^N h_\lambda^{1/2}\| \leq \|(I + h_\lambda^{-1/2} V_M^\perp h_\lambda^{-1/2})^{-1}\| \leq \frac{1}{1 - 4\rho_M^{-r} \|V\|_r}, \quad (59)$$

and using that

$$\|h_{-1}^{1/2-r/2} h_\lambda^{-1/2} P_M^N\| \leq 2\rho_M^{-r/2}, \quad (60)$$

we obtain

$$\|h_{-1}^{1/2-r/2} P_M^N (\mathcal{H}_M^N - \lambda)^{-1} P_M^N h_{-1}^{1/2-r/2}\| \leq \frac{4\rho_M^{-r}}{1 - 4\rho_M^{-r} \|V\|_r}. \quad (61)$$

Second,

$$\|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\| = \|h_{-1}^{1/2-r/2} P_N^\perp [(\mathcal{H}_N^\perp - \lambda P_N^\perp) - P_N^\perp V P_M^N (\mathcal{H}_M^\perp - \lambda)^{-1} P_M^N V P_N^\perp]^{-1} P_N^\perp h_{-1}^{1/2-r/2}\|.$$

Noting that

$$\|h_{-1}^{1/2-r/2} h_\lambda^{-1/2} P_N^\perp\| \leq 2\rho_N^{-r/2},$$

there holds

$$\|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\| \leq 4\rho_N^{-r} \|h_\lambda^{1/2} P_N^\perp [(\mathcal{H}_N^\perp - \lambda P_N^\perp) - P_N^\perp V P_M^N (\mathcal{H}_M^\perp - \lambda)^{-1} P_M^N V P_N^\perp]^{-1} P_N^\perp h_\lambda^{1/2}\|.$$

Factorizing  $h_\lambda$ , we deduce

$$\|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\| \leq 4\rho_N^{-r} \|[I + h_\lambda^{-1/2} V_N^\perp h_\lambda^{-1/2} - h_\lambda^{-1/2} P_N^\perp V P_M^N (\mathcal{H}_M^\perp - \lambda)^{-1} P_M^N V P_N^\perp h_\lambda^{-1/2}]^{-1}\|.$$

From (49) with  $N$  in place of  $M$ , we obtain

$$\|h_\lambda^{-1/2} V_N^\perp h_\lambda^{-1/2}\| \leq 4\rho_N^{-r} \|V\|_r. \quad (62)$$

Moreover,

$$\|h_\lambda^{-1/2} P_N^\perp V P_M^N (\mathcal{H}_M^\perp - \lambda)^{-1} P_M^N V P_N^\perp h_\lambda^{-1/2}\| \leq \|h_\lambda^{-1/2} P_N^\perp V P_M^N h_\lambda^{-1/2}\|^2 \|h_\lambda^{1/2} P_M^N (\mathcal{H}_M^\perp - \lambda)^{-1} P_M^N h_\lambda^{1/2}\|,$$

which, from (49) and (59), leads to

$$\|h_\lambda^{-1/2} P_N^\perp V P_M^N (\mathcal{H}_M^\perp - \lambda)^{-1} P_M^N V P_N^\perp h_\lambda^{-1/2}\| \leq \frac{16\rho_M^{-2r} \|V\|_r^2}{1 - 4\rho_M^{-r} \|V\|_r}.$$

Combining this last line with (62), we obtain the bound

$$\|h_{-1}^{1/2-r/2} A h_{-1}^{1/2-r/2}\| \leq 4\rho_N^{-r} \frac{1}{1 - 4\rho_N^{-r} \|V\|_r - \frac{16\rho_M^{-2r} \|V\|_r^2}{1 - 4\rho_M^{-r} \|V\|_r}}.$$

This leads to the following bound for the difference  $U_M(\lambda) - U_{MN}(\lambda)$

$$\|U_M(\lambda) - U_{MN}(\lambda)\|_r \leq \rho_N^{-r} \frac{4\|V\|_r^2}{1 - 4\rho_N^{-r} \|V\|_r - \frac{16\rho_M^{-2r} \|V\|_r^2}{1 - 4\rho_M^{-r} \|V\|_r}} \left[ 1 + \frac{4\rho_M^{-r} \|V\|_r}{1 - 4\rho_M^{-r} \|V\|_r} \right]^2. \quad (63)$$

For the second term on the right handside of (57), we write  $U_{MN}(\lambda) - U_\sigma(\lambda)$  as

$$U_{MN}(\lambda) - U_\sigma(\lambda) = - \sum_{k=K+1}^{\infty} \mathsf{P}_M V \mathsf{P}_M^N h_\lambda^{-1/2} \left[ -h_\lambda^{-1/2} V_M^N h_\lambda^{-1/2} \right]^k h_\lambda^{-1/2} \mathsf{P}_M^N V \mathsf{P}_M. \quad (64)$$

Using (49) and (50), we obtain

$$\begin{aligned} \|U_{MN}(\lambda) - U_\sigma(\lambda)\|_r &\leq \|V\|_r^2 \|h_{-1}^{1/2-r/2} h_\lambda^{-1/2} P_M^\perp\|^2 \sum_{k=K+1}^{\infty} \|h_\lambda^{-1/2} V_M^N h_\lambda^{-1/2}\|^k \\ &\leq 4\|V\|_r^2 \rho_M^{-r} \sum_{k=K+1}^{\infty} [4\rho_M^{-r} \|V\|_r]^k, \end{aligned}$$

from which we deduce that

$$\|U_{MN}(\lambda) - U_\sigma(\lambda)\|_r \leq \frac{4\|V\|_r^2}{1 - 4\rho_M^{-r} \|V\|_r} \rho_M^{-r} [4\rho_M^{-r} \|V\|_r]^{K+1}. \quad (65)$$

Combining (63) and (65), we obtain (56).  $\square$

**Lemma 8.** *For  $\lambda < \mu < \min(\kappa_M, \frac{1}{2}\rho_M)$ ,  $\rho_M \geq 1$  and if  $4\rho_M^{-r} \|V\|_r < 1$ , there holds*

$$\|U_M(\mu) - U_M(\lambda)\|_r \leq |\mu - \lambda| \frac{\rho_M^{-r}}{\pi(\kappa_M - 2\lambda)} \frac{4\|V\|_r^2}{1 - 4\rho_M^{-r} \|V\|_r} \quad (66)$$

*Proof of Lemma 8.* Without loss of generality, we assume that  $\lambda < \mu$ . Writing

$$U_M(\mu) - U_M(\lambda) = \int_\lambda^\mu U'_M(s) ds,$$

yields

$$\|U_M(\mu) - U_M(\lambda)\|_r \leq |\mu - \lambda| \max_{s \in [\lambda, \mu]} \|U'_M(s)\|_r.$$

We conclude by applying (55) of Lemma 6.  $\square$

## 5. PROOF OF THE MAIN RESULTS

The goal of this section is to provide the proof for Theorem 1.

We first prove the following technical lemmas which will be useful later.

**Lemma 9.** *For  $H, W$  such that  $H = -\Delta + W$ , there holds for  $\alpha \geq (2\|W\|_r)^{1/r}$  such that  $H + \alpha > 0$*

$$\|A\|_{H,\alpha} \leq 2\|A\|_{-\Delta,\alpha} \leq 2\alpha^{-r}\|A\|_r.$$

*Proof.* First, denoting again  $h_{-\alpha} := -\Delta + \alpha$ , note that

$$\|A\|_{H,\alpha} \leq \|A\|_{-\Delta,\alpha} \|h_{-\alpha}^{1/2}(H + \alpha)^{-1/2}\|^2.$$

Let  $v$  be arbitrary and define  $u = (H + \alpha)^{-1/2}v$ . Note that

$$\|h_{-\alpha}^{1/2}u\|^2 = \langle u, h_{-\alpha}u \rangle = \langle u, (H + \alpha - W)u \rangle = \|v\|^2 - \langle u, Wu \rangle.$$

Further, there holds

$$\langle u, Wu \rangle \leq \|W\|_{-\Delta,\alpha} \|h_{-\alpha}^{1/2}u\|^2,$$

and using the inequality  $\|W\|_{-\Delta,\alpha} \leq \alpha^{-r}\|W\|_r$  we obtain that  $\|W\|_{-\Delta,\alpha} \leq \frac{1}{2}$  for all  $\alpha \geq (2\|W\|_r)^{1/r}$ , and that  $\|h_{-\alpha}^{1/2}(H + \alpha)^{-1/2}\|^2 \leq 2$  yielding the first inequality.

The second inequality results from using, once again, the inequality  $\|A\|_{-\Delta,\alpha} \leq \alpha^{-r}\|A\|_r$ .  $\square$

Before starting the proof of Theorem 1, we analyze the relation of the spectra of  $\mathcal{H}$  and  $\mathcal{H}_M(\lambda)$ .

**Lemma 10.** *Let  $\lambda_i < \lambda_j$  be the  $i$ -th resp.  $j$ -the eigenvalue of  $\mathcal{H}$  and let  $\nu_{Mi}(\lambda)$  denote the  $i$ -th eigenvalue of the operator  $\mathcal{H}_M(\lambda)$ . Then, we have the following  $M$ -independent lower bound*

$$\lambda_j - \lambda_i \leq \min(\nu_{Mj}(\lambda_i) - \nu_{Mi}(\lambda_i), \nu_{Mj}(\lambda_j) - \nu_{Mi}(\lambda_j)). \quad (67)$$

*Proof.* Since  $\lambda_i = \nu_{Mi}(\lambda_i)$  and  $\lambda_j = \nu_{Mj}(\lambda_j)$ , we have  $\lambda_j - \lambda_i = \nu_{Mj}(\lambda_j) - \nu_{Mi}(\lambda_i)$ . By definition (17) and Proposition 2, the family  $\mathcal{H}_M(\lambda)$  is monotonically decreasing with  $\lambda$ . Hence  $\nu_{Mi}(\lambda_j) \leq \nu_N^i(\lambda_i)$  and  $\nu_{Mj}(\lambda_j) \leq \nu_{Mj}(\lambda_i)$ , so that  $\lambda_j - \lambda_i = \nu_{Mj}(\lambda_j) - \nu_{Mi}(\lambda_i)$  implies (67).  $\square$

**Corollary 9.** *Let  $\lambda_i$  be an isolated eigenvalue of  $H$ . Then, the gap  $\gamma_{Mi}$  of  $\lambda_i$  to the rest of the spectrum of  $\mathcal{H}_M(\lambda_i)$  is bounded below by the gap  $\gamma_0$  of  $\lambda_i$  to the rest of the spectrum of  $H$ .*

**Lemma 11.** *Let be  $\alpha > 0$  such that*

$$\mathcal{H}_M(\lambda_\star) + \alpha > 0, \quad \text{and} \quad \alpha > (2\|U_M(\lambda_\star) - U_\sigma(\lambda_{\sigma i})\|_r)^{1/r}, \quad (68)$$

*M large enough such that  $\lambda_\star, \lambda_{\sigma i} < \min(\kappa_M, \frac{1}{2}\rho_M)$ ,  $\rho_M \geq 1$  and  $4\rho_M^{-r}\|V\|_r \leq \frac{1}{3}$ , and  $N \geq M$ .*

Then, there holds

$$\|U_M(\lambda_\star) - U_\sigma(\lambda_{\sigma i})\|_{\mathcal{H}_M(\lambda_\star), \alpha} \leq |\lambda_{\sigma i} - \lambda_\star| \frac{8\alpha^{-r}\rho_M^{-r}\|V\|_r^2}{\pi(\kappa_M - 2\min(\lambda_\star, \lambda_{\sigma i}))} + 36 \frac{\|V\|_r^2}{\alpha^r} \varepsilon(\sigma, r, V) \quad (69)$$

with, using the notation  $\sigma = (N, M, K)$ ,

$$\varepsilon(\sigma, r, V) := \rho_N^{-r} + \rho_M^{-r} [4\rho_M^{-r}\|V\|_r]^{K+1}.$$

*Proof.* Applying Lemma 9 to the present case, we obtain

$$\|U_M(\lambda_\star) - U_\sigma(\lambda_{\sigma i})\|_{\mathcal{H}_M(\lambda_\star), \alpha} \leq 2\alpha^{-r}\|U_M(\lambda_\star) - U_\sigma(\lambda_{\sigma i})\|_r.$$

Employing the triangle inequality, there holds

$$\|U_M(\lambda_\star) - U_\sigma(\lambda_{\sigma i})\|_r \leq \|U_M(\lambda_\star) - U_M(\lambda_{\sigma i})\|_r + \|U_M(\lambda_{\sigma i}) - U_\sigma(\lambda_{\sigma i})\|_r. \quad (70)$$

For the first term of the right handside, Lemma 8 yields

$$\|U_M(\lambda_{\sigma i}) - U_M(\lambda_\star)\|_r \leq |\lambda_{\sigma i} - \lambda_\star| \frac{8\rho_M^{-r}\|V\|_r^2}{\pi(\kappa_M - 2\min(\lambda_\star, \lambda_{\sigma i}))}.$$

Finally, we use Lemma 7 in order to bound the second term of (70), we conclude that

$$\|U_M(\lambda_{\sigma i}) - U_\sigma(\lambda_{\sigma i})\|_r \leq \|V\|_r^2 \left[ 18\rho_N^{-r} + 6\rho_M^{-r} [4\rho_M^{-r}\|V\|_r]^{K+1} \right].$$

Combining all estimates concludes the proof.  $\square$

Finally, we are now ready to prove Theorem 1.

*Proof of Theorem 1.* Note that  $\lambda_\star = \nu_{Mi}(\lambda_\star)$ ,  $\lambda_{\sigma i} = \nu_{\sigma i}(\lambda_{\sigma i})$  where  $\nu_{Mi}(\lambda)$  and  $\nu_{\sigma i}(\lambda)$  represents the  $i$ -th eigenvalue of  $\mathcal{H}_M(\lambda)$  and  $\mathcal{H}_\sigma(\lambda)$  respectively.

We choose  $\alpha > 0$  in order to satisfy the condition (68) of Lemma 11. By the expression (69) of Lemma 11 we can choose  $\widetilde{M}_0 \in \mathbb{N}$  such that

$$\|U_M(\lambda_\star) - U_\sigma(\lambda_{\sigma i})\|_{\mathcal{H}_M(\lambda_\star), \alpha} \leq \frac{1}{4} \frac{\gamma_0}{\lambda_\circ},$$

for any  $M \geq \widetilde{M}_0$ , and where  $\gamma_0$  denotes the gap of  $\lambda_\star$  to the rest of the spectrum of  $\mathcal{H}$ , which is a lower bound of  $\gamma_{Mi}$  by Corollary 9, and  $\lambda_\circ = \lambda_\star + \gamma_0 + \alpha$ . In consequence, we can apply Theorem 6 with

$$H_0 = \mathcal{H}_M(\lambda_\star), \quad W = U_\sigma(\lambda_{\sigma i}) - U_M(\lambda_\star), \quad \lambda_0 = \lambda_\star, \quad (71)$$

and thus  $H = \mathcal{H}_\sigma(\lambda_{\sigma i})$ , yielding

$$|\lambda_\star - \lambda_{\sigma i}| = |\nu_{Mi}(\lambda_\star) - \nu_{\sigma i}(\lambda_{\sigma i})| \leq (\lambda_\star + \alpha) \|U_M(\lambda_\star) - U_\sigma(\lambda_{\sigma i})\|_{\mathcal{H}_M(\lambda_\star), \alpha} \quad (72)$$

$$\leq (\lambda_\star + \alpha) \left[ |\lambda_{\sigma i} - \lambda_\star| \frac{16\alpha^{-r}\rho_M^{-r}\|V\|_r^2}{\pi(\kappa_M - 2\min(\lambda_\star, \lambda_{\sigma i}))} + 54 \frac{\|V\|_r^2}{\alpha^r} \varepsilon(\sigma, r, V) \right] \quad (73)$$

It exists a  $M_0 \in \mathbb{N}$ , with  $M_0 \geq \widetilde{M}_0$ , such that

$$4\rho_M^{-r}\|V\|_r \leq \frac{1}{3} \quad \text{and} \quad (\lambda_\star + \alpha) \frac{16\alpha^{-r}\rho_M^{-r}\|V\|_r^2}{\pi(\kappa_M - 2\min(\lambda_\star, \lambda_{\sigma i}))} \leq \frac{1}{2}, \quad (74)$$

for all  $M \geq M_0$  and thus

$$|\lambda_\star - \lambda_{\sigma i}| \lesssim (\lambda_\star + \alpha) \frac{\|V\|_r^2}{\alpha^r} \varepsilon(\sigma, r, V). \quad (75)$$

A similar development for the eigenfunctions based on the estimates (25)–(26) of Theorem 6 can be applied. Indeed, we denote by  $\varphi_{\sigma i}$  the  $i$ -th normalized eigenfunction of  $\mathcal{H}_\sigma(\lambda_{\sigma i})$ , thus, in terms of the notation used in Theorem 6, we have  $\psi_i = \varphi_{\sigma i}$ . Then, the corresponding eigenfunction in the span of all eigenfunctions corresponding to  $\lambda_\star$  is given by  $\varphi_{Mi} = \mathsf{P}_0 \varphi_{\sigma i}$ , where  $\mathsf{P}_0$  is the projector onto this span of all eigenfunctions of  $\mathcal{H}_M(\lambda_\star)$  corresponding to  $\lambda_\star$ . Thus,  $\varphi_{Mi}$  is an eigenfunction of  $\mathcal{H}_M(\lambda_\star)$  associated to the eigenvalue  $\lambda_\star$ . Taking, again, Corollary 9 into account yields

$$\|\varphi_{Mi} - \varphi_{\sigma i}\| \leq 4 \frac{\lambda_\circ}{\gamma_0} \|U_M(\lambda_\star) - U_\sigma(\lambda_{\sigma i})\|_{\mathcal{H}_M(\lambda_\star), \alpha}, \quad (76)$$

Using the bounds of  $\|U_M(\lambda_\star) - U_\sigma(\lambda_{\sigma i})\|_{\mathcal{H}_M(\lambda_\star), \alpha}$  expression (69) of Lemma 11 and combining with (75) yields the auxiliary result (see (14)): for  $s \geq 0$  and for all  $M \geq M_0$ ,

$$\|(-\Delta + \text{Id})^s(\varphi_{Mi} - \varphi_{\sigma i})\| \lesssim \rho_M^s \|\varphi_{Mi} - \varphi_{\sigma i}\| \lesssim \frac{\lambda_\circ}{\gamma_0} \frac{\|V\|_r^2}{\alpha^r} \rho_M^s \varepsilon(\sigma, r, V). \quad (77)$$

Now we define the eigenfunction  $\varphi_i := \mathsf{Q}_M(\lambda_\star)\varphi_{Mi}$  where  $\mathsf{Q}_M$  is defined by (18). Following Corollary 5,  $\varphi_i$  is an eigenfunction of  $\mathcal{H}$  associated to the eigenvalue  $\lambda_\star$ . Note that

$$\varphi_i - Q_\sigma(\lambda_{\sigma i})\varphi_{\sigma i} = \varphi_{Mi} - \varphi_{\sigma i} - [S_M(\lambda_\star)\varphi_{Mi} - S_\sigma(\lambda_{\sigma i})\varphi_{\sigma i}],$$

with  $S_M(\lambda) := \mathsf{P}_M^\perp (\mathcal{H}_M^\perp - \lambda)^{-1} \mathsf{P}_M^\perp V \mathsf{P}_M$ ,  $S_\sigma(\lambda) := R_\sigma(\lambda) \mathsf{P}_M^N V \mathsf{P}_M$ . Applying the triangle inequality several times yields

$$\|\varphi_i - Q_\sigma(\lambda_{\sigma i})\varphi_{\sigma i}\| \leq (1 + I_1) \|\varphi_{Mi} - \varphi_{\sigma i}\| + I_2 + I_3, \quad (78)$$

with

$$I_1 = \|S_M(\lambda_\star)\|, \quad I_2 = \|S_M(\lambda_\star) - S_M(\lambda_{\sigma i})\|, \quad I_3 = \|S_M(\lambda_{\sigma i}) - S_\sigma(\lambda_{\sigma i})\|.$$

For  $I_1$ , proceeding as in the proof of Lemma 5, writing

$$S_M(\lambda) = \sum_{k=0}^{\infty} \mathsf{P}_M^\perp h_\lambda^{-1/2} \left[ -h_\lambda^{-1/2} V_M^\perp h_\lambda^{-1/2} \right]^k h_\lambda^{-1/2} \mathsf{P}_M^\perp V \mathsf{P}_M.$$

and under the assumptions  $\lambda \leq \min(\frac{1}{2}\rho_M, \kappa_M)$ ,  $\rho_M \geq 2$ ,  $4\rho_M^{-r}\|V\|_r \leq \frac{1}{2}$ , using the estimates of Lemma 4  $\|\mathbf{P}_M^\perp h_\lambda^{-1/2}\| \leq \sqrt{2}\rho_M^{-1/2}$ ,  $\|\mathbf{P}_M h_{-1}^s\| \leq (\rho_M + 1)^s \leq (3\rho_M/2)^s$  for  $s \geq 0$ , one obtains

$$\|S_M(\lambda)\| \leq 2 \|\mathbf{P}_M^\perp h_\lambda^{-1/2}\| \|\mathbf{P}_M^\perp h_\lambda^{-1/2} \mathbf{P}_M^\perp V \mathbf{P}_M h_{-1}^{-1/2+r/2}\| \|\mathbf{P}_M h_{-1}^{1/2-r/2}\| \quad (79)$$

$$\leq 2\sqrt{2}(3/2)^{1/2-r/2} \rho_M^{-1/2} \rho_M^{-r/2} \|V\|_r \rho_M^{1/2-r/2} \leq 12\rho_M^{-r} \|V\|_r. \quad (80)$$

For  $I_2$ , we proceed as in Lemma 8 based on the results of Lemma 6 but substituting  $U_M(\lambda)$  by  $S_M(\lambda)$  in order to obtain

$$I_2 \leq \frac{|\lambda_* - \lambda_{\sigma i}|}{\pi(\kappa_M - 2\min(\lambda_*, \lambda_{\sigma i}))} 12\rho_M^{-r} \|V\|_r, \quad (81)$$

based upon the estimate  $\|S_M(\lambda)\| \leq 12\rho_M^{-r} \|V\|_r$  from (79)–(80).

Finally for  $I_3$ , we proceed as in the proof of Lemma 7. Indeed, we apply, once again, the triangle inequality to obtain  $I_3 \leq I_{3,1} + I_{3,2}$  with

$$I_{3,1} = \|S_M(\lambda_{\sigma i}) - S_{MN}(\lambda_{\sigma i})\| \quad I_{3,2} = \|S_{MN}(\lambda_{\sigma i}) - S_\sigma(\lambda_{\sigma i})\|.$$

with  $S_{MN}(\lambda) := \mathbf{P}_M^N(\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N V \mathbf{P}_M$ . To estimate  $I_{3,1}$ , we first note that

$$\begin{aligned} \|S_M(\lambda) - S_{MN}(\lambda)\| &\leq \|h_{-1}^{-1/2+r/2} \mathbf{P}_M^N\| \|h_{-1}^{1/2-r/2} [\mathbf{P}_M^\perp(\mathcal{H}_M^\perp - \lambda)^{-1} \mathbf{P}_M^\perp - \mathbf{P}_M^N(\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N] h_{-1}^{1/2-r/2}\| \\ &\quad \|h_{-1}^{-1/2+r/2} V \mathbf{P}_M\| \end{aligned}$$

Using that  $\|h_{-1}^{-1/2+r/2} \mathbf{P}_M^N\| \leq \rho_M^{-1/2+r/2}$  and  $\|h_{-1}^{-1/2+r/2} V \mathbf{P}_M\| \leq \|V\|_r \rho_M^{1/2-r/2}$ , we obtain

$$\|S_M(\lambda) - S_{MN}(\lambda)\| \leq \|V\|_r \|h_{-1}^{1/2-r/2} [\mathbf{P}_M^\perp(\mathcal{H}_M^\perp - \lambda)^{-1} \mathbf{P}_M^\perp - \mathbf{P}_M^N(\mathcal{H}_M^N - \lambda)^{-1} \mathbf{P}_M^N] h_{-1}^{1/2-r/2}\|.$$

Adapting the proof of (63), and noting that  $4\rho_N^{-r} \|V\|_r - \frac{16\rho_M^{-2r} \|V\|_r^2}{1-4\rho_M^{-r} \|V\|_r} < 1$ , and  $4\rho_M^{-r} \|V\|_r < 1$ , we obtain

$$I_{3,1} \lesssim \rho_N^{-r} \|V\|_r. \quad (82)$$

Finally, for  $I_{3,2}$ , we proceed as in the derivation of (65), starting from a similar ansatz than (64)

$$S_{MN}(\lambda) - S_\sigma(\lambda) = \sum_{k=K+1}^{\infty} \mathbf{P}_M^N h_\lambda^{-1/2} \left[ -h_\lambda^{-1/2} V_M^N h_\lambda^{-1/2} \right]^k h_\lambda^{-1/2} \mathbf{P}_M^N V \mathbf{P}_M,$$

yielding, together with  $\|\mathbf{P}_M^N h_\lambda^{-1/2}\| \lesssim \rho_M^{-1/2}$  and  $\|h_\lambda^{-1/2} \mathbf{P}_M^N V \mathbf{P}_M\| \lesssim \rho_M^{1/2-r} \|V\|_r$ ,

$$I_{3,2} = \|S_{MN}(\lambda_{\sigma i}) - S_\sigma(\lambda_{\sigma i})\| \lesssim \rho_M^{-r} \|V\|_r [4\rho_M^{-r} \|V\|_r]^{K+1}. \quad (83)$$

Starting from (78) and combining (79)–(80), (81), (82), (83) with the estimates (77) (with  $s = 0$ ), (75) and the bound (74) concludes the proof.  $\square$

**Remark 10.** Having now understood how to apply Theorem 6 in this context, i.e. using (71), and seen the abstract theory of Section 3 we can now make a statement about the convergence of the fixed-point iteration schemes (19) and (20): Following Corollary 8, we note that if  $\nu_{\sigma i}(\lambda)$  parametrizes an isolated branch on some interval  $I_i$  containing  $\lambda_{\sigma i}$ ,  $M \geq M_0$  and, using (72)–(73) and (75), if  $\varepsilon(\sigma, r, V)$  is small enough, then, the fixed-point iterations converge for any starting point in  $I_i$ . Unfortunately, it is difficult to assess whether the isolated branch property holds in practical application and this remains an abstract result.

## 6. NUMERICAL RESULTS

In this section, we test the theoretical estimates developed in this article, first in a one-dimensional case, and second, in a three-dimensional case for a Coulomb potential.

**6.1. One-dimensional case.** We first consider a one-dimensional test case with  $\Omega = (0, 1)$  and a potential  $V_t$  given by its Fourier coefficients:

$$\widehat{V}_0 = -10, \quad \widehat{V}_n = -\frac{5}{|n|^t},$$

so that  $V_t \in H_{\text{per}}^{t-\frac{1}{2}-\varepsilon}$  for any  $\varepsilon > 0$ . We then consider the two different values  $t = 1$  or  $t = 0$ , see Figure 2 for a graphical illustration for the case  $t = 1$ , so that including the embedding expressed under the constraints (87) yields that  $\|V\|_r < \infty$  for  $r = 1$  if  $t = 1$  and for all  $r < 1/2$  if  $t = 0$ .

Note that, to the best of our knowledge, the classical convergence analysis does not cover such low regularities of the potentials. While the standard analysis can probably be extended to potentials in  $L^2$  and thus covering the case  $t = 1$  (although we are not aware of any published analysis in this case), it certainly does not hold without further developments for  $t = 0$ .

In Figure 3, we illustrate the convergence of the discrete solutions with respect to  $K$  for different values of  $M$  and  $t = 1$  and  $t = 0$ , and for fixed  $N = 500$ . The error in the eigenvalue and eigenvector are defined as

$$\text{err}_{\text{val}} = |\lambda_* - \lambda_{\sigma i}| \quad \text{and} \quad \text{err}_{\text{vec}} = \|\varphi_i - Q_\sigma(\lambda_{\sigma i})\varphi_{\sigma i}\|,$$

where  $(\lambda_*, \varphi_{Mi})$ ,  $(\lambda_{\sigma i}, \varphi_{\sigma i})$  are the  $i$ -th solution to (6) and (10) respectively, using the computational Strategy 1 defined by (19) targeting the smallest eigenvalue. The “exact” solution is obtained by computing the variational approximation for  $N_e = 1000$ .

We observe that, in agreement with the theory for small enough  $K, M$ , the convergence rate with respect to  $K$  for different values of  $M$  is the same for the eigenvalue and eigenvector error and that the convergence rate improves with increased values of  $M$ . In this example, we observe that the condition  $M \geq M_0$  is not restrictive and convergence can be observed for all values of  $M$ . It is also noted, in particular for the higher values of  $M$  (but still very moderate), that




FIGURE 2. Illustration of the potential  $V_{t=1}$ .

the number on the truncation order  $K$  can be kept very low to achieve a good accuracy. This, in turn, means that the number of computations on the fine grid (essentially matrix-vector products involving the fine grid for each matrix-vector product involving the coarse Hamiltonian  $\mathcal{H}_\sigma(\lambda)$ ) can be kept to a minimum.

A similar behaviour is reported in Figure 5 for the third eigenvalue using the potential  $V_{t=1}$  and again  $N = 500$ .

The number of required SCF-iterations (19) to converge to an increment in the eigenvalue smaller than  $10^{-12}$ , thus very tight, is stable and very moderate over all test cases as reported in Tables 1 and 2 (the case of  $t = 0$  behaves similarly and is not reported here).

Finally, Figure 4 illustrates the error in the eigenvalue and eigenvector with respect to  $N$  for different values of  $K$  and  $M$  for the first eigenvalue with  $V_{t=1}$  and  $V_{t=0}$ . We observe two regimes. First, for small values of  $N$ , the error is limited by the small size of the fine grid  $X_N$ , i.e. by a small  $N$ , and decreases with increasing  $N \in [25, 500]$ . Second, when  $N$  is large enough, the error due to the moderate values of  $M, K$  is dominating and the error stagnates. As  $M$  or  $K$  increase, the transition of the two regimes moves to lower accuracy. This agrees well with the theoretical result presented in Theorem 1. For the potential  $V_{t=1}$ , we observe that the convergence rate in  $N$  (for  $M, K$  large enough) is roughly 3 for the eigenvalue error and 2.5 for the eigenvector error, which are the rates predicted by the standard analysis as outlined in Corollary 2. For the less regular case  $V_{t=0}$ , and thus  $\|V\|_r < \infty$  for any  $r < 1/2$ , we observe a rate of roughly 1 in both cases, eigenvalue and eigenvector  $L^2$ -error, which is exactly as predicted by Theorem 1.

**6.2. Three-dimensional case.** We then performed similar numerical tests for a three-dimensional problem, in order to test cases closer to real applications. Note that, as in the rest of the article, we consider a linear eigenvalue problem. We focus on the lowest eigenvalue of the Schrödinger hamiltonian. To do so, we used the code Density Functional Theory Toolkit (DFTK.jl) [19], which provides a




FIGURE 3. The convergence of the eigenvalue error  $\text{err}_{\text{val}}$  (left) and the eigenvector error  $\text{err}_{\text{vec}}$  (right) corresponding to the first eigenvalue with respect to  $K$  for different values of  $M$  and for fixed  $N = 500$  in the one-dimensional case. The top row corresponds to a potential  $V_{t=1}$  whereas the bottom row corresponds to  $V_{t=0}$ .

spectral Fourier discretization for Density Functional Theory eigenvalue problems, and which can easily be adapted to treat linear eigenvalue problems. Since our method is valid for a large range of potentials, including low-regularity ones, we provide numerical simulations for the Coulomb potential, that is given in Fourier space as

$$\widehat{V}_0 = 10, \quad \widehat{V}_{\mathbf{k}} = -\frac{10}{|\mathbf{k}|^2}.$$

The lattice is chosen as a cubic lattice with a lattice constant of 10 Bohr. Since the electronic structure theory uses a different convention to define the approximation

$M = 1$		$M = 2$		$M = 3$		$M = 4$		$M = 5$	
$K$	# scf								
1	6	1	5	1	4	1	4	1	4
2	6	2	5	2	4	2	4	2	4
3	6	3	5	3	4	3	4	3	4
4	6	4	5	4	4	4	4	4	4

$M = 6$	
$K$	# scf
1	4
2	4
3	4
4	4

TABLE 1. The number of SCF-like iterations of the solution strategy (19) to reach an increment of  $10^{-12}$  in the eigenvalue for the approximation of the first eigenvalue with  $V_{t=1}$  using  $N = 500$  in the one-dimensional case.

$M = 1$		$M = 2$		$M = 3$		$M = 4$		$M = 5$	
$K$	# scf								
1	6	1	4	1	4	1	4	1	3
2	6	2	4	2	4	2	4	2	3
3	6	3	4	3	4	3	4	3	3
4	6	4	4	4	4	4	4	4	3

$M = 6$	
$K$	# scf
1	3
2	3
3	3
4	3

TABLE 2. The number of SCF-like iterations of the solution strategy (19) to reach an increment of  $10^{-12}$  in the eigenvalue for the approximation of the third eigenvalue with  $V_{t=1}$  using  $N = 500$  in the one-dimensional case.

space  $X_N$ , we now switch to a slightly different terminology and refer to the energy cutoff  $\text{EcN} = \frac{1}{2}\rho_N$ , corresponding to the largest eigenvalue/kinetic energy of the operator  $-\frac{1}{2}\Delta$  in the approximation space  $X_N$ , to fix the values of  $N$ .

In order to compute the reference (“exact”) solution an energy cutoff  $\text{EcN}_{\text{ref}} = 225$  Ha is taken, which corresponds to choosing a reference grid of size  $N_{\text{ref}} \simeq 21.2$ , with 161’235 Fourier coefficients.

We obtain similar results as in the one-dimensional case. Namely, we observe in Figure 6 that for small values of  $K$ , the convergence rate with respect to the




FIGURE 4. The convergence of the eigenvalue error  $\text{err}_{\text{val}}$  (straight lines) and the eigenvector error  $\text{err}_{\text{vec}}$  (dotted lines) corresponding to the first eigenvalue with respect to  $N$  for different values of  $K$  (left) with  $M = 2$  (top) and  $M = 1$  (bottom) and different values of  $M$  (right) with  $K = 2$  (top) and  $K = 1$  (bottom) for the potential  $V_{t=1}$  (top) and  $V_{t=0}$  (bottom), in the one-dimensional case.

energy cutoff  $\text{EcM}$  is the same for the eigenvalue and the eigenvector and increases when  $\text{EcM}$  increases, as predicted by the theory. For large values of  $K$ , we observe a saturation of the error, which is due to the error with respect to  $N$  dominating, as stated in Theorem 1.

In Figure 7, we show the analogous of Figure 4 for the one-dimensional case, which is quite similar, as we also observe two regimes. For small values of  $\text{EcN}$ , most of the error comes from the error in  $N$ , while for large values of  $\text{EcN}$ , the error mainly comes from the error in  $M$  and  $K$ . However, due to the slow convergence




FIGURE 5. The convergence of the eigenvalue error  $\text{err}_{\text{val}}$  (left) and the eigenvector error  $\text{err}_{\text{vec}}$  (right) corresponding to the third eigenvalue with respect to  $K$  for different values of  $M$  and for fixed  $N = 500$  and with  $V_{t=1}$ , in the one-dimensional case.

with respect to the grid size, we do not observe the convergence rates in EcN as precisely as in the one-dimensional case, this being mostly due to the huge size of the reference basis needed to achieve good convergence.

Moreover, the iterative algorithm used to solve the problem converges very fast. In practice, we do not need to go beyond 7 SCF iterations in all tested cases to reach a  $10^{-12}$  increment on the eigenvalue.

## 7. CONCLUSION AND PERSPECTIVES

In this paper, we have proposed a new numerical method based on the Feshbach-Schur map in combination with the spectral Fourier discretizations for linear Schrödinger eigenvalue problems. The method does not rely on the variational principle but reformulates the infinite-dimensional problem as an equivalent problem, non-linear in the spectral parameter, on a finite dimensional grid whose unknowns are the exact eigenvalue and the best-approximation of the exact eigenfunctions on the given grid. Such a problem can then be approximated by evaluating the Feshbach-Schur map on a second finer grid. The substantial contribution of this paper is an analysis in order to provide error estimates of the proposed method in all discretization parameters.

For this, we developed in Section 3 a version of perturbation theory that relies on the notion of form-boundedness with increased regularity, as stated by Assumption 1.

Having established the method and its analysis, its full benefits shall be further analyzed in future. At the present stage, it is worth to mention that, for the




FIGURE 6. The convergence of the eigenvalue error  $\text{err}_{\text{val}}$  (left) and the eigenvector error  $\text{err}_{\text{vec}}$  (right) corresponding to the first eigenvalue with respect to  $K$  for different values of energy cutoffs  $\text{EcM}$  and for fixed  $\text{EcN} = 155$ , for a (three-dimensional) Coulomb potential.




FIGURE 7. The convergence of the eigenvalue error  $\text{err}_{\text{val}}$  (straight lines) and the eigenvector error  $\text{err}_{\text{vec}}$  (dotted lines) corresponding to the first eigenvalue with respect to the energy cutoff  $\text{EcN}$  for different values of  $\text{EcM}$  (left) with  $K = 1$  and different values of  $K$  (right) with  $\text{EcM} = 4$  for a (three-dimensional) Coulomb potential.

considered one- and three-dimensional (with a Coulomb potential) problems, the

contraction in the perturbation is rather small and the non-linear iteration converge rapidly. Also, in view of more sophisticated non-linear eigenvalue problems, the artificial extra non-linearity does not seem to be much of a burden.

The future developments include the extension of Section 3 to a more general family of operators, including non-symmetric perturbations of self-adjoint operators as well as extending the numerical method and its analysis to cluster of eigenvalues using a density-matrix based formulation.

## REFERENCES

- [1] I. Babuška and J. Osborn. Eigenvalue problems. In *Handbook of Numerical Analysis*, volume 2, pages 641–787. Elsevier, Jan. 1991.
- [2] I. Babuška and J. E. Osborn. Finite element-galerkin approximation of the eigenvalues and eigenvectors of selfadjoint problems. *Mathematics of computation*, 52(186):275–297, 1989.
- [3] D. Boffi. Finite element approximation of eigenvalue problems. *Acta Numer.*, 19:1–120, 2010.
- [4] D. Brust. Electronic Spectra of Crystalline Germanium and Silicon. *Phys. Rev.*, 134(5A):A1337–A1353, June 1964.
- [5] E. Cancès, R. Chakir, L. He, and Y. Maday. Two-grid methods for a class of nonlinear elliptic eigenvalue problems. *IMA J. Numer. Anal.*, 38(2):605–645, Apr. 2018.
- [6] E. Cancès, R. Chakir, and Y. Maday. Numerical analysis of nonlinear eigenvalue problems. *Journal of Scientific Computing*, 45(1-3):90–117, 2010.
- [7] E. Cancès, R. Coyaud, and L. R. Scott. Van der Waals interactions between two hydrogen atoms: The next orders. *arXiv preprint arXiv:2007.04227*, 2020.
- [8] E. Cancès, G. Dusson, Y. Maday, B. Stamm, and M. Vohralík. A perturbation-method-based a posteriori estimator for the planewave discretization of nonlinear Schrödinger equations. *C. R. Math.*, 352(11):941–946, Nov. 2014.
- [9] E. Cancès, G. Dusson, Y. Maday, B. Stamm, and M. Vohralík. A perturbation-method-based post-processing for the planewave discretization of Kohn–Sham models. *J. Comput. Phys.*, 307:446–459, Feb. 2016.
- [10] E. Cancès, G. Dusson, Y. Maday, B. Stamm, and M. Vohralík. Post-processing of the planewave approximation of Schrödinger equations. Part I: linear operators. *IMA Journal of Numerical Analysis*, 09 2020.
- [11] E. Cancès and L. R. Scott. Van der Waals Interactions Between Two Hydrogen Atoms: The Slater–Kirkwood Method Revisited. *SIAM Journal on Mathematical Analysis*, 50(1):381–410, 2018.
- [12] F. Chatelin. *Spectral approximation of linear operators*. SIAM, 2011.
- [13] H. Cycon, R. Froese, W. Kirsch, and B. Simon. *Schrödinger operators with application to quantum mechanics and global geometry*. Springer-Verlag, 1987.
- [14] X. Dai and A. Zhou. Three-scale finite element discretizations for quantum eigenvalue problems. *SIAM journal on numerical analysis*, 46(1):295–324, 2008.
- [15] G. Dusson. Post-processing of the plane-wave approximation of Schrödinger equations. Part II: Kohn–Sham models. *IMA Journal of Numerical Analysis*, 09 2020.
- [16] M. Griesemer and D. Hasler. On the smooth Feshbach–Schur map. *J. Funct. Anal.*, 254(9):2329–2335, May 2008.
- [17] P. Grisvard. *Elliptic problems in nonsmooth domains*, volume 24 of *Monographs and Studies in Mathematics*. Pitman Advanced Publishing Program, Boston, MA, 1985.
- [18] S. J. Gustafson and I. M. Sigal. *Mathematical Concepts of Quantum Mechanics*. Springer Science & Business Media, Sept. 2011.

- [19] M. F. Herbst, A. Levitt, and E. Cancès. DFTK: A julian approach for simulating electrons in solids. *JuliaCon Proceedings*, 3(26):69, May 2021.
- [20] P. D. Hislop and I. M. Sigal. *Introduction to spectral theory: With applications to Schrödinger operators*. Springer-Verlag, 1996.
- [21] T. Kato. *Perturbation theory for linear operators*. Springer Berlin Heidelberg, 1976.
- [22] C. Möller and M. S. Plesset. Note on an approximation treatment for many-electron systems. *Physical Review*, 46(7):618–622, Oct. 1934.
- [23] R. Norton and R. Scheichl. Convergence analysis of planewave expansion methods for 2d schrödinger operators with discontinuous periodic potentials. *SIAM journal on numerical analysis*, 47(6):4356–4380, 2010.
- [24] M. Reed and B. Simon. *Methods of Modern Mathematical Physics Vol. 2: Fourier Analysis, Self-Adjointness*. Elsevier, 1975.
- [25] M. Reed and B. Simon. *Methods of Modern Mathematical Physics. Vol. 4. Operator Analysis*. Academic Press, New York, 1979.
- [26] F. Rellich and J. Berkowitz. *Perturbation theory of eigenvalue problems*. CRC Press, 1969.
- [27] J. Xu and A. Zhou. A two-grid discretization scheme for eigenvalue problems. *Math. Comput.*, 70(233):17–26, Aug. 1999.

## APPENDIX A. TECHNICAL RESULTS AND PROOFS

We present proofs of some technical statements made in the previous sections and some additional technical results. In what follows, we denote  $V_N^\perp := \mathsf{P}_N^\perp V \mathsf{P}_N^\perp$ .

**Lemma 12.** *Under Assumption 1, (i) the norm  $\|V\|_{0,\alpha} := \|(-\Delta + \alpha)^{-1/2}V(-\Delta + \alpha)^{-1/2}\|$  is well-defined for  $\alpha > 0$  and  $\|V\|_{0,\alpha} \rightarrow 0$  when  $\alpha \rightarrow +\infty$  and (ii)  $\mathcal{H}$  is bounded below as*

$$\mathcal{H} \geq [1 - \|V\|_{0,\alpha}](-\Delta + \alpha) - \alpha, \quad (84)$$

*Proof.* The first statement is obvious and the second one follows from the identity

$$\mathcal{H} = (-\Delta + \alpha)^{1/2}[\text{Id} + (-\Delta + \alpha)^{-1/2}V(-\Delta + \alpha)^{-1/2}](-\Delta + \alpha)^{1/2} - \alpha \quad (85)$$

and the fact that the expression in square braces is bounded below by  $1 - \|V\|_{0,\alpha}$ .  $\square$

*Proof of Lemma 1.* Proceeding as in (84), we obtain for any  $\alpha \in \mathbb{R}$  large enough so that  $\|V_M^\perp\|_{0,\alpha} < 1$

$$-\Delta + V_M^\perp \geq [1 - \|V_M^\perp\|_{0,\alpha}](-\Delta + \alpha) - \alpha.$$

Using that  $-\Delta \geq (2\pi M/L)^2 = \rho_M$  on  $\text{Ran } \mathsf{P}_M^\perp$  and the estimate

$$\begin{aligned} \|V_M^\perp\|_{0,\alpha} &\leq \|(1 - \Delta)^{1/2}(-\Delta + \alpha)^{1/2}\mathsf{P}_M^\perp\|^2 \|(1 - \Delta)^{-1/2}\mathsf{P}_M^\perp V \mathsf{P}_M^\perp(1 - \Delta)^{-1/2}\| \\ &\leq \frac{\rho_M + 1}{\rho_M + \alpha} \rho_M^{-r} \|V\|_r, \end{aligned}$$

we deduce

$$\begin{aligned} -\Delta + V_M^\perp &\geq \left(1 - \frac{\rho_M + 1}{\rho_M + \alpha} \rho_M^{-r} \|V\|_r\right) (\rho_M + \alpha) - \alpha, \\ &\geq \rho_M + \alpha - (\rho_M + 1) \rho_M^{-r} \|V\|_r - \alpha, \\ &\geq \rho_M - (\rho_M + 1) \rho_M^{-r} \|V\|_r, \end{aligned}$$

from which we obtain the result.  $\square$

Now, we introduce the Sobolev spaces  $H_{\text{per}}^s$  of periodic real functions resp. distributions which can be characterized in a simple way using Fourier series: for  $s \in \mathbb{R}$ , we have

$$H_{\text{per}}^s := \left\{ v = \sum_{\mathbf{k} \in \mathcal{R}^*} \widehat{v}_{\mathbf{k}} e_{\mathbf{k}} \mid \forall \mathbf{k} \in \mathcal{R}^* : \quad \widehat{v}_{-\mathbf{k}} = \overline{\widehat{v}_{\mathbf{k}}}, \quad \|v\|_{H_{\text{per}}^s} < \infty \right\}, \quad (86)$$

where the norm  $\|v\|_{H_{\text{per}}^s}$  is given by the  $H_{\text{per}}^s$  inner product is defined by

$$\forall u, v \in H_{\text{per}}^s, \quad (u, v)_{H_{\text{per}}^s} := \sum_{\mathbf{k} \in \mathcal{R}^*} (1 + |\mathbf{k}|^2)^s \overline{u_{\mathbf{k}}} v_{\mathbf{k}}.$$

**Lemma 13.** *If  $V \in H_{\text{per}}^s$  for some  $s \in \mathbb{R}$ , then  $V$  satisfies Assumption 1 for all  $r \geq 0$  satisfying  $s \geq r - 1$  and  $s > -2(1 - r) + d/2$  resp.*

$$r \leq s + 1 \quad \text{and} \quad r < 1 + \frac{s}{2} - \frac{d}{4}, \quad (87)$$

where  $d$  is the spatial dimension. Moreover,

$$\|V\|_r \leq C_{r,s} \|V\|_{H_{\text{per}}^s}.$$

*Proof.* From [17, Theorem 1.4.4.2], for any  $s_1, s_2 \geq t$ , such that  $s_1 + s_2 > t + d/2$ , there exists  $C_{s_1, s_2, t} > 0$  such that for all  $u \in H_{\text{per}}^{s_1}, v \in H_{\text{per}}^{s_2}$ , then  $uv \in H_{\text{per}}^t$  and

$$\|uv\|_{H_{\text{per}}^t} \leq C_{s_1, s_2, t} \|u\|_{H_{\text{per}}^{s_1}} \|v\|_{H_{\text{per}}^{s_2}}.$$

Hence if  $V \in H_{\text{per}}^s$ ,  $s > -2(1 - r) + d/2$ , and  $s \geq r - 1$ , then there exists  $C_{r,s}$  such that

$$\|V\|_r = \sup_{\psi \in H_{\text{per}}^{1-r}} \frac{\|V\psi\|_{H_{\text{per}}^{-1+r}}}{\|\psi\|_{H_{\text{per}}^{1-r}}} \leq C_{r,s} \|V\|_{H_{\text{per}}^s},$$

which implies the estimate in the lemma.  $\square$

*Proof of Proposition 1.* For each  $\lambda$  such that  $\lambda < \kappa_M$ ,  $U_M(\lambda)$  can be written as the product of five bounded operators:  $-\mathsf{P}_M(-\Delta + 1)^{1/2-r/2}$ ,  $(-\Delta + 1)^{-1/2+r/2}V(-\Delta + 1)^{-1/2+r/2}$ ,  $(-\Delta + 1)^{1/2-r/2}\mathsf{P}_M^\perp(\mathcal{H}_M^\perp - \lambda)^{-1}\mathsf{P}_M^\perp(-\Delta + 1)^{1/2-r/2}$ ,  $(-\Delta + 1)^{-1/2+r/2}V(-\Delta + 1)^{-1/2+r/2}$ ,  $(-\Delta + 1)^{1/2-r/2}\mathsf{P}_M$ . Therefore,  $U_M(\lambda)$  is a well-defined operator.  $\square$

LABORATOIRE DE MATHÉMATIQUES DE BESANÇON, UMR CNRS 6623, UNIVERSITÉ BOURGOGNE FRANCHE-COMTÉ, 16 ROUTE DE GRAY, 25030 BESANÇON, FRANCE

*Email address:* genevieve.dusson@math.cnrs.fr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, 40 ST. GEORGE STREET,  
BAHEN CENTRE, TORONTO, ON M5S 2E4, CANADA

*Email address:* im.sigal@utoronto.ca

APPLIED AND COMPUTATIONAL MATHEMATICS, DEPARTMENT OF MATHEMATICS, RWTH  
AACHEN UNIVERSITY, SCHINKELSTR. 2, 52062 AACHEN, GERMANY

*Email address:* best@acom.rwth-aachen.de