



**HAL**  
open science

## Towards a noncommutative Picard-Vessiot theory

G. Duchamp, Viin cel Hoang Ngoc Minh, Vu Nguyen Dinh

► **To cite this version:**

G. Duchamp, Viin cel Hoang Ngoc Minh, Vu Nguyen Dinh. Towards a noncommutative Picard-Vessiot theory. 2020. hal-02921131v1

**HAL Id: hal-02921131**

**<https://hal.science/hal-02921131v1>**

Preprint submitted on 24 Aug 2020 (v1), last revised 24 Oct 2021 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Towards a noncommutative Picard-Vessiot theory

G.H.E. Duchamp

*University Paris 13, Sorbonne Paris City, 93430 Villetaneuse, France,*

V. Hoang Ngoc Minh

*University of Lille, 1 Place Déliot, 59024 Lille, France,*

V. Nguyen Dinh

*Institute of Mathematics, Vietnam Academy of Science and Technology,  
18 Hoang Quoc Viet, 10307 Hanoi, Vietnam.*

August 24, 2020 (Draft)

---

## **Abstract**

A Chen generating series, along a path and with respect to  $m$  differential forms, is a non-commutative series on  $m$  letters and with coefficients which are holomorphic functions over a simply connected manifold in other words a series with variable (holomorphic) coefficients. Such a series satisfies a first order noncommutative differential equation which is considered, by some authors, as the universal differential equation, *i.e.* universality can be seen by replacing each letter by constant matrices (resp. analytic vector fields) and then solving a system of linear (resp. nonlinear) differential equations.

Via rational series, on noncommutative indeterminates and with coefficients in rings, and their non-trivial combinatorial Hopf algebras, we give the first step of a noncommutative Picard-Vessiot theory and we illustrate it with the case of linear differential equations with singular regular singularities thanks to the universal equation previously mentioned.

---

## Contents

1	Introduction	2
2	Combinatorial framework	4
2.1	Factorization in bialgebras	4
2.2	Representative series	8
3	Triangularity, solvability and rationality	12
3.1	Syntactically exchangeable rational series	12
3.2	Exchangeable rational series and their linear representations	14
4	Towards a noncommutative Picard-Vessiot theory	19
4.1	Noncommutative differential equations	19
4.2	First step of a noncommutative Picard-Vessiot theory	22
5	Conclusion	24
	References	24

## 1 Introduction

Combinatorial Picard-Vessiot (PV for short) theory of bilinear systems<sup>1</sup> was realized by Fliess and Reutenauer [29], as an application of differential algebra [42,47]. This theory allows to employ, with success, linear algebraic groups in control theory (*i.e.* as symmetry groups of linear differential equations), for which some questions were solved thanks to the theory of Hopf algebras [11] and some combinatorial and effective aspects were set in [46].

Let us, for instance, consider the following nonlinear dynamical system

$$\dot{q}(z) = A_0(q)u_0(z) + \dots + A_m(q)u_m(z), q(z_0) = q_0, y(z) = f(q(z)), \quad (1)$$

where

- (i)  $y$  is the output,
- (ii) the vector state  $q = (q_1, \dots, q_n)$  belongs to a complex analytic manifold  $\mathcal{M}$  of dimension  $n$ ,
- (iii) the observation  $f$  is defined within a fixed connected neighbourhood<sup>2</sup>  $U$  of the initial state  $q_0$ .
- (iv) the vector fields  $(A_i)_{i=0, \dots, m}$  are defined with respect to the coordinates as follows

<sup>1</sup> Namely - locally - linear of the states  $q_1, \dots, q_n$  and linear of the inputs  $u_0, \dots, u_m$  [29].

<sup>2</sup> In this introductory description the points are loosely identified with their coordinates through some chart  $\varphi_U : U \rightarrow \mathbb{C}^n$  likewise, in [45], the space of analytic functions  $\mathcal{H}(U)$  is described by  $\mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]$ .

$$A_i = \sum_{j=1}^n A_i^j(q) \frac{\partial}{\partial q_j}, \text{ with } A_i^j(q) \in \mathcal{H}(U), \quad (2)$$

(v) the inputs  $(u_i)_{i=0,\dots,m}$ , as well as their inverses  $(u_i^{-1})_{i=0,\dots,m}$ , belong to a subring,  $\mathcal{C}_0$ , of the ring of holomorphic functions  $\mathcal{H}(\Omega)$  with neutral element  $1_{\mathcal{H}(\Omega)}$  over the simply connected manifold<sup>3</sup>  $\Omega$ .

It is convenient to separate the contribution of the vector fields  $(A_i)_{i=0,\dots,m}$  and that of the differential forms  $(\omega_i)_{i=0,\dots,m}$ , defined by the inputs  $\omega_i(z) = u_i(z)dz$ , through the encoding alphabet  $X = \{x_i\}_{i=0,\dots,m}$  which generates the monoid  $X^*$  with neutral element  $1_{X^*}$ . Indeed, the output  $y$  can be computed by

$$y(z_0, z) = \langle C_{z_0 \rightsquigarrow z} | \sigma f|_{q_0} \rangle = \sum_{w \in X^*} \alpha_{z_0}^z(w) \mathcal{Y}(w) [f]|_{q_0}, \quad (3)$$

as the pairing (under suitable convergence conditions [31,34,36,45]) between the Chen series<sup>4</sup> of  $(\omega_i)_{i=0,\dots,m}$  along the path  $z_0 \rightsquigarrow z$  over  $\Omega$ ,  $C_{z_0 \rightsquigarrow z} \in \mathcal{H}(\Omega) \langle \langle X \rangle \rangle$  [10], and the generating series of (1),  $\sigma f|_{q_0} \in \mathcal{H}(U) \langle \langle X \rangle \rangle$  [31], defined as follows

$$C_{z_0 \rightsquigarrow z} := \sum_{w \in X^*} \alpha_{z_0}^z(w) w \text{ and } \sigma f|_{q_0} := \sum_{w \in X^*} \mathcal{Y}(w) [f]|_{q_0} w, \quad (4)$$

where, in (3)–(4), the iterated integral  $\alpha_{z_0}^z(w)$  and the differential operator  $\mathcal{Y}(w)$ , are decoded, from the word  $w \in X^*$ , recursively as follows

$$\begin{cases} \alpha_{z_0}^z(w) = 1_{\mathcal{H}(\Omega)} & \text{and } \mathcal{Y}(w) = \text{Id}, & \text{for } w = 1_{X^*}, \\ \alpha_{z_0}^z(w) = \int_{z_0}^z \omega_i(s) \alpha_{z_0}^s(v) & \text{and } \mathcal{Y}(w) = A_i \circ \mathcal{Y}(v), & \text{for } w = x_i v, \\ & & x_i \in X, v \in X^*. \end{cases} \quad (5)$$

In this work, following this route, considering the differential ring  $(\mathcal{H}(\Omega), \partial)$  and equipping  $\mathcal{H}(\Omega) \langle \langle X \rangle \rangle$  with the derivation defined, for any  $S \in \mathcal{H}(\Omega) \langle \langle X \rangle \rangle$ , by

$$\mathbf{d}S = \sum_{w \in X^*} (\partial \langle S | w \rangle) w, \quad (6)$$

the Chen series satisfies the following noncommutative differential equation

$$\mathbf{d}S = MS \text{ with } M = u_0 x_0 + \dots + u_m x_m, \quad (7)$$

considered by many authors as the universal differential equation [10,14,17,18,39], *i.e.* universality can be seen by specialization, *i.e.* replacing the letters by constant matrices (resp. analytic vector fields) obtaining linear (resp. nonlinear) differential equations (see Remark 4.9 below) and their solutions. From the equation (7), it follows (see, for example, [9]) that a PV theory of nonlinear systems (1) should be

<sup>3</sup> This (usually one dimensional) manifold will be the support of the iterated integrals below.

<sup>4</sup> By a Ree's theorem [44], there is a primitive series  $L_{z_0 \rightsquigarrow z} = \sum_{n \geq 1} L_n \in \widehat{\mathcal{H}(\Omega) \langle \langle X \rangle \rangle}$  s.t.  $e^{L_{z_0 \rightsquigarrow z}} = C_{z_0 \rightsquigarrow z}$ , meaning that  $C_{z_0 \rightsquigarrow z}$  is group-like and  $L_n$  is (homogenous of degree  $n \geq 1$ ) primitive series.

intimately connected with (7) (remark that, due to the fact that  $\Omega$  is connected, the constants of  $(\mathcal{H}(\Omega)\langle\langle X \rangle\rangle, \mathbf{d})$  are such that

$$\text{Const}(\mathcal{H}(\Omega)\langle\langle X \rangle\rangle) = \ker \mathbf{d} = \mathbb{C} \cdot 1_{\mathcal{H}(\Omega)\langle\langle X \rangle\rangle}. \quad (8)$$

It culminates with the fact that the coefficients of any suitable<sup>5</sup> solution is group-like, *i.e.* satisfies<sup>6</sup>, for any  $u, v \in X^*$  and  $x_i \in X$ ,

$$\partial \langle S | x_i u \rangle = u_i \langle S | v \rangle \text{ and } \langle S | u \sqcup v \rangle = \langle S | u \rangle \langle S | v \rangle \quad (9)$$

and, due to the fact that  $\Omega$  is simply connected, its values only depend on the endpoints and not on paths belonging to  $\Omega$ . Denoting the subalgebra of  $(\mathcal{H}(\Omega), \partial)$  generated by the family  $(f_i)_{i \in I}$  and derivatives by  $\mathbb{C}\{\{(f_i)_{i \in I}\}\}$  [49] (*i.e.* the differential algebra generated by  $(f_i)_{i \in I}$ ), it follows that [29]

$$\text{span}_{\mathbb{C}}\{\langle \mathbf{d}^l S | w \rangle\}_{w \in X^*, l \geq 0} \subset \text{span}_{\mathbb{C}\{\{(u_i)_{i=0, \dots, m}\}\}}\{\langle S | w \rangle\}_{w \in X^*} \quad (10)$$

$$\subset \text{span}_{\mathbb{C}\{\{(u_i^{\pm 1})_{i=0, \dots, m}\}\}}\{\langle S | w \rangle\}_{w \in X^*} \quad (11)$$

and then, in Section 4, the isomorphism between  $\text{span}_{\mathbb{C}\{\{(u_i^{\pm 1})_{i=0, \dots, m}\}\}}\{\alpha_{z_0}^z(w)\}_{w \in X^*}$  and  $\mathbb{C}\{\{(u_i^{\pm 1})_{i=0, \dots, m}\}\} \otimes_{\mathbb{C}} \text{span}_{\mathbb{C}}\{\alpha_{z_0}^z(w)\}_{w \in X^*}$  will be examined (Theorem 4.4) via the PV-extension related to (7) and, on the other hand, the output of (1) will be computed (Theorem 4.8) by pairing the series given in (4). As example, this calculation will be achieved according to the algebraic combinatorics of rational series, established beforehand in Sections 2 (Theorems 2.1, 2.3) and 3 (Theorems 3.2, 3.7).

## 2 Combinatorial framework

In this section, coefficients are taken in a commutative ring<sup>7</sup>  $A$  and, unless explicitly stated, all tensor products will be considered over the ambient ring (or field).

### 2.1 Factorization in bialgebras

In section 1, the encoding alphabet  $X$  was already introduced. In particular, for  $m = 1$  (*i.e.*  $X = \{x_0, x_1\}$ ), let us note that there are one-to-one correspondences

$$(s_1, \dots, s_r) \in \mathbb{N}_+^r \leftrightarrow x_0^{s_1-1} x_1 \dots x_0^{s_r-1} x_1 \in X^* x_1 \xrightleftharpoons[\pi_X]{\pi_Y} y_{s_1} \dots y_{s_r} \in Y^*, \quad (12)$$

where  $Y := \{y_k\}_{k \geq 1}$  and  $\pi_X$  is the conc morphism, from  $A\langle Y \rangle$  to  $A\langle X \rangle$ , mapping  $y_k$  to  $x_0^{k-1} x_1$ . This morphism  $\pi_X$  admits an adjoint  $\pi_Y$  for the two standard scalar

<sup>5</sup> *i.e.* group-like at one - interior or frontier - point.

<sup>6</sup> In the first identity, also called Friedrichs criterion, is involved the shuffle product ( $\sqcup$ ) [10,30,46].

<sup>7</sup> although some of the properties already hold for a general commutative semiring [1].

products<sup>8</sup> which has a simple combinatorial description: the restriction of  $\pi_Y$  to the subalgebra  $(A1_{X^*} \oplus A\langle Y \rangle_{x_1}, \cdot)$ , is an isomorphism given by  $\pi_Y(x_0^{k-1}x_1) = y_k$  (and the kernel of the non-restricted  $\pi_Y$  is  $A\langle X \rangle_{x_0}$ ). For all matters concerning finite ( $X$  and similar) or infinite ( $Y$  and similar) alphabets, we will use a generic model noted  $\mathcal{X}$  in order to state their common combinatorial features.

As an algebra the  $A$ -module  $A\langle \mathcal{X} \rangle$  is equipped with the associative unital concatenation and the associative commutative and unital shuffle product. The latter being defined, for any  $x, y \in \mathcal{X}$  and  $u, v, w \in \mathcal{X}^*$ , by the following recursion

$$w \sqcup 1_{\mathcal{X}^*} = 1_{\mathcal{X}^*} \sqcup w = w \text{ and } xu \sqcup yv = x(u \sqcup yv) + y(xu \sqcup v) \quad (13)$$

or, equivalently, by its dual comultiplication (which is a morphism for concatenations<sup>9</sup>), defined, for each letter  $x \in \mathcal{X}$ , by

$$\Delta_{\sqcup}(x) = 1_{\mathcal{X}^*} \otimes x + x \otimes 1_{\mathcal{X}^*}. \quad (14)$$

Once  $\mathcal{X}$  has been totally ordered<sup>10</sup>, the set of Lyndon words over  $\mathcal{X}$  will be denoted by  $\mathcal{Lyn}\mathcal{X}$ . A pair of Lyndon words  $(l_1, l_2)$  is called the standard factorization of a Lyndon  $l$  (and will be noted  $(l_1, l_2) = st(l)$ ) if  $l = l_1l_2$  and  $l_2$  is the longest nontrivial proper right factor of  $l$  or, equivalently, its smallest such (for the lexicographic ordering, see [43] for proofs and details). According to a theorem by Radford, the set of Lyndon words form a pure transcendence basis of the  $A$ -shuffle algebras  $(A\langle \mathcal{X} \rangle, \sqcup, 1_{\mathcal{X}^*})$ .

It is well known that the enveloping algebra  $\mathcal{U}(\mathcal{L}ie_A\langle \mathcal{X} \rangle)$  is isomorphic to the (connected, graded and co-commutative) bialgebra<sup>11</sup>  $\mathcal{H}_{\sqcup}(\mathcal{X}) = (A\langle \mathcal{X} \rangle, \text{conc}, 1_{\mathcal{X}^*}, \Delta_{\sqcup}, \mathbf{e})$  (the counit being here  $\mathbf{e}(P) = \langle P | 1_{\mathcal{X}^*} \rangle$ ) and, via the pairing

$$A\langle \langle \mathcal{X} \rangle \rangle \otimes_A A\langle \mathcal{X} \rangle \longrightarrow A, \quad (15)$$

$$T \otimes P \longrightarrow \langle T | P \rangle := \sum_{w \in \mathcal{X}^*} \langle T | w \rangle \langle P | w \rangle, \quad (16)$$

we can, classically, endow  $A\langle \mathcal{X} \rangle$  with the graded<sup>12</sup> linear basis  $\{P_w\}_{w \in \mathcal{X}^*}$  (expanded after any homogeneous basis  $\{P_l\}_{l \in \mathcal{Lyn}\mathcal{X}}$  of  $\mathcal{L}ie_A\langle \mathcal{X} \rangle$ ) and its graded dual basis  $\{S_w\}_{w \in \mathcal{X}^*}$  (containing the pure transcendence basis  $\{S_l\}_{l \in \mathcal{Lyn}\mathcal{X}}$  of the

<sup>8</sup> That is to say  $(\forall p \in A\langle X \rangle) (\forall q \in A\langle Y \rangle) (\langle \pi_Y p | q \rangle_Y = \langle p | \pi_X q \rangle_X)$ .

<sup>9</sup> On  $A\langle \mathcal{X} \rangle$  and  $A\langle \mathcal{X} \rangle \otimes A\langle \mathcal{X} \rangle$ , respectively.

<sup>10</sup> For technical reasons, the orders  $x_0 < x_1$  (for  $X$ ) and  $y_1 > \dots y_n > y_{n+1} > \dots$  (for  $Y$ ) are usual.

<sup>11</sup> In case  $A$  is a  $\mathbb{Q}$ -algebra, the isomorphism  $\mathcal{U}(\mathcal{L}ie_A\langle \mathcal{X} \rangle) \simeq \mathcal{H}_{\sqcup}(\mathcal{X})$  can also be seen as an easy application of the CQMM theorem.

<sup>12</sup> For  $\mathcal{X} = X$  or  $= Y$  the corresponding monoids are equipped with length functions, for  $X$  we consider the length of words and for  $Y$  the length is given by the weight  $\ell(y_{i_1} \dots y_{i_n}) = i_1 + \dots + i_n$ . This naturally induces a grading of  $A\langle \mathcal{X} \rangle$  and  $\mathcal{L}ie_A\langle \mathcal{X} \rangle$  in free modules of finite dimensions. For general  $\mathcal{X}$ , we consider the fine grading [46] *i.e.* the grading by all partial degrees which, as well, induces a grading of  $A\langle \mathcal{X} \rangle$  and  $\mathcal{L}ie_A\langle \mathcal{X} \rangle$  in free modules of finite dimensions.

$A$ -shuffle algebra). In the case when  $A$  is a  $\mathbb{Q}$ -algebra, we also have the following factorization<sup>13</sup> of the diagonal series, *i.e.* [46] (here all tensor products are over  $A$ )

$$\mathcal{D}_{\mathcal{X}} := \sum_{w \in \mathcal{X}^*} w \otimes w = \sum_{w \in \mathcal{X}^*} S_w \otimes P_w = \prod_{l \in \mathcal{L}yn\mathcal{X}} \overrightarrow{e^{S_l \otimes P_l}} \quad (17)$$

and (still in case  $A$  is a  $\mathbb{Q}$ -algebra) dual bases of homogenous polynomials  $\{P_w\}_{w \in \mathcal{X}^*}$  and  $\{S_w\}_{w \in \mathcal{X}^*}$  can be constructed recursively as follows

$$\begin{cases} P_x = x, & S_x = x & \text{for } x \in \mathcal{X}, \\ P_l = [P_{l_1}, P_{l_2}], & S_l = yS_{l'}, & \text{for } l = yl' \in \mathcal{L}yn\mathcal{X} - \mathcal{X} \\ & & st(l) = (l_1, l_2), \\ P_w = P_{l_1}^{i_1} \dots P_{l_k}^{i_k}, & S_w = \frac{S_{l_1}^{\sqcup i_1} \sqcup \dots \sqcup S_{l_k}^{\sqcup i_k}}{i_1! \dots i_k!}, & \text{for } w = l_1^{i_1} \dots l_k^{i_k}, \text{ with } l_1, \dots, \\ & & l_k \in \mathcal{L}yn\mathcal{X}, l_1 > \dots > l_k. \end{cases} \quad (18)$$

The graded dual of  $\mathcal{H}_{\sqcup}(\mathcal{X})$  is  $\mathcal{H}_{\sqcup}^{\vee}(\mathcal{X}) = (A\langle \mathcal{X} \rangle, \sqcup, 1_{\mathcal{X}^*}, \Delta_{\text{conc}}, \varepsilon)$ .

As an algebra, the module  $A\langle Y \rangle$  is also equipped with the associative commutative and unital quasi-shuffle product defined, for  $u, v, w \in Y^*$  and  $y_i, y_j \in Y$ , by

$$w \sqcup 1_{Y^*} = 1_{Y^*} \sqcup w = w, \quad (19)$$

$$y_i u \sqcup y_j v = y_i (u \sqcup y_j v) + y_j (y_i u \sqcup v) + y_{i+j} (u \sqcup v). \quad (20)$$

This product also can be dualized according to  $(y_k \in Y)$

$$\Delta_{\sqcup}(y_k) := y_k \otimes 1_{Y^*} + 1_{Y^*} \otimes y_k + \sum_{i+j=k} y_i \otimes y_j \quad (21)$$

which is also a conc-morphism (see [28]). We then get another (connected, graded and co-commutative) bialgebra which, in case  $A$  is a  $\mathbb{Q}$ -algebra, is isomorphic to the enveloping algebra of the Lie algebra of its primitive elements<sup>14</sup>,

$$\mathcal{H}_{\sqcup}(Y) = (A\langle Y \rangle, \cdot, 1_{Y^*}, \Delta_{\sqcup}, \varepsilon) \cong \mathcal{U}(\text{Prim}(\mathcal{H}_{\sqcup}(Y))), \quad (22)$$

where  $\text{Prim}(\mathcal{H}_{\sqcup}(Y)) = \text{Im}(\pi_1) = \text{span}_A\{\pi_1(w) | w \in Y^*\}$  and  $\pi_1$  is the eulerian projector defined by [37,38]

$$\forall w \in Y^*, \pi_1(w) = w + \sum_{k=2}^{(w)} \frac{(-1)^{k-1}}{k} \sum_{u_1, \dots, u_k \in Y^+} \langle w | u_1 \sqcup \dots \sqcup u_k \rangle u_1 \dots u_k, \quad (23)$$

and, for any  $w = y_{i_1} \dots y_{i_k} \in Y^*$ ,  $(w)$  denotes the number  $i_1 + \dots + i_k$ .

Now, let  $\{\Pi_w\}_{w \in Y^*}$  be the linear basis, expanded by decreasing Poincaré-Birkhoff-Witt (PBW for short) after any basis  $\{\Pi_l\}_{l \in \mathcal{L}ynY}$  of  $\text{Prim}(\mathcal{H}_{\sqcup}(Y))$  homogeneous in weight<sup>15</sup>, and let  $\{\Sigma_w\}_{w \in Y^*}$  be its dual basis which contains the

<sup>13</sup> Also called MSR factorization after the names of Mélançon, Schützenberger and Reutenauer.

<sup>14</sup> Remark that, for  $\Delta_{\sqcup}(y_k)$ ,  $y_1 \in Y$  is the only primitive letter, see (21).

<sup>15</sup> Factorization (24) will be true in particular for the basis (25) explicitly constructed there.



## 2.2 Representative series

Representative (or rational) series are the representative functions on the free monoid<sup>17</sup> [22] and their magic is that it rests on four (apparently distant) pillars:

- Separated coproduct (SC)<sup>18</sup>,
- Finite orbit by shifts (FS),
- Result of a rational expression (RE),
- Linear representation (LR).

We first define what shifts, for (FS), and the Kleene star, for (RE) are, and then state the equivalence:

**Definition 2.2** Let  $S \in A\langle\langle\mathcal{X}\rangle\rangle$  (resp.  $A\langle\mathcal{X}\rangle$ ) and  $P \in A\langle\mathcal{X}\rangle$  (resp.  $A\langle\langle\mathcal{X}\rangle\rangle$ ).

- (i) The *left* (resp. *right*) *shift*<sup>19</sup> of  $S$  by  $P$ , is  $P \triangleright S$  (resp.  $S \triangleleft P$ ) defined by<sup>20</sup>

$$\forall w \in \mathcal{X}^*, \langle P \triangleright S \mid w \rangle = \langle S \mid wP \rangle \text{ (resp. } \langle S \triangleleft P \mid w \rangle = \langle S \mid Pw \rangle).$$

- (ii) For any  $S \in A\langle\langle\mathcal{X}\rangle\rangle$  such that  $\langle S \mid 1_{\mathcal{X}^*} \rangle = 0$ , the Kleene star of  $S$  is defined as<sup>21</sup>  $S^* = (1 - S)^{-1}$ .
- (iii) In case  $A = K$  is a field, one can define also the Sweedler's dual  $\mathcal{H}_{\sqcup}^{\circ}(\mathcal{X})$  of  $\mathcal{H}_{\sqcup}(\mathcal{X})$  by  $S \in \mathcal{H}_{\sqcup}^{\circ}(\mathcal{X}) \iff \Delta_{\text{conc}}(S) = \sum_{i \in I} G_i \otimes D_i$  [46], for some  $I$  finite,  $\{G_i\}_{i \in I}, \{D_i\}_{i \in I}$  being series (as a matter of fact, it can be shown that they even can be chosen in  $\mathcal{H}_{\sqcup}^{\circ}(\mathcal{X})$ , see [19,39])

**Theorem 2.3** ([20,22,35,46]) For  $S \in A\langle\langle\mathcal{X}\rangle\rangle$ , the following assertions are equivalent<sup>22</sup>

- (i) The shifts  $\{S \triangleleft w\}_{w \in \mathcal{X}^*}$  (resp.  $\{w \triangleright S\}_{w \in \mathcal{X}^*}$ ) lie in a finitely generated shift-

<sup>17</sup> These functions were considered on groups in [11,12].

<sup>18</sup> Uniquely for fields.

<sup>19</sup> Some schools (as Jacob one, see [40,32]) used to call this a *residual*. These actions are none other than the shifts of functions of harmonic analysis.

<sup>20</sup> They are associative, commute with each other:  $S \triangleleft (PR) = (S \triangleleft P) \triangleleft R, P \triangleright (R \triangleright S) = (P \triangleright R) \triangleright S$  and  $(P \triangleleft S) \triangleright R = P \triangleleft (S \triangleright R)$  and, for  $x, y \in \mathcal{X}, w \in \mathcal{X}^*, x \triangleright (wy) = (yw) \triangleleft x = \delta_x^y w$  (Kronecker delta).

<sup>21</sup> Using one of the topologies of section 4.2 (adapted with  $A$  replacing  $\mathcal{H}(\Omega)$ ), we have  $S^* = \sum_{n \geq 0} S^n$ . We also get the fact that the space  $\widehat{A \cdot \mathcal{X}}$  (used below) of series of degree 1, *i.e.* the set  $\{\sum_{x \in \mathcal{X}} \alpha(x)x\}_{\alpha \in A^{\mathcal{X}}}$  is the closure of the  $A$ -module  $A \cdot \mathcal{X}$  generated by letters. In the case of a finite alphabet however (here  $\mathcal{X} = X$ ) [22],  $\widehat{A \cdot \mathcal{X}} = A \cdot \mathcal{X}$ .

<sup>22</sup> When  $A$  is noetherian, first condition is equivalent to the fact that the module generated by  $\{S \triangleleft w\}_{w \in \mathcal{X}^*}$  (resp.  $\{w \triangleright S\}_{w \in \mathcal{X}^*}$ ) is finitely generated (and more precisely, in this case, by a finite number of those shifts). Unfortunately we are not in this case here, but our ring being without zero divisors (analytic functions), we can use the fraction field, here being realized by germs [15].

invariant  $A$ -module<sup>23</sup>.

- (ii) The series  $S$  belongs to the (algebraic) closure of  $\widehat{A \langle \mathcal{X} \rangle}$  by the operations  $\{\text{conc}, +, *\}$  (within  $A \langle \langle \mathcal{X} \rangle \rangle$ ).
- (iii) There is a linear representation  $(\mathbf{v}, \mu, \eta)$ , of rank  $n$ , for  $S$  with  $\mathbf{v} \in M_{1,n}(A)$ ,  $\eta \in M_{n,1}(A)$  and a morphism of monoids  $\mu : \mathcal{X}^* \rightarrow M_{n,n}(A)$  such that

$$S = \sum_{w \in \mathcal{X}^*} (\mathbf{v} \mu(w) \eta) w.$$

A series satisfying one of the conditions of Theorem 2.3 is called *rational*. The set of these series, a  $A$ -module<sup>24</sup>, is denoted by  $A^{\text{rat}} \langle \langle \mathcal{X} \rangle \rangle$  and is closed by  $\{\text{conc}, +, *\}$ . We also have

**Proposition 2.4** (see also [21,40]) *The module  $A^{\text{rat}} \langle \langle \mathcal{X} \rangle \rangle$  (resp.  $A^{\text{rat}} \langle \langle Y \rangle \rangle$ ) is closed by  $\sqcup$  (resp.  $\sqcup$ ). Moreover, for  $i = 1, 2$ , let  $R_i \in A^{\text{rat}} \langle \langle \mathcal{X} \rangle \rangle$  and  $(\mathbf{v}_i, \mu_i, \eta_i)$  be its representation of dimension  $n_i$ . Then the linear representation of<sup>25</sup>*

$$R_i^* \text{ is } \left( \begin{pmatrix} 0 & 1 \end{pmatrix}, \left\{ \begin{pmatrix} \mu_i(x) + \eta_i \mathbf{v}_i \mu_i(x) & 0 \\ \mathbf{v}_i \eta_i & 0 \end{pmatrix} \right\}_{x \in \mathcal{X}}, \begin{pmatrix} \eta_i \\ 1 \end{pmatrix} \right),$$


$$\text{that of } R_1 + R_2 \text{ is } \left( \begin{pmatrix} \mathbf{v}_1 & \mathbf{v}_2 \end{pmatrix}, \left\{ \begin{pmatrix} \mu_1(x) & \mathbf{0} \\ \mathbf{0} & \mu_2(x) \end{pmatrix} \right\}_{x \in \mathcal{X}}, \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix} \right),$$

$$\text{that of } R_1 \cdot R_2 \text{ is } \left( \begin{pmatrix} \mathbf{v}_1 & 0 \end{pmatrix}, \left\{ \begin{pmatrix} \mu_1(x) & \eta_1 \mathbf{v}_2 \mu_2(x) \\ 0 & \mu_2(x) \end{pmatrix} \right\}_{x \in \mathcal{X}}, \begin{pmatrix} \eta_1 \mu_2 \eta_2 \\ \eta_2 \end{pmatrix} \right),$$

$$\text{that of } R_1 \sqcup R_2 \text{ is } (\mathbf{v}_1 \otimes \mathbf{v}_2, \{ \mu_1(x) \otimes \mathbf{I}_{n_2} + \mathbf{I}_{n_1} \otimes \mu_2(x) \}_{x \in \mathcal{X}}, \eta_1 \otimes \eta_2),$$

$$\text{that of } R_1 \sqcup R_2 \text{ is } (\mathbf{v}_1 \otimes \mathbf{v}_2, \{ \mu_1(y_k) \otimes \mathbf{I}_{n_2} + \mathbf{I}_{n_1} \otimes \mu_2(y_k) \\ + \sum_{i+j=k} \mu_1(y_i) \otimes \mu_2(y_j) \}_{k \geq 1}, \eta_1 \otimes \eta_2).$$

**Example 2.5** [Identity  $(-t^2 x_0 x_1)^* \sqcup (t^2 x_0 x_1)^* = (-4t^4 x_0^2 x_1^2)^*$ , [34,35]]



$$(-t^2 x_0 x_1)^* \leftrightarrow (\mathbf{v}_2, \{ \mu_2(x_0), \mu_2(x_1) \}, \eta_2) \quad (t^2 x_0 x_1)^* \leftrightarrow (\mathbf{v}_1, \{ \mu_1(x_0), \mu_1(x_1) \}, \eta_1).$$


<sup>23</sup> see [41].

<sup>24</sup> In fact (we will see it) a unital  $A$ -algebra for  $\text{conc}$  and  $\sqcup$ .

<sup>25</sup> The first constructions are already treated in [21,40], only the last one is new.

$$\begin{aligned} \mathbf{v}_1 &= \begin{pmatrix} 1 & 0 \end{pmatrix}, \mu_1(x_0) = \begin{pmatrix} 0 & t \\ 0 & 0 \end{pmatrix}, \mu_1(x_1) = \begin{pmatrix} 0 & 0 \\ t & 0 \end{pmatrix}, \eta_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \\ \mathbf{v}_2 &= \begin{pmatrix} 1 & 0 \end{pmatrix}, \mu_2(x_0) = \begin{pmatrix} 0 & it \\ 0 & 0 \end{pmatrix}, \mu_2(x_1) = \begin{pmatrix} 0 & 0 \\ it & 0 \end{pmatrix}, \eta_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \end{aligned}$$

$$\begin{aligned} (-t^2x_0x_1)^* \sqcup (t^2x_0x_1)^* &\leftrightarrow (\mathbf{v}, \{\mu(x_0), \mu(x_1)\}, \eta) \\ &= (\mathbf{v}_1 \otimes \mathbf{v}_2, \{\mu_1(x_0) \otimes \mathbf{I}_{n_2} + \mathbf{I}_{n_1} \otimes \mu_2(x_0), \\ &\quad \mu_1(x_1) \otimes \mathbf{I}_{n_2} + \mathbf{I}_{n_1} \otimes \mu_2(x_1), \eta_1 \otimes \eta_2\}). \end{aligned}$$



$$\begin{aligned} \mathbf{v} &= \begin{pmatrix} 1 & 0 & 0 & 0 \end{pmatrix}, \\ \mu(x_0) &= \begin{pmatrix} 0 & 0 & t & 0 \\ 0 & 0 & 0 & t \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & it & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & it \\ 0 & 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & it & t & 0 \\ 0 & 0 & 0 & t \\ 0 & 0 & 0 & it \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ \mu(x_1) &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ t & 0 & 0 & 0 \\ 0 & t & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ it & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & it & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ it & 0 & 0 & 0 \\ 0 & t & it & 0 \end{pmatrix}, \quad \eta = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}. \end{aligned}$$

With the notations of Definition 2.2.(iii) and from Theorem 2.3, it follows that

**Proposition 2.6** Suppose  $A$  to be a field  $K$ . We have

- (a) Assertions of Theorem 2.3 are equivalent to  
 (iv) There exists a finite double family of series  $(G_i, D_i)_{i \in F}$  such that<sup>26</sup>

$$\Delta_{\text{conc}}(S) = \sum_{i \in F} G_i \otimes D_i$$

- (b) For  $S \in \mathcal{H}_{\sqcup}^{\circ}(\mathcal{X})$ , since  $A$  is a field then the previous identity is equivalent to

$$\forall P, Q \in \mathcal{H}_{\sqcup}(\mathcal{X}), \langle S \mid PQ \rangle = \sum_{i \in I} \langle G_i \mid P \rangle \langle D_i \mid Q \rangle.$$

Therefore,  $(K^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle, \sqcup, 1_{\mathcal{X}^*}, \Delta_{\text{conc}}, \mathbf{e})$  (resp.  $(K^{\text{rat}}\langle\langle Y \rangle\rangle, \sqcup, 1_{\mathcal{X}^*}, \Delta_{\text{conc}}, \mathbf{e})$ ) is the Sweedler's dual of  $\mathcal{H}_{\sqcup}(\mathcal{X})$  (resp.  $\mathcal{H}_{\sqcup}(Y)$ ).

Now, let us characterize characters of  $(A\langle X \rangle, \text{conc}, 1_{X^*})$ .

**Proposition 2.7 (Kleene stars of the plane)** Let  $R \in A^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle$ ,  $\langle R \mid 1_{\mathcal{X}^*} \rangle = 1_A$ . The following assertions are equivalent


- (i)  $\langle R \mid \bullet \rangle$  realizes a character<sup>27</sup> of  $(A\langle X \rangle, \text{conc}, 1_{X^*})$ .  
 (ii) There is a family of coefficients  $(c_x)_{x \in \mathcal{X}}$  such that  $R = (\sum_{x \in \mathcal{X}} c_x x)^*$ .  
 (iii) The series  $R$  admits a linear representation of dimension one<sup>28</sup>.

Moreover, we have<sup>29</sup>

$$\begin{aligned} (\alpha_0 x_0 + \alpha_1 x_1)^* \sqcup (\beta_0 x_0 + \beta_1 x_1)^* &= ((\alpha_0 + \beta_0)x_0 + (\alpha_1 + \beta_1)x_1)^* \\ \left( \sum_{s \geq 1} a_s y_s \right)^* \sqcup \left( \sum_{s \geq 1} b_s y_s \right)^* &= \left( \sum_{s \geq 1} (a_s + b_s) y_s + \sum_{r, s \geq 1} a_s b_r y_{s+r} \right)^*, \end{aligned}$$

where, for any  $i = 0, 1$  and  $s \geq 1$ ,  $\alpha_i, \beta_i, a_s, b_s \in \mathbb{C}$ .

**Example 2.8** [Identity  $(-t^2 y_2)^* \sqcup (t^2 y_2)^* = (-4t^4 y_4)^*$ , [34,35]]



$$\begin{aligned} (-t^2 y_2)^* &\leftrightarrow (v_2, \mu_2(y_2), \eta_2) & (t^2 y_2)^* &\leftrightarrow (v_1, \mu_1(y_2), \eta_1) & (-t^4 y_4)^* &\leftrightarrow (v, \mu(y_4), \eta) \\ &= (1, -t^2, 1), & &= (1, t^2, 1), & &= (1, -t^4, 1). \end{aligned}$$

<sup>26</sup> See [39] for a way to obtain this finite double family of series  $(G_i, D_i)_{i \in F}$ .

<sup>27</sup> For  $A = K$  being a field, this can be rephrased as “ $R$  is a group like element of  $K^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle$ ”.

<sup>28</sup> The dimension is here (as in [1]) the size of the matrices.

<sup>29</sup> In particular,  $(a_s y_s)^* \sqcup (a_r y_r)^* = (a_s y_s + a_r y_r + a_s a_r y_{s+r})^*$  and  $(a_s y_s)^* \sqcup (-a_s y_s)^* = (-a_s^2 y_{2s})^*$ .

### 3 Triangularity, solvability and rationality

#### 3.1 Syntactically exchangeable rational series

Now, we have to study a special set of series in order to work with the rational series of this class: a series  $S \in A\langle\langle\mathcal{X}\rangle\rangle$  is called *syntactically exchangeable* if and only if it is constant on multi-homogeneous classes, *i.e.*

$$(\forall u, v \in \mathcal{X}^*)((\forall x \in \mathcal{X})(|u|_x = |v|_x)) \Rightarrow \langle S | u \rangle = \langle S | v \rangle. \quad (26)$$

A series  $S \in A\langle\langle\mathcal{X}\rangle\rangle$  is syntactically exchangeable iff it is of the following form

$$S = \sum_{\alpha \in \mathbb{N}(\mathcal{X}), \text{supp}(\alpha) = \{x_1, \dots, x_k\}} s_{\alpha, x_1}^{\alpha(x_1)} \sqcup \dots \sqcup x_k^{\alpha(x_k)}. \quad (27)$$

The set of these series, a shuffle subalgebra of  $A\langle\langle X \rangle\rangle$ , will be denoted  $A_{\text{exc}}^{\text{synt}}\langle\langle\mathcal{X}\rangle\rangle$ .

When  $A$  is a field, the rational and exchangeable series are exactly those who admit a representation with commuting matrices (at least the minimal one is such, see Theorem 3.2 below). We will take this as a definition as, even for rings, this property implies syntactic exchangeability.

**Definition 3.1** A series  $S \in A^{\text{rat}}\langle\langle\mathcal{X}\rangle\rangle$  will be called *rationally exchangeable* if it admits a representation  $(v, \mu, \eta)$  such that  $\{\mu(x)\}_{x \in \mathcal{X}}$  is a set of commuting matrices, the set of these series, a shuffle subalgebra of  $A\langle\langle X \rangle\rangle$ , will be denoted  $A_{\text{exc}}^{\text{rat}}\langle\langle\mathcal{X}\rangle\rangle$ .

**Theorem 3.2 (See [24,39])** Let  $A_{\text{exc}}^{\text{synt}}\langle\langle\mathcal{X}\rangle\rangle$  denote the set of (syntactically) exchangeable series. Then

- (i) In all cases, one has  $A_{\text{exc}}^{\text{rat}}\langle\langle\mathcal{X}\rangle\rangle \subset A^{\text{rat}}\langle\langle\mathcal{X}\rangle\rangle \cap A_{\text{exc}}^{\text{synt}}\langle\langle\mathcal{X}\rangle\rangle$ . The equality holds when  $A$  is a field and

$$A_{\text{exc}}^{\text{rat}}\langle\langle X \rangle\rangle = A^{\text{rat}}\langle\langle x_0 \rangle\rangle \sqcup A^{\text{rat}}\langle\langle x_1 \rangle\rangle = \bigsqcup_{x \in X} A^{\text{rat}}\langle\langle x \rangle\rangle,$$

$$A_{\text{exc}}^{\text{rat}}\langle\langle Y \rangle\rangle \cap A_{\text{fin}}^{\text{rat}}\langle\langle Y \rangle\rangle = \bigcup_{k \geq 0} A^{\text{rat}}\langle\langle y_1 \rangle\rangle \sqcup \dots \sqcup A^{\text{rat}}\langle\langle y_k \rangle\rangle \subsetneq A_{\text{exc}}^{\text{rat}}\langle\langle Y \rangle\rangle,$$

where  $A_{\text{fin}}^{\text{rat}}\langle\langle Y \rangle\rangle = \bigcup_{F \subset_{\text{finite}} Y} A^{\text{rat}}\langle\langle F \rangle\rangle$ , the algebra of series over finite subalphabets<sup>30</sup>.

- (ii) (Kronecker's theorem [1,51]) One has  $A^{\text{rat}}\langle\langle x \rangle\rangle = \{P(1 - xQ)^{-1}\}_{P, Q \in A[x]}$  (for  $x \in \mathcal{X}$ ) and if  $A = K$  is an algebraically closed field of characteristic zero one

<sup>30</sup> The last inclusion is strict as shows the example of the following identity [6]

$$(ty_1 + t^2y_2 + \dots)^* = \lim_{k \rightarrow +\infty} (ty_1 + \dots + t^ky_k)^* = \lim_{k \rightarrow +\infty} (ty_1)^* \sqcup \dots \sqcup (t^ky_k)^* = \bigsqcup_{k \geq 1} (t^ky_k)^*$$

which lives in  $A_{\text{exc}}^{\text{rat}}\langle\langle Y \rangle\rangle$  but not in  $A_{\text{exc}}^{\text{rat}}\langle\langle Y \rangle\rangle \cap A_{\text{fin}}^{\text{rat}}\langle\langle Y \rangle\rangle$ .

also has  $K^{\text{rat}}\langle\langle x \rangle\rangle = \text{span}_K\{(ax)^* \sqcup K\langle x \rangle \mid a \in K\}$ .

- (iii) The rational series  $(\sum_{x \in \mathcal{X}} \alpha_x x)^*$  are conc-characters and any conc-character is of this form.
- (iv) Let us suppose that  $A$  is without zero divisors and let  $(\varphi_i)_{i \in I}$  be a family within  $\widehat{A\mathcal{X}}$  which is  $\mathbb{Z}$ -linearly independent then, the family  $\mathcal{Lyn}(\mathcal{X}) \uplus \{\varphi_i^*\}_{i \in I}$  is algebraically free over  $A$  within  $(A^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle, \sqcup, 1_{\mathcal{X}^*})$ .
- (v) In particular, if  $A$  is a ring without zero divisors  $\{x^*\}_{x \in \mathcal{X}}$  (resp.  $\{y^*\}_{y \in Y}$ ) are algebraically independent over  $(A\langle\mathcal{X}\rangle, \sqcup, 1_{\mathcal{X}^*})$  (resp.  $(A\langle Y \rangle, \sqcup, 1_{Y^*})$ ) within  $(A^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle, \sqcup, 1_{\mathcal{X}^*})$  (resp.  $(A^{\text{rat}}\langle\langle Y \rangle\rangle, \sqcup, 1_{Y^*})$ ).

**Proof.**

- (i) The inclusion is obvious in view of (27). For the equality, it suffices to prove that, when  $A$  is a field, every rational and exchangeable series admits a representation with commuting matrices. This is true of any minimal representation as shows the computation of shifts (see [20,24,39]).

Now, if  $\mathcal{X}$  is finite, as all matrices commute, we have

$$\sum_{w \in \mathcal{X}^*} \mu(w)w = \left( \sum_{x \in \mathcal{X}} \mu(x)x \right)^* = \sqcup_{x \in \mathcal{X}} (\mu(x)x)^*$$

and the result comes from the fact that  $R$  is a linear combination of matrix elements. As regards the second equality, inclusion  $\supset$  is straightforward. We remark that the union  $\bigcup_{k \geq 1} A^{\text{rat}}\langle\langle y_1 \rangle\rangle \sqcup \dots \sqcup A^{\text{rat}}\langle\langle y_k \rangle\rangle$  is directed as these algebras are nested in one another. With this in view, the reverse inclusion comes from the fact that every  $S \in A_{\text{fin}}^{\text{rat}}\langle\langle Y \rangle\rangle$  is a series over a finite alphabet and the result follows from the first equality.

- (ii) Let  $\mathcal{A} = \{P(1-xQ)^{-1}\}_{P,Q \in A[x]}$ . Since  $P(1-xQ)^{-1} = P(xQ)^*$  then it is obvious that  $\mathcal{A} \subset A^{\text{rat}}\langle\langle x \rangle\rangle$ . Next, it is easy to check that  $\mathcal{A}$  contains  $A\langle x \rangle (= A[x])$  and it is closed by  $+$ , conc as, for instance,

$$(1-xQ_1)(1-xQ_2) = (1-x(Q_1+Q_2-xQ_1Q_2)).$$

We also have to prove that  $\mathcal{A}$  is closed for  $*$ . For this to be applied to  $P(1-xQ)^{-1}$ , we must suppose that  $P(0) = 0$  (as, indeed,  $\langle P(1-xQ)^{-1} \mid 1_{x^*} \rangle = P(0)$ ) and, in this case,  $P = xP_1$ . Now

$$\left( \frac{P}{1-xQ} \right)^* = \left( 1 - \frac{P}{1-xQ} \right)^{-1} = \frac{1-xQ}{1-x(Q+P_1)} \in \mathcal{A}.$$

- (iii) Let  $S = (\sum_{x \in \mathcal{X}} \alpha_x x)^*$  and note that  $S = 1 + (\sum_{x \in \mathcal{X}} \alpha_x x)S$ . Then  $\langle S \mid 1_{\mathcal{X}^*} \rangle = 1_A$  and, if  $w = xu$ , we have  $\langle S \mid xu \rangle = \alpha_x \langle S \mid u \rangle$ , then by recurrence on the length,  $\langle S \mid x_1 \dots x_k \rangle = \prod_{i=1}^k \alpha_{x_i}$  which shows that  $S$  is a conc-character. For the converse, we have Schützenberger's reconstruction lemma which says

that, for every series  $S$

$$S = \langle S \mid 1_{\mathcal{X}^*} \rangle \cdot 1_A + \sum_{x \in \mathcal{X}} x \cdot x^{-1} S$$

but, if  $S$  is a conc-character,  $\langle S \mid 1_{\mathcal{X}^*} \rangle = 1$  and  $x^{-1} S = \langle S \mid x \rangle S$ , then the previous expression reads

$$S = 1_A + \left( \sum_{x \in \mathcal{X}} \langle S \mid x \rangle x \right) S$$

this last equality being equivalent to  $S = (\sum_{x \in \mathcal{X}} \langle S \mid x \rangle \cdot x)^*$  proving the claim.

- (iv) As  $(A\langle \mathcal{X} \rangle, \sqcup, 1_{\mathcal{X}^*})$  and  $(A\langle Y \rangle, \sqcup, 1_{Y^*})$  are enveloping algebras, this property is an application of the fact that, on an enveloping  $\mathcal{U}$ , the characters are linearly independent w.r.t. to the convolution algebra  $\mathcal{U}_\infty^*$  (see the general construction and proof in [25] or [27]). Here, this convolution algebra ( $\mathcal{U}_\infty^*$ ) contains the polynomials (is equal in case of finite  $\mathcal{X}$ ). Now, consider a monomial

$$(\varphi_{i_1}^*)^{\sqcup} \alpha_1 \dots (\varphi_{i_n}^*)^{\sqcup} \alpha_n = \left( \sum_{k=1}^n \alpha_{i_k} \varphi_{i_k} \right)^*$$

The  $\mathbb{Z}$ -linear independence of the monomials in  $(\varphi_i)_{i \in I}$  implies that all these monomials are linearly independent over  $A\langle \mathcal{X} \rangle$  which proves algebraic independence of the family  $(\varphi_i)_{i \in I}$ .

To end with, the fact that  $\mathcal{Lyn}(\mathcal{X}) \sqcup \{\varphi_i^*\}_{i \in I}$  is algebraically free comes from Radford theorem  $(A\langle \mathcal{X} \rangle, \sqcup, 1_{\mathcal{X}^*}) \simeq A[\mathcal{Lyn}(\mathcal{X})]$  and the transitivity of polynomial algebras (see [3] ch III.2 Proposition 8).

- (v) Comes directly as an application of the preceding point. □

**Remark 3.3** (Point (ii) of Theorem 3.2 above) Kronecker's theorem which can be rephrased in terms of stars as  $A^{\text{rat}}\langle\langle x \rangle\rangle = \{P(xQ)^*\}_{P, Q \in A[x]}$  holds for every ring and is therefore characteristic free, unlike the shuffle version requiring algebraic closure and denominators.

### 3.2 Exchangeable rational series and their linear representations

As examples, one can consider the following forms  $(F_0)$ ,  $(F_1)$  and  $(F_2)$  of rational series in  $A^{\text{rat}}\langle\langle X \rangle\rangle$  [33, 39]:

$$\begin{aligned} (F_0) & E_1 x_{i_1} \dots E_j x_{i_j} E_{j+1}, \text{ where } x_{i_1}, \dots, x_{i_j} \in X, E_1, \dots, E_j \in A^{\text{rat}}\langle\langle x_0 \rangle\rangle, \\ (F_1) & E_1 x_{i_1} \dots E_j x_{i_j} E_{j+1}, \text{ where } x_{i_1}, \dots, x_{i_j} \in X, E_1, \dots, E_j \in A^{\text{rat}}\langle\langle x_1 \rangle\rangle, \\ (F_2) & E_1 x_{i_1} \dots E_j x_{i_j} E_{j+1}, \text{ where } x_{i_1}, \dots, x_{i_j} \in X, E_1, \dots, E_j \in A_{\text{exc}}^{\text{rat}}\langle\langle X \rangle\rangle. \end{aligned}$$

Using linear representations, we also have

**Theorem 3.4 (Triangular sub bialgebras of  $(A^{\text{rat}}\langle\langle\mathcal{X}\rangle\rangle, \sqcup, 1_{X^*}, \Delta_{\text{conc}}, \epsilon)$ , [39])**  
Let  $\rho = (\nu, \mu, \eta)$  a representation of  $R \in A^{\text{rat}}\langle\langle\mathcal{X}\rangle\rangle$ . Then

- (i) If the matrices  $\{\mu(x)\}_{x \in \mathcal{X}}$  commute between themselves and if the alphabet is finite, every rational exchangeable series decomposes as

$$R = \sum_{i=1}^n \sqcup_{x \in \mathcal{X}} R_x^{(i)} \text{ with } R_x^{(i)} \in A^{\text{rat}}\langle\langle x \rangle\rangle.$$

- (ii) If  $\mathcal{L}$  consists of upper-triangular matrices then  $R \in A_{\text{exc}}^{\text{rat}}\langle\langle\mathcal{X}\rangle\rangle \sqcup A\langle\mathcal{X}\rangle$ .  
(iii) For any  $x \in \mathcal{X}$ , letting  $M(x) := \mu(x)x$  and then extending, in the obvious way, this representation to  $A\langle\langle\mathcal{X}\rangle\rangle$  by  $M(S) = \sum_{w \in \mathcal{X}^*} \langle S | w \rangle \mu(w)w$ , we have

$$R = \nu M(\mathcal{X}^*) \eta.$$

Moreover, we have

- (a) If  $\{\mu(x)\}_{x \in \mathcal{X}}$  are upper-triangular then  $M(\mathcal{X}) = D(\mathcal{X}) + N(\mathcal{X})$ , where  $D(\mathcal{X})$  and  $N(\mathcal{X})$  are diagonal and strictly upper-triangular letter matrices, respectively, such that<sup>31</sup>

$$M(\mathcal{X}^*) = ((D(\mathcal{X}^*)N(\mathcal{X}))^* D(\mathcal{X}^*)).$$

- (b) We get<sup>32</sup> (for  $\mathcal{X} = X$ )

$$M((x_0 + x_1)^*) = (M(x_1^*)M(x_0))^* M(x_1^*) = (M(x_0^*)M(x_1))^* M(x_0^*)$$

and the modules generated by the families  $(F_0)$ ,  $(F_1)$  and  $(F_2)$  are closed by  $\text{conc}$ ,  $\sqcup$  (and coproducts if  $A = K$  is a field). From this, it follows that  $R$  is a linear combination of expressions in the form  $(F_0)$  (resp.  $(F_1)$ ) if  $M(x_1^*)M(x_0)$  (resp.  $M(x_0^*)M(x_1)$ ) is strictly upper-triangular.

- (c) If  $A$  is a  $\mathbb{Q}$ -algebra then

$$M(\mathcal{X}^*) = \prod_{l \in \mathcal{L}_{\text{yn}} \mathcal{X}} e^{\mathcal{S}_l \mu(P_l)}.$$

**Remark 3.5** (i) The point (i) of Theorem 3.4 is no longer true for an infinite alphabet as shows the example of the series  $S = \sum_{k \geq 1} y_k$  in  $A^{\text{rat}}\langle\langle Y \rangle\rangle$ .

- (ii) On a general ring it can happen that  $R$  is exchangeable,  $\rho$  minimal and nevertheless  $\mathcal{L}$  is noncommutative, as shows the case of  $A = \mathbb{Q}[x, t]/t^3\mathbb{Q}[x, t]$  and

$$X = \{a, b\}, \mu(a) = t \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}, \mu(b) = t \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}, \nu = \begin{pmatrix} 1 & 1 \end{pmatrix}, \eta = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

<sup>31</sup> by Lazard factorization [43,50].

<sup>32</sup> *idem*.

With these data,  $R = 2 + (xt + 2t)(a + b) + (x^2t^2 + 2xt^2 + 2t^2)(ab + ba)$  which is an exchangeable polynomial but

$$\mu(a)\mu(b) = \begin{pmatrix} t^2 & xt^2 \\ xt^2 & x^2t^2 + t^2 \end{pmatrix}, \mu(b)\mu(a) = \begin{pmatrix} x^2t^2 + t^2 & xt^2 \\ xt^2 & t^2 \end{pmatrix}$$

Now the representation is minimal because if it were of dimension 1,  $\frac{1}{2}R$  would be a conc-character, which is not the case. Otherwise, if it were of dimension 0,  $R$  would be zero.

In order to establish Theorem 3.7 below, we will use the following

**Lemma 3.6** *Let  $(\mathbf{v}, \tau, \eta)$  a representation of  $S$  of dimension  $r$  such that, for all  $x \in \mathcal{X}$ ,  $(\tau(x) - c(x)I_r)$  is strictly upper triangular, then  $S \in K_{\text{exc}}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle \sqcup K\langle \mathcal{X} \rangle$ .*

**Proof.** Let  $(e_i)_{1 \leq i \leq r}$  be the canonical basis of  $K^{1 \times r}$ . We construct the representations  $\rho_1 = (\mathbf{v}, (x \mapsto \tau(x) - c(x)I_r), \eta)$ ,  $\rho_2 = (e_1, (x \mapsto c(x)I_r), e_1^*)$  of  $S_1$  and  $S_2$  and remark that  $S_1 \sqcup S_2$  admits the representation

$$\rho_3 = (\mathbf{v} \otimes e_1, ((\tau(x) - c(x)I_r) \otimes I_r + I_r \otimes c(x)I_r)_{x \in \mathcal{X}}, \eta \otimes e_1^*)$$

as  $I_r \otimes c(x)I_r = c(x)I_r \otimes I_r$ ,  $\rho_3$  is, in fact,  $(\mathbf{v} \otimes e_1, (\tau(x) \otimes I_r)_{x \in \mathcal{X}}, \eta \otimes e_1^*)$  which represents  $S$ , the result now comes from the fact that  $S_1 \in K\langle \mathcal{X} \rangle$  and  $S_2 = (\sum_{x \in \mathcal{X}} c(x)x)^* \in K_{\text{exc}}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle$ .  $\square$

We first begin by properties essentially true over algebraically closed fields.

**Theorem 3.7 (Triangular sub bialgebras of  $(K^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle, \sqcup, 1_{X^*}, \Delta_{\text{conc}}, \mathbf{e})$ , [39])**  
*We suppose that  $K$  is an algebraically closed field and that  $\rho = (\mathbf{v}, \mu, \eta)$  is a linear representation of  $R \in K^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle$  of minimal dimension  $n$ , we note  $\mathcal{L} = \mathcal{L}(\mu) \subset K^{n \times n}$  the Lie algebra generated by the matrices  $(\mu(x))_{x \in \mathcal{X}}$ . Then*

- (i)  $\mathcal{L}$  is commutative iff  $R \in K_{\text{exc}}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle$ ,
- (ii)  $\mathcal{L}$  is nilpotent iff  $R \in K_{\text{exc}}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle \sqcup K\langle \mathcal{X} \rangle$ ,
- (iii)  $\mathcal{L}$  is solvable iff  $R$  is a linear combination of expressions in the form  $(F_2)$ .

Moreover, denoting  $K_{\text{nil}}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle$  (resp.  $K_{\text{sol}}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle$ ), the set of rational series such that  $\mathcal{L}(\mu)$  is nilpotent (resp. solvable), we get a tower of sub Hopf algebras of the Sweedler's dual,  $K_{\text{nil}}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle \subset K_{\text{sol}}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle \subset \mathcal{H}_{\sqcup}^{\circ}(\mathcal{X})$ .

**Proof.**

- (i) Let us remark that, for  $x, y \in \mathcal{X}$ ,  $p, s \in \mathcal{X}^*$ , we have  $\langle R \mid pxys \rangle = \langle R \mid pyxs \rangle$  which is due to the commutation of matrices. Conversely, since  $\rho$  is minimal then there is  $P_i, Q_i \in K\langle \mathcal{X} \rangle$ ,  $i = 1 \dots n$  such that (see [1,20,48])

$$\forall u \in \mathcal{X}^*, \mu(u) = (\langle P_i \triangleright R \triangleleft Q_i \mid u \rangle)_{1 \leq i, j \leq n} = (\langle R \mid Q_i u P_i \rangle)_{1 \leq i, j \leq n}.$$

Now, for  $x, y \in \mathcal{X}$ , we have

$$\mu(xy) = (\langle R \mid Q_i x y P_i \rangle)_{1 \leq i, j \leq n} \stackrel{*}{=} (\langle R \mid Q_i y x P_i \rangle)_{1 \leq i, j \leq n} = \mu(yx)$$

equality  $\stackrel{*}{=}$  being due to exchangeability.

- (ii) Let us consider  $K^n$  as the space of the representation of  $\mathcal{L}$  given by  $\mu$ . Let  $K^n = \bigoplus_{j=1}^m V_j$  be a decomposition of  $K^n$  into indecomposable  $\mathcal{L}$ -modules (see [16], Theorem 1.3.19 where it is done for  $ch(K) = 0$ , or [5] Chapter VII §1 Propoposition 9 for arbitrary characteristic), we know that each  $V_j$  is a  $\mathcal{L}$ -module and that the action of  $\mathcal{L}$  is triangularizable with constant diagonals inside each sector  $V_j$ . Thus, it is an invertible matrix  $P \in GL(n, K)$  such that

$$\forall x \in \mathcal{X}, P\mu(x)P^{-1} = \text{blockdiag}(T_1, T_2, \dots, T_k) = \begin{pmatrix} T_1 & 0 & 0 & \dots & 0 \\ 0 & T_2 & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & T_k \end{pmatrix}$$

where the  $T_j$  are upper triangular matrices with scalar diagonal *i.e.* is of the form  $T_j(x) = \lambda(x)I + N(x)$  where  $N(x)$  is strictly upper-triangular<sup>33</sup>. Set  $d_j$  to be the dimension of  $T_j$  (so that  $n = \sum_{j=1}^m d_j$ ), partitioning  $vP^{-1} = v'$  (resp.  $P\eta = \eta'$ ) with these dimensions we get blocks so that each  $(v'_j, T_j, \eta'_j)$  is the representation of a series  $R_j$  and  $R = \sum_{j=1}^m R_j$ . It suffices then to prove that, for all  $j$ ,  $R_j \in K_{\text{exc}}^{\text{rat}} \langle\langle \mathcal{X} \rangle\rangle \sqcup K \langle \mathcal{X} \rangle$ . This is a consequence of Lemma 3.6.

Conversely, if  $\rho_i = (v_i, \tau_i, \eta_i), i = 1, 2$ , are two representations then  $[\tau_1(x) \otimes I_r + I_r \otimes \tau_2(x), \tau_1(y) \otimes I_r + I_r \otimes \tau_2(y)] = [\tau_1(x), \tau_1(y)] \otimes I_r + I_r \otimes [\tau_2(x), \tau_2(y)]$  and a similar formula holds for  $m$ -fold brackets (Dynkin combs), so that if  $\mathcal{L}(\tau_i)$ 's are nilpotent, the Lie algebra  $\mathcal{L}(\tau_1 \otimes I_r + I_r \otimes \tau_2)$  is also nilpotent. The point here comes from the fact that series in  $K_{\text{exc}}^{\text{rat}} \langle\langle \mathcal{X} \rangle\rangle$  as well as in  $K \langle \mathcal{X} \rangle$  admit nilpotent representations, so, let  $(\alpha, \tau, \beta)$  such a representation and  $(\alpha', \tau', \beta')$  its minimal quotient (obtained by minimization, see [1]), then  $\mathcal{L}(\tau')$  is nilpotent as a quotient of  $\mathcal{L}(\tau)$ . Now two minimal representations being isomorphic,  $\mathcal{L}(\mu)$  is isomorphic to  $\mathcal{L}(\tau)$  and then it is nilpotent.

- (iii) As  $\mathcal{L}$  is solvable and  $K$  algebraically closed, using Lie's theorem, we can find a conjugate form of  $\rho = (v, \mu, \eta)$  such that the matrices  $\mu(x)$  are upper-triangular. Since this form also represents  $R$ , letting  $D(\mathcal{X})$  (resp.  $N(\mathcal{X})$ ) be

<sup>33</sup> Even, as  $K$  is infinite, there is a global linear form on  $\mathcal{L}$ ,  $\lambda_{in}$  such that, for all  $g \in \mathcal{L}$ ,  $PgP^{-1} - \lambda_{in}(g)I$  is strictly upper-triangular.

the diagonal (rep. strictly upper-triangular) letter matrix such that  $M(\mathcal{X}) = D(\mathcal{X}) + N(\mathcal{X})$  then

$$R = \mathbf{v}M(\mathcal{X}^*)\boldsymbol{\eta} = \mathbf{v}(D(\mathcal{X}^*)N(\mathcal{X}))^*D(\mathcal{X}^*)\boldsymbol{\eta}.$$

Since  $D(\mathcal{X}^*)N(\mathcal{X})$  being nilpotent of order  $n$  then  $(D(\mathcal{X}^*)N(\mathcal{X}))^* = \sum_{j=0}^{n-1} (D(\mathcal{X}^*)N(\mathcal{X}))^j$ . Hence, letting  $\mathcal{S}$  be the vector space generated by forms of type  $(F_2)$  which is closed by concatenation, we have  $D(\mathcal{X}^*)N(\mathcal{X}) \in \mathcal{S}^{n \times n}$  and then  $(D(\mathcal{X}^*)N(\mathcal{X}))^* \in \mathcal{S}^{n \times n}$ . Finally,  $R = \mathbf{v}M(\mathcal{X}^*)\boldsymbol{\eta} \in \mathcal{S}$  which is the claim.

Conversely, as sums and quotients of solvable representations are solvable it suffices to show that a single form of type  $F_2$  admits a solvable representation and end by quotient and isomorphism as in (ii). From Proposition (2.4), we get the fact that, if  $R_i$  admit solvable representations so does  $R_1R_2$ , then the claim follows from the fact that, firstly, single letters admit solvable (even nilpotent) representations and secondly series of  $\sqcup\{K^{\text{rat}}\langle\langle x \rangle\rangle\}_{x \in \mathcal{X}}$  admit solvable representations. Finally, we choose (or construct) a solvable representation of  $R$ , call it  $(\alpha, \tau, \beta)$  and  $(\alpha', \tau', \beta')$  its minimal quotient, then  $\mathcal{L}(\tau')$  is solvable as a quotient of  $\mathcal{L}(\tau)$ . Now two minimal representations being isomorphic,  $\mathcal{L}(\mu)$  is isomorphic to  $\mathcal{L}(\tau)$ , hence solvable.

Moreover and ff.] Comes from the computation of the coproduct by insertion of identity  $\sum_{i=1}^n e_i^* e_i$ .

□

**Remark 3.8** For an example of series  $S$  with solvable representation but such that  $S \notin K_{\text{exc}}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle \sqcup K\langle \mathcal{X} \rangle$ . One can take  $\mathcal{X} = \{a, b\}$  and  $S = a^*b(-a)^*$ .

To end this section (of combinatorial framework), for a need of the proof of Theorem 4.8 below, let us extent the pairing (15) as follows

$$A\langle\langle \mathcal{X} \rangle\rangle \otimes A\langle\langle \mathcal{X} \rangle\rangle \longrightarrow A, \quad (28)$$

$$T \otimes S \longrightarrow \langle T \| S \rangle := \sum_{w \in \mathcal{X}^*} \langle T | w \rangle \langle S | w \rangle. \quad (29)$$

Here, the family  $\sum_{w \in \mathcal{X}^*} \langle T | w \rangle \langle S | w \rangle$  is summable, for a topology on  $A$ . Its sum is denoted by  $\langle T \| S \rangle$  and the set of these series  $S$  is denoted by  $\text{Dom}_{\text{word}}(T)$ . This proof will also use

**Lemma 3.9** For any without zero divisors ring  $A$ , let  $R \in A^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle$  of linear representation  $(\mathbf{v}, \boldsymbol{\mu}, \boldsymbol{\eta})$  of dimension  $n$ . Then any family  $\{R \triangleleft P_i\}_{i=1 \dots m > n}$  is linearly dependent, i.e. there is  $\{\alpha_i\}_{i=1 \dots m}$  in  $A$ , not all zero, such that  $\sum_{i=1}^m \alpha_i (R \triangleleft P_i) = 0$ .

## 4 Towards a noncommutative Picard-Vessiot theory

Let  $(\mathcal{A}, d)$  be a commutative associative differential ring ( $\ker(d) = k$  being a field),  $\mathcal{C}_0$  be a differential subring of  $\mathcal{A}$  ( $d(\mathcal{C}_0) \subset \mathcal{C}_0$ ) which is an integral domain containing the field of constants and  $\mathbb{C}\{(g_i)_{i \in I}\}$  be the differential subalgebra of  $\mathcal{A}$  generated by  $(g_i)_{i \in I}$ , i.e. the  $k$ -algebra generated by  $g_i$ 's and their derivatives [49].

### 4.1 Noncommutative differential equations

Let us consider the following differential equation, with homogeneous series of degree 1 as multiplier (a polynomial in the case of finite alphabet).

$$\mathbf{d}S = MS; \langle S \mid 1 \rangle = 1, \text{ where } M = \sum_{x \in \mathcal{X}} u_x x \in \mathcal{C}_0 \langle\langle \mathcal{X} \rangle\rangle \quad (30)$$

**Example 4.1** [Drinfel'd equation]  $X = \{x_0, x_1\}$  and  $\Omega = \mathbb{C} \setminus (]-\infty, 0] \cup [1, +\infty[)$ .

$$(KZ_3) \quad \mathbf{d}S = (x_0 u_{x_0} + x_1 u_{x_1})S \text{ with } u_{x_0}(z) = z^{-1}, u_{x_1}(z) = (1-z)^{-1}.$$

This equation, with asymptotic initial conditions, was introduced in [17,18] and a complete study was presented in [39] (solutions and special values).

**Example 4.2**  $Y = \{y_i\}_{i \geq 1}$  and  $\Omega = \{z \in \mathbb{C} \mid |z| < 1\}$ .

$$\mathbf{d}S = \left( \sum_{i \geq 1} y_i u_{y_i} \right) S \text{ with } u_{y_i}(z) = \partial \ell_i(z).$$

where, denoting  $\gamma$  the Euler's constant and  $\zeta(s)$  the Riemann zeta function,

$$\ell_1(z) := \gamma z - \sum_{k \geq 2} \zeta(k) \frac{(-z)^k}{k} \text{ and for } r \geq 2, \ell_r(z) := - \sum_{k \geq 1} \zeta(kr) \frac{(-z^r)^k}{k}.$$

This equation was introduced in [9] to study the independence of a family of eulorian functions.

Let us also recall the following useful result for proving Theorem 4.8 bellow.

**Proposition 4.3** ([34,35,37]) *Let  $S \in \mathcal{A} \langle\langle \mathcal{X} \rangle\rangle$  be solution of (30). Then  $S$  satisfies the differential equations  $\mathbf{d}^l S = Q_l S$ , for  $l \geq 0$ , where  $Q_l \in \mathbb{C}\{(u_i)_{i \geq 0}\} \langle\langle \mathcal{X} \rangle\rangle$  satisfying the recursion  $Q_0 = 1$  and  $Q_l = Q_{l-1} M + \mathbf{d}Q_{l-1}$ .*

*More explicitly,  $Q_l$  can be computed as follows (suming over words  $w = x_{i_1} \dots x_{i_l}$  and derivation multi-indices  $\mathbf{r} = (r_1, \dots, r_l)$  of degree  $\text{deg } \mathbf{r} = |w| = l$  and of weight  $\text{wgt } \mathbf{r} = l + r_1 + \dots + r_l$ )*

$$Q_l = \sum_{\substack{\text{wgt } \mathbf{r} = l \\ w \in \mathcal{X}^{\text{deg } \mathbf{r}}}} \prod_{l=1}^{\text{deg } \mathbf{r}} \binom{\sum_{j=1}^l r_j + j - 1}{r_l} \tau_{\mathbf{r}}(w) \text{ and } \begin{cases} \tau_{\mathbf{r}}(w) = \tau_{r_1}(x_{i_1}) \dots \tau_{r_l}(x_{i_l}) = \\ (\partial_z^{r_1} u_{x_{i_1}})_{x_{i_1}} \dots (\partial_z^{r_l} u_{x_{i_l}})_{x_{i_l}}. \end{cases}$$

**Theorem 4.4** Suppose that the  $\mathbb{C}$ -commutative ring  $\mathcal{A}$  is without zero divisors and equipped with a differential operator  $\partial$  such that  $\mathbb{C} = \ker \partial$ .

Let  $S \in \mathcal{A}\langle\langle \mathcal{X} \rangle\rangle$  be a group-like solution of (30), in the following form

$$S = 1_{\mathcal{X}^*} + \sum_{w \in \mathcal{X}^* \mathcal{X}} \langle S | w \rangle w = 1_{\mathcal{X}^*} + \sum_{w \in \mathcal{X}^* \mathcal{X}} \langle S | S_w \rangle P_w = \prod_{l \in \mathcal{L}_{\text{yn}} \mathcal{X}} e^{\langle S | S_l \rangle P_l}.$$

Then

- (i) If  $H \in \mathcal{A}\langle\langle \mathcal{X} \rangle\rangle$  is another group-like solution of (30) then there exists  $C \in \text{Lie}_{\mathcal{A}}\langle\langle \mathcal{X} \rangle\rangle$  such that  $S = He^C$  (and conversely).
- (ii) The following assertions are equivalent
  - (a)  $\{\langle S | w \rangle\}_{w \in \mathcal{X}^*}$  is  $\mathcal{C}_0$ -linearly independent,
  - (b)  $\{\langle S | l \rangle\}_{l \in \mathcal{L}_{\text{yn}} \mathcal{X}}$  is  $\mathcal{C}_0$ -algebraically independent,
  - (c)  $\{\langle S | x \rangle\}_{x \in \mathcal{X}}$  is  $\mathcal{C}_0$ -algebraically independent,
  - (d)  $\{\langle S | x \rangle\}_{x \in \mathcal{X} \cup \{1_{\mathcal{X}^*}\}}$  is  $\mathcal{C}_0$ -linearly independent,
  - (e) The family  $\{u_x\}_{x \in \mathcal{X}}$  is such that, for  $f \in \text{Frac}(\mathcal{C}_0)$  and  $(c_x)_{x \in \mathcal{X}} \in \mathbb{C}^{(\mathcal{X})}$ ,

$$\sum_{x \in \mathcal{X}} c_x u_x = \partial f \implies (\forall x \in \mathcal{X})(c_x = 0).$$

- (f) The family  $(u_x)_{x \in \mathcal{X}}$  is free over  $\mathbb{C}$  and  $\partial \text{Frac}(\mathcal{C}_0) \cap \text{span}_{\mathbb{C}}\{u_x\}_{x \in \mathcal{X}} = \{0\}$ .

**Proof.** [Sketch] The first item has been treated in [35]. The second is a group-like version of the abstract form of Theorem 1 of [15]. It goes as follows

- due to the fact that  $\mathcal{A}$  is without zero divisors, we have the following embeddings  $\mathcal{C}_0 \subset \text{Frac}(\mathcal{C}_0) \subset \text{Frac}(\mathcal{A})$ ,  $\text{Frac}(\mathcal{A})$  is a differential field, and its derivation can still be denoted by  $\partial$  as it induces the previous one on  $\mathcal{A}$ ,
- the same holds for  $\mathcal{A}\langle\langle \mathcal{X} \rangle\rangle \subset \text{Frac}(\mathcal{A})\langle\langle \mathcal{X} \rangle\rangle$  and **d**
- therefore, equation (30) can be transported in  $\text{Frac}(\mathcal{A})\langle\langle \mathcal{X} \rangle\rangle$  and  $M$  satisfies the same condition as previously.
- Equivalence between **a-d** comes from the fact that  $\mathcal{C}_0$  is without zero divisors and then, by denominator chasing, linear independances w.r.t  $\mathcal{C}_0$  and  $\text{Frac}(\mathcal{C}_0)$  are equivalent. In particular, supposing condition **d**, the family  $\{\langle S | x \rangle\}_{x \in \mathcal{X} \cup \{1_{\mathcal{X}^*}\}}$  (basic triangle) is  $\text{Frac}(\mathcal{C}_0)$ -linearly independent which imply, by the Theorem 1 of [15], condition **e**,
- still by Theorem 1 of [15], **e** is equivalent to **f**, implying that  $\{\langle S | w \rangle\}_{w \in \mathcal{X}^*}$  is  $\text{Frac}(\mathcal{C}_0)$ -linearly independent which induces  $\mathcal{C}_0$ -linear independence (i.e. **a**).  $\square$

Now, let us go back to notations of Section 1 and equip the differential rings of

- (i) analytical functions over a simply connected domain  $\Omega$ ,  $(\mathcal{H}(\Omega), \partial)$ , with the topology of compact convergence (CC),
- (ii) formal series over  $\mathcal{X}$  and with coefficients in  $\mathcal{H}(\Omega)$ ,  $(\mathcal{H}(\Omega) \langle\langle \mathcal{X} \rangle\rangle, \mathbf{d})$ , with the ultrametric distance defined by<sup>34</sup>  $\delta(S, T) = 2^{-\varpi(S-T)}$ .

Let us also consider again the Chen series of the differential forms  $(\omega_i)_{i \geq 1}$  defined by the inputs  $\omega_i = u_{x_i} dz$  along a path  $z_0 \rightsquigarrow z$  on  $\Omega$ . By (17), it follows that

$$C_{z_0 \rightsquigarrow z} = \sum_{w \in \mathcal{X}^*} \alpha_{z_0}^z(w) w = (\alpha_{z_0}^z \otimes \text{Id}) \mathcal{D} \mathcal{X} = \prod_{l \in \mathcal{L}_{\text{yn}} \mathcal{X}} e^{\alpha_{z_0}^z(S_l) P_l}. \quad (31)$$

This series satisfies (30) and is obtained as limit, for the topology of (discrete) point-wise convergence, of Picard iteration process initialized at  $\langle C_{z_0 \rightsquigarrow z} | 1_{\mathcal{X}^*} \rangle = 1_{\mathcal{H}(\Omega)}$ .

Let us illustrate Theorem 4.4, with simple examples, for which  $\mathcal{C}_0$  contains  $\mathbb{C} \{ \{ (u_x^{\pm 1})_{x \in \mathcal{X}} \} \} = \mathbb{C} [u_x^{\pm 1}, \partial^i u_x]_{i \geq 1, x \in \mathcal{X}} \subset \mathcal{A} = (\mathcal{H}(\Omega), \partial)$ . In these examples, we use

**Proposition 4.5 ([33])** For  $\mathcal{X} = \{x\}$ , since  $x^n = x^{\sqcup n} / n!$  then

$$\alpha_{z_0}^z(x^n) = \frac{(\alpha_{z_0}^z(x))^n}{n!}, C_{0 \rightsquigarrow z} = \sum_{n \geq 0} \alpha_{z_0}^z(x^n) x^n = e^{\alpha_{z_0}^z(x)x}, \alpha_0^z(x^*) = e^{\alpha_{z_0}^z(x)}.$$

**Example 4.6** Let us consider two positive cases over  $\mathcal{X} = \{x\}$ .

- (i)  $\Omega = \mathbb{C}, u_x(z) = 1_\Omega, \mathcal{C}_0 = \mathbb{C}$ . Since  $\alpha_0^z(x^n) = z^n / n!$  then, by Proposition 4.5,

$$C_{0 \rightsquigarrow z} = e^{zx} \text{ and } \mathbf{d}C_{0 \rightsquigarrow z} = x C_{0 \rightsquigarrow z}.$$

Moreover,  $\alpha_0^z(x) = z$  which is transcendent over  $\mathcal{C}_0$  and  $\{\alpha_0^z(x^n)\}_{n \geq 0}$  is  $\mathcal{C}_0$ -free. Now, let  $f \in \mathcal{C}_0$  then  $\partial = 0$ . Hence, if  $\partial f = cu_x$  then  $c = 0$ .

- (ii)  $\Omega = \mathbb{C} \setminus ]-\infty, 0], u_x(z) = z^{-1}, \mathcal{C}_0 = \mathbb{C}[z^{\pm 1}] \subset \mathbb{C}(z)$ . Since  $\alpha_1^z(x^n) = \log^n(z) / n!$  then, by Proposition 4.5,

$$C_{1 \rightsquigarrow z} = z^x \text{ and } \mathbf{d}C_{1 \rightsquigarrow z} = z^{-1} x C_{1 \rightsquigarrow z}.$$

Moreover,  $\alpha_1^z(x) = \log(z)$  which is transcendent over  $\mathbb{C}(z)$  then over  $\mathcal{C}_0$  and  $\{\alpha_1^z(x^n)\}_{n \geq 0}$  is  $\mathcal{C}_0$ -free. Now, let  $f \in \mathcal{C}_0$  then  $\partial f \in \text{span}_{\mathbb{C}} \{z^{-n}\}_{n \in \mathbb{Z}, n \neq 1}$ . Hence, if  $\partial f = cu_x$  then  $c = 0$ .

**Example 4.7** Let us consider two negative cases over  $\mathcal{X} = \{x\}$ .

- (i)  $\Omega = \mathbb{C}, u_x(z) = e^z, \mathcal{C}_0 = \mathbb{C}[e^{\pm z}]$ . Since  $\alpha_0^z(x^n) = (e^z - 1)^n / n!$  then, by Proposition 4.5,

$$C_{0 \rightsquigarrow z} = e^{(e^z - 1)x} \text{ and } \mathbf{d}C_{0 \rightsquigarrow z} = e^z x C_{0 \rightsquigarrow z}.$$

<sup>34</sup>  $\forall S \in \mathcal{H}(\Omega) \langle\langle \mathcal{X} \rangle\rangle$ , if  $S = 0$  then  $\varpi(S) = -\infty$  else  $\min_{w \in \text{supp}(S)} \{|w| \text{ or } (w)\}$  [1].

Moreover,  $\alpha_0^z(x) = e^z - 1$  which is not transcendent over  $\mathcal{C}_0$  and  $\{\alpha_0^z(x^n)\}_{n \geq 0}$  is not  $\mathcal{C}_0$ -free. If  $f(z) = ce^z \in \mathcal{C}_0$  ( $c \neq 0$ ) then  $\partial f(z) = ce^z = cu_x(z)$ .

- (ii)  $\Omega = \mathbb{C} \setminus ]-\infty, 0]$ ,  $u_x(z) = z^a$ ,  $a \in \mathbb{C} \setminus \mathbb{Q}$ ,  $\mathcal{C}_0 = \mathbb{C}\{\{z, z^{\pm a}\}\} = \text{span}_{\mathbb{C}}\{z^{ka+l}\}_{k,l \in \mathbb{Z}}$ . Since  $\alpha_0^z(x^n) = (a+1)^{-n} z^{n(a+1)}/n!$  then, by Proposition 4.5,

$$\mathbf{C}_{0 \rightsquigarrow z} = e^{(a+1)^{-1} z^{(a+1)} x} \text{ and } \mathbf{d}\mathbf{C}_{0 \rightsquigarrow z} = z^a x \mathbf{C}_{0 \rightsquigarrow z}.$$

Moreover,  $\alpha_0^z(x) = z^{a+1}/(a+1)$  which is not transcendent over  $\mathcal{C}_0$  and  $\{\alpha_0^z(x^n)\}_{n \geq 0}$  is not  $\mathcal{C}_0$ -free. If  $f(z) = cz^{a+1}/(a+1) \in \mathcal{C}_0$  ( $c \neq 0$ ) then  $\partial f(z) = cz^a = cu_x(z)$ .

#### 4.2 First step of a noncommutative Picard-Vessiot theory

Let us recall that the vector space of solutions of (30) is a free  $(\mathbb{C}\langle\langle \mathcal{X} \rangle\rangle)$ -right module of dimension one<sup>35</sup> generated by  $C_{z_0 \rightsquigarrow z}$  [35]. Hence, by Lemma 4.4, we have common traits with the ordinary case of first order differential equations,

- (i) the differential Galois group of (30) + group-like is the Hausdorff group  $\{e^{\mathcal{C}}\}_{\mathcal{C} \in \mathcal{L}ie_{\mathbb{C}.1, \mathcal{H}(\Omega)}\langle\langle \mathcal{X} \rangle\rangle}$  (group of characters of  $\mathcal{H}_{\sqcup}(\mathcal{X})$ ).
- (ii) the PV extension related to (30) is  $\mathcal{C}\langle\langle \mathcal{X} \rangle\rangle(\mathbf{C}_{z_0 \rightsquigarrow z})$ , where  $\mathcal{C} \subset \mathcal{A} = (\mathcal{H}(\Omega), \partial)$  such that  $\text{Const}(\mathcal{C}\langle\langle \mathcal{X} \rangle\rangle) = \ker \mathbf{d} = \mathbb{C}.1_{\mathcal{H}(\Omega)}\langle\langle \mathcal{X} \rangle\rangle$ .

**Theorem 4.8** ([34,35,37]) *Let  $R \in \mathbb{C}.1_{\mathcal{H}(\Omega)}^{\text{rat}}\langle\langle \mathcal{X} \rangle\rangle$ . Then, for any path  $z_0 \rightsquigarrow z$  over  $\Omega$ , we have<sup>36</sup>  $R \in \text{Dom}_{\text{word}}(\mathbf{C}_{z_0 \rightsquigarrow z})$  and the output of (1) can be computed by*

$$y(z_0, z) = \alpha_{z_0}^z(R) = \sum_{w \in \mathcal{X}^*} (v\mu(w)\eta) \alpha_{z_0}^z(w) = \langle \mathbf{C}_{z_0 \rightsquigarrow z} \| R \rangle.$$

Now, let  $N$  be the least integer  $n$  such that  $y$  satisfies a (non-trivial) differential equation of order  $N$  (with coefficients in  $\mathcal{C}$ ), the family  $\{\partial^k y\}_{0 \leq k \leq N-1}$  is  $\mathcal{C}$ -linearly independent, i.e.

$$(a_n \partial^N + \dots + a_1 \partial + a_0)y = 0, \text{ with } a_N, \dots, a_0 \in \mathcal{C}.$$

and, from what precedes, we have  $N \leq n = \text{rk}(R)$ .

**Proof.** Due to this strong convergence condition, we have

- (i) for any  $T \in \mathcal{H}(\Omega)\langle\langle \mathcal{X} \rangle\rangle$  and  $P \in \mathcal{H}(\Omega)\langle \mathcal{X} \rangle$ ,  $S \in \text{Dom}_{\text{word}}(T)$ , we have  $S \in \text{Dom}_{\text{word}}(PT)$ ,  $S \triangleleft P \in \text{Dom}_{\text{word}}(T)$  and  $\langle PT \| S \rangle = \langle T \| S \triangleleft P \rangle$ ,

<sup>35</sup> In fact, we will see that it is the  $\mathbb{C}\langle\langle \mathcal{X} \rangle\rangle$ -right module  $C_{z_0 \rightsquigarrow z}. \mathbb{C}.1_{\mathcal{H}(\Omega)}\langle\langle \mathcal{X} \rangle\rangle$ .

<sup>36</sup> Once  $(z, z_0)$  is fixed on  $\Omega$ ,  $\text{Dom}_{\text{word}}(\mathbf{C}_{z_0 \rightsquigarrow z})$  is the subset of  $A\langle\langle \mathcal{X} \rangle\rangle$  of series  $R$  such that  $\sum_{n \geq 0} \alpha_{z_0}^z(R_n)$  is convergent for the standard topology, where  $R_n = \sum_{|w|=n} \langle R | w \rangle w$  is a homogeneous component (we need to check that this series is convergent via *majoration morphisms* [34,35,37]).

(ii) from the continuity of  $\partial$ , for any  $T \in \mathcal{H}(\Omega) \langle\langle \mathcal{X} \rangle\rangle$  and  $S \in \text{Dom}_{\text{word}}(T)$ , we have  $\partial(\langle T \| S \rangle) = \langle \mathbf{d}T \| S \rangle + \langle T \| \mathbf{d}S \rangle$ .

Now, let  $(\mathbf{v}, \mu, \eta)$  be a representation of  $R \in \mathbb{C}.1_{\mathcal{H}(\Omega)}^{\text{rat}} \langle\langle \mathcal{X} \rangle\rangle$  of rank  $n$ . Let us see that the family  $(\langle C_{z_0 \rightsquigarrow z} | w \rangle \langle R | w \rangle)_{w \in \mathcal{X}^*}$  is summable in  $\mathcal{H}(\Omega)$ . Indeed, since the matrix norm is multiplicative then, for any  $w \in \mathcal{X}^*$  and  $B_1 > 0$ , we have<sup>37</sup>

$$\|\mu(w)\| \leq B_1^{|w|} \text{ and } |\mathbf{v}\mu(w)\eta| \leq k_1 \|\mathbf{v}\|_r \|\mu(w)\| \|\eta\|_c.$$

The Chen series  $C_{z_0 \rightsquigarrow z}$  is exponentially bounded from above<sup>38</sup>, *i.e.* for all compact  $\kappa \subset \Omega$ , there is  $k_2, B_2 > 0$  such that<sup>39</sup> [34,35,37]

$$\forall w \in \mathcal{X}^*, \|\langle C_{z_0 \rightsquigarrow z} | w \rangle\|_{\kappa} \leq k_2 B_2^{|w|} / |w|!$$

Hence, choosing a compact  $\kappa \subset \Omega$ , we obtain

$$\begin{aligned} \sum_{w \in \mathcal{X}^*} \|\langle C_{z_0 \rightsquigarrow z} | w \rangle \langle R | w \rangle\|_{\kappa} &\leq \sum_{w \in \mathcal{X}^*} \|\langle C_{z_0 \rightsquigarrow z} | w \rangle\|_{\kappa} |\langle R | w \rangle| \\ &\leq \sum_{w \in \mathcal{X}^*} k_2 \frac{B_2^{|w|}}{|w|!} (k_1 \|\mathbf{v}\|_r B_1^{|w|} \|\eta\|_c) < +\infty. \end{aligned}$$

Since  $y = y(z_0, z) = \alpha_{z_0}^z(R)$  and  $\partial$  is continuous for (CC) then, by Proposition 4.3,

$$\partial^l y(z_0, z) = \langle \mathbf{d}^l C_{z_0 \rightsquigarrow z} \| R \rangle \text{ and for } l \leq n, \mathbf{d}^l C_{z_0 \rightsquigarrow z} = Q_l(z) C_{z_0 \rightsquigarrow z}$$

and then [34,35,37]

$$\partial^l y(z_0, z) = \langle Q_l(z) C_{z_0 \rightsquigarrow z} \| R \rangle = \langle C_{z_0 \rightsquigarrow z} \| R \triangleleft Q_l(z) \rangle.$$

By Lemma 3.9, there is  $\{a_k\}_{k=0, \dots, n}$  in  $\mathcal{C}$ , not all zero, such that  $\sum_{k=0}^n a_k (R \triangleleft Q_k) = 0$  yielding the expected result. This linear independence holds in any module whatever the ring.  $\square$

**Remark 4.9** (i) The rational series in Theorem 4.8 is the generating series of the first order linear differential system,  $\partial q = (u_0 \mu(x_0) + \dots + u_m \mu(x_m))q$ ,  $y = \mathbf{v}q$ , initialized at  $y(z_0) = \eta$ . From [30],  $y(z) = \alpha_{z_0}^z(R)$ . The  $N$ th-order differential equation in Theorem 4.8 is then the result, obtained by eliminating the states  $\{q_i\}_{i=0, \dots, m}$  in this system.

(ii) The converse process is also possible thanks to the *compagnion form*.

(iii) Analogue results for nonlinear equations can be found in [34,35,37].

<sup>37</sup> We choose a matrix norm (*i.e.* multiplicative) on  $\mathbb{C}^{n \times n}$ , denoted  $\|M\|$ , and two norms  $\|\mathbf{v}\|_r, \|\eta\|_c$  on  $\mathbb{C}^{1 \times n}, \mathbb{C}^{n \times 1}$ , respectively, and there is classically  $k_1 > 0$  such that  $|\mathbf{v}.M.\eta| \leq k_1 \|\mathbf{v}\|_r \|M\| \|\eta\|_c$ .

<sup>38</sup> In the references the bounding is finer and adapted as well to infinite alphabet.

<sup>39</sup> For any  $f \in \mathcal{H}(\Omega)$ , we denote  $\|f\|_{\kappa} := \sup_{s \in \kappa} |f(s)|$ .

## 5 Conclusion

In this work, we gave a first step to construct a Picard-Vessiot theory for a class of noncommutative differential equations satisfied by the Chen series  $C_{z_0 \rightsquigarrow z}$  over the alphabet  $\mathcal{X} = \{x_i\}_{i \geq 0}$  (along paths  $z_0 \rightsquigarrow z$  belonging to a simply connected manifold  $\Omega$  and with respect to the differential forms  $(u_i dz)_{i \geq 0}$ ):

- (i) The coefficients of these noncommutative generating series belong to the differential ring  $\mathbb{C}\{\{(u_i)_{i \geq 0}\}\}\{C_{z_0 \rightsquigarrow z} \mid w\}_{w \in \mathcal{X}^*}$  which is closed by integration with respect to  $(u_i dz)_{i \geq 0}$ .
- (ii) The Picard-Vessiot extension of these noncommutative differential equations is the module  $C_{z_0 \rightsquigarrow z} \mathbb{C}1_{\mathcal{H}(\Omega)}$  and the Hausdorff group  $\{e^C\}_{C \in \mathcal{L}ie_C(\langle \mathcal{X} \rangle)}$  plays the rôle of differential Galois group associated with this extension.
- (iii) These differential equations were considered as universal differential equations [10,14,17,18,39], *i.e.* the universality can be seen by replacing each letter by constant matrices (resp. analytic vector field, given in (2)) and then solving a system of linear (resp. nonlinear) differential equations, given in (1).
- (iv) These solutions are obtained as a pairing between the series  $C_{z_0 \rightsquigarrow z}$  and the generating series of finite Hankel (resp. Lie-Hankel) rank [31,29,30,45], for linear (resp. nonlinear) differential equations explained by Remark 4.9
- (v) Via rational series (on noncommutative indeterminates and with coefficients in rings) [1,46] and their non-trivial combinatorial Hopf algebras (Theorems 2.1, 2.3, 3.2, 3.4 and 3.7), we illustrated this theory, still under construction, with the case of linear differential equations with singular regular singularities (Theorem 4.8) thanks to the equation satisfied by the Chen generating series.

This practical study allowed also to treat the noncommutative generating series of multiindexed polylogarithms and harmonic sums and as well as those of their special values (polyzetas). In particular, we proved the existence of well defined infinite sums of these polylogarithms and harmonic sums [9] in order to describe solutions of differential equations (Theorem 4.8).

## References

- [1] J. Berstel, C. Reutenauer. — *Rational series and their languages*, Spr.-Ver., 1988.
- [2] N. Bourbaki. — *General Topology, Ch 1-3*, Hermann (1966)
- [3] N. Bourbaki. — *Algebra I-III*, Springer-Verlag Berlin and Heidelberg GmbH & Co. K; (2nd printing 1989)

- [4] N. Bourbaki. — *Commutative Algebra: Chapters 1-7*, Springer (1998)
- [5] N. Bourbaki. — *Groupes et algèbres de Lie, Ch 7-8*, N. Bourbaki et Springer-Verlag Berlin Heidelberg 2006
- [6] V.C. Bui, G.H.E. Duchamp, Hoang Ngoc Minh, L. Kane, C. Tollu. — *Dual bases for non commutative symmetric and quasi-symmetric functions via monoidal factorization*, Journal of Symbolic Computation, 1. **75**, pp 56–73 (2016).
- [7] V.C. Bui, G.H.E. Duchamp, V. Hoang Ngoc Minh, Q.H. Ngo and C. Tollu. — *(Pure) Transcendence Bases in  $\varphi$ -Deformed Shuffle Bialgebras*, 74ème Sém. Lotharingien de Comb., Haus Schönenberg, Ellwangen (2018).
- [8] V.C. Bui, G.H.E. Duchamp, Hoang Ngoc Minh. — *Structure of Polyzetas and Explicit Representation on Transcendence Bases of Shuffle and Stuffle Algebras*, J. of Sym. Comp., Volume 83, November–December 2017, Pages 93-111.
- [9] V.C. Bui, Hoang Ngoc Minh, Q.H. Ngo. — *Families of eulerian functions involved in regularization of divergent polyzetas*, in preparation.
- [10] P. Cartier. — *Jacobiennes généralisées, monodromie unipotente et intégrales itérées*, Séminaire Bourbaki, 687 (1987), 31–52.
- [11] P. Cartier. — *A primer of Hopf algebras*, in: Cartier P., Moussa P., Julia B., Vanhove P. (eds) *Frontiers in Number Theory, Physics, and Geometry II*, (2007).
- [12] R. Chari & A. Pressley. — *A guide to quantum group*, Cambridge (1994)
- [13] C. Chevalley. — *Fundamental Concepts of Algebra*, Acad. Press, NY, 1956.
- [14] P. Deligne. — *Equations Différentielles à Points Singuliers Réguliers*, Lecture Notes in Math, 163, Springer-Verlag (1970).
- [15] M. Deneufchâtel, G.H.E. Duchamp, Hoang Ngoc Minh, A.I. Solomon. — *Independence of hyperlogarithms over function fields via algebraic combinatorics*, in *Lecture Notes in Computer Science* (2011), Volume 6742/2011, 127-139.
- [16] J. Dixmier. — *Enveloping algebras*, North-Holland Publishing Company (1977)
- [17] V. Drinfel'd– *Quasi-Hopf Algebras*, *Len. Math. J.*, 1, 1419-1457, 1990.
- [18] V. Drinfel'd– *On quasitriangular quasi-hopf algebra and a group closely connected with  $\text{gal}(\bar{q}/q)$* , *Leningrad Math. J.*, 4, 829-860, 1991.
- [19] G. DUCHAMP, D. KROB. — *Computing with P.B.W. in enveloping algebras*, in Jacob G., Lamnabhi-Lagarrigue F. (eds) *Algebraic Computing in Control. Lecture Notes in Control and Information Sciences* (1991), vol 165. Springer, Berlin, Heidelberg.

- [20] G. Duchamp, C. Reutenauer. — *Un critère de rationalité provenant de la géométrie non-commutative*, *Inventiones Mathematicae*, 128, 613-622, (1997)
- [21] G. Duchamp, M. Flouret, É. Laugerotte, J. G. Luque. — *Direct and dual laws for automata with multiplicities*, *Theoretical Computer Science*, 267, 105-120, (2001)
- [22] G. H. E. Duchamp and C. Tollu. — *Sweedler's duals and Schützenberger's calculus*, In *Combinatorics and Physics*, p. 67-78, Amer. Math. Soc. (Contemporary Mathematics, vol. 539), 2011.
- [23] G.H.E. Duchamp, V. Hoang Ngoc Minh, Q.H. Ngo. — *Harmonic sums and polylogarithms at negative multi-indices*, *J. of Sym. Comp.*, Volume 83, November–December 2017, Pages 166-186.
- [24] G.H.E. Duchamp, V. Hoang Ngoc Minh, Q.H. Ngo. — *Kleene stars of the plane, polylogarithms and symmetries*, *Theoretical Computer Science*, 800, p. 52-72, 2019.
- [25] G. Duchamp, D. Grinberg, Hoang Ngoc Minh. — *Bialgebraic generalizations of linear independence of characters*, In preparation.
- [26] GÉRARD H. E. DUCHAMP. — *MathOverflow question #203771: Local coordinates on (infinite dimensional) Lie groups, factorization of Riemann zeta functions*, <https://mathoverflow.net/a/203771>
- [27] GÉRARD H. E. DUCHAMP. — *MathOverflow question #310354: Independence of characters with respect to polynomials*, <https://mathoverflow.net/questions/310354>
- [28] J.Y. Enjalbert, G.H.E. Duchamp, V. Hoang Ngoc Minh, C. Tollu. — *The contrivances of shuffle products and their siblings*, *Discrete Mathematics* 340(9): 2286-2300 (2017).
- [29] M. Fliess, C. Reutenauer. — *Théorie de Picard-Vessiot des systèmes réguliers (ou bilinéaires)*, dans "Outils et modèles mathématiques pour l'automatique, l'analyse de systèmes et le traitement du signal, CNRS-RCP 567 (1980).
- [30] M. Fliess. — *Fonctionnelles causales non linéaires et indéterminées non commutatives*, *Bull. Soc. Math. France*, N°109, 1981, pp. 3-40.
- [31] M. Fliess. — *Réalisation locale des systèmes non linéaires, algèbres de Lie filtrées transitives et séries génératrices*, *Inv. Math.*, t 71, 1983, pp. 521-537.
- [32] C. Hespel. — *Une étude des séries formelles noncommutatives pour l'Approximation et l'Identification des systèmes dynamiques*, thèse docteur d'état, Université Lille (1998).
- [33] V. Hoang Ngoc Minh. — *Summations of Polylogarithms via Evaluation Transform*, in *Math. & Computers in Simulations*, 1336, pp 707-728, 1996.

- [34] V. Hoang Ngoc Minh. — *Finite polyzêtas, Poly-Bernoulli numbers, identities of polyzêtas and noncommutative rational power series*, Proceedings of 4<sup>th</sup> International Conference on Words, pp. 232-250, 2003.
- [35] V. Hoang Ngoc Minh. — *Differential Galois groups and noncommutative generating series of polylogarithms*, in “Automata, Combinatorics and Geometry”. 7th World Multi-conference on Systemics, Cybernetics and Informatics, Florida, 2003.
- [36] V. Hoang Ngoc Minh. — *Algebraic combinatoric aspects of asymptotic analysis of nonlinear dynamical system with singular inputs*. Acta Academiae Aboensis, Ser. B **67**(2), 117-126 (2007)
- [37] V. Hoang Ngoc Minh. — *On a conjecture by Pierre Cartier about a group of associators*, Acta Math. Vietnamica (2013), 38, Issue 3, 339-398.
- [38] V. Hoang Ngoc Minh. — *Structure of polyzetas and Lyndon words*, Vietnamese Math. J. (2013), 41, Issue 4, 409-450.
- [39] V. Hoang Ngoc Minh. — *On the solutions of universal differential equation with three singularities*, in Confluentes Mathematici, Tome 11 (2019) no. 2, p. 25-64.
- [40] G. Jacob. — *Réalisation des systèmes réguliers (ou bilinéaires) et séries génératrices non commutatives*, dans “Outils et modèles mathématiques pour l’automatique, l’analyse de systèmes et le traitement du signal, CNRS-RCP 567 (1980).
- [41] G. Jacob. — *Représentations et substitutions matricielles dans la théorie algébrique des transductions*, thèse d’Etat, Univ. Paris 7, 1975.
- [42] E.R. Kolchin. — *Differential Algebra and Algebraic Groups*, New York: Academic, 1973.
- [43] M. Lothaire. — *Combinatorics on Words*, Encyclopedia of Mathematics and its Applications, Addison-Wesley, 1983.
- [44] R. Ree. — *Lie elements and an algebra associated with shuffles* Ann. Math **68** 210–220, 1958.
- [45] C. Reutenauer. — *The local realisation of generating series of finite Lie rank*. Algebraic and Geometric Methods In Nonlinear Control Theory, 33-43
- [46] C. Reutenauer. — *Free Lie Algebras*, London Math. Soc. Monographs (1993).
- [47] J.F. Ritt. — *Differential Algebra*, New York: AMS, 1950.
- [48] M.-P., Schützenberger. — *On the definition of a family of automata*. Information and Control **4**, 245–270, 1961

- [49] M. Van der Put, M. F. Singer. — *Galois Theory of Linear Differential Equations*, Springer (2003)
- [50] G. Viennot. — *Algèbres de Lie libres et monoïdes libres*, Lecture Notes in Mathematics, Springer-Verlag, 691, 1978.
- [51] A. Zygmund. — *Trigonometric series*, Cambridge University Press 2002