Antimicrobial properties against *Aeromonas hydrophila* and immunostimulant effect on *Clarias gariepinus* of *Piper betle*, *Psidium guajava*, and *Tithonia diversifolia* plants

Nunak Nafiqoh, S. Sukenda, Muhammad Zairin Junior, A. Alimuddin, Angela Maria Lusiastuti, Samira Sarter, Domenico Caruso, Jean-Christophe Avarre

► To cite this version:

Nunak Nafiqoh, S. Sukenda, Muhammad Zairin Junior, A. Alimuddin, Angela Maria Lusiastuti, et al.. Antimicrobial properties against *Aeromonas hydrophila* and immunostimulant effect on *Clarias gariepinus* of *Piper betle*, *Psidium guajava*, and *Tithonia diversifolia* plants. *Aquaculture International*, 2020, 28 (1), pp.1-13. 10.1007/s10499-019-00439-6 . hal-02921646

HAL Id: hal-02921646

<https://hal.science/hal-02921646>

Submitted on 26 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

**Antimicrobial properties against *Aeromonas hydrophila* and immunostimulant effect on
Clarias gariepinus of *Piper betle*, *Psidium guajava* and *Tithonia diversifolia* plants**

Nunak Nafiqoh¹, Sukenda², M. Zairin Jr², Alimudin², Angela Lusiastuti¹, Samira Sarter³,
Domenico Caruso³, Jean-Christophe Avarre^{3*}

¹*Research Institute for Freshwater Aquaculture and Fisheries Extension, Bogor, Indonesia*

²*Bogor Agriculture University, Bogor, Indonesia*

³*ISEM, IRD, CIRAD, CNRS, EPHE, University of Montpellier, Montpellier, France*

***Corresponding author**

Jean-Christophe Avarre

Institut des Sciences de l'Evolution de Montpellier

University of Montpellier, Place Eugène Bataillon, cc065

34095 Montpellier cedex 5, France

+33467143748

jean-christophe.avarre@ird.fr

ORCID 0000-0001-6899-7052

Abstract

Aquaculture of the African catfish *Clarias gariepinus* has rapidly increased in South-East Asia over recent years, which has now become one of the most cultivated species. This culture intensification has resulted in the development of various diseases, among which the Motile Aeromonas Septicaemia (MAS) caused by *Aeromonas hydrophila*. The present study aimed at investigating the potential of three plants, *Piper betle*, *Psidium guajava* and *Tithonia diversifolia*, to prevent this disease using an enriched fish diet during four weeks. Though the minimum inhibitory and bactericidal concentrations of acetone-extracts were relatively high (2-32 mg mL⁻¹ and 8-64 mg mL⁻¹, respectively), all 3 plant powders imbedded in feed significantly reduced fish mortalities following an experimental infection with *A. hydrophila*, as indicated by relative percent survivals (RPS) ranging from 77% to 79%. When used in combination, the 3 plant extracts showed no interaction *in vitro*. Even though they improved RPS in comparison with controls, they were, however, less efficient in reducing mortalities than single plants. None of the plant-enriched diet had a clear effect on blood cell count or on reactive oxygen species (ROS) production (except for *P. betle*, which significantly increased ROS production after infection); however, both individual plants and their combinations significantly limited the drop of haematocrit following infection. Altogether, these results indicate that selection of plants for herbal therapy is complex and might depend on several *in vivo* criteria, and as such should not solely rely upon measurement of their antimicrobial activity.

Key-words: *Clarias*, Motile Aeromonas Septicaemia, plants, antimicrobial properties

1 **Introduction**

2
3 Aquaculture provides nearly half of the fish consumed in the world, and developing countries
4 account for 61 % of the world's traded sea food (Fao 2018). Diseases of aquatic animals are
5 considered as a severe threat to the sustainability of aquaculture, as they jeopardize the efforts
6 and outcomes of millions of small fish farmers throughout the world. Global outbreaks in
7 aquaculture contribute to the loss of several billion US dollars per year (Assefa and Abunna
8 2018).

9 Aquaculture intensification over the world has often been associated with a considerable use
10 of antibiotics, widely recognized for their adverse effects on the environment microbial
11 ecosystems, as well as on public and animal health. Antimicrobial resistance has been
12 reported in both European and tropical aquaculture, not only in pathogenic bacteria, but also
13 in commensal and environmental isolates (Cabello et al. 2013). Moreover, resistant bacteria
14 or their resistance genes may spread among the aquatic communities, which in turn may
15 enter human food chain (Chen et al. 2015).

16 Eco-friendly alternatives for therapeutic and prophylactic purposes in health management of
17 aquatic animals are therefore of high priority, as they meet both environmental and societal
18 demands (a high sanitary quality of seafood products). In this regard, the use of plants may
19 provide an eco-friendly solution for the development of a sustainable health management of
20 fish farms. There is growing scientific evidence about the beneficial effects of diets enriched
21 with plant extracts or natural compounds on fish health and prevention of disease outbreaks
22 (Reverter et al. 2014). The active molecules contained in plants (such as alkaloids,
23 terpenoids, saponins and flavonoids) may provide inhibitory activity against pathogenic
24 microorganisms, without eliciting bacterial resistance as is often the case with antibiotics
25 (Chandra et al. 2017). For these reasons, plants may be considered as a promising alternative
26 to the use of chemo-therapeutants, particularly antibiotics (Wink 2015). However, the
27 effectiveness of plant extracts used as antimicrobial depends on many factors linked to their
28 chemical composition, the hydrophobicity of bacteria tested, the exposure time and bacterial
29 cell concentration (Patra and Mohanta 2014). On the other hand, the use of plant
30 combinations may increase their effectiveness through synergistic effects (Ji et al. 2007).
31 Finally, in addition to their potential antibacterial activity, plant extracts may improve the
32 non-specific immunity of several cultured fish, as well as the digestibility and availability of
33 nutrients, thereby resulting in an increase in feed conversion efficiency and protein synthesis
34 (Talpur and Ikhwanuddin 2013).

Although the use of plants in freshwater aquaculture is widespread in Indonesia, the reasons for their choice by the farmers should be developed further. A comprehensive ethno-botanic survey carried out in West Java revealed that improvement of water quality and disease management were the major drivers (63 % of cases) of plant use (Caruso et al. 2013). This study also highlighted that *Clarias spp.* farming was strongly associated with the use of local plants. In addition to their frequent usage by fish farmers, the three plants investigated in the present study (*Piper betle*, *Psidium guajava* and *Tithonia diversifolia*) display antibacterial activity against *Aeromonas hydrophila* (Caruso et al. 2016; Muniruzzaman and Chowdhury 2004), and may therefore be useful for preventing fish infection by this major pathogen in tropical regions.

Clarias gariepinus was introduced to Indonesia in the early 1980s and its culture quickly expanded to the whole country. Like other members of this genus (e.g. *C. batrachus*), it is easy to breed and can grow in most aquaculture systems; as such, it has become a popular food source to fulfil protein needs in several developing countries, including Indonesia, where it is now the second bred freshwater species with a production of more than 1 M tons y^{-1} (Poumogne 2019). Due to its low oxygen demand, *C. gariepinus* is often cultured in high density, which favors the development of various diseases, among which the Motile Aeromonas Septicaemia (MAS) caused by *Aeromonas hydrophila* and may cause around 70-90% mortality during an outbreak (Hanson et al. 2014). The most common symptoms associated with MAS disease are loss of appetite, sores around the mouth, skin haemorrhages and lesions. As in other countries, this bacterial disease is mostly treated in Indonesia with antibiotics such as penicilline, oxytetracycline or ciprofloxacin (Sukarni et al. 2012).

In order to promote an ecological alternative to antibiotics against MAS in *Clarias* aquaculture, the present study aimed at investigating the antibacterial activities, hematological and immunostimulant effects of three plants widely used by fish farmers: *Piper betle*, *Psidium guajava* and *Tithonia diversifolia*. Leaves of the 3 plants were tested alone and in combination, both for their *in vitro* and *in vivo* activity.

Materials and methods

Plant extractions

The plants were all purchased as leaf powder from the Indonesian Spices and Medicinal Crops Research Institute (ISMERI) and selected due to their wide availability in Indonesia and frequent use as medicinal purpose both for humans and animals. Extracts were obtained

from these plants using a maceration method because of its simplicity and low cost. For maceration, one part of plant leave powder was immersed in 10 parts of acetone (g/v) as the best solvent described by Eloff (1998), incubated for 24 hours at room temperature in an orbital shaker (50 rpm) and then filtered through Whatmann (N°41) paper filters. Filtered solutions were then evaporated by rotary evaporation (Heidolph WB 200) at 40°C and 60 rpm until complete solvent removal. The resulting plant extracts were stored in dark glass bottles at 4°C until use.

MIC and MBC tests

Each of the three plant extracts was tested against the Gram negative *Aeromonas hydrophila* fish pathogenic bacteria to determine its ability to inhibit bacterial growth *in vitro*. The bacterial strain used in this study was isolated from a diseased catfish (*Clarias sp.*) by the fish health management laboratory, Research Institute of Freshwater Aquaculture and Extension Fisheries, West Java, Indonesia. To confirm its pathogenicity, it was reinjected into a healthy *C. gariepinus* and collected after the appearance of clinical signs, twice consecutively. It was then identified as *Aeromonas hydrophila* with the kit API 20 NE (BioMérieux) and archived at -80°C in 20% glycerol. For the present assays, the strain was thawed, passaged 4 times on liquid medium and tested again twice on catfish as described above. It was incubated overnight in Mueller Hinton Broth (MHB) medium (Himedia) at 30°C, such that the optical density at 630 nm was around 0.08 to 0.09, corresponding to 10^7 CFU mL⁻¹. Serial 1:2 dilutions of plant extracts, from 64 mg mL⁻¹ to 0.031 mg mL⁻¹, were prepared in water supplemented with 1% DMSO. Using 96-microwell plate, 20 µL of each diluted plant extract were mixed with 20 µL of the bacterial suspension and 160 µL of MHB, and incubated overnight at 30°C. Controls consisted of either 20 µL of bacterial suspension and 180 µL of MHB (positive control), 200 µL of MHB alone (negative control) or 20 µL of plant extract and 180 µL MHB (extract control as a blank). Each control was tested in triplicate and each dilution was tested in 4 replicates. Optical density was measured after 24h of incubation, and the lowest concentration of plant extract that inhibit the bacterial growth was determined as the minimum inhibitory concentration (MIC). To measure the minimum bactericidal concentration (MBC), full loops of the culture were striked onto MHA plates (Mueller Hinton Agar) and incubated overnight at 30°C. The MBC was determined as the lowest concentration that did not show any colony growth after sub-culturing (Fankam et al. 2015).

Bacterial growth interaction tests

To test a potential synergistic effect between the three plant extracts, two combinations with varying proportions were tested: combination 1 (C1) contained an equal amount of each plant extract (33%), whereas combination 2 (C2) was based on the individual MIC values that were measured for each plant, and contained 19% of *P. guajava*, 5% of *P. betle* and 76% of *T. diversifolia*. MIC and MBC tests for plant extract combinations were determined as describe above. The fractional inhibitory concentration (FIC), defined as the ratio between the MIC of the test sample in combination and the MIC of the sample used alone, were calculated for each plant combination. FIC index, which corresponds to the sum of fractional inhibitory concentrations, was calculated following Shafiei et al. (2016), and determined the interactions among the plant extracts:

$$\sum FIC = (FIC I + FIC II + FIC III)$$

It was defined as synergistic when $\sum FIC \leq 1$, antagonistic when $\sum FIC \geq 4$, or non-interactive when $1 < \sum FIC < 4$ (van Vuuren and Viljoen 2011).

Experimental fish, feeding trials and bacterial challenge

Juvenile fish *Clarias gariepinus* (7±0.6 cm and 3.2±0.8 g) were purchased from local fish farmer, placed in 30 L aquaria (30 fish / aquarium) filled with room temperature water (28-30°C) and treated with NaCl (10 ppm) for 3 days after arrival. They were acclimated to such laboratory conditions for two weeks during which they were fed *ad libitum* with commercial diet (Chaoroen Phokpand Prima Indonesia), and during which their health status was frequently checked by visual inspection. Fish were treated according to Indonesian regulations, and all experiments were carried out under the accreditation SNI 7583:2010. Fish were split into six experimental groups: one group for each individual plant (*P. guajava*, *P. betle* and *T. diversifolia* groups), two groups corresponding to the 2 combinations (C1 and C2 groups) and 1 control group. Each group comprised three replicates consisting of 20 fish reared in a single 30L aquarium.

For the feeding trials, the same commercial feed as above was ground and mixed with plant powder. For single plant treatments, 80 g of leave powder were mixed to 1 kg of grinded pellets supplemented with 1% of carboxymethylcellulose as binder. Three hundred mL of water were added to this mix, which were then repelleted. For C1 and C2 treatments, the same proportions of dry leaves as used in *in vitro* combination tests were applied. Fish of the control group were fed with dry pellets without plant supplementation. Fish from all groups

were fed with the experimental diet for four weeks, twice daily, representing ~5% of biomass. Every 3 days, 40-50% of water was renewed.

***In vivo* challenge test**

For the challenge test, *A. hydrophila* was grown in Tryptic Soy Broth (TSB) medium overnight at 30°C. The actual bacterial concentration was determined by measuring the optical density at 630 nm (as described above) and confirmed by colony counting. The culture was diluted in Phosphate Buffer Saline (PBS) to obtain a concentration of 10⁷ CFU mL⁻¹. One hundred µL of diluted culture were intramuscularly injected into the dorsal part of the fish at the end of the 4-week feeding trial. During the challenge, fish continued to receive the same experimental diet, and mortality was recorded two times per day for 10 days.

Blood sampling procedure

Five fish from each treatment group were sampled at the beginning (day 1) and at the end of the feeding period (day 28) as well as 24 hours post-challenge. Approximately 200 µL of blood were withdrawn from the caudal vein of each individual fish with a 1mL syringe. The withdrawn blood was treated as follow: one drop was smeared onto a microscope slide for leucocyte differentiation, 10µL were injected into a heparinized microhaematocrit capillary tube, whereas the remainder was mixed with Na-citrate (3.8%) as anticoagulant for reactive oxygen species measurement.

Measurement of haematological and innate immunity parameters

Reactive oxygen species (ROS) production was measured by quantifying the reduction of nitroblue tetrazolium into formazan, following the protocol described by Biller-Takahashi et al. (2013), except that blood was incubated at 27°C instead of 25°C. Haematocrit percentage was measured according to Sirimanapong et al. (2014). For leukocyte differentiation, the smeared blood was air dried, stained with Giemsa and observed under microscope with 1000x magnification assisted with immersion oil. One hundred leukocytes were counted and percentage of neutrophils, monocytes and lymphocytes was then calculated according to Adel et al. (2015).

Statistical analyses

Normality and equal variance of data were checked using Shapiro-Wilk test and Brown-Forsythe test, respectively. When both criteria were respected, a one-way Anova followed by a post hoc test (Holm-Sidak method) was applied to evaluate the effect of the different herbal treatments; otherwise, a Kruskal-Wallis one-way Anova on Ranks was used. These analyses were used to study the effect of the different herbal treatments on blood parameters between the last day of feeding trial (day 28, before infection) and 1 day after *A. hydrophila* infection. Comparison before infection and after infection within each treatment group was carried out using a t-test. Mortalities resulting from the experimental infection were compared with Chi square followed by a Kaplan-Meier survival analysis and post hoc analysis using Holm-Sidak method. Differences between treatments were considered significant when $P < 0.05$. Relative Percent Survival (RPS) was calculated according to Amend (1981) as follows:

$$\text{Relative percent survival} = 1 - \frac{(\% \text{ of treated group})}{(\% \text{ of control group})} \times 100.$$

Results

Antimicrobial activity

The minimum inhibitory concentrations and minimum bactericidal concentrations were measured for the three plant extracts against the pathogenic bacteria *A. hydrophila*. Overall, *P. betle* extracts showed the highest ability to inhibit bacterial growth, with MIC and MBC of 2 and 8 mg mL⁻¹, respectively. *P. guajava* and *T. diversifolia* displayed higher MIC (8 and 32 mg mL⁻¹, respectively) and MBC (64 mg mL⁻¹ each) values (Table 1). The FIC index of tested combinations among the 3 plant extracts indicated the absence of antagonistic or synergistic effect between them.

Fish growth during feeding trials

After replacement of the regular feed by the experimental diet, fish from all groups stopped eating. Whereas control fish recovered appetite after 2 days, fish from the other treatment groups started to eat again after 3 days only. Body weight of each fish was measured every 7 days during the feeding trial. After one week of experimental diet, the average weight was comparable in all groups, ranging from 2.90 ± 0.21 g (*P. betle* group) to 3.49 ± 0.17 g (C2 group). After 4 weeks, however, fish from the control group displayed a significantly higher ($P < 0.05$) body weight (7.00 ± 0.57 g) than those from all the other treatment groups ($3.56 \pm$

0.27 g for *P. betle* group to 5.14 ± 0.39 g for *P. guajava* group). Moreover, no visible change in vigour/health was observed in any group following administration of the experimental diet.

Haematological and immune parameters

Since almost no significant difference was found between day 1 and day 28, figures will only illustrate the differences before and after experimental infection.

After 4 weeks of feeding, there were no differences in haematocrit values, except for fish fed with *P. betle*, which had a significantly lower haematocrit than control animals (Anova on Rank $P=0.009$). After infection, haematocrit was significantly lower in survivor fish of control group than in fish fed with *T. diversifolia* leaves (One way Anova, $P=0.03$) (Fig.1). Most notably, infection had no significant effect on the haematocrit of the treated groups, whereas it significantly decreased that of the control group (t-test, $P < 0.001$) (Fig 1).

Likewise, there was no variation in blood cell count after 4 weeks of herbal administration, except in fish fed with *T. divesifolia*, which showed the highest percentage of monocytes; however this difference was significant (one way Anova, $P=0.021$) only compared to group C1 (not shown). Following experimental infection, significant variations in blood cell count were observed in survivors of all groups; notably with a marked neutrophilia, monocytosis and lymphocytopenia (Fig. 2). However, none of these modifications could be ascribed to herbal treatment and no difference was observed with the control group in all three categories of blood cells.

Reactive oxygen production, reflected by NBT test, showed no significant difference after 4 weeks between the different herbal treatments and the control (Anova on Rank, $P=0.161$). Following infection, NBT increased for all treatments, including the control; however, this increase was significantly higher in fish fed with *P. betle* than in the control animals (Fig. 3).

Experimental infection

Clinical signs of all moribund and dying fish consisted of red sores, skin ulcers, internal hemorrhages and hemorrhagic ascites, compliant with an infection by *A. hydrophila*. Data of cumulative mortality and RPS are summarized in Table 2. Chi-square analysis revealed a significant difference between treatments (78.631 with 5 degrees of freedom, $P = < 0.001$). The highest cumulative mortality was observed in the control group (83.6%). All mortalities occurred during the first 7 days post-infection (Fig. 4). Except for fish fed with C1 combination, mortalities were significantly lower in all other herbal treatments. Survival analysis (Fig. 4) showed that best survival rates were obtained in fish fed with individual

plants, and in a lesser extent with C2 combination. No synergistic effect on survival was observed when the three plants were administrated together, compared to single plant treatments.

Discussion

The aim of the present work was to evaluate the effect of three plants against MAS, caused by *Aeromonas hydrophila*, by testing both *in vitro* and *in vivo* approaches in *Clarias gariepinus*. MIC values of the 3 plants were all above 1 mg mL⁻¹. Even though a compound that displays a MIC value higher than 100 µg mL⁻¹ has been reported to have a low antibacterial activity (Kuethe 2010), it is worth noticing that this threshold concerns isolated compounds of plants and might vary depending on the methods used, which are not standardized in the literature for plant extracts. For crude extracts, this threshold might be too high, as attested by our study and other published work (Bussmann et al. 2010). The ability of plant extracts to inhibit bacterial growth indicates that some of their natural compounds have antimicrobial properties. *P. betle* main active compounds belong to phenylpropene class, which includes eugenol, chavicol or chavibetol (Bajpai et al. 2010). In contrast, *Tithonia diversifolia* and *Psidium guajava* share a number of common compounds, such as α-pinene, β-pinene, germacrene D and β-caryophyllene, though at different percentages (Rajan and Thenmozhi 2015; Wanzala et al. 2016). These natural compounds may have various effects against different bacteria, depending on their biochemical characteristics. The common modes of action of plant extracts against bacteria are not specific, and include for instance increased cell permeability, cytotoxicity, gene silencing, cell filamentation, membrane disruption, genomic DNA binding etc. (He et al. 2018; Wink 2015). The high survival rates obtained on treated fish after the bacterial challenge tend to indicate that the *in vitro* properties of a plant are not directly correlated with its *in vivo* effects, as was already reported (Anusha et al. 2014). Under these conditions, selection of active plants should not solely rely upon measurement of their MIC. A recent review highlighted that all antimicrobial tests carried out *in vitro* are difficult to reproduce in clinical trials, and may have a limited predictive value (Mundy et al. 2016). In spite of this, *in vitro* antimicrobial tests remain essential in the evaluation of potential antimicrobial candidates, mainly because they do not require the sacrifice of animals. However, additional information, such as ethnobotanical data or comparative usage in human or veterinary pharmacopeia, is likely to help the discovery of efficacious plants for fish.

Application of combined plants may be considered as an alternative way for preventing diseases. It is indeed acknowledged that the combination of two or more plants will provide a better performance of plants as medicine (Che et al. 2013). For illustration, the use of four herbs in combination significantly increased the survival of red seabream juveniles compared to single herb application (Ji et al. 2007). In the present study, the three individual plant extracts significantly reduced the mortality caused by *A. hydrophila*. However, when used in combination, their efficiency was lower, as indicated by higher RPS values. Interestingly, the two combinations used *in vivo* led to significantly different survival rates, suggesting that the proportion of each active compound in the diet is an important factor to consider. A previous study on the combination of *P. betle* with *P. guajava* and *Andrographis paniculata* showed an ability to prevent MAS infection caused by *A. hydrophila* on *Clarias* sp. when supplemented to fish diet (Wahjuningrum et al. 2007). In this study, the combination 0.25:0.75:1 elicited the best survival rate compared to other combinations. In herbal therapy, complex interactions may appear when plants are formulated together with varying proportions. An example of unwanted interaction between herbs is the presence of tannins. These substances may hinder the absorption of proteins and alkaloids, or induce enzymes such as cytochrome P450, which may in turn accelerate drug catabolism and result in blood levels of active substances that are too low to elicit a therapeutic effect (Williamson 2001). Surprisingly, growth of herbal-treated fish was lower than that of control animals. As already reported (Bulfon et al. 2015), the impact of plants on fish growth may greatly vary. The reduced growth that was observed after herbal treatment could be linked to the presence of antinutritional compounds, or more simply to a loss of feed palatability that resulted in a reduced feed intake. Though additional studies are needed to address this loss of growth, the lower weight of animals fed with plants did not prevent their beneficial effect against *A. hydrophila* infection. These results suggest that such herbal treatments should not be used as long-term diets.

Nitroblue tetrazolium assay, which measures the reactive oxygen species (ROS) produced by phagocytic cells, is a good indicator of phagocytic activity in fish (Biller-Takahashi et al. 2013). An increase in ROS production was observed in all fish groups only after the bacterial challenge, and this increase was significantly higher in animals treated with *P. betle* compared to controls. Concomitantly, the proportion of neutrophils and leucocytes also increased after infection, yet in the same manner for all treatments. However, none of the plant extracts tested here had a direct effect on ROS production before infection, as was observed with date palm extract applied to European seabass (Guardiola et al. 2016) or with

different herbs and spices administered to the tilapia *Oreochromis mossambicus* (Gültepe et al. 2014), and in contrast with the oral administration of *Eclipta alca* to *O. mossambicus*, where ROS production increased after one week post-feeding, although fish were not challenged with any pathogen (Christybapita et al. 2007).

Haemolysis is retained as a key factor of *A. hydrophila* virulence (González-Serrano et al. 2002). As a consequence of infection, a drop of red blood cell counts was observed in the control fish, and in a lesser extent in the C1 group. However, the fact that haematocrit of all treated fish did not significantly change after bacterial infection suggests that the addition of plants in the diet contributed to prevent blood cell lysis during infection. These results are in good agreement with the survival rates measured after infection, and indicate that leave powder of these plants may inhibit, at least partially, the haemolytic activity of *A. hydrophila* or prevent red blood cell from haemolysis.

To conclude, in spite of contrasted MIC values, the 3 species of plants tested in this study showed a high and nearly identical efficiency in reducing the mortality due to an infection with *A. hydrophila*. Although the two tested combinations had no antagonistic interaction *in vitro* and elicited higher survival rates than controls, they were less efficient in reducing mortalities than the individual plants. In order to get better insights into the beneficial use of these plants, additional combinations leading to synergistic interactions should be searched and tested for their ability to increase survival rates.

Acknowledgements

Nunak Nafiqoh benefited from an IRD fellowship under the BEST program. We are thankful to the Indonesian Ministry of Fisheries and Marine Affairs for providing PhD scholarship to Mrs. Nafiqoh as well as experimental facilities. This is publication ISEM-DIVA-XXX.

Compliance with Ethical Standards

The authors declare that they have no conflict of interest.). All experiments involving fish complied with animal welfare and were carried out under the Indonesian accreditation SNI 7583:2010.

References

Adel M, Abedian Amiri A, Zorriehzahra J, Nematolahi A, Esteban Mn (2015) Effects of dietary peppermint (*Mentha piperita*) on growth performance, chemical body composition and hematological and immune parameters of fry Caspian white fish

334 (*Rutilus frisii kutum*) Fish Shellfish Immunol 45:841-847.
 335 <https://doi.org/10.1016/j.fsi.2015.06.010>

336 Amend DF (1981) Potency testing of fish vaccines. International Symposium in Fish
 337 Biologics: Serodiagnostics and Vaccines. Dev Biol Stand 49:447-454

338 Anusha P, Thangaviji V, Velmurugan S, Michaelbabu M, Citarasu T (2014) Protection of
 339 ornamental goldfish *Carassius auratus* against *Aeromonas hydrophila* by treating
 340 *Ixora coccinea* active principles. Fish Shellfish Immunol 36:485-493.
 341 <https://doi.org/10.1016/j.fsi.2013.12.006>

342 Assefa A, Abunna F (2018) Maintenance of fish health in aquaculture: Review of
 343 epidemiological approaches for prevention and control of infectious disease of fish.
 344 Vet Med International 2018:1-10. <https://doi.org/10.1155/2018/5432497>

345 Bajpai V, Sharma D, Kumar B, Madhusudanan KP (2010) Profiling of *Piper betle* Linn.
 346 cultivars by direct analysis in real time mass spectrometric technique. Biomed
 347 Chromatogr 24:1283-1286. <https://doi.org/10.1002/bmc.1437>

348 Biller-Takahashi JD, Takahashi LS, Saita MV, Gimbo RY, Urbinati EC (2013) Leukocytes
 349 respiratory burst activity as indicator of innate immunity of pacu *Piaractus*
 350 *mesopotamicus*. Braz J Biol 73:425-429. [https://doi.org/10.1590/S1519-](https://doi.org/10.1590/S1519-69842013000200026)
 351 [69842013000200026](https://doi.org/10.1590/S1519-69842013000200026)

352 Bulfon C, Volpatti D, Galeotti M (2015) Current research on the use of plant-derived
 353 products in farmed fish. Aquac Res 46:513-551. <https://doi.org/10.1111/are.12238>

354 Bussmann RW et al. (2010) Minimum Inhibitory concentration of medicinal plants used in
 355 Northern Peru as antibacterial remedies. J Ethnopharmacol 132:101-108.
 356 <https://doi.org/10.1016/j.jep.2010.07.0>

357 Cabello FC, Godfrey HP, Tomova A, Ivanova L, Dölz H, Millanao A, Buschmann AH
 358 (2013) Antimicrobial use in aquaculture re-examined: Its relevance to antimicrobial
 359 resistance and to animal and human health. Environ Microbiol 15:1917-1942.
 360 <https://doi.org/10.1111/1462-2920.12134>

361 Caruso D, Lusiastuti AM, Slembrouck J, Komarudin O, Legendre M (2013) Traditional
 362 pharmacopeia in small scale freshwater fish farms in West Java , Indonesia : An
 363 ethnoveterinary approach. Aquaculture 416-417:334-345.
 364 <https://doi.org/10.1016/j.aquaculture.2013.09.048>

365 Caruso D, Lusiastuti AM, Taukhid T, Avarre J-C, Yuhana M, Sarter S (2016) Ethnobotanical
 366 uses and antimicrobial properties of plants in small-scale tropical fish farms: The case

- of Indonesian fish farmers in Java (Indonesia). J World Aquacult Soc.
<https://doi.org/10.1111/jwas.12345>
- Chandra H, Bishnoi P, Yadav A, Patni B, Mishra AP, Nautiyal dAR (2017) Antimicrobial resistance and the alternative resources with special emphasis on plant-based antimicrobials—A Review. Plants 6:11. <https://doi.org/10.3390/plants6020016>
- Che CT, Wang ZJ, Sing M, Chow S, Wai C, Lam K (2013) Herb-herb combination for therapeutic enhancement and advancement: Theory, practice and future perspectives. Molecules 18:5125-5141. <https://doi.org/10.3390/molecules18055125>
- Chen H et al. (2015) Antibiotics in typical marine aquaculture farms surrounding Hailing Island, South China: Occurrence, bioaccumulation and human dietary exposure. Mar Pollut Bull 90:181-187. <https://doi.org/10.1016/j.marpolbul.2014.10.053>
- Christybapita D, Divyagnaneswari M, Dinakaran Michael R (2007) Oral administration of *Eclipta alba* leaf aqueous extract enhances the non-specific immune responses and disease resistance of *Oreochromis mossambicus*. Fish Shellfish Immunol 23:840-852. <https://doi.org/10.1016/j.fsi.2007.03.010>
- Eloff JN (1998) Which extractant should be used for the screening and isolation of antimicrobial components from plants ? J Ethnopharmacol 60:1-8. [https://doi.org/10.1016/S0378-8741\(97\)00123-2](https://doi.org/10.1016/S0378-8741(97)00123-2)
- Fankam AG, Kuate JR, Kuete V (2015) Antibacterial and antibiotic resistance modifying activity of the extracts from *Allanblackia gabonensis*, *Combretum molle* and *Gladiolus quartinianus* against Gram-negative bacteria including multi-drug resistant phenotypes. BMC Complement Altern Med 15:1-12. <https://doi.org/10.1186/s12906-015-0726-0>
- Fao (2018) The state of world fisheries and aquaculture 2018 - Meeting the sustainable development goals. FAO. <http://www.fao.org/documents/card/en/c/I9540EN>. Accessed 1 July 2019
- González-Serrano CJ, Santos JA, Garcí-Lopéz M-L, Otero A (2002) Virulence markers in *Aeromonas hydrophila* and *Aeromonas veronii* biovar *sobria* isolates from freshwater fish and from a diarrhoea case. J Appl Microbiol 93:414-419. <https://doi.org/10.1046/j.1365-2672.2002.01705.x>
- Guardiola FA, Porcino C, Cerezuela R, Cuesta A, Faggio C, Esteban MA (2016) Impact of date palm fruits extracts and probiotic enriched diet on antioxidant status, innate immune response and immune-related gene expression of European seabass

400 (*Dicentrarchus labrax*). Fish Shellfish Immunol 52:298-308.
 401 <https://doi.org/10.1016/j.fsi.2016.03.152>
 402 Gültepe N, Bilen S, Yılmaz S, Güroy D, Aydın S (2014) Effects of herbs and spice on health
 403 status of tilapia (*Oreochromis mossambicus*) challenged with *Streptococcus iniae*.
 404 Acta Vet Brno 83:129-136. <https://doi.org/10.2754/avb201483020000>
 405 Hanson LA, Liles MR, Hossain MJ, Griffin MJ, Hemstreet WG (2014) Motile *Aeromonas*
 406 Septicemia. In: Fish Health Section Blue Book: suggested procedures for the
 407 detection and identification of certain finfish and shellfish pathogens., vol 1.2.9. 2014
 408 edn. AFS-FHS, Bethesda, Maryland, pp 1-11
 409 He T-F, Zhang Z-H, Zeng X-A, Wang L-H, Brennan CS (2018) Determination of membrane
 410 disruption and genomic DNA binding of cinnamaldehyde to *Escherichia coli* by use
 411 of microbiological and spectroscopic techniques. J Photochem Photobiol B-Biol
 412 178:623-630
 413 Ji SC, Takaoka O, Jeong GS, Lee SW, Ishimaru K, Seoka M, Takii K (2007) Dietary
 414 medicinal herbs improve growth and some non-specific immunity of red sea bream
 415 *Pagrus major*. Fish Sci 73:63-69. <https://doi.org/10.1111/j.1444-2906.2007.01302.x>
 416 Kuete V (2010) Potential of Cameroonian plants and derived products against microbial
 417 infections : A review. Planta Med:1479-1491
 418 Mundy L, Pendry B, Rahman M (2016) Antimicrobial resistance and synergy in herbal
 419 medicine. J Herb Med 6:53-58. <https://doi.org/10.1016/j.hermed.2016.03.001>
 420 Muniruzzaman M, Chowdhury MBR (2004) Sensitivity of fish pathogenic bacteria to various
 421 medicinal herbs. Bangl J Vet Med 2:75-84
 422 Patra JK, Mohanta YK (2014) Antimicrobial compounds from mangrove plants: A
 423 pharmaceutical prospective. Chin J Integr Med 20:311-320.
 424 <https://doi.org/10.1007/s11655-014-1747-0>
 425 Poumogne V (2019) Cultured Aquatic Species Information Programme. *Clarias gariepinus*.
 426 FAO. http://www.fao.org/fishery/culturedspecies/Clarias_gariepinus/en. Accessed 1
 427 July 2019
 428 Rajan S, Thenmozhi S (2015) GC-MS analysis of bioactive compounds in *Psidium guajava*
 429 leaves. J Pharmacogn Phytochem 3:162-166
 430 Reverter M, Bontemps N, Lecchini D, Banaigs B, Sasal P (2014) Use of plant extracts in fish
 431 aquaculture as an alternative to chemotherapy: Current status and future perspectives.
 432 Aquaculture 433:50-61. <https://doi.org/10.1016/j.aquaculture.2014.05.048>

- Shafiei Z, Haji Abdul Rahim Z, Philip K, Thurairajah N (2016) Antibacterial and anti-adherence effects of a plant extract mixture (PEM) and its individual constituent extracts (*Psidium sp.*, *Mangifera sp.*, and *Mentha sp.*) on single- and dual-species biofilms. *PeerJ* 4:e2519-e2519. <https://doi.org/10.7717/peerj.2519>
- Sirimanapong W, Thompson KD, Kledmanee K, Thaijongrak P, Collet B, Ooi EL, Adams A (2014) Optimisation and standardisation of functional immune assays for striped catfish (*Pangasianodon hypophthalmus*) to compare their immune response to live and heat killed *Aeromonas hydrophila* as models of infection and vaccination. *Fish Shellfish Immunol* 40:374-383. <https://doi.org/10.1016/j.fsi.2014.07.021>
- Sukarni, Maftuch, Nursya H (2012) Kajian penggunaan ciprofloxacin terhadap histologi insang dan hati ikan botia (*Botia macracanthus*, Bleeker) yang diinfeksi bakteri *Aeromonas hydrophila*. *J Exp Life Sci* 2:6-12
- Talpur AD, Ikhwanuddin M (2013) *Azadirachta indica* (neem) leaf dietary effects on the immunity response and disease resistance of Asian seabass, *Lates calcarifer* challenged with *Vibrio harveyi*. *Fish Shellfish Immunol* 34:254-264. <https://doi.org/10.1016/j.fsi.2012.11.003>
- van Vuuren S, Viljoen A (2011) Plant-based antimicrobial studies – Methods and approaches to study the interaction between natural products. *Planta Med* 77:1168-1182. <https://doi.org/10.1055/s-0030-1250736>
- Wahjuningrum D, Taroni, Angka SL (2007) Efficacy of *Andrographis paniculata*, *Psidium guajava* and *Piper betle* as prevention on Motile Aeromonad Septicaemia infection in African catfish (*Clarias sp.*). *J Akuakultur Indonesia* 6:127-133
- Wanzala W, Osundwa EM, Alwala J, Gakuubi MM (2016) Chemical composition of essential oil of *Tithonia diversifolia* (Hemsl.) A. Gray from the Southern slopes of Mount Elgon in Western Kenya. *IJEPP* 2:72-83
- Williamson EM (2001) Synergy and other interactions in phytomedicines. *Phytomedicine* 8:401-409
- Wink M (2015) Modes of action of herbal medicines and plant secondary metabolites. *Medicines* 2:251-286. <https://doi.org/10.3390/medicines2030251>

Table 1. MIC, MBC and FIC index for the three plant extracts and their combination tested against *A. hydrophila*

	<i>P. betle</i>	<i>P. guajava</i>	<i>T. diversifolia</i>	Σ FIC
MIC (mg mL ⁻¹)	2	8	32	
MBC (mg mL ⁻¹)	8	64	64	
FIC for C1 (mg mL ⁻¹)	2	0.5	0.125	2.625
FIC for C2 (mg mL ⁻¹)	2	0.5	0.125	2.625

Table 2. Cumulative mortality rate, expressed as % $\pm$ CI ($\alpha=0.05$), and relative percent survival (RPS) of *Clarias gariepinus* according to herbal treatments following an experimental infection with *A. hydrophila* (N=60)

Treatment	% mortality	Confidence interval	RPS
<i>P. betle</i>	17.2%	$\pm 9.7\%$	79%
<i>P. guajava</i>	18.2%	$\pm 11.4\%$	78%
<i>T. diversifolia</i>	19.5%	$\pm 12.1\%$	77%
C2	27.9%	$\pm 13.4\%$	67%
C1	52.1%	$\pm 14.1\%$	38%
Control	83.6%	$\pm 9.8\%$	0%

Figure 1. Haematocrit, expressed as Mean % $\pm$ SE, before and after infection (N=5). Asterisks indicate significant differences with the respective control group at the same treatment time, whilst double daggers indicate significant differences before and after experimental infection.

Figure 2. Differential leucocyte count (%) of *Clarias gariepinus* before and after infection. The proportion of lymphocytes (A), neutrophils (B) and monocytes (C), expressed as a Mean $\pm$ SE, amounts 100% (N=5). Except for control animals, each group of fish was fed for 4 weeks prior to infection with a diet enriched with plant extracts. Different letters indicate a statistically significant difference (P<0.05).

Figure 3. Reactive oxygen production (Mean $\pm$ SE) of *Clarias gariepinus* before and after infection (N=5). Except for control animals, each group of fish was fed for 4 weeks prior to infection with a diet enriched with plant extracts. Different letters indicate a statistically significant difference (P<0.05).

Figure 4. Kaplan-Meier Survival Analysis (N=60). Except for control animals, each group of fish was fed for 4 weeks prior to infection with a diet enriched with plant extracts. Different letters indicate a statistically significant difference (P<0.05).

497 Figure 1

498

499

502 Figure 3

503

504

505 Figure 4

506
507