Frogs model man: In vivo thyroid hormone signaling during development

Laurent Sachs, Daniel R Buchholz

► To cite this version:

Laurent Sachs, Daniel R Buchholz. Frogs model man: In vivo thyroid hormone signaling during development. *Genesis*, 2017, 55 (1-2), pp.e23000. 10.1002/dvg.23000 . hal-02926081

HAL Id: hal-02926081

<https://hal.science/hal-02926081>

Submitted on 31 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Frogs Model Man: *In-vivo* thyroid hormone signaling during development

Laurent M. Sachs¹ and Daniel R. Buchholz²

¹UMR 7221 CNRS, Muséum National d'histoire Naturelle, Dépt. Régulation,
Développement et Diversité Moléculaire, Sorbonne Universités, 75005, Paris, France.

Fax: 33140793618

²Dept. Biological Sciences, University of Cincinnati, 312 Clifton Ct., Cincinnati, Ohio
45221. Fax: (513) 556-5299

Short title: Frog model of developmental TH signaling

Key words: amphibian, *Xenopus*, metamorphosis, perinatal endocrinology, gene
regulation, molecular mechanism

Abstract:

Thyroid hormone (TH) signaling comprises TH transport across cell membranes, metabolism by deiodinases, and molecular mechanisms of gene regulation. Proper TH signaling is essential for normal perinatal development, most notably for neurogenesis and fetal growth. Knowledge of perinatal TH endocrinology needs improvement to provide better treatments for premature infants and endocrine diseases during gestation and to counteract effects of endocrine disrupting chemicals. Studies in amphibians have provided major insights to understand *in-vivo* mechanisms of TH signaling. The frog model boasts dramatic TH-dependent changes directly observable in free-living tadpoles with precise and easy experimental control of the TH response at developmental stages comparable to fetal stages in mammals. The hormones, their receptors, molecular mechanisms, and developmental roles of TH signaling are conserved to a high degree in humans and amphibians, such that with respect to developmental TH signaling "frogs are just little people that hop". The frog model is exceptionally illustrative of fundamental molecular mechanisms of *in-vivo* TH action involving TH receptors, transcriptional cofactors, and chromatin remodeling. This review highlights the current need, recent successes, and future prospects using amphibians as a model to elucidate molecular mechanisms and functional roles of TH signaling during post-embryonic development.

1) Thyroid hormone (TH) signaling during human perinatal development

Lack of thyroid hormone (TH) during perinatal development causes debilitating mental deficits and short stature known as cretinism (Delange, 2005), but even subtle reductions in TH signaling can cause significantly lower IQ (Biondi and Cooper, 2008). Mutations in components of TH signaling, such as TH receptors, plasma membrane TH transporters, deiodinases, and cytosolic TH binding proteins cause various forms of hypo- and hyperthyroidism (Abe et al., 2003; Hernandez et al., 2006; Refetoff, 2005; Visser et al., 2010). Endocrine disrupting chemicals that interfere with any of these TH signaling components can also compromise short-term and long-term health and fitness (Gore et al., 2015).

The critical dependence on appropriate TH signaling for proper development of the brain and other organs motivates efforts to look for treatment options during pregnancy with jeopardized TH signaling, such as in Hashimoto's thyroiditis, Grave's disease, premature birth, and endocrine disruption. Unfortunately, many mechanisms pertinent to perinatal endocrinology are not well understood, and studies are needed to reduce known and unknown treatment consequences for fetal growth and development (Forhead and Fowden, 2014). The fact that the rate of preterm delivery with incomplete TH-dependent organ maturation is increasing despite extensive efforts to stop it and the increasing prevalence of endocrine disrupting chemicals points to the importance of broadening the net of basic research to better diagnose, treat, and/or prevent medical issues that may affect perinatal development (Goldenberg et al., 2008; Wang et al., 2014).

2) The value of *Xenopus* to study TH signaling during development

Elucidating the critical, complex, and experimentally intractable period of perinatal development in humans requires studies in model organisms. Model organisms provide simplified systems that are accessible and easily manipulated and can reveal the basic operating principles and disease etiologies associated with TH signaling that are nearly the same across vertebrates. Mice are the most commonly used model to study human health and disease because of their small size, ease of breeding in lab, and evolutionary closeness to humans. However, studies to elucidate developmental mechanisms of TH signaling are constrained in mammalian systems by the difficulty of observing relatively subtle or cryptic TH-dependent changes and of obtaining samples from fetuses *in utero*. An additional difficulty is that fetal tissues are constantly exposed to maternal hormones through the placenta (Forhead and Fowden, 2014), such that manipulation of fetal endocrine signaling to examine receptor function in plus or minus hormonal states is difficult to achieve without potentially introducing artefacts from altered maternal endocrine physiology. Furthermore, as with all model systems, none can fully recapitulate the human situation. For example, mutations in the TH transporter MCT8 in humans cause the crippling disorder, Allan-Herndon-Dudley syndrome (Friesema et al., 2004), whereas equivalent mutations in mice are nearly asymptomatic (Heuer and Visser, 2013).

For elucidating the molecular mechanisms of TH signaling during development, amphibians have intrinsic experimental advantages that make them the model of choice for several reasons. First, the dramatic TH-dependent molecular and morphological changes that occur during metamorphosis are unrivaled among terrestrial vertebrates

(Dent, 1968). Second, tadpoles are large and accessible throughout their development, including TH-dependent stages, comparable to perinatal stages in humans (Buchholz, 2015). Third, signaling via TH and their receptors is necessary and sufficient to initiate virtually all developmental events during metamorphosis (Das et al., 2010; Dodd and Dodd, 1976). Fourth, plasma levels of TH undetectable by radioimmunoassay occur naturally during the frog larval period prior to metamorphosis (Leloup and Buscaglia, 1977), indicating that virtually all of the receptors *in vivo* are in the unliganded condition. Fifth, endogenous TH levels increase to a peak at metamorphic climax (Leloup and Buscaglia, 1977), such that simple exogenous addition of TH to the rearing water during premetamorphosis enables precise temporal control of TH receptors to the liganded state that can mimic natural metamorphosis (Dodd and Dodd, 1976). Sixth, mechanisms of TH signaling in gene regulation and development are highly conserved between frogs and humans (see below), such that fundamental processes can be worked out in frogs and then applied to specific situations in mice and humans. Seventh, amphibians produce large numbers of free-living eggs and embryos that are easy to culture without specialized media or temperature requirements making tadpole studies fast, easy, and cheap with respect to comparable stages in mice (i.e., perinatal stages). Thus, because frogs are easy to breed and maintain in the laboratory, are the closest relatives to humans with easily accessible embryos, and have all the modern tools of a genetic model system, such as a sequenced genome (Hellsten et al., 2010; Session et al., 2016), an ORFeome (Grant et al., 2015), and established methods for gene knockout (Tandon et al., 2016) and transgenesis (Buchholz, 2012; Ishibashi et al., 2012), amphibian development is a particularly compelling model system from

experimental and fiscal perspectives for use in elucidating TH signaling applicable to human perinatal development.

3) Conservation in TH signaling between frogs and man

Many developmental events are regulated by TH in common in humans and frogs, including central and peripheral nervous systems (Kollros, 1981; Patel et al., 2011; Préau et al., 2016; Thompson and Cline, 2016), musculoskeletal growth and development (Dodd and Dodd, 1976; Van Vliet, 2005), and interaction with glucocorticoids to promote terminal organ maturation (Buchholz, 2015; Fowden and Forhead, 2013). Underlying these conserved TH-dependent developmental events are conserved molecular components (Buchholz et al., 2006; Furlow and Neff, 2006). As in mammals, amphibians have two types of TH receptors (TR α and TR β) (Helbing et al., 1992; Schneider et al., 1993; Wang et al., 2008; Yaoita et al., 1990) with similar alternative splicing of multiple mRNA isoforms and high amino acid sequence similarity (Yaoita et al., 1990). In addition, mammals and frogs share similar TR heterodimer partners (retinoic acid receptors alpha, beta, gamma), TR-associated co-repressors and co-activators, and canonical TH response elements in enhancer or promoter regions of TH target genes (Furlow and Neff, 2006). Furthermore, all known vertebrate TRs commonly bind genomic DNA sequences consisting of two direct repeats of a consensus hexameric AGGTCA sequence separate by four nucleotides (Das et al., 2009). Prior to TR binding, TH enters cells in mammals and frogs via conserved TH transporters (Lat1, MCT8, MCT10, OATP1c1 (Choi et al., 2015a; Connors et al., 2010; Ritchie et al., 2003)), followed by deiodinase-mediated metabolism (deiodinases I, II, III,

(Brown, 2005; Kuiper et al., 2006)) and possibly binding cytoplasmic proteins (CRYM, PKM2, and others (Choi et al., 2015a; Shi et al., 1994; Yamauchi and Tata, 2001)). Thus, amphibian TH signaling behaves as in humans to regulate many developmental events in common via comparable mechanisms with similar cytoplasmic TH signaling components, TR secondary structure, ligand and DNA binding properties, and requirement to heterodimerize with RXR (Wong and Shi, 1995).

4) Recent advances elucidated first in frogs

Role of unliganded TH receptor in post-embryonic development

The action of TH receptors (TRs) on gene regulation depend on TH, such that TRs actively repress genes in the absence of TH and induce those same genes in the presence of TH (Cheng et al., 2010). Such ligand-dependent gene regulation was derived from *in-vitro* cell culture studies, but the dual function model for how TRs might act during development was first proposed in frogs to account for TR action during critical periods of tissue differentiation when TRs transition from unliganded to liganded states (Buchholz et al., 2006; Sachs et al., 2000; Shi, 2009; Yaoita and Brown, 1990). Numerous studies in frogs and mice established that TR-mediated gene induction in response to TH is required for tissue differentiation, but only recently were two groups, working in frogs, able to address the role of unliganded TR α during development (Choi et al., 2015b; Wen and Shi, 2015). Using TALENs, they showed that TR α knockout led to the earlier initiation of metamorphosis and acceleration of development and further showed higher expression of TH response genes important for developmental progression resulting from lack of repression by unliganded TR α . Equivalent knockout

models were previously made in mice (Flamant and Samarut, 2003), but the more subtle murine TH-dependent phenotypes were not previously observed revealing the importance of the frog model to detect activities of TR during development. The production of a knockout of TR β is highly expected to define the relative contribution of each TR isoform.

Identification of TH direct response genes in developing tissues

The effects of TH on development are mediated mainly through transcriptional regulation of TH response genes (Buchholz et al., 2004). Identification of such direct response genes is thus of critical importance in understanding TH action during development. However, only a limited number of direct T3 response genes are known in different model systems, thus hampering our understanding of how TH regulates development *in vivo*. The first global analysis to identify such genes in development was conducted in frogs using cycloheximide-treated tadpoles to block translation thereby blocking secondary transcriptional regulation by TH-induced transcription factors (Das et al., 2009). They identified 188 up-regulated and 249 down-regulated genes by TH in the absence of new protein synthesis in whole animals, and gene ontology analysis showed that the direct up-regulated genes are enriched in categories important for transcriptional regulation and protein degradation-dependent signaling processes but not DNA replication. These findings thus revealed the pathways induced by TH at the earliest step of TH-dependent development.

Another experimental approach is to use chromatin immunoprecipitation (ChIP) to identify direct response genes (Grøntved et al., 2015). With the ChIP-Seq technology,

high throughput sequencing of purified DNA fragments enriched in TR binding sites followed by location of sequencing reads on the genome sequence supply genome-wide TR binding profiles. However, ChIP-seq suffers from a major limitation, as it provides no evidence of the functional connection between the TR binding sites and the target genes (only nearby locations can be inferred). Furthermore, it is now clear that transcription factors can act over large distances through DNA looping with their target promoter. To take into account these concerns, chromatin interaction analysis by paired-end tags sequencing (ChIA-PET) was carried out to map genome-wide TR binding sites and to link them with their target promoter by resolving long-range interaction between enhancer and promoter (Buisine et al., 2015). This study is the first to benefit from such technology using chromatin material isolated directly from animal tissues (Buisine et al., 2015). ChIA-PET simultaneously provides the equivalent of both ChIP-Seq plus chromosome conformation capture analyses combined. However, ChIA-PET technology requires high-resolution genome (assembly and annotation) that is achieved in *Xenopus tropicalis* (Buisine et al., 2015; Hellsten et al., 2010). The preliminary data show that indeed, TR can act over large genomic distances and that it optimizes identification of TR direct target genes (Buisine et al., 2015). Analyses of the TR ChIA-PET data are ongoing. This will represent a big step in the understanding of TH and TR action in a way not yet addressed in mammals.

Recruitment of cofactors to TH-response genes in vivo

Numerous cofactors (corepressors and coactivators) have been found to interact with TR using various biochemical approaches *in vitro* (Glass and Rosenfeld, 2000). Genetic

studies and the first use of ChIP from living tissue was performed in frogs to elucidate cofactor recruitment of TH/TR action *in vivo* during development (Sachs and Shi, 2000). Nuclear receptor corepressor (NCoR), and histone acetylase 3 (HDAC3), are major corepressor components and were found to be recruited to TREs in the absence of TH during early development and are released following TH treatment. Experimental confirmation of the importance of NCoR in TH-mediated gene repression *in vivo* was first obtained in frogs using overexpression of dominant negative NCoR by *in vivo* gene transfer into tadpole tail muscle (Sachs et al., 2002) and later by transgenesis (Sato et al., 2007) which led to the loss of repression by unliganded TR and faster tadpole development than wild type siblings. As for corepressors, coactivators, namely steroid receptor coactivator (SRC/p160), histone acetyltransferase (p300), coactivator associated arginine methyltransferase 1 (CARM1), and protein arginine methyltransferase 1 (PRMT1), were first shown to be recruited in a TH-dependent manner to TH-response genes in frogs *in vivo* during development (Matsuda et al., 2007; Matsuda et al., 2009; Paul et al., 2005; Paul et al., 2005; Paul et al., 2007). The value of *in-vivo* studies is demonstrated in that SRC3 was recruited in a gene and tissue dependent manner to TRE (Havis et al., 2003; Paul et al., 2005; Paul et al., 2005), which is difficult to determine in other systems. Transgenic overexpression of dominant negative SRC3 and dominant negative p300, which compete for the recruitment of endogenous coactivators, prevented TH-dependent gene regulation and delayed or arrested metamorphosis (Paul et al., 2005; Paul et al., 2007).

TH-dependent chromatin modifications in vivo

Chromatin modifications (covalent changes to DNA or histones) include DNA methylation, post-translational histone modifications, and altered chromatin structure, including histone composition and nucleosome displacement or removal. All these aspects of chromatin modification were analyzed in the context of TH gene regulation in amphibians (Grimaldi et al., 2013). Antibodies for studying chromatin modifications are against antigens highly conserved among vertebrates, which when coupled with advantages of metamorphosis make the frog system a superior model for elucidating mechanisms of TH-dependent chromatin remodeling *in vivo*. Several studies have shown that liganded TR induces chromatin remodeling. First, analysis in the reconstituted frog oocyte system suggests that TR is able to recognize a TRE within chromatin, TR makes use of chromatin assembly process to silence transcription efficiently, and TR directs the disruption of TRE chromatin structure in response to TH (Li et al., 1999; Wong et al., 1997a; Wong et al., 1997b; Wong et al., 1995; Wong et al., 1998). The disruption corresponds to the loss of 2 to 3 nucleosomes, a process that has been observed during metamorphosis (Matsuura et al., 2012). BRG1 and BAF57, recruited by liganded TR (Heimeier et al., 2008), are good candidates to remove the nucleosomes at the target genes. In the oocyte-reconstituted system, p300 recruitment by liganded TR modifies histones to initiate the recruitment of BRG1 (Huang et al., 2003). Before metamorphosis, histones H3 and H4 are deacetylated around TRE loci in the absence of TH and are later acetylated when TH level rises (Sachs and Shi, 2000). Many studies *in vivo* in frogs have provided more detail than the analyses done in mammals and have highlighted the importance of histone acetylation, i.e., a correlation

with the levels of gene expression, TR binding and RNA polymerase II recruitment (Bilesimo et al., 2011; Matsuura et al., 2012), treatment with HDAC inhibitor (Sachs and Shi, 2000; Sachs et al., 2001a; Sachs et al., 2001b), specific lysine acetylation in agreement with cofactor recruitment HAT or HDAC activity (Havis et al., 2006), transgenic overexpression of dominant positive TR (Buchholz et al., 2004) or dominant negative TR (Buchholz et al., 2003) and finally transgenic overexpression of a dominant negative SRC3 (Paul et al., 2005).

Histone methylation is a more complex type of modification, which can correlate with transcriptional silencing or activation (Kouzarides, 2007). ChIP assays were carried out with antibodies against different histone modifications in several tissues (brain, intestine and/or tail fin) in premetamorphic tadpole and during TH induced or natural metamorphosis. First, the repressive marks, H3K9 dimethylation and trimethylation, are not involved in TR mediated T_3 response gene regulation (Bilesimo et al., 2011; Matsuura et al., 2012). Second, Me3H3K27, another repressive mark, showed a tissue specific deposition in premetamorphic tadpoles inversely correlated with the basal low level of TH response gene expression measured in absence of TH (Bilesimo et al., 2011). Following TH treatment, Me3H3K27 level at TRE decreased pointing to a role of Polycomb and Thrithorax for regulating TH response genes (Bilesimo et al., 2011). Next, the level of Me3H3K79 and Me2H3R17, two active marks, increased upon TH treatment (Matsuura et al., 2012) suggesting the recruitment of Dot1L to methylate H3K79 and correlating with CARM1 recruitment for H3R17 (Matsuda et al., 2007). Dot1L knockdown with TALENs revealed its important role together with Me3H3K79 leading to retarded growth and lethality prior to metamorphosis (Wen et al., 2015). Finally, H3K4

methylation marks are more ambiguous because they can correlate with either activation or repression and have gene and tissue specific variations in the context of TR and TH dependent regulation of transcription in premetamorphic tadpoles following TH treatment (Bilesimo et al., 2011). Histone methylation is under intense investigation to clarify their *in-vivo* roles in gene regulation, with frog studies pushing the field, showing that H3K4 methylation levels correlated with TR binding to TRE (Bilesimo et al., 2011), an observation in accordance with its association with mammal enhancer or promoter regions (He et al., 2010; Heintzman et al., 2009) and the possible co-occurrence of the active Me3H3K4 mark with the repressive Me3H3K27 mark (Bilesimo et al., 2011).

Tool for screening TH-active compounds

The extreme sensitivity and responsivity of tadpoles to TH signaling provides a great platform for assaying chemicals that affect TH signaling. At the organismal level, the thyroid system represents an important target of endocrine disruption (Boas et al., 2012; Brucker-Davis, 1998), thereby prompting development of the *Xenopus* metamorphosis assay for screening TH-disrupting chemicals (Opitz et al., 2005). A high-throughput, whole-tadpole method using transgenic animals that express green fluorescent protein upon exposure to TH-disrupting chemicals was also developed (Castillo et al., 2013; Fini et al., 2007). At the receptor level, the two TR isoforms, α and β , have distinct tissue-specific expression patterns (Cheng et al., 2010), such that use of TR antagonists and TR isoform-selective agonists would benefit treatments desiring to block TR signaling or target specific tissues and avoid affecting off-target tissue-specific

side-effects. Tadpoles were the *in-vivo* model of choice to assess chemicals to modify TR function, including NH₃ (TR antagonist (Lim et al., 2002)), CO₂3 (TR α -selective agonist (Ocasio and Scanlan, 2006)), and GC1 (TR β selective agonist (Furlow et al., 2004)). The compounds GC1 and CO₂3 were then used to show that in tadpole brain development, TR α and not TR β is responsible for neural progenitor proliferation (Denver et al., 2009). In a further use of frog metamorphosis to reveal new knowledge, microarrays were used to identify targets of endocrine disruptors (Kulkarni and Buchholz, 2013). For example, the known xenoestrogen bisphenol A (BPA) was shown to antagonize the regulation of most TH-response genes, thereby suggesting that BPA predominantly affected TH-signaling pathways during development (Heimeier et al., 2009). More importantly, this result provides molecular evidence for the likely deleterious effects of BPA on human development and the importance of studying endocrine disruption in a developmental context *in vivo*.

5) Future potential to elucidate *in-vivo* TH signaling mechanisms using frogs

Future potential benefits regarding the preceding topics were covered in the sections above. Beyond those topics, TH signaling is such a dominant feature of frog post-embryonic development that many aspects of development related to or dependent on TH signaling are also powerfully studied using the frog system. Such TH-dependent processes where fundamental discoveries are being made include neurogenesis (Denver et al., 2009; Préau et al., 2016; Thompson and Cline, 2016), origin of adult intestinal stem cells (Ishizuya-Oka and Shi, 2005; Ishizuya-Oka et al., 2009), early eye development (Havis et al., 2006), spinal cord regeneration (Bhumika and Darras, 2014;

Gibbs et al., 2011; Lee-Liu et al., 2014), and interaction between TH and stress hormones at global and local scales (Bagamasbad et al., 2015; Kulkarni and Buchholz, 2012).

Of special note, because it has the potential to impact studies not only TH signaling but also gene regulation more generally, is the use of the frog model in network analysis to integrate transcriptome analyses, transcription factor binding, and epigenetic analyses to dramatically change how fundamental questions in biology are addressed. Such ambitious studies depend on the exploitation of the sequenced *Xenopus tropicalis* genome (Hellsten et al., 2010) and *Xenopus laevis* genome (Session et al., 2016) and the rise of high-throughput sequencing technologies and their applications to RNA (RNA-Seq). The execution of biological processes induced by TH (or any other bioregulatory molecule) requires the interaction and regulation of thousands of molecules. Systematic approaches to study large numbers of genes have revealed complex molecular networks as well as novel insights in understanding basic mechanisms controlling normal biological processes and pathologies (Zhu et al., 2007). Collection of large-scale data sets has begun and is being assembled into a network format whose topological structure contains significant biological properties (Sachs unpubl.). The integration of all interactions/modifications along with their dynamics will reveal a more complete description of how complex biological processes occur and can be controlled. These data based on work in frogs will allow more precise modeling human disease and to test more accurately corrective actions.

6) Conclusions

Starting with the discovery that feeding thyroid extracts to tadpoles induces metamorphosis (Gudernatsch, 1912), a century of discoveries stemmed from metamorphosis research, highlighted by hypothalamic and pituitary control of TH secretion (Allen, 1938; Dodd and Dodd, 1976), TH induction of mRNA and protein synthesis pointing to a nuclear action for the receptor (Tata, 1965; Tata, 1966), biochemical changes induced by TH during development (Frieden and Just, 1970), and identification of TH response gene regulation cascades (Brown et al., 1995; Brown et al., 1996; Brown et al., 2005; Buckbinder and Brown, 1992; Shi and Brown, 1990; Wang and Brown, 1993). Since these discoveries, the genome sequence, genetic methods (*in-vivo* gene transfer, transgenesis, gene knockout) have been added to the intrinsic benefits of frog biology (Buchholz, 2012; Grant et al., 2015; Hellsten et al., 2010; Ishibashi et al., 2012; Session et al., 2016; Tandon et al., 2016). Additional discoveries in fundamental biology are to be revealed about upstream TH signaling components, molecular mechanisms to gene regulation, and developmental roles of downstream TH-response genes in the gene regulation cascade. Potential use of fundamental discoveries in the treatment of disease can also be tested using the *in-vivo* frog model. Furthermore, endocrine disrupting chemicals are not going away and more are being produced and released into the environment contaminating drinking water. Elucidating the complexity of TH signaling at molecular and organismal levels will continue to require exemplary animal models such as frog development, with its inherent biological advantages and continuing development of experimental and bioinformatics tools. The frog system has been and will be a full-service model at the forefront in the study of TH-

dependent development. Another century of discovery awaits as we utilize the amazingly dramatic frog metamorphosis model to study *in-vivo* mechanisms of TH signaling during development.

7) Acknowledgements

Thanks to L. Shewade for help with the illustrations and to Muséum National d'Histoire Naturelle and the Centre National de la Recherche Scientifique for LMS.

References

- Abe, S., Katagiri, T., Saito-Hisaminato, A., Usami, S.-., Inoue, Y., Tsunoda, T., Nakamura, Y., 2003. Identification of CRYM as a candidate responsible for non-syndromic deafness through cDNA microarray analysis of human cochlear and vestibular tissues. *Am. J. Human Genet.* 72, 73-82.
- Allen, B.M., 1938. The Endocrine Control of Amphibian Metamorphosis. *Biol. Rev.* 13, 1-19.
- Bagamasbad, P.D., Bonett, R.M., Sachs, L., Buisine, N., Raj, S., Knoedler, J.R., Kyono, Y., Ruan, Y., Ruan, X., Denver, R.J., 2015. Dicephering the regulatory logic of an ancient, ultraconserved nuclear receptor enhancer module. *Mol. Endocrinol.* 29, 856-872.
- Bhumika, S., Darras, V.M., 2014. Role of thyroid hormones in different aspects of nervous system regeneration in vertebrates. *Gen. Comp. Endocrinol.* 203, 86-94.

- Bilesimo, P., Jolivet, P., Alfama, G., Buisine, N., Le Mevel, S., Havis, E., Demeneix, B.A., Sachs, L.M., 2011. Specific histone lysine 4 methylation patterns define TR-binding capacity and differentiate direct T3 responses. *Mol. Endocrinol.* 25, 225-237.
- Biondi, B., Cooper, D.S., 2008. The Clinical Significance of Subclinical Thyroid Dysfunction. *Endocrine Reviews* 29, 76-131.
- Boas, M., Feldt-Rasmussen, U., Main, K.M., 2012. Thyroid effects of endocrine disrupting chemicals. *Mol. Cell. Endocrinol.* 355, 240-248.
- Brown, D.D., 2005. The role of deiodinases in amphibian metamorphosis. *Thyroid* 15, 815-821.
- Brown, D.D., Cai, L., Das, B., Marsh-Armstrong, N., Schreiber, A.M., Juste, R., 2005. Thyroid hormone controls multiple independent programs required for limb development in *Xenopus laevis* metamorphosis. *Proc. Natl. Acad. Sci. U. S. A.* 102, 12455-12458.
- Brown, D.D., Wang, Z., Furlow, J.D., Kanamori, A., Schwartzman, R.A., Remo, B.F., Pinder, A., 1996. The thyroid hormone-induced tail resorption program during *Xenopus laevis* metamorphosis. *Proc. Natl. Acad. Sci. U. S. A.* 93, 1924-1929.
- Brown, D.D., Wang, Z., Kanamori, A., Eliceiri, B., Furlow, J.D., Schwartzman, R., 1995. Amphibian metamorphosis: a complex program of gene expression changes controlled by the thyroid hormone. *Recent Prog. Horm. Res.* 50, 309-315.
- Brucker-Davis, F., 1998. Effects of environmental synthetic chemicals on thyroid function. *Thyroid* 8, 827-856.

- Buchholz, D.R., 2012. Tet-On binary systems for tissue-specific and inducible transgene expression. *Methods in Mol. Biol.* 917, 265-275.
- Buchholz, D.R., 2015. More similar than you think: Frog metamorphosis as a model of human perinatal endocrinology. *Dev. Biol.* 408, 188-195.
- Buchholz, D.R., Hsia, S.C., Fu, L., Shi, Y.B., 2003. A dominant-negative thyroid hormone receptor blocks amphibian metamorphosis by retaining corepressors at target genes. *Mol. Cell. Biol.* 23, 6750-6758.
- Buchholz, D.R., Paul, B.D., Fu, L., Shi, Y.B., 2006. Molecular and developmental analyses of thyroid hormone receptor function in *Xenopus laevis*, the African clawed frog. *Gen. Comp. Endocrinol.* 145, 1-19.
- Buchholz, D.R., Tomita, A., Fu, L., Paul, B.D., Shi, Y.B., 2004. Transgenic analysis reveals that thyroid hormone receptor is sufficient to mediate the thyroid hormone signal in frog metamorphosis. *Mol. Cell. Biol.* 24, 9026-9037.
- Buckbinder, L., Brown, D.D., 1992. Thyroid hormone-induced gene expression changes in the developing frog limb. *J. Biol. Chem.* 267, 25786-25791.
- Buisine, N., Ruan, X., Bilesimo, P., Grimaldi, A., Alfama, G., Ariyaratne, P., Mulawadi, F., Chen, J., Sung, W.K., Liu, E.T., Demeneix, B.A., Ruan, Y., Sachs, L.M., 2015. *Xenopus tropicalis* genome re-scaffolding and re-annotation reach the resolution required for in vivo ChIA-PET analysis. *PloS One* 10, e0137526.
- Castillo, L., Seriki, K., Mateos, S., Loire, N., Guédon, N., Lemkine, G.F., Demeneix, B.A., Tindall, A.J., 2013. In vivo endocrine disruption assessment of wastewater treatment plant effluents with small organisms. *Water Sci. Technol.* 68, 261-268.

- Cheng, S.-., Leonard, J.L., Davis, P.J., 2010. Molecular Aspects of Thyroid Hormone Actions. *Endocr. Rev.* 31, 139-170.
- Choi, J., Moskalik, C.L., Ng, A., Matter, S.F., Buchholz, D.R., 2015a. Regulation of thyroid hormone-induced development in vivo by thyroid hormone transporters and cytosolic binding proteins. *Gen. Comp. Endocrinol.* 222, 69-80.
- Choi, J., Suzuki, K.T., Sakuma, T., Shewade, L., Yamamoto, T., Buchholz, D.R., 2015b. Unliganded thyroid hormone receptor alpha regulates developmental timing via gene repression as revealed by gene disruption in *Xenopus tropicalis*. *Endocrinol.* 156, 735-744.
- Connors, K.A., Korte, J.J., Anderson, G.W., Degitz, S.J., 2010. Characterization of thyroid hormone transporter expression during tissue-specific metamorphic events in *Xenopus tropicalis*. *Gen. Comp. Endocrinol.* 168, 149-159.
- Das, B., Heimeier, R.A., Buchholz, D.R., Shi, Y.B., 2009. Identification of direct thyroid hormone response genes reveals the earliest gene regulation programs during frog metamorphosis. *J. Biol. Chem.* 284, 34167-34178.
- Das, B., Matsuda, H., Fujimoto, K., Sun, G., Matsuura, K., Shi, Y.B., 2010. Molecular and genetic studies suggest that thyroid hormone receptor is both necessary and sufficient to mediate the developmental effects of thyroid hormone. *Gen. Comp. Endocrinol.* 168, 174-180.
- Delange, F.M., 2005. Endemic Cretinism. In: Braverman, L.E., Utiger, R.D. (Eds.), *Werner & Ingbar's The Thyroid: A fundamental and clinical text*, 9th ed. Lippincott Williams and Wilkins, Philadelphia, pp. 731-744.

- Dent, J.N., 1968. A survey of amphibian metamorphosis In: W. Etkin, L.I.G. (Ed.), Metamorphosis. A Problem in Developmental Biology. Appleton-Crofts, New York, pp. 271-311.
- Denver, R.J., Hu, F., Scanlan, T.S., Furlow, J.D., 2009. Thyroid hormone receptor subtype specificity for hormone-dependent neurogenesis in *Xenopus laevis*. Dev. Biol. 326, 155-168.
- Dodd, M.H.I., Dodd, J.M. 1976. The biology of metamorphosis. In: Lofts, B. (Ed.), Physiology of the Amphibia, New York, pp. 467-599.
- Finì, J.-., Le Mevel, S., Turque, N., Palmier, K., Zalko, D., Cravedi, J.-., Demeneix, B.A., 2007. An In Vivo Multiwell-Based Fluorescent Screen for Monitoring Vertebrate Thyroid Hormone Disruption. Environ. Sci. Technol. 41, 5908-5914.
- Flamant, F., Samarut, J., 2003. Thyroid hormone receptors: lessons from knockout and knock-in mutant mice. Trends Endocrinol. Metab. 14, 85-90.
- Forhead, A.J., Fowden, A.L., 2014. Thyroid hormones in fetal growth and prepartum maturation. J. Endocrinol. 221, R87-R103.
- Fowden, A.L., Forhead, A.J. 2013. Endocrine interactions in the control of fetal growth. In: Bhatia, J., Bhutta, Z.A., Kalhan, S.C. (Eds.), Maternal and Child Nutrition: The First 1,000 Days. Karger, Basel, pp. 91-102.
- Frieden, E., Just, J.J. 1970. Hormonal Responses in Amphibian Metamorphosis. In: Litwack, G. (Ed.), Biochemical Actions of Hormones. Academic Press, New York, pp. 1-52.
- Friesema, E.C., Grueters, A., Biebermann, H., Krude, H., Von Moers, A., Reeser, M., Barrett, T.G., Mancilla, E.E., Svensson, J., Kester, M.H., Kuiper, G.G., Balkassmi,

- S., Uitterlinden, A.G., Koehle, J., Rodien, P., Halestrap, A.P., Visser, T.J., 2004. Association between mutations in a thyroid hormone transporter and severe X-linked psychomotor retardation. *Lancet* 364, 1435-1437.
- Furlow, J.D., Neff, E.S., 2006. A developmental switch induced by thyroid hormone: *Xenopus laevis* metamorphosis. *Trends Endocrinol. Metab.* 17, 40-47.
- Furlow, J.D., Yang, H.Y., Hsu, M., Lim, W., Ermio, D.J., Chiellini, G., Scanlan, T.S., 2004. Induction of larval tissue resorption in *Xenopus laevis* tadpoles by the thyroid hormone receptor agonist GC-1. *J. Biol. Chem.* 279, 26555-26562.
- Gibbs, K.M., Chittur, S.V., Szaro, B.G., 2011. Metamorphosis and the regenerative capacity of spinal cord axons in *Xenopus laevis*. *European J. Neuroscience* 33, 9-25.
- Glass, C.K., Rosenfeld, M.G., 2000. The coregulator exchange in transcriptional functions of the nuclear receptors. *Genes Dev.* 14: 121-141. *Genes Dev.* 14, 121-141.
- Goldenberg, R.L., Culhane, J.F., Iams, J.D., Romero, R., 2008. Epidemiology and causes of preterm birth. *The Lancet* 371, 75-84.
- Gore, A.C., Chappell, V.A., Fenton, S.E., Flaws, J.A., Nadal, A., Prins, G.S., Toppari, J., Zoeller, R.T., 2015. EDC-2: The Endocrine Society's Second Scientific Statement on Endocrine-Disrupting Chemicals. *Endocr. Rev.* 36, 1-150.
- Grant, I.M., Balcha, D., Hao, T., Shen, Y., Trivedi, P., Patrushev, I., Fortriede, J.D., Karpinka, J.B., Liu, L., Zorn, A.M., Stukenberg, P.T., Hill, D.E., Gilchrist, M.J., 2015. The *Xenopus* ORFeome: A resource that enables functional genomics. *Dev. Biol.* 408, 345-357.

- Grimaldi, A., Buisine, N., Miller, T., Shi, Y.-., Sachs, L.M., 2013. Mechanisms of thyroid hormone receptor action during development: Lessons from amphibian studies. *Biochimica Et Biophysica Acta* 1830, 3882-3892.
- Grøntved, L., Waterfall, J.J., Kim, D.W., Baek, S., Sung, M.H., Zhao, L., Park, J.W., Nielsen, R., Walker, R.L., Zhu, Y.J., Meltzer, P.S., Hager, G.L., Cheng, S.-., 2015. Transcriptional activation by the thyroid hormone receptor through ligand-dependent receptor recruitment and chromatin remodeling. *Nat. Commun.* 6: 7048. *Nat. Commun.* 6, 7048.
- Gudernatsch, J.F., 1912. Feeding Experiments on Tadpoles. I. Tile influence of specific organs given as food on growth and differentiation. *Arch. f. Entwicklungsmechn, d. Organ* 35, 457-457-484.
- Havis, E., Le Mevel, S., Morvan Dubois, G., Shi, D.L., Scanlan, T.S., Demeneix, B.A., Sachs, L.M., 2006. Unliganded thyroid hormone receptor is essential for *Xenopus laevis* eye development. *Embo j.* 25, 4943-4951.
- Havis, E., Sachs, L.M., Demeneix, B.A., 2003. Metamorphic T3-response genes have specific co-regulator requirements. *EMBO Rep.* 4, 883-888.
- He, H.H., Meyer, C.A., Shin, H., Bailey, S.T., Wei, G., Wang, Q., Zhang, Y., Xu, K., Ni, M., Lupien, M., Mieczkowski, P., Lieb, J.D., Zhao, K., Borwn, M., Liu, X.S., 2010. Nucleosome dynamics define transcriptional enhancers. *Nat. Genet.* 42, 343-348.
- Heimeier, R.A., Hsia, V.S., Shi, Y.-., 2008. Participation of Brahma-related gene 1 (BRG1)-associated factor 57 and BRG1-containing chromatin remodeling complexes in thyroid hormone-dependent gene activation during vertebrate development. *Mol. Endocrinol.* 22, 1065-1077.

- Heimeier, R.A., Das, B., Buchholz, D.R., Shi, Y.B., 2009. The xenoestrogen bisphenol A inhibits postembryonic vertebrate development by antagonizing gene regulation by thyroid hormone. *Endocrinology* 150, 2964-2964-2973.
- Heintzman, N.D., Hon, G.C., Hawkins, R.D., Kheradpour, P., Stark, A., Harp, L.F., Ye, Z., Lee, L.K., Stuart, R.K., Ching, C.W., Ching, K.A., Antosiewicz-Bourget, J.E., Liu, H., Green, R.D., Lobanenkov, V.V., Stewart, R., Thomson, J.A., Crawford, G.E., Kellis, M., Ren, B., 2009. Histone modifications at human enhancers reflect global cell-type-specific gene expression. *Nature* 459, 108-112.
- Helbing, C.C., Gergely, G., Atkinson, B.G., 1992. Sequential up-regulation of thyroid hormone β receptor, ornithine transcarbamylase, and carbamyl phosphate synthetase mRNAs in the liver of *Rana catesbeiana* tadpoles during spontaneous and thyroid hormone-induced metamorphosis. *Dev. Genet.* 13, 289-289-301.
- Hellsten, U., Harland, R.M., Gilchrist, M.J., Hendrix, D., Jurka, J., Kapitonov, V., Ovcharenko, I., Putnam, N.H., Shu, S., Taher, L., Blitz, I.L., Blumberg, B., Dichmann, D.S., Dubchak, I., Amaya, E., Detter, J.C., Fletcher, R., Gerhard, D.S., Goodstein, D., Graves, T., Grigoriev, I.V., Grimwood, J., Kawashima, T., Lindquist, E., Lucas, S.M., Mead, P.E., Mitros, T., Ogino, H., Ohta, Y., Poliakov, A.V., Pollet, N., Robert, J., Salamov, A., Sater, A.K., Schmutz, J., Terry, A., Vize, P.D., Warren, W.C., Wells, D., Wills, A., Wilson, R.K., Zimmerman, L.B., Zorn, A.M., Grainger, R., Grammer, T., Khokha, M.K., Richardson, P.M., Rokhsar, D.S., 2010. The genome of the Western clawed frog *Xenopus tropicalis*. *Science* 328, 633-636.

- Hernandez, A., Martinez, M.E., Fiering, S., Galton, V.A., St. Germain, D., 2006. Type 3 deiodinase is critical for the maturation and function of the thyroid axis. *J. Clin. Investigation* 116, 476-484.
- Heuer, H., Visser, T.J., 2013. The pathophysiological consequences of thyroid hormone transporter deficiencies: Insights from mouse models. *Biochimica Et Biophysica Acta - General Subjects* 1830, 3974-3978.
- Huang, Z.-., Li, J., Sachs, L.M., Cole, P.A., Wong, J., 2003. A role for cofactor-cofactor and -histone interactions in targeting p300, SWI/SNF and Mediator for transcription. *Embo j.* 22, 2146-2155.
- Ishibashi, S., Love, N.R., Amaya, E., 2012. A simple method of transgenesis using I-SceI meganuclease in *Xenopus*. *Methods in Mol. Biol.* 917, 205-218.
- Ishizuya-Oka, A., Hasebe, T., Buchholz, D.R., Kajita, M., Fu, L., Shi, Y.B., 2009. Origin of the adult intestinal stem cells induced by thyroid hormone in *Xenopus laevis*. *Faseb j.*
- Ishizuya-Oka, A., Shi, Y.B., 2005. Molecular mechanisms for thyroid hormone-induced remodeling in the amphibian digestive tract: a model for studying organ regeneration. *Dev. Growth Differ.* 47, 601-607.
- Kollros, J.J., 1981. Chapter 13: Transitions in the nervous system during amphibian metamorphosis. In: Gilbert, L.I., Frieden, E. (Eds.), *Metamorphosis: A problem in developmental biology*, 2nd Edition. Plenum Press, New York, pp. 445-460.
- Kouzarides, T., 2007. Chromatin modifications and their function. *Cell* 128, 693-705.
- Kuiper, G.G., Klootwijk, W., Morvan Dubois, G., Destree, O., Darras, V.M., Van der Geyten, S., Demeneix, B., Visser, T.J., 2006. Characterization of recombinant

- Xenopus laevis* type I iodothyronine deiodinase: substitution of a proline residue in the catalytic center by serine (Pro132Ser) restores sensitivity to 6-propyl-2-thiouracil. *Endocrinology* 147, 3519-3529.
- Kulkarni, S.S., Buchholz, D.R., 2012. Beyond Synergy: Corticosterone and thyroid hormone have numerous interaction effects on gene regulation in *Xenopus tropicalis* tadpoles. *Endocrinol.* 153, 5309-5324.
- Kulkarni, S.S., Buchholz, D.R., 2013. Developmental Programs and Endocrine Disruption in Frog Metamorphosis: The Perspective from Microarray Analysis. *Current Topics in Developmental Biology* 103, 329-364.
- Lee-Liu, D., Moreno, M., Almonacid, L.I., Tapia, V.S., Muñoz, R., von Marées, J., Gaete, M., Melo, F., Larraín, J., 2014. Genome-wide expression profile of the response to spinal cord injury in *Xenopus laevis* reveals extensive differences between regenerative and non-regenerative stages. *Neural Develop.* 9, 12.
- Leloup, J., Buscaglia, M., 1977. La triiodothyronine, hormone de la metamorphose des Amphibies. *C. R. Acad. Sci. Paris, Ser. D* 284, 2261-2263.
- Li, Q., Imhof, A., Collingwood, T.N., Urnov, F.D., Wolffe, A.P., 1999. p300 stimulates transcription instigated by ligand-bound thyroid hormone receptor at a step subsequent to chromatin disruption. *Embo j.* 18, 5634-5652.
- Lim, W., Nguyen, N.H., Yang, H.Y., Scanlan, T.S., Furlow, J.D., 2002. A thyroid hormone antagonist that inhibits thyroid hormone action in vivo. *J. Biol. Chem.* 277, 35664-35670.
- Matsuda, H., Paul, B.D., Choi, C.Y., Hasebe, T., Shi, Y.B., 2009. Novel functions of protein arginine methyltransferase 1 in thyroid hormone receptor-mediated

transcription and in the regulation of metamorphic rate in *Xenopus laevis*. *Mol. Cell. Biol.* 29, 745-757.

Matsuda, H., Paul, B.D., Choi, C.Y., Shi, Y.B., 2007. Contrasting effects of two alternative splicing forms of coactivator-associated arginine methyltransferase 1 on thyroid hormone receptor-mediated transcription in *Xenopus laevis*. *Mol. Endocrinol.* 21, 1082-1094.

Matsuura, K., Fujimoto, K., Fu, L., Shi, Y.-., 2012.

Liganded thyroid hormone receptor induces nucleosome removal and histone modifications to activate transcription during larval intestinal cell death and adult stem cell development. *Endocrinol.* 153, 961-972.

Ocasio, C.A., Scanlan, T.S., 2006. Design and characterization of a thyroid hormone receptor alpha (TRalpha)-specific agonist. *ACS Chemical Biology* 1, 585-593.

Opitz, R., Braunbeck, T., Bögi, C., Pickford, D.B., Nentwig, G., Oehlmann, J., Tooi, O., Lutz, I., Kloas, W., 2005.

Description and initial evaluation of a *Xenopus* metamorphosis assay for detection of thyroid system-disrupting activities of environmental compounds. *Environ. Toxicol. Chem.* 24, 653-664.

Patel, J., Landers, K., Li, H., Mortimer, R.H., Richard, K., 2011. Thyroid hormones and fetal neurological development. *J. Endocrinol.* 209, 1-8.

Paul, B.D., Buchholz, D.R., Fu, L., Shi, Y.B., 2005. Tissue- and gene-specific recruitment of steroid receptor coactivator-3 by thyroid hormone receptor during development. *J. Biol. Chem.* 280, 27165-27172.

- Paul, B.D., Buchholz, D.R., Fu, L., Shi, Y.B., 2007. SRC-p300 coactivator complex is required for thyroid hormone-induced amphibian metamorphosis. *J. Biol. Chem.* 282, 7472-7481.
- Paul, B.D., Fu, L., Buchholz, D.R., Shi, Y.B., 2005. Coactivator recruitment is essential for liganded thyroid hormone receptor to initiate amphibian metamorphosis. *Mol. Cell. Biol.* 25, 5712-5724.
- Préau, L., Le Blay, K., Saint Paul, E., Morvan-Dubois, G., Demeneix, B.A., 2016. Differential thyroid hormone sensitivity of fast cycling progenitors in the neurogenic niches of tadpoles and juvenile frogs. *Mol. Cell. Endocrinol.* 420, 138-151.
- Refetoff, S., 2005. Resistance to thyroid hormone. In: Braverman, L.E., Utiger, R.D. (Eds.), *Werner & Ingbar's The Thyroid: A fundamental and clinical text*, 9th ed. Lippincott Williams and Wilkins, Philadelphia, pp. 1109-1129.
- Ritchie, J.W., Shi, Y.B., Hayashi, Y., Baird, F.E., Muchekeh, R.W., Christie, G.R., Taylor, P.M., 2003. A role for thyroid hormone transporters in transcriptional regulation by thyroid hormone receptors. *Mol. Endocrinol.* 17, 653-661.
- Sachs, L.M., Amano, A., Shi, Y.-., 2001a. An essential role of histone deacetylases in postembryonic organ transformations in *Xenopus laevis*. *J. Mol. Med.* 8, 595-601.
- Sachs, L.M., Amano, T., Rouse, N., Shi, Y.-., 2001b. Requirement of histone deacetylase at two distinct steps in thyroid hormone receptor mediated gene regulation during amphibian development. *Dev. Dynamics* 222, 280-291.
- Sachs, L.M., Shi, Y.B., 2000. Targeted chromatin binding and histone acetylation in vivo by thyroid hormone receptor during amphibian development. *PNAS* 97, 13138-13143.

- Sachs, L.M., Damjanovski, S., Jones, P.L., Li, Q., Amano, T., Ueda, S., Shi, Y.B., Ishizuya-Oka, A., 2000. Dual functions of thyroid hormone receptors during *Xenopus* development. *Comp. Biochem. Physiol. B. Biochem. Mol. Biol.* 126, 199-211.
- Sachs, L.M., Jones, P.L., Havis, E., Rouse, N., Demeneix, B.A., Shi, Y.B., 2002. Nuclear receptor corepressor recruitment by unliganded thyroid hormone receptor in gene repression during *Xenopus laevis* development. *Mol. Cell. Biol.* 22, 8527-8538.
- Sato, Y., Buchholz, D.R., Paul, B.D., Shi, Y.B., 2007. A role of unliganded thyroid hormone receptor in postembryonic development in *Xenopus laevis*. *Mech. Dev.* 124, 476-488.
- Schneider, M.J., Davey, J.C., Galton, V.A., 1993. *Rana catesbeiana* tadpole red blood cells express an alpha, but not a beta, c-erbA gene. *Endocrinology* 133, 2488-2495.
- Session, A.M., Uno, Y., Kwon, T., Chapman, J.A., Toyoda, A., Takahashi, S., Fukui, A., Hikosaka, A., Suzuki, A., Kondo, M., van Heeringen, S.J., Quigley, I., Heinz, S., Ogino, H., Ochi, H., Hellsten, U., Lyons, J.B., Simakov, O., Putnam, N., Stites, J., Kuroki, Y., Tanaka, T., Michiue, T., Watanabe, M., Bogdanovic, O., Lister, R., Georgiou, G., Paranjpe, S.S., van Kruijsbergen, I., Shu, S., Carlson, J., Kinoshita, T., Ohta, Y., Mawaribuchi, S., Jenkins, J., Grimwood, J., Schmutz, J., Mitros, T., Mozaffari, S.V., Suzuki, Y., Haramoto, Y., Yamamoto, T.S., Takagi, C., Heald, R., Miller, K., Haudenschield, C., Kitzman, J., Nakayama, T., Izutsu, Y., Robert, J., Fortriede, J., Burns, K., Lotay, V., Karimi, K., Yasuoka, Y., Dichmann, D.S., Flajnik, M.F., Houston, D.W., Shendure, J., DuPasquier, L., Vize, P.D., Zorn, A.M., Ito, M., Marcotte, E.M., Wallingford, J.B., Ito, Y., Asashima, M., Ueno, N., Matsuda, Y.,

- Veenstra, G.J., Fujiyama, A., Harland, R.M., Taira, M., Rokhsar, D.S., 2016. Genome evolution in the allotetraploid frog *Xenopus laevis*. *Nature* 538, 336-343.
- Shi, Y.B., 2009. Dual functions of thyroid hormone receptors in vertebrate development: the roles of histone-modifying cofactor complexes. *Thyroid* 19, 987-999.
- Shi, Y.B., Brown, D.D., 1990. Developmental and thyroid hormone-dependent regulation of pancreatic genes in *Xenopus laevis*. *Genes Dev.* 4, 1107-1113.
- Shi, Y.B., Liang, V.C., Parkison, C., Cheng, S.Y., 1994. Tissue-dependent developmental expression of a cytosolic thyroid hormone protein gene in *Xenopus*: its role in the regulation of amphibian metamorphosis. *FEBS Lett.* 355, 61-64.
- Tandon, P., Conlon, F., Furlow, J.D., Horb, M.E., 2016. Expanding the genetic toolkit in *Xenopus*: Approaches and opportunities for human disease modeling. *Dev. Biol.* (Epub ahead of print).
- Tata, J.R., 1965. Turnover of nuclear and cytoplasmic ribonucleic acid at the onset of induced amphibian metamorphosis. *Nature* 204, 378-381.
- Tata, J.R., 1966. Requirement for RNA and protein synthesis for induced regression of the tadpole tail. *Develop. Biol.* 13, 77-94.
- Thompson, C.K., Cline, H.T., 2016. Thyroid hormone acts locally to increase neurogenesis, neuronal differentiation, and dendritic arbor elaboration in tadpole visual system. *J. Neurosci.* 36, 10356-10375.
- Van Vliet, G., 2005. Hypothyroidism in infants and children: Congenital hypothyroidism. In: Braverman, L.E., Utiger, R.D. (Eds.), *Werner & Ingbar's The Thyroid A fundamental and clinical text*, 9th ed. Lippincott Williams & Wilkins, Philadelphia, PA, pp. 1033-1041.

- Visser, W.E., Friesema, E.C., Visser, T.J., 2010. Minireview: Thyroid Hormone Transporters: The Knowns and the Unknowns. *Mol. Endocrinol.*
- Wang, L.-., Chen, W.-., Chen, C.-., 2014. Preterm birth trend in Taiwan from 2001 to 2009. *J. Obstet. Gynaecol. Res.* 40, 1547-1554.
- Wang, X., Matsuda, H., Shi, Y.B., 2008. Developmental regulation and function of thyroid hormone receptors and 9-cis retinoic acid receptors during *Xenopus tropicalis* metamorphosis. *Endocrinology* 149, 5610-5618.
- Wang, Z., Brown, D.D., 1993. Thyroid hormone-induced gene expression program for amphibian tail resorption. *J. Biol. Chem.* 268, 16270-16278.
- Wen, L., Fu, L., Guo, X., Chen, Y., Shi, Y.-., 2015. Histone methyltransferase Dot1L plays a role in postembryonic development in *Xenopus tropicalis*. *Faseb j.* 29, 385-393.
- Wen, L., Shi, Y.-., 2015. Unliganded thyroid hormone receptor α controls developmental timing in *Xenopus tropicalis*. *Endocrinol.* 156, 721-734.
- Wong, J., Li, Q., Levi, B.-., Shi, Y.-., Wolffe, A.P., 1997a. Structural and functional features of a specific nucleosome containing a recognition element for the thyroid hormone receptor. *Embo j.* 16, 7130-7145.
- Wong, J., Shi, Y.-., Wolffe, A.P., 1997b. Determinants of chromatin disruption and transcriptional regulation initiated by the thyroid hormone receptor: hormone regulated chromatin disruption is not sufficient for transcriptional activation. *Embo j.* 16, 3158-3171.
- Wong, J., Liang, V.C., Sachs, L.M., Shi, Y.B., 1998. Transcription from the thyroid hormone-dependent promoter of the *Xenopus laevis* thyroid hormone receptor betaA

- gene requires a novel upstream element and the initiator, but not a TATA Box. *J. Biol. Chem.* 273, 14186-14193.
- Wong, J., Shi, Y.B., 1995. Coordinated regulation of and transcriptional activation by *Xenopus* thyroid hormone and retinoid X receptors. *J. Biol. Chem.* 270, 18479-18483.
- Wong, J., Shi, Y.B., Wolffe, A.P., 1995. A role for nucleosome assembly in both silencing and activation of the *Xenopus* TR beta A gene by the thyroid hormone receptor. *Genes Dev.* 9, 2696-2711.
- Yamauchi, K., Tata, J.R., 2001. Characterization of *Xenopus* cytosolic thyroid-hormone-binding protein (xCTBP) with aldehyde dehydrogenase activity. *Chem. Biol. Interact.* 130-132, 309-321.
- Yaoita, Y., Brown, D.D., 1990. A correlation of thyroid hormone receptor gene expression with amphibian metamorphosis. *Genes Dev.* 4, 1917-1924.
- Yaoita, Y., Shi, Y., Brown, D., 1990. *Xenopus laevis* alpha and beta thyroid hormone receptors. *Proc. Natl. Acad. Sci. U. S. A.* 87, 8684.
- Zhu, X., Gerstein, M., Snyder, M., 2007. Getting connected: analysis and principles of biological networks. *Genes Dev.* 21, 1010-1024.

Figure Legends

Figure 1: Conservation in endocrine mechanisms of TH signaling during development in mammals and amphibians. (a) During the aquatic to terrestrial transitions at birth and metamorphosis, humans and frogs each have a peak in plasma TH level, which most notably regulates neurogenesis and skeletal growth and development (G & D). (b) TH affects these processes in frogs to a higher degree than in mammals in order to accomplish the dramatic changes in diet (herbivorous to carnivorous) and locomotion (swimming to jumping). (c) A notable difference between groups is that TH is virtually absent in tadpole plasma until after limb bud development, whereas human and rodent fetuses are exposed to maternal TH from neural tube closure throughout fetal life (Forhead2014JOE), such that a distinct, experimentally advantageous switch from an unliganded to liganded condition of the TH receptors may occur naturally only during frog development.

Figure 2: Conservation in molecular mechanisms of TH signaling in mammals and amphibians. Processes from TH transport into the cell to altered gene expression share homologous proteins and mechanisms in humans and frogs. 1) TH transporters, such as LAT1 and MCT8, enable TH entry into cells where 2) deiodinase type I, II, and III function to remove iodine atoms from TH to activate or deactivate it. Before entry into the nucleus, cytoplasmic TH binding proteins (CTHBPs), e.g., mu-crystallin, modulate cytoplasmic occupancy. In the cell nucleus, 4) TH receptor (TR) heterodimerizes with retinoid-X-receptor (RXR) and binds to DNA at 5) TH response elements (TREs), where either 6) co-repressors, e.g., NCoR or SMRT, or co-activators, e.g., SRC, p300,

PRMT1, CARM1, are recruited depending on the 7) absence or presence of TH. The cofactors alter the 8) state of chromatin ultimately leading to 9) induced expression of TH response genes, e.g., *klf9*, *TR β* , in mammals and frogs.

Figure 3: Comparison of mouse and frog models for study of *in-vivo* TH signaling during development. Several unique aspects in frogs provide advantages for serving as a model to elucidate molecular mechanisms of TH signaling during development in vertebrates including humans. The key natural history assets for frogs are their exaggerated TH-dependent development, large clutch size, and free-living embryos/tadpoles giving rise to 1) ease of observation, tissue accessibility, and hormone manipulation, 2) lack of influence of maternal endocrine system on fetal development, and 3) natural development with unliganded TRs. The shorter generation time, historical advantage of mice for genome manipulation, and evolutionary closeness to humans are powerful aspects of the mouse model for TH signaling studies. However, with advances in gene disruption technology and sequencing of *X. tropicalis* and *X. laevis* genomes, the disparity in genomics tools between frogs and mice is closing.

Figure 1

Figure 2

Figure 3

		Frog advantage
		Model Feature		
Natural History	Morphological change	subtle	dramatic	✓
	Brood size/throughput	low (3-14)	high (>1000)	✓
	Generation time	2-3 mo.	4-6 mo.	—
	Similarity to humans	closer	farther	—
Free-living vs <i>in utero</i>	Accessibility	difficult	easy	✓
	Natural apo-TR	-/+	yes	✓
	Maternal influence	yes	no	✓
	Manipulate TH	problematic	easy	✓
Resources	Genetic resources	yes+	yes	—
	Genomic resources	yes	yes	=