



Peri-operative hyperglycemia in kidney transplantation

K. Monthé-Sagan, J Morera, F Fobe, V. Châtelet, M. Joubert, B. Hurault de Ligny, J L Hanouz, C. Gakuba

► To cite this version:

K. Monthé-Sagan, J Morera, F Fobe, V. Châtelet, M. Joubert, et al.. Peri-operative hyperglycemia in kidney transplantation. World Congress of Anesthesia, Aug 2016, Hong-Kong, China. hal-02927132

HAL Id: hal-02927132

<https://hal.science/hal-02927132>

Submitted on 1 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Peri-operative hyperglycemia in kidney transplantation PR391

K.Monthé-Sagan*, J.Morera, F.Fobe, V.Chatelet, M.Joubert, B.Hurault de Ligny, J.L.Hanouz, C.Gakuba
Pôle Réanimations Anesthésie SAMU/SMUR, CHU de Caen, Avenue de la Côte de Nacre, CS 30001, F-14 000 Caen, France
*monthesagan-k@chu-caen.fr

Introduction

Kidney transplantation is frequently related to peri-operative hyperglycemia due to surgical stress response and the introduction of immunosuppressive therapy. But these hyperglycemia could enhance the risk of delayed graft function. Our goal was, after establishment of a dynamic glycemic protocol, to assess its impact on the incidence of postoperative hyperglycemia.

Methods

This is an observational study. Over a period of 24 months, all patients who received a kidney transplant at the University Hospital of Caen were included. The glycemic values were recorded from the day before the surgery (D-1) to seven days (D7) after transplantation. After the first twelve months (1st period), a dynamic glucose protocol has been established (2nd period). Statistics: Fisher exact test, Student test or Mann-Whitney test (SPSS Version 21 software).

Results

One hundred and nine patients were enrolled. After creating a dynamic protocol for insulin therapy we saw a significant decrease in blood glucose values at day 1 (8.5 vs 7.4 mmol/L, $p = 0.001$) with more frequent use of subcutaneous insulin therapy (5 (9%) vs 20 (37%), $p = 0.001$). There is no significant difference in intravenous insulin therapy ($p = 0.51$). The peri-operative infection rate was reduced in the second period benefiting from the glycemic control protocol (14 (25%) vs 3 (6%) patients, $p = 0.004$) but no significant difference in rate of kidney graft dysfunction defined by the need for dialysis before J7 was observed ($p = 0.95$) (Table 1).

Conclusion

This work suggests that the introduction of a protocol for glycemic control in renal graft patient would allow a reduction of perioperative hyperglycemia. However glycemic control does not affect the rate of renal graft dysfunction.

Table 1: Perioperative adverse events

	1st Period n=55	2 nd Period n=54	p value
Hypoglycemia <3,3mmol.L ⁻¹	1(2)	1(2)	0.75
Hyperglycemia> 10mmol.L ⁻¹	25(46)	23(43)	0.76
Hypoglycemia & hyperglycemia	1(2)	0(0)	0.51
Diabetes	3(15)	20(37)	<0.001*
Infectious complications	14(25)	3(6)	0.004*
Surgical complications	13(24)	19(35)	0.19
Dialysis before the 7th day	27(49)	28(52)	0.95
Creatinine>250 μmol.L ⁻¹ beyond the 7th day	10(18)	13(24)	0.45

Values are expressed as number (%), median [minimum-maximum],
n number of patients, p* significant <0.05