Siphonodella leiosa (Conodonta), a new unornamented species from the Tournaisian (lower Carboniferous) of Puech de la Suque (Montagne Noire, France)

Louise Souquet, Carlo Corradini, Catherine Girard

► To cite this version:

Louise Souquet, Carlo Corradini, Catherine Girard. *Siphonodella leiosa* (Conodonta), a new unornamented species from the Tournaisian (lower Carboniferous) of Puech de la Suque (Montagne Noire, France). *Geobios*, 2020, 61, pp.55-60. 10.1016/j.geobios.2020.06.004 . hal-02934218

HAL Id: hal-02934218

<https://hal.science/hal-02934218>

Submitted on 6 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

***Siphonodella leiosi* (Conodonta), a new unornamented species from the Tournaisian (lower Carboniferous) of Puech de la Suque (Montagne Noire, France)**

Louise Souquet^{a, b}, Carlo Corradini^c and Catherine Girard^{d, *}

^a Centre de Recherche en Paléontologie - Paris, UMR CNRS 7207, CP 38, Muséum National d'Histoire Naturelle, 75005 Paris, France

^b Mécanismes Adaptatifs et Evolution, UMR CNRS 7179, Bâtiment Anatomie Comparée, CP 55, Muséum National d'Histoire Naturelle, 75005 Paris, France

^c Dipartimento di Matematica e Geoscienze, Università di Trieste, via Weiss 2, 34128 Trieste, Italy

^d ISEM, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France. (* corresponding author: Catherine.Girard@umontpellier.fr)

Abstract

A new conodont species, *Siphonodella leiosi*, is described from the lower Carboniferous pelagic limestones of the Montagne Noire (France), deposited on North Gondwana in outer platform environment. Specimens were obtained from one level dated to the *Siphonodella jii* conodont Zone. The major difference from other siphonodellid conodonts known in this area is that the elements of this new species have a practically entirely smooth and unornamented platform, apart from the development of one or two low rostral ridge-like nodes. Similar morphologies were generally observed in shallow marine deposits of the same time frame from China, Russia and East and Central European areas. The new discovery reinforces the idea that ornamentation of siphonodellids is not only related to bathymetry, but that temperature could play an important role in the diversification and radiation of unornamented species during the *Siphonodella jii* conodont Zone.

Keywords

Carboniferous, conodonts, *Siphonodella*, Montagne Noire

1. Introduction

The study of the Puech de la Suque pelagic section in the Montagne Noire (France) for high resolution stratigraphy through the Devonian/Carboniferous boundary resulted in an abundant and diverse conodont fauna collection (Feist et al., 2020). Conodonts are marine jawless vertebrates that lived from the late Cambrian to the Late Triassic. They are mainly known in the fossil record by their small teeth-like feeding structures called conodont elements. These elements are widely recognized as valuable tools for biostratigraphy, due to their abundance, high evolutionary rate, and good spatial and temporal record over long periods of time (Sweet, 1988).

Among conodonts, *Siphonodella* species are key stratigraphic markers to date the base of the Carboniferous. The first appearance datum of *Siphonodella sulcata* (Huddle) defines the base of the Carboniferous (Paproth et al., 1991). Other species of *Siphonodella* are used as index fossils for Tournaisian (lower Carboniferous) conodont zonations (Sandberg et al., 1978; Jii, 1985; Kaiser et al., 2009; Becker et al., 2016; Corradini et al., 2017; Hogancamp et al., 2019).

Siphonodella platform elements are usually ornamented, meaning that they bear nodes and ridges on the oral surface. The general shape of the platform and ornamentation are criteria used for taxonomy. *Siphonodella* is commonly associated with offshore open-marine environments with mesopelagic habitat (e.g. Austin, 1976; Ziegler & Sandberg, 1984; Savoy & Harris, 1993; Kalvoda et al., 1999; Kaiser et al., 2008). The ornamented *Siphonodella*: *Siphonodella bransoni* Ji, *Si. carinthiaca* Schönlaub, *Si. cooperi* Hass, *Si. crenulata* Cooper, *Si. duplicata* (Branson & Mehl), *Si. isosticha* Cooper, *Si. jii* Becker et al., *Si. lobata* (Branson & Mehl), *Si. obsoleta* Hass, *Si. praesulcata* Sandberg, *Si. quadruplicata* (Branson & Mehl), *Si. sandbergi* Klapper and *Si. sulcata* (Huddle) are considered pelagic species.

In Eastern and Central Europe, Russia, and China, species of *Siphonodella* that have a smooth oral surface without ornamentation have been described. These include: *Siphonodella belkai* Dzik, *Si. bella* Kononova & Migdisova, *Si. dasaibaensis* Ji, *Si. eurylobata* Ji, *Si. homosimplex* Ji & Ziegler, *Si. kalvodai* Kaiser, Kumpan & Cigler, *Si. levis* Ni, *Si. puchovki* Zhuravlev, *Si. quasinuda* Gagiev Kononova & Pazuhin, *Si. simplex* Ji, *Si. sinensis* Ji and *Si. uralica* Zhuravlev (e.g. Coen & Groessens, 1996; Dzik, 1997; Gagiev et al., 1987; Ji, 1987; Ji &

Ziegler, 1992; Kaiser et al., 2017; Kononova & Lipnjagov, 1976; Li et al., 2014; Lipnjagov, 1979; Malec, 2014; Matyja, 1976; Qie et al., 2014; Qin et al., 1988; Zhuravlev & Plotitsyn, 2017; Zhuravlev, 2019). They are found in Tournaisian neritic carbonate platform and slope facies, especially in the eastern and northern part of the Paleotethys realm. A “shallow-water” conodont zonation utilizing unornamented siphonodellids was proposed (Ji, 1987; 1988; Qie et al., 2014). In China (Ji, 1987; 1989), as well as in Russia (Zhuravlev, 2017), ornamented siphonodellids co-occur with unornamented species, allowing the correlation of “deep” and “shallow” water zonations.

Recently, Kaiser et al. (2017) found unornamented species (*Si. belkai* and *Si. kalvodai*) from the *jii* Zone in western part of Paleotethys. *Siphonodella belkai* was described from Kowala, Poland (Dzik, 1997 ; Malec, 2014). Kaiser et al. (2017) concluded that unornamented siphonodellids were able to invade the western part of the Paleotethys, but not north Gondwanan areas, due to the presence of the large Paleotethys ocean, nor eastern Laurussia, probably due to a terrestrial barrier.

One layer of strata in the *Siphonodella jii* Zone from the north Gondwana area in the Puech de la Suque section (Montagne Noire, France) yielded siphonodellid specimens that have an outline close to ornamented siphonodellids from this age, but differed in having a smooth and unornamented platform. This occurrence is coincident with the increased diversity within the genus *Siphonodella* observed at a global scale during the *Si. jii* Zone. This paper describes a new unornamented *Siphonodella* species, as well as the ecological and environmental significance of this taxon.

2. Geological setting

The Puech de la Suque section is located on the southeastern slope of the Puech de la Suque hill, near St Nazaire de Ladarez in the Montagne Noire (France) 43°29'59.7"N 3°05'54.8"E (Fig. 1(A)). This outcrop is tectonically inverted (Fig. 1(B)). The end-Famennian shales deposited during the regressive event of the Hangenberg Crisis are overlain by the Tournaisian succession recording transgressive trend from middle platform to basinal settings (Girard, 1994; Feist et al., 2020). The shales are superseded by the reappearance of carbonate sedimentation with 2m of gray-yellowish, nodular, dm-thick bedded mudstone to wackestone. These limestones contain ostracods (Lethiers & Feist, 1991) and conodonts

(Boyer et al., 1968; Girard, 1994, Kaiser, 2005; Feist et al., 2020), and are topped by well-bedded thin intercalations of cherts.

3. Material

New samples from the Puech de la Suque section yielded unornamented siphonodellid specimens from bed PS20, exclusively, located 1.2 meter above the base of the Carboniferous (Girard, 1994; Feist et al., 2020). The conodonts were extracted from this level in the laboratory of the University of Montpellier (France). Two kilograms of rock were dissolved using formic acid (10%) and then rinsed through 100µm and 1mm sieves. Conodonts were removed from the insoluble residue using a binocular microscope (Nikon SM8). Among the fifty *Siphonodella* specimens, nine were unornamented. The nine specimens were imaged using an X-ray microtomograph (µCT) Phoenix nanotomeS on the AniRA-Immos platform of the SFR Biosciences (UMS 3444, ENS Lyon) at a cubic voxel resolution of 1 µm. All specimens, holotypes and paratypes (UM PSQ 1 to UM PSQ 9), were housed in the collections of the University of Montpellier, and their 3D reconstructions (UM PSQ 1 to UM PSQ 9) were deposited in MorphoMuseum (Souquet et al., 2020).

4. Systematic Paleontology

Phylum Chordata Bateson, 1886

Class Conodonta Pander, 1856

Order Ozarkodinida Dzik, 1976

Family Elictognathidae Austin & Rhodes, 1981

Genus *Siphonodella* Branson & Mehl, 1944

Type species: *Siphonognathus duplicata* Branson & Mehl, 1934

Siphonodella leiosi sp. nov.

Figure 2

HOLOTYPE: dextral P1 element (UM PSQ 2), figured in Fig. 2(B).

PARATYPES: Figured P1 elements (UM PSQ 1, 3 to 9).

STRATUM TYPICUM: Bed of sample PS20.

TYPE LOCALITY: Puech de la Suque section, Montagne Noire, France.

DERIVATIO NOMINIS: From the Greek *leios*: slick, smooth

OCCURRENCE: *Siphonodella jii* Zone (lower Tournaisian).

MATERIAL: 9 specimens from Puech de la Suque section.

DIAGNOSIS: A species of *Siphonodella* whose P1 element is characterized by an asymmetric unornamented platform with a unilateral rostral ridge composed of nodes on the inner side. The lower side bears a wide pseudokeel.

DESCRIPTION: The P1 element of *Siphonodella leiosi* sp. nov. is characterized by an asymmetric smooth platform arched downward with apex of arch at position of the pit. Inner platform lanceolate and outer platform ovate. The short rostral ridge on the inner platform is composed of 3 to 5 discrete nodes. The anterior part of outer platform is strongly reduced. The short free blade bears partially fused and laterally flattened sharp denticles. Low incurved carina with nodes on posterior part of the platform. The aboral side of the element (lower view) is typical of the *Siphonodella* genus. It presents a wide pseudokeel which narrows anteriorly to the pit, the latter being placed slightly more posteriorly than the middle of the element.

REMARKS: The species differs from other taxa of *Siphonodella* by the lack of ornamentation, the shape of the very large and unornamented platform, and presence of the lateral rostrum ridge composed of nodes. Very reduced nodes may be developed on some specimens on the outer platform. This new species shows similar outline compared to ornamented species from Montagne Noire, but does not have ornamentation on the platform excepting one or two rostral ridge(s) that converge toward the carina. *Siphonodella leiosi* sp. nov. differs from unornamented species from the eastern part of the Paleotethys by the larger rostrum with well-differentiated nodes which converge towards the carina. Kalvoda et al. (2015, fig. 5(3)) illustrated an element from the *jii* Zone of Lesní lom, Moravia, that is similar to our element illustrated in Fig. 2(I), being slightly different by the more strongly developed rostral ridges.

This form may enter in the variability of *Si. leiosi*. The occurrence of the species in Moravia is confirmed by the recovery of another specimen from the same section and biozone (T. Kumpan, pers.com).

5. Discussion

Due to their global geographical distribution most of the species of the genus *Siphonodella* are used for stratigraphic subdivision of the uppermost Famennian and lower Tournaisian (e.g. Sandberg et al., 1978). The ornamented taxa are interpreted as inhabitants of offshore environments. Distinctive features include platform outline, development of a keel, nature of the rostrum and rostral ridges, as well as the nature of nodes and ridges on the inner and outer platform.

At the beginning of the Tournaisian, unornamented *Siphonodella* species are present on the eastern border of the Paleotethys. Two regions of endemism are recognized: South China and western Asia (Tajikistan, Kazakhstan, and Urals) (Figure 3(A)).

From the *jii* Zone, around 1.5 My after the Devonian/Carboniferous boundary (Davydov et al. 2012) an important radiation of ornamented siphonodellids (Sandberg et al., 1978) occurs worldwide (Kaiser et al., 2017 for a review). A diversification of endemic unornamented species was also observed in China and Russia (Ji, 1985 ; Zhuravlev, 2017). At the same time, unornamented species appeared on the western border of the Paleotethys realm for the first time with the recognition of *Si. belkai*, *Si. kalvodai*, and *Si. leiosi* sp. nov. (Figure 3(B)).

The *jii* Zone is a period of radiation of ornamented siphonodellids with the first occurrence of the cosmopolitan species *Si. jii*, *Si. cooperi*, *Si. obsoleta* and *Si. carinthiaca*. In China and Russia (s. l.), the unornamented species also radiated and diversified, with *Si. belkai* and *Si. dasaibaensis* in China (Ji & Ziegler, 1992., Li et al., 2014); and *Si. semichatovae* and *Si. ludmilae* in Russia (Zhuravlev, 2017).

Based on the stratigraphic distribution of taxa in different regions it is possible to suppose the migration route of some species along the Palaeotethys: *Si. belkai* (Kaiser et al., 2017), is known in China and is present from the base of the *jii* Zone in western and eastern

parts of the Paleotethys, and becomes abundant in eastern Europe only in the upper part of the *jii* Zone (Kaiser et al., 2017).

The Devonian / Carboniferous boundary coincides with a major regressive phase followed by a deepening during the Tournaisian (Girard, 1994; Haq & Schutter, 2008). The migration and dispersion of smooth species could be synchronous with transgressive periods. This scenario was suggested by Narkiewicz & Sculc (2004) and Kaiser et al. (2017) to explain dispersion of conodont faunas throughout the Tethys.

The *Siphonodella jii* Zone is also characterized by the first occurrence of two new species on the western part of the Paleotethys: *Si. leiosi* (this study), and *Si. kalvodai*. Kaiser et al. (2017, p. 17) noted that *Si. kalvodai* resembles *Si. belkai* (M1 and M2), « but differs in having a crenulated outer margin and by rostral ridges only on the anterior part of the platform ». It cannot be excluded that *Si. kalvodai* evolved from *Si. belkai* or from the same ancestor on western part of Paleotethys when local environmental conditions became favorable.

The history of *Siphonodella leiosi* is different as it does not resemble any other described species (except minor similarities in the outline with *Si. duplicata* ?), nor from east Paleotethys. As smooth species are supposed to prefer shallow-waters, the radiation and development of unornamented species all around the Paleotethys could be due to a global regression. However, this contradicts the scenario that suggests a transgressive phase allowing the migration of *Si. belkai* from the eastern part of the Paleotethys to the western part. Furthermore, biofacies and sedimentological analyses of sections in Montagne Noire (Girard, 1994; Feist et al., 2020) confirm a deepening through the basal Tournaisian after the major regression associated with Hangenberg Crisis just before the Devonian/Carboniferous boundary.

From the Devonian/Carboniferous boundary to the *jii* Zone, there is an increase in temperature of approximately 2-3°C, which remained stable thereafter (Buggisch et al., 2008). The diversification of unornamented siphonodellids was coincident with the warmest time. Shallow seas must be warmer than deep seas, and unornamented species have an equatorial repartition in the beginning of the Carboniferous. Associated, or not, with depth, the temperature could have played a role in development and selection of smooth siphonodellids.

Conclusions

A smooth siphonodellid was found in North Gondwana in the Montagne Noire (France) for the first time. This new species *Siphonodella leiosi* is characterized by an asymmetric smooth platform, a short rostral ridge on the inner platform composed of 3 to 5 discrete nodes, an anterior part of outer platform strongly reduced. The radiation and development of unornamented species all around the Paleotethys could be due to global changes in temperatures, associated, or not, to change in water depth.

References

- Austin, R.L., Rhodes, F.H.T., 1981. Systematic descriptions, in: Moore, R.C., Robison, R.A. (Eds.), Treatise on Invertebrate Paleontology. Geological Society of America, Boulder, p. 196.
- Austin, R.L., 1976. Evidence from Great Britain and Ireland concerning West European Dinantian conodont paleoecology, in: Barnes, C.R. (Ed.), Conodont paleoecology. Geological Association of Canada Special Paper, Montreal, pp. 201-224.
- Bateson, W., 1886. Memoirs: the ancestry of the Chordata. Journal of Cell Science 2, 535-572.
- Becker, R.T., Kaiser, S.I., Aretz, M., 2016. Review of chrono-, litho- and biostratigraphy across the global Hangenberg Crisis and Devonian–Carboniferous Boundary, in: Becker, R.T., Koenigshof, P., Brett, C.E. (Eds.), Devonian Climate, Sea Level and Evolutionary Events. The Geological Society of London, London, pp. 355-386.
- Boyer, F., Krylatov, S., Le Fèvre, J., Stoppel, D., 1968. Le Dévonien supérieur et la limite Dévonien/Carbonifère en Montagne Noire (France). Lithostratigraphie. Biostratigraphie (conodontes). Bulletin du Centre de Recherches de Pau-SPNA 2, 5-33.
- Branson, E.B., Mehl, M.G., 1934. Conodont studies number three. The University of Missouri studies. A quarterly of research VIII, 171-259.
- Branson, E.B., Mehl, M.G., 1944. Conodonts, in: Shimer, H.W. and RR Shrock, R.R. (Ed.), Index Fossils of North America.
- Buggisch, W., Joachimski, M.M., Sevastopulo, G., Morrow, J.R., 2008. Mississippian $\delta^{13}\text{C}_{\text{carb}}$ and conodont apatite $\delta^{18}\text{O}$ records — Their relation to the Late Palaeozoic Glaciation. Palaeogeography Palaeoclimatology Palaeoecology 268, 273-292.
- Coen, M., Groessens, E., 1996. Conodonts from the Devonian-Carboniferous transition beds of central Hunan, South China. Mémoires de l'Institut Géologique de l'Université de Louvain 36, 21-28.
- Corradini, C., Spalletta, C., Mossoni, A., Matyja, H., Over, J.D., 2017. Conodonts across the Devonian/Carboniferous boundary: A review and implication for the redefinition of the boundary and a proposal for an updated conodont zonation. Geological Magazine 154, 888-902.
- Davydov, V.I., Korn, D., Schmitz, M.D., 2012. The Carboniferous period, in: Gradstein, F.M., Ogg, J.G., Schmitz, M.D., Ogg, G.M. (Eds.), The Geologic Time scale 2012. Elsevier, pp. 603-651.
- Dzik, J., 1976. Remarks on the evolution of Ordovician conodonts. Acta Paleontologica Polonica 21, 395-455.
- Dzik, J., 1997. Emergence and succession of Carboniferous conodont and ammonoid communities in the Polish part of the Variscan sea. Acta Paleontologica Polonica 42, 57-170.
- Feist, R., Cornée, J.-J., Corradini, C., Hartenfels, S., Aretz, M., Girard, C., 2020. The Devonian–Carboniferous boundary in the stratotype area (SE Montagne Noire, France). Palaeobiodiversity and Palaeoenvironments. <https://doi.org/10.1007/s12549-019-00402-6>.

- Gagiev, M.H., Kononova, L.I., Pazukhin, V.N., 1987. Conodonts, in: Maslov, V.A. (Ed.), Fauna and Biostratigraphy of the Devonian-Carboniferous boundary deposits in Bertschogura (Mugodzhary), pp. 91-97.
- Girard, C., 1994. Conodont biofacies and event stratigraphy across the D/C boundary in the stratotype area (Montagne Noire, France), in: Königshof, P., Werner, R. (Eds.), Willi Ziegler-Festschrift I. Courier Forschung-Institut Senckenberg, Frankfurt a. M., pp. 299-309.
- Haq, B.U., Schutter, S.R., 2008. A chronology of Paleozoic sea-level changes. *Science* 322, 64-68.
- Hogancamp, N.J., Stolfus, B.M., Cramer, B.D., Day, J.E., 2019. A revised conodont zonation of the Tournaisian (Kinderhookian to lower Osagean) and implication for stratigraphic correlation in North America. In: Cramer, B.D., Clark, R.J., Day, J.E. (Eds.), The Devonian-Carboniferous boundary in the type area of the Mississippian. Iowa Geological Survey Guidebook 30, 11-17.
- Ji, Q., 1985. Study on the phylogeny, taxonomy, zonation and biofacies of *Siphonodella* (conodonta). *Bulletin of the Institute of Geology, Chinese Academy of Geological Sciences* 11, 51-75.
- Ji, Q., 1987. The Devonian–Carboniferous boundary in shallow-water facies areas of China as based on conodonts. *Acta Geologica Sinica* 61, 11-22.
- Ji, Q., et al. 1989. The Dapoushang section: an excellent section for the Devonian-Carboniferous boundary stratotype in China. *Science Press, Beijing, China*.
- Ji, Q., Ziegler, W., 1992. Introduction to some Late Devonian sequences in the Guilin area of Guangxi, South China. *Courier Forschung-Institut Senckenberg* 154, 149-177.
- Kaiser, S.I., 2005. Mass extinctions, climatic and oceanographic changes at the Devonian–Carboniferous boundary, Fakultät für Geowissenschaften. Ruhr-Universität Bochum, p. 156.
- Kaiser, S.I., Becker, R.T., Spalletta, C., Steuber, T., 2009. High-resolution conodont stratigraphy, biofacies and extinctions around the Hangenberg Event in pelagic successions from Austria, Italy and France. *Paleontographica Americana* 63, 97-139.
- Kaiser, S.I., Kumpan, T., Cíglér, V., 2017. New unornamented siphonodellids (Conodonta) of the lower Tournaisian from the Rhenish Massif and Moravian Karst (Germany and Czech Republic). *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen* 286, 1-33.
- Kaiser, S.I., Steuber, T., Becker, R.T., 2008. Environmental change during the Late Famennian and Early Tournaisian (Late Devonian–Early Carboniferous): implications from stable isotopes and conodont biofacies in southern Europe. *Geological Journal* 43, 241-260.
- Kalvoda, J., Bábek, O., Malovaná, A., 1999. Sedimentary and Biofacies Record in Calciturbidites at the Devonian-Carboniferous Boundary in Moravia (Moravian-Silesian Zone, Middle Europe). *Facies* 41, 141-158.
- Kalvoda, J., Kumpan, T., Babek, O., 2015. Upper Famennian and Lower Tournaisian sections of the Moravian Karst (Moravo-Silesian Zone, Czech Republic): a proposed key area for correlation of the conodont and foraminiferal zonations. *Geological Journal* 50, 17-38.
- Kononova, L.I., Lipnjagov, O.M., 1976. Occurrence of new conodonts from Lower Carboniferous of the Russian Platform and Don Basin. *Paleontologičeskij Žurnal* 4, 118-119 (in Russian).
- Lethiers, F., Feist, R., 1991. Ostracodes, stratigraphie et bathymétrie du passage Dévonien-Carbonifère au Viséen inférieur en Montagne Noire, France. *Geobios* 24, 71-104.
- Li, Z., Peng, Z., Cheng, L., Wang, C., Wang, B., 2014. New material of the early Carboniferous Conodonts from the lower member of the Baping Formation in Nandong, Guangxi, China. *Acta Micropalaeontologica Sinica* 31, 271-284.
- Lipnjagov, O.M., 1979. The conodonts of Ct1a and Ct1b of the Donetz Basin. *Service Geologique de Belgique, Professional Papers* 162, 41-49.
- Malec, J., 2014. The Devonian/Carboniferous boundary in the Holy Cross Mountains (Poland). *Geological Quarterly* 58, 217-234.
- Matyja, H., 1976. Biostratigraphy of the Devonian–Carboniferous passage beds from some selected profiles of NW Poland. *Acta Geologica Polonica* 26, 489-539.

- Narkiewicz, K., Szulc, J., 2004. Controls on migration of conodont fauna in peripheral oceanic areas. An example from the Middle Triassic of the Northern Peri-Tethys. *Geobios* 37, 425-436.
- Pander, C.H., 1856. Monographie der fossilen Fische des Silurischen Systems der russisch-baltischen Gouvernements. Königlich Preussischen Akademie der Wissenschaften, St Petersburg.
- Paproth, E., Feist, R., Flajs, G., 1991. Decision on the Devonian-Carboniferous boundary stratotype. *Episodes* 14, 331-336.
- Qie, W.K., Zhang, X., Du, Y., Yang, B., Ji, W., Luo, G., 2014. Conodont biostratigraphy of Tournaisian shallow-water carbonates in central Guangxi, South China. *Geobios* 47, 389-401.
- Qin, G., Zhao, R., Ji, Q., 1988. Late Devonian and Early Carboniferous conodonts from northern Guangdong and their stratigraphic significance. *Acta Micropalaeontologica Sinica* 5, 57-71.
- Sandberg, C.A., Ziegler, W., Leuteritz, K., Brill, S.M., 1978. Phylogeny, speciation and zonation of *Siphonodella* (Conodonta, Upper Devonian and Lower Carboniferous). *Newsletters on Stratigraphy* 7(2), 102-120.
- Savoy, L.E., Harris, A.G., 1993. Conodont biofacies and taphonomy along a carbonate ramp to black shale basin (latest Devonian and earliest Carboniferous), southernmost Canadian Cordillera and adjacent Montana. *Canadian Journal of Earth Sciences* 30, 2404-2422.
- Scotese, C.R., 2016. Tutorial: PALEOMAP PaleoAtlas for GPlates and the PaleoData Plotter Program, <http://www.earthbyte.org/paleomap-paleoatlas-for-gplates/>.
- Souquet L., Corradini C., Girard C., 2020. 3D models related to the publication: *Siphonodella leiosi* (Conodonta), a new unornamented species from the Tournaisian (lower Carboniferous) of Puech de la Suque (Montagne Noire, France). *MorphoMuseuM* e115. doi: 10.18563/journal.m3.115
- Sweet, W.C., 1988. The Conodonta: Morphology, taxonomy, paleoecology, and evolutionary history of a long-extinct animal phylum. *Oxford Monographs on Geology and Geophysics*, Oxford, New York, p. 212.
- Zhuravlev, A.V., 2017. A new species of the conodont genus *Siphonodella* Branson & Mehl (late Tournaisian). *Estonian Journal of Earth Sciences* 66, 188-192.
- Zhuravlev, A.V., Plotitsyn, A.N., 2017. The symmetry of the rostrum as a key to taxonomy of advanced *Siphonodella* (Conodonta, Early Carboniferous). *Stratigraphy* 14, 457-474.
- Zhuravlev, A.V., 2019. A new species of shallow-water *Siphonodella* (Conodonta) from the Tournaisian (Lower Carboniferous) of Pechora-Zozhva upliuft, Timan-Pechora basin. *Vestnik IG Komi SC UB RAS* 4. doi: 10.19110/2221-1381-2019-4-25-28
- Ziegler, W., Sandberg, C.A., 1984. *Palmatolepis*-based revision of upper part of standard Late Devonian conodont zonation, in: Clark, D.L. (Ed.), *Conodont biofacies and provincialism*. Geological Society of American special Paper, Boulder/Colorado, pp. 179-194.

Acknowledgment

We warmly thank Anne-Lise Charruault and Jean-Jacques Cornée for their fieldwork, and Sabrina Renaud (LBBE, Lyon) for discussions. We acknowledge the contribution of SFR Biosciences (UMS3444/CNRS, US8/ Inserm, ENS de Lyon, UCBL) AniRa-ImmOs facility. We particularly thank Mathilde Bouchet for her kind assistance during the scanning sessions. The research was supported by the Project Marcon (Labex CeMEB). This is contribution ISEM 2020-109.

357

358 **Figure captions**

359 Figure 1: A. Location of the studied area, the Montagne Noire (France). B. View of the
 360 inverted beds exposed in Puech de la Suque. C. Log of the lithological succession in the
 361 Puech de la Suque (PS) with occurrence of the species of *Siphonodella*. Conodont Zones after
 362 Corradini et al. (2017).

363

364 Figure 2: *Siphonodella leiosi* new species. From right to left: upper, lower and lateral views of
 365 P1 element (UM PSQ 1 to 9),. A: Paratype UM PSQ 1; B: Holotype UM PSQ 2; C: Paratype UM
 366 PSQ 3; D: UM PSQ 4; E: UM PSQ 5; F: UM PSQ 6; G: UM PSQ 7; H: UM PSQ 8; I: UM PSQ 9.
 367 Scale bar: 500 µm.

368

369 Figure 3: Tournaisian paleogeographic maps with the main currents and their directions
 370 (Scotese, 2016) showing the paleogeographic distribution of unornamented siphonodellids
 371 (A) from the *Protognathodus kockeli* Zone to the *Si. duplicata* Zone), and (B) during the
 372 *Siphonodella jii* Zone (Corradini et al., 2017). In bold, species not present before the *Si. jii*
 373 Zone. Illustrations of the siphonodellids from Ji & Ziegler (1992), Kaiser et al. (2017);
 374 Zhuravlev & Plotitsyn (2017), and this study. Rep.: Republic, *Pr.*: *Protognathodus*, *Si.*:
 375 *Siphonodella*.

Figure 1

Figure 3