**Ultrasonic bioacoustics and stridulum morphology reveal cryptic species among
Lipotactes big-eyed katydids (Orthoptera: Tettigoniidae: Lipotactinae) from Borneo**

Ming Kai Tan^{1*}, Sigfrid Ingrisch², Rodzay bin Haji Abdul Wahab³, Razy Japir⁴, Arthur Y. C. Chung⁴

¹ Institut de Systématique, Evolution et Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, SU, EPHE, UA, 57 rue Cuvier, CP 50, 75231 Paris Cedex 05, France

² Zoological Research Museum Alexander Koenig, Adenauerallee 160, D-53113 Bonn, Germany.

³ Institute for Biodiversity and Environmental Research, Universiti Brunei Darussalam, Jalan Universiti, BE1410, Brunei Darussalam

⁴ Forest Research Centre (Sepilok), Sabah Forestry Department, P.O. Box 1407, 90715 Sandakan, Sabah.

*Corresponding author: orthoptera.mingkai@gmail.com

Running title: Ultrasonic bioacoustics and taxonomy of Lipotactinae

Abstract

Lipotactinae is an elusive monophyletic subfamily of katydids (Orthoptera: Tettigoniidae) unique to Asia and comprising two genera—*Lipotactes* and *Mortoniellus*. Nearly nothing is known beyond their original descriptions. The stridulum morphology is rarely examined and described in taxonomy and their acoustics are only known for six species, none of which is from Borneo. New *Lipotactes* specimens collected from Borneo—Belait and Kuala Belalong in Brunei and Sandakan in Sabah—were initially identified as *Lipotactes alienus* and/or

Lipotactes virescens based on traditional morphological characters. However, the structures of male calling songs of individuals from Kuala Belalong and Sandakan were profoundly different from each other and from the known songs of *L. virescens* from Thailand and Peninsular Malaysia. This led us to examine the morphology of the Bornean specimens more closely. By integrating call structures, stridulum morphology and traditional morphology, we establish that the specimens from Borneo are different from *L. virescens* and that the populations from Kuala Belalong, Belait and Sandakan represent different ethospecies. Here, we describe the species from Sandakan as *Lipotactes kabili* **n. sp.** We also demonstrate that both stridulum morphology and call structures can be useful in separating Lipotactinae species because inter-specific differences are larger than intra-specific differences.

Key words: Brunei Darussalam, Sabah, Southeast Asia, ethospecies, new species, taxonomy

Introduction

Katydids communicate by stridulating with their forewings to produce sound (Montealegre-Z, 2009). More than 70% of the species with known songs sing at ultrasonic frequencies (>20 kHz) (Montealegre-Z et al., 2006, 2017). The ability to communicate at ultrasonic frequencies has drawn academic interest to study their biomechanics by investigating the stridulatory anatomy of the wings (e.g., Montealegre-Z, 2009, 2012; Montealegre-Z et al., 2017). However, ultrasonic-singing katydids are still poorly known in many aspects. These include whether their songs can be used to differentiate the species by bioacoustic signals because inter-individuals to inter-species variations are rarely examined (partly owing to insufficient specimens). Particularly, knowledge of bioacoustics of katydids from the hyper-diverse Southeast Asia remains remarkably poor—owing to the sheer number of species, difficulty of identifying them and relative lack of regional expertise—until recently (Tan et al., 2019).

The katydids of the subfamily Lipotactinae Ingrisch, 1995 are examples in which their bioacoustics are understudied. This small monophyletic subfamily is nowadays endemic to Asia (but see Gorochoy, 2010) and comprises two genera, *Lipotactes* Brunner von Wattenwyl, 1898 and *Mortoniellus* Griffini, 1909 with 24 species in *Lipotactes* and seven species in *Mortoniellus* (Ingrisch, 1995; Mugleston et al., 2018; Cigliano et al., 2019). These katydids are characterised by disproportionately large heads (and eyes), highly reduced and modified tegmina for singing, and legs armed with predatory spurs. Most species are recognised by their genitalia morphology and colour patterns on the pronotum and hind femur. Ingrisch (1995) erected the subfamily and provided the most thorough treatment on the biology and acoustics of this group to date. Since then, a few additions to this subfamily were published,

but Lipotactinae is still poorly known with very few references to them beyond the original descriptions (Gorochov, 1993, 1996, 1998; Chang et al., 2005; Shi & Li, 2009; Feng et al., 2017).

The bioacoustics of Lipotactinae is known from only five *Lipotactes* and one *Mortoniellus* species from Thailand, West Malaysia, Singapore and Sumatra, all of which were reported by Ingrisch (1995). More recently, Wang & Shi (2020) also report the songs of one *Lipotactes* from China. The calling songs of Lipotactinae typically consist of verses (often crescendoing) or continuous trains, and peak at ultrasonic frequencies (Ingrisch, 1995). However, stridulation was considered to have limited use in separating species (Ingrisch, 1995). The stridulum anatomy is also generally precluded in species descriptions and taxonomic studies (except for *Lipotactes montanus* Ingrisch, 1990 [Ingrisch, 1995]), even though the subfamily comprises only species with highly reduced tegmina modified for singing. This may be because the wings are concealed under the pronotum and are required to be dissected to examine the stridulatory organs

The tegmina of Lipotactinae are highly modified for singing, which in turn is imperative to communicate and attract conspecific females. Thus, the stridulum morphology, along with the acoustic parameters of the songs, can be useful for differentiating species, especially morphologically cryptic ones, and help resolve problems with delimitation of species. During recent orthopteran samplings in Borneo (i.e., Brunei Darussalam and Sabah), we collected males and females of *Lipotactes* (Fig. 1) that were tentatively identified to either *Lipotactes alienus* Brunner von Wattenwyl, 1898 or *L. virescens* Ingrisch, 1995. *L. alienus* is the type species of *Lipotactes* but was originally described from only one female from Borneo, River Baram (Brunner, 1898). Ingrisch (1995) described *L. virescens* from South Thailand/

Malaysia but also included specimens from Borneo. When the new specimens from Borneo were compared with the description and images in Ingrisch (1995) using traditional characters (i.e., genitalia and patterns), we were led to believe that these two species are morphologically very similar, and that *L. virescens* specimens from Borneo (as listed in Ingrisch, 1995) may belong to a different species (may or may not be *L. alienus*). The characters also exhibit population variations, suggesting that there may exist a species complex of *L. alienus-cum-virescens* from Borneo.

To elucidate these morphologically cryptic species, we examined previously unknown characters, specifically the acoustic parameters in the frequency (e.g., peak frequency) and time (e.g., pulse repetition rates) domains and stridulum morphology (i.e., mirror area and harp area, stridulatory file). We evaluated whether these characters can reveal information on the species boundaries of *Lipotactes*. Our prediction is that these new characters can help unravel the *Lipotactes* from Borneo into different putative ethospecies which are different from *L. virescens* from Thai-Malay Peninsula. Here, we also describe previously unknown acoustic signatures and stridulum morphology of the Bornean species.

Material and methods

Collection and depositories of katydids

Field collections and observations were made in the Belait and Temburong Districts of Brunei Darussalam between 23 February and 3 March 2019 and between 6 and 18 July 2019; as well as in Sandakan, Sabah between 8 and 12 January 2019 and between 30 September and 4 October 2019. Specimens were collected by sight during night and day. Whenever

possible, in-situ images were taken using a Canon EOS 500D digital SLR camera with a compact-macro lens EF 100 mm f/2.8 Macro USM and Canon Macro Twin Lite MT-24EX was used for lighting and flash. Specimens collected by Tan & Wahab (2018) from Kuala Belalong Field Studies Centre (Temburong, Brunei Darussalam) and Tan & Kamaruddin (2016) from Bukit Larut (Perak, Peninsular Malaysia) were also examined.

Specimen identification and curation

Identification was done using descriptions of all known *Lipotactes* species: Gorochov (1993, 1996, 1998); Ingrisch (1995); Chang et al. (2005); Shi & Li (2009). The specimens were preserved in absolute analytical-grade ethanol and later pinned and dry-preserved. For studying and documenting the distribution of teeth at the stridulatory file, the left tegmen was removed from fresh specimens using micro-scissors. For ethanol-preserved specimens, small amount of dilute KOH was added to relax the specimens before dissection. A single hind leg from each specimen was also preserved in absolute analytic-grade ethanol for future molecular work.

Close-up images of morphological features (including stridulum morphology) were done using a Canon macro photo lens MP-E 65 mm f/2.8 USM (1–5×). Image-editing was accomplished using Adobe Photoshop CC 2014 (Adobe Systems Incorporated, San Jose, CA, USA).

Acoustic recordings and analysis

Acoustic recording and analysis generally followed that in Tan et al. (2019). In the biological stations in Brunei Darussalam and Sandakan, the katydids were kept in insect cages with nylon netting. Acoustic recording was done using a sampling frequency of 256 kHz-

samples/s Echo Meter Touch (based on Knowles FG sensor) placed horizontally and at about 2–5 m away from the cage. The katydid usually (but not always) sings while clinging onto the side of the cage (thus positioning with dorsal surface facing the recording device).

Ambient temperature was logged using a HOBO 8K Pendant® Temperature logger (model: UA-001-08, Onset, Bourne, MA). The recorded signals were saved in 12-bit WAV format.

Acoustic parameters (e.g., peak frequency, mean frequency, repetition rate, song duration) were extracted using the ‘specan’ function in the open source R package WarbleR version 1.1.14 (Araya-Salas and Wright, 2017) in the R software version 3.5.1 (R Development Core Team, 2018). Sample size, from which the arithmetic means and standard deviations were calculated from, was denoted by the number of sound files. To filter out low or high background noise before performing measurements, the lower and upper limits of a frequency bandpass filter (in kHz) were set at 15 kHz and 120 kHz respectively so measurements of the frequency domain were obtained only between the frequency range. To generate power spectra using FFT, we used the function ‘spec’ at 256,000 sampling frequency, using Hanning window of window length 512. To compare the acoustic parameters, we fitted linear mixed effects models using the acoustic parameters as the response variables, population at which the katydid was collected as the fixed effect and temperature at the time of recording as random intercept to determine least-square means of each acoustic parameters to account for temperature effect. This was done using the ‘lmer’ function from the R package ‘lme4’ (Bates et al., 2014).

Measurement of morphological traits

We measured the maximum length and width and area of mirror and harp; inter-tooth distance and tooth length. Measurements were accomplished using ImageJ 1.51j8 (Wayne

166 Rasband, Research Services Branch, National Institute of Mental Health, Bethesda, MD,
167 USA).

168

169 *Terminologies*

170 For the description of songs, terminologies follow that by Ingrisch (1995) and Tan et al.
171 (2019): verse = complete sequence of syllables which is separated from other verse by a
172 distinct and prolonged pause; syllable = one complete cycle of tegminal movement; pulse = a
173 single unbroken wave train isolated in time by significant amplitude; peak frequency =
174 frequency with highest energy from the mean spectrum; mean frequency = weighted average
175 of frequency by amplitude; pulse duration = duration of the pulse; pulse repetition rate =
176 number of discrete pulses per second.

177

178 For the description of stridulum morphology, terminologies follow that by Béthoux (2012)
179 and Chivers et al. (2017): CuPb = posterior branch of posterior cubitus (CuP); h1 = anterior
180 portion of harp area; h2 = median portion of harp area; h3 = posterior part of harp area.

181

182 For the description of stridulatory file, terminologies follow that of Montealegre-Z and
183 Mason (2005) and Tan et al. (2019): inter-tooth distance = distance from the edge of the cusp
184 of one tooth to the cusp of the next one; tooth length = distance from the anterior to the
185 posterior ends of the cusp of each tooth.

186

187 For the measurements, the following abbreviations are used: BL = body length; HL = head
188 length; HW = head width (include eyes); PL = pronotum length; PW = pronotum width (at
189 metazona); TL = tegmen length; HFL = hind femur length; HTL = hind tibia length; OL =
190 ovipositor length.

191

192 ***Depositories***

193 All sound files were uploaded to the Orthoptera Species File Online Version 5.0/5.0

194 (Cigliano et al., 2019).

195

196 The dry-pinned specimens were deposited:

197 FRC Forest Research Centre, Sepilok, Sandakan, East Malaysia

198 UBDM Universiti Brunei Darussalam Museum, Brunei Darussalam

199 ZRC Zoological Reference Collection, Lee Kong Chian Natural History Museum,

200 Singapore

201

202

203 **Results**

204

205 ***Part I: taxonomy***

206 ***Lipotactes alienus* Brunner von Wattenwyl, 1898** (Figs. 1.1, 2.1, 2.5, 3.1, 5.1, 6.1)

207 Material examined. — Brunei Darussalam, Temburong District: 1 female (KB.16.37), Ulu

208 Temburong National Park, canopy walk, primary dipterocarp forest, N4.55196 E115.15990,

209 234.4±5.9 m, 1851h, on foliage of shrub, 25 September 2016, coll. M.K. Tan; 1 female

210 (KB.16.62), Temburong District, Kuala Belalong Field Studies Centre, Ashton Trail, primary

211 dipterocarp forest, N4.54712 E115.15726, 123.7±9.5 m, 2152h, on branch of tree after rain,

212 26 September 2016, coll. M.K. Tan; 1 male (KB.19.11) Ashton Trail, primary dipterocarp

213 forest, N4.54561 E115.15679, 109.3±11.9 m, 1948h, on foliage near ground, 13 July 2019,

214 coll. M.K. Tan & H. Yeo (UBDM and ZRC). Brunei Darussalam, Belait District: 1 male

215 (BR.19.37), Andulau Forest Reserve, N4.62420 E114.51322, 84.6±7.0 m, 1818h, under

216 foliage 28 February 2019, coll. M.K. Tan & H. Yeo (UBDM) (note that any reference to the
217 population/ individuals from Belait in the text does not include this specimen but instead refer
218 to *Lipotactes* ethospecies [Belait] [see below]).

219 Remarks. — The females are sufficiently described in Ingrisch (1995). Females collected
220 from Kuala Belalong resemble *L. alienus* by colour patterns on hind femur (including black
221 hind knee); dark markings on pronotum dorsal disk, metanotum and abdominal tergite; shape
222 of subgenital plate. We inferred that the males collected from the same trail Kuala Belalong
223 are therefore probably the undescribed males of *L. alienus*. However, we recommend that
224 DNA data may be needed to match the females from Kuala Belalong with the holotype (and
225 specimens from Long Pa Sia [see Ingrisch 1995]) to be completely certain.

226 Description of male. — Very similar to *L. virescens*: size, colouration; pronotum with
227 anterior margin and transverse sulcus black and M- or U-shaped black markings on dorsal
228 disk similar to *L. virescens* males (Fig. 2.1, see also Ingrisch 1995: Fig. 65); but differs by:
229 generally lighter colouration. Tenth abdominal tergite prolonged behind in middle,
230 prolongation more broadly rounded (Fig. 3.1). Cercus more slender at apex (instead of
231 uniformly conical in *L. virescens*), basal lobe of internal process distinctly larger, length
232 nearly reaching mid-point of internal process. Subgenital plate with two pronounced lateral
233 lobes at apex (where styli are inserted); apex between base of styli strongly emarginated.
234 Stylus dark, long and slender (Fig. 3.1).

235

236 ***Lipotactes virescens* Ingrisch, 1995** (Figs. 1.2, 2.2, 2.6, 3.2, 5.2, 6.2)

237 Material examined. — 1 male (LAR.15.135), Peninsular Malaysia, Perak, Taiping, near
238 Sungei Larut, young secondary forest and forest edge, N4.85418 E100.76057, 62.7±5.3 m, on
239 branch, 19 June 2015, coll. M.K. Tan & S.T. Toh (ZRC).

240 Remarks. — Specimens from Borneo previously examined and identified as *L. virescens* in
241 Ingrisch (1995) probably belong to either one of our ethospecies collected from Sandakan
242 and Belait or other currently unknown ethospecies. We consider *L. virescens* to occur only in
243 Thai-Malay Peninsula and not in Borneo.

244

245 ***Lipotactes kabili* Tan, Japir and Chung, new species** (Figs. 1.3, 1.4, 2.3, 2.7, 3.3, 4, 5.3, 6.3)

246 Material examined. — Holotype (male, SDK.19.101), East Malaysia, Sabah, Sandakan,
247 Sepilok, Kabili Sepilok Forest Reserve, primary and old secondary dipterocarp forest,
248 N5.87014 E117.93778, 81.8±6.4 m, 2029h, on foliage near ground, 3 October 2019, coll.
249 M.K. Tan & John Lee Yukang (FRC).

250 Paratypes (2 males and 1 female): same locality, 1 female (SDK.19.74), N5.86952
251 E117.93882, 49.3±5.9 m, 2005h, on foliage, 1 October 2019, coll. M.K. Tan, Razy Japir &
252 John Lee Yukang; 1 male (SDK.19.98), N5.86926 E117.93940, 52.7±6.9 m, 1936h, on
253 foliage, 3 October 2019, coll. M.K. Tan & John Lee Yukang; 1 male (SDK.19.102),
254 N5.87039 E117.93772, 82.0±6.2 m, 2042h, on branch, 3 October 2019, coll. M.K. Tan &
255 John Lee Yukang (ZRC).

256 Remarks. — The individuals from Sandakan are remarkably different from other Bornean
257 species in their acoustic signals and in shape of the stridulatory apparatus (especially the
258 stridulatory file). We also compared with *Mortoniellus macrognathus* Ingrisch, 1995 from
259 Poring and Danum Valley in Sabah but differs by shape of cercus (despite the presence of
260 inner process), pronotum and hind femoral colouration and shape of female subgenital plate.
261 We are confident that this population represents a valid species of Lipotactinae.

262 Diagnosis. — This new species differs from all known congeners by the following characters:
263 pronotum with dense black patterns and with metazona brown; male cercus conical, with
264 inner process and with apex styliform; inner process of male cercus elongated and somewhat

265 lamellated, with posterior margin convexly curved but anterior margin straight, with basal
266 lobe large and cylindrical; female subgenital plate distinctly wider than long with apex
267 roundly and narrowly emarginated in middle.

268 Description of male. — Pronotum with metazona prolonged and slightly elevated (Fig. 2.7).
269 Pattern on pronotum dorsal disk similar to that of *L. alienus* and *L. virescens*: anterior margin
270 and transverse sulcus black, a circa M-shaped figure behind sulcus; but differs by: the figure
271 behind sulcus thicker (bolder), and the anterior area of the metazona with a broad U-shaped
272 figure that fades posteriorly into the darkened apex of pronotum (Fig. 2.3). Posterior femur
273 with a reddish stroke that reaches the base and dorsal margin; hind knee darkened, but not as
274 dark as female.

275 Tegmen with dorsal field 3.0 mm long and 2.8–2.9 mm wide; lateral field infumated (Fig.
276 5.3). h1 distinctly less wide than mirror width, h2 narrowly pyriform, with anal end rounded
277 and with frame nearly touching CuPb at the basal end; mirror large and distinctly wider than
278 long (Fig. 5.3). Stridulatory file on CuPb crescent shaped, ca. 2 mm long, with 84 evenly
279 distributed teeth (Fig. 6.3). Apical area of tegmen narrowly truncated (Fig. 5.3).

280 Tenth abdominal tergite and epiproct as in *L. virescens* (Fig. 3.3): tenth abdominal tergite
281 incised in the middle; epiproct triangular, apex obtuse, with a faint medio-longitudinal furrow.

282 Male cercus large and conical, with inner process, apex characteristically styloform; inner
283 process of male cercus elongate and somewhat lamellate, with posterior margin convexly
284 curved but anterior margin straight, apex obtuse; inner process with basal lobe large and
285 cylindrical (Fig. 3.3). Subgenital plate with lateral lobes at apex greatly spaced, posterior
286 margin broadly emarginated. Stylus relatively stout, dark coloured, cylindrical, with apex
287 obtuse (Fig. 3.3). Phallus membranous.

288 Description of female. — Pronotum dorsal disk pattern similar to males, but M-shaped figure
289 even more prominent (Figs. 4.1, 4.2). Wings absent. Posterior femur with a strong black

stroke that reaches the base and dorsal margin; hind knee darkened. Abdominal tergite black laterally. Tenth abdominal tergite with posterior margin truncated in the middle; epiproct tongue-shaped with apex obtuse (Fig. 4.3). Subgenital plate concealed at base by preceding sternite, transverse (distinctly wider than long), tapering towards apex; apex roundly and narrowly emarginated in middle (Fig. 4.4). Ovipositor slightly surpassing posterior femur, straight in basal third, then feebly narrowed and curved dorsad, apical third widening feebly before tapering into acute apex; ventral valve finely serrulated at ventro-apical margin (Fig. 4.5).

Measurements. — See Table 1.

Distribution. — This species is so far known only from Sandakan.

Etymology. — This species name is named after the type locality, Kabili; noun in apposition.

***Lipotactes ethospecies* (Belait)** (Figs. 1.5, 1.6, 2.4, 2.8, 3.4, 5.4, 6.4)

Material examined. — Brunei Darussalam, Belait District: 1 female (BR.19.6), Jalan (= Road) Labi near Andulau Forest Reserve, N4.63354 E114.51073, 76.8±7.9 m, 1947h, under foliage, 24 February 2019, coll. M.K. Tan & H. Yeo; 1 male (BR.19.51), Jalan Labi at Teraja, N4.28456 E114.41866, 40.0±5.7 m, 2005h, on tree sapling branch, 1 March 2019, coll. M.K. Tan & H. Yeo; 1 male (BR.19.56), Jalan Labi at Teraja, N4.28485 E114.41828, 50.1±5.5 m, 2045h, on tree sapling branch, 1 March 2019, coll. M.K. Tan & H. Yeo (UBDM and ZRC).

Remarks. — The two populations from Brunei may represent either two subspecies or species. The males can be distinguished by stridulum morphology, shape of cerci and subgenital plate, patterns on pronotum. However, females from Belait are almost indistinguishable from those from Kuala Belalong. Since we did not have acoustic data for this population, we prefer to putatively consider this population as tentative ethospecies instead of describing a new species to avoid future synonymy or taxonomic confusion.

Part II: stridulum morphology

Tegmen in *Lipotactes* is highly modified: dorsal field with large and well-defined mirror area (among other well-defined resonators) and few veins, usually transparent (although with some darkened veins in *L. kabili n. sp.* and *L. ethospecies* (Belait) , Figs. 5.3, 5.4); apical area of dorsal field and lateral field more infumated (Fig. 5).

Harp area differs between different species in size and shape. h1 can be slender (distinctly less wide than mirror width) in *L. kabili n. sp.* (Fig. 5.3) but very broad (nearly as wide as mirror width) in *L. ethospecies* (Belait) (Fig. 5.4). h2 differs between different species in size: smallest in *L. ethospecies* (Belait) (0.08 mm²) (Fig. 5.4) and largest in *L. kabili n. sp.* (0.12–0.15 mm²) (Fig. 5.3) and *L. alienus* (0.12 mm²) (Fig. 5.1). h2 in *L. kabili n. sp.* are exceptionally narrower and more transverse than other species (Fig. 5.3). We did not compare area of h3 because the vein demarcating the area at the anal end is not always clear.

Mirrors in all species are always wider than long but range from 1.16 times wider than long in *L. alienus* (Fig. 5.1) to 1.46 in *L. kabili n. sp.* (Fig. 5.3); the mirror area also varies in size: ranging from 0.60 mm² in *L. ethospecies* (Belait) (Fig. 5.4) to 1.24–1.27 mm² in *L. kabili n. sp.* (Fig. 5.3). As expected, venation is typically conserved although variation between species is greater than within (Fig. 5).

The stridulatory file on the left tegmen differs drastically between different species. Specifically, it differs by number and distribution teeth in both, inter-tooth distance and tooth length, even though the stridulatory files are generally crescent shaped (Figs. 6, 7). Tooth number is lowest in *L. ethospecies* (Belait) (14), followed by *L. alienus* (28) and *L. virescens*

(39). *L. kabili* n. sp. have distinctively more teeth (84) but the teeth are more densely and evenly distributed than in the other populations (Fig. 7).

Part III: calling songs

The calling songs of *Lipotactes* from Borneo consist of verses (Fig. 8) in the near-ultrasonic to ultrasonic frequencies, thus being faintly audible to human ear. The songs are non-resonant owing to broad distribution of frequencies; but usually consist of two peaks, one higher than the other (Fig. 9). These characters are somewhat similar to the known songs of other Lipotactinae. We also observed that *Lipotactes* exhibit intra-individual variation in their calling songs, as in the case of *L. alienus*. Two song types were considered as the songs have vastly different time domain.

Calling song type I of *L. alienus* from Kuala Belalong (n = 9 songs, 1 male): At 24.4–29.2 °C, the male calling song consists of a verse, average verse duration 790 ± 130 ms (507–997 ms), with variable amplitude modulation in the time domain (Fig. 8.1). Each verse consists of an average of 48 ± 12 (27–70) pulses (probably syllables) and its repetition rate is 61.2 ± 13.0 s⁻¹ (53.2–89.8 s⁻¹). Average pulse duration is 16.9 ± 2.8 ms (11.1–18.8 ms) (Fig. 8.2). Each pulse is made up of a series of closely-spaced impulses (Fig. 8.3). Arithmetic mean frequency is 36.9 ± 0.4 kHz, arithmetic mean peak frequency is 35.5 ± 0.9 kHz and arithmetic mean fundamental frequency is 23.8 ± 3.1 kHz (Fig. 9.1).

Calling song type II of *L. alienus* from Kuala Belalong (n = 8 series of songs, 1 male):

Another song type was identified from the same individual. At 24.4 °C, the song also consists of a continuous series of verses, albeit each verse duration is shorter at an average of 267 ± 80 ms (175–434 ms). Each verse consists of an average of 13 ± 5 (8–24) pulses (probably

syllables) and its repetition rate is $49.5 \pm 2.8 \text{ s}^{-1}$ ($45.7\text{--}55.3 \text{ s}^{-1}$). Average pulse duration is $20.3 \pm 1.1 \text{ ms}$ ($18.1\text{--}21.9 \text{ ms}$). Arithmetic mean frequency is $37.4 \pm 0.4 \text{ kHz}$, arithmetic mean peak frequency is $36.6 \pm 1.2 \text{ kHz}$ and arithmetic mean fundamental frequency is $16.8 \pm 1.1 \text{ kHz}$.

Calling song of *L. kabili* n. sp. from Sandakan (n = 19 songs, 1 male): The song is unique among the known songs of Lipotactinae. At 30.1°C , each male calling song consists of a verse of 8 ± 4 ($3\text{--}15$) crescendoing syllables closely-spaced together in time, song average duration $1222 \pm 565 \text{ ms}$ ($470\text{--}2350 \text{ ms}$) (Fig. 8.4). Each average syllable duration is $14.3 \pm 1.5 \text{ ms}$ ($12.5\text{--}19.9 \text{ ms}$) (Fig. 8.5). Each syllable consists of 26 ± 6 ($21\text{--}45$) rapid-decaying pulses. Average rapid-decay duration is $5.6 \pm 0.7 \text{ ms}$ ($3.2\text{--}6.6 \text{ ms}$) and pulse repetition rate is $183.5 \pm 33.6 \text{ s}^{-1}$ ($152.5\text{--}310.9 \text{ s}^{-1}$) (Fig. 8.6). Arithmetic mean frequency is $38.1 \pm 0.3 \text{ kHz}$, arithmetic mean peak frequency is $28.7 \pm 0.3 \text{ kHz}$ and arithmetic mean fundamental frequency is $20.5 \pm 1.2 \text{ kHz}$ (Fig. 9).

When the temperature at the time of recording was accounted for in the models, the mean peak frequency and mean frequency of the songs are still markedly different between *L. alienus* from Kuala Belalong and *L. kabili* n. sp. from Sandakan (Figs. 9, 10.1, 10.2). As expected, the frequency domain of the two song types belonging to *L. alienus* are not different (Figs. 9, 10.1, 10.2). For the time domain of the song, verse/ syllable duration is drastically different between the two song types of *L. alienus* from Kuala Belalong and *L. kabili* n. sp. from Sandakan (Fig. 10.3). The two song types of *L. alienus* can be differentiated mainly verse/ syllable duration (Fig. 10.3). The pulse duration, and hence pulse repetition rate, is markedly different between *L. alienus* and *L. kabili* n. sp., but not so between the two song types of *L. alienus* (Figs. 10.4, 10.5).

Discussion

Integrative taxonomy of Bornean Lipotactes

Prior to the comparison of call and stridulatory structures between *L. virescens* from South Thailand and Malaysia and the populations from Borneo, we thought that individuals from Borneo are merely variants of *L. virescens* (based on new specimens and Ingrisch [1995]). This is owing to the characteristic male cerci that have a stout lobe at the base of the inner process, seen previously only in *Lipotactes virescens* but also in some *Mortoniellus* species (Ingrisch, 1995). *Lipotactes* specimens from Borneo also exhibit this character with variation in shape and size. We also observed a gradient in colouration and patterns on the pronotum dorsal disk among the different populations: individuals from Belait being most prominent and darkest, followed by those from Sandakan (i.e., *L. kabili* n. sp.), *L. virescens* and individuals from Kuala Belalong being lightest. Although *Lipotactes* congeners display differences in colour intensity and pattern on pronotum, such characters usually call for great caution when used to differentiate species. Other traditional characters such as subgenital plate and styli also show variations among the Bornean populations but it was difficult to ascertain if they represent species differences.

For the first time, the male calling songs of Lipotactinae were recorded from individuals collected in Kuala Belalong (Brunei) and Sandakan (Sabah)—both in Borneo. The songs are very different from one another and from those in South Thailand and Malaysia. Specifically, both the time and the frequency domains of the songs are clearly distinguishable between songs from Kuala Belalong (Brunei) and Sandakan (Sabah). By comparing the new acoustic data with the literature (i.e., Ingrisch, 1995), it was clear that the time domain of the call structure between the Bornean individuals and *L. virescens* from South Thailand and Peninsular Malaysia, even though we could not precisely compare the frequency domain (see

415 Ingrisch, 1995). Specifically, the reported (sensu Ingrisch, 1995) repetition rate of *L.*
 416 *virescens* (ca. 71–101 s⁻¹) is drastically higher than in *L. alienus* but lower than individuals
 417 from Sandakan. The peak frequency is at 35–40 kHz, which is comparable to *L. alienus* but
 418 higher than individuals from Sandakan.
 419
 420 That the bioacoustics signals differ profoundly shed new lights into the species boundaries of
 421 *L. virescens*. This prompted us to re-evaluate the morphology of the Bornean specimens more
 422 closely and eventually helped us cluster the specimens into three groups: (1) population from
 423 Belait, (2) population from Kuala Belalong (probably *L. alienus*) and (3) population from
 424 Sandakan (here described as *L. kabili* n. sp.). Upon closer examination, the specimens from
 425 Borneo consistently differ from *L. virescens* from the Thai-Malay Peninsula by larger inner
 426 process and basal lobe, in addition to the call structures. Individuals from Belait also
 427 invariably have the inner process more strongly curved than other populations. All
 428 individuals from Sandakan have distinctly styliform cerci and the inner process more
 429 lamellate than other populations. The styli in individuals from Belait and Kuala Belalong are
 430 more slender whereas in Sandakan individuals they are most the stout among all populations.
 431
 432 ***Stridulation and stridulum morphology in Lipotactes***
 433 The stridulum morphology of *L. virescens* and the Bornean groups are found to be different
 434 and can be used to differentiate species even though the characters were not formerly used for
 435 taxonomy of Lipotactinae. The dissimilarities in the radiating structure, including the harp
 436 and mirror, may have implications on the differences in call structure. The properties of the
 437 membrane structure and frame structure of the mirror can dictate the spectral energy, and thus
 438 the resulting frequencies, of the song in katydids (Bennett-Clark, 2003; Sarria-S et al., 2016;
 439 Chivers et al., 2017). In *Lipotactes*, the mirror on the left tegmen is seemingly functional.

440 This illustrates that the asymmetrical tegminal system in which a single radiating structure on
441 the right tegmen facilitates the production of tonal, extreme ultrasonic songs (Montealegre-Z
442 & Postles, 2010; Sarria-S et al., 2016; Chivers et al., 2017), is not necessarily typical in
443 modern katydids. Instead, a symmetrical tegminal system may be typical for short-winged
444 species with tegmina covered by pronotum (e.g., *Afroanthracites* in Hemp et al., 2015). Now
445 that we have established empirical evidence that both the frequency domain of the call
446 structure and the resonator can be different between populations of *Lipotactes*, the functions
447 of the left mirror and harp area warrant further investigation to enhance our knowledge on the
448 biomechanics of sound production in Lipotactinae.

449

450 The morphology of the stridulatory files in Bornean *Lipotactes* is also diverse. Most
451 interestingly, stridulatory files can range from those with a few large and separated teeth (i.e.,
452 *L. alienus* and individuals from Belait) to those with systematically distributed teeth (i.e.,
453 individuals from Sandakan). For *L. alienus* and individuals from Belait, the few teeth may
454 help distort and recoil scraper at high speed to produce ultrasonic songs, typical of katydids
455 that sing at extreme ultrasonic frequencies (Montealegre-Z & Mason, 2005; Montealegre-Z,
456 et al., 2006; Tan et al., 2019). For the individuals from Sandakan, the file morphology is more
457 representative of many ensiferans in which the systematic distribution of teeth helps maintain
458 constant tooth rate strike (e.g., Robillard et al., 2013; Heller et al., 2017; Heller & Hemp,
459 2018). This may explain why the time domain of the song of Sandakan individuals is distinct
460 from that of *L. alienus* even though we did not find both extreme ultrasonic songs and lower
461 frequency songs respectively. We postulate that there may exist more than one type of sound
462 production mechanism among morphologically cryptic *Lipotactes* from Borneo and
463 recommend further investigation into the biomechanics of the stridulatory file. This may

include examining the wing movements using high-speed cameras and detecting for resilin in scraper (Montealegre-Z et al., 2006).

We observed that there is a marked disparity in the call structure and stridulum morphology among different Bornean *Lipotactes* as compared to the cryptic morphology of their male cerci. This may be explained by potentially complicated communication preceding mating. We speculate that the males sing for the females to recognise and locate them (probably with aid of the visual sense during dusk time or day time as known from other Lipotactini [e.g., some *Mortoniellus* species] [Ingrisch, 1995]) and assess the mate quality as typical in other katydids (Heller et al., 2017). Furthermore, vibrational communication in ensiferans is also more common than previously thought (Morris et al., 1995; ter Hofstede, 2015). It will not be a surprise that some Lipotactinae exhibit such communication given that some of them are leaf-dwellers (and leaf are excellent medium for such communication).

Conclusions

Our findings offered insights into the usefulness of stridulation and stridulum morphology for separating species of *Lipotactes*. We found that inter-population differences in stridulation and stridulum morphology are generally larger than intra-population differences, even though our sample size is somewhat small (attributed to the difficulty of collecting males). This oppose to postulation by Ingrisch (1995) that there is probably a lack of evolutionary pressure in Lipotactinae for the displacement of call structures, especially since the visual sense is seemingly well developed. The morphological, anatomical and acoustic evidence support the assumption that the populations from Belait, Kuala Belalong and Sandakan probably belong to a species-complex that also includes the described species *L. alienus* and *L. virescens*. We

are also confident enough to consider the individuals from Sandakan as a unique species: *L. kabili* n. sp.

However, our findings also raised new questions with regard to the Lipotactinae. The shape of male cercus with the long inner process and shorter lobe is typical of some *Mortoniellus* from Borneo and Sulawesi and not unique for the *L. alienus-virescens* species complex. This led to whether the shape of the male cercus is a good character to differentiate between species and whether similarity in the shape of the male cercus is congruent with the phylogeny. If not, we will need to consider whether the arrangement into genera needs to be evaluated anew. The study of more specimens from other parts of Borneo and a wider area of Southeast Asia, along with the incorporation of phylogenetic analyses will further validate the status of Bornean *Lipotactes*.

Acknowledgements

The authors thank Fernando Montealegre-Zapata for providing an Echo Meter Touch 1 unit for acoustic recording and for providing feedback to the manuscript prior to submission; Huiqing Yeo (in Brunei Darussalam), Siew Tin Toh (in Bukit Larut and Sandakan), Momin Binti, John Lee Yukang and Saudi Bintang (in Sandakan) for field assistance. The permissions for collecting specimens were granted by the Forestry Department, Ministry of Primary Resources and Tourism, Brunei Darussalam (JPH/PDK/01 Pt 2) and the Sabah Biodiversity Centre (JKM/MBS.1000-2/3 JLD.3 (99)) (for Sandakan). The work of MKT was supported by the Orthoptera Species File Grant 2019 under the taxonomic research project titled “Contribution to the species diversity and acoustic data on Orthoptera from Sandakan

(Borneo, East Malaysia, Sabah)”; and Percy Sladen Memorial Fund (The Linnean Society of London) under the project titled “Advancing biodiversity informatics of Orthoptera from Brunei Darussalam”. The bioacoustics component was supported by the Wildlife Acoustics Scientific Product Grant 2019 under the project titled “Discovery of Ultrasonic Singing Katydid in Southeast Asia”.

References

- Araya-Salas, M., & Smith-Vidaurre, G. (2017). warbleR: an R package to streamline analysis of animal acoustic signals. *Methods in Ecology and Evolution*, 8(2), 184–191.
- Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., & Dai, B. (2014). lme4: linear mixed-effects models using Eigen and S4 (Version 1.1-7).
- Bennet-Clark, H. C. (2003). Wing resonances in the Australian field cricket *Teleogryllus oceanicus*. *Journal of Experimental Biology*, 206, 1479–1496.
- Béthoux, O. (2012). Grylloptera—a unique origin of the stridulatory file in katydids, crickets, and their kin (Archaeorthoptera). *Arthropod Systematics & Phylogeny*, 70(1), 43–68.
- Chang, Y. L., Shi, F. M., & Ran, J. C. (2005). Descriptions of two new species of Lipotactes Brunner v. Watt. (Orthoptera: Tettigoniidae) from China. *Oriental Insects*, 39, 353–357.
- Chivers, B. D., Béthoux, O., Sarria-S, F. A., Jonsson, T., Mason, A. C., & Montealegre-Z, F. (2017). Functional morphology of tegmina-based stridulation in the relict species *Cyphoderris monstrosa* (Orthoptera: Ensifera: Prophalangopsidae). *Journal of Experimental Biology*, 220(6), 1112–1121.
- Cigliano, M. M., Braun, H., Eades, D. C., & Otte, D. (2019). Orthoptera Species File Online. Version 5 (5.0). Retrieved from:

538 <http://orthoptera.speciesfile.org/HomePage/Orthoptera/HomePage.aspx> (accessed 10
539 October 2019).

540 Feng, J. Y., Zhou, Z. J., Chang, Y. L., & Shi, F. M. (2017). Remarks on the genus *Lipotactes*
541 Brunner v. W., 1898 (Orthoptera: Tettigoniidae: Lipotactinae) from China. *Zootaxa*,
542 4291(1), 183–191.

543 Gorochov, A. V. (1993). Two new species of the genus *Lipotactes* from Vietnam (Orthoptera:
544 Tettigoniidae). *Zoosystematica Rossica*, 2(1), 59–62.

545 Gorochov, A. V. (1996). New and little known species of the genus *Lipotactes* Br.-W.
546 (Orthoptera, Tettigoniidae) from Vietnam. *Entomologicheskoe Obozrenie*, 75(1), 32–38.

547 Gorochov, A. V. (1998). A new species of the genus *Lipotactes* from Cambodia (Orthoptera:
548 Tettigoniidae). *Zoosystematica Rossica*, 7(1), 132.

549 Gorochov, A. V. (2010). New and little-known Orthopteroid insects (Polyneoptera) from
550 fossil resins: Communication 4. *Paleontological Journal*, 44(6), 657–671.

551 Heller, K. G., & Hemp, C. (2019). Extremely divergent song types in the genus *Aerotegmina*
552 Hemp (Orthoptera: Tettigoniidae: Hexacentrinae) and the description of a new species
553 from the Eastern Arc Mountains of Tanzania (East Africa). *Bioacoustics*, 28(3), 269–285.

554 Heller, K. G., Ingrisch, S., Liu, C. X., Shi, F. M., Hemp, C., Warchałowska-Śliwa, E., &
555 Rentz, D. C. (2017). Complex songs and cryptic ethospecies: the case of the *Ducetia*
556 *japonica* group (Orthoptera: Tettigoniidae: Phaneropteridae: Phaneropterinae).
557 *Zoological Journal of the Linnean Society*, 181(2), 286–307.

558 Hemp, C., Heller, K. G., Warchałowska-Śliwa, E., Grzywacz, B., & Hemp, A. (2015).
559 Ecology, acoustics and chromosomes of the East African genus *Afroanthracites* Hemp &
560 Ingrisch (Orthoptera, Tettigoniidae, Conocephalinae, Agraeciini) with the description of
561 new species. *Organisms Diversity & Evolution*, 15(2), 351–368.

562 Ingrisch, S. (1995). Revision of the Lipotactinae, a new subfamily of Tettigoniioidea
563 (Ensifera). *Entomologica Scandinavica*, 26, 273–320.

564 Montealegre-Z, F. (2009). Scale effects and constraints for sound production in katydids
565 (Orthoptera: Tettigoniidae): generator size constrains signal parameters. *Journal of*
566 *Evolutionary Biology*, 22(2), 355–366.

567 Montealegre-Z, F. (2012). Reverse stridulatory wing motion produces highly resonant calls in
568 a neotropical katydid (Orthoptera: Tettigoniidae: Pseudophyllinae). *Journal of Insect*
569 *Physiology*, 58, 116–124.

570 Montealegre-Z, F., & Mason, A. C. (2005). The mechanics of sound production in
571 *Panacanthus pallicornis* (Orthoptera: Tettigoniidae: Conocephalinae): the stridulatory
572 motor patterns. *Journal of Experimental Biology*, 208(7), 1219–1237.

573 Montealegre-Z, F., Morris, G. K., & Mason, A. C. (2006). Generation of extreme ultrasonics
574 in rainforest katydids. *Journal of Experimental Biology*, 209, 4923–4937.

575 Montealegre-Z, F., Ogden, J., Jonsson, T., & Soulsbury, C. D. (2017). Morphological
576 determinants of carrier frequency signal in katydids (Orthoptera): a comparative analysis
577 using biophysical evidence of wing vibration. *Journal of Evolutionary Biology*, 30(11),
578 2068–2078.

579 Montealegre-Z, F., & Postles, M. (2010). Resonant sound production in *Copiphora*
580 *gorgonensis* (Tettigoniidae: Copiphorini), an endemic species from Parque Nacional
581 Natural Gorgona, Colombia. *Journal of Orthoptera Research*, 19, 347–355.

582 Morris, G. K., Mason, A. C., Wall, P., & Belwood, J. J. (1994). High ultrasonic and
583 tremulation signals in neotropical katydids (Orthoptera: Tettigoniidae). *Journal of*
584 *Zoology*, 233(1), 129–163.

585 Mugleston, J. D., Naegle, M., Song, H., & Whiting, M. F. (2018). A comprehensive
586 phylogeny of Tettigoniidae (Orthoptera: Ensifera) reveals extensive ecomorph

587 convergence and widespread taxonomic incongruence. *Insect Systematics and Diversity*,
588 2(4), 5.

589 R Core Team. (2018). R: A language and environment for statistical computing. R
590 Foundation for Statistical Computing, Vienna, Austria.

591 Robillard, T., Montealegre-Z, F., Desutter-Grandcolas, L., Grandcolas, P., & Robert, D.
592 (2013). Mechanisms of high-frequency song generation in brachypterous crickets and the
593 role of ghost frequencies. *Journal of Experimental Biology*, 216(11), 2001–2011.

594 Sarria-S, F. A., Buxton, K., Jonsson, T., & Montealegre-Z, F. (2016). Wing mechanics,
595 vibrational and acoustic communication in a new bush-cricket species of the genus
596 *Copiphora* (Orthoptera: Tettigoniidae) from Colombia. *Zoologischer Anzeiger-A Journal*
597 *of Comparative Zoology*, 263, 55–65.

598 Shi, F. M., & Li, R. L. (2009). A review of the genus *Lipotactes* Brunner v. W., 1898
599 (Orthoptera, Tettigoniidae, Lipotactinae) from China. *Zootaxa*, 2152, 36–42

600 Tan, M. K., & Kamaruddin, K. N. (2016). A contribution to the knowledge of Orthoptera
601 diversity from Peninsular Malaysia: Bukit Larut, Perak. *Zootaxa*, 4111(1), 21–40.

602 Tan, M. K., Montealegre-Z, F., Wahab, R. A., Lee, C. Y., Belabut, D. M., Japir, R., & Chung
603 A. Y. C. (2019). Ultrasonic songs and stridulum anatomy of *Asiophlugis* crystal
604 predatory katydids (Tettigoniioidea: Meconematinae: Phlugidini). *Bioacoustics*.

605 Tan, M. K., & Wahab, R. A. (2018). Preliminary study on the diversity of Orthoptera from
606 Kuala Belalong Field Studies Centre, Brunei Darussalam, Borneo. *Journal of Orthoptera*
607 *Research*, 27(2), 119–142.

608 ter Hofstede, H. M., Schöneich, S., Robillard, T., & Hedwig, B. (2015). Evolution of a
609 communication system by sensory exploitation of startle behaviour. *Current Biology*,
610 25(24), 3245–3252.

Wang, T., & Shi, F. (2020). Potential effect of the global warming to the subfamily
 Lipotactinae (Orthoptera: Tettigoniidae) in China and acoustics data of *Lipotactes*
truncatus Shi & Li, 2009. *Journal of Asia-Pacific Entomology*, 23(1), 146–151.

Table 1. Measurements (all in mm) of *Lipotactes kabili* n. sp.

	BL	HL	HW	PL	PW	TL	HFL	HTL	OL
Male holotype	10.1	1.6	4.9	4.1	3.2	2.4	10.7	9.4	—
Male paratypes (n = 2)	9.6– 10.9	1.5– 2.0	5.0– 5.2	4.2– 4.4	3.3– 3.6	2.4	10.9	9.8	—
Female paratype (n = 1)	9.9	2.0	5.0	2.6	3.2	—	11.5	9.5	7.1

Figure captions

Figure 1. Morphologically similar *Lipotactes* in the natural environment from different parts of Borneo: *L. alienus* female from Kuala Belalong (1), *L. virescens* male from Bukit Larut (2), *L. kabili* **n. sp.** male (3) and female (4), and *L. ethospecies* (Belait) males (5, 6).

Figure 2. Habitus of dry-pinned male *Lipotactes* in dorsal (1–4) and lateral (5–8) views: *L. alienus* (1, 5), *L. virescens* (2, 6), *L. kabili* **n. sp.** (3, 7), and *L. ethospecies* (Belait) (4, 8). Scale bars: 5 mm.

Figure 3. Abdominal apex of male *Lipotactes*: *L. alienus* (1), *L. virescens* (2), *L. kabili* **n. sp.** (3), and *L. ethospecies* (Belait) (4). Scale bars: 1 mm.

Figure 4. *Lipotactes kabili* **n. sp.** female: habitus in dorsal (1) and lateral (2) views; abdominal apex in dorsal (3) and ventral (4) views; ovipositor in lateral view (5). Scale bars: 5 mm (1, 2); 2 mm (3–6).

Figure 5. Left tegmen of *Lipotactes* in ventral view: *L. alienus* (1), *L. virescens* (2), *L. kabili* **n. sp.** (3), and *L. ethospecies* (Belait) (4). Scale bars: 1 mm.

Figure 6. Stridulatory file on left tegmen of *Lipotactes* in ventral view: *L. alienus* (1), *L. virescens* (2), *L. kabili* **n. sp.** (3), and *L. ethospecies* (Belait) (4). Scale bars: 0.2 mm.

Figure 7. Tooth distribution on the stridulatory file of *Lipotactes* based on inter-tooth distance (1) and tooth length variations (2). Trend lines are generalised additive model fits. The tooth number corresponds to the first tooth from the anal end to the last tooth at the basal end of the left stridulatory file.

Figure 8. Oscillograms of the *Lipotactes* songs: *L. alienus* (1–3) and *L. kabili* **n. sp.** (4–6).

Figure 9. Power spectra of the *Lipotactes* songs: *L. alienus* (1) and *L. kabili* **n. sp.** (2). The relative amplitude scale is scaled between 0 and 1.

Figure 10. Mean plots on the comparison of the time and frequency domains of the *Lipotactes* song. The dot represents the least-square means and the horizontal line represents the 95% confidence interval (CI).