

**EXISTENCE OF ALMOST AUTOMORPHIC
SOLUTION IN DISTRIBUTION FOR A CLASS OF
STOCHASTIC INTEGRO-DIFFERENTIAL
EQUATION DRIVEN BY LÉVY NOISE**

Mamadou Moustapha Mbaye, Solym Manou-Abi

► **To cite this version:**

Mamadou Moustapha Mbaye, Solym Manou-Abi. EXISTENCE OF ALMOST AUTOMORPHIC SOLUTION IN DISTRIBUTION FOR A CLASS OF STOCHASTIC INTEGRO-DIFFERENTIAL EQUATION DRIVEN BY LÉVY NOISE. 2020. hal-02974673

HAL Id: hal-02974673

<https://hal.archives-ouvertes.fr/hal-02974673>

Preprint submitted on 22 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**EXISTENCE OF ALMOST AUTOMORPHIC SOLUTION IN DISTRIBUTION
FOR A CLASS OF STOCHASTIC INTEGRO-DIFFERENTIAL EQUATION
DRIVEN BY LÉVY NOISE**

MAMADOU MOUSTAPHA MBAYE AND SOLYM MAWAKI MANOU-ABI

ABSTRACT. We investigate a class of stochastic integro-differential equations driven by Lévy noise. Under some appropriate assumptions, we establish the existence of a square-mean almost automorphic solutions in distribution. Particularly, based on Schauder's fixed point theorem, the existence of square-mean almost automorphic mild solution distribution is obtained by using the condition which is weaker than Lipschitz conditions. We provide an example to illustrate our results.

1. INTRODUCTION

The aim of this work is to study the existence and uniqueness of the square-mean almost automorphic mild solutions in distribution to the following class of nonlinear stochastic integro-differential equations driven by Lévy noise in a separable Hilbert space H

$$(1.1) \quad \begin{aligned} x'(t) &= Ax(t) + g(t, x(t)) + \int_{-\infty}^t B_1(t-s)f(s, x(s))ds \\ &+ \int_{-\infty}^t B_2(t-s)h(s, x(s))dW(s) \\ &+ \int_{-\infty}^t B_2(t-s) \int_{|y|_V < 1} F(s, x(s-), y)\tilde{N}(ds, dy) \\ &+ \int_{-\infty}^t B_2(t-s) \int_{|y|_V \geq 1} G(s, x(s-), y)N(ds, dy) \quad \text{for all } t \in \mathbb{R}, \end{aligned}$$

where $A : D(A) \subset H$ is the infinitesimal generator of a C_0 -semigroup $(T(t))_{t \geq 0}$, B_1 and B_2 are convolution-type kernels in $L^1(0, \infty)$ and $L^2(0, \infty)$ respectively. $g, f : \mathbb{R} \times L^2(P, H) \rightarrow L^2(P, H)$ $h : \mathbb{R} \times L^2(P, H) \rightarrow L(V, L^2(P, H))$ $F, G : \mathbb{R} \times L^2(P, H) \times V \rightarrow L^2(P, H)$; W and N are the Lévy-Itô decomposition components of the two-sided Lévy process L (with assumptions stated in Section 2.).

Throughout this work, we assume $(H, \|\cdot\|)$ and $(V, |\cdot|)$ are real separable Hilbert spaces. We denote by $L(V, H)$ the family of bounded linear operators from V to H and $L^2(P, H)$ is the space of all H -valued random variables x such that

$$\mathbb{E}\|x\|^2 = \int_{\Omega} \|x\|^2 dP < +\infty.$$

2000 *Mathematics Subject Classification.* 34C27; 34K14; 34K30; 34K50; 35B15; 35K55; 43A60; 60G20.

Key words and phrases. almost automorphic solution; stochastic processes; stochastic evolution equations; Lévy noise.

The concept of almost automorphic is a natural generalization of the almost periodicity that was introduced by Bochner [6]. The basic aspects of the theory of almost automorphic functions can be found for instance in to the book [21].

In recent years, the study of almost periodic or almost automorphic solutions to some stochastic differential equations have been considerably investigated in lots of publications [2, 3, 4, 5, 7, 8, 9, 10, 11, 13, 24] because of its significance and applications in physics, mechanics and mathematical biology. The concept of square-mean almost automorphic stochastic processes was introduced by Fu and Liu [14]. As indicated in [15, 16], it appears that almost periodicity or automorphy in distribution sense is a more appropriate concept relatively to solutions of stochastic differential equations. Recently, the concept of Poisson square-mean almost automorphy was introduced by Liu and Sun [17] to deal with some stochastic evolution equations driven by Lévy noise. For the almost automorphy in distribution, its various extensions in distribution sense and the applications in stochastic differential equations, one can see [12, 19, 26] for more details.

One should point out that other slightly different versions of equation (1.1) have been considered in the literature. In particular, Bezandry [3], Xia [25] and Mbaye [18] investigated the existence and uniqueness of the solution of equation (1.1) in the case when $F = G = 0$ based on another fixed point theorem.

The rest of this work is organized as follows. In Section 2, we make a recalling on Lévy process. In Section 3, we review some concepts and basic properties on almost automorphic and Poisson almost automorphic processes. In Section 4, by Schauder's fixed point theorem, we prove the existence of an square-mean almost automorphic mild solution in distribution of equation (1.1). In Section 5, we provide an example to illustrate our results.

2. LÉVY PROCESS

Definition 2.1. A V -valued stochastic process $L = (L(t), t \geq 0)$ is called Lévy process if:

- (1) $L(0) = 0$ almost surely;
- (2) L has independent and stationary increments;
- (3) L is stochastically continuous, i.e. for all $\epsilon > 0$ and for all $s > 0$

$$\lim_{t \rightarrow s} P(|L(t) - L(s)| > \epsilon) = 0.$$

Every Lévy process is càdlàg.

Let L be a Lévy process.

Definition 2.2. [17]

- (1) A borel B in $V - \{0\}$ is bounded below if $0 \notin \bar{B}$, where \bar{B} is the closure of B .
- (2) $\nu(\cdot) = \mathbb{E}(N(1, \cdot))$ is called the intensity measure associated with L , where $\Delta L(t) = L(t) - L(t-)$ for each $t \geq 0$ and

$$N(t, B)(\omega) := \#\left\{0 \leq s \leq t : \Delta L(s)(\omega) \in B\right\} := \sum_{0 \leq s \leq t} \chi_B(\Delta L(s)(\omega))$$

with $L(t-) = \lim_{t \nearrow s} L(s)$ and χ_B being the indicator function for any Borel set B in $V - \{0\}$.

- (3) $N(t, B)$ is called Poisson random measure if B is bounded below, for each $t \geq 0$.
- (4) For each $t \geq 0$ and B bounded below, we define the compensated Poisson random measure by

$$\tilde{N}(t, B) = N(t, B) - t\nu(B).$$

Proposition 2.1. [1, 22] *Let L be the V -valued Lévy process. Then there exist $a \in V$, V -valued Wiener process W with covariance operator Q , and an independent Poisson random measure on $\mathbb{R}^+ \times (V - \{0\})$ such that for each $t \geq 0$*

$$L(t) = at + W(t) + \int_{|x|<1} x\tilde{N}(t, dx) + \int_{|x|\geq 1} xN(t, dx),$$

where the Poisson random measure N has the intensity measure ν satisfying

$$(2.1) \quad \int_V (|x|^2 \wedge 1)\nu(dx) < \infty$$

and \tilde{N} is the compensated Poisson random measure of N .

Let $L_1(t)$ and $L_2(t)$, $t \geq 0$ be two independent and identically distributed Lévy processes. Let

$$L(t) = \begin{cases} L_1(t) & \text{for } t \geq 0, \\ -L_2(-t) & \text{for } t < 0 \end{cases}$$

Remark 2.1. By (2.1), it follows that $\int_{|x|\geq 1} \nu(dx) < \infty$. For convenience, we denote

$$b := \int_{|x|\geq 1} \nu(dx).$$

Then L is a two-sided Lévy process defined on the filtered probability space $(\Omega, \mathcal{F}, P, (\mathcal{F}_t)_{t \in \mathbb{R}})$. We assume that Q is a positive, self-adjoint and trace class operator on V , see [23] for more details. The stochastic process $\tilde{L} = (\tilde{L}(t), t \in \mathbb{R})$ given by $\tilde{L}(t) := L(t+s) - L(s)$ for some $s \in \mathbb{R}$ is also a two-sided Lévy process which shares the same law as L . For more details about the Lévy process, we refer to [1, 17, 22].

3. SQUARE-MEAN ALMOST AUTOMORPHIC PROCESS

In this section, we recall the concepts of square-mean almost automorphic process and there basic properties.

Definition 3.1. Let $x : \mathbb{R} \rightarrow L^2(P, H)$ be a stochastic process.

- (1) x is said to be stochastically bounded if there exists $M > 0$ such that

$$\mathbb{E}\|x(t)\|^2 \leq M \quad \text{for all } t \in \mathbb{R}.$$

- (2) x is said to be stochastically continuous if

$$\lim_{t \rightarrow s} \mathbb{E}\|x(t) - x(s)\|^2 = 0 \quad \text{for all } s \in \mathbb{R}.$$

Denote by $SB C(\mathbb{R}, L^2(P, H))$ the space of all the stochastically bounded and continuous processes. Clearly, the space $SB C(\mathbb{R}, L^2(P, H))$ is a Banach space equipped with the following norm

$$\|x\|_\infty = \sup_{t \in \mathbb{R}} (\mathbb{E}\|x(t)\|^2)^{\frac{1}{2}}.$$

Definition 3.2. [17] Let $J : \mathbb{R} \times V \rightarrow L^2(P, H)$ be a stochastic process.

- (1) J is said to be Poisson stochastically bounded if there exists $M > 0$ such that

$$\int_V \mathbb{E}\|J(t, x)\|^2 \nu(dx) \leq M \quad \text{for all } t \in \mathbb{R}.$$

(2) J is said to be Poisson stochastically continuous if

$$\lim_{t \rightarrow s} \int_V \mathbb{E} \|J(t, x) - J(s, x)\|^2 \nu(dx) = 0 \quad \text{for all } s \in \mathbb{R}.$$

Denote by $PSBC(\mathbb{R} \times V, L^2(P, H))$ the space of all the stochastically bounded and continuous processes.

Definition 3.3. [14] Let $x : \mathbb{R} \rightarrow L^2(P, H)$ be a continuous stochastic process. x is said be square-mean almost automorphic process if for every sequence of real numbers $(t'_n)_n$ we can extract a subsequence $(t_n)_n$ such that, for some stochastic process $y : \mathbb{R} \rightarrow L^2(P, H)$, we have

$$\lim_{n \rightarrow +\infty} \mathbb{E} \|x(t + t_n) - y(t)\|^2 = 0 \quad \text{for all } t \in \mathbb{R}$$

and

$$\lim_{n \rightarrow +\infty} \mathbb{E} \|y(t - t_n) - x(t)\|^2 = 0 \quad \text{for all } t \in \mathbb{R}.$$

We denote the space off all such stochastic processes by $SAA(\mathbb{R}, L^2(P, H))$.

Theorem 3.1. [14] $SAA(\mathbb{R}, L^2(P, H))$ equipped with the norm $\|\cdot\|_\infty$ is a Banach space.

Definition 3.4. [17] Let $D : \mathbb{R} \times V \rightarrow L^2(P, H)$ be stochastic process. D is said be Poisson square-mean almost automorphic process in $t \in \mathbb{R}$ if D is Poisson continuous and for every sequence of real numbers $(t'_n)_n$ we can extract a subsequence $(t_n)_n$ such that, for some stochastic process $\tilde{D} : \mathbb{R} \times V \rightarrow L^2(P, H)$ with $\int_V \mathbb{E} \|\tilde{D}(t, x)\|^2 \nu(dx) < \infty$ such that

$$\lim_{n \rightarrow +\infty} \int_V \mathbb{E} \|D(t + t_n, x) - \tilde{D}(t, x)\|^2 \nu(dx) = 0 \quad \text{for all } t \in \mathbb{R}$$

and

$$\lim_{n \rightarrow +\infty} \int_V \mathbb{E} \|\tilde{D}(t - t_n, x) - D(t, x)\|^2 \nu(dx) = 0 \quad \text{for all } t \in \mathbb{R}.$$

We denote the space off all such stochastic processes by $PSAA(\mathbb{R} \times V, L^2(P, H))$.

Definition 3.5. [17] Let $F : \mathbb{R} \times L^2(P, H) \times V \rightarrow L^2(P, H)$ be stochastic process. F is said be Poisson square-mean almost automorphic process in $t \in \mathbb{R}$ for each $Y \in L^2(P, H)$ if F is Poisson continuous and for every sequence of real numbers $(t'_n)_n$ we can extract a subsequence $(t_n)_n$ such that, for some stochastic process $\tilde{F} : \mathbb{R} \times L^2(P, H) \times V \rightarrow L^2(P, H)$ with $\int_V \mathbb{E} \|\tilde{F}(t, Y, x)\|^2 \nu(dx) < \infty$ such that

$$\lim_{n \rightarrow +\infty} \int_V \mathbb{E} \|F(t + t_n, Y, x) - \tilde{F}(t, Y, x)\|^2 \nu(dx) = 0 \quad \text{for all } t \in \mathbb{R}$$

and

$$\lim_{n \rightarrow +\infty} \int_V \mathbb{E} \|\tilde{F}(t - t_n, Y, x) - F(t, Y, x)\|^2 \nu(dx) = 0 \quad \text{for all } t \in \mathbb{R}.$$

We denote the space off all such stochastic processes by $PSAA(\mathbb{R} \times L^2(P, H) \times V, L^2(P, H))$.

Let $\mathcal{P}(H)$ be the space of all Borel probability measures on H with the β metric.

$$\beta(\mu, \nu) := \sup \left\{ \left| \int f d\mu - \int f d\nu \right| : \|f\|_{BL} \leq 1 \right\}, \quad \mu, \nu \in \mathcal{P}(H),$$

where f are Lipschitz continuous real-valued functions on H with

$$\|f\|_{BL} = \|f\|_L + \|f\|_\infty, \quad \|f\|_L = \sup_{x \neq y} \frac{|f(x) - f(y)|}{\|x - y\|}, \quad \|f\|_\infty = \sup_{x \in H} |f(x)|.$$

Definition 3.6. [17] An H -valued stochastic process $Y(t)$ is said to be almost automorphic in distribution if its law $\mu(t)$ is a $\mathcal{P}(H)$ -valued almost automorphic mapping, i.e. for every sequence of real numbers $(s'_n)_n$, there exist a subsequence $(s_n)_n$ and a $\mathcal{P}(H)$ -valued mapping $\tilde{\mu}(t)$ such that

$$\lim_{n \rightarrow \infty} \beta(\mu(t + s_n), \tilde{\mu}(t)) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \beta(\tilde{\mu}(t - s_n), \mu(t)) = 0$$

hold for each $t \in \mathbb{R}$.

To study the existence of mild solutions to the stochastic evolution equations (1.1). we will need the following assumptions,

(H.1) The semigroup $T(t)$ is compact for $t > 0$ and is exponentially stable, i.e., there exists constants $K, \omega > 0$ such that

$$(3.1) \quad \|T(t)\| \leq K e^{-\omega t} \quad \text{for all } t \geq 0.$$

(H.2) The functions f, g and h are uniformly continuous on any bounded subset K of $L^2(\Omega, H)$ for each $t \in \mathbb{R}$. F, G are uniformly continuous on any bounded subset K of $L^2(\Omega, H)$ for each $t \in \mathbb{R}$ and $x \in V$. For each bounded subset $K \subset L^2(\Omega, H)$, $g(\mathbb{R}, K)$, $f(\mathbb{R}, K)$, $h(\mathbb{R}, K)$ are bounded and $F(\mathbb{R}, K, V)$ and $G(\mathbb{R}, K, V)$ are Poisson stochastically bounded. Moreover we suppose that there exists $r > 0$ such that

$$(3.2) \quad \Delta_r \leq \frac{\omega^2 r}{20\theta K^2}$$

where

$$\Delta_r = \max \left\{ \sup_{t \in \mathbb{R}, \|u\|_{L^2} \leq r} \|f(t, u)\|_{L^2}, \sup_{t \in \mathbb{R}, \|u\|_{L^2} \leq r} \|g(t, u)\|_{L^2}, \sup_{t \in \mathbb{R}, \|u\|_{L^2} \leq r} \|h(t, u)\|_{L^2}, \right. \\ \left. \sup_{t \in \mathbb{R}, \|u\|_{L^2} \leq r} \int_{|y|_V < 1} \|F(t, u, x)\|_{L^2}^2 \nu(dx), \sup_{t \in \mathbb{R}, \|u\|_{L^2} \leq r} \int_{|y|_V \geq 1} \|G(t, u, x)\|_{L^2}^2 \nu(dx) \right\}$$

and

$$\theta = \max \left(1, \|B_1\|_{L^1(0, \infty)}^2, 4\|B_2\|_{L^2(0, \infty)}^2, 2b\|B_2\|_{L^1(0, \infty)}^2 \right)$$

(H.3) we suppose that there exist measurable functions $m_g, m_f, m_h, m_F, m_G : \mathbb{R} \rightarrow [0, \infty)$ such that

$$(3.3) \quad \mathbb{E} \|g(t, Y) - g(t, Z)\|^2 \leq m_g(t) \cdot \mathbb{E} \|Y - Z\|^2$$

$$(3.4) \quad \mathbb{E} \|f(t, Y) - f(t, Z)\|^2 \leq m_f(t) \cdot \mathbb{E} \|Y - Z\|^2$$

$$(3.5) \quad \mathbb{E} \|(h(t, Y) - h(t, Z))Q^{\frac{1}{2}}\|_{L(V, L^2(P, H))}^2 \leq m_h(t) \cdot \mathbb{E} \|Y - Z\|^2$$

$$(3.6) \quad \int_{|x|_V < 1} \mathbb{E} \|F(t, Y, x) - F(t, Z, x)\|^2 \nu(dx) \leq m_F(t) \cdot \mathbb{E} \|Y - Z\|^2$$

$$(3.7) \quad \int_{|x|_V \geq 1} \mathbb{E} \|G(t, Y, x) - G(t, Z, x)\|^2 \nu(dx) \leq m_G(t) \cdot \mathbb{E} \|Y - Z\|^2$$

for all $t \in \mathbb{R}$ and for any $Y, Z \in L^2(P, H)$.

(H.4) If $(u_n)_{n \in \mathbb{N}} \subset SBC(\mathbb{R}, L^2(P, H))$ is uniformly bounded and uniformly convergent upon every compact subset of \mathbb{R} , then $g(\cdot, u_n(\cdot)), f(\cdot, u_n(\cdot)), h(\cdot, u_n(\cdot)),$ and $F(\cdot, u_n(\cdot), \cdot), G(\cdot, u_n(\cdot), \cdot)$ are relatively compact in $SBC(\mathbb{R}, L^2(P, H)), PSBC(\mathbb{R} \times V, L^2(P, H))$, respectively.

Definition 3.7. An \mathcal{F}_t -progressively measurable process $\{x(t)\}_{t \in \mathbb{R}}$ is called a mild solution on \mathbb{R} of equation (1.1) if it satisfies the corresponding stochastic integral equation

$$\begin{aligned}
(3.8) \quad x(t) &= T(t-a)x(a) + \int_a^t T(t-s)g(s, x(s))ds \\
&+ \int_a^t T(t-\sigma) \int_a^\sigma B_1(\sigma-s)f(s, x(s))dsd\sigma \\
&+ \int_a^t T(t-\sigma) \int_a^\sigma B_2(\sigma-s)h(s, x(s))dW(s)d\sigma \\
&+ \int_a^t T(t-\sigma) \int_a^\sigma B_2(\sigma-s) \int_{|y|_V < 1} F(s, x(s-), y)\tilde{N}(ds, dy)d\sigma \\
&+ \int_a^t T(t-\sigma) \int_a^\sigma B_2(\sigma-s) \int_{|y|_V \geq 1} G(s, x(s-), y)N(ds, dy)d\sigma
\end{aligned}$$

for all $t \geq a$.

Remark 3.1. If we let $a \rightarrow -\infty$ in the stochastic integral equation (3.8), by the exponential dissipation condition of $(T(t))_{t \geq 0}$, then we obtain the stochastic process $x : \mathbb{R} \rightarrow L^2(\Omega, \mathbb{H})$ is a mild solution of the equation (3.8) if and only if x satisfies the stochastic integral equation

$$\begin{aligned}
(3.9) \quad x(t) &= \int_{-\infty}^t T(t-s)g(s, x(s))ds + \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^\sigma B_1(\sigma-s)f(s, x(s))dsd\sigma \\
&+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^\sigma B_2(\sigma-s)h(s, x(s))dW(s)d\sigma \\
&+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^\sigma B_2(\sigma-s) \int_{|y|_V < 1} F(s, x(s-), y)\tilde{N}(ds, dy)d\sigma \\
&+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^\sigma B_2(\sigma-s) \int_{|y|_V \geq 1} G(s, x(s-), y)N(ds, dy)d\sigma.
\end{aligned}$$

4. SQUARE-MEAN ALMOST AUTOMORPHIC SOLUTIONS

This section is devoted to the existence and the uniqueness of the square-mean almost automorphic mild solution in distribution on \mathbb{R} of Eq. (1.1).

Define the following integral operator,

$$\begin{aligned}
(\Lambda x)(t) &= \int_{-\infty}^t T(t-s)g(s, x(s))ds + \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^\sigma B_1(\sigma-s)f(s, x(s))dsd\sigma \\
&+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^\sigma B_2(\sigma-s)h(s, x(s))dW(s)d\sigma \\
&+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^\sigma B_2(\sigma-s) \int_{|y|_V < 1} F(s, x(s-), y)\tilde{N}(ds, dy)d\sigma \\
&+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^\sigma B_2(\sigma-s) \int_{|y|_V \geq 1} G(s, x(s-), y)N(ds, dy)d\sigma.
\end{aligned}$$

We have

Lemma 4.1. *If the semigroup $T(t)$ verifies (3.1) in assumption (H.1) and if the functions f, g and h are uniformly continuous on any bounded subset K of $L^2(\Omega, H)$ for each $t \in \mathbb{R}$. F, G are uniformly continuous on any bounded subset K of $L^2(\Omega, H)$ for each $t \in \mathbb{R}$ and $x \in V$. For each bounded subset $K \subset L^2(\Omega, H)$, $g(\mathbb{R}, K)$, $f(\mathbb{R}, K)$, $h(\mathbb{R}, K)$ are bounded and $F(\mathbb{R}, K, V)$ and $G(\mathbb{R}, K, V)$ are Poisson stochastically bounded., then the mapping $\Lambda : SBC(\mathbb{R}, L^2(P, H)) \rightarrow SBC(\mathbb{R}, L^2(P, H))$ is well-defined and continuous.*

Proof. It is easy to see that S is well-defined. To complete the proof it remains to show that Λ is continuous. Consider an arbitrary sequence of functions $u_n \in SBC(\mathbb{R}, L^2(P, H))$ that converges uniformly to some $u \in SBC(\mathbb{R}, L^2(P, H))$, that is, $\|u_n - u\|_\infty \rightarrow 0$ as $n \rightarrow \infty$. There exists a bounded subset K of $L^2(\Omega, H)$ such that $u_n(t), u(t) \in K$ for each $t \in \mathbb{R}$ and $n = 1, 2, \dots$. By assumptions, given $\epsilon > 0$, there exist $\delta > 0$ and $N > 0$ such that $\mathbb{E}\|u_n(t) - u(t)\|^2 < \delta$ imply that

$$\mathbb{E} \| g(t, u_n(t)) - g(t, u(t)) \|^2 < \frac{\omega^2 \epsilon}{25K^2}$$

$$\mathbb{E} \| f(t, u_n(t)) - f(t, u(t)) \|^2 < \frac{\omega^2 \epsilon}{25K^2(\|B_1\|_{L^1(0, \infty)}^2 + 1)}$$

$$\mathbb{E} \| h(t, u_n(t)) - h(t, u(t)) Q^{\frac{1}{2}} \|^2 < \frac{\omega^2 \epsilon}{25K^2(\|B_2\|_{L^2(0, \infty)}^2 + 1)}$$

$$\int_{|x|_V < 1} \mathbb{E} \| F(t, u_n(t), x) - F(t, u(t), x) \|^2 \nu(dx) < \frac{\omega^2 \epsilon}{25K^2(\|B_2\|_{L^2(0, \infty)}^2 + 1)}$$

$$\int_{|x|_V < 1} \mathbb{E} \| G(t, u_n(t), x) - G(t, u(t), x) \|^2 \nu(dx) < \frac{\omega^2 \epsilon}{50K^2(\|B_2\|_{L^2(0, \infty)}^2 + b\|B_2\|_{L^1(0, \infty)}^2 + 1)}$$

Hence

$$\begin{aligned} \mathbb{E} \| (\Lambda u_n)(t) - (\Lambda u)(t) \|^2 &= \mathbb{E} \left\| \int_{-\infty}^t T(t-s) \left(g(s, u_n(s)) - g(s, u(s)) \right) ds \right. \\ &\quad + \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s) \left(f(s, u_n(s)) - f(s, u(s)) \right) ds d\sigma \\ &\quad + \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \left(h(s, u_n(s)) - h(s, u(s)) \right) dW(s) d\sigma \\ &\quad + \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} \left(F(s, u_n(s-), y) - F(s, u(s-), y) \right) \tilde{N}(ds, dy) d\sigma \\ &\quad \left. + \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(G(s, u_n(s-), y) - G(s, u(s-), y) \right) N(ds, dy) d\sigma \right\|^2 \end{aligned}$$

$$\begin{aligned}
&\leq 5\mathbb{E}\left\|\int_{-\infty}^t T(t-s)\left(g(s, u_n(s)) - g(s, u(s))\right)ds\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{-\infty}^t T(t-\sigma)\int_{-\infty}^{\sigma} B_1(\sigma-s)\left(f(s, u_n(s)) - f(s, u(s))\right)dsd\sigma\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{-\infty}^t T(t-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\left(h(s, u_n(s)) - h(s, u(s))\right)dW(s)d\sigma\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{-\infty}^t T(t-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|<1}\left(F(s, u_n(s-), y) - F(s, u(s-), y)\right)\tilde{N}(ds, dy)d\sigma\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{-\infty}^t T(t-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|\geq 1}\left(G(s, u_n(s-), y) - G(s, u(s-), y)\right)N(ds, dy)d\sigma\right\|^2 \\
&\leq 5(I_1 + I_2 + I_3 + I_4 + I_5).
\end{aligned}$$

Using Cauchy-Schwartz's inequality, we get

$$I_1 < \frac{\epsilon}{25} \quad I_2 < \frac{\epsilon}{25}$$

For I_3 , using Cauchy-Schwartz's inequality and Ito's isometry property, we obtain

$$I_3 \leq \frac{K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \mathbb{E}\left\|B_2(\sigma-s)\left(h(s, u_n(s)) - h(s, u(s))\right)Q^{\frac{1}{2}}\right\|^2 dsd\sigma < \frac{\epsilon}{25}$$

As to I_4 and I_5 , by Cauchy-Schwartz's inequality and the properties of the integral for the Poisson random measure, we have

$$\begin{aligned}
I_4 &= \mathbb{E}\left\|\int_{-\infty}^t T(t-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|<1}\left(F(s, u_n(s-), y) - F(s, u(s-), y)\right)\tilde{N}(ds, dy)d\sigma\right\|^2 \\
&\leq \frac{K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \mathbb{E}\left\|\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|<1}\left(F(s, u_n(s-), y) - F(s, u(s-), y)\right)\tilde{N}(ds, dy)\right\|^2 d\sigma \\
&\leq \frac{K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 \int_{|y|<1} \mathbb{E}\left\|F(s, u_n(s-), y) - F(s, u(s-), y)\right\|^2 \nu(dy)d\sigma \\
&< \frac{\epsilon}{25}.
\end{aligned}$$

And

$$\begin{aligned}
I_5 &= \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(F(s, u_n(s-), y) - F(s, u(s-), y) \right) N(ds, dy) d\sigma \right\|^2 \\
&\leq 2\mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(F(s, u_n(s-), y) - F(s, u(s-), y) \right) \tilde{N}(ds, dy) d\sigma \right\|^2 \\
&+ 2\mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(F(s, u_n(s-), y) - F(s, u(s-), y) \right) \nu(dy) d\sigma \right\|^2 \\
&\leq \frac{2K^2}{\omega^2} \|B_2\|_{L^2(0, \infty)}^2 \left(\frac{\omega^2 \epsilon}{50K^2(\|B_2\|_{L^2(0, \infty)}^2 + b\|B_2\|_{L^1(0, \infty)}^2 + 1)} \right) \\
&+ 2\frac{K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \mathbb{E} \left\| \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(F(s, u_n(s-), y) - F(s, u(s-), y) \right) \nu(dy) ds \right\|^2 d\sigma \\
&\leq \frac{2K^2}{\omega^2} \|B_2\|_{L^2(0, \infty)}^2 \left(\frac{\omega^2 \epsilon}{50K^2(\|B_2\|_{L^2(0, \infty)}^2 + b\|B_2\|_{L^1(0, \infty)}^2 + 1)} \right) + 2\frac{K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \left(\int_{-\infty}^{\sigma} \|B_2(\sigma-s)\| ds \int_{|y|_V \geq 1} \nu(dy) \right. \\
&\quad \cdot \left. \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\| \int_{|y|_V \geq 1} \mathbb{E} \left\| F(s, u_n(s-), y) - F(s, u(s-), y) \right\|^2 \nu(dy) ds \right) d\sigma \\
&\leq \frac{2K^2}{\omega^2} \|B_2\|_{L^2(0, \infty)}^2 \left(\frac{\omega^2 \epsilon}{50K^2(\|B_2\|_{L^2(0, \infty)}^2 + b\|B_2\|_{L^1(0, \infty)}^2 + 1)} \right) \\
&+ 2b\frac{K^2}{\omega} \|B_2\|_{L^1(0, \infty)}^2 \left(\frac{\omega^2 \epsilon}{50K^2(\|B_2\|_{L^2(0, \infty)}^2 + b\|B_2\|_{L^1(0, \infty)}^2 + 1)} \right) \\
&< \frac{\epsilon}{25}.
\end{aligned}$$

Thus, by combining $I_1 - I_5$, it follows that for each $t \in \mathbb{R}$ and $n > N$

$$\mathbb{E} \|(\Lambda u_n)(t) - (\Lambda u)(t)\|^2 < \epsilon.$$

This implies that Λ is continuous. The proof is complete. \square

Theorem 4.1. Assume that assumptions (H.1) – (H.4) hold and

- 1- $g, f \in SAA(\mathbb{R} \times L^2(P, H), L^2(P, H))$,
- 2- $h \in SAA(\mathbb{R} \times L^2(P, H), L(V, L^2(P, H)))$,
- 3- $F \in PSAA(\mathbb{R} \times L^2(P, H) \times V, L^2(P, H))$
- 4- $G \in PSAA(\mathbb{R} \times L^2(P, H) \times V, L^2(P, H))$.

then Eq. (1.1) has at least one almost automorphic in distribution mild solution on \mathbb{R} provided that

$$\begin{aligned}
L_g &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t e^{-\omega(t-s)} m_g(s) ds < \infty, & L_f &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_1(t-s)\| m_f(s) ds < \infty, \\
L_h &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_2(t-s)\|^2 m_h(s) ds < \infty, & L_F &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_2(t-s)\|^2 m_F(s) ds < \infty, \\
L_G &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_2(t-s)\|^2 m_G(s) ds < \infty
\end{aligned}$$

and

$$(4.1) \quad \vartheta := 10\frac{K^2}{\omega^2} \left[\omega L_g + L_f \|B_1\|_{L^1(0, \infty)} + L_h + L_F + 2 \left(1 + b \|B_2\|_{L^1(0, \infty)} \right) L_G \right] < 1.$$

Proof. Let $B = \{u \in SBC(\mathbb{R}, L^2(P, H)) : \|u\|_\infty^2 \leq r\}$. By Lemma 4.1 and (3.2), it follows that B is a convex and closed set satisfying $\Lambda B \subset B$. To complete the proof, we have to prove the following statements:

- a) That $V = \{\Lambda u(t) : u \in B\}$ is a relatively compact subset of $L^2(P, H)$ for each $t \in \mathbb{R}$;
- b) That $U = \{\Lambda u : u \in B\} \subset SBC(\mathbb{R}, L^2(P, H))$ is equi-continuous.
- c) The mild solution is almost automorphic in distribution.

To prove a), fix $t \in \mathbb{R}$ and consider an arbitrary $\varepsilon > 0$. Then

$$\begin{aligned}
(\Lambda_\varepsilon x)(t) &= \int_{-\infty}^{t-\varepsilon} T(t-s)g(s, x(s))ds + \int_{-\infty}^{t-\varepsilon} T(t-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s)f(s, x(s))dsd\sigma \\
&+ \int_{-\infty}^{t-\varepsilon} T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s)h(s, x(s))dW(s)d\sigma \\
&+ \int_{-\infty}^{t-\varepsilon} T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} F(s, x(s-), y)\tilde{N}(ds, dy)d\sigma \\
&+ \int_{-\infty}^{t-\varepsilon} T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} G(s, x(s-), y)N(ds, dy)d\sigma \\
&= T(\varepsilon) \int_{-\infty}^{t-\varepsilon} T(t-\varepsilon-s)g(s, x(s))ds + \int_{-\infty}^{t-\varepsilon} T(t-\varepsilon-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s)f(s, x(s))dsd\sigma \\
&+ \int_{-\infty}^{t-\varepsilon} T(t-\varepsilon-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s)h(s, x(s))dW(s)d\sigma \\
&+ \int_{-\infty}^{t-\varepsilon} T(t-\varepsilon-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} F(s, x(s-), y)\tilde{N}(ds, dy)d\sigma \\
&+ \int_{-\infty}^{t-\varepsilon} T(t-\varepsilon-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} G(s, x(s-), y)N(ds, dy)d\sigma \\
&= T(\varepsilon)(\Lambda x)(t-\varepsilon)
\end{aligned}$$

and hence $V_\varepsilon := \{(\Lambda_\varepsilon)x(t) : x \in B\}$ is relatively compact in $L^2(P, H)$ as the semigroup family $T(\varepsilon)$ is compact by assumption.

Now

$$\begin{aligned}
\mathbb{E}\|(\Lambda u)(t) - (\Lambda_\varepsilon u)(t)\|^2 &= \mathbb{E}\left\| \int_{t-\varepsilon}^t T(t-s)g(s, u(s))ds + \int_{t-\varepsilon}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s)f(s, u(s))dsd\sigma \right. \\
&+ \int_{t-\varepsilon}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \left(h(s, u_n(s)) - h(s, u(s))dW(s) \right) d\sigma \\
&+ \int_{t-\varepsilon}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} F(s, u(s-), y)\tilde{N}(ds, dy)d\sigma \\
&\left. + \int_{t-\varepsilon}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} G(s, u(s-), y)N(ds, dy)d\sigma \right\|^2
\end{aligned}$$

$$\begin{aligned}
&\leq 5\mathbb{E}\left\|\int_{t-\epsilon}^t T(t-s)g(s, u(s))ds\right\|^2 + 5\mathbb{E}\left\|\int_{t-\epsilon}^t T(t-\sigma)\int_{-\infty}^{\sigma} B_1(\sigma-s)f(s, u(s))dsd\sigma\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{t-\epsilon}^t T(t-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)h(s, u(s))dW(s)d\sigma\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{t-\epsilon}^t T(t-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|_V < 1} F(s, u(s-), y)\tilde{N}(ds, dy)d\sigma\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{t-\epsilon}^t T(t-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|_V \geq 1} G(s, u(s-), y)N(ds, dy)d\sigma\right\|^2 \\
&\leq 5\left[\Delta_r K^2\left(\int_{t-\epsilon}^t e^{-\omega(t-\sigma)}d\sigma\right)^2 + \Delta_r K^2\|B_1\|_{L^1(0, \infty)}^2\left(\int_{t-\epsilon}^t e^{-\omega(t-\sigma)}d\sigma\right)^2\right. \\
&+ \Delta_r K^2\|B_2\|_{L^2(0, \infty)}^2\left(\int_{t-\epsilon}^t e^{-\omega(t-\sigma)}d\sigma\right)^2 + \Delta_r K^2\|B_2\|_{L^2(0, \infty)}^2\left(\int_{t-\epsilon}^t e^{-\omega(t-\sigma)}d\sigma\right)^2 \\
&\left. + \left(2\Delta_r K^2\|B_2\|_{L^2(0, \infty)}^2 + 2b\Delta_r K^2\|B_2\|_{L^1(0, \infty)}^2\right)\left(\int_{t-\epsilon}^t e^{-\omega(t-\sigma)}d\sigma\right)^2\right] \\
&\leq 5\Delta_r K^2\left[1 + \|B_1\|_{L^1(0, \infty)}^2 + 4\|B_2\|_{L^2(0, \infty)}^2 + 2b\|B_2\|_{L^1(0, \infty)}^2\right]\epsilon^2
\end{aligned}$$

from which it follows that $V = \{\Lambda u(t) : u \in B\}$ is a relatively compact subset of $L^2(P, H)$ for each $t \in \mathbb{R}$.

We now show that $b)$ holds. Let $u \in B$ and $t_1, t_2 \in \mathbb{R}$ such that $t_1 < t_2$. Similar computation as that in Lemma 4.1, we have

$$\begin{aligned}
\mathbb{E}\|(\Lambda u)(t_1) - (\Lambda u)(t_2)\|^2 &= \mathbb{E}\left\|\int_{-\infty}^{t_2} T(t_2-s)g(s, x(s))ds + \int_{-\infty}^{t_2} T(t_2-\sigma)\int_{-\infty}^{\sigma} B_1(\sigma-s)f(s, x(s))dsd\sigma\right. \\
&+ \int_{-\infty}^{t_2} T(t_2-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)h(s, x(s))dW(s)d\sigma \\
&+ \int_{-\infty}^{t_2} T(t_2-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|_V < 1} F(s, x(s-), y)\tilde{N}(ds, dy)d\sigma \\
&+ \int_{-\infty}^{t_2} T(t_2-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|_V \geq 1} G(s, x(s-), y)N(ds, dy)d\sigma \\
&- \left(\int_{-\infty}^{t_1} T(t_1-s)g(s, x(s))ds + \int_{-\infty}^{t_1} T(t_1-\sigma)\int_{-\infty}^{\sigma} B_1(\sigma-s)f(s, x(s))dsd\sigma\right. \\
&+ \int_{-\infty}^{t_1} T(t_1-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)h(s, x(s))dW(s)d\sigma \\
&+ \int_{-\infty}^{t_1} T(t_1-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|_V < 1} F(s, x(s-), y)\tilde{N}(ds, dy)d\sigma \\
&\left. + \int_{-\infty}^{t_1} T(t_1-\sigma)\int_{-\infty}^{\sigma} B_2(\sigma-s)\int_{|y|_V \geq 1} G(s, x(s-), y)N(ds, dy)d\sigma\right)\|^2
\end{aligned}$$

$$\begin{aligned}
&= \mathbb{E} \left[\left\| \left(T(t_2 - t_1) - I \right) \left[\int_{-\infty}^{t_1} T(t_1 - s) g(s, x(s)) ds + \int_{-\infty}^{t_1} T(t_1 - \sigma) \int_{-\infty}^{\sigma} B_1(\sigma - s) f(s, x(s)) ds d\sigma \right. \right. \right. \\
&+ \int_{-\infty}^{t_1} T(t_1 - \sigma) \int_{-\infty}^{\sigma} B_2(\sigma - s) h(s, x(s)) dW(s) d\sigma \\
&+ \int_{-\infty}^{t_1} T(t_1 - \sigma) \int_{-\infty}^{\sigma} B_2(\sigma - s) \int_{|y|_V < 1} F(s, x(s-), y) \tilde{N}(ds, dy) d\sigma \\
&+ \left. \left. \left. \int_{-\infty}^{t_1} T(t_1 - \sigma) \int_{-\infty}^{\sigma} B_2(\sigma - s) \int_{|y|_V \geq 1} G(s, x(s-), y) N(ds, dy) d\sigma \right] \right\|^2 \right. \\
&+ \int_{t_1}^{t_2} T(t_2 - s) g(s, x(s)) ds + \int_{t_1}^{t_2} T(t_2 - \sigma) \int_{-\infty}^{\sigma} B_1(\sigma - s) f(s, x(s)) ds d\sigma \\
&+ \int_{t_1}^{t_2} T(t_2 - \sigma) \int_{-\infty}^{\sigma} B_2(\sigma - s) h(s, x(s)) dW(s) d\sigma \\
&+ \int_{t_1}^{t_2} T(t_2 - \sigma) \int_{-\infty}^{\sigma} B_2(\sigma - s) \int_{|y|_V < 1} F(s, x(s-), y) \tilde{N}(ds, dy) d\sigma \\
&+ \left. \left. \left. \int_{t_1}^{t_2} T(t_2 - \sigma) \int_{-\infty}^{\sigma} B_2(\sigma - s) \int_{|y|_V \geq 1} G(s, x(s-), y) N(ds, dy) d\sigma \right\|^2 \right\} \right. \\
&\leq 6 \sup_{y \in V} \mathbb{E} \left[\left\| (T(t_2 - t_1) - I)y \right\|^2 \right] + 6 \mathbb{E} \left[\left\| \int_{t_1}^{t_2} T(t_2 - s) g(s, x(s)) ds \right\|^2 \right] \\
&+ 6 \mathbb{E} \left[\left\| \int_{t_1}^{t_2} T(t_2 - \sigma) \int_{-\infty}^{\sigma} B_1(\sigma - s) f(s, x(s)) ds d\sigma \right\|^2 \right] \\
&+ 6 \mathbb{E} \left[\left\| \int_{t_1}^{t_2} T(t_2 - \sigma) \int_{-\infty}^{\sigma} B_2(\sigma - s) h(s, x(s)) dW(s) d\sigma \right\|^2 \right] \\
&+ 6 \mathbb{E} \left[\left\| \int_{t_1}^{t_2} T(t_2 - \sigma) \int_{-\infty}^{\sigma} B_2(\sigma - s) \int_{|y|_V < 1} F(s, x(s-), y) \tilde{N}(ds, dy) d\sigma \right\|^2 \right] \\
&+ 6 \mathbb{E} \left[\left\| \int_{t_1}^{t_2} T(t_2 - \sigma) \int_{-\infty}^{\sigma} B_2(\sigma - s) \int_{|y|_V \geq 1} G(s, x(s-), y) N(ds, dy) d\sigma \right\|^2 \right] \\
&\leq 6 \sup_{y \in V} \mathbb{E} \left[\left\| (T(t_2 - t_1) - I)y \right\|^2 \right] + 6 \left[\Delta_r K^2 \left(\int_{t_1}^{t_2} e^{-\omega(t_2 - \sigma)} d\sigma \right)^2 \right. \\
&+ \Delta_r K^2 \|B_1\|_{L^1(0, \infty)}^2 \left(\int_{t_1}^{t_2} e^{-\omega(t_2 - \sigma)} d\sigma \right)^2 \\
&+ \Delta_r K^2 \|B_2\|_{L^2(0, \infty)}^2 \left(\int_{t_1}^{t_2} e^{-\omega(t_2 - \sigma)} d\sigma \right)^2 + \Delta_r K^2 \|B_2\|_{L^2(0, \infty)}^2 \left(\int_{t_1}^{t_2} e^{-\omega(t_2 - \sigma)} d\sigma \right)^2 \\
&+ \left. \left(2\Delta_r K^2 \|B_2\|_{L^2(0, \infty)}^2 + 2b\Delta_r K^2 \|B_2\|_{L^1(0, \infty)}^2 \right) \left(\int_{t_1}^{t_2} e^{-\omega(t_2 - \sigma)} d\sigma \right)^2 \right] \\
&\leq 6 \sup_{y \in V} \mathbb{E} \left[\left\| (T(t_2 - t_1) - I)y \right\|^2 \right] \\
&+ 6\Delta_r K^2 \left[1 + \|B_1\|_{L^1(0, \infty)}^2 + 4\|B_2\|_{L^2(0, \infty)}^2 + 2b\|B_2\|_{L^1(0, \infty)}^2 \right] \left(\int_{t_1}^{t_2} e^{-\omega(t_2 - \sigma)} d\sigma \right)^2.
\end{aligned}$$

The right-hand side tends to 0 independently to $u \in B$ as $t_2 \rightarrow t_1$ which implies that U is right equi-continuous at t . Similarly, we can show that U is left equi-continuous at t .

Denote the closed convex hull of ΛB by $\overline{c\partial} \Lambda B$. Since $\overline{c\partial} \Lambda B \subset B$ and B is a closed convex, it follows that

$$\Lambda(\overline{c\partial} \Lambda B) \subset \Lambda B \subset \overline{c\partial} \Lambda B.$$

Further, it is not hard to see that $\overline{c\partial} \Lambda B$ is relatively compact in $L^2(P, H)$. Using Arzelà-Ascoli theorem, we deduce that the restriction of $\overline{c\partial} \Lambda B$ to any compact subset I of \mathbb{R} is relatively compact in $C(I, L^2(P, H))$. Thus condition (H.4) implies that $\Lambda : \overline{c\partial} \Lambda B \mapsto \overline{c\partial} \Lambda B$ is a compact operator. In summary, $S : \overline{c\partial} \Lambda B \mapsto \overline{c\partial} \Lambda B$ is continuous and compact. Using the Schauder fixed point it follows that Λ has a fixed-point.

To end the proof, we have to check this the fixed-point is almost automorphic in distribution. Since g, f, h are almost automorphic and F, G are Poisson almost automorphic, then for every sequence of real numbers $(t_n)_n$ we can extract a subsequence (t_n) such that, for some stochastic processes $\tilde{g}, \tilde{f}, \tilde{h}, \tilde{F}, \tilde{G}$

$$(4.2) \quad \lim_{n \rightarrow +\infty} \mathbb{E} \|g(s+t_n, X) - \tilde{g}(s)\|^2 = 0, \quad \lim_{n \rightarrow +\infty} \mathbb{E} \|\tilde{g}(s-t_n, X) - g(s, X)\|^2 = 0;$$

$$(4.3) \quad \lim_{n \rightarrow +\infty} \mathbb{E} \|f(s+t_n, X) - \tilde{f}(s, X)\|^2 = 0, \quad \lim_{n \rightarrow +\infty} \mathbb{E} \|\tilde{f}(s-t_n, X) - f(s, X)\|^2 = 0;$$

$$(4.4)$$

$$\lim_{n \rightarrow +\infty} \mathbb{E} \left\| \left(h(s+t_n, X) - \tilde{h}(s, X) \right) \mathcal{Q}^{\frac{1}{2}} \right\|_{L(V, L^2(P, H))}^2 = 0, \quad \lim_{n \rightarrow +\infty} \mathbb{E} \left\| \left(\tilde{h}(s-t_n, X) - h(s, X) \right) \mathcal{Q}^{\frac{1}{2}} \right\|_{L(V, L^2(P, H))}^2 = 0;$$

$$(4.5) \quad \lim_{n \rightarrow +\infty} \int_{|y|_V < 1} \mathbb{E} \|F(s+t_n, X, y) - \tilde{F}(s, X, y)\|^2 \nu(dy) = 0,$$

$$\lim_{n \rightarrow +\infty} \int_{|y|_V < 1} \mathbb{E} \|\tilde{F}(s-t_n, X, y) - F(s, X, y)\|^2 \nu(dy) = 0$$

and

$$(4.6) \quad \lim_{n \rightarrow +\infty} \int_{|y|_V \geq 1} \mathbb{E} \|G(s+t_n, X, y) - \tilde{G}(s, X, y)\|^2 \nu(dy) = 0,$$

$$\lim_{n \rightarrow +\infty} \int_{|y|_V \geq 1} \mathbb{E} \|\tilde{G}(s-t_n, X, y) - G(s, X, y)\|^2 \nu(dy) = 0$$

hold for each $s \in \mathbb{R}$ and $X \in L^2(P, H)$.

For $t \in \mathbb{R}$, we define

$$\begin{aligned} \tilde{X}(t) &= \int_{-\infty}^t T(t-s) \tilde{g}(s, \tilde{X}(s)) ds + \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s) \tilde{f}(s, \tilde{X}(s)) ds d\sigma \\ &+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \tilde{h}(s, \tilde{X}(s)) dW(s) d\sigma \\ &+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} \tilde{F}(s, \tilde{X}(s-), y) \tilde{N}(ds, dy) d\sigma \\ &+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \tilde{G}(s, \tilde{X}(s-), y) N(ds, dy) d\sigma. \end{aligned}$$

Let $W_n(s) = W(s+t_n) - W(t_n)$, $N_n(s, y) = N(s+t_n, y) - N(t_n, y)$ and $\tilde{N}_n(s, y) = N(s+t_n, y) - N(t_n, y)$ for each $s \in \mathbb{R}$. Then W_n is also a Q-Wiener process having the same distribution as W and N_n have the same distribution as N with compensated Poisson random measure \tilde{N}_n .

Consider the process define as follows

$$\begin{aligned}
X_n(t) &= \int_{-\infty}^t T(t-s)g(s+t_n, X_n(s))ds + \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s)f(s+t_n, X_n(s))dsd\sigma \\
&+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s)h(s+t_n, X_n(s))dW(s)d\sigma \\
&+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} F(s+t_n, X_n(s-), y)\tilde{N}(ds, dy)d\sigma \\
&+ \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} G(s+t_n, X_n(s-), y)N(ds, dy)d\sigma.
\end{aligned}$$

Note that $X(t+t_n)$ and X_n have the same law and since the convergence in L^2 implies convergence in distribution, then we have

$$\begin{aligned}
&\mathbb{E}\|X_n(t) - \tilde{X}(t)\|^2 \\
&\leq 5\mathbb{E}\left\|\int_{-\infty}^t T(t-s)\left(g(s+t_n, X_n(s)) - \tilde{g}(s, \tilde{X}(s))\right)ds\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s)\left(f(s+t_n, X_n(s)) - \tilde{f}(s, \tilde{X}(s))\right)dsd\sigma\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s)\left(h(s+t_n, X_n(s)) - \tilde{h}(s, \tilde{X}(s))\right)dW(s)d\sigma\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} \left(F(s+t_n, X_n(s-), y) - \tilde{F}(s, \tilde{X}(s-), y)\right)\tilde{N}(ds, dy)d\sigma\right\|^2 \\
&+ 5\mathbb{E}\left\|\int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(G(s+t_n, X_n(s-), y) - \tilde{G}(s, \tilde{X}(s-), y)\right)N(ds, dy)d\sigma\right\|^2 \\
&\leq 5(J_1 + J_2 + J_3 + J_4 + J_5).
\end{aligned}$$

For J_1 , we have

$$\begin{aligned}
J_1 &= \mathbb{E}\left\|\int_{-\infty}^t T(t-s)\left(g(s+t_n, X_n(s)) - \tilde{g}(s, \tilde{X}(s))\right)ds\right\|^2 \\
&\leq 2\mathbb{E}\left\|\int_{-\infty}^t T(t-s)\left(g(s+t_n, X_n(s)) - g(s+t_n, \tilde{X}(s))\right)ds\right\|^2 \\
&+ 2\mathbb{E}\left\|\int_{-\infty}^t T(t-s)\left(g(s+t_n, \tilde{X}(s)) - \tilde{g}(s, \tilde{X}(s))\right)ds\right\|^2 \\
&\leq \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-s)} m_g(s+t_n) \mathbb{E}\|X_n(s) - \tilde{X}(s)\|^2 ds \\
&+ \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-s)} \mathbb{E}\|g(s+t_n, \tilde{X}(s)) - \tilde{g}(s, \tilde{X}(s))\|^2 ds \\
&\leq \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-s)} m_g(s+t_n) \mathbb{E}\|X_n(s) - \tilde{X}(s)\|^2 ds + \zeta_1^n,
\end{aligned}$$

where $\zeta_1^n := \frac{2K^2}{\omega^2} \sup_{s \in \mathbb{R}} \mathbb{E}\|g(s+t_n, \tilde{X}(s)) - \tilde{g}(s, \tilde{X}(s))\|^2 ds$. Since $\tilde{X}(\cdot)$ is bounded in $L^2(P, H)$, it follows by (4.2), that $\zeta_1^n \rightarrow 0$ as $n \rightarrow \infty$.

For J_2 , we have

$$\begin{aligned}
J_2 &= \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s) \left(f(s+t_n, X_n(s)) - \tilde{f}(s, \tilde{X}(s)) \right) ds d\sigma \right\|^2 \\
&\leq 2\mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s) \left(f(s+t_n, X_n(s)) - f(s+t_n, \tilde{X}(s)) \right) ds d\sigma \right\|^2 \\
&\quad + 2\mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_1(\sigma-s) \left(f(s+t_n, \tilde{X}(s)) - \tilde{f}(s, \tilde{X}(s)) \right) ds d\sigma \right\|^2 \\
&\leq \frac{2K^2}{\omega} \|B_1\|_{L^1(0,\infty)} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_1(\sigma-s)\| m_f(s+t_n) \mathbb{E} \|X_n(s) - \tilde{X}(s)\|^2 ds d\sigma \\
&\quad + \frac{2K^2}{\omega} \|B_1\|_{L^1(0,\infty)} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_1(\sigma-s)\| \mathbb{E} \|f(s+t_n, \tilde{X}(s)) - \tilde{f}(s, \tilde{X}(s))\|^2 ds d\sigma \\
&\leq \frac{2K^2}{\omega} \|B_1\|_{L^1(0,\infty)} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_1(\sigma-s)\| m_f(s+t_n) \mathbb{E} \|X_n(s) - \tilde{X}(s)\|^2 ds d\sigma + \zeta_2^n,
\end{aligned}$$

where $\zeta_2^n := \frac{2K^2}{\omega^2} \|B_1\|_{L^1(0,\infty)}^2 \sup_{s \in \mathbb{R}} \mathbb{E} \|f(s+t_n, \tilde{X}(s)) - \tilde{f}(s, \tilde{X}(s))\|^2$. For the same reason as for ζ_1^n , $\zeta_2^n \rightarrow 0$ as $n \rightarrow \infty$.

For J_3 , using Cauchy-Schwartz's inequality and the Ito's isometry, we have

$$\begin{aligned}
J_3 &= \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \left(h(s+t_n, X_n(s)) - \tilde{h}(s, \tilde{X}(s)) \right) dW(s) d\sigma \right\|^2 \\
&\leq 2\mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \left(h(s+t_n, X_n(s)) - h(s+t_n, \tilde{X}(s)) \right) dW(s) d\sigma \right\|^2 \\
&\quad + 2\mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \left(h(s+t_n, \tilde{X}(s)) - \tilde{h}(s, \tilde{X}(s)) \right) dW(s) d\sigma \right\|^2 \\
&\leq \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 m_h(s+t_n) \mathbb{E} \|X_n(s) - \tilde{X}(s)\|^2 ds d\sigma \\
&\quad + \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 \mathbb{E} \left\| \left(h(s+t_n, \tilde{X}(s)) - \tilde{h}(s, \tilde{X}(s)) \right) \mathcal{Q}^{\frac{1}{2}} \right\|_{L(V, L^2(P, H))}^2 ds d\sigma \\
&\leq \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 m_h(s+t_n) \mathbb{E} \|X_n(s) - \tilde{X}(s)\|^2 ds d\sigma + \zeta_3^n,
\end{aligned}$$

where $\zeta_3^n := \frac{2K^2}{\omega^2} \|B_2\|_{L^2(0,\infty)}^2 \sup_{s \in \mathbb{R}} \mathbb{E} \left\| \left(h(s+t_n, \tilde{X}(s)) - \tilde{h}(s, \tilde{X}(s)) \right) \mathcal{Q}^{\frac{1}{2}} \right\|_{L(V, L^2(P, H))}^2$. By (4.4), it follows that $\zeta_3^n \rightarrow 0$ as $n \rightarrow \infty$, like ζ_1^n .

For J_4 , using Cauchy-Schwartz's inequality and the properties of the integral for the Poisson random measure, we have

$$\begin{aligned}
J_4 &= \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} \left(F(s+t_n, X_n(s-), y) - \tilde{F}(s, \tilde{X}(s-), y) \right) \tilde{N}(ds, dy) d\sigma \right\|^2 \\
&\leq 2\mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} \left(F(s+t_n, X_n(s-), y) - F(s+t_n, \tilde{X}(s-), y) \right) \tilde{N}(ds, dy) d\sigma \right\|^2 \\
&\quad + 2\mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V < 1} \left(F(s+t_n, \tilde{X}(s-), y) - \tilde{F}(s, \tilde{X}(s-), y) \right) \tilde{N}(ds, dy) d\sigma \right\|^2
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 m_F(s+t_n) \mathbb{E} \left\| X_n(s) - \tilde{X}(s) \right\|^2 ds d\sigma \\
&+ \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 \int_{|y|_V < 1} \mathbb{E} \left\| F(s+t_n, \tilde{X}(s-), y) - \tilde{F}(s, \tilde{X}(s-), y) \right\|^2 \nu(dy) ds d\sigma \\
&\leq \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 m_F(s+t_n) \mathbb{E} \left\| X_n(s) - \tilde{X}(s) \right\|^2 ds d\sigma + \zeta_4^n,
\end{aligned}$$

where $\zeta_4^n := \frac{2K^2}{\omega^2} \|B_2\|_{L^2(0,\infty)}^2 \sup_{s \in \mathbb{R}} \left(\int_{|y|_V < 1} \mathbb{E} \left\| F(s+t_n, \tilde{X}(s-), y) - \tilde{F}(s, \tilde{X}(s-), y) \right\|^2 \nu(dy) \right)$.

Using (4.5), it follows that $\zeta_4^n \rightarrow 0$ as $n \rightarrow \infty$, like ζ_1^n .

For J_5 , using Cauchy-Schwartz's inequality and the properties of the integral for the Poisson random measure, we have

$$\begin{aligned}
J_5 &= \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(G(s+t_n, X_n(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right) N(ds, dy) d\sigma \right\|^2 \\
&\leq 2 \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(G(s+t_n, X_n(s-), y) - G(s+t_n, \tilde{X}(s-), y) \right) N(ds, dy) d\sigma \right\|^2 \\
&+ 2 \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(G(s+t_n, \tilde{X}(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right) N(ds, dy) d\sigma \right\|^2 \\
&\leq 4 \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(G(s+t_n, X_n(s-), y) - G(s+t_n, \tilde{X}(s-), y) \right) \tilde{N}(ds, dy) d\sigma \right\|^2 \\
&+ 4 \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(G(s+t_n, X_n(s-), y) - G(s+t_n, \tilde{X}(s-), y) \right) \nu(dy) ds d\sigma \right\|^2 \\
&+ 4 \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(G(s+t_n, \tilde{X}(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right) \tilde{N}(ds, dy) d\sigma \right\|^2 \\
&+ 4 \mathbb{E} \left\| \int_{-\infty}^t T(t-\sigma) \int_{-\infty}^{\sigma} B_2(\sigma-s) \int_{|y|_V \geq 1} \left(G(s+t_n, \tilde{X}(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right) \nu(dy) ds d\sigma \right\|^2 \\
&\leq \frac{4K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 m_G(s+t_n) \mathbb{E} \left\| X_n(s) - \tilde{X}(s) \right\|^2 ds d\sigma \\
&+ \frac{4K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \left(\int_{-\infty}^{\sigma} \|B_2(\sigma-s)\| \int_{|y|_V \geq 1} \nu(dy) ds \right. \\
&\quad \times \left. \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\| \int_{|y|_V \geq 1} \mathbb{E} \left\| G(s+t_n, X_n(s-), y) - G(s+t_n, \tilde{X}(s-), y) \right\|^2 \nu(dy) ds \right) d\sigma \\
&+ \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 \int_{|y|_V \geq 1} \mathbb{E} \left\| G(s+t_n, \tilde{X}(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right\|^2 \nu(dy) ds d\sigma \\
&+ \frac{4K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \left(\int_{-\infty}^{\sigma} \|B_2(\sigma-s)\| \int_{|y|_V \geq 1} \nu(dy) ds \right. \\
&\quad \times \left. \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\| \int_{|y|_V \geq 1} \mathbb{E} \left\| G(s+t_n, \tilde{X}(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right\|^2 \nu(dy) ds \right) d\sigma
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{4K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 m_G(s+t_n) \mathbb{E} \left\| X_n(s) - \tilde{X}(s) \right\|^2 ds d\sigma \\
&+ \frac{4bK^2}{\omega} \|B_2\|_{L^1(0,\infty)} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\| m_G(s+t_n) \mathbb{E} \left\| X_n(s) - \tilde{X}(s) \right\|^2 ds d\sigma \\
&+ \frac{2K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 \int_{|y|v \geq 1} \mathbb{E} \left\| G(s+t_n, \tilde{X}(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right\|^2 v(dy) ds d\sigma \\
&+ \frac{4bK^2}{\omega} \|B_2\|_{L^2(0,\infty)} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\| \int_{|y|v \geq 1} \mathbb{E} \left\| G(s+t_n, \tilde{X}(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right\|^2 v(dy) ds d\sigma \\
&\leq \frac{4K^2}{\omega} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\|^2 m_G(s+t_n) \mathbb{E} \left\| X_n(s) - \tilde{X}(s) \right\|^2 ds d\sigma \\
&+ \frac{4bK^2}{\omega} \|B_2\|_{L^1(0,\infty)} \int_{-\infty}^t e^{-\omega(t-\sigma)} \int_{-\infty}^{\sigma} \|B_2(\sigma-s)\| m_G(s+t_n) \mathbb{E} \left\| X_n(s) - \tilde{X}(s) \right\|^2 ds d\sigma + \zeta_5^n(t)
\end{aligned}$$

where

$$\begin{aligned}
\zeta_5^n &= \frac{2K^2}{\omega^2} \|B_2\|_{L^2(0,\infty)}^2 \sup_{s \in \mathbb{R}} \left(\int_{|y|v \geq 1} \mathbb{E} \left\| G(s+t_n, \tilde{X}(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right\|^2 v(dy) \right) \\
&+ \frac{4bK^2}{\omega^2} \|B_2\|_{L^1(0,\infty)}^2 \sup_{s \in \mathbb{R}} \left(\int_{|y|v \geq 1} \mathbb{E} \left\| G(s+t_n, \tilde{X}(s-), y) - \tilde{G}(s, \tilde{X}(s-), y) \right\|^2 v(dy) \right)
\end{aligned}$$

Using (4.6), it follows that $\zeta_5^n \rightarrow 0$ as $n \rightarrow \infty$, like ζ_1^n . By combining the estimations $J_1 - J_5$, we get for all $t \in \mathbb{R}$

$$\mathbb{E} \left\| X_n(t) - \tilde{X}(t) \right\|^2 \leq \zeta^n + \vartheta \sup_{s \in \mathbb{R}} \mathbb{E} \left\| X_n(s) - \tilde{X}(s) \right\|^2$$

where $\zeta^n = \sum_{i=1}^5 \zeta_i^n$. It follows that

$$\mathbb{E} \left\| X_n(t) - \tilde{X}(t) \right\|^2 \leq \frac{\zeta^n}{1-\vartheta}.$$

Since $\vartheta < 1$ and $\zeta^n \rightarrow 0$ as $n \rightarrow \infty$ for all $t \in \mathbb{R}$ then one has

$$\mathbb{E} \left\| X_n(t) - \tilde{X}(t) \right\|^2 \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad \text{for each } t \in \mathbb{R}.$$

Hence we deduce that $X(t+t_n)$ converge to $\tilde{X}(t)$ in distribution. Similarly, one can get that $\tilde{X}(t-t_n)$ converge to $X(t)$ in distribution too. Therefore this fixed-point solution of the equation (1.1) is square-mean almost automorphic in distribution. which completes the proof. \square

5. EXAMPLE

Consider the following stochastic integro-differential equations

$$(5.1) \quad \begin{cases} \frac{\partial u(t,x)}{\partial t} = \frac{\partial^2}{\partial x^2} u(t,x) + g(t, u(t,x)) + \int_{-\infty}^t e^{-\omega(t-s)} f(s, u(s,x)) ds + \int_{-\infty}^t e^{-\omega(t-s)} h(s, u(s,x)) dW(s) \\ \quad + \int_{-\infty}^t e^{-\omega(t-s)} \theta(s, u(s,x)) Z(ds), \quad (t,x) \in \mathbb{R} \times (0,1), \\ u(t,0) = u(t,1) = 0, \quad t \in \mathbb{R} \\ u(0,x) = u_0(x) = 0, \quad x \in (0,1) \end{cases}$$

where W is a Q -Wiener process on $L^2(0, 1)$ with $\text{Tr } Q < \infty$ and Z is a Lévy pure jump process on $L^2(0, 1)$ which is independent of W and $\omega > 0$. The forcing terms are follows:

$$g(t, u) = \delta \sin u(\sin t + \sin \sqrt{2}t), \quad f(t, u) = \delta \sin u(\sin t + \sin \sqrt{3}t),$$

$$h(t, u) = \delta \sin u(\sin t + \sin \sqrt{5}t), \quad \theta(t, u) = \delta \sin u(\sin t + \sin \pi t)$$

with $\delta > 0$. Denote $H = V = L^2(0, 1)$. In order to write the system (5.1) on the abstract form (1.1), we consider the linear operator $A : D(A) \subset L^2(0, 1) \rightarrow L^2(0, 1)$, given by

$$\begin{aligned} D(A) &= H^2(0, 1) \cap H_0^1(0, 1), \\ Ax(\xi) &= x''(\xi) \quad \text{for } \xi \in (0, 1) \quad \text{and } x \in D(A). \end{aligned}$$

It is well-known that A generates a C_0 -semigroup $(T(t))_{t \geq 0}$ on $L^2(0, 1)$ defined by

$$(T(t)x)(r) = \sum_{n=1}^{\infty} e^{-n^2\pi^2 t} \langle x, e_n \rangle_{L^2} e_n(r),$$

where $e_n(r) = \sqrt{2} \sin(n\pi r)$ for $n = 1, 2, \dots$, and $\|T(t)\| \leq e^{-\pi^2 t}$ for all $t \geq 0$. Then the system (5.1) takes the following abstract form

$$\begin{aligned} u'(t) &= Au(t) + g(t, u(t)) + \int_{-\infty}^t B_1(t-s)f(s, u(s))ds \\ &+ \int_{-\infty}^t B_2(t-s)h(s, u(s))dW(s) \\ &+ \int_{-\infty}^t B_2(t-s) \int_{|y|_V < 1} F(s, u((s-), y))\tilde{N}(ds, dy) \\ &+ \int_{-\infty}^t B_2(t-s) \int_{|y|_V \geq 1} G(s, u((s-), y))N(ds, dy) \quad \text{for all } t \in \mathbb{R}, \end{aligned}$$

where $u(t) = u(t, \cdot)$, $B_1(t) = B_2(t) = e^{\omega t}$ for $t \geq 0$ and

$$\theta(t, u)Z(dt) = \int_{|y|_V < 1} F(t, u(t-), y)\tilde{N}(dt, dy) + \int_{|y|_V \geq 1} G(t, u(t-), y)N(dt, dy)$$

with

$$Z(t) = \int_{|y|_V < 1} y\tilde{N}(t, dy) + \int_{|y|_V \geq 1} yN(t, dy),$$

$$F(t, u, y) = h(t, u)y \cdot 1_{\{|y|_V < 1\}} \quad \text{and} \quad G(t, u, y) = h(t, u)y \cdot 1_{\{|y|_V \geq 1\}}.$$

Here, we assume that the Lévy pure jump process Z on $L^2(0, 1)$ is decomposed as above by the Lévy-It decomposition theorem.

Clearly, f, θ are almost automorphic, and F, G Poisson almost automorphic and satisfying H_1 – H_3 . Moreover, it is easy to see that the conditions (3.3)–(3.7) are satisfied with

$$\begin{aligned} m_g(t) &= \delta^2 \left(\sin t + \sin \sqrt{2}t \right)^2, \quad m_f(t) = \delta^2 \left(\sin t + \sin \sqrt{3}t \right)^2 \\ m_h(t) &= \delta^2 \|Q\|_{L(V, V)} \left(\sin t + \sin \sqrt{5}t \right)^2, \quad m_F(t) = \delta^2 \nu(B_1(0)) \left(\sin t + \sin \sqrt{3}t \right)^2 \\ m_G(t) &= \delta^2 b \left(\sin t + \sin \pi t \right)^2, \end{aligned}$$

where $B_1(0)$ is the ball in V centered at the origin with radius 1.

Obviously,

$$\begin{aligned}
L_g &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t e^{-\omega(t-s)} m_g(s) ds = \delta^2 \sup_{t \in \mathbb{R}} \int_{-\infty}^t e^{-\omega(t-s)} m_g(s) ds \left(\sin s + \sin \sqrt{2}s \right)^2 < \infty, \\
L_f &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_1(t-s)\| m_f(s) ds = \delta^2 \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_1(t-s)\| \left(\sin s + \sin \sqrt{3}s \right)^2 ds < \infty, \\
L_h &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_2(t-s)\| m_h(s) ds = \delta^2 \|Q\|_{L(V,V)} \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_2(t-s)\| \left(\sin s + \sin \sqrt{5}s \right)^2 ds < \infty, \\
L_F &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_2(t-s)\|^2 m_F(s) ds = \delta^2 \nu(B_1(0)) \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_2(t-s)\|^2 \left(\sin s + \sin \pi s \right)^2 ds < \infty, \\
L_G &= \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_2(t-s)\|^2 m_G(s) ds = \delta^2 b \sup_{t \in \mathbb{R}} \int_{-\infty}^t \|B_2(t-s)\|^2 \left(\sin s + \sin \pi s \right)^2 ds < \infty
\end{aligned}$$

Therefore, by Theorem 4.1, the equation (5.1) has a square-mean almost automorphic in distribution mild solution on \mathbb{R} whenever δ is small enough.

REFERENCES

- [1] D. Applebaum, Lévy Process and Stochastic Calculus, second edition, Cambridge University Press, (2009).
- [2] P. Bezandry and T. Diagana, Existence of almost periodic solutions to some stochastic differential equations, *Appl. Anal.*, 86, (2007), 819–827.
- [3] P. Bezandry, Existence of almost periodic solutions to some functional integro-differential stochastic evolution equations, *Statist. Probab. Lett.* 78, (2008), 2844–2849.
- [4] P. H. Bezandry and T. Diagana, Square-mean almost periodic solutions nonautonomous stochastic differential equations, *Electron. J. Differential Equations*, 2007, No. 117, 10 pp.
- [5] P. Bezandry and T. Diagana, Existence of S^2 -almost periodic solutions to a class of nonautonomous stochastic evolution equations, *Electron. J. Qual. Theory Differ. Equ.* 35, (2008), 1–19.
- [6] S. Bochner, Uniform convergence of monotone sequences of functions, *Proc. Natl. Acad. Sci. USA*, 47, (1961), 582–585.
- [7] Y. K. Chang, Z. H. Zhao, G.M. N?Gu?r?k?ata and R. Mab, Stepanov-like almost automorphy for stochastic processes and applications to stochastic differential equations, *Nonlinear Analysis: Real World Applications* 12, (2011), 1130–1139.
- [8] T. Diagana, Weighted pseudo almost periodic functions and applications, *Comptes Rendus de l’Académie des Sciences de Paris, S?rie I* 343, (10), (2006), 643–646.
- [9] T. Diagana, Weighted pseudo-almost periodic solutions to some differential equations, *Nonlinear Analysis, Theory, Methods and Applications*, 68, (8), (2008), 2250–2260.
- [10] M. A. Diop, K. Ezzinbi and M. M. Mbaye, Measure theory and S^2 - pseudo almost periodic and automorphic process: Application to stochastic evolution equations, *Afrika Matematika*, 26, (5), (2015), 779–812.
- [11] M. A. Diop, K. Ezzinbi and M. M. Mbaye, Existence and global attractiveness of a pseudo almost periodic solution in the p -th mean sense for stochastic evolution equation driven by a fractional Brownian, *Stochastics: An International Journal Of Probability And Stochastic Processes*, 87, (6), (2015), 1061–1093.
- [12] M. A. Diop, K. Ezzinbi and M. M. Mbaye, Existence and global attractiveness of a square-mean μ -pseudo almost automorphic solution for some stochastic evolution equation driven by Lévy noise, *Math. Nachr.*, 290, (8-9), 1260–1280.
- [13] M. M. Fu, Almost automorphic solutions for nonautonomous stochastic differential equations, *Journal of Mathematical Analysis and Applications*. 393, (2012), 231–238.
- [14] M. M. Fu and Z.X. Liu, Square-mean almost automorphic solutions for some stochastic differential equations, *Proc. Amer. Math. Soc.* 133, (2010), 3689–3701.
- [15] M. Kamenskii, O. Mellah, P. Raynaud de Fitte, Weak averaging of semilinear stochastic differential equations with almost periodic coefficients, *J. Math. Anal. Appl.* 427 (2015) 336–364
- [16] O. Mellah, P. Raynaud de Fitte, Counterexamples to mean square almost periodicity of the solutions of some SDEs with almost periodic coefficients, *Electron. J. Differential Equations* 2013 (91) (2013) 1–7.
- [17] Z. Liu and K. Sun, Almost automorphic solutions for stochastic differential equations driven by Lévy noise, *J. Funct. Anal.*, 266, (3), (2014), 1115–1149.
- [18] M. M. Mbaye, Square-mean μ -pseudo almost periodic and automorphic solutions for a class of semilinear integro-differential stochastic evolution equations, *Afrika Matematika*, 28, (3-4), (2017), 643–660.

- [19] M. M. Mbaye, Almost automorphic solution for some stochastic evolution equation driven by Lévy noise with coefficients S^2 -almost automorphic, *Nonauton. Dyn. Syst.* 2016, 3:85–103
- [20] G.M. N'Guérékata, Almost automorphic solutions to second-order semilinear evolution equations, *Nonlinear Analysis, Theory, Methods and Applications* 71, (12), (2009), e432-e435.
- [21] G.M. N'Guérékata, *Topics in almost Automorphy*, Springer-verlag, New York, 2005.
- [22] S. Peszat, J. Zabczyk, *Stochastic Partial Differential Equations with Lévy Noise*, Cambridge University Press, (2007).
- [23] G.D. Prato and J. Zabczyk, *Stochastic Equations in Infinite Dimensions*, *Encyclopedia of Mathematics and its Applications* 44, Cambridge University Press, Cambridge, (1992).
- [24] C. Tudor, Almost periodic solutions of affine stochastic evolutions equations, *Stoch. Stoch. Rep.* 38, (1992), 251–266.
- [25] Zhinan Xia, Pseudo almost automorphic in distribution solutions of semilinear stochastic integro-differential equations by measure theory, *Int. J. Math.*, doi.org/10.1142/S0129167X15501128
- [26] Y. Wang, Z.X. Liu, Almost periodic solutions for stochastic differential equations with Lévy noise, *Nonlinearity*, 25, (2012), 2803–2821.

DÉPARTEMENT DE MATHÉMATIQUES, FACULTÉ DES SCIENCES ET TECHNIQUE, UNIVERSITÉ CHEIKH ANTA DIOP, BP 5005, DAKAR-FANN, SENEGAL

E-mail address: mamadoumoustapha3.mbaye@ucad.edu.sn

INSTITUT MONTELLIÉRAIN ALEXANDER GROTHENDIECK, UMR CNRS 5149, UNIVERSITÉ DE MONTPELLIER

E-mail address: solym-mawaki.manou-abi@umontpellier.fr

DÉPARTEMENT SCIENCES ET TECHNOLOGIES, CUFR DE MAYOTTE, 3 ROUTE NATIONALE, 97660 DEMBNI

E-mail address: solym.manou-abi@univ-mayotte.fr