Reconstructing the formation history of top-shaped asteroids from the surface boulder distribution

Bin Cheng, Yang Yu, Erik Asphaug, Patrick Michel, Derek Richardson,
Masatoshi Hirabayashi, Makoto Yoshikawa, Hexi Baoyin

► To cite this version:

Bin Cheng, Yang Yu, Erik Asphaug, Patrick Michel, Derek Richardson, et al.. Reconstructing the formation history of top-shaped asteroids from the surface boulder distribution. *Nature Astronomy*, 2020, 10.1038/s41550-020-01226-7 . hal-02986154

HAL Id: hal-02986154

<https://hal.science/hal-02986154>

Submitted on 28 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reconstructing the formation history of top-shaped asteroids by the surface boulders distribution

Bin Cheng^{1,3}, Yang Yu^{2,*}, Erik Asphaug³, Patrick Michel⁴, Derek C. Richardson⁵, Masatoshi Hirabayashi⁶, Makoto Yoshikawa^{7,8} & Hexi Baoyin^{1,+}

¹*School of Aerospace Engineering, Tsinghua University, Beijing, China.*

²*School of Aeronautic Science and Engineering, Beihang University, Beijing, China.*

³*Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ, USA.*

⁴*Université Côte d’Azur, Observatoire de la Côte d’Azur, CNRS, Laboratoire Lagrange, Nice, France.*

⁵*Department of Astronomy, University of Maryland, College Park, MD, USA.*

⁶*Department of Aerospace Engineering, Auburn University, Auburn, AL, USA.*

⁷*Institute of Space and Astronautical Science (ISAS), Japan Aerospace Exploration Agency (JAXA), Sagamihara, Japan.*

⁸*The Graduate University for Advanced Studies (SOKENDAI), Kanagawa, Japan.*

**yuyang.thu@gmail.com*

+baoyin@tsinghua.edu.cn

Finding the basic mechanism governing the surface history of asteroids of various shapes is essential for understanding their origin and evolution. In particular, the currently visited asteroids (162173) Ryugu¹ and (101955) Bennu² by Hayabusa2 and OSIRIS-REx appear to be top-shaped. This distinctive shape, characterized by a raised equatorial bulge, is shared

by other similarly sized asteroids, including Didymos A³, 2008 EV5⁴ and 1999 KW4 Alpha⁵. However, the possibly common formation mechanism that causes the top-like shape is still under debate. One clue may lie in the boulders on their surfaces. The distribution of these boulders, that was precisely measured in unprecedented detail by the two spacecrafts^{1,2}, constitutes a record of the geological evolution of the surface regolith since the origin of these asteroids. Here, we show that during the regolith migration driven by YORP spin-up⁶⁻⁹, the surface boulders coevolve with the underlying regolith and exhibit diverse dynamical behaviors: they can remain undisturbed, sink into the regolith layer and become tilted, or totally be buried by the downslope deposition, depending on their latitudes. The predominant geological features commonly observed on top-shaped asteroids, including the boulder-rich region near the pole^{1,10}, the deficiency of large boulders at the equatorial area^{10,11} and partially buried, oblique boulders exposed on the regolith surface^{12,13}, are commensurate with this coevolution scenario. The surface regolith migration thus is the prevalent mechanism for the formation history of the top-shaped asteroids with stiffer cores.

Our investigation of the quasi-static process of boulder-regolith evolution by YORP spin-up was performed using numerical code based on the soft-sphere discrete element method¹⁴⁻¹⁶ (SSDEM). The considered structure and surface of a top-shaped asteroid consist of a “two-layer structure” with a shallow mantle of loose granular regolith covering a stiffer interior^{7,8}. This model is supported by multiple observations of the surface geologic features on Ryugu and Bennu, e.g., the longitudinal ridges and long linear grooves denoting internal stiffness and structural rigidity with a weak strength of less than a few Pa¹⁷; the measured thermal inertia and substantial material

42 flows evidencing loose and unconsolidated surface regolith^{1,13}. The pit feature of the artificial
 43 impact crater on Ryugu provides further evidence of a stronger substrate covered by cohesionless
 44 regolith¹⁸. This heterogeneity could appear as a result of stronger interparticle cohesion, grains
 45 of different materials, higher packing fraction, a big inner monolith, etc., or a combination of
 46 those particular characteristics⁸. The “granular shell” model is shown in Fig. 1a, i.e., the asteroid
 47 interior is modelled as a rigid spheroid (~ 400 m in radius), and a longitudinal slice of the regolith
 48 layer (~ 30 m in thickness) is considered for the SSDEM simulation, consisting of 2,063,044
 49 pebble-size particles. The boulders of various sizes, modelled as massive polyhedra, are located at
 50 four latitudes $\{15^\circ, 45^\circ, 75^\circ, 87^\circ\}$ as representatives of low, mid-low, mid-high, and high latitudes,
 51 respectively. We assumed these boulders are initially on top of the regolith layer, which could be
 52 the direct result of the re-accumulation that formed the asteroids after the catastrophic disruption
 53 of their parent bodies^{12,13}, or much later by ballistic sorting¹⁹ or Brazil-nut effect²⁰ during impact-
 54 induced seismic events. Two different types of regolith grains were used: one with moderate
 55 friction (internal friction angle of $\sim 25^\circ$) and one with gravel-like friction (internal friction angle
 56 of $\sim 30^\circ$). We then used two types of regolith grains for each of the four latitudes, and each of the
 57 three boulders, giving a total of 24 simulations (all simulation parameters are summarized in the
 58 supplementary information).

59 The time evolution of this regolith migration during the spin-up process is shown in Fig. 1c.
 60 The regolith remains stable at first. When the progenitor reaches the critical spin period $T_c = 4.35$ hr,
 61 a localized landslide first occurs at mid-latitudes^{9,21}. Note that T_c here is for surface landslide
 62 failure, which is different from the shedding limits and internal failure limits^{7,8}. The regolith mi-

63 gration forms a narrow zone with intense shear strain (Fig. 1c, first panel). Within the shear band,
 64 particle contacts break and remake. Such rearrangement events soften the shear strength of the
 65 granular material^{22,23}, i.e., the particles in this zone rotate and slip, transporting the regolith above
 66 this weakened band downslope along the failure surface like a frozen ice block, with an average
 67 creeping speed of less than 1 mm/s (Fig. 1b). Regolith above the slip region remains stable and
 68 undisturbed due to the low surface slope near the pole⁹. With increasing spin rate Ω , the scarp
 69 at the head of the slip zone retreats towards high latitudes, leading to the shrinking of the stable
 70 region, as shown by the kinetic energy distribution change in Supplementary Fig. 3, in parallel
 71 with the massive mass wasting towards low latitudes. The retreat of the headscarp to high latitudes
 72 and the expansion of the depletion region at mid-latitudes are both evident from the growth of the
 73 failure surface (Fig. 1c, second panel). In the later stage of the migration, due to the existence
 74 of a stable region near the equator (low surface slope), the barely moved grains here increase the
 75 resisting force along the failure surface in the landslide foot, blocking the material in the head from
 76 moving towards the foot. Thus, the creeping debris moves uphill and then overlies the pre-existing
 77 materials, forming a distinct deposit bulge near the equator. The original shear band propagates to
 78 low latitudes and subsequently creates a rupture surface inside the deposition (Fig. 1c, third panel).
 79 It is notable that the elevated geologic upheaval that marks the foot of the landslides is not at the
 80 equator at first, i.e., the collapse is limited to the mid-latitudes due to the resistance force between
 81 regolith grains. Then, as the accumulation of downslope materials from the depletion zone goes
 82 on, this bulge swells upward, leading to a steeper local slope that eventually collapses towards
 83 the equator. The raised, pronounced equatorial ridge denotes a fully sped-up progenitor for which

the top-shaped morphology emerges (Fig. 2a). The simulated topography, with low elevation at mid-latitudes, mean elevation at the polar region and high elevation at the equator, resembles the observed shapes of typical top-shaped asteroids (Fig. 2b), which is consistent with previous findings that regolith migrations are a major force that has reshaped these asteroids if the interior is stronger⁷⁻⁹. Simulations with higher friction give a similar picture, except that the stable region near the pole is larger and that the equatorial ridge is steeper (Supplementary Video 6).

The motion of boulders across the surface shows three distinct dynamical behaviours (see Supplementary Videos 2-5). The 15° and 45° boulders are located in the early migration zone, thus they start to creep downslope in synchronization with the first landslide of granular regolith. During the creeping-down process of the 45° boulder, the strength of its supporting force chain network is weakened due to the grain agitations induced by the shear deformation²⁴. These weak grain bonds generate a fluid-like behaviour; thus this boulder sinks into the regolith layer by a few metres and becomes tilted, reminiscent of the abundant partially buried boulders on Ryugu and Bennu^{12,13}. This flow-induced submergence phenomenon has also been observed in granular experiments^{22,24}. At lower latitudes, the 15° boulder is first elevated by the extruded underlying sediment, subsequently transferred downslope to the foot of the deposit upheaval, and eventually buried entirely by the accumulated debris from the collapsed bulge hill. This mechanism, i.e., boulders initially located at low latitudes being totally buried by the downslope deposition, may have caused the deficiency of large boulders at the equator as observed on Ryugu¹⁰ and Bennu²⁵. At higher latitudes, the 75° boulder remains static until the regressive headscarp of the depletion zone reaches 75° latitude. The growth of this scarp removes support from adjacent grains, forming

a fragile force-chain network downstream (Fig. 3). The weak support increases the equivalent local slope and thus triggers a downslope migration of the boulder. Finally, this boulder sinks into the granular mantle and leaves only a partially exposed body above the regolith layer, generating a distinct outcrop-like landform (Fig. 4). The 87° boulder, however, is located in a stable region (see Fig. 2a) and thus does not move throughout the reshaping process of the top-shaped asteroids. This result is consistent with the geologic indication of a stable pole deduced from the disorganized boulder structure near Bennu's pole and the non-slipping polar boulder Otohime on Ryugu^{1,12}. We infer that Otohime could be a primordial relic of the original disruption/reaccumulation of Ryugu's parent body. Future studies on the geologic age of Otohime from thermal or fracture properties would provide significant constraints on when the catastrophic impact event that formed Ryugu occurred. Note that we do not rule out the possibility that other exhumation processes could contribute to the subsequent evolution of these boulders. Some of the buried boulders in the equatorial region on Ryugu could be exhumed again by migrating grains from the ridge towards the current geopotential lows at higher latitudes¹². Other mechanisms including Brazil nut effect²⁰ and ballistic sorting¹⁹ could also further modify these landforms.

Our results show that the top shapes of rubble pile asteroids can be evolved via regolith migrations from mid-latitudes towards the equator driven by the YORP spin-up effect. If the interior is stronger than the outer regolith, then miniature surface landslides could be frequent as the spin rate approaches the critical spin limit^{9,21}. These landslides are to be distinguished from their counterpart on large-gravity bodies like Mars, the Moon or Earth. On such planets the large boulders usually bounce and deposit near the toe of the avalanche sediment²⁶. By contrast, regolith migra-

tion on low gravity rubble piles, driven by YORP, once near the spin limit, creeps imperceptibly at glacial speeds. A detailed look into these creep-like landslides shows that the boulders are in a “co-evolving” relationship with the underlying granular regolith, which may account for the unique geological landforms observed on top-shaped asteroids, i.e., the stable boulders near the pole, the abundant partially buried boulders and the deficiency of large boulders at the equator. We find consistency of the derived shapes and boulder features between the observations and the numerical results, which lends support to our model as a way to decipher the evolution history of top-shaped asteroids. Further simulations show that the large craters that overlay the equator of Ryugu would have been eroded by the migrating regolith if they were formed before the equatorial ridges (Supplementary Fig. 7), implying an early-stage formation of the overall shapes¹. This would appear to be in tension with the long timescales over which YORP effect could significantly modify the spins²⁷. A plausible explanation is that the progenitor bodies that emerged from the catastrophic disruptions are already oblate shapes with relatively high spin rates, which is a possible result of the direct reaccumulation of fragments in the case of large starting angular momentum²⁷. These early-formed “diamond-shaped” rubble piles with high spin rates could serve as a midway starting point that facilitates the formation of top shapes, i.e., the YORP-induced migrations further reshaped the bodies and remodeled their surface features like large boulders to the current geomorphology. Another possibility is these craters could be created by detachment of a small chunk from the equatorial ridge at a more rapid rotation state²⁸. This mechanism seems to be plausible given that the past spin period of Ryugu could be up to around 3 hours as deduced from its east-west dichotomy²⁹. In such a scenario, the circum-equatorial ridge and overlying craters could both be

landscapes sculpted by YORP spin-up.

Methods

We performed discrete element simulations^{14–16} using an original soft-particle N -body code DEMBody³⁰, which is capable of tracing the quasi-static behaviour of particle-particle and particle-polyhedron interactions. In this code, the nonlinear contact forces between two contacting objects (sphere-sphere, sphere-polyhedron, sphere-wall) are calculated based on Hertz-Mindlin contact theory³¹ according to the mutual overlap (typically $<1\%$ of their radii). A dimensionless coefficient μ is used to control the stick-slip friction between colliding particles. A physically based rotational resistance model incorporating rolling and twisting friction was implemented into this code, parameterized by a quantity β that represents a statistical measure of real particle nonsphericity^{30,32}. To calculate the contact between non-spherical boulders and grains, the boulder is represented as an assembly of movable walls with the same geometry and inertia tensor³⁰, which allows it to behave as a rigid body. A second-order leapfrog integrator is used for integration of the equations of motion of both particles and boulders.

Our simulation scheme is as follows. First, we prepared a polydisperse granular medium consisting of 2,063,044 cohesionless particles with properties chosen according to Ryugu’s regolith¹, i.e., material density of 2.4 g/cc. A slight dispersion in the grain radius was considered, with a random distribution ranging from 0.4 m to 0.6 m to avoid any crystallization of the packing. The particles were dropped onto a rigid, frictional shell with a radius of 400 m and width of 100 m confined by two vertical frictionless walls⁹, as shown in Fig. 1a, forming a longitudinal slice of

the asteroid surface with a thickness of ~ 30 m. A cuboid boulder, the size of which is similar to that of typical boulders on Ryugu, was initially located above the interface of the granular media. The whole system is under the two-component gravity⁹ exerted by the rocky interior (bulk density of 1.2 g/cc)¹ and the regolith shell. After a settling-down phase of the granular-polyhedron system, we started to accelerate the asteroid linearly following the prescribed spin-up path shown in Fig. 1b. The acceleration was performed until massive debris shedding occurred at the equator. The angular acceleration used in this study is $2.5 \times 10^{-10} \text{ rad/s}^2$, which is much smaller than the reference values used in previous works^{8,32}. By using this slow spin-up acceleration, we confirm that the motion of the granular grains would immediately stop at any point in time when the spin-up is stopped, which means that the regolith layer maintains quasi-static equilibrium during the spin-up process (see Supplementary Information).

Regolith bed geometry. We investigated the dependency of the boulder evolution on the geometry of the regolith bed. By using granular beds with depth of 45 m and of 20 m, which consists of 3,246,524 particles and 1,452,565 particles, respectively, we tested the influence of regolith bed depth (Supplementary Figure 4-5). Simulations give results similar to those described above, in which tilted boulders and buried boulders emerge with the surface migration driven by YORP spin-up. By using a granular bed with width of 160 m, which consists of 3,332,716 particles, we tested the influence of confining sidewalls (Supplementary Video 7). Simulations indicate the size of surrounding walls is large enough to eliminate the boundary effect. By using a spherical wedge-shaped bed with angular width of 20° , which consists of 2,063,045 particles, we tested the influence of the boundary shape (Supplementary Figure 6). Simulations indicate the shrinkage effect from

surrounding walls has only a minor influence on the boulder dynamics for our simulation settings.

By using a granular bed of both hemispheres, which consists of 4,126,088 particles, we tested the symmetry of the regolith migration. Simulations show the sliding debris meet and accumulate symmetrically at the equator (Supplementary Video 9). The overall geomorphology is similar to that in hemisphere cases.

Equatorial crater. We investigated the evolution of equatorial craters during the YORP spin-up. A crater with radius of 80 m was generated at the equator before rotational acceleration (Supplementary Fig. 7). Simulations show that the migrating debris would have filled the equatorial craters if these craters formed before the equatorial ridges, which implies an early-stage formation of the overall morphology.

Boulders configuration. We investigated the dependency of the boulder evolution on their configurations. Simulations with boulders ten meters below the regolith surface show these initially underground boulders keep buried throughout the creeping evolution (Supplementary Videos 10-11). We also tested the interaction between multiple boulders. Given that the number density of boulders¹⁰ larger than 15 m on Ryugu is $\sim 100 \text{ km}^{-2}$, we randomly placed eight boulders, either on the surface or embedded in regolith, at middle and low latitudes of the simulated longitudinal slice. Simulations give results similar to those described above, in which surface boulders sink into or are buried by the creeping regolith, while underground boulders remain buried (Supplementary Video 12).

Data availability. The data that support the plots within this paper and other findings of this study are available from the corresponding authors upon reasonable request.

Code availability. The code used to generate the datasets is available from the corresponding author on reasonable request.

1. Watanabe, S. *et al.* Hayabusa2 arrives at the carbonaceous asteroid 162173 Ryugu—A spinning top-shaped rubble pile. *Science* **364**, 268–272 (2019).
2. Lauretta, D. S. *et al.* The unexpected surface of asteroid (101955) Bennu. *Nature* **568**, 55–60 (2019).
3. Benner, L. A. *et al.* Radar imaging and a physical model of binary asteroid 65803 Didymos. In *Bull. Am Astron. Soc.*, vol. 42, 1056 (2010).
4. Busch, M. W. *et al.* Radar observations and the shape of near-Earth asteroid 2008 EV5. *Icarus* **212**, 649–660 (2011).
5. Ostro, S. J. *et al.* Radar imaging of binary near-Earth asteroid (66391) 1999 KW4. *Science* **314**, 1276–1280 (2006).
6. Walsh, K. J., Richardson, D. C. & Michel, P. Rotational breakup as the origin of small binary asteroids. *Nature* **454**, 188–191 (2008).
7. Hirabayashi, M., Sánchez, P. & Scheeres, D. J. Internal structure of asteroids having surface shedding due to rotational instability. *Astrophys. J.* **808**, 63 (2015).

- 226 8. Sánchez, P. & Scheeres, D. J. Rotational evolution of self-gravitating aggregates with cores of
227 variable strength. *Planet. Space. Sci.* **157**, 39–47 (2018).
- 228 9. Sánchez, P. & Scheeres, D. J. Cohesive regolith on fast rotating asteroids. *Icarus* **338**, 113443
229 (2020).
- 230 10. Michikami, T. *et al.* Boulder size and shape distributions on asteroid Ryugu. *Icarus* **331**,
231 179–191 (2019).
- 232 11. DellaGiustina, D. N. *et al.* Properties of rubble-pile asteroid (101955) Bennu from OSIRIS-
233 REx imaging and thermal analysis. *Nat. Astron.* **3**, 341–351 (2019).
- 234 12. Sugita, S. *et al.* The geomorphology, color, and thermal properties of Ryugu: Implications for
235 parent-body processes. *Science* **364**, 252 (2019).
- 236 13. Walsh, K. J. *et al.* Craters, boulders and regolith of (101955) Bennu indicative of an old and
237 dynamic surface. *Nat. Geosci.* **12**, 242–246 (2019).
- 238 14. Cundall, P. A. & Strack, O. D. A discrete numerical model for granular assemblies. *Geotech-*
239 *nique* **29**, 47–65 (1979).
- 240 15. Sánchez, P. & Scheeres, D. J. Simulating asteroid rubble piles with a self-gravitating soft-
241 sphere distinct element method model. *Astrophys. J.* **727**, 120 (2011).
- 242 16. Schwartz, S. R., Richardson, D. C. & Michel, P. An implementation of the soft-sphere discrete
243 element method in a high-performance parallel gravity tree-code. *Granul. Matter* **14**, 363–380
244 (2012).

- 245 17. Barnouin, O. S. *et al.* Shape of (101955) Bennu indicative of a rubble pile with internal
246 stiffness. *Nat. Geosci.* **12**, 247–252 (2019).
- 247 18. Arakawa, M. *et al.* An artificial impact on the asteroid (162173) Ryugu formed a crater in the
248 gravity-dominated regime. *Science* **368**, 67–71 (2020).
- 249 19. Wright, E. *et al.* Boulder stranding in ejecta launched by an impact generated seismic pulse.
250 *Icarus* **337**, 113424 (2020).
- 251 20. Matsumura, S. *et al.* The brazil nut effect and its application to asteroids. *Mon. Not. R. Astron.*
252 *Soc.* **443**, 3368–3380 (2014).
- 253 21. Scheeres, D. J. *et al.* The dynamic geophysical environment of (101955) Bennu based on
254 OSIRIS-REx measurements. *Nat. Astron.* **3**, 352–361 (2019).
- 255 22. Reddy, K., Forterre, Y. & Pouliquen, O. Evidence of mechanically activated processes in slow
256 granular flows. *Phys. Rev. Lett.* **106**, 108301 (2011).
- 257 23. Germanovich, L. N., Kim, S. & Puzrin, A. M. Dynamic growth of slip surfaces in catastrophic
258 landslides. *Proc. Math. Phys. Eng. Sci.* **472**, 20150758 (2016).
- 259 24. Nichol, K., Zanin, A., Bastien, R., Wandersman, E. & van Hecke, M. Flow-induced agitations
260 create a granular fluid. *Phys. Rev. Lett.* **104**, 078302 (2010).
- 261 25. Walsh, K. J. *et al.* Geology of Bennu’s equatorial ridge. In *Asteroid Science 2019* (2019).
- 262 26. Gray, J. M. N. T. Particle segregation in dense granular flows. *Annu. Rev. Fluid. Mech.* **50**,
263 407–433 (2018).

- 264 27. Michel, P. *et al.* Collisional formation of top-shaped asteroids and implications for the origins
265 of Ryugu and Bennu. *Nat. Commun.* **11**, 1–11 (2020).
- 266 28. Tardivel, S., Sánchez, P. & Scheeres, D. J. Equatorial cavities on asteroids, an evidence of
267 fission events. *Icarus* **304**, 192–208 (2018).
- 268 29. Hirabayashi, M. *et al.* The western bulge of 162173 Ryugu formed as a result of a rotationally
269 driven deformation process. *Astrophys. J. Lett.* **874**, L10 (2019).
- 270 30. Cheng, B., Yu, Y. & Baoyin, H. Numerical simulations of the controlled motion of a hopping
271 asteroid lander on the regolith surface. *Mon. Not. R. Astron. Soc.* **485**, 3088–3096 (2019).
- 272 31. Somfai, E., Roux, J.-N., Snoeijer, J. H., Van Hecke, M. & Van Saarloos, W. Elastic wave
273 propagation in confined granular systems. *Phys. Rev. E.* **72**, 021301 (2005).
- 274 32. Zhang, Y. *et al.* Rotational failure of rubble-pile bodies: Influences of shear and cohesive
275 strengths. *Astrophys. J.* **857**, 15 (2018).
- 276 33. Peters, J. F., Muthuswamy, M., Wibowo, J. & Tordesillas, A. Characterization of force chains
277 in granular material. *Phys. Rev. E.* **72**, 041307 (2005).
- 278 34. Hayabusa2 Optical Navigation Camera (ONC) dataset at JAXA Data Archives and Transmis-
279 sion System (DARTS),
280 http://darts.isas.jaxa.jp/pub/hayabusa2/onc_bundle

281 **Corresponding Author** Correspondence and inquiries for materials should be addressed to H.B. and Y.Y.

Acknowledgements This work is funded by the National Science Fund for Distinguished Young Scholars of China (No. 11525208). Y.Y. acknowledges support from the Natural Science Foundation of China (No. 11702009). P.M. acknowledges funding from the French space agency CNES, from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 870377 (project NEO-MAPP) and from Academies of Excellence: Complex systems and Space, environment, risk, and resilience, part of the IDEX JEDI of the Université Côte d'Azur. The authors acknowledge supports from JSPS Core-to-Core program "International Network of Planetary Sciences". We are grateful to the Hayabusa2 ONC team for providing images which are available at the JAXA Data Archives and Transmission System (DARTS) at http://darts.isas.jaxa.jp/pub/hayabusa2/onc_bundle.

Author contributions B.C. performed the soft-sphere numerical simulations and analysed the numerical results. B.H. and Y.Y. initiated the project, designed the simulations and led the research. E.A. and D.C.R. contributed to the discussion of the two layer model and the creeping process. M.H. provides essential comments on the interior structure and deformation. P.M. initiated the collaboration between the institutions with Y.Y. and provided outstanding questions on scope of the research with M.Y. All authors contributed to interpretation of the results and preparation of the manuscript.

Competing Interests The authors declare no competing interests.

Figure 1 Asteroid morphological change during the YORP spin-up process. **a**, “Two-layer structure” with a shallow mantle of loose granular regolith covering a stiffer interior. A large boulder lies on the regolith surface. **b**, Spin-up path and consequent creep speed of global regolith migrations for a 75° boulder with moderate-friction regolith case. **c**, Four selected frames of the regolith and boulder at different spin periods showing the regolith migration under the YORP effect. The cuboid represents the boulder and is located at 75° latitude initially. The regolith has moderate friction. When the progenitor reaches the critical spin period $T_c = 4.35$ hr, the centrifugal tension induces a narrow zone with intense shear strain (dashed line in first panel; see Supplementary Fig. 2 for the shear heat of regolith particles). This shear band expands towards high latitudes, and creeping debris moves to low latitudes along the rupture surface in the deposition (dashed line in third panel). At the end of the simulation, deposit sediment accumulates at low-latitudes, forming a pronounced ridge along the equator (fourth panel). The failure surface acts as a dynamical divider separating the static zone from the depletion zone, whose growth controls the landslides across the regolith bed. **Animation of this figure: Supplementary Video 1.**

Figure 2 The simulated geomorphology. **a**, The topography changes during the YORP spin-up process, showing the barely moved pole and migrating debris from mid-latitudes overlying the pre-existing materials near the equator. Here the elevation is the radial height of particles above the interior shell. Data are for a moderate-friction case. **b**, Simulated morphological profile (black line) compared with Ryugu's (blue)¹, Bennu's (red)²¹, 1999 KW4's (green)⁵ and 2008 EV5's (yellow)⁴ average shapes. All profiles are normalized by the corresponding equivalent radius. Shaded areas stand for the 1σ error bands.

Figure 3 Visualization of the force networks excited by the creeping boulder. Colour indicates the elastic energy, E , stored in each particle relative to the average elastic energy, E_0 . Note that only those particles with $E > E_0$ are considered part of the force chains³³. The white-curve indicates the free surface of the creeping regolith. Red arrows show the sinking and tilting directions of the boulder. The chain structure downstream is significantly fragmented and spatially sparse due to the removal of adjacent grains by the retrogressive landslide, leading to subduction migration of the boulder. Data are for a 75° boulder with moderate-friction regolith case.

Figure 4 Typical landforms of boulders sculpted by YORP evolution. The 87° boulder remains stable, consistent with the non-slipping polar boulder Otohime on Ryugu¹². The boulders at mid-latitudes sink into the regolith layer by a few metres and become tilted. These landforms generated by regolith migrations are likely the geophysical origins of the partially buried/oblique boulders on Ryugu¹² and Bennu¹³. (panel d, hyb2_onc_20180831_101059_ panel e, hyb2_onc_20180801_160642.tnf_l2c and panel f, hyb2_onc_20180720_075208.tvf_l2b are from Hayabusa2 ONC dataset³⁴.)