Brønsted Acid and H-Bond Activation in Boronic Acid Catalysis

Shaofei Zhang, David Lebœuf, Joseph Moran

► To cite this version:

Shaofei Zhang, David Lebœuf, Joseph Moran. Brønsted Acid and H-Bond Activation in Boronic Acid Catalysis. *Chemistry - A European Journal*, 2020, 26 (44), pp.9883-9888. 10.1002/chem.202001902 . hal-02990292

HAL Id: hal-02990292

<https://hal.science/hal-02990292>

Submitted on 30 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Brønsted Acid and H-Bond Activation in Boronic Acid Catalysis

Shaofei Zhang, David Lebœuf and Joseph Moran*

S. Zhang, Dr. D. Lebœuf and Prof. Dr. J. Moran
University of Strasbourg, CNRS, ISIS UMR 7006
67000 Strasbourg, France
E-mail: moran@unistra.fr

Supporting information for this article is given via a link at the end of the document.

Abstract: Boronic acid catalysis has emerged as a mild method for promoting a wide variety of reactions. It has been proposed that the mode of catalysis involves Lewis acid or covalent activation of hydroxyl groups by boron, but limited mechanistic evidence exists. Here, we reinvestigate representative boronic acid-catalyzed reactions of alcohols and oximes. We interpret a series of control experiments with boronic and Brønsted acids along with correlations between their reactivity and their acidity measured by the Gutmann-Beckett method. Overall, we conclude that the major modes of catalysis involve either dual H-bond catalysis or Brønsted acid catalysis. We show that strong Brønsted acids are generated *in situ* from covalent assembly of the boronic acids with hexafluoroisopropanol, explaining why the solvent had such a major impact on the reactivity. This new insight should guide the future development of boronic acid catalysis, where the diverse and solvent-specific nature of catalytic modes has been overlooked.

Boronic acids have emerged as a promising class of catalysts^[1] that enable dehydrative nucleophilic substitution or rearrangements of alcohols,^[2] Beckmann rearrangement of oximes,^[3] and various reactions involving either carboxylic acids^[4] or epoxides^[5] under mild conditions. In depth studies into the catalytic mechanism have been performed in the case of carboxylic acids,^[4c,4h] whereas only preliminary mechanistic evidence exists for the reactions of alcohols and oximes.^[2d,2,3b] The arylboronic acid catalyst systems required for reactions involving alcohols and oximes (**B1-B3**, Scheme 1) are substantially more electrophilic than those used for the activation of carboxylic acids, the former requiring either multiple electron-withdrawing groups, cationic boronic acids or complexation with highly electronically deactivated diols. Another critical parameter in these reactions is the solvent. Our group as well as many others have pointed out the enabling effect of solvents, such as hexafluoroisopropanol (HFIP)^[6-7] and nitromethane (MeNO₂),^[8] on Brønsted and Lewis acid-catalyzed reactions through the formation of an H-bond network. In the case of HFIP, we emphasized that the role of the catalysts was to significantly increase the acidity of an H-bond cluster of HFIP, which was the true catalytically active species.^[7i,7m] Our reflections on the reactivity of arylboronic acids started during our investigations on the TfOH-catalyzed arylative ring-opening of unactivated cyclopropanes in HFIP (Table 1, entry 1.1).^[7m] We were puzzled by the reactivity of two catalyst systems typically used for activation of alcohols and oximes (**B1** and **B3**), as they were able to trigger the ring-opening of phenylcyclopropane to generate product **1** (entries 1.2 and 1.5). Given the absence of an OH functional group in the substrate and the lack of Frustrated Lewis

Scheme 1. Representative boronic acid catalyzed transformations of alcohols and oximes.

Pairs,^[9] we surmised that it would be unlikely that this reaction features covalent or Lewis acid catalysis. One plausible mechanism would entail catalysis by a Brønsted acid, which would be generated through the coordination of the boronic acid (in the case of **B1**) or the boronate ester (in the case of **B3**) with HFIP.^[10] Indeed, the presence of 2,6-di-*tert*-butylpyridine, a bulky Brønsted base commonly used to distinguish between Lewis and Brønsted acid catalysis,^[11] completely shut down the reactivity in both cases, consistent with Brønsted acid catalysis (entries 1.3 and 1.6). Additionally, Brønsted acids weaker than TfOH were not effective catalysts (entries 1.10-1.13), leading us to suspect that very strong Brønsted acids might have been produced from the boronic acids under the reaction conditions. These observations led us to wonder whether certain previously reported boronic acid catalyzed reactions might also be the result of Brønsted acid

COMMUNICATION

Table 1. Comparison between boronic and Brønsted acids for the catalytic ring-opening hydroarylation of phenylcyclopropane.

Entry	Catalyst	Yield [%] ^[a]
1.1	TfOH	97
1.2	B1	68
1.3	B1	<1 ^[b]
1.4	B2	<1
1.5	B3	96
1.6	B3	<1 ^[b]
1.7	HCl	94
1.8	H ₂ SO ₄	92
1.9	CSA	95
1.10	CF ₃ CO ₂ H	36
1.11	(COOH) ₂	12
1.12	CH ₃ CO ₂ H	<5
1.13	B(OH) ₃	<5

[a] Yields were determined by ¹H NMR using hexamethyldisiloxane as an external standard. [b] In the presence of 15 mol% 2,6-di-*tert*-butylpyridine. CSA = camphorsulfonic acid.

catalysis.

Herein, we address the above issues by comparing boronic and Brønsted acid catalysts with nine formerly disclosed boronic acid catalyzed reactions spanning the four representative studies summarized in Scheme 1. We will provide a correlation between the reactivity profiles generated by the various boronic and Brønsted acids and their influence on triethylphosphine oxide (Gutmann-Beckett method).^[12] We found that in eight of the nine reactions that were probed, Brønsted acid or H-bond catalysis, rather than a Lewis acid or covalent activation mechanism, is likely the main source of reactivity.

At the outset, to correlate the catalytic effects observed for boronic and Brønsted acids with their physicochemical properties, we elected to compare their interaction with triethylphosphine oxide (TEPO), a representative weak Lewis base, in various solvents (Figure 1). The strength of the interaction between the additive and TEPO can be inferred from the change in chemical shift in the corresponding ³¹P NMR spectrum, compared to that obtained from a control experiment performed in the absence of additive and HFIP (i.e., 46.1 ppm in toluene-*d*₈). The control experiments confirmed our hypothesis regarding the pivotal role of the solvent. Indeed, in the presence of MeNO₂ or HFIP/MeNO₂ (4:1), we observed substantial shifts in the ³¹P NMR signal (53.0 and 67.1 ppm, respectively), indicating that the solvents are non-innocent in the activation of alcohols, even in the absence of Lewis or Brønsted acids. This does not come as a surprise since we and others have noticed similar reactivity trends in the past for HFIP and MeNO₂. The former solvent forms aggregates that are excellent H-bond donors,^[7,8] while the latter templates the formation of similar aggregates through interactions with molecules such as water.^[8a,13] In the case of the HFIP/MeNO₂ mixture, adding 2,6-di-*tert*-butylpyridine did not affect the ³¹P NMR shift, confirming that, without catalyst, no Brønsted acid is generated. Mixing catalyst system **B3** with TEPO gave rise to a few new resonances, the highest frequency of which (90.3 ppm) is

12.1 ppm further downfield than the signal generated due to the same experiment carried out with B(C₆F₅)₃·H₂O (78.2 ppm). The influence of **B3** in HFIP/MeNO₂ (4:1) thus appears to be stronger than B(C₆F₅)₃·H₂O, and comparable to that of HCl (90.7 ppm) or CSA (91.3 ppm), in close agreement with the ability of **B3** to promote the opening of unactivated cyclopropanes (see Table 1). This is very different from the situation in toluene, where CSA shifts the signal of TEPO nearly 20 ppm further downfield than **B3**. At least one species produced from the components of **B3**, presumably a highly electrophilic hexafluoropinacol boronate ester, can serve to generate a strong Brønsted acid in HFIP. It should be highlighted that the diol component of **B3** is essential here, as no shift was observed with the boronic acid alone (**B3'**). In line with these suggestions, it was established that a strongly Brønsted acidic species is formed from the covalent assembly of pentafluorophenylboronic acid and oxalic acid, another electron-poor bidentate species.^[14] Likewise, the shift produced by the addition of **B1** (84.5 ppm) is significantly higher than that produced by B(C₆F₅)₃·H₂O, congruent with its demonstrated reactivity (see Table 1). Although this experiment does not distinguish whether Brønsted or Lewis acids are causing the observed shifts *per se*, strong boron Lewis acids such as B(C₆F₅)₃ are well known to rapidly react with adventitious water to form hydrates that are strong Brønsted acids.^[15] In a similar way, the large magnitude of the observed shift in the ³¹P NMR means that strong Brønsted acids are almost certainly produced by **B1** and **B3** in HFIP. For these reasons, the mild *pK_a* values established for these boronic acids in DMSO or water cannot be transposed to reactions carried out in HFIP and HFIP/MeNO₂ in order to predict reactivity. Indeed, none of the shifts corresponding to the boronic acids in the absence of HFIP exceeded 61.5 ppm (see the Supporting Information for details). Lastly, we found that boronic acid **B2** (66.8 ppm in HFIP/MeNO₂ 4:1) induces a shift in the ³¹P NMR comparable to oxalic acid (69.5 ppm), in agreement with the lack of reactivity observed in the ring-opening transformation. Of note, in the absence of the diol component of **B3**, the shift is similar to that of **B2** (66.9 ppm).

Figure 1. Gutmann-Beckett plot showing the influence of an additive (3 equiv) on TEPO (1 equiv) in HFIP/MeNO₂ (4:1) as expressed by the variations in chemical shift of the highest frequency signal observed in the ³¹P NMR spectrum when compared to the reference TEPO in toluene-*d*₈. **B3'** = **B3** in the absence of diol.

Based on these results, the first set of transformations examined was the Friedel-Crafts reaction of primary benzylic alcohols **2a-2c** catalyzed by hexafluoroantimonate salt **B1** in HFIP/MeNO₂

COMMUNICATION

(4:1).^[29] The three substrates reacted as previously described (Table 2, entry 2.1) but not in the presence of 2,6-di-*tert*-butylpyridine (entry 2.2).^[16] Replacing the cationic boronic acid by Brønsted acids spanning a wide pK_a range revealed that, in all cases, the boronic acid may be substituted with a Brønsted acid with similar results, although the strength of the required acid varies with the nature of the electronic substituents on the benzylic alcohol. The conjugate acid of the boronic acid catalyst, HSbF_6 , catalyzed the reaction of all three substrates (entry 2.3), albeit in lower yields. In the case of benzylic alcohols **2a** and **2b**, the weak Brønsted acid oxalic acid was surprisingly able to promote the reaction under otherwise identical conditions (entries 2.9-2.11). Electronically deactivated alcohol **2c** needed a stronger Brønsted acid but its transformation could still be effectively catalyzed by H_2SO_4 (entry 2.6). Previously, the possibility of Brønsted acid catalysis was ruled out on the basis of: 1) the lack of reactivity of **2b** with $\text{CF}_3\text{CO}_2\text{H}$ and 2) the fact that a different boronic acid (**B2**), which has a comparable pK_a in H_2O and DMSO to the catalyst used (**B1**),^[2b,17] does not facilitate the reaction in $\text{HFIP}/\text{MeNO}_2$. However, as the data provided in Figure 1 emphasized, **B2** and $\text{CF}_3\text{CO}_2\text{H}$ do not produce a suitably strong Brønsted acid in $\text{HFIP}/\text{MeNO}_2$ and thus are not suitable points of reference. The reduced reactivity of **B3** compared to **B1**, can be explained by the sequestration of the diol component of **B3** by H-bonding with an excess of benzylic alcohol substrate,^[18] leaving behind boronic acid **B3'** as a much weaker catalyst as expected from Figure 1.

Table 2. Comparison between boronic and Brønsted acids for the catalytic dehydrative Friedel-Crafts reactions of benzylic alcohols.^[a]

2a: X = H, Ar = *m*-xylene **2b:** X = Br, Ar = mesitylene **2c:** X = NO_2 , Ar = *p*-xylene

Entry	Catalyst	Yield 3a [%] ^[b]	Yield 3b [%] ^[b]	Yield 3c [%] ^[c]
2.1	B1	95	94	40
2.2	B1	<1 ^[d]	<1 ^[d]	<1 ^[e]
2.3	B2	<1	<1	<1
2.4	B3	62	19	5
2.5	$\text{HSbF}_6 \cdot 6\text{H}_2\text{O}$	79	40	45
2.6	TiOH	98	98	95
2.7	HCl	90	92	<1
2.8	H_2SO_4	95	94	90
2.9	CSA	95	96	<1
2.10	$\text{CF}_3\text{CO}_2\text{H}$	92	<1	<1
2.11	$(\text{COOH})_2$	92	47	<1
2.12	$\text{CH}_3\text{CO}_2\text{H}$	85	<1	<1
2.13	$\text{B}(\text{OH})_3$	15	<1	<1

[a] Yields were determined by ^1H NMR using hexamethyldisiloxane as an external standard. [b] 10 mol% catalyst, 50 °C, 24 h. [c] 20 mol% catalyst, 80 °C, 48 h. [d] With 15 mol% 2,6-di-*tert*-butylpyridine. [e] With 30 mol% 2,6-di-*tert*-butylpyridine.

The second transformation that we examined was the Beckmann rearrangement of oximes to amides catalyzed by **B3**. Previous studies on acetophenone oxime (**4a**) supported a mechanism involving slow formation of a catalytically competent O-boronyl oxime ester.^[3b] We tested three representative aryl-alkyl (**4b**), aryl-aryl (**4c**) and alkyl-alkyl oximes (**4d**) from the original publication, as well as **4a**, all under the reported conditions. The

transformations catalyzed by **B3** were efficient in each instance (Table 3, entry 3.3) but not in the presence of 2,6-di-*tert*-butylpyridine (entry 3.4). In the case of **4a**, none of the Brønsted acids tested promoted the rearrangement. In the catalytic experiments with **B3**, the reaction kinetics for **4a** did display an induction period consistent with a slow catalyst formation and the previously proposed covalent mechanism. However, since all prior optimization and mechanistic studies were conducted on **4a**, this might have led the authors to conclusions about the mechanism which do not hold for other substrates. For substrates **4b-4d**, a screen of Brønsted acids revealed several to be capable of promoting the reaction with either a similar efficiency or even more effectively than the **B3** catalyst system (entries 3.5-3.9). CSA proved to be particularly effective in the case of substrates **4b** and **4c**, while $\text{CF}_3\text{CO}_2\text{H}$ was able to promote the reaction with **4d** (entries 3.8-3.9). Indeed, the kinetic profile of the reaction of oxime **4b** catalyzed by **B3** proved to be nearly identical to that catalyzed by CSA, with no observation of an induction period expected for a mechanism involving slow formation of a catalytically active acyl oxime species (see Supporting Information for details). These experiments thus support Brønsted acid catalysis, rather than covalent catalysis, being the dominant mechanism for substrates **4b-4d**. Covalent activation and Brønsted acid catalysis therefore appear to be competitive catalytic mechanisms in the Beckmann rearrangement, with three of the four representative substrates dominated by Brønsted acid catalysis.

Table 3. Comparison between boronic acids and Brønsted acids for the catalytic Beckmann rearrangement.^[a]

4a: $\text{R}_1 = \text{C}_6\text{H}_5$, $\text{R}_2 = \text{Me}$

4b: $\text{R}_1 = 4\text{-OH-(C}_6\text{H}_4\text{)}$, $\text{R}_2 = \text{Me}$

4c: $\text{R}_1 = \text{Ph}$, $\text{R}_2 = \text{Ph}$

4d: $\text{R}_1 = \text{R}_2 = \text{-(CH}_2\text{)}_5\text{-}$

Entry	Catalyst	Yield 5a [%] ^[b]	Yield 5b [%] ^[c]	Yield 5c [%] ^[c]	Yield 5d [%] ^[b]
3.1	B1	<1	20	<10	<1
3.2	B2	<1	<1	<1	<1
3.3	B3	98	94	85	45
3.4	B3	<1 ^[d]	<1 ^[e]	<1 ^[e]	<1 ^[d]
3.5	TiOH	<1	40	96	<1
3.6	HCl	<1	16	26	<1
3.7	H_2SO_4	<1	<1	<1	<1
3.8	CSA	<1	90	86	20
3.9	$\text{CF}_3\text{CO}_2\text{H}$	<1	36	21	78
3.10	$(\text{COOH})_2$	<1	<1	<1	<1
3.11	$\text{CH}_3\text{CO}_2\text{H}$	<1	<1	<1	<1
3.12	$\text{B}(\text{OH})_3$	<1	<1	<1	<1

[a] Yields were determined by ^1H NMR using hexamethyldisiloxane as an external standard. [b] 30 mol% catalyst and perfluoropinacol, 80 °C, 24 h. [c] 5 mol% catalyst and perfluoropinacol, 25 °C, 24 h. [d] 30 mol% catalyst and perfluoropinacol, 80 °C, 24 h. [e] With 45 mol% 2,6-di-*tert*-butylpyridine. [f] With 7.5 mol% 2,6-di-*tert*-butylpyridine.

The third reaction that we analyzed was the carbocyclization of allylic alcohols, reported to be catalyzed by boronic acid **B2** in MeNO_2 .^[2f] The transformation occurred as described (Table 4,

COMMUNICATION

entry 4.2), but was inefficient in the presence of 2,6-di-*tert*-butylpyridine (entry 4.3). Likewise, **B1** and **B3** have a relatively similar efficacy (entries 4.1 and 4.4). In contrast, a Brønsted acid as weak as oxalic acid enabled the reaction under otherwise identical conditions (entries 4.11). Both oxalic acid and boronic acids are known to act as dual H-bond donors,^[19] and likely act as H-bond catalysts here. The possibility of H-bond activation may have been previously overlooked, since in the original disclosure, control experiments designed to compare **B2** to Brønsted acids were performed with *p*-TsOH only (entry 4.8).

Table 4. Comparison between boronic acids and Brønsted acids for the catalytic carbocyclization of allylic alcohols.

Entry	Catalyst	Yield [%]
4.1	B1	72
4.2	B2	52
4.3	B2	<1 ^[a]
4.4	B3	67
4.5	TfOH	81
4.6	HCl	<1
4.7	H ₂ SO ₄	36
4.8	<i>p</i> -TsOH	24
4.9	CSA	<1
4.10	CF ₃ CO ₂ H	12
4.11	(COOH) ₂	50
4.12	CH ₃ CO ₂ H	12
4.13	B(OH) ₃	<1

[a] Yields were determined by ¹H NMR using hexamethyldisiloxane as an external standard. [b] In the presence of 15 mol% 2,6-di-*tert*-butylpyridine.

The last process explored was the 1,3-allylic transposition of 1,1-diphenyl allyl alcohol, also reported to be catalyzed by **B2** (Table 5).^[2d] This reaction proved to be more challenging to rationalize as we faced a major hurdle to reproduce the published results. Using either commercially available or freshly prepared and recrystallized catalyst **B2**, yields never exceeded 20%. We suspect that the reported success of this transformation might result from the presence of an impurity in the way that **B2** was synthesized (a borinic acid or traces of another acid, for instance). Even a trace impurity might not be negligible given the catalyst loading of 20 mol%. In turn, stronger Brønsted acids enabled the reaction to a limited extent (entries 5.5-5.7), but CSA was highly effective, delivering the product in 85% yield (entry 5.8). This result is similar to those reported in the literature, which again strongly suggest Brønsted acid catalysis.

Table 5. Comparison between boronic acids and Brønsted acids for the catalytic 1,3-allylic transposition of allylic alcohols.^[a]

Entry	Catalyst	Yield [%]
5.1	B1	46
5.2	B2	20 (80) ^[b]
5.3	B2	<1 ^[c]

5.4	B3	<1
5.5	TfOH	6
5.6	HCl	16
5.7	H ₂ SO ₄	44
5.8	CSA	85
5.9	CF ₃ CO ₂ H	18
5.10	(COOH) ₂	11
5.11	CH ₃ CO ₂ H	<1
5.12	B(OH) ₃	<1

[a] Yields were determined by ¹H NMR using hexamethyldisiloxane as an external standard. [b] Yield reported in reference 2d. [c] In the presence of 15 mol% 2,6-di-*tert*-butylpyridine.

To summarize, this study sheds light on the activation mode of boronic acid catalysis of alcohols and oximes, showing that Brønsted acid and H-bond catalysis, rather than Lewis acid or covalent activation, are likely responsible for the observed reactivity in nearly all the representative examples studied. Specifically, catalysts **B1** and **B3** produce strong Brønsted acids in the presence of HFIP, and catalyst **B2** likely acts as a H-bond catalyst in MeNO₂. We have concluded the above on the basis of the following key findings: 1) Boronic acids **B1** and **B3** are able to rupture unactivated cyclopropanes in HFIP. 2) The hindered Brønsted base 2,6-di-*tert*-butylpyridine entirely inhibits reactivity of the boronic acids in the reactions investigated. 3) Boronic acid catalyzed reactions of alcohols and oximes can be promoted by Brønsted acids or H-bond catalysts. 4) Brønsted acids that facilitate those transformations exert a deshielding influence on TEPO comparable to the boronic acids that promote the same transformations. 5) For three of the four representative oximes studied, the data collected are inconsistent with a covalent mechanism. Moving forward, these insights should be useful for the rational design of second-generation catalysts for dehydrative nucleophilic substitution or for oxime rearrangements, whether or not they are based on boron. Finally, this work cautions that, before novel catalytic mechanisms are proposed, a wide range of control experiments are necessary to rule out a catalytic role for Brønsted acid and H-bond donors, taking into consideration the important numerous roles played by the solvent.

Acknowledgements

J.M. thanks the European Research Council (ERC) (grant agreement n° 639170), ANR LabEx "Chemistry of Complex Systems" (ANR-10-LABX-0026 CSC) and CNRS. S. Z. thanks the China Scholarship Council for a fellowship. We thank Prof. D. G. Hall for discussion and comments.

Notes

The authors declare no competing financial interest.

Acknowledgements

J.M. thanks the European Research Council (ERC) (grant agreement n° 639170), ANR LabEx "Chemistry of Complex Systems" (ANR-10-LABX-0026 CSC) and CNRS. S. Z. thanks the China Scholarship Council for a fellowship. We thank Prof. D. G. Hall for discussion and comments.

Keywords: boronic acid • hexafluoroisopropanol • Bønsted acid • Lewis acid • dual H-bond

- [1] For reviews on boronic acid catalysis, see: a) E. Dimitrijević, M. S. Taylor, *ACS Catal.* **2013**, *3*, 945-962; b) H. C. Zhang, D. G. Hall, *Aldrichim. Acta.* **2014**, *47*, 41-51; c) D. G. Hall, *Chem. Soc. Rev.* **2019**, *48*, 3475-3496.
- [2] For catalytic activation of alcohols by boronic acids, see: a) J. A. McCubbin, O. V. Krokhnin, *Tetrahedron Lett.* **2010**, *51*, 2447-2449; b) J. A. McCubbin, H. Hosseini, O. V. Krokhnin, *J. Org. Chem.* **2010**, *75*, 959-962; c) J. A. McCubbin, C. Nassar, O. V. Krokhnin, *Synthesis* **2011**, *19*, 3152-3160; d) H. Zheng, M. Lejkowski, D. G. Hall, *Chem. Sci.* **2011**, *2*, 1305-1310; e) J. A. McCubbin, C. Nassar, O. V. Krokhnin, *Synthesis* **2011**, 3152-3160; f) H. C. Zheng, S. Ghanbari, S. Nakamura, D. G. Hall, *Angew. Chem. Int. Ed.* **2012**, *51*, 6187-6190; g) X. B. Mo, J. Yakiwchuk, J. Dansereau, J. A. McCubbin, D. G. Hall, *J. Am. Chem. Soc.* **2015**, *137*, 9694-9703; h) C. L. Ricardo, X. B. Mo, J. A. McCubbin, D. G. Hall, *Chem. Eur. J.* **2015**, *21*, 4218-4223; i) W.-B. Tang, K.-S. Cao, S.-S. Meng, W.-H. Zheng, *Synthesis* **2017**, *49*, 3670-3675; j) H. T. Ang, J. P. G. Rygus, D. G. Hall, *Org. Biomol. Chem.* **2019**, *17*, 6007-6014.
- [3] For catalytic activation of oximes by boronic acids, see: a) S. Chandrasekhar, K. Gopalaiah, *Tetrahedron Lett.* **2002**, *43*, 2455-2457; b) X. B. Mo, T. D. R. Morgan, H. T. Ang, D. G. Hall, *J. Am. Chem. Soc.* **2018**, *140*, 5264-5271.
- [4] For catalytic activation of carboxylic acids by boronic acids, see: a) R. M. Al-Zoubi, O. Marion, D. G. Hall, *Angew. Chem. Int. Ed.* **2008**, *47*, 2876-2879; b) A. Sakakura, T. Ohkubo, R. Yamashita, M. Akakura, K. Ishihara, *Org. Lett.* **2011**, *13*, 892-895; c) N. Gernigon, R. M. Al-Zoubi, D. G. Hall, *J. Org. Chem.* **2012**, *77*, 8386-8400; d) C. Wang, H. Z. Yu, Y. Fu, Q. X. Guo, *Org. Biomol. Chem.* **2013**, *11*, 2140-2146; e) N. Gernigon, H. C. Zheng, D. G. Hall, *Tetrahedron Lett.* **2013**, *54*, 4475-4478; f) K. Ishihara, Y. H. Lu, *Chem. Sci.* **2016**, *7*, 1276-1280; g) K. Wang, Y. H. Lu, K. Ishihara, *Chem. Commun.* **2018**, *54*, 5410-5413; h) S. Arkhipenko, M. T. Sabatini, A. S. Batsanov, V. Karaluka, T. D. Sheppard, H. S. Rzepa, A. Whiting, *Chem. Sci.* **2018**, *9*, 1058-1072.
- [5] a) J. Liu, H. Yao, C. Wang, *ACS Catal.* **2018**, *8*, 9376-9381; b) H. Yao, J. Liu, C. Wang, *Org. Biomol. Chem.* **2019**, *17*, 1901-1905.
- [6] For recent reviews on HFIP, see: a) T. Sugiishi, M. Matsugi, H. Hamamoto, H. Amii, *RSC Adv.* **2015**, *5*, 17269-17282; b) J. Wencel-Delord, F. Colobert, *Org. Chem. Front.* **2016**, *3*, 394-400; c) I. Colomer, A. E. R. Chamberlain, M. B. Haughey, T. J. Donohoe, *Nat. Rev. Chem.* **2017**, *1*, 0088; d) S. K. Sinha, T. Bhattacharya, D. Maiti, *React. Chem. Engl.* **2019**, *4*, 244-253.
- [7] For recent examples of enhancement of reactivity via the use of HFIP, see: a) A. Berkessel, J. A. Adrio, D. Hüttenhain, J. M. Neudörfl, *J. Am. Chem. Soc.* **2006**, *128*, 8421-8426; b) A. Berkessel, J. A. Adrio, *J. Am. Chem. Soc.* **2006**, *128*, 13412-13420; c) H. F. Motiwala, C. Fehl, S.-W. Li, E. Hirt, P. Porubsky, J. Aubé, *J. Am. Chem. Soc.* **2013**, *135*, 9000-9009; d) R. H. Vekariya, J. Aubé, *Org. Lett.* **2016**, *18*, 3534-3537; e) X. Zeng, S. Liu, B. Xu, *Org. Lett.* **2016**, *18*, 4770-4773; f) I. Colomer, R. C. Barcelos, K. E. Christensen, T. J. Donohoe, *Org. Lett.* **2016**, *18*, 5880-5883; g) T. Kamitanaka, K. Morimoto, K. Tsuboshima, D. Koseki, H. Takamuro, T. Dohi, Y. Kita, *Angew. Chem. Int. Ed.* **2016**, *55*, 15535-15538; h) I. Colomer, C. Batchelor-McAuley, B. Odell, T. J. Donohoe, R. G. Compton, *J. Am. Chem. Soc.* **2016**, *138*, 8855-8861; i) D. Lebœuf, L. Marin, B. Michelet, A. Perez-Luna, R. Guillot, E. Schulz, V. Gandon, *Chem. Eur. J.* **2016**, *22*, 16165-16171; j) V. D. Vuković, E. Richmond, E. Wolf, J. Moran, *Angew. Chem. Int. Ed.* **2017**, *56*, 3085-3089; k) C. Qi, F. Hasenmaile, V. Gandon, D. Lebœuf, *ACS Catal.* **2018**, *8*, 1734-1739; l) Z. Tao, K. A. Robb, K. Zhao, S. E. Denmark, *J. Am. Chem. Soc.* **2018**, *140*, 3569-3573; m) E. Richmond, J. Yi, V. D. Vuković, F. Sajadi, C. N. Rowley, J. Moran, *Chem. Sci.* **2018**, *9*, 6411; n) C. Qi, V. Gandon, D. Lebœuf, *Angew. Chem. Int. Ed.* **2018**, *57*, 14245-14249; o) Y. Zhu, I. Colomer, A. L. Thompson, T. J. Donohoe, *J. Am. Chem. Soc.* **2019**, *141*, 6489-6493.
- [8] a) M. Dryzhakov, M. Hellal, E. Wolf, F. C. Falk, J. Moran, *J. Am. Chem. Soc.* **2015**, *137*, 9555-9558; b) M. Dryzhakov, J. Moran, *ACS Catal.* **2016**, *6*, 3670-3673.
- [9] J. G. M. Morton, M. A. Dureen, D. W. Stephan, *Chem. Commun.* **2010**, *46*, 8947-8949.
- [10] For selected papers where transition metal-promoted reactions were later found to be Brønsted acid catalyzed processes, see: a) B. Schlummer, J. F. Hartwig, *Org. Lett.* **2002**, *4*, 1471-1474; b) T. C. Wabnitz, J.-Q. Yu, J. B. Spencer, *Chem. Eur. J.* **2004**, *10*, 484-493; c) D. C. Rosenfeld, S. Shekhar, A. Takemiyu, M. Utsunomiya, J. F. Hartwig, *Org. Lett.* **2006**, *8*, 4179-4182; d) P. N. Liu, Z. Y. Zhou, C. P. Lau, *Chem. Eur. J.* **2007**, *13*, 8610-8619; e) T. Jin, F. Yang, C. Liu, Y. Yamamoto, *Chem. Commun.* **2009**, 3533-3535; f) T. T. Dang, F. Boeck, L. Hintermann, *J. Org. Chem.* **2011**, *76*, 9353-9361; g) H. J. Jin, J. H. Kim, E. J. Kang, *Synthesis* **2017**, *49*, 3137-3144; h) E. T. Sletten, Y.-J. Tu, H. B. Schlegel, H. M. Nguyen, *ACS Catal.* **2019**, *9*, 2110-2123.
- [11] The bulky pyridine 2,6-di-*tert*-butylpyridine (DTBP) is commonly used to distinguish between boron or other Lewis acids and Brønsted acids because it does not complex most Lewis acids, including boron: a) H. C. Brown, B. Kanner, *J. Am. Chem. Soc.* **1966**, *88*, 986-992; b) P. G. Gassman, D. A. Singleton, *J. Am. Chem. Soc.* **1984**, *106*, 7993-7994; c) C. Bergquist, B. M. Bridgewater, C. J. Harlan, J. R. Norton, R. A. Friesner, G. Parkin, *J. Am. Chem. Soc.* **2000**, *122*, 10581-10590; d) M. Yasuda, T. Somyo, A. Baba, *Angew. Chem. Int. Ed.* **2006**, *45*, 793-796; e) M. Vayer, R. Guillot, C. Bour, V. Gandon, *Chem. Eur. J.* **2017**, *23*, 13901-13905.
- [12] a) V. Gutmann, *Coordination Chemistry Reviews* **1976**, *18*, 225-255; b) H. Großekappenberg, M. Reißmann, M. Schmidtman, T. Müller, *Organometallics* **2015**, *34*, 4952-4958; c) M. A. Beckett, D. S. Brassington, S. J. Coles, M. B. Hursthouse, *Inorg. Chem. Commun.* **2000**, *3*, 530-533; d) G. C. Welch, L. Cabrera, P. A. Chase, E. Hollink, J. D. Masuda, P. R. Wei, D. W. Stephan, *Dalton Trans.* **2007**, 3407-3414; e) J. Alarcón-Espósito, R. Contreras, R. A. Tapia, P. R. Campodónico, *Chem. Eur. J.* **2016**, *22*, 13347-13351.
- [13] J. J. Montalvo-Acosta, M. Dryzhakov, E. Richmond, M. Cecchini, J. Moran, *under revisions*.
- [14] a) S. Estopiña-Durán, L. J. Donnelly, E. B. Lclean, B. M. Hockin, A. M. Z. Slawin, J. E. Taylor, *Chem. Eur. J.* **2019**, *25*, 3950-3956; b) E. Wolf, E. Richmond, J. Moran, *Chem. Sci.* **2015**, *6*, 2501-2505.
- [15] T. Beringhelli, D. Maggioni, G. D'Alfonso, *Organometallics* **2001**, *20*, 4927-4938.
- [16] The possibility that the inhibiting effect of DTBP might be due to poisoning of the boronic acid by alkoxide generated from deprotonation of HFIP seems unlikely due to the following experiment: we found the boronic acid catalyzed reaction also occurred in the absence of DTBP in the aprotic solvent DCE under slightly higher temperatures. It is also inhibited by DTBP under these conditions, where no alkoxides are present (see the Supporting Information for more information).
- [17] A. N. J. Moore, D. D. M. Wayner, *Can. J. Chem.* **1999**, *77*, 681-686.
- [18] A. Shokri, X.-B. Wang, S. R. Kass, *J. Am. Chem. Soc.* **2013**, *135*, 9525-9530.
- [19] a) K. M. Diemoz, A. K. Franz, *J. Org. Chem.* **2019**, *84*, 1126-1138; b) G. Lv, X. Sun, C. Zhang, M. Lei, *Atmos. Chem. Phys.* **2019**, *19*, 2833-2844.

Entry for the Table of Contents

To B or not to B: We shed new light on the mechanism of boronic acid catalysis for reactions of alcohols or oximes, which were thought to involve the boron atom acting as a Lewis acid or to feature the transient formation of covalent bonds to boron. The present study suggests they simply act as H-bond donors or Brønsted acids.

Institute and/or researcher Twitter usernames: @djpleboeuf and @MoranLabUdS