Thermochronological, petrographic and geochemical characteristics of the Combia Formation, Amagá basin, Colombia

Matthias Bernet^{1*}, Juliana Mesa Garcia^{2,3}, Catherine Chauvel^{1,4}
and Maria Isabel Marín-Cerón²,

¹Institut des Sciences de la Terre, CNRS, Université Grenoble Alpes, Grenoble, France

²Departemento de Geociencias, Universidad EAFIT, Medellín, Colombia

³present address: Geology Department, University of Michigan, Ann Arbor, MI, USA

⁴Université de Paris, Institut de Physique du Globe de Paris, CNRS, F-75005 Paris, France

*corresponding author, email: matthias.bernet@univ-grenoble-alpes.fr,
ORCID: 0000-0001-5046-7520

Abstract

The Amagá basin between the Western and Central Cordilleras of the Northern Andes of Colombia host the Neogene volcanic and volcanoclastic Combia Formation. At this stage it is not clear how the formation of this unit is related to arc volcanism and which role the Nazca plate subduction beneath the western margin of South America plays. The timing, petrography and geochemical characteristics of Combia Formation rocks were studied in the western and eastern parts of the Amagá basin, in order to gain more information on the type of magma generation and volcanic activity that led to the deposition of the Combia Formation.

Apatite and zircon fission-track dating largely confirm a 12-6 Ma age for the deposition of the Combia Formation. Petrographic and major element analyses show that mainly trachy-andesite ignimbrites with a calc-alkaline composition were deposited in the western Amagá basin, whereas the volcanic rocks of the eastern

Amagá basin are lava flow and fall-out deposits of basaltic andesites or of tholeiitic composition. Trace element and isotopic analyses show that slab dehydration and sediment melting were important in primary magma generation in the mantle wedge, but the primary magma was mixed with lower continental crustal melts, resulting in characteristic isotope signatures in the western and eastern Amagá basin. All this points to subduction driven arc volcanism with slab dehydration, sediment melting magma mixing.

Introduction

The late Paleogene to present-day magmatism of northwestern South America can be divided into four major phases of activity at about 24-20 Ma, 12-6 Ma, 6-3 Ma, and 3 Ma to the Present (e.g. Sierra, 1994; Toro et al., 1999; Gonzalez, 2001; Ramírez et al., 2006; Cediél et al., 2011; Pérez et al., 2013; Lesage et al., 2013). These different magmatic phases are related to the complex tectonic setting in which the Caribbean, Nazca and South American plates interact with each other (Fig. 1). The break-up of the Farallón plate into the Nazca and Cocos plates between 26 and 24 Ma (Marriner and Millward, 1984), the reorientation of subduction direction (Pardo-Casas and Molnar, 1987), and collision of the Panamá-Choco block with northwestern South America at about 25 Ma drove the first magmatic pulse (e.g. McCourt et al., 1984; Aspden et al., 1987; Kellogg and Vega, 1995; Trenkamp et al., 2002; Cediél et al., 2003; Lonsdale, 2005; Restrepo-Moreno et al., 2010; Farris et al., 2011). Second, since the late Paleogene the Nazca plate subduction zone was subjected to changes in subduction angle and direction over time, resulting in Miocene-Pliocene magmatic intrusions in the Western and Central Cordillera and

deposition of the Combia Formation in the Amagá basin (e.g. Pardo-Casas and Molnar, 1987; Taboada et al., 2000; Cediel et al., 2003; Vargas and Mann, 2013). At the same time, subduction of the Caribbean plate beneath the northern (Caribbean) margin of South America caused isolated late Miocene-Pliocene volcanic activity in the Eastern Cordillera (e.g. Vargas and Mann, 2013), such as in the Vetás-California gold-mining district of the Santander Massif (Mantilla et al., 2013), or the Paipa-Iza complex 150 km to the north-east of Bogotá (Fig. 1; Padro et al., 2005; Bernet et al., 2016). Today the main volcanic activity in Colombia is focused on the Central Cordillera with for example the Nevado del Ruiz, Nevado del Tolima, Cerro Machín, Nevado del Huila, Azufral, Cumbal, etc. well to the south of the study area (Fig. 1; e.g. Marín-Cerón et al., 2010, 2019; Leal-Mejía, 2011).

Different techniques have been used for more than a century to understand the genesis, age and evolution of the Combia Formation, including petrography, heavy mineral analysis, X-ray diffraction, geochemistry, geochronology, thermochronology and stratigraphic analyses (e.g. Grosse, 1926; Jaramillo, 1976; Calle and González, 1980; Álvarez, 1983; Marriner and Millward, 1984; Rios and Sierra, 2004; Pérez, 2005; López et al., 2006; Ramírez et al., 2006), but the evolution of the Nazca plate subduction zone magmatism still remains poorly constrained. Here we present a study of a suite of samples collected from three sections, the Cerro Amarillo section in the eastern Amagá basin, and the Anzá-Bolombolo and La Metida Creek sections in the western Amagá basin (Fig. 2), in order to improve the knowledge gained so far about the Combia Formation. The volcanoclastic, tuff/lapilli and flow deposits of the Combia Formation were examined with a) apatite fission-track (AFT) and zircon fission-track (ZFT) thermochronology, b) petrographic analyses, and c) major and trace element analysis, as well as Sr, Nd

and Pb isotope analyses. All this was done with the objective of a) characterizing and comparing the eastern and western volcanic deposits, and b) to better understand the mid-late Miocene evolution of the Nazca subduction zone magmatism manifested between the Western and Central Cordilleras.

Geological setting

The Northern Andes of northwestern South America consist in Colombia of the Western, Central and Eastern Cordilleras (Fig. 1). Each of these mountain belts reflects a particular part of the long-term evolution of the Northern Andes, which is characterized by magmatic episodes since the Precambrian, during the Triassic, Jurassic, Late Cretaceous, and since the late Paleogene/Neogene until today (Aspden et al., 1987). In general, these magmatic phases have been related to Farallón/Nazca plate subduction beneath the western margin of the South American plate (e.g. Marriner and Millward, 1984; McCourt et al., 1984; Cediél et al., 2003; Saenz, 2003; Restrepo-Moreno et al., 2009; Rodríguez et al., 2012). Accretion of tectonic blocks or terranes of oceanic affinity to the continental margin during the late Mesozoic and early Cenozoic did not cause Andean-type subduction volcanism, because of their relatively young age and high buoyancy preventing subduction (Cediél et al., 2003), and forcing surface uplift and the formation of the Western and Central Cordilleras during the Pre-Andean and Andean orogenies (e.g. Van der Hammen, 1960; Taboada et al., 2000; Cediél et al., 2003).

The present-day Andean volcanism is commonly divided into four volcanic zones, the Northern Volcanic Zone (NVZ), Central Volcanic Zone (CVZ), Southern Volcanic Zone (SVZ), and Austral Volcanic Zone (AVZ) (e.g. Thorpe and Francis, 1979; Thorpe et al., 1982; Stern, 2004; Marín-Cerón et al., 2019). These segments

have been distinguished based on differences in petrographic features and geochemical signatures, and they are separated from each other by volcanic gaps (e.g. Thorpe and Francis, 1979; Stern, 2004). The NVZ is located in north-western South America and encompasses the region of present-day volcanism in the Northern Andes of Ecuador and Colombia.

Geology of the Amagá basin

The Amagá basin forms the northern part of the much larger Amagá-Cauca-Patía basin located between the Western and Central Cordilleras of the Northern Andes in western Colombia (Fig. 1; Sierra and Marín-Cerón, 2011). Dextral strike-slip faulting along the Cauca and Romeral fault systems to the west and east respectively is responsible for development of the Amagá basin, which is tectonically a pull – apart basin (e.g. Cediel et al., 2003). Basin evolution started possibly during the Eocene (?) – Oligocene, with surface uplift and erosion of the Central Cordillera from the Late Cretaceous to Eocene and deposition of clastic sediments of the Lower Amagá Formation in the basin (e.g. Restrepo-Moreno et al., 2009). The Lower Amagá Formation is known for its quartz-rich sandstones and mainly sub-bituminous but locally anthracite grade coal (Silva et al., 2008; Bandon et al., 2008). The Lower Amagá Formation is separated from the Oligocene to Miocene Upper Amagá Formation by an unconformity and a change to a lithic arenite composition with sedimentary and metamorphic lithoclasts derived from the Central Cordillera (Paez Acuna, 2012). During the mid to late Miocene, subduction of the Nazca plate below the South American plate allowed the genesis of the Combia Formation in the Amagá basin (e.g. Grosse, 1926; Marriner and Millward, 1984; González, 2001; Cediel et al., 2003; Ramírez et al., 2006; Leal-Mejía, 2011; Cediel et al., 2011).

Therefore, the Upper Amagá Formation is overlain by volcanic and volcanoclastic deposits of the Combia Formation. Here we focus on the Cerro Amarillo section in the Eastern Amagá basin and the Anzá – Bolombolo and Le Metida Creek sections in the western Amagá basin.

The Cerro Amarillo section

This section is located between the towns of Damasco and La Pintada (Fig. 2). It has a total thickness of 193 m and comprises 34 layers of welded tuff, pyroclastic and agglomerate breccia, lapilli tuff breccia, basalt and scoria (Fig. 3). The layers vary in thickness from a 20 m pyroclastic and agglomerate breccia to 0.3 m lapilli tuff breccia, both at the top (Mesa-Garcia, 2015). There is also a 19.2 m thick welded tuff at the bottom. However, it is most common to find layers of 1 – 7 m in thickness. The layers are characterized by a tabular geometry. No evidence of pinchout or lenses were observed in the outcrops. The bottom of the stratigraphic succession is mainly characterized by lava flows and welded tuff while, the top of the sequence mainly consists in coarse to very coarse grained pyroclastic flows. The bottom layers have a strike and dip of S05°E/25°SW in average; towards the middle of the section the layers strike and dip N70°E/18°SE. Finally the top breccia and lapilli tuff layers strike and dip N15°W/19SW. Many basalt and welded tuffs layers have randomly distributed vesicles and amygdulites of variable size shapes, some are elongated indicating lava flow directions.

The Anzá – Bolombolo section

This section is also located on the western bank of the Cauca River (Fig. 2) between the villages of Anzá and Bolombolo. The stratigraphy of the 11.04 m thick

succession consists of tuffs and lapilli tuffs in the lower part of the section, which are separated laterally from andesitic basalts and ash flow deposits by an erosional unconformity (Fig. 4; Grosse, 1926; González, 2001; Sierra and Marín-Cerón, 2011). In the upper part of the succession are a lapilli tuff breccia and a non-differentiated lava flow, which cover the underlying units and the unconformity. No particular sedimentary structures were observed at this location.

The La Metida Creek section

This section is located on the western bank of the Cauca River to the west of Bolombolo (Fig. 2). The Combia Formation crops out along the stream bed (e.g. González, 2001) and the exposed stratigraphic sequence has a thickness of 45 m and is composed of 33 layers of tuff, lapilli tuff, lapilli tuff breccia, pyroclastic flows and volcano-clastic sandstones (Fig. 5). Sedimentary structures found throughout the sequence include lenticular bedding, load structures, ripple lamination and cross-bedding. The layers at the bottom are mainly grain-supported, whereas the top layers are matrix-supported. In addition, nodules and organic matter are commonly observed at the bottom of the section, and the middle section is characterized by bioturbation and fossilized plants (e.g. leaves). No such material was observed towards the top of the section.

Analytical methods

Apatite and zircon fission-track analyses

AFT analysis was done on two samples from the AB section and six samples from the MC section. In addition, ten samples of the MC section were analyzed with

the ZFT method. Unfortunately, the apatite and zircon yield of the CA section samples was too low for fission-track analyses.

Sample preparation and analyses were performed at the thermochronology laboratory of the Universidad EAFIT at Medellín and the ISTerre thermochronology laboratory at the Université Grenoble Alpes. Rock samples were crushed and sieved and heavy mineral fractions were separated using standard hydraulic, magnetic and heavy liquid separation techniques. The apatite crystals were mounted in epoxy and the zircon crystals in Teflon® sheets, polished and etched. Apatite grains were etched for 20 seconds at 21°C with 5.5 mol HNO₃, and zircons were etched at 228°C for 10-40 h in a NaOH-KOH melt to reveal fission tracks. A white mica sheet was mounted as the external detector. All samples were irradiated with thermal neutrons at the FRM II reactor in Garching, Germany, with a nominal fluence of 8×10^{15} n/cm² for apatite and 0.5×10^{15} n/cm² for zircon, together with IRMM540R dosimeter glasses and Durango age standards for apatite, and IRMM541 dosimeter glasses and Buluk and Fish Canyon Tuff age standards for zircon. After irradiation external detectors were etched in 48% HF for 18 min at 21°C. Fission tracks were counted dry at 1250x using an Olympus BX51 microscope and the FTStage 4.04 system. Fission-track ages for each sample were calculated using the Binomfit software of Brandon (see Ehlers et al., 2005) and the RadialPlotter program of Vermeesch (2009).

Petrographic analyses

Petrographic analyses were performed on twenty samples, seven from the Cerro Amarillo section, three from the Anzá-Bolombolo section, and ten from the La Metida Creek section. Petrographic thin sections were prepared at Geoensayos

S.A.S., Medellín, Colombia. Some of the pyroclastic samples had to be impregnated with epoxy, as these deposits were not well consolidated.

The samples were analyzed using an Olympus BX41TF petrographic microscope at the Geology department of EAFIT University. Modal analysis was performed counting 300 – 500 points per sample, using a point counter. The description of mineral assemblages and textures was done according to Mackenzie et al. (1984) and Ehlers (1987). The samples were classified using modal mineral composition and the QAP diagram of Streckeisen (1976) for volcanic rocks and based on the size of the material after Pettijohn (1975) for pyroclastic rocks.

Geochemical analyses

The rock samples collected in the field were crushed to 1 mm chips at the Laboratory of Solid Materials at EAFIT University. Two hundred grams per sample were separated thoroughly, choosing the rock chips that were the least weathered and geochemical analyses for major and trace elements were performed in the clean laboratory at the Institut des Sciences de la Terre (ISTerre) – Université Grenoble Alpes, France. Samples were finely powdered in an agate mortar previous to analyses, except for Pb isotope analysis for which rock chips were used directly. The sample preparation and analytical procedures were executed according to Chauvel et al. (2011), as summarized below.

For major elements, 50 mg of powdered sample were dissolved in 800 µl of concentrated HNO₃ and 15 drops of concentrated HF and heated in a Savillex beaker for two days at 90°C on a hot plate. After cooling, 20 ml of H₃BO₃ (25 g/l) were added to the solution to neutralize excess HF, 10 g of HNO₃ and 250 ml of milliQ water for further dilution. Five standards (BR 24, BEN, BHVO2, AGV-1 and

BCR–1), a duplicate and a blank were as well prepared for analysis. The solutions were analyzed using Inductively Coupled Plasma Atomic Emission Spectroscopy (ICP AES) at ISTerre to determine the major element composition of each sample. Concentrations were obtained using the international rock standard BR to calibrate the signal and the values recommended by Chauvel et al. (2011). Loss on ignition (LOI) was calculated for all samples by heating one gram of sample at 1000°C for one hour.

Sixteen samples, five standards (BR 24, BEN, BHVO2, AGV–1 and BCR–1), three duplicates and one blank were analyzed for trace element contents. One hundred milligrams of samples and standards were dissolved in 20 drops of concentrated HNO₃ and 3 ml of HF for three days at 120°C in Savillex beakers. Samples were further taken up in 7N HNO₃ and finally dissolved in 2% HNO₃ with traces of HF, to obtain a dilution factor of about 5000 (Chauvel et al., 2011). The sample was analyzed using an Inductively Coupled Plasma Mass Spectrometry (ICP MS) Agilent 7500 at ISTerre.

Ten samples were prepared for Nd, Pb and Sr isotope ratios analyses. Two sets of Sr samples, unleached and leached, were prepared following the same procedures as for the Nd samples and Pb samples, respectively. The samples were dissolved and conditioned in Savillex beakers. Rock chips were leached according to McDonough and Chauvel (1991), to eliminate as much as possible Sr and Pb superficial contamination. For Nd and unleached Sr isotopic measurements, 100 mg of sample were dissolved using HNO₃ and HF and isolated using the same procedure as Chauvel et al. (2011). For Pb and leached Sr, 1 g of rock chips was leached prior to dissolution using HCl and then isolated as in Chauvel et al. (2011). Nd and Pb isotopic ratios were measured using a Nu Plasma HR MC–ICPMS at

ENS Lyon, France, while Sr isotopic ratios were measured using a Thermo Scientist Triton MS at the University of Brest, France.

Results

Fission-track results

Anzá – Bolombolo section

The tuff layer at the bottom of the Anzá-Bolombolo section (sample JJ22) has an AFT central age of 8.4 ± 3.1 Ma, and the pyroclastic flow sampled towards the top of the section (sample JJ13) has an AFT central age of 7.9 ± 1 Ma (Table 1; see the supplementary data for individual grain ages and radial plots of all samples). The combined Combia Formation AFT data are shown in a radial plot in Figure 6A with a central age of 8.6 ± 1.4 Ma based on 302 grain ages.

La Metida Creek section

In total six samples were analyzed with the AFT method and ten with the ZFT method. AFT central ages range between 15.9 ± 11.1 and 5.1 ± 2.5 Ma. Nonetheless, inherited single grains with apparent cooling ages of between 374 and 49 Ma can also be observed in all samples (Table 1). Zircon crystals were mainly found in the middle and bottom of the La Metida Creek section. The euhedral to subhedral zircon crystals range in color from colorless, yellow, pink to red. The ZFT central ages are between 12.7 ± 2.4 Ma and 6.1 ± 1.1 Ma (Table 2). Except for the volcanoclastic sandstone sample (JJ6), no strong evidence exist of contamination with inherited zircons derived from surrounding country rock (see the supplementary data for individual grain ages and radial plots of all samples). The combined Combia

Formation ZFT data are shown in Figure 6B with a central age of 9.1 ± 1.1 Ma based on 346 grain ages.

Petrographic results

Cerro Amarillo section

In general, the seven samples analyzed for this section share similar characteristics being hypocrystalline, porphyritic rocks. The mineral assemblage is represented by plagioclase + pyroxene $\pm$ amphibole $\pm$ olivine (Table 3). Other minerals present in the samples are secondary calcite, biotite and opaque minerals. The rock fragments found in the ignimbrites are mainly basalts (aphanitic textures and volcanic glass) or andesites (plagioclase crystals in volcanic glass). Some of these fragments are fractured. There is also evidence of oxidation and sericitization, even though carbonate minerals are the main alteration product in these samples.

The ignimbrites are dominated by crystals and rock fragments ranging from 55 – 70 %, whereas the two basalt samples have different percentage relationships between the matrix and phenocrysts, CA – 14 rich in crystals and CA – 18 rich in matrix. The crystal and/or rock fragments are in-equigranular, which is observed in the presence of seriate, glomeroporphyritic, poikilitic, and interstitial textures. Overgrowth textures such as skeletal, corona and crystal zoning are also found in the samples, mainly in the ignimbrites. The crystals have subhedral to anhedral shapes in most samples; euhedral-shaped crystals are rare or unable to identify due to alteration. The matrix is generally altered to secondary calcite, but in some cases a non-altered volcanic glass composition is observed with some embedded microlites of plagioclase and less common pyroxene.

Based on the lack of quartz and potassic feldspar, all samples from the Cerro Amarillo section plot in the basalt/andesite field of the classification diagram of Streckeisen (1976) shown in Figure 7A. Color plates of thin section photographs are given in the supplementary data archive.

Anzá – Bolombolo section

The analyzed samples are hypocrystalline, un-equigranular volcanic rocks. The mineral assemblages are mainly plagioclase + pyroxene (Table 3). The matrix consists of volcanic glass. Other rock components are spherulites, found in samples AB–6 and AB–7, and olivine, found in sample AB–6. These samples are located to the east of the unconformity described in the stratigraphic section. The samples are characterized by having skeletal and crystal zoning overgrowth, spherulites and vesicular textures. The lava flows can be classified as basalt/andesite in the Streckeisen (1976) diagram (Fig. 7a), and the pyroclastic flow deposits as crystal tuffs after Pettijohn (1975), as shown in Figure 7b. Color plates of thin section photographs are given in the supplementary data.

La Metida Creek section

Nine samples from this section mainly plot in the crystal tuff field of the Pettijohn (1975) classification (Fig. 7b). Only one sample (QML – 9) is defined as tuffaceous sandstone because of the presence of cross-bedding in the layer. Petrographic characteristics of all samples are similar and the classification as pyroclastic and epiclastic rocks is based on sedimentary structures observed in the field. The mineral assemblage is mainly plagioclase + pyroxene. Some amphiboles are present at the bottom layers.

The sampled rocks are mainly hypocrystalline due to the presence of phenocryst, and microlites in the matrix. The minerals have un-equigranular and seriate fabrics. In general, skeletal textures are found in all analyzed samples. Other common textures are glomero-porphyritic and crystal zoning, although the latter is not present in the samples analyzed from the middle section of the La Metida Creek section. Similar to the Anzá-Bolombolo section spherulites are found in the samples from the bottom layers. Color plates of thin section photographs are given in the supplementary data.

Geochemical results

Mayor element analyses of 16 samples are presented in Table 4. Loss on ignition (LOI) is generally below 3%. For the Cerro Amarillo section samples, SiO₂ contents range between 51 and 53 wt%; they plot in the tholeiitic field of the AFM diagram (Fig. 8A) and correspond to basaltic andesite according to Figures 8B. Samples coming from the Anza-Bolombolo area have slightly higher SiO₂ content at about 56 wt%; they plot in the alkaline field in Fig. 8A and correspond to trachy-andesites in Fig. 8B.

Trace element contents of all samples are given in Table 5 and plotted in primitive mantle-normalized spider diagrams in Fig. 9A. The Cerro Amarillo samples display an enrichment in large-ion lithophile elements (Rb, Ba, Cs, Sr) and a strong depletion in Nb and Ta. The Anzá-Bolombolo samples are even more enriched in Rb, Ba, Cs and Sr, and with similar depletion in Nb and Ta (Fig. 9A). The main difference between the Cerro Amarillo and Anzá-Bolombolo samples is the Li enrichment of the Anzá-Bolombolo section samples. Samples from both sections are enriched in Light Rare Earth Elements (LREE) relative to the Heavy Rare Earth

Elements (HREE) (Fig. 9B) with a stronger fractionation in the Anzá-Bolombolo section samples. No significant Eu anomaly exists for all samples.

Nd, Sr, and Pb isotope compositions

Nd, Sr, and Pb isotope analyses of all samples are given in Table 6. Strontium isotopic ratios measured on leached and unleached samples are similar within errors. $^{87}\text{Sr}/^{86}\text{Sr}$ ratios range from 0.703862 to 0.703931 for the Cerro Amarillo section samples, but are higher at about 0.70417 for the Anzá-Bolombolo section samples (Figure 10A). $^{143}\text{Nd}/^{144}\text{Nd}$ ratios vary between 0.51292 and 0.51298 for the Cerro Amarillo section samples, and are somewhat lower at 0.51290 for the AB section samples (Figure 10B and C). Finally, the Cerro Amarillo section samples define a small range in Pb isotopic ratios ($^{208}\text{Pb}/^{204}\text{Pb}$: 38.68 to 38.80, $^{207}\text{Pb}/^{204}\text{Pb}$: 15.61 to 15.62 and $^{206}\text{Pb}/^{204}\text{Pb}$: 18.91 to 19.07), and the Anzá Bolombolo section samples fall in the middle of the range (Figure 11a and b)

Discussion

Constraints from low-temperature thermochronology

The AFT and ZFT data presented in this study can in principle be used for constraining the age of deposition of the volcanic and volcanoclastic deposits of the Combia Formation (Kowallis et al., 1986; Bernet et al., 2016). The AFT and ZFT data central age values ranging from 15.9 – 5.1 Ma for AFT and 12.7 – 6.1 Ma for ZFT, bracket the known 12-6 Ma age of Combia Formation volcanic activity (e.g. Leal-Mejía, 2011). Our fission-track data highlight two important aspects that are

challenging in dating relatively young volcanic and volcanoclastic deposits with the fission-track method. As to be expected the ZFT data correspond more closely to the known 12-6 Ma age range determined from zircon U-Pb analyses (Leal-Mejía, 2011), because of the higher U concentration and the better track counting statistics resulting in higher precision results. The AFT data suffer under the very low (in general <10 ppm) U concentration of the apatites in the Combia Formation. As can be seen in the single grain data provided in the supplementary data archive, many apatite grains are zero-track grains, resulting in a very high single grain age uncertainty, as the induced track counts also tend to be because of the low U concentrations.

In addition, dealing with volcanic and particularly volcanoclastic deposits the risk of contamination with apatites and zircons recycled from the country rock is high, and has been shown to be the case for certain volcanic deposits of the Paipa-Iza volcanic complex (Bernet et al., 2016). Here we also think that zircons with >12 Ma apparent cooling ages were most likely recycled from the Amagá Formation (Piedrahita et al., 2017). Pre-Miocene cooling ages are common in apatites of the Combia Formation deposits (Table 1), and are considered to be derived from underlying basement rock and recycling of the Amagá Formation.

In summary, based on the AFT and ZFT data, volcanic activity occurred between 12 Ma and 6 Ma. The main phase of activity for these deposits may have been at around 9 Ma, as suggested by the central ages of the combined AFT and ZFT data sets shown in Figure 6. This confirms a late Miocene depositional age, which has previously been proposed based on stratigraphic position and whole rock K-Ar dating of hypabyssal porphyries (e.g. Grosse, 1926; Restrepo et al., 1981;

Marriner and Millward, 1984; González, 2001, Pérez, 2005; Ramírez et al., 2006; Leal-Mejía, 2011).

Shallow-level processes prior to eruption

The results presented in this study indicate that there are two different petrographic trends among the Combia Formation rocks. In general, the rocks have porphyritic textures, and plagioclase is one of the main mineral components, both as phenocrysts and microlites. Similar results have been reported by Marriner and Millward (1984), López et al. (2006) and Ramírez et al. (2006).

The matrix for most samples is comprised of volcanic glass with microlites of plagioclase and to a lesser extent pyroxene. Devitrification of the volcanic glass matrix is common in the samples. The samples show evidence of alteration (e.g. secondary calcite, oxidation, argillitization) which may indicate metasomatic to surficial processes related to hydrothermal alterations and mineralization processes in the hypabyssal porphyries of the study area (e.g. Tassinari et al., 2008; Leal-Mejía, 2011; Lesage et al., 2013; Uribe-Mogollón, 2013).

All samples show disequilibrium textures such as skeletal, sieve and spherulitic textures, embayments, reaction rims, coronas and crystal zoning. These textures are attributed to different conditions and magmatic processes such as pressure variations, zoned magma chambers, decompression, magma mixing, phenocrysts recycling and fractional crystallization (e.g. Nixon and Pearce, 1987; Nelson and Montana, 1992; Singer et al., 1995; Perugini et al., 2003; Aldanmaz, 2006; Maro and Remesal, 2011). Thus, our petrographic results favor mainly magma mixing and fractional crystallization to be the main causes for disequilibrium in the magma chamber.

Magma genesis

Magmatic arcs are the result of subduction of oceanic crust beneath continental crust and this complex setting has impact on magmatic processes such as partial melting, fractional crystallization, changes in pressure and temperature, sediment melting, dehydration, and decarbonation (e.g. Rollinson, 1993; Albarade, 1995; White, 2013). In addition, fluids added to magma play an important role, as fluids allow the transport of incompatible elements from the subducted oceanic basalt and sediments to the magma, leading to enrichment in specific mobile elements and modifying the isotopic composition of the asthenospheric magma source (Tatsumi, 2005; Tatsumi and Stern, 2006; Tatsumi and Takahashi, 2006; Nakamura et al., 1985).

Our new geochemical data for the Cerro Amarillo area are generally similar to what was previously published by Marriner and Millward (1984) and by Ordoñez (2001). Major elements allow identification of a tholeiitic trend in the samples from the eastern Amagá basin while the samples from the western Amagá basin follow a calc-alkaline trend (Fig. 8A). Both magma suites have been previously recognized at other sampling sites of the Combia Formation (Fig. 8A; e.g. Álvarez, 1983; Marriner and Millward, 1984; Ordoñez, 2001; Leal-Mejía, 2011). With SiO₂ contents between 47 – 59 wt% our Cerro Amarillo samples plot in the basaltic andesite field of Fig. 8B and the samples from Anzá Bolombolo plot in the trachy-andesite field of LeMaitre et al. (1989) and Cox et al. (1979) (Fig. 8B). For comparison, published data for the Combia Formation are also shown in this figure. It appears clearly that the AB section samples from the western Amagá basin are much more alkaline than others (Fig. 8B), and that samples from both sections have lower SiO₂ contents than the

volcanic rocks of the 24-20 Ma magmatic phase and most of the 17-9 Ma magmatic phase rocks analyzed by Leal-Mejía (2011). Despite these differences, so far no precise division of the basin with respect to the magmatic suites can be given at the moment, and more detailed mapping and geochemical analyses are necessary.

The trace elements provide clear evidence of a subduction-related geochemical signature, with Nb and Ta depletion (e.g. Wilson, 1989; White, 2013). Samples also all have very high Ba, U, Pb and Sr contents as already noticed for older samples from the Combia Formation (Fig. 9C, Leal-Mejía, 2011 data summarized in Marín-Cerón et al., 2019). Medium to slightly elevated Ba/Th values (a proxy for slab dehydration Fig. 9E; Labanieh et al., 2012) characterize the Cerro Amarillo samples from the eastern Amagá basin, whereas the AB samples of the western Amagá basin have low ratio consistent with a sediment melting trend.

In general, the trace element results of the Amagá basin resemble those of the northern volcanic zone (NVZ) of the Andes, as summarized in Marín-Cerón et al. (2019), even though the Combia Formation has higher LILE (Rb, Sr, Ba) contents and lower Nb and Ta contents than those reported for the NVZ (Thorpe et al., 1982; Marriner and Millward, 1984).

The REE patterns of samples from the Anza Bolombolo and Cerro Amarillo sections are somewhat different, with higher fractionation in the former group (Fig. 9b). In comparison to the REE spectra of the 17-6 Ma porphyritic intrusions and recent volcanism, the Combia Formation shows a pattern closer to the latest stage of volcanic activity (Fig. 9D; data from Leal-Mejía (2011), summarized by Marín-Cerón et al. (2019)). The difference in the slope of the REE patterns of eastern and western Amagá basin samples suggest differences in terms of magma formation, most probably during melting processes in the mantle source. Overall, the absence of Eu

anomaly suggests that melting occurred relatively deep, below the plagioclase stability level (~40 km depth).

Magma source composition

Combining trace element data with isotopic data can help understanding the origin of magmas and the potential role of subducted slab addition to the mantle wedge (e.g. Tatsumi, 2005; Tatsumi and Stern, 2006). The Cerro Amarillo and Anzá-Bolombolo samples have medium to slightly elevated Sr/Th and rather low $^{87}\text{Sr}/^{86}\text{Sr}$ ratios (Fig. 10A), indicating transfer of elements from the subducted slab by NVZ aqueous fluids. Because no systematic difference in $^{87}\text{Sr}/^{86}\text{Sr}$ is observed between leached and unleached samples (Table 6), post-emplacement alteration can be excluded, and the measured ratios can be considered as representative of the magma source (e.g. Tamura and Nakamura, 1996; Shibata and Nakamura, 1997). Similarly, the large range of Ba/Th at almost constant $^{143}\text{Nd}/^{144}\text{Nd}$ indicates addition of Ba through slab dehydration (Fig. 10B), with a stronger effect in the eastern Amagá basin than in the western part of the basin. In a $^{87}\text{Sr}/^{86}\text{Sr}$ vs $^{143}\text{Nd}/^{144}\text{Nd}$ isotopic space (Fig. 10C), the Amaga basin samples fall as expected within the field defined by the NVZ (see compilation of Marín-Cerón et al., 2019). It is worth noting that the Amaga Basin samples correspond to the most depleted part of the isotopic range observed along the Andes, suggesting limited crustal contamination during magma ascent and/or limited contribution of subducted sedimentary material to the mantle wedge. It is also worth noting that the Anzá-Bolombolo samples have slightly more enriched isotopic characteristics than the Cerro Amarillo samples, a feature consistent with the difference seen in Figure 9E since sediment addition tends to lower the Nd isotopic composition and increase the Sr isotopes.

Lead isotopes provide complementary and useful information. As was the case for Sr and Nd isotopic data, the Amaga basin samples fall in the field defined by the northern volcanic zone and differ drastically from fields defined by central and southern volcanic zones (Fig. 11A) (see compilations made by Marín-Cerón, 2007 and Marín-Cerón et al., 2010; 2019). The rather radiogenic values of Pb isotopes for the northern volcanic zone have been interpreted as being due either to an enriched mantle reservoir or to continental crust assimilation. The enriched mantle hypothesis was suggested by Rodríguez-Vargas et al. (2005) on the basis of the Nd and Sr isotopic characteristics of xenoliths from the Mercaderes region in SW Colombia. The authors invoked the potential involvement of magma material coming from the Galapagos plume, but such influence under the Amaga basin seems quite unlikely given the distance between the SW Colombian arc and the Galapagos (>450 km, Pedersen and Furnes, 2001). However, the Amaga Basin samples analyzed in this study provide new information because they define a tight correlation in $^{208}\text{Pb}/^{204}\text{Pb}$ vs $^{206}\text{Pb}/^{204}\text{Pb}$ space (see Fig 11B). Such linear array implies the involvement and mixture of two endmembers whose compositions remain unchanged during the entire volcanic sequence. The enriched end-member seems to correspond to the local lower continental crust (see Fig. 11B) while the less radiogenic endmember is more ambiguous. Following Marín-Cerón's model (2019), this 'depleted' endmember could correspond to the mantle wedge whose composition would be affected by the presence of material originating from the subducted slab (see Fig. 11B). Nonetheless, there might be differences in the subduction mechanisms, proportion of end-members interaction, and magma source evolution between magmatism at 12 – 6 Ma and magmatism at 3 Ma – present, despite assuming the same end-members. In summary, magma was generated by slab dehydration, sediment melting and

interactions with the LCC, resulting in the mixing of at least two end-member sources. Due to differences in depth of melting and other magmatic processes (AFM and MASH), as shown in Figure 12, the late Miocene volcanic rocks of the western and eastern Amagá basin show distinct petrologic and geochemical signatures.

Conclusions

The late Miocene volcanic and volcanoclastic Combia Formation of the Amagá basin between in the Central and Western Cordillera developed above the Nazca plate subduction zone in western Colombia. Volcanic and volcanoclastic rocks of the Combia Formation are characterized by a sequence of ignimbrites and lava flows of tholeiitic affinity at the bottom and pyroclastic flows to the top of the formation in the eastern Amaga basin and a succession of pyroclastic and epiclastic flows with calc-alkaline affinity in the western Amagá basin. Using apatite and zircon fission-track dating the timing of volcanic activity during the deposition of the Combia Formation was confirmed between 12 and 6 Ma.

Trace element, REE and Nd, Sr and Pb isotopic analyses show that the surface weathering did not modify the geochemical signatures and that the geochemical composition of the samples results of several magmatic processes including slab dehydration and sediment melting to form the primary magma in the mantle wedge and mixing of this primary magma with lower continental crustal before eruption. In contrast, contamination by upper crustal rocks could not be detected. Finally, the new geochemical results confirm that volcanism in the Combia area between 12 and 6 Ma was similar to what is known about the NVZ of the Andes in South America.

552

553 **Acknowledgements**

554 We acknowledge support of this study from ECOS-NORD/Colciencias/ICETEX
555 project C12U01 of M. Bernet and M.I. Marín-Cerón, as well as a BQR SUD grant at
556 ISTERre, awarded to M. Bernet. We thank Wilton, Francois Senebier and Francis
557 Coeur for help with sample preparation and mineral separation.

558

559

References

- Albarade, F. 1995. Introduction to geochemical modelling. Cambridge University Press. 543 pp.
- Aldanmaz, E. 2006. Mineral – chemical constraints in the Miocene calc-alkaline and shoshonitic volcanic rocks of Western Turkey: Disequilibrium phenocryst assemblages as indicators of magma storage and mixing conditions. Turkish Journal of Earth Sciences. 15: 47 – 73.
- Álvarez, A. 1983. Geología de la cordillera Central y el Occidente colombiano y petroquímica de los intrusivos granitoides Mesocenoicoicos. Boletín Geológico. 26: 175 pp.
- Aspden, J.A., McCourt, W.J., and Brook, M., 1987. Geometrical control of subduction-related magmatism: the Mesozoic and Cenozoic plutonic history of Western Colombia. Journal of the Geological Society, London. 144: 893 – 905.
- Barret, T.J. 1983. Strontium- and lead-isotope composition of some basalts from Deep Sea Drilling Project Hole 504B, Costa Rica Rift, Legs 69 and 70. In Cann, J.R., Langseth, M.G., Honnorez, J., VonHerzen, R.P., White, S.M., et al., Init. Repts. DSDP, 69. Washington (U.S. Govt. Printing Office). 643 – 650.
- Barret T. J., Taylor P. N., Lugowski J. 1987. Metalliferous sediments from DSDP leg 92: the East Pacific Rise transects. Geochim. Cosmochim. Acta. 46:651 – 666.
- Bernet, M., Uruena, C., Amaya, S.,Pena, M.L., 2016. New thermo- and geochronological constraints on the Pliocene-Pleistocene eruption history of the Paipa-Iza volcanic complex, Eastern Cordillera, Colombia, Journal of Volcanology and Geothermal Research , 327:299-309.

583 Bandon, A., Parra, N., Gorin, G.E., Arango, F. 2008. Adapting palynological
 584 preparation methods in subbituminous and bituminous coals from Colombia to
 585 improve palynofacies and hydrocarbon source rock evaluations. *International Journal*
 586 *of Coal Geology*, 73:99–114.

587 Calle, B., González, H. 1980. *Geología y Geoquímica de la Plancha 166, Jericó*.
 588 Informe no. 1822. Medellín. INGEOMINAS. 232 pp.

589 Cediél, F., Shaw, R.P., Caceres, C. 2003. Tectonic assembly of the Northern
 590 Andean Block, in C. Bartolini, R. T. Buffler, and J. Blickwede, eds., *The Circum-Gulf*
 591 *of Mexico and the Caribbean: Hydrocarbon habitats, basin formation, and plate*
 592 *tectonics*.- AAPG Memoir 79: 815-848.

593 Cediél, F., Leal-Mejía, H., Shaw, R.P., Melgarego, J.C., Restrepo-Pace, P.A. 2011.
 594 *Petroleum Geology of Colombia: Regional Geology of Colombia*. ANH – Colombia.
 595 1: 220 pp.

596 Chauvel, C., Bureau, S., Poggi, C. 2011. Comprehensive Chemical and Isotopic
 597 Analyses of Basalt and Sediment Reference Materials. *Geostandards and*
 598 *geoanalytical research*. 35 (1): 125 – 143.

599 Chiaradia, M., Fontbote, L. 2002. Lead isotope systematics of Late Cretaceous
 600 Tertiary Andean arc magmas and associated ores between 8°N and 40°S: evidence
 601 for latitudinal mantle heterogeneity beneath the Andes. *Terra Nova*. 14 (5): 337 –
 602 342.

603 Cox, K.G., Bell, J.D., Pankhurst, R.J. 1979. *The Interpretation of Igneous Rocks*:
 604 London, George Allen & Unwin. 464 pp.

605 Dasch, E.J. 1981. Lead isotopic composition of metalliferous sediments from the
 606 Nazca Plate. *Mem. Geol. Soc. Am.* 154: 199 – 209.

607 Davidson, J.P., de Silva, S.L. 1992. Volcanic rocks from the Bolivian Altiplano:
 608 Insights into crustal structure, contamination, and magma genesis in the central
 609 Andes. *Geology*. 20: 1127 – 1130.

610 de Silva, S. 1991. Styles of zoning in Central Andean ignimbrites-insights into
 611 magma chamber processes: In: *Andean magmatism and its tectonic setting*. Special
 612 paper 265: 217 – 232.

613 DePaolo, D. J. (1981). Trace-element and isotopic effects of combined wallrock
 614 assimilation and fractional crystallisation. *Earth and Planetary Science Letters* 53,
 615 189–202.

616 Ehlers, E.G. 1987. *Optical Mineralogy: Vol. 2. Mineral Descriptions*. Blackwell
 617 Scientific Pub. Palo Alto. 286 pp.

618 Ehlers, T.A., Chaudhri, T., Kumar, S., Fuller, C., Willett, S.D., Ketcham, R., Brandon,
 619 M.T. 2005. Computational tools for low-temperature thermochronometer
 620 interpretation. In: Reiners, P.W., Ehlers, T.A. (Eds.), *Low-temperature*
 621 *Thermochronology. Techniques, Interpretations and Applications*. *Reviews in*
 622 *Mineralogy and Geochemistry*, Mineralogical Society of America 58:pp. 589–622.
 623 <http://dx.doi.org/10.2138/rmg.2005.58.22>.

624 Evensen, N. M., Hamilton, P. J., O’Nions, R. K. 1978. Rare-earth abundances in
 625 chondritic meteorites. *Geochimica et Cosmochimica Acta*. 42: 1199 – 1212.

626 Farris, D.W., Jaramillo, C., Bayona, G., Restrepo-Moreno, S.A., Montes, C.,
 627 Cardona, A., Mora, A., Speakman, R.J., Glascock, M.D., Valencia, V. 2011.

628 Fracturing of the Panamanian Isthmus during initial collision with South America.
629 *Geology*. 39 (11): 1007 – 1010.

630 Frey, F.A., Gerlach, D.C., Hickey, R.L., López-Escobar, L., Minizaga-Villavicencio, F.
631 1984. Petrogenesis of the Laguna del Maule volcanic complex, Chile (36°S). *Contrib.*
632 *Mineral Petrol.* 88: 133 – 149.

633 González, H. 2001. Memoria Explicativa del Mapa Geológico del Departamento de
634 Antioquia. Escala 1:400.000. Medellín, INGEOMINAS. 240 pp.

635 Grosse, E., 1926. Estudio Geológico del Terciario carbonífero de Antioquia en la
636 parte occidental de la Cordillera Central de Colombia: Berlín, Verlag Von Dietrich
637 Reimer. 361 pp.

638 Halliday, A.N., Lee, D-C., Tommasini, S., Davies, G.R., Paslick, C.R., Fitton, J.G.,
639 James, D.E. 1995. Incompatible trace elements in OIB and MORB and source
640 enrichment in the sub-oceanic mantle. *Earth Planet. Sci. Lett.* 133: 379 – 395.

641 Harmon, R.S., Barreiro, B., Moorbath, S., Hoefs, J., Francis, P.W., Thorpe, R.S.,
642 Deruelle, B., McHugh, J. and Viglino, J.A. 1984. Regional O-, Sr-, and Pb- isotope
643 relationships in late Cenozoic calc-alkaline lavas of the Andean Cordillera: *Journal of*
644 *the Geological Society*. 141 (5): 803 – 822.

645 Hawkesworth, C.J., Powell, M. 1980. Magma genesis in the Lesser Antilles island
646 arc. *Earth and Planetary Science Letters*. 51: 297 – 308.

647 Hawkesworth, C.J., Norry, M.J., Roddick, J.C., Baker, P.E., Francis, P.W., Thorpe,
648 R.S., 1979. $^{143}\text{Nd}/^{144}\text{Nd}$, $^{87}\text{Sr}/^{86}\text{Sr}$, and incompatible trace element variations in
649 calc-alkaline andesitic and plateau lavas from South America. *Earth Planet. Sci.*
650 *Letters*. 42: 45 – 57.

651 Hemming S.R, McLeannan, S.M. 2001. Pb isotopic compositions of modern deep
 652 sea turbidites. *Earth Planet. Sci. Let.* 184, 489 – 503.

653 Hickey, R.L., Frey, F.A., Gerlach, D.C. López-Escobar, L. 1986. Multiple sources for
 654 basaltic arc rocks from the southern volcanic zone of the Andes (34° - 41° S): Trace
 655 element and isotopic evidence for contributions from subducted oceanic crust,
 656 mantle, and continental crust. *J. Geophys. Res.* 91 (B6): 5963 – 5983.

657 Hildreth, W.; Moorbath, S. 1988. Crustal contribution to arc magmatism in the Andes
 658 of central Chile. *Contribution to Mineralogy and Petrology.* 98: 455 – 489.

659 Irvine, T.N., Baragar, W.R.A. 1971. A guide to the chemical classification of the
 660 common volcanic rocks: *Canadian Journal of Earth Sciences.* 8 (5): 523 – 548.

661 Ishikawa T., Nakamura E. 1994. Origin of the slab component in arc lavas from
 662 across-arc variation of B and Pb isotopes. *Nature.* 370: 205 – 208.

663 James, D.E. 1982. A combined O, Sr, Nd, and Pb isotopic and trace element study
 664 of crustal contamination in central Andean lavas: I. Local geochemical variations.
 665 *Earth and Planetary Science Letter.* 57: 47 – 62.

666 James, D.E., Brooks, C., Cuyubamba, A. 1976. Andean Cenozoic volcanism:
 667 Magma genesis in the light of strontium isotopic composition and trace-element
 668 geochemistry. *Geol. Soc. Amer. Bull.* 87: 592 – 600.

669 Jaramillo, J. M., 1976. Volcanic rocks of the Río Cauca valley, Colombia S.A. Thesis
 670 Degree of Master of Arts, Rice University, Houston.

671 Kay, S., Mpodozis, C., Ramos, V.A., Munizaga, F. 1991. Magma source variations
 672 for mid-Tertiary magmatic rocks associated with a shallowing subduction zone and a

673 thickening crust in the Central Andes (28 – 33°S). In: Andean Magmatism and its
674 Tectonic Setting, Boulder, Colorado. Harmon, R.S., Rapela, C.W., eds. Spec. Pap.
675 Geol. Soc. Am. 265: 113 – 137.

676 Kellogg, J., Vega, V. 1995. Tectonic development of Panamá, Costa Rica, and the
677 Colombian Andes: Constraints from global positioning system geodetic studies and
678 gravity. Geological Society of America. Special Paper 295: 75 – 90.

679 Kerr, A.C. 2003. Oceanic Plateaus. In: Rudnick, R. (ed.) The Crust. Elsevier
680 Science: Oxford. Treatise on Geochemistry. 3: 537 – 565.

681 Kowallis, B.J., Heaton, J.S., Bringham, K., 1986. Fission-track dating of volcanically
682 derived sedimentary rocks. *Geology* 14, 19–22.

683 Labanieh, S., Chauvel, C., Germa, A., Quidelleur, X. 2012. Martinique: a clear case
684 for sediment melting and slab dehydration as a function of distance to the trench.
685 *Journal of Petrology*. 53: 2441 – 2464.

686 Leal-Mejía, H. 2011. Phanerozoic Gold Metallogeny in the Colombian Andes – A
687 tectono-magmatic approach: Ph.D. thesis, Barcelona (Catalonia), Spain, University
688 of Barcelona, 1000p.

689 LeMaitre, R. W., Bateman, P., Dudek, A., Keller, J., Lameyre-LeBas, M. J., Sabine,
690 P. A., Schmid, R., Sorensen, H., Streckeisen, A., Woolley, A. R. & Zanettin, B., A.
691 1989. *Classification of Igneous Rocks and Glossary of Terms* Oxford: Blackwell.

692 Lesage, G., Richards, J.P., Muehlenbachs, K., Spell, T.L. 2013. Geochronology,
693 geochemistry, and Fluid Characterization of the Late Miocene Buriticá Gold Deposit,
694 Antioquia Department, Colombia. *Economic Geology*. 108: 1067 – 1097.

695 Lonsdale, P. 2005, Creation of the Cocos and Nazca plates by fission of the
 696 Farallón plate: Tectonophysics. 404 (3 – 4): 237 – 264

697 López, A., Sierra, G.M., Ramírez, S. 2006. Vulcanismo Neógeno en el suroccidente
 698 antioqueño y sus implicaciones tectónicas. Boletín Ciencias de la Tierra. 19: 27 – 41.

699 Mackenzie, W.S., Donaldson, C.H., Guilford, C. 1984. Atlas of igneous rocks and
 700 their textures. Longman Scientific and Technical, UK.

701 Mantilla, L.C., Bissig, T., Valencia, V., Hart, C.J.R., 2013. The magmatic history of
 702 the Vetás–California mining district, Santander Massif, Eastern Cordillera, Colombia.
 703 Journal of South American Earth Sciences 45, 235–249.

704 Marín-Cerón, M.I. 2007. Major, trace element and multi-isotopic systematics of SW
 705 Colombian volcanic arc, northern Andes: Contributions of slab fluid, mantle wedge
 706 and lower crust to the origin of Quaternary andesites. Doctoral thesis. Okayama
 707 University, Japan. 133 pp.

708 Marín-Cerón, M.I., Moriguti, T., Makishima, A., Nakamura, E. 2010. Slab
 709 decarbonation and CO₂ recycling in the Southwestern Colombian volcanic arc.
 710 Geochimica et Cosmochimica Acta. 74: 1104 – 1121.

711 Marín-Cerón, M.I., Leal-Mejía, H., Bernet, M., Mesa-García, J., 2019. Late Cenozoic
 712 to Modern-Day Volcanism in the Northern Andes: A Geochronological,
 713 Petrographical, and Geochemical Review. In Cedié, F., Shaw, R.P. (eds.) Geology
 714 and Tectonics of Northwestern South America, Frontiers in Earth Sciences, Springer
 715 Nature Switzerland., p. 603-648.

716 Maro, G., Remesal, M.B. 2011. El vulcanismo de la Alta Sierra de Somún Curá: El
 717 Cerro Corona, Provincia de Río Negro, Argentina. Abstract. XVIII Congreso
 718 Geológico Argentino. Neuquén. 2 pp.

719 Marriner, G.F., Millward, D. 1984. Petrochemistry of Cretaceous to recent Vulcanism
 720 in Colombia. Journal of Geological Society of London. 141: 473 – 486.

721 McCourt, W.J., Aspden, J.A., Brook, M. 1984. New geological and geochronological
 722 data from the Colombian Andes: continental growth by multiple accretion. Journal of
 723 Geological Society, London. 141: 831 – 845.

724 McDonough, W.F., Chauvel, C. 1991. Sample contamination explains the Pb isotopic
 725 composition of some Rurutu island and Sasha seamount basalts. Earth Planet. Sci.
 726 Lett. 105: 397 – 404.

727 McDonough, W. F., Sun, S. S. 1995. Composition of the Earth. Chemical Geology.
 728 120: 223 – 253.

729 Mesa-Garcia, J. 2015. Combia Formation: a Miocene immature volcanic arc? EAFIT
 730 University Master thesis. 244 pp.

731 Nakamura, E., Campbell, I.H., Sun, S.S. 1985. The influence of subduction
 732 processes on the geochemistry of Japanese alkaline basalt. Nature. 316: 55 – 58.

733 Nelson, S.T., Montana, A. 1992. Sieve-textured plagioclase in volcanic rocks
 734 produced by rapid decompression. American Mineralogist. 77: 1242 – 1249.

735 Nixon, G.T., Pearce, T. H. 1987. Laser-interferometry study of oscillatory zoning in
 736 plagioclase: The record of magma mixing and phenocryst recycling in calc- alkaline
 737 magma chambers, Iztaccihuatl Volcano, Mexico. Amer. Mineral. 72: 1144 – 1162.

738 Ordoñez, O. 2002. Caracterizacáo isotópica Rb-Sr E Sm-Nd dos principais
 739 eventos magmáticos nos Andes Colombianos. Doctoral thesis. Brazilia University.
 740 165 pp.

741 Páez Acuna, L. 2012. Análisis estratigráfico y de proveniencia del Miembro Superior
 742 de la Formación Amagá en los sectores de la Pintada y Valparaiso (cuenca Amagá,
 743 andes noroccidentales). Master thesis on Earth Sciences. EAFIT University.

744 Pardo, N., Cepeda, H., Jaramillo, J.M., 2005. The Paipa volcano, Eastern Cordillera
 745 of Colombia, South America: volcanic stratigraphy. *Earth Sci. Res. J.* 9, 3–18.

746 Pardo-Casas, F., Molnar, P. 1987. Relative motion of the Nazca (Farallón) and
 747 South American plates since Late Cretaceous time. *Tectonics* 6 (3): 223 – 248.

748 Patiño, L. C, Carr, M., Feigenson, M. 2000. Local and regional variations in Central
 749 American arc lavas controlled by variations in subducted sediment input. *Contrib.*
 750 *Mineral Petrol.* 138, 265-283.

751 Pedersen R., Furnes, H. 2001. Nd- and Pb-isotopic variations through the upper
 752 oceanic crust in DSDP/ODP Hole 504B, Costa Rica Rift. *Earth and Planetary Sc.*
 753 *Lett.* 189: 221 – 235.

754 Pérez, A. L., 2005. Formación Combia: Litofacies y aproximación a su edad con
 755 base en estudios palinológicos, suroeste antioqueño. Undergraduate thesis.
 756 Medellín, EAFIT University.

757 Pérez, A.M., Marín-Cerón, M.I., Bernet, M., Sierra, G., Moreno, N. 2013.
 758 Resultados preliminares de AFT en la Formación Amagá, Pozos el Cinco- 1B y
 759 Venecia-1. XIV Congreso Colombiano de Geología Libro: XIV Congreso
 760 Colombiano de Geología. Resúmenes.

761 Perugini, D., Busa, T., Poli, G., Nazzareni, S. 2003. The role of chaotic dynamics
 762 and flow fields in the development of disequilibrium textures in volcanic rocks.
 763 *Journal of Petrology*. 44 (4): 733 – 756.

764 Pettijohn, F. J. 1975. *Sedimentary rocks*, third edition: New York, Harper & Row. 628
 765 pp.

766 Piedrahita, V.A., Bernet, M., Chadima, M., Sierra, G.M., Marín-Cerón, M.I., Toro,
 767 G.E., 2017. Detrital zircon fission-track thermochronology and magnetic fabric of the
 768 Amagá Formation (Colombia): Intracontinental deformation and exhumation events
 769 in the northwestern Andes. *Sedimentary Geology*, 356, 26-42.

770 Plank, T., Langmuir, C. 1998. The chemical composition of subducting sediment and
 771 its consequences for the crust and mantle, *Chem. Geol.* 145, 325 – 394.

772 Ramírez, D.A., López, A., Sierra, G.M., Toro, G.E. 2006. Edad y proveniencia de las
 773 rocas volcánicas sedimentarias de la Formación Combia en el suroccidente
 774 Antioqueño- Colombia. *Boletín Ciencias de la Tierra*. 19: 9 – 26.

775 Restrepo, J.J., Toussaint, J.F., González, H., 1981. Edades MioPliocenas del
 776 magmatismo asociado a la Formación Combia. Departamentos de
 777 Antioquia y Caldas, Colombia. *Geología Norandina*. 3: 2126.

778 Restrepo-Moreno, S.A., Foster, D.A., Stockli, D.F., Parra-Sánchez, L.N. 2009. Long-
 779 term erosion and exhumation of the „Altiplano Antioqueño“, Northern Andes
 780 (Colombia) from apatite (U-Th)/He thermochronology. *Earth and Planetary Science*
 781 *Letters*. 278: 1 – 12.

782 Restrepo-Moreno, S.A., Cardona, A., Jaramillo, C., Bayona, G., Montes, C., Farris,
 783 D.W. 2010. Constraining Cenozoic uplift/exhumation of the Panamá-Chocó Block by

784 apatite and zircon low-temperature thermochronology: insights on the onset of
 785 collision and the morphotectonic history of the region. Abstract. GSA Denver Annual
 786 Meeting. Geological Society of America Abstracts with Programs. 42 (5): 521.

787 Rios, A.M., Sierra, M.I. 2004. La Formación Combia: Registro de la relación entre el
 788 volcanismo Neógeno y la sedimentación fluvial, sección Guineales – Bolombolo,
 789 suroeste antioqueño. Undergraduate thesis. EAFIT University. 122 pp.

790 Rodríguez-Vargas, A., Koester, E., Mallmann, G., Conceicao, R.V., Kawashita, K.,
 791 Weber, M.B.I. 2005. Mantle diversity beneath the Colombian Andes, Northern
 792 Volcanic Zone: Constraints from Sr and Nd isotope. Lithos 82: 471 – 484.

793 Rodríguez, G., Arango, M.I., Bermúdez, J.G. 2012. Batolito de Sabanalarga,
 794 plutonismo de arco en la zona de sutura entre las cortezas oceánica y continental de
 795 los Andes del Norte. Boletín Ciencias de la Tierra. (32): 81 – 98.

796 Rollinson, H.R. 1993. Using geochemical data: evaluation, presentation,
 797 interpretation. Longman group UK. 352 pp.

798 Saenz, E.A. 2003. Fission track thermochronology and denudational response to
 799 tectonics in the north of the Colombian Central Cordillera. Master thesis. Shimane
 800 Un Salters, V. & Stracke, A. 2004. Composition of depleted mantle, Geochem,
 801 Geophys, Geosyst 5 (5): 27 pp.

802 Saunders, A.D., Norry M.J., Tarney, J. 1988. Origin of MORB and chemically
 803 depleted mantle reservoirs: trace element constraints. J. Petrology, Special
 804 Lithosphere Issue. 415 – 445.iversity. 131 pp.

805 Shibata T., Nakamura, E. 1997. Across-arc variations of isotope and trace element
 806 compositions from Quaternary basaltic volcanic rocks in northeastern Japan:

807 implications for interaction between subducted oceanic slab and mantle wedge. J.
808 Geophys. Res. 102: 8051 – 8064.

809 Sierra, G. 1994. Structural and sedimentary evolution of the Irra Basin, northern
810 Colombian Andes. Master thesis, Department of Geological Science, State
811 University of New York, Binghamton, NY. 102 pp.

812 Sierra, G.M., Marín Cerón, M.I., 2011. Petroleum Geology of Colombia. Amagá,
813 Cauca and Patía Basins. Agencia Nacional de Hidrocarburos. 104 pp.

814 Silva, J.C., Sierra, G.M., Correa, L.G. 2008. Tectonic and climate driven fluctuations
815 in the stratigraphic base level of a Cenozoic continental coal basin, northwestern
816 Andes. Journal of South American Earth Sciences. 26: 369 – 382.

817 Singer, B.S., Dungan, M.A., Layne, G.D. 1995. Textures and Sr, Ba, Mg, Fe, K, and
818 Ti compositional profiles in volcanic plagioclase: Clues to the dynamics of calc-
819 alkaline magma chambers. American Mineralogist. 80: 776 – 798.

820 Stern, C.R. 2004. Active Andean volcanism: its geologic and tectonic setting.
821 Andean Geology. 31 (2): 161 – 206.

822 Stern, C.R., Moreno, H., López-Escobar, L., Clavero, J.E., Lara, L.E., Naranjo, J.A.,
823 Parada, M.A., Skewes, M.A. 2007. Chilean volcanoes. In: The Geology of Chile.
824 Moreno, T., Gibbons, W. (eds) The Geological Society, UK. 147 – 178.

825 Streckeisen, A. 1976. To Each Plutonic Rock its Proper Name. Earth Science
826 Review. 12: 1-33.

827 Sun, S.-s. and McDonough, W.F.. 1989. Chemical and isotopic systematics of
828 oceanic basalts: implications for mantle composition and processes. In: 4.D.

829 Saunders and M.J. Norry (Editors). *Magmatism in the Ocean Basins*. Geological
830 Society. London. pp. 3 13-345.

831 Taboada, A., Rivera, Luís A., Fuenzalida, A., Cisternas, A., Philip, Hervé, Bijwaard,
832 H., Olaya, J., and Rivera, C., 2000. Geodynamics of the northern Andes:
833 Subductions and intracontinental deformation (Colombia). *Tectonics*. 19 (5): 787 –
834 813.

835 Tamura, Y., Nakamura, E. 1996. The arc lavas of the Shirahama Group Japan: Sr
836 and Nd isotopic data indicate mantle-derived bimodal magmatism. *Journal of*
837 *petrology*. 37 (6): 1307 – 1319.

838 Tassinari, C.C.G., Díaz, F., Buena, J. 2008. Age and sources of gold mineralization
839 in the Marmato mining district, NW Colombia: A Miocene – Pliocene epizonal
840 gold deposit. *Ore Geology Reviews*. 33: 505 – 518

841 Tatsumi, Y. 2005. The subduction factory: how it operates in the evolving earth. *GSA*
842 *today*. 15 (7): 4 – 10.

843 Tatsumi, Y., Stern, R.J. 2006. Manufacturing continental crust in the subduction
844 factory. *Oceanography*. 19 (4): 104 – 112.

845 Tatsumi, Y., Takahashi, T. 2006. Operation of the subduction factory and
846 production of andesite. *Journal of Mineralogical and Petrographical Sciences*. 101:
847 145 – 153.

848 Thorpe, R.S. 1984. The tectonic setting of active Andean volcanism. In *Andean*
849 *magmatism: Chemical and Isotopic Constraints* (Harmon, R.S.; Barreiro, B.A.;
850 editors). Shiva Geological Series, Shiva Publications, Nantwich, U.K. 4 – 8.

851 Thorpe, R.S., Francis, P.W. 1979. Variations in Andean andesite composition and
 852 their petrogenetic significance. *Tectonophysics*. 57: 53 – 70.

853 Thorpe, R. S., Francis, P. W, Hammill, M., Baker, M.C.W., 1982. The Andes.
 854 Andesites. Ed Thorpe, R.S. 187 – 205.

855 Toro, G; Restrepo, JJ; Poupeau, G; Saenz, E y Azdimousa, A. 1999. Datación por
 856 trazas de fisión de circones rosados asociados a la secuencia volcano –
 857 sedimentaria de Irra (Caldas). *Boletín de Ciencias de la Tierra*. 13: 28 – 34.

858 Trenkamp, R., Kellogg, J.N., Freymueller, J.T., Mora, H.P., 2002. Wide plate margin
 859 deformation, southern Central America and northwestern South America, CASA
 860 GPS observations. *Journal of South American Earth Sciences*. 15: 157 – 171.

861 Uribe – Mogollón, C.A., 2013. Hydrothermal evolution of the Titiribí mining district.
 862 Undergraduate thesis. EAFIT University. 127 pp.

863 van der Hammen, T., 1960. Estratigrafía del Terciario y Maastrichtiano continentales
 864 y tectogénesis de los Andes Colombianos, Informe No. 1279. Servicio Geológico
 865 Nacional, Bogotá. 128 pp.

866 Vargas, C., Mann, P.2013. Tearing and breaking off of subducted slabs as the result
 867 of collision of the Panama arc-indenter with northwestern South America. *Bull.*
 868 *Seismol. Soc. Am.* 103:2025–2046.

869 Vermeesch, P. 2009. RadialPlotter: A Java application for fission track,
 870 luminescence and other radial plots. *Radiation Measurements*. 44: 409 – 410.

871 Walker, G.P.L., Wilson, C.J.N., Froggat, P.C. 1991. An ignimbrite veneer deposits;
 872 the trail marker of a pyroclastic flow. *Journal of Volcanology and Geothermal*
 873 *Research*. 9: 409 – 421.

874 Weber, M.B.I., Tarney, J., Kempton, P.D. and Kent, R. W., 2002, Crustal make-up of
 875 the northern Andes: evidence based on deep crustal xenolith suites, Mercaderes,
 876 SW Colombia. *Tectonophysics*. 345: 49 – 82.

877 White, W.M. 2013. *Geochemistry* (1st ed). Hoboken, NJ, Wiley-Blackwell. 660 pp.

878 White, W.M., Dupre, B., Vidal, P. 1985. Isotope and trace element geochemistry of
 879 sediments from the Barbados Ridge – Demerara Plain region, Atlantic Ocean.
 880 *Geochim. Cosmochim. Acta*. 49: 1875 – 1886.

881 Wilson, M. 1989. *Igneous petrogenesis: a global tectonic approach*. London, UK,
 882 Chapman & Hall. 466 pp.

883 Winter, J.D. 2001. *Introduction to igneous and metamorphic petrology*. Upper Saddle
 884 River, New Jersey, Prentice Hall. 796 pp.

885 Wörner, G.; Davidson, J.; Moorbath, S.; Turner, T.L.; McMillan, N.; Nye, C.; López-
 886 Escobar, L.; Moreno, H. 1988. The Nevados de Payachata Volcanic Region
 887 18°S/69°W, Northern Chile. I. Geological, geochemical and isotopic observations.
 888 *Bulletin of Volcanology*. 30: 287 – 303.

889

890

891 **Figure captions**

892 Fig. 1 Overview map of the Colombia Andes, showing the subduction of the Nazca
 893 and Caribbean plates beneath the South American plate. Also shown are the

894 Western, Central and Eastern Cordilleras and areas of Pliocene to present volcanic
895 activity, as well as the location of the study area (map from GeoMapApp,
896 <http://www.geomapapp.org/>).

897 Fig. 2 Geological map of the Amagá basin with the locations of the Cerro Amarillo
898 (CA), Anzá-Bolombolo (AB) and La Metida Creek (MC) sections (modified after
899 Sierra and Marín-Cerón, 2011).

900 Fig. 3 Cerro Amarillo stratigraphic of the eastern Amagá basin.

901 Fig. 4 Anzá Bolombolo stratigraphic section of the western Amagá basin.

902 Fig. 5 La Mertida Creek stratigraphic section of the western Amagá basin.

903 Fig. 6 Combined fission-track data radial plots of A) apatite fission-track data and B)
904 zircon fission track data with central ages.

905 Fig. 7 A) Streckeisen (1976) volcanic rock classification diagram. Both the Anzá-
906 Bolombolo and Cerro Amarillo samples plot in the andesite, basalt field. B)
907 Pyroclastic rock classification diagram of Pettijohn (1975), in which almost all La
908 Metida Creek and Anzó-Bolombolo pyroclastic rock samples plot in the crystal tuff
909 field.

910 Fig. 8 A) AFM diagram plot (Irvine and Baragar, 1971) for volcanic rocks from the
911 Cerro Amarillo and Anzá- Bolombolo sections. For comparison, data of Marriner and
912 Millward (1984), Ordoñez (2001) and Leal-Mejía (2011) from the Combia Formation
913 are also shown. B) Total alkalis versus silica (TAS) diagram of volcanic rocks from
914 the Cerro Amarillo and Anzá Bolombolo sections; boundaries in the total alkalis are
915 from LeMaitre et al. (1989) for rock classification and Irvine and Baragar (1971) for
916 magma series classification (red solid line). For comparison, data of Marriner and
917 Millward (1984), Ordoñez (2001) and Leal-Mejía (2011) are shown.

918 Fig. 9 A) Primordial mantle (McDonough and Sun, 1995) normalized trace element
919 spider diagrams of volcanic rocks from the Cerro Amarillo and Anzá-Bolombolo
920 sections. B) Chondrite (Evensen et al., 1978) normalized Rare Earth Element (REE)
921 patterns of the volcanic rocks from the Cerro Amarillo section. C) For comparison
922 trace element patterns of published data by Leall-Mejia (2011) from Combia
923 Formation and hypabyssal porphyritic intrusions in the study area, as summarized in

Marín-Cerón et al. (2019). Trace element data normalized after Wood et al. (1979). D) For comparison, published REE data of Leal-Mejía (2011) normalized after Sun and McDonough (1989) for hypabyssal porphyritic intrusions, Combia Formation volcanic rocks and recent to present volcanism in the Central Cordillera, as summarized in Marín-Cerón et al. (2019). E) The La/Sm versus Ba/Th plot indicates that the Cerro Amarillo volcanic rocks were more derived from magma related to slab dehydration, whereas the Anzá-Bolombolo samples trend more towards sediment melting derived magma.

Fig. 10 Comparison between trace elements and isotopic systematics: A) Sr/Th versus $^{87}\text{Sr}/^{86}\text{Sr}$, and B) Ba/Th versus $^{143}\text{Nd}/^{144}\text{Nd}$, for volcanic rocks of the CA section (blue circles) and the AB section (red circles). The dashed line represents altered oceanic crust values (AOC) (after Barret, 1983). Both diagrams show that addition of fluids to the magma affected the geochemistry to the analyzed rocks and that the slab dehydration effects are more pronounced in the eastern than the western Amagá basin. C) Nd vs. Sr isotope ratio plot, showing the fields for the Northern Volcanic Zone (NVZ), Central Volcanic Zone (CVZ) and Southern Volcanic Zone (SVZ) as well as typical MORB composition, based on data from James et al. (1976); Hawkesworth et al. (1979); James (1982); Harmon et al. (1984); Frey et al. (1984); Thorpe (1984); Hickey et al. (1986); Hildreth and Moorbath (1988); Wörner et al. (1988); Walker et al. (1991); de Silva (1991); Kay et al. (1991); Davidson and de Silva (1992). Winter (2001); Marín-Cerón (2007), as summarized by and plot modified from Marín-Cerón et al. (2019). The data of our study are shown for the Cerro Amarillo section (yellow circles) and the Anzá-Bolombolo section (red circles), plotted over the NVZ field.

Fig. 11 Lead isotopic systematics of the Combia Formation shown for the Cerro Amarillo section (yellow circles) and the Anzá-Bolombolo section (red circles). A) Plots of $^{208}\text{Pb}/^{204}\text{Pb}$ vs $^{206}\text{Pb}/^{204}\text{Pb}$ for the Andean volcanic zones (Northern Volcanic Zone – NVZ; Central Volcanic Zone – CVZ; and Southern Volcanic Zone – SVZ) and the pre-Andean basement (plot modified from data compilation plot of Marín-Cerón (2019). Pacific sediments (Dasch, 1981; White et al., 1985); Paleozoic basement (Chiaradia and Fontboté, 2002); metalliferous sediments from DSDP leg 92 (Barret et al., 1987). B) Zoom on the $^{208}\text{Pb}/^{204}\text{Pb}$ vs $^{206}\text{Pb}/^{204}\text{Pb}$ diagram. Squares represent possible end-members and the respective trends of the interaction between each of

the components involved during magma formation. AOC: altered oceanic crust; HS: hemipelagic sediments; CS: carbonaceous sediments; LCC: lower continental crust. The solid blue line represents the linear trend of the samples suggesting bimodal mixing between primary magma (yellow star) and LCC. This plot is based on the compilation of Marín-Cerón et al. (2019); Cretaceous Domain (Kerr, 2003); Lower crust xenoliths (Weber et al., 2002); and ACC from Hole 504 (Pedersen and Furnes, 2001); NVZ data from Marín-Cerón (2007). The data from our study are shown for the Cerro Amarillo section (yellow circles) and the Anzá-Bolombolo section (red circles) and the plotted over the Northern Volcanic Zone field.

Fig. 12 Schematic diagram for magma source genesis of Combia Formation volcanism. AOC: altered oceanic crust; HS: hemipelagic sediments; CS: carbonaceous sediments; AFC: assimilation fraction crystallization processes (DePaolo, 1981); MASH: melting, assimilation, storage, and homogenization processes (Hildreth & Moorbath, 1988); LCC: lower continental crust; UCC: upper continental crust. Illustration based on model proposed for southwestern Colombian volcanism (Marín-Cerón, 2007) and is compatible with models presented in Marín-Cerón et al. (2019).

Tables

Table 1 Apatite fission-track data of the Combia Formation

Table 2 Zircon fission-track data of the La Metida Creek Formation

Table 3 Petrographic modal analyses

Table 4 Major elements (wt%) of the Cerro Amarillo and Anzá-Bolombolo section samples

Table 5 Trace elements (ppm) of volcanic rock samples of the Cerro Amarillo and Anzá-Bolombolo sections

Table 6 Isotopic compositions of volcanic rocks from the Cerro Amarillo and Anzá-Bolombolo sections.

986 Table 7 Estimated isotopic compositions and parameters for end-member involved in
987 magma genesis

Table 1 Apatite fission-track data of the Combia Formation

Sample number	Lithology	Number of grains	Single grain age range (Ma)	RhoS ($\times 10^5 \text{t/cm}^2$)	Ns	RhoI ($\times 10^5 \text{t/cm}^2$)	Ni	RhoD ($\times 10^5 \text{t/cm}^2$)	P(χ^2) (%)	Age dispersion (%)	Central age* (Ma) ± 2 SE	U (ppm) ± 2 SE
<i>AB section</i>												
JJ13	Pyroclastic agglomerate	60	2.8 – 36.9	0.28	67	4.88	1183	9.72	44.9	2.7	7.9 $\pm$ 2.1	6 $\pm$ 0
JJ22	Tuff	35	3.2 – 38.5	0.29	35	4.96	592	9.72	36.3	24.0	8.4 $\pm$ 3.1	7 $\pm$ 1
<i>MC section</i>												
JJ17	Lapilli – tuff	60	3.6 – 71.5	0.26	36	4.57	624	9.72	7.3	50.8	8.3 $\pm$ 3.1	6 $\pm$ 0
JJ14	Tuff	33	4.9 – 374.9	0.29	18	3.34	209	9.72	0.0	141.1	15.9 $\pm$ 11.1	4 $\pm$ 1
JJ21	Tuff	51	3.5 – 74.2	0.17	19	4.56	517	9.72	66.9	2.0	5.1 $\pm$ 2.5	6 $\pm$ 1
JJ18	Volcaniclastic sandstone	16	4.7 – 48.9	0.23	7	4.59	139	9.72	0.9	155.4	7.3 $\pm$ 9.4	6 $\pm$ 1
JJ19	Tuff	20	6.6 – 138.7	0.24	10	3.23	133	9.72	45.7	2.1	10.5 $\pm$ 7.3	4 $\pm$ 1
JJ20	Lapilli tuff breccia	27	4.9 – 138.5	0.50	28	5.68	317	9.72	6.9	57.1	15.8 $\pm$ 9.2	9 $\pm$ 1
combined		302	2.8-374.9					9.72	0.0	48.6	8.6 $\pm$ 1.4	6 $\pm$ 0

Note – RhoS: spontaneous track density. RhoI: induced track density; P(χ^2): Chi² probability. Fission-track ages were calculated using a Zeta value of 288.36 $\pm$ 7.90. AFT data calculated with Binomfit of Brandon (see Ehlers et al., 2005).

Table 2 Zircon fission-track data of the La Metida Creek Formation

Sample number	Lithology	Number of grains	Single grain age range (Ma)	RhoS (x10 ⁵ t/cm ²)	Ns	RhoI (x10 ⁵ t/cm ²)	Ni	RhoD (x10 ⁵ t/cm ²)	P(χ^2) (%)	Age dispersion (%)	Central age* (Ma) ± 2 SE	U (ppm) ± 2 SE
JJ1	Lapilli tuff breccia	30	1.0 – 14.3	4.65	175	17.6	662	2.71	24.4	24.2	6.7 $\pm$ 1.5	258 $\pm$ 21
JJ8	Tuff	12	2.1 – 19.4	5.89	58	24.1	237	2.67	3.3	44.3	6.2 $\pm$ 2.6	359 $\pm$ 47
JJ3	Tuff	25	4.0 – 19.5	10.6	383	27.5	993	2.71	47.5	6.6	10.0 $\pm$ 1.7	405 $\pm$ 28
JJ9	Tuff	28	0.9 – 25.5	13.2	570	33.1	1432	2.67	0.0	35.3	10.1 $\pm$ 2.1	493 $\pm$ 29
JJ10	Tuff	22	5.0 – 20.0	10.5	363	28.6	989	2.66	68.0	0.6	9.3 $\pm$ 1.5	427 $\pm$ 29
JJ4	Tuff	31	2.9 – 25.8	17.7	585	38.8	1280	2.70	0.3	24.1	11.2 $\pm$ 2.0	572 $\pm$ 36
JJ6	Volcaniclastic sandstone	51	2.9 – 46.7	10.6	909	22.1	1844	2.69	0.0	44.1	12.7 $\pm$ 2.4	327 $\pm$ 17
JJ5	Tuff	31	2.6 – 14.8	14.1	515	3.85	1405	2.69	73.1	6.5	9.4 $\pm$ 1.4	569 $\pm$ 14
JJ11	Lapilli tuff breccia	48	1.3 – 16.1	6.27	362	2.63	1517	2.65	14.2	19.6	6.1 $\pm$ 1.1	394 $\pm$ 22
JJ7	Lapilli tuff breccia	68	4.3 – 14.8	9.00	824	29.5	2705	2.68	94.0	0.7	7.8 $\pm$ 1.1	439 $\pm$ 20
Combined		346	0.9 – 46.7	10.2	4744	28.5	13206	2.68	0.0	33.9	9.1 $\pm$ 1.1	418 $\pm$ 14

Note – RhoS: spontaneous track density. RhoI: induced track density; P(χ^2): Chi² probability. Fission-track ages were calculated using a Zeta value of 191.6 $\pm$ 10.25. ZFT data calculated with Binomfit of Brandon (see Ehlers et al., 2005).

[illegible]

Table 4 Major elements (wt %) of the Cerro Amarillo and Anzó – Bolombolo section samples

Oxides (wt %)	JJ1 - 3	JJ1 - 6	JJ1 - 9	JJ1 - 13	JJ1 - 17	JJ1 - 18	JJ1 - 20	JJ1 - 23	JJ3 - 2	JJ3 - 5	JJ3 - 6	JJ3 - 9	JJ4 - 2	JJ4 - 3	JJ2-1- 10	JJ2-1- 14
SiO₂	53.20	52.68	52.56	52.62	52.55	52.10	52.49	52.64	52.09	51.19	52.07	52.22	51.78	52.34	55.45	55.78
Al₂O₃	14.80	14.70	14.82	14.71	14.66	14.61	14.59	14.87	18.30	15.29	15.05	15.22	17.49	17.66	19.02	18.93
Fe₂O₃ t	12.03	11.77	12.13	12.82	12.67	12.55	12.93	12.75	10.40	13.44	13.48	13.63	10.49	10.23	5.01	5.53
MnO	0.19	0.17	0.17	0.19	0.19	0.18	0.19	0.18	0.18	0.21	0.21	0.21	0.17	0.17	0.09	0.09
MgO	3.56	3.51	3.32	3.48	3.43	3.32	3.49	3.30	2.99	3.44	3.50	3.46	3.48	3.05	1.05	1.12
CaO	7.68	7.82	7.56	7.57	7.58	7.47	7.36	7.50	9.33	8.28	8.35	8.24	8.80	8.84	4.51	5.28
Na₂O	2.95	2.74	2.63	2.60	2.69	2.70	2.76	2.80	2.66	2.76	2.89	2.80	2.94	2.98	3.58	3.73
K₂O	1.89	1.75	1.94	1.88	1.98	1.84	1.79	1.82	1.20	1.40	1.25	1.45	1.47	1.48	3.58	4.60
TiO₂	1.31	1.30	1.38	1.39	1.38	1.38	1.38	1.38	0.86	1.03	1.06	1.06	0.96	0.99	5.10	0.52
P₂O₅	0.48	0.48		0.50	0.50	0.51	0.52	0.51	0.28	0.39	0.39	0.39	0.44	0.46	0.47	0.49
LOI	1.48	2.75	2.60	2.19	1.77	1.98	1.47	1.74	1.12	1.63	1.63	1.65	2.08	2.31	5.18	2.70
SUM	99.67	99.11	99.95	99.4	98.64	98.97	99.49	99.41	99.06	99.88	100.33	100.1	100.51	98.45	98.77	99.67

Table 5 Trace elements (ppm) of volcanic rock samples of the Cerro Amarillo and Anzó Bolombolo sections.

	JJ1-3	JJ1-6	JJ1-9	JJ1-13	JJ1-17	JJ1-18	JJ1-20	JJ1-23	JJ3-2	JJ3-5	JJ3-6	JJ3-9	JJ4-2	JJ4-3	JJ2-1-10	J22-1-14
Li	8.52	8.33	7.58	8.46	8.73	9.75	8.71	10.4	8.98	8.76	7.64	9.39	8.07	8.38	13.6	15.5
Sc	31	30.7	30.2	30.5	30.1	29.5	30.1	30.4	27.4	33	32.8	33.1	24	24.8	8	8.75
Ti	7840	7760	8270	8340	8210	8110	8250	8240	5070	6170	6160	6200	5690	5900	2710	3020
V	343	340	367	369	361	365	366	367	296	331	338	337	268	278	96.9	100
Cr	12.7	11.6	13.5	13.1	13.2	13.5	13.5	13.3	10.9	15.7	16.5	17.2	31.4	32.6	1.4	1.02
Co	33.1	30.7	33.7	34.5	34.9	35.2	34.7	35.5	25.4	34.7	34.2	35.6	28	26.3	9.58	11.8
Ni	14.9	13	16	16.5	17.1	17.3	16.6	18	7.18	9.99	9.79	10.3	20.2	17.7	2.35	2.63
Cu	224	203	227	220	223	240	226	238	127	184	169	179	191	187	117	143
Zn	107	136	116	120	114	116	115	116	86.6	111	112	114	100	96.9	70.3	74.2
As	25.1	6.22	6.78	3.1	7.12	10.5	6.82	8.08	3.19	4.37	4.68	4.61	5.37	5.99	4.38	3.9
Rb	43.9	42.3	49.9	51.4	49.2	51.5	49.3	50.1	25.8	32.9	34.7	34	37.2	38.3	169	130
Sr	475	478	470	457	459	461	453	457	558	497	457	473	483	489	1870	1170
Y	26	26	28	27.5	27.2	27.6	27.3	27.4	18.9	23.3	23.5	24	21	22.2	15.5	17
Zr	110	108	116	115	114	116	114	115	55.7	75	76.2	77	65.7	68.5	106	118
Nb	5.97	5.83	6.18	6.25	6.11	6.26	6.19	6.16	2.63	3.03	3.07	3.06	3.05	3.16	4.86	5.54

	JJ1-3	JJ1-6	JJ1-9	JJ1-13	JJ1-17	JJ1-18	JJ1-20	JJ1-23	JJ3-2	JJ3-5	JJ3-6	JJ3-9	JJ4-2	JJ4-3	JJ2-1-10	J22-1-14
Cd	0.0498	0.108	0.0574	0.0541	0.0487	0.0538	0.055	0.058	0.0441	0.0513	0.0469	0.0488	0.045	0.0461	0.0391	0.049
Cs	1.51	1.73	1.86	1.97	1.76	1.96	1.84	2	0.884	1.12	1.23	1.19	1.35	1.59	3	1.11
Ba	967	948	1030	990	1030	1010	1000	1010	607	842	823	847	717	742	1350	1250
La	10.5	10.5	11.2	11.2	11.1	11.3	11.2	11.2	5.7	7.46	7.54	7.62	7.14	7.39	19.7	21.4
Ce	22.7	22.6	23.9	24.3	23.8	24	23.9	23.9	12.4	16.2	16.4	16.5	15.4	15.7	36.5	39.9
Pr	3.21	3.21	3.42	3.46	3.36	3.42	3.41	3.41	1.83	2.39	2.41	2.42	2.22	2.29	4.69	5.13
Nd	14.7	14.6	15.5	15.8	15.5	15.7	15.4	15.4	8.64	11.3	11.5	11.7	10.3	10.7	18.4	20.2
Sm	3.87	3.94	4.24	4.29	4.16	4.2	4.25	4.18	2.48	3.24	3.33	3.37	2.81	2.92	3.74	4.13
Eu	1.21	1.23	1.27	1.27	1.26	1.28	1.28	1.24	0.858	1.09	1.07	1.1	0.98	1.04	1.2	1.29
Gd	4.53	4.46	4.74	4.71	4.73	4.67	4.66	4.73	2.96	3.87	3.92	3.96	3.34	3.52	3.3	3.59
Tb	0.713	0.717	0.747	0.758	0.745	0.754	0.755	0.753	0.479	0.623	0.645	0.66	0.542	0.58	0.456	0.487
Dy	4.5	4.55	4.65	4.77	4.68	4.85	4.78	4.78	3.19	4.08	4.18	4.23	3.58	3.77	2.54	2.84
Ho	0.925	0.929	1	1.01	0.984	0.986	0.975	0.984	0.665	0.844	0.869	0.868	0.746	0.789	0.503	0.548
Er	2.75	2.65	2.84	2.89	2.84	2.89	2.86	2.87	1.99	2.48	2.53	2.57	2.23	2.32	1.44	1.62
Tm																
Yb	2.53	2.55	2.64	2.71	2.64	2.69	2.65	2.67	1.87	2.32	2.36	2.4	2.13	2.18	1.39	1.59

	JJ1-3	JJ1-6	JJ1-9	JJ1-13	JJ1-17	JJ1-18	JJ1-20	JJ1-23	JJ3-2	JJ3-5	JJ3-6	JJ3-9	JJ4-2	JJ4-3	JJ2-1-10	J22-1-14
Lu	0.375	0.368	0.397	0.403	0.394	0.398	0.394	0.395	0.271	0.331	0.347	0.349	0.313	0.329	0.211	0.239
Hf	2.99	2.92	3.17	3.18	3.15	3.15	3.1	3.12	1.61	2.18	2.25	2.27	1.81	1.87	2.55	2.81
Ta	0.368	0.37	0.39	0.39	0.388	0.392	0.385	0.384	0.166	0.191	0.189	0.194	0.192	0.193	0.282	0.303
Tl	0.298	0.234	0.322	0.36	0.361	0.297	0.245	0.272	0.2	0.268	0.317	0.222	0.282	0.297	0.351	0.138
Pb	9.64	9.7	10.2	10.5	10.1	10.2	10.2	10.1	6	9.58	9.9	9.67	4.63	4.74	10.9	12.1
Th	1.98	1.97	2.07	2.11	2.06	2.1	2.08	2.09	1.03	1.39	1.4	1.41	1.19	1.23	4.46	4.95
U	1.08	1.08	1.18	1.11	1.12	1.11	1.1	1.11	0.517	0.854	0.857	0.843	0.528	0.532	1.22	2.19

Table 6 Isotopic compositions of volcanic rocks from the Cerro Amarillo and Anz Bolombolo sections.

	JJ1-3	JJ1-9	JJ1-17	JJ1-23	JJ3-2	JJ3-5	JJ3-9	JJ4-2	JJ2-1-10	JJ2-1-14
$^{87}\text{Sr}/^{86}\text{Sr}$	0.703911	0.703930	0.703931	0.703921	0.703916	0.703876	0.70387	0.703862	0.70417	0.70416
2σ	0.000006	0.000008	0.000008	0.000008	0.000006	0.000006	0.000006	0.000006	0.000006	0.000006
$^{87}\text{Sr}/^{86}\text{Sr}$ (leached)	0.70391	0.70393	0.70393	0.70393	0.70392	0.70388	0.70387	0.70387	0.70417	0.70417
2σ	0.000006	0.000006	0.000006	0.000008	0.000006	0.000006	0.000006	0.000006	0.000006	0.000006
$^{143}\text{Nd}/^{144}\text{Nd}$	0.51294	0.51293	0.51292	0.51295	0.51293	0.51296	0.51293	0.51298	0.51290	0.51289
2σ	0.000009	0.000018	0.000017	0.000010	0.000010	0.000007	0.000013	0.000008	0.000007	0.000010
ϵNd	5.959	5.930	5.684	6.223	5.846	6.420	5.905	6.882	5.255	5.191
$^{208}\text{Pb}/^{204}\text{Pb}$	38.72422	38.73927	38.73610	38.73702	38.71385	38.67951	38.68273	38.79674	38.74386	38.75921
2σ	0.00418	0.00418	0.00418	0.00418	0.00418	0.00417	0.00417	0.00418	0.004179	0.004180
$^{207}\text{Pb}/^{204}\text{Pb}$	15.61989	15.62459	15.62410	15.62365	15.62449	15.61989	15.62098	15.61286	15.60328	15.61037
2σ	0.00167	0.00167	0.00167	0.00167	0.00167	0.00167	0.00167	0.00167	0.001668	0.001669
$^{206}\text{Pb}/^{204}\text{Pb}$	18.96667	18.97333	18.97308	18.97264	18.93568	18.91002	18.91054	19.06522	19.00789	19.018005
2σ	0.00365	0.00365	0.00365	0.00365	0.00365	0.00364	0.00364	0.00367	0.003662	0.003666

Table 7 Estimated isotopic compositions and parameters for end-member involved in magma genesis

	$^{208}\text{Pb}/^{204}\text{Pb}$	$^{206}\text{Pb}/^{204}\text{Pb}$	Pb (ppm)	$^{142}\text{Nd}/^{144}\text{Nd}$	Nd (ppm)	$^{87}\text{Sr}/^{86}\text{Sr}$	Sr (ppm)
AOC^{1,9}	38.14	18.59	0.53	0.51280	4.71	0.70381	61.40
HS^{2,3,4}	38.86	18.64	9.59	0.51247	17.0	0.70763	336.16
CS^{2,3,4,5}	38.16	18.46	3.70	0.51242	0.89	0.70858	1 504.12
SC_1*	38.62	18.62	1.43	0.51285	5.94	0.70454	75.138
SC_2*	38.57	18.60	1.54	0.51284	1.15	0.70661	146.412
MW^{6,7}	37.90	18.50	0.02	0.51310	0.71	0.70270	9.80
PM*	38.45	18.59	0.82	0.51307	5.81	0.70469	114.95
LCC⁸	38.73	19.02	8.81	0.51309	9.50	0.70423	173.00

* Estimated values. Kd from Halliday et al. (1995). AOC: Altered oceanic crust; HS: Hemipelagic sediments; CS: Carbonaceous sediments; SC_1: Subduction component 1 (AOC+HS); SC_2: Subduction component 2 (SC_1 + CS); MW: Mantle wedge (5% cpx, 25% opx, 70% ol); PM: Primary magma; and LCC: Lower continental crust. Values are calculated based on procedures followed by Marín-Cerón (2007) for the SW Colombian volcanic arc. Values for AOC, HS, CS, MW, and LCC, are given according to data presented by Marín-Cerón (2007). SC_2 represents the interaction between AOC (85%) and sediments (HS – 10% and CS – 5%). PM is calculated assuming 40% of metasomatized mantle by subduction component. It is considered 10% of LCC interacts with primary magma for magma source genesis. ¹Pedersen et al. (2001); ²Plank and Langmuir (2000); ³Patiño et al. (2000); ⁴Vervoort et al. (1999); ⁵Hemming and McLennan (2001); ⁶Salters and Stracke (2004); ⁷Saunders et al. (1988); ⁸Weber et al. (2002); ⁹Barret (1983); and ¹⁰Halliday et al. (1995).

Fig. 1 Bernet et al.

Lithological units

Cenozoic	Quaternary	Qar	Recent alluvial deposits
		Qd	Depris deposits
		Qtl	Talus deposits
		Qt	Terrace deposits
	Oligocene-Miocene	Td	Dacitic porphyry
		Tadh	Andesitic porphyry
		Tada	Augite-andesitic porphyry
		Tdsa	Andesitic dikes and sills
		Tsc	Combia Fm. volcanoclastic member
		Tmc	Combia Fm. volcanic member
		Tos	Upper Amagá Fm.
		Toi	Lower Amagá Fm.
		Tdc	Cambumbia stock
Mesozoic	Cretaceous	Kcdu	Ursula stock
		Kgh	Hispania gabbro
		Kg	Gabbro
		Ksga	Altamira gabbro
		Ksta	Antioquia batholith
		Kdha	Heliconia quartzdiorite
		Kdhh	Heliconia hornblende diorite
		Kvb	Barroso Fm.
		Kvc	Quebradagrande Fm. volcanic member
		Ksc	Quebradagrande Fm. sedimentary member
		Kuh	Ultramafic harzburgite
		Kaa	Penderisco Fm.
		Kida	Altavista batholith
Paleozoic	Jurassic	Jus	Ultramafic rocks
		Jgr	Roneral gabbro
		Jdp	Pueblito diorite
		TRa	Amagá stock
	Paleozoic	Pgnp	Palmitas gneissic granite
		Pbsd	Ayurá Montebello Gr. clastic texture
		Pev	Ayurá Montebello Gr. green schists
Paleozoic	Paleozoic	Pes	Ayurá Montebello Gr. phyllite
		Pei	Ayurá Montebello Gr. interbedded
		Peni	La Iguana micaceous gneiss
		Png	Migmatite
		PEa	Caldas amphibolite

AB = Anzá and Bolombolo
MC = La Metdia Creek
CA = Cerro Amarillo

Fig. 2 Bernet et al.

CERRO AMARILLO SECTION

Fig. 3 Bernet et al.

ANZA - BOLOMBOLO SECTION

Trachy-andesite and pyroclastic sequence

Tuffaceous sequence

Fig. 4 Berner et al.

5 m

5 m

Fig 6 Berner et al.

Fig 7 Bernet et al.

Fig. 8 Bernet et al.

Fig. 9 Bernet et al.

Fig 10 Bernet et al.

Fig. 11 Bernet et al.

Fig. 12 Bernet al.