Ultra-depleted 2.05 Ga komatiites of Finnish Lapland: Products of grainy late accretion or core-mantle interaction?

Igor S Puchtel, Andrea Mundl-Petermeier, Mary Horan, Eero J Hanski, Janne Blichert-Toft, Richard J Walker

► To cite this version:

Igor S Puchtel, Andrea Mundl-Petermeier, Mary Horan, Eero J Hanski, Janne Blichert-Toft, et al.. Ultra-depleted 2.05 Ga komatiites of Finnish Lapland: Products of grainy late accretion or core-mantle interaction?. Chemical Geology, 2020, 554, pp.119801. 10.1016/j.chemgeo.2020.119801 . hal-02991119

HAL Id: hal-02991119

<https://hal.science/hal-02991119>

Submitted on 5 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Ultra-depleted 2.05 Ga komatiites of Finnish Lapland: Products of grainy
late accretion or core-mantle interaction?**

Igor S. Puchtel^{1*}, Andrea Mundl-Petermeier^{1,2}, Mary Horan³, Eero J. Hanski⁴,
Janne Blichert-Toft⁵, and Richard J. Walker¹,

¹Department of Geology, University of Maryland, 8000 Regents Drive, College Park, MD 20742, USA

²Department of Lithospheric Research, University of Vienna, Althanstraße 14, 1090 Vienna, Austria

³Department of Terrestrial Magnetism, Carnegie Institution for Science, 5241 Broad Branch Rd. NW,
Washington, DC 20015, USA

⁴Oulu Mining School, P.O. Box 3000, FI-90014 University of Oulu, Finland

⁵Laboratoire de Géologie de Lyon, Ecole Normale Supérieure de Lyon, CNRS UMR 5276, Université de Lyon,
46 Allée d'Italie, 69007 Lyon, France

*Corresponding author: ipuchtel@umd.edu

Revised for:
Chemical Geology

Revised for *Chemical Geology*
Version: June 24, 2020

Keywords: Hf-W, Sm-Nd, Lu-Hf, Re-Os and Pt-Os isotope systems; highly siderophile
elements; Paleoproterozoic komatiites; grainy late accretion; core-mantle interaction; Finnish
Lapland

Abstract

Rhenium-Os, Pt-Os, Sm-Nd, Lu-Hf, and Hf-W isotope data, together with lithophile trace element and highly siderophile element (HSE: Re, Os, Ir, Ru, Pt, and Pd) abundances, are reported for 2.05 Ga Jeesiörova and Kevitsa komatiites from the Central Lapland Greenstone Belt, Fennoscandia, Finland. Both komatiites are closely genetically related, with the Kevitsa dikes having served as feeding magma conduits to the Jeesiörova pillowed and massive lavas. The parental komatiite magma is estimated to have contained ~25 wt.% MgO and was, thus, derived from a mantle source at least as hot as those of some of its late Archean counterparts. A suite of Jeesiörova and Kevitsa whole-rock komatiite samples and olivine and chromite separates define an internal Re-Os isochron with an age of 2049 ± 13 Ma and an initial $\gamma^{187}\text{Os} = -0.2 \pm 0.2$ (2SE), indicating long-term chondritic Re/Os in the mantle source. By contrast, Pt-Os data for a set of Jeesiörova chromite separates define an average initial $\mu^{186}\text{Os} = +29 \pm 2$ (2SE), indicating a long-term history of suprachondritic Pt/Os in the mantle source. The absolute HSE abundances in the mantle source of the Jeesiörova-Kevitsa komatiite system are estimated to have been $120 \pm 5\%$ of the present-day Bulk Silicate Earth (BSE). This is the first komatiite system for which excess HSE in the mantle source, relative to modern BSE, has been documented.

The ^{147}Sm - ^{143}Nd and ^{176}Lu - ^{176}Hf data yield isochron ages and initial ratios of, respectively, 2046 ± 22 Ma with $\epsilon^{143}\text{Nd} = +3.7 \pm 0.3$, and 2072 ± 20 Ma with $\epsilon^{176}\text{Hf} = +8.7 \pm 0.4$ (2SE), indicating a long-term history of depletions of Nd relative to Sm, and Hf relative to Lu. The measured $\mu^{182}\text{W} = +1.5 \pm 3.3$ is indistinguishable from the modern mantle value. Despite being strongly depleted in highly incompatible lithophile trace elements, the Th-Nb-La systematics of the komatiites indicate ~1% crustal contamination of the original komatiite magma, assuming the contaminant was similar in composition to the calculated Fennoscandian Tonalite Average (FTA). This level of contamination would have also significantly modified the Nd, Hf, and W isotope compositions of the original komatiitic magma, but not the Os isotope compositions or HSE abundances. The calculated original komatiite magma, corrected for the effects of crustal contamination, would have had initial $\epsilon^{143}\text{Nd} \sim +4.9$, $\epsilon^{176}\text{Hf} \sim +10.2$, and $\mu^{182}\text{W} \sim -10$.

Our modeling indicates that the initial $^{186,187}\text{Os}/^{188}\text{Os}$ isotopic compositions and suprachondritic HSE abundances, coupled with the projected negative $\mu^{182}\text{W}$, are best explained by either (1) derivation from a mantle domain characterized by an excess of late accreted, differentiated planetesimal core metal, i.e., “grainy” late accretion, or (2) addition of chemically fractionated terrestrial core metal to the mantle source domain of the komatiites. The presence of these characteristics in the Jeesiörova-Kevitsa komatiite mantle source provides further evidence for the early creation and long-term survival of chemically diverse domains within the mantle.

1. Introduction

The chemically and isotopically heterogeneous nature of the mantle has long been established, largely through the study of modern rocks (e.g., Gast et al., 1964; Hart and Brooks, 1977; Zindler et al., 1982; White and Hofmann, 1982; White and Patchett, 1984; Hofmann, 1984, 1997; Jacobsen, 1988; Galer and Goldstein, 1991). However, the origin of the heterogeneities for different elements and isotopic systems (e.g., $^{146,147}\text{Sm}$ - $^{142,143}\text{Nd}$, ^{176}Lu - ^{176}Hf , ^{182}Hf - ^{182}W), as well as their length scales and residence times, remain topics of debate. Some of the heterogeneities have been argued to be primordial, reflecting initial planetary accretion/differentiation, including magma ocean crystallization processes (e.g., Goldstein and Galer, 1992; Albarède et al., 2000; Boyet and Carlson, 2005; Touboul et al., 2012; Jacobsen and Yu, 2015; Puchtel et al., 2013; 2016a; Rizo et al., 2016b; Mundl et al., 2017; Mundl-Petermeier et al., 2019; Tusch et al., 2019), while others may have originated as a result of a protracted accretion history (Willbold et al., 2011; Kruijer et al., 2015; Touboul et al., 2015; Puchtel et al., 2018; Archer et al., 2019).

Osmium isotope and highly siderophile element (HSE, including Re, Os, Ir, Ru, Pt, and Pd) abundance systematics of the mantle through time provide information about certain aspects of early Earth processes. For example, the observation that the HSE occur in approximately chondritic relative proportions in the Bulk Silicate Earth (BSE), and that absolute abundances of at least some of the HSE are higher than would be expected from metal-silicate equilibration during core formation, have led to the concept of late accretion, which purports that 0.5-1% of Earth's mass was accreted to the mantle subsequent to cessation of core formation (Kimura et al., 1974; Chou et al., 1983; Morgan, 1985, 1986). Issues related to the nature of late accretion include the composition of the late accreted materials and the time frame within which they were delivered to Earth and homogenized within the mantle (e.g., Bennett et al., 2002; Maier et al., 2009; Coggon et al., 2013; Walker et al., 2015; van de Löcht et al., 2018).

The Os isotopic composition and absolute HSE abundances of portions of the Archean mantle have been the subject of considerable prior study (e.g., Brüggmann et al., 1987; Walker et al., 1988; Foster et al., 1996; Gangopadhyay and Walker, 2003; Puchtel et al., 2014, 2016a,b; 2018; Maier et al., 2009). These studies have mostly focused on komatiites because komatiites are valuable probes of absolute HSE abundances and Os isotopic compositions in some portions of the mantle. This stems from the fact that most komatiites were formed by sufficiently high degrees of partial melting that the dominant sulfide hosts of HSE in the mantle were dissolved into the melts. Under such conditions, the HSE abundances in

komatiitic melts are much more similar to the initial mantle source compositions than those in basaltic melts. These high degrees of partial melting also led to comparatively high Os abundances in komatiitic melts, which make their Os isotopic compositions insensitive to crustal contamination. The high Os concentrations in komatiites, coupled with typically low Re/Os ratios, mean that they can provide precise initial Os isotopic compositions of mantle sources.

Compared to Archean komatiites, HSE and Os isotopes have been much less extensively studied in their Proterozoic counterparts, primarily due to the paucity of Proterozoic komatiites in the rock record. Here, we report combined ^{147}Sm - ^{143}Nd , ^{176}Lu - ^{176}Hf , ^{187}Re - ^{187}Os , ^{190}Pt - ^{186}Os , ^{182}Hf - ^{182}W , and HSE and lithophile trace element abundance data for the Proterozoic Kevitsa komatiitic dikes and associated Jeesiörova komatiites, all part of the Central Lapland Greenstone Belt in Fennoscandia, Finland. The Kevitsa komatiite dikes in particular display strong chemical differentiation trends, excellent preservation of igneous minerals, including olivine, and no signs of interaction with their wall rocks (Mutanen, 1997; Huhma et al., 2018). They hence provide ample opportunity for dating and obtaining information on the composition of the original komatiitic magmas. We use these data to (i) constrain the long-term evolution of the mantle domain from which the Fennoscandian Shield was produced and which also gave rise to the komatiite parental magmas, and (ii) further evaluate the extent of Archean-Paleoproterozoic mantle heterogeneity.

2. Geological background

Paleoproterozoic komatiites of Finnish Lapland are part of the Central Lapland Greenstone Belt (**Fig. 1**), which consists of a thick supracrustal rock series spanning 570 Ma (Hanski and Huhma, 2005). The komatiites are known to occur at two stratigraphic levels and, as such, have been referred to as the lower and upper komatiites (Hanski et al., 2001). The lower komatiites are found near the bottom of the succession and are assigned to the Kuusamo Group. They were emplaced subaerially and, locally, lie directly on Archean granitoid basement. A type occurrence has been described from the Möykkelmä area, where komatiites form part of a 250-m-thick komatiite-tholeiite sequence with a strong upper crustal signature in their chemical composition (Hanski and Huhma, 2005). Stratigraphically, the lower komatiites can be correlated with the komatiitic lavas of the Vetreny Belt in the SW part of the Fennoscandian Shield, which have an internal Re-Os isochron age of 2407 ± 6 Ma (Puchtel et al., 2016b).

The upper komatiites are associated with phyllites and black schists of the Savukoski Group whose sedimentary precursors were deposited on cratonic siliciclastic sediments of the Sodankylä Group (Hanski and Huhma, 2005). The upper komatiitic volcanic rocks extend from Finnish Lapland to northern Norway over a distance of ~400 km and are associated with Ti-rich picrites and basalts (Hanski et al., 2001). Unlike typical Archean komatiites, the upper komatiites are characterized by the abundance of volcanoclastic eruptions (Saverikko, 1985; Barnes and Often, 1990). They also occur as massive lava flows, pillow lavas, tuffs, and rare layered flows. The volcanic structures and types of the associated sedimentary rocks attest to submarine conditions of eruption in a deepened sedimentary basin. Although belonging to the Al-undepleted, or Munro-type of lavas of Nesbitt et al. (1979) typical of late Archean komatiite sequences, the upper komatiites are further distinct from Archean komatiites in having elevated TiO₂ contents and, therefore, they were classified as a Ti-enriched komatiite type (Hanski et al., 2001).

The upper komatiites are best preserved in the Jeesiörova area studied by Hanski et al. (2001). Magmatic olivine has not been reported to have been preserved in komatiitic lavas anywhere in Finnish Lapland, but magmatic clinopyroxene is well preserved at Jeesiörova, occurring as poikilitic intergrowths in olivine cumulates, as prismatic grains in gabbroic rocks, and as needle-like crystals in pillowed and massive lavas. The presence of magmatic clinopyroxene in the lavas allowed Hanski et al. (2001) to obtain a weighted average Sm-Nd isochron age of 2056±25 Ma for clinopyroxene-whole-rock pairs for the Jeesiörova komatiites. The initial $\epsilon^{143}\text{Nd}$ values range from +2 to +4 for individual samples, with the most LREE-depleted samples having the highest initial $\epsilon^{143}\text{Nd}$ values. This inverse correlation between the La/Sm and $\epsilon^{143}\text{Nd}$ values was interpreted to be the result of contamination by material of the upper crustal rocks through which the Jeesiörova komatiite magma ascended (Hanski et al., 2001).

Gangopadhyay et al. (2006) analyzed a set of whole-rock Jeesiörova komatiite samples and chromite separates for Re-Os isotope systematics. Although the Re-Os system in the whole-rock samples was found to be disturbed by post-magmatic processes, by analyzing chromite separates, the authors constrained the initial $^{187}\text{Os}/^{188}\text{Os}$ ratio for the komatiite lavas to be chondritic ($\gamma^{187}\text{Os}(\text{T}) = +0.1 \pm 0.6$).

The Kevitsa komatiitic dikes were discovered by Mutanen (1997) during an exploration drilling campaign of the Kevitsa Ni-Cu sulfide deposit. The dikes were penetrated by at least a dozen drill holes in an area of ~1 km², covering the ore-bearing part of the Kevitsa mafic-

ultramafic intrusion (**Fig. 1c**). This intrusion is located approximately 50 km east of the Jeesiörova area within pelitic metasediments of the Savukoski Group (**Fig. 1a**) and has a U-Pb zircon age of 2058 ± 4 Ma (Mutanen and Huhma, 2001), which is identical, within uncertainty, to the Sm-Nd age of the Jeesiörova komatiites. Each of the drill cores contains from one to four dike intersections at depths varying from 12 to 870 m beneath the surface. Although the true thickness of the dikes is less than 7 m, the lengths of the intersections reach up to 16 m because the drill holes cut through the dikes at sharp angles ($\leq 30^\circ$).

The dikes are generally subvertical, with the dip varying between 65° and 90° , and strike approximately ENE (Mutanen, 2005). They are characterized by strong internal differentiation related to olivine fractionation, having an olivine-enriched central part and finer-grained chilled zones near contacts with the wall rocks, which are composed of olivine-pyroxene cumulates of the Kevitsa intrusion. The differentiation is rather symmetrical across the dikes, suggesting that originally they were also steeply dipping. The dikes are chemically similar to the Jeesiörova komatiites and have been interpreted to represent their feeder magma conduits (Huhma et al., 2018); the absolute age of the dikes, however, has so far remained unknown.

3. Samples

Sampling of the Kevitsa komatiite dikes was done at the National Drill Core Depot of the Geological Survey of Finland, Loppi, Finland. Samples for chemical and isotopic analyses were selected from drill core DDH 814, which contains two dike intersections at depths of 152.10–159.55 and 175.00–188.50 m, with the corresponding estimated thicknesses of the dikes being ~1.3 and ~6.8 m.

For this study, we also prepared powders from hand specimens of the best preserved and most primitive komatiite lava samples 12D-PPR, 13-EJH, and 17.1-PPR from the Jeesiörova area (Hanski et al., 2001). In addition, we powdered sample LP-10 that was recently collected from an olivine-phyric massive komatiite lava flow from the same area.

4. Analytical techniques

Details of the majority of the analytical techniques used in this study have been reported in a number of previous publications (e.g., Puchtel et al., 2016a,b; 2018); these, therefore, are only briefly summarized here. In contrast, full details are provided for those techniques implemented here for the first time.

4.1. Analysis of mineral compositions

Mineral compositions were determined using a *JEOL JXA-8530F Plus Hyper Probe* instrument at the Center of Material Analysis, University of Oulu. The analytical conditions were an accelerating voltage of 15 kV and a beam current of 15 nA. Peak and background counting times were 10 and 5 s, respectively. The standard built-in ZAF correction routine was used.

4.2. Re-Os isotopic compositions and HSE abundances

The measurements of Ru, Pd, Re, Ir, and Pt were performed at the *Plasma Laboratory (PL)* on Faraday cups of a *ThermoFisher Neptune Plus* ICP-MS in static mode using 10^{13} Ohm amplifiers. Isotopic mass fractionation was monitored and corrected for by interspersing samples and standards. The external precision of the analyses was estimated, on the basis of standard measurements performed during the period of the analytical campaign, to be $^{185}\text{Re}/^{187}\text{Re} = 0.25\%$, $^{99}\text{Ru}/^{101}\text{Ru} = 0.26\%$, $^{191}\text{Ir}/^{193}\text{Ir} = 0.15\%$, $^{194}\text{Pt}/^{196}\text{Pt} = 0.10\%$, and $^{105}\text{Pd}/^{106}\text{Pd} = 0.08\%$ relative (2SD). The accuracy of the data was assessed by comparing the results for the reference materials IAG MUH-1 (Austrian harzburgite), IAG OKUM (ultramafic komatiite), and NRC TDB-1 (Diabase PGE Rock Material) obtained at the *Isotope Geochemistry Laboratory (IGL)* with the reference values. Concentrations of all HSE and Os isotopic compositions obtained at the *IGL* are within the uncertainties of the certified reference values (**Supplementary Table A1**).

The average total analytical blank (TAB) measured during the present analytical campaign was (in pg): Ru 6.0, Pd 17, Re 0.53, Os 0.42, Ir 1.2, and Pt 174 ($N = 9$). For the whole-rock komatiite samples, the average TAB constituted less than 0.1% for Os, Re, Ir, Ru, and Pd, and less than 1% for Pt of the total amount of element analyzed. For the olivine and chromite separates, the TAB for Os and Ir constituted less than 0.1%, for Ru less than 0.4%, for Re between 0.1 and 3%, for Pt between 14 and 19%, and for Pd less than 1% of the total amount of element analyzed. We, therefore, cite $\pm 0.1\%$ as the uncertainty on the concentrations of Os, $\pm 0.2\%$ for Ir, between 0.3 and 0.4% for Ru, between 0.1 and 1% for Pd, between 0.1 and 3% for Re, and between 1.0 and 19% for Pt, of the total amount of element analyzed. The uncertainty on the Re/Os ratio for each sample was calculated by multiplying the estimated uncertainties on the Re and Os abundances. These uncertainties vary between 0.30 and 3.0% relative.

4.3. Pt-Os isotopic data.

In order to obtain the amount of Os required for high-precision measurements of $^{186}\text{Os}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ (~100 ng), ~1.0 g of pure chromite separates from Jeesiörova komatiite samples 12D-PPR and 13-EJH, representing different fractions in terms of magnetic susceptibility and density, were digested in Carius tubes. For the initial unspiked digestions, six ~1 g fractions of chromite separate 12D-PPR and two fractions of 13-EJH were digested. After the digestion was complete, the tubes were chilled, opened, and ~2% of the acid sample solution from each CT were transferred into a 25 mL Pyrex™ CT for precise determination of Pt/Os, Re/Os, and Ir/Os. The spiked aliquots were processed using the same procedure utilized for the Re-Os and HSE analyses. From the remaining part of the unspiked acid sample solutions, Os was extracted and purified using the same protocol utilized for the Re-Os work. The Os cuts from the batch of two CT containing sample 13-EJH were combined into one cut and used for the precise measurements of $^{186}\text{Os}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ in this sample. For sample 12D-PPR, every Os cut from six individual CT was used for the precise measurements of Os isotopic compositions. For TAB-corrections of the data from the spiked aliquots, since the ID digestion introduced ~98% of the total blank, the TAB of the initial (IC) digestion was considered negligible.

The high-precision measurements of the $^{186}\text{Os}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios were performed by *N-TIMS* in static mode on a *ThermoFisher Triton*[®] mass spectrometer at the *IGL*. During each run, between 1200 and 2400 ratios were collected for each sample load; the in-run uncertainties on the measured $^{186}\text{Os}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios are quoted as 2SE. The possible isobaric interference of $^{186}\text{W}^{16}\text{O}_3^-$ on $^{186}\text{Os}^{16}\text{O}_3^-$ was monitored and corrected for by measuring masses $^{184}\text{OsO}_3^-$ and $^{183}\text{W}^{16}\text{O}_3^-$ using the electron multiplier.

The mean of the Johnson-Matthey Os standard runs during the period of data collection was 0.001302 ± 2 for $^{184}\text{Os}/^{188}\text{Os}$, 0.1198432 ± 18 (± 15 ppm) for $^{186}\text{Os}/^{188}\text{Os}$, and 0.1137950 ± 18 (16 ppm) for $^{187}\text{Os}/^{188}\text{Os}$ (2SD, $N = 14$); these long-term reproducibilities were used to assess the true uncertainty on the measured $^{186}\text{Os}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios for individual samples.

4.4. Tungsten isotopic compositions and abundances

The W isotope analyses were carried out at the *IGL*. For each sample, approximately 100 grams of powder were processed to obtain the ~ 1 μg of W necessary for high-precision W isotope measurements. The sample powders for each sample were digested in four 300 mL Savillex Teflon screw-cap vials using a 5:1 mixture of double-distilled concentrated HF and HNO_3 on a hot plate at 150°C for one week. Tungsten was separated and purified using the four-stage ion-exchange chromatography protocol described in Peters et al. (2019), with minor modifications. The third stage involving a 1.5 mL anion-exchange column was repeated to improve the separation of Ti from W, and this step significantly increased W ionization efficiency. Tungsten recovery using this procedure was better than 90%.

Tungsten isotopic compositions were measured by *N-TIMS* on a *ThermoFisher Triton* mass spectrometer at the *IGL* using a 2-line multi-static acquisition protocol and following a slightly modified technique described by Archer et al. (2017). All data are reported as $\mu^{182}\text{W}$ and $\mu^{183}\text{W}$, which are the part per million (ppm) deviations of $^{182}\text{W}/^{184}\text{W}$ and $^{183}\text{W}/^{184}\text{W}$, respectively, in a given sample from those of the in-house *Alfa Aesar* laboratory W standard. Uncertainties on $\mu^{182}\text{W}$, based on the long-term 2SD of our *Alfa Aesar* laboratory standard, were ± 4.5 ppm. The measured $^{183}\text{W}/^{184}\text{W}$ ratios were identical within uncertainties (± 6 $\mu^{183}\text{W}$ units) to those in the average *Alfa Aesar* standard data.

Tungsten abundances were determined by isotope dilution ICP-MS. Tungsten was purified using a previously established anion-exchange chromatography technique (e.g., Kleine et al., 2004). Concentrations were measured using the *Element 2* single-collector ICP-MS at the *PL*. Typical uncertainties were $\sim 5\%$ relative.

4.5. Sm-Nd isotopic compositions and abundances

The Sm-Nd isotopic data were collected at the *IGL*. Approximately 300 mg of whole-rock komatiite sample powder, or between 300 and 400 mg of pure clinopyroxene separates, were digested. Measurements of the Nd isotopic compositions were performed on Faraday cups of the *ThermoFisher Neptune Plus* ICP-MS at the *PL* in static mode followed by off-line correction for spike contributions. For each sample, 200 to 300 ratios were collected with 8 s integration times in blocks of 100 ratios each. For every block of data collected, the peaks were centered, and a 30 s baseline measurement was performed for each Faraday cup/amplifier pair by beam deflection. The effects of instrumental mass fractionation were corrected for relative to $^{146}\text{Nd}/^{144}\text{Nd} = 0.7219$ using an exponential law. A total of five to six 60 ppb AMES Nd standard solutions were run at the beginning and end of the analytical session, with 200 to 300 ratios collected during each measurement. The in-run precision

of the measured $^{143}\text{Nd}/^{144}\text{Nd}$ ratio for both samples and standards was between 5 and 7 ppm (2SE). During the course of the present analytical campaign, the external reproducibility of the 60 ppb AMES Nd standard solution measurements for $^{143}\text{Nd}/^{144}\text{Nd}$ was ± 7.0 ppm (2SD). The average $^{143}\text{Nd}/^{144}\text{Nd}$ measured for the AMES Nd standard during each analytical session was used to calculate the instrumental mass bias coefficient for correction of the measured $^{143}\text{Nd}/^{144}\text{Nd}$ in the samples using the long-term average $^{143}\text{Nd}/^{144}\text{Nd}$ value measured precisely for the AMES Nd standard on the *IGL ThermoFisher Triton* (0.512152 ± 2 ; Puchtel et al., 2018).

In order to assess the accuracy of the analyses, several separate powder aliquots of USGS GRM BIR-1 and BCR-1 were processed and analyzed using the same analytical protocol. The measured $^{143}\text{Nd}/^{144}\text{Nd}$ ratios for both GRM are identical, within their respective uncertainties, to the GeoRem preferred values (**Supplementary Table A2**), as well as to the average high-precision value for BCR-1 obtained on the *IGL ThermoFisher Triton* (0.512645 ± 1 ; Puchtel et al., 2018).

Measurements of the Sm isotopic compositions were also performed on Faraday cups of the *ThermoFisher Neptune Plus* ICP-MS at the *PL* in static mode. For each sample, 20 ratios were collected with 8 s integration times in blocks of 10 ratios each. The effects of instrumental mass fractionation were corrected for relative to $^{147}\text{Sm}/^{152}\text{Sm} = 0.56081$ using an exponential law. The uncertainty on the Sm/Nd ratio in the samples analyzed was estimated on the basis of replicate analyses of the USGS GRM BCR-1 ($^{147}\text{Sm}/^{144}\text{Nd} = 0.13941 \pm 0.00014$) to be 0.10% relative (2SD).

4.6. Lu-Hf isotopic compositions and abundances

The Hf isotopic compositions and Lu and Hf concentrations in the Jejsiörova komatiites were determined at the *IGL* and the Department of Terrestrial Magnetism (*DTM*), Carnegie Institution for Science. At the *IGL*, approximately 500 mg of the whole-rock sample powder and pure clinopyroxene separate were weighted out in 15 mL Teflon inserts of Parr bombs with 7 mL double-distilled concentrated HF, 1.5 mL HNO_3 , and an appropriate amount of mixed ^{176}Lu - ^{178}Hf spike, sealed in stainless steel jackets and digested in an oven at 170°C for one week. The solutions were dried down, 0.7 mL of concentrated *SeaStar* HClO_4 added to the residue, the inserts re-sealed and kept on a hotplate at 180°C for 24 hours. The solutions were then dried down at $\sim 230^\circ\text{C}$, and the residues converted into the chloride form using 6M HCl. This step was repeated several times until clear solutions were obtained, and then dried down.

At *DTM*, the residue was re-dissolved in 5 mL of a 1M HCl – 0.1M HF mixture, and loaded onto a 0.6×20 cm column filled with AG50W-X8 200-400 mesh cation-exchange resin. High field strength elements, including Hf, were eluted in the loading solution and an additional 5 mL of 1M HCl – 0.1M HF. After eluting 44 mL of 2.5M HCl, heavy REE, including Lu, were eluted in 12 mL of 4M HCl (Blichert-Toft, 2001). Hafnium was purified on a 0.6×10 cm column of Eichrom LN resin (100-150 μm bead size) by loading the sample in 5 mL of 2.5N HCl, followed by washes of 10 mL 2.5M HCl, 10 mL 6M HCl, 4 mL H_2O , 40 mL of a mixture consisting of 0.09M Citric acid – 0.45M HNO_3 – 1% H_2O_2 , 5 mL of the same solution without H_2O_2 , and, finally, 20 mL of 6M HCl – 0.06N HF. Hafnium was then eluted in 8 mL of 6M HCl – 0.4M HF. Lutetium was purified using Eichrom LN resin (50-100 μm bead size) on a 0.4×7 cm column. The HREE fraction from the first column was loaded in 2.5M HCl, followed by a wash of 30 mL 2.5M HCl to remove much of the other HREE, then Lu was collected in 10 mL 6M HCl. The resultant Hf and Lu cuts were used for the measurements of the Hf and Lu isotopic compositions.

Measurements of the Hf isotopic compositions were performed on the Faraday cups of the *ThermoFisher Neptune Plus* ICP-MS at the *PL* in static mode followed by off-line correction for spike contributions. For each sample, 200 to 300 ratios were collected with 8 s integration time in blocks of 100 ratios each. Before each sample measurement, a 60 s baseline measurement was performed for each Faraday cup/amplifier pair without beam deflection, and this baseline was then automatically subtracted from the sample beam. Instrumental mass fractionation was corrected for relative to $^{179}\text{Hf}/^{177}\text{Hf} = 0.7325$ using an exponential law. The Yb, Lu, Ta, and W isobaric interferences were corrected for using $^{173}\text{Yb}/^{176}\text{Yb} = 1.256$, $^{175}\text{Lu}/^{176}\text{Lu} = 37.70$, $^{181}\text{Ta}/^{180}\text{Ta} = 8129$, and $^{183}\text{W}/^{180}\text{W} = 109.0$. A total of five 30 ppb JMC-475 Hf standard solutions were run at the beginning and end of the analytical session, with 200 ratios collected during each measurement. During the measurements, the signal intensities for both the standard and the samples were kept at constant levels, between 1.8 and 2.6 V on the ^{178}Hf mass; the in-run precision of the measured $^{176}\text{Hf}/^{177}\text{Hf}$ ratio for both samples and standards was between 11 and 13 ppm (2SE). During the course of the present analytical campaign, the external reproducibility of the 30 ppb JMC-475 Hf standard solution measurements for $^{176}\text{Hf}/^{177}\text{Hf}$ was ± 14 ppm (2SD); this value was used to estimate the true uncertainty on the Hf isotopic analyses. The average $^{176}\text{Hf}/^{177}\text{Hf}$ ratio measured for the JMC-475 Hf standard solution during the analytical session was used to calculate the instrumental mass bias coefficient for correction of the measured $^{176}\text{Hf}/^{177}\text{Hf}$ ratio in the samples using the true value of $^{176}\text{Hf}/^{177}\text{Hf} = 0.282163 \pm 0.000009$ as determined by a multiple dynamic analysis protocol of Blichert-Toft et al. (1997).

In order to assess the accuracy of the analyses, an aliquot of USGS GRM BCR-1 was processed and analyzed using the same protocol as that used for the samples. The measured $^{176}\text{Hf}/^{177}\text{Hf}$ ratio of BCR-1 is identical, within the uncertainty, to the GeoRem preferred value.

Measurements of the Lu isotopic compositions were also performed on the Faraday cups of the *ThermoFisher Neptune Plus* ICP-MS at the *PL* in static mode. For each sample, 40 ratios were collected with 4 s integration time in blocks of 20 ratios each. Before each sample measurement, a 60 s baseline measurement was performed for each Faraday cup/amplifier pair without beam deflection, and this baseline was then automatically subtracted from the sample beam. Instrumental mass fractionation was corrected for by analysis of a 10 ppb Lu standard solution before and after every 4 samples. The Yb and Hf isobaric interferences were corrected for on-line using $^{176}\text{Yb}/^{173}\text{Yb} = 0.7962$ and $^{176}\text{Hf}/^{177}\text{Hf} = 0.2822$. The Lu concentration obtained for the SRM BCR-1 in this study was identical, within the uncertainty, to the GeoRem recommended value. The propagated uncertainty on the Lu/Hf ratio in the samples analyzed is estimated to be 0.5%. The total analytical blanks were 70 pg for Hf and 2 pg for Lu.

5. Results

5.1. Petrography of the komatiites

The studied samples of the Jeesiörova komatiites are massive rocks with up to 1 mm-sized euhedral, completely serpentinized olivine phenocrysts in a groundmass of prismatic or needle-like clinopyroxene crystals and devitrified glass (Fig. 2a in Hanski and Kamenetsky, 2003). The rocks also contain euhedral chromite grains up to 0.3 mm in size, often containing melt inclusions (Figs. 2b and 2d in Hanski and Kamenetsky, 2003).

Compared to the Jeesiörova rocks, the Kevitsa komatiites analyzed in this study are characterized by a much better preserved magmatic mineralogy. The rocks are olivine-porphyrific, with the most coarse-grained varieties found in dike centers, due to the presence of olivine phenocrysts (**Fig. 2A, B, E**). The size of the phenocrysts is commonly less than 6 mm in the center of the dikes, decreasing gradually to 0.1-0.2 mm near the dike margins. Occasionally, large amoeboid and embayed megacrysts are observed, with a maximum grain size of ~2 cm (**Fig. 3**). Olivine phenocrysts show various morphologies; they are usually euhedral to subhedral, but sometimes have irregular shapes with ragged outlines (**Fig. 2D**). The latter are interpreted to be broken pieces of larger phenocrysts. Also observed are elongated olivine bars up to 1 cm in length, with an aspect ratio of up to 20 (**Fig. 2B**). Although olivine phenocrysts contain ubiquitous chrome spinel inclusions 0.1-0.3 mm in size, some of the largest phenocrysts are nearly devoid of them; in these cases, chrome spinel grains are concentrated along the olivine grain boundaries.

The holocrystalline groundmass is composed of clinopyroxene, orthopyroxene, olivine, plagioclase, brown amphibole, and opaque minerals, including two kinds of spinels, chrome spinel, and lamellar ilmenomagnetite (**Fig. 2C**). These oxides occur as individual or composite grains, in which chrome spinel is partly surrounded by ilmenomagnetite. Plagioclase forms lath-like grains up to 1 cm in length, but other silicates are anhedral or equant, with their grain size ranging commonly between 0.1 and 0.5 mm.

The chilled margins of the dikes contain olivine microphenocrysts (<0.5 mm in size) and some chrome spinel grains in a fine-grained to cryptocrystalline groundmass composed mostly of clinopyroxene and plagioclase. It is noteworthy that no vesicles are found in the contact zones, indicating that either the pressure of emplacement was too high for volatiles to form a gas phase, or the volatile content was very low.

5.2. Mineral chemistry

Representative electron probe analyses of igneous minerals from the Kevitsa komatiites are listed in the **Supplementary Table A3**.

5.2.1. Olivine. Olivine exhibits a continuous compositional range from Fo_{94.1} to Fo_{73.8}, with the highest forsterite content reported for large phenocrysts and the lowest for small grains in the groundmass. Microphenocrysts in the chilled margin samples vary from Fo_{83.5} to Fo_{93.2}. The CaO content of all analyzed olivine grains is 0.20–0.32 wt.%, typical of magmatic olivine. This, together with a negative correlation between MgO and MnO, indicates the

absence of xenocrystic mantle olivine among the grains analyzed. Instead, all olivine grains are considered to be cognate phases crystallized from the same komatiitic parental magma.

5.2.2. Pyroxenes and plagioclase. Groundmass clinopyroxene has an augitic average composition of $\text{En}_{49}\text{Fs}_{10}\text{Wo}_{41}$ with Al_2O_3 ranging from 2.4 to 5.8 wt.% and TiO_2 from 0.12 to 0.71 wt.%. Orthopyroxene grains average $\text{En}_{71}\text{Fs}_{25}\text{Wo}_4$ and have 3.4–4.5 wt.% Al_2O_3 and 0.40–0.57 wt.% TiO_2 . The anorthite content of the analyzed plagioclase grains in the groundmass ranges from 57.7 to 72.1 mol.%.

5.2.3. Chrome spinel. The most primitive chrome spinel grains have MgO contents of around 15–16 wt.% and corresponding Mg# of 0.71–0.76, whereas in the most evolved grains, MgO and Mg# drop to 2.7 wt.% and 0.14, respectively. The MgO-rich grains show an average Al_2O_3 content of 14.7 wt.% and an average TiO_2 concentration of 0.38 wt.%, yielding $\text{Al}_2\text{O}_3/\text{TiO}_2$ typical of well-preserved chrome spinel grains in komatiitic lavas in Central Lapland, but distinct from the more Ti-rich spinels that occur in the associated picrites (Hanski and Kamenetsky, 2013).

5.2.4. Amphibole. The compositions of the brown groundmass amphibole (**Fig. 2B, C**) straddle the boundary between Ti-rich pargasite and Ti-rich magnesio-hastingsite. The TiO_2 contents fall between 2.9 and 4.0 wt.%, consistent with the magmatic nature of the mineral. Amphibole shows moderate Na_2O contents of 2.3–3.0 wt.%, whereas K_2O is low, 0.10–0.26 wt.%, consistent with the low level of K_2O in the parental komatiite magma.

5.3. Major and lithophile trace element abundances

The major and trace element concentration data for the Jeesiörova and Kevitsa komatiites are listed in **Tables 1 and 2**, and selected elements are plotted against MgO contents on variation diagrams in **Fig. 4** and as BSE-normalized values in **Fig. 5**.

The MgO abundances range between 25.3 and 28.6 wt.% in the Jeesiörova komatiites and between 16.2 and 28.3 wt.% in the Kevitsa komatiites. In the Kevitsa komatiite dikes, the lowest abundances are observed in the chilled margin samples and the highest abundances in the samples from the central parts. The abundances of TiO_2 , CaO, Al_2O_3 and incompatible lithophile trace elements in the dikes plot on well-constrained trends with negative slopes that intersect the MgO axes at ~51 wt.%; these intersections correspond to the composition of liquidus olivine obtained in this study and also reported for the dikes by Nicklas et al. (2019). The abundances of the compatible elements Ni and Cr show positive correlations with MgO

contents (**Fig. 4**). These strong correlations reflect olivine and chromite control during magmatic differentiation of the dikes, and also indicate immobile behavior of the elements analyzed during secondary alteration processes.

The four samples of the Jeesiörova komatiites analyzed in this study have slightly higher Ti, Cr, La, Ce, Th, and U concentrations compared to the Kevitsa komatiites (**Fig. 4**), but for the rest of major and trace elements, plot on the same trends. The Jeesiörova and Kevitsa komatiites have $\text{Al}_2\text{O}_3/\text{TiO}_2$ (11.7 ± 0.3 and 13.9 ± 0.1 , respectively, 2SE) that are lower, and $(\text{Gd}/\text{Yb})_{\text{N}}$ (1.53 ± 0.08 and 1.49 ± 0.01 , respectively, 2SE) that are higher, than the BSE values of 22.4 and 1.00, respectively. This is due to enrichment of the parental komatiite magma in Ti and the middle REE, rather than to depletion in Al and the heavy REE, as is evident from comparison with the typical Al-undepleted komatiites from Pyke Hill, Ontario, Canada (**Fig. 4A**). As such, they are classified as the Al-undepleted komatiite type of Nesbitt et al. (1979).

The BSE-normalized lithophile trace element abundances (**Fig. 5**) are characterized by strong depletions in light REE ($(\text{La}/\text{Sm})_{\text{N}} = 0.347\pm0.022$ and 0.264 ± 0.002 in the Jeesiörova and Kevitsa komatiites, respectively; 2SE), sloping HREE ($(\text{Gd}/\text{Yb})_{\text{N}} = 1.53\pm0.08$ and 1.49 ± 0.01 , respectively; 2SE), and enrichments in Th and U relative to the neighboring element with similar incompatibility, i.e., Nb ($(\text{Nb}/\text{Th})_{\text{N}} = 0.595\pm0.019$ and 0.583 ± 0.011 , respectively).

5.4. Re-Os isotopic compositions and HSE abundances

The Re-Os isotopic and HSE abundance data for the Jeesiörova and Kevitsa komatiites and olivine and chromite separates are listed in **Table 3** and plotted on the $^{187}\text{Re}/^{188}\text{Os}$ versus $^{187}\text{Os}/^{188}\text{Os}$ diagram in **Fig. 6**, on a CI chondrite-normalized plot in **Fig. 7**, and on MgO versus HSE variation diagrams in **Fig. 8**.

Thirty-three whole-rock samples and olivine and chromite separates define a regression line with a slope corresponding to an ISOPLOT Model 3 age of 2049 ± 13 Ma and a chondritic initial $^{187}\text{Os}/^{188}\text{Os} = 0.11284\pm28$ ($\gamma^{187}\text{Os} = -0.2\pm0.2$). This age is in good agreement with the internal Sm-Nd isochron age of 2056 ± 25 Ma obtained by Hanski et al. (2001) for the Jeesiörova komatiites. The initial $\gamma^{187}\text{Os} = -0.2\pm0.2$ is identical, within uncertainty, to the less precise average initial $\gamma^{187}\text{Os} = +0.1\pm0.6$ obtained for chromite separates from the Jeesiörova komatiites by Gangopadhyay et al. (2006).

In CI chondrite-normalized HSE diagrams (**Fig. 7**), the Jeesiörova and Kevitsa komatiite samples display typical komatiitic patterns with moderate enrichments in incompatible Pt, Pd,

and Re relative to compatible Os, Ir, and Ru ($(\text{Pd}/\text{Ir})_{\text{N}} = 2.84\text{--}10.1$). By contrast, the two olivine separates analyzed show an inverse relationship, being enriched in compatible relative to incompatible HSE, with the exception of the elevated Re abundance in one of the separates. This elevated Re concentration is, however, supported by the correspondingly more radiogenic Os isotopic composition of this separate, indicating that the olivine likely contained inclusions of interstitial melt enriched in incompatible elements.

In the MgO versus HSE diagrams (**Fig. 8**), variations in the Os and Ir contents show broad positive correlations with the MgO abundances, with the large degree of scatter most likely due to the presence of Os-Ir alloy on the liquidus, along with olivine. The Ru abundances display a more regular behavior and were likely mostly controlled by chromite fractionation during differentiation of the komatiites.

The Pt, Pd, and Re abundances show strong negative correlations with MgO contents and, with the exception of some scatter in the Re content, especially among the Jeesiörova komatiites, plot on the trends that pass through the olivine compositions; these variations are consistent with olivine control, indicating both incompatible behavior of these HSE during komatiite differentiation and their immobile behavior during post-magmatic processes. Of note, the Jeesiörova and Kevitsa rocks show a subchondritic $(\text{Pt}/\text{Pd})_{\text{N}} = 0.78 \pm 0.01$, which is typical of komatiites (e.g., Puchtel et al., 2009b), but is ~20% lower than the near-chondritic ratios in the Ni-Cu-PGE sulfide deposits of Finnish Lapland (Mutanen 1997; Törmänen et al., 2016).

5.5. Pt-Os isotope systematics

The high-precision Os isotopic data for the chromite separates from the Jeesiörova komatiite samples 12D-PPR and 13 EJH are presented in **Table 4** and plotted on the $^{190}\text{Pt}/^{188}\text{Os}$ versus $^{186}\text{Os}/^{188}\text{Os}$ diagram in **Fig. 6B**. The data for the seven chromite separates define a regression line with a slope corresponding to an age of ~2.0 Ga and an average initial $^{186}\text{Os}/^{188}\text{Os} = 0.1198369 \pm 3$ ($\mu^{186}\text{Os} = +29 \pm 2$, 2SE) calculated for the time of the komatiite emplacement derived from the Re-Os isochron (2049 Ma).

5.6. W isotopic compositions and abundances

The W abundances and isotopic compositions of the Jeesiörova and Kevitsa komatiites are listed in **Tables 2 and 5** and plotted in **Fig. 4B** and as BSE-normalized values in **Fig. 5**. The W abundances vary between 8.8 and 13 ppb in the Kevitsa komatiites; the only analyzed sample of the Jeesiörova komatiite contains 27 ppb W. In the MgO versus W variation

diagram (**Fig. 4B**), the data for the Kevitsa komatiites plot with some scatter around a nearly horizontal trend line, whereas the only data point for the Jeesiörova komatiite plots above the trend line.

The BSE-normalized trace element abundances (**Fig. 5**) for the Kevitsa komatiites are characterized by small negative to positive W abundance offsets relative to neighboring elements (i.e., Th and U) with similar incompatibility during mantle melting ($W/W^* = 0.95 \pm 0.13$, where $W/W^* = W_N / (\sqrt{[Th_N \times U_N]})$ and N are BSE-normalized values from Arevalo and McDonough (2008) and Hofmann (1988)). The W/W^* ratio in the only Jeesiörova komatiite sample analyzed (LP10) is 2.1.

The two Kevitsa komatiite samples and two replicates analyzed in this study for W isotopic composition, are characterized by $^{182}\text{W}/^{184}\text{W}$ ratios that are indistinguishable from $^{182}\text{W}/^{184}\text{W}$ measured in the laboratory reference material, with an average $\mu^{182}\text{W} = +1.5 \pm 3.3$ (2SE, $n = 4$), where $\mu^{182}\text{W}$ is the parts per million deviation of $^{182}\text{W}/^{184}\text{W}$ of a given sample from that of the reference material, which, by definition, is equal to zero (**Table 5**).

5.7. Sm-Nd and Lu-Hf isotopic compositions and abundances

The ^{147}Sm - ^{143}Nd isotopic and concentration data for the whole-rock samples and clinopyroxene separates of the Jeesiörova and Kevitsa komatiites are listed in **Table 6** and plotted in **Fig. 9**. Regression of the combined Sm-Nd isotopic data (**Fig. 9a**) for 13 samples yields a Model 1 ISOPLOT internal isochron age of 2046 ± 22 Ma (MSWD = 2.4), which is identical, within the uncertainty, to the Re-Os isochron age of 2049 ± 13 Ma. The initial $\epsilon^{143}\text{Nd}$ derived from the isochron is $+3.8 \pm 0.9$. The individual samples have very uniform initial Nd isotopic compositions, with initial $\epsilon^{143}\text{Nd}$ ranging between +3.5 and +3.9. A more precise initial $\epsilon^{143}\text{Nd} = +3.7 \pm 0.3$ (2SD, $n = 13$) is obtained by averaging the initial $^{143}\text{Nd}/^{144}\text{Nd}$ ratios calculated for each sample using the measured $^{147}\text{Sm}/^{144}\text{Nd}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ ratios. Our results are consistent with those obtained earlier for the most LREE-depleted Jeesiörova komatiites (Hanski et al., 2001).

The ^{176}Lu - ^{176}Hf data (**Fig. 9b**) for the Jeesiörova whole-rock komatiite sample 13-EJH and clinopyroxene separate define a regression line in $^{176}\text{Lu}/^{177}\text{Hf}$ - $^{176}\text{Hf}/^{177}\text{Hf}$ space with a slope corresponding to an ISOPLOT Model 1 age of 2072 ± 37 Ma, which also overlaps, within uncertainties, the Re-Os emplacement age of the Jeesiörova and Kevitsa komatiites. The initial $\epsilon^{176}\text{Hf}$ value derived from the regression is $+8.7 \pm 0.7$. The average initial $\epsilon^{176}\text{Hf}$

value calculated for the two samples analyzed ($+8.7 \pm 0.3$, 2SD, $n = 2$) is identical to the one derived from the isochron.

Plotted in **Fig. 9c** are the Nd and Hf isotopic data for the Jeesiörova komatiites, together with the Nd-Hf terrestrial array from Blichert-Toft and Puchtel (2010). In contrast to other late Archean and post-Archean komatiite systems (e.g., Kostomuksha: Blichert-Toft and Puchtel, 2010; Abitibi: Puchtel et al., 2018; Vetryny: Puchtel et al., 2016a; Gorgona: Thompson et al., 2003), the calculated initial $\varepsilon^{143}\text{Nd}$ and $\varepsilon^{176}\text{Hf}$ values of the Jeesiörova komatiites (**Fig. 9c**) do not plot on the terrestrial array of Blichert-Toft and Puchtel (2010).

6. Discussion

6.1. Jeesiörova-Kevitsa emplaced komatiite magma composition

Nicklas et al. (2019) reported major and trace element abundance data for olivine phenocrysts and whole-rock samples of the Kevitsa komatiites. Based on the Mg-Fe olivine-komatiite melt equilibria, these authors estimated that the parental magma to the Kevitsa komatiites contained 27.4 ± 2.4 wt.% MgO. By comparison for this study, in order to estimate the parental magma composition, we used an approach different from that of Nicklas et al. (2019). We performed mathematical incremental step-wise addition of olivine to a melt composition represented by sample KD-14 (MgO = 16.3 wt.%) until the melt was in equilibrium with olivine Fo_{94.1}, the most MgO-rich olivine population found in the Kevitsa komatiites (Nicklas et al., 2019, and this study). To select appropriate olivine-melt Fe-Mg distribution coefficients (K_D), we utilized the results of the isothermal crystallization experiments of komatiitic liquids carried out by Sossi and O'Neill (2016). Using the empirical K_D defined as $(\text{Fe}^{2+}/\text{Mg})_{\text{ol}}/(\Sigma\text{Fe}/\text{Mg})_{\text{melt}}$ and the T-MgO- K_D -log f_{O_2} relationships from their study, we adjusted the K_D values at each step of olivine addition. Because K_D defined in this way using bulk Fe in the melt is dependent on the oxygen fugacity, two different redox conditions were considered, $\Delta\text{FMQ} = +0.9$ and $\Delta\text{FMQ} = 0.0$. The former choice is based on the observed V partitioning between olivine and melt in the Kevitsa komatiites (Nicklas et al., 2019) and the latter on the composition of chrome spinel ($\text{Fe}^{3+}/(\text{Fe}^{3+} + \text{Al} + \text{Cr}) = 0.06$). The K_D values were varied between 0.28–0.26 and 0.29–0.27 in these two cases, respectively, and the required additions of olivine were 34 and 40%.

The two resulting major element compositions of the emplaced komatiite melt are similar, with the most important difference being the MgO content. The more oxidized conditions yielded an MgO content of 25.2 wt.%, while the FMQ buffer conditions gave an MgO content

of 26.3 wt.%. These estimates are identical, within the uncertainties, to those obtained by Nicklas et al. (2019) using an independent technique, and corroborate the komatiitic composition of the emplaced Kevitsa komatiite magma. The results indicate that the parental magmas to the Kevitsa, and, therefore, also Jeesiörova, komatiites approached those of their late Archean counterparts from the Abitibi and Belingwe greenstone belts (Puchtel et al., 2009b) in terms of MgO content, and, therefore, their liquidus temperatures.

6.2. *Lithophile trace element and isotope systematics of the komatiites*

The Jeesiörova and Kevitsa komatiites have low $(\text{La}/\text{Sm})_{\text{N}}$ (0.347 ± 0.022 and 0.264 ± 0.002 , respectively), and high positive initial ϵ_{Nd} and ϵ_{Hf} ($+3.7 \pm 0.3$ and $+8.7 \pm 0.3$, respectively), indicating derivation from a mantle source that was strongly depleted in highly incompatible, lithophile trace elements. These komatiites are ~2.5 times more strongly depleted in the light REE than, e.g., modern N-MORB (average $(\text{La}/\text{Sm})_{\text{N}} = 0.654$; Hofmann, 1988). Yet, the lavas are also characterized by enrichments in Th, U, and W relative to Nb, an element with similar incompatibility during melting of spinel peridotite ($(\text{Nb}/\text{Th})_{\text{N}} = 0.583 \pm 0.011$), and a negative Nb anomaly ($\text{Nb}/\text{Nb}^* = 0.582 \pm 0.010$, where $\text{Nb}/\text{Nb}^* = \text{Nb}_{\text{N}}/\sqrt{(\text{Th}_{\text{N}} \cdot \text{La}_{\text{N}})}$). This is most likely indicative of minor, yet significant crustal contamination.

Upper crustal rocks are generally characterized by low $(\text{Nb}/\text{Th})_{\text{N}}$ and $(\text{Nb}/\text{La})_{\text{N}}$ ratios and, as a result, display pronounced negative Nb anomalies relative to the elements with similar incompatibility during mantle melting (Th and La) on BSE-normalized trace element diagrams, i.e., their $\text{Nb}/\text{Nb}^* \ll 1.0$ (Rudnick and Fountain, 1995; Rudnick and Gao, 2014). By contrast, primary uncontaminated komatiitic magmas commonly do not show a Nb anomaly, i.e., their $\text{Nb}/\text{Nb}^* = 1.00$ (Jochum et al., 1991; Puchtel et al., 1998). Jochum et al. (1991) noted that the elements most sensitive to crustal contamination in komatiite magmas include Th, U, and light REE and, to a lesser extent, Nb. As a result, the Nb/Nb^* ratio can be utilized as a measure of the degree of crustal contamination and, when applied in combination with the lithophile element isotopic systems, to discriminate between contaminated and uncontaminated lavas. Based on these observations, it is likely that the original Jeesiörova-Kevitsa komatiite magma was contaminated with upper crustal rocks of the Fennoscandian Shield *en route* to the surface.

In order to test this hypothesis and calculate the major and trace element and Nd, Hf, and W isotopic compositions of the original Jeesiörova-Kevitsa komatiite magma, we performed model mixing calculations, the results of which are presented in **Table 8**. For the crustal contaminant endmember, we assumed an average composition of the Vodla Block tonalitic

gneisses that are the dominant upper crustal rock type within the adjacent Karelian granite-greenstone terrane (Puchtel et al., 2016b). These Karelian tonalites are similar in age and major- and trace element composition to their counterparts from Finnish Lapland (e.g., Huhma et al., 2012), but, unlike the latter, have been better characterized in terms of the isotopic systematics of interest (**Table 8**). This average tonalite composition from Puchtel et al. (2016b) is henceforth referred to as the Fennoscandian Tonalite Average (FTA). In order to calculate the degree of contamination and the trace element abundances and Nd and Hf isotopic compositions of the original, uncontaminated Jeesiörova-Kevitsa komatiite magma, we assumed that it had a $Nb/Nb^* = 1.00$. Calculations show that to produce the emplaced Jeesiörova-Kevitsa komatiite magma with $Nb/Nb^* = 0.582$, the original komatiite magma had to be mixed with ~1% of the average FTA ($Nb/Nb^* = 0.118$).

Using 1% contamination and the Nd and Hf abundances and initial Nd and Hf isotopic compositions of the FTA calculated at the time of the Kevitsa komatiite magma emplacement, the original Kevitsa komatiite magma and, thus, its source is calculated to have had initial ϵNd and ϵHf values of +4.9 and +10.2, respectively.

On the ϵNd versus ϵHf evolution diagram in **Fig. 9c**, the calculated datum for the Jeesiörova-Kevitsa komatiite source plots within uncertainty of the terrestrial evolution curve of Blichert-Toft and Puchtel (2010). This indicates coupled, or congruent, behavior of the two isotope systems in the source of the original komatiite magma. It appears, therefore, that the Nd-Hf isotope decoupling observed in the Jeesiörova-Kevitsa emplaced komatiite magma was caused by contamination with the older tonalitic crust that had isotopically evolved away from the terrestrial evolution curve by the time it was mixed with the original komatiite magma (**Table 8**). The datum for the Jeesiörova-Kevitsa komatiite source also plots near the Depleted MORB Mantle (DMM) value of Goldstein and Jacobsen (1988) at 2049 Ma (**Fig. 9c**), indicating derivation from a mantle source that was already as long-term depleted in incompatible lithophile trace elements as DMM at 2.05 Ga.

6.3. Osmium isotope - HSE systematics of the Jeesiörova-Kevitsa komatiite source

Although the Os isotopic compositions and absolute and relative HSE abundances in the mantle sources of early and late Archean komatiites have been the subject of numerous studies over nearly four decades (e.g., Brüggemann et al., 1987; Walker et al., 1988; Foster et al., 1996; Gangopadhyay and Walker, 2003; Puchtel et al., 2009a; 2014, 2016b; Maier et al., 2009), studies of post-Archean komatiites have been rare, in part because the number of Proterozoic and younger komatiites is far more limited.

In order to characterize the HSE present in the Paleoproterozoic mantle source of the Jeziörova-Kevitsa komatiites, the effects of crustal contamination must first be evaluated. We again use the average FTA composition from Puchtel et al. (2016b) to assess the effects of 1% contamination of the original Jeziörova-Kevitsa komatiite magma on its Os isotopic composition and HSE abundances. Because of the very low concentrations of these elements in the FTA, the effects are negligible. Contamination of this magnitude would have increased $\mu^{186}\text{Os}$ and $\gamma^{187}\text{Os}$ values by only 0.004 (based on the calculated $^{186}\text{Os}/^{188}\text{Os} = 0.1198767$ at 2.05 Ga for the FTA assuming its derivation from a BSE-type mafic source at 3.2 Ga, per Puchtel et al., 2016b) and 0.01 (based on the calculated $^{187}\text{Os}/^{188}\text{Os} = 1.2091$ at 2.05 Ga for the FTA assuming its derivation from a BSE-type mafic source at 3.2 Ga, per Puchtel et al., 2016b) units, respectively, which are well within the uncertainties of the current estimates for these values. Similarly, effects on absolute HSE abundances would have been $\ll 1\%$.

Estimation of the HSE abundances in the mantle source of the Jeziörova-Kevitsa komatiites is achieved by the bootstrap method of Puchtel et al. (2004b) that sequentially combines Os isotopic and HSE concentration data. First, we used the initial $^{187}\text{Os}/^{188}\text{Os}$ and $^{186}\text{Os}/^{188}\text{Os}$ ratios calculated for the Jeziörova-Kevitsa komatiites to estimate the time-integrated evolution of Re/Os and Pt/Os in their mantle source. In order to place minimum constraints on these long-term source ratios, we calculated the parent/daughter elemental ratios necessary to arrive at the Os isotopic composition at 2049 Ma by assuming generation of this mantle domain soon after Solar System formation. It is estimated that this source evolved from an early Solar System $^{187}\text{Os}/^{188}\text{Os} = 0.09517$ at 4567 Ma (Archer et al., 2014) to the initial $^{187}\text{Os}/^{188}\text{Os} = 0.11285 \pm 23$ at 2049 Ma with $^{187}\text{Re}/^{188}\text{Os} = 0.397 \pm 5$. This ratio is well within the range of bulk chondritic meteorites. Using the early Solar System $^{186}\text{Os}/^{188}\text{Os} = 0.1198269$ at 4567 Ma (Brandon et al., 2006) requires a source with a $^{190}\text{Pt}/^{188}\text{Os} = 0.00269 \pm 8$ to have evolved to its $^{186}\text{Os}/^{188}\text{Os} = 0.1198369 \pm 3$ at 2049 Ma. In contrast to Re/Os, the required minimum $^{190}\text{Pt}/^{188}\text{Os}$ ratio is $\sim 50\%$ higher than the average $^{190}\text{Pt}/^{188}\text{Os} = 0.00180 \pm 17$ (2SD) in bulk chondritic meteorites (as compiled from the data of Horan et al. (2003), Brandon et al. (2005, 2006), Fischer-Gödde et al. (2010), and van Acken et al. (2011)).

The calculated initial $\gamma^{187}\text{Os}$ of the Jeziörova-Kevitsa komatiite source is also within the range of those for the majority of Archean and Paleoproterozoic komatiite sources, which were characterized by initial $\gamma^{187}\text{Os}$ values between -0.1 and $+1.3$, also well within the range of 95% of chondritic meteorites (**Fig. 10a**). The $^{186}\text{Os}/^{188}\text{Os}$ isotope data available for Archean and Paleoproterozoic komatiite sources indicate that mantle sources of the 2.7 Ga

Abitibi (Canada), 2.7 Ga Belingwe (South Africa), and 2.4 Ga Vetreny (Fennoscandia) komatiite systems evolved with time-integrated Pt/Os within the chondritic range (**Fig. 10b**), whereas the coupled high initial $^{186,187}\text{Os}/^{188}\text{Os}$ of the 2.8 Ga Kostomuksha komatiite system (Fennoscandia) require long-term suprachondritic Pt/Os and Re/Os (Puchtel et al., 2005). By contrast, the 3.5 Ga Komati and 3.3 Ga Weltevreden komatiite systems (South Africa) evolved with non-chondritic Pt/Os, but chondritic Re/Os ratios (Puchtel et al., 2014), displaying decoupling of the Re-Os and Pt-Os isotopic systems. The calculated suprachondritic initial $\mu^{186}\text{Os} = +29 \pm 2$ in the Jeesiörova-Kevitsa komatiite source obtained in this study is similar to $\mu^{186}\text{Os} = +22 \pm 7$ in the source of the Weltevreden komatiite system (Puchtel et al., 2014).

Because of the availability of high-precision Os isotopic data and immobile behavior of HSE in the Kevitsa komatiites during post-magmatic alteration, absolute abundances of incompatible Re, Pt, and Pd and compatible Os in the Jeesiörova-Kevitsa komatiite source can be estimated using the bootstrap protocol of Puchtel et al. (2004b).

As was shown in a number of previous studies (e.g., Alard et al., 2000; Lorand and Alard, 2001; Luguet et al., 2007), $\geq 90\%$ of the HSE budget of the mantle resides in two major types of sulfides. The high-temperature Os-Ir-Ru-rich Fe-Ni monosulfide solid solution (*mss*) usually occurs as inclusions in liquidus phases, primarily olivine, whereas lower-temperature Cu-Ni sulfides occupy interstitial space between the liquidus phases. During partial melting of mantle peridotite, Cu-Ni sulfides enter the melt, whereas *mss* remains trapped in the melting residue. It takes $\sim 18\text{--}25\%$ partial melting (Barnes et al., 1985; Keays, 1995; Luguet et al., 2007; Mungall and Naldrett, 2008; Naldrett, 2010; Fonseca et al., 2011; 2012; Kiseeva et al., 2017) for all the low-temperature Cu-Ni sulfides to get exhausted in the mantle. As the degree of melting continues to increase, the magma becomes sulfide-undersaturated. It has also recently been shown that decrease in $f\text{S}_2$ with increase in the degree of melting triggers exsolution of Os-Ir alloys from the refractory *mss* in the residue (Fonseca et al., 2011; 2012). All low-degree (basalts) and the majority of higher-degree (picrites and komatiites) partial melts are characterized by compatible behavior of Os and Ir during magmatic differentiation, indicating that their parental magmas remained saturated in Os-Ir alloys (Puchtel et al., 2004b; Barnes and Fiorentini, 2008). However, some lavas, such as the 2.8 Ga Kostomuksha and the 3.55 Ga Schapenburg komatiites, exhibited an incompatible behavior of Os and Ir during magma differentiation, likely indicating near-complete exhaustion of Os-Ir alloys in the mantle sources of these komatiites (Puchtel and Humayun, 2005; Puchtel et al., 2009a).

688 Apparently, the sulfide saturation status of komatiite magmas is a crucial parameter that
689 plays a key role in determining whether their HSE abundances accurately reflect those in their
690 mantle sources. In the following discussion, we evaluate the sulfide saturation status of the
691 Jeesiörova-Kevitsa komatiite magma during both mantle melting and after emplacement.

692 Sulfide saturation of a mafic magma is determined by several key parameters, the most
693 important of which are pressure (P), temperature (T), and FeO content. In their recent study,
694 Steenstra et al. (2020) combined the results of their own work with previously published
695 experimental data of Mavrogenes and O'Neill (1999), Li and Agee (2001), Holzheid and
696 Grove (2002), O'Neill and Mavrogenes (2002), Jugo et al. (2005), Liu et al. (2007), Brennan
697 (2008), Kiseeva and Wood (2013, 2015), Ding et al. (2014, 2018), Fortin et al., (2015), Wood
698 and Kiseeva (2015), Smythe et al. (2017), and Steenstra et al. (2018) to produce a refined
699 parameterization model for the sulfur concentration at sulfide saturation (SCSS) as a function
700 of P, T, and major and trace element abundances in silicate liquids (Eq. 4 in Steenstra et al.,
701 2020).

702 In order to calculate the P-T parameters of the Jeesiörova-Kevitsa komatiite magma
703 generation, we used the mantle melting parameterization model of McKenzie and Bickle
704 (1988). These calculations yield the following parameters: potential mantle temperature of the
705 Jeesiörova-Kevitsa komatiite mantle source $T_{pot} = 1763^{\circ}\text{C}$, liquidus temperature upon
706 emplacement $T_{liq} = 1544^{\circ}\text{C}$, depth of melting initiation 294 km, which translates into a
707 pressure of 9.5 GPa. Based on these parameters, the temperatures and pressures at which
708 melting terminated for the Jeesiörova-Kevitsa komatiite magma was calculated using the
709 model of Herzberg et al. (2010) to be 1730°C and 5.0 GPa. Using these parameters, the
710 chemical composition of the original Jeesiörova-Kevitsa komatiite magma from Table 8, and
711 Eq. 4 of Steenstra et al. (2020), the SCSS of the Jeesiörova-Kevitsa komatiite magma upon
712 separation from the melting residue and ensuing emplacement are calculated to be 2270 and
713 5200 ppm, respectively.

714 In order to estimate the S content in the Jeesiörova-Kevitsa komatiite magma and, thus, its
715 sulfide saturation status, knowledge of the sulfur content of its mantle source, the S partition
716 coefficient, and the degree of melting is required. The average S content of modern DMM is
717 estimated to be 206 ± 25 ppm (Sun et al., 2020). Since the Jeesiörova-Kevitsa komatiite source
718 was as strongly melt-depleted as contemporary DMM, we assume it also had a similar S
719 content.

720 To estimate the degree of partial melting that produced the original Jeesiörova-Kevitsa
721 komatiite magma, we used a batch melting model and the calculated abundances of

moderately incompatible lithophile elements (e.g., Al, Ti, Gd) in this magma and its garnet-free mantle source of spinel lherzolite composition, as indicated by their Al-undepleted nature and the depth of melting initiation of ~290 km. These calculations yield 49% partial melting. Assuming that sulfur was highly incompatible during komatiite melting, with the degree of incompatibility similar to that of La, the original Jeesiörova-Kevitsa komatiite magma is calculated to have contained 420 ppm S. This S content is, thus, ~5 times lower than the SCSS of the Jeesiörova-Kevitsa komatiite magma upon separation from the melting residue and ~12 times lower than the SCSS of the Jeesiörova-Kevitsa komatiite magma upon emplacement, indicating that it was strongly sulfur-undersaturated upon separating from the source and remained as such all the way to the surface.

The sulfur-undersaturated nature of the Jeesiörova-Kevitsa komatiite magma upon emplacement is also attested to by the incompatible behavior of the highly chalcophile element Cu during magma differentiation after emplacement. The Cu abundances in the Kevitsa komatiites strongly correlate with the MgO content and plot on an olivine control line in the MgO versus Cu diagram (**Fig. 4a**), indicating that sulfide liquid was not a fractionating phase within the compositional range represented by the samples of this study.

To estimate the abundances of incompatible Pt and Pd in the Jeesiörova-Kevitsa komatiite source, we extrapolated the differentiation trends for each of these elements to a mantle MgO = 38 wt.% using ISOPLOT. The concentrations of Pt and Pd in the source were, thus, calculated to be 9.54 ± 0.22 and 8.13 ± 0.19 ppb, respectively, and the total Pt and Pd abundances to be 17.7 ± 0.8 ppb, or $120 \pm 5\%$ of those in the average estimates for modern BSE of 14.7 ± 2.0 ppb (Becker et al., 2006).

The concentration of compatible Os in the source was estimated to be 3.4 ± 0.1 ppb using the calculated Pt content in the source and the long-term $^{190}\text{Pt}/^{188}\text{Os}$ derived from the initial $^{186}\text{Os}/^{188}\text{Os}$. Further, using the calculated Os concentration and the estimated long-term $^{187}\text{Re}/^{188}\text{Os}$ derived from the initial $^{187}\text{Os}/^{188}\text{Os}$, the Re concentration in the source is calculated to be 0.285 ± 0.008 ppb. Unlike the Pt and Pd concentrations, therefore, the calculated Os and Re concentrations in the source of the Jeesiörova-Kevitsa komatiites are, respectively, 13% and 19% lower than the average estimates for BSE.

The presence of residual Os/Ir alloys in the melting residue, as evidenced by the compatible behavior of Os and Ir during the Jeesiörova-Kevitsa magma differentiation, makes it possible for the Os/Ir ratio to fractionate between the source and the komatiitic magma. Furthermore, as discussed earlier, chromite fractionation likely had a strong effect on Ru concentrations in the Jeesiörova-Kevitsa komatiite magma. Based on these observations, we

were unable to constrain the abundances of Ir and Ru in the source with any meaningful degree of accuracy.

The calculated total Pt and Pd abundances in the Jeesiörova-Kevitsa komatiite source are also plotted as a function of age in **Fig. 11** and compared with the data for other Archean and Paleoproterozoic komatiite systems studied to date and the BSE estimate. The calculated total Pt and Pd abundances in the source of the Jeesiörova-Kevitsa komatiites are significantly higher than in any other komatiite system examined so far. The significance of this observation will be discussed in the following sections.

6.4. The siderophile and lithophile element isotope conundrum

There is an apparent decoupling between the radiogenic lithophile trace element isotope systematics (e.g., Sm-Nd and Lu-Hf) and Re-Os isotope systematics in komatiite mantle sources worldwide. Many komatiite systems, such as those from the Abitibi (Canada) and Belingwe (South Africa) greenstone belts (Puchtel et al., 2009b; Puchtel et al., 2018), the Weltevreden and Komati (South Africa) Formations (Puchtel et al., 2014), and the Sumozero-Kenozero (Fennoscandia) greenstone belt (Puchtel et al., 2007), are characterized by strong, long-term depletions in incompatible lithophile trace elements, e.g., Nd versus Sm, but chondritic Re/Os. By contrast, certain other komatiite systems, such as the Kostomuksha in Fennoscandia (Puchtel et al., 2005) and the Schapenburg in South Africa (Puchtel et al., 2016a), while being long-term depleted in light REE, evolved with long-term suprachondritic Re/Os ratios. In this study, we address this apparent conundrum using the Sm-Nd, Lu-Hf, and Re-Os isotope and lithophile trace element abundance data for the Jeesiörova-Kevitsa komatiite system.

In Section 6.2, we calculated that the mantle source of the Jeesiörova-Kevitsa komatiites had the initial $\epsilon^{143}\text{Nd} = +4.9$ and $\epsilon^{176}\text{Hf} = +10.2$. This source was similar to DMM at 2.05 Ga (Goldstein and Jacobsen, 1988) and had long-term coupled Sm/Nd and Lu/Hf isotope systematics (Blichert-Toft and Puchtel, 2010). If the source had originally chondritic Sm/Nd and Lu/Hf, then it experienced a melt-extraction event a minimum of several hundred million years prior to komatiite emplacement. The same melt extraction event might also have been expected to result in depletion of incompatible Re relative to compatible Os, as well as more modest depletion in Pt relative to Os. Over time, these chemical effects would have led to the development of negative $\gamma^{187}\text{Os}$ and $\mu^{186}\text{Os}$ values. Instead, as shown in Section 6.3, the

Jeesiörova-Kevitsa komatiite source evolved within the range of chondritic Re/Os and suprachondritic Pt/Os.

6.5. Tungsten isotopic composition of the Jeesiörova-Kevitsa komatiite source

The Paleoproterozoic mantle was recently shown to have been characterized by an oxidation state similar to that of the modern mantle (Nicklas et al., 2019). As a result, during melting of the Paleoproterozoic mantle, W likely was characterized by a degree of incompatibility similar to that of Th and U (Newsom et al., 1996; Arevalo and McDonough, 2008; König *et al.*, 2011). Therefore, the BSE-normalized W abundance in the emplaced Jeesiörova-Kevitsa komatiite magma is expected to be similar to those of Th and U. This is the case, with an average calculated $W/W^* = 1.0$. This also implies that W, similarly to Th and U, was characterized by immobile behavior during the limited postmagmatic alteration of the Jeesiörova-Kevitsa komatiites. Further, similarly to the other lithophile trace elements, such as Nd and Hf, upper crustal rocks have significantly higher W abundances than komatiites; they also likely had a different W isotopic composition compared to komatiites, and, therefore, addition of even a small amount of upper crustal material can significantly change both the W isotopic composition and W abundances of a komatiite magma.

In this study, we used the measured W isotopic composition and W abundance of the emplaced Jeesiörova-Kevitsa komatiite magma to calculate those of the original, non-contaminated komatiite magma. In order to remove the effect of the 1% crustal contamination estimated on the basis of the lithophile trace element systematics, we used the W isotopic composition and W abundances of the average FTA from Puchtel et al. (2016b). The endmember parameters and the results of the mixing calculations are given in **Table 8**. The calculations suggest that the original Jeesiörova-Kevitsa komatiite magma contained ~5.8 ppb W, implying that ~50% of the total W budget in the emplaced Jeesiörova-Kevitsa komatiite magma was derived from the crustal contaminant.

The average $\mu^{182}\text{W}$ value of the FTA reported by Puchtel et al. (2016b) was $+12.6 \pm 4.5$, so contamination of the original komatiite magma with W-rich crust would have led to an increase in the $\mu^{182}\text{W}$ value of the original magma. If the assumptions regarding the original uncontaminated komatiite magma concentration of W, and concentration and ^{182}W isotopic composition of the crustal contaminant are accurate, the W isotopic composition of the original Jeesiörova-Kevitsa komatiite magma and, by extension, of its mantle source, after correction for crustal contamination, is calculated to be $\mu^{182}\text{W} = -10 \pm 5$. Here, the uncertainty

solely reflects propagated measurement uncertainties for the komatiite and the FTA. Because it is a projected original melt composition, this $\mu^{182}\text{W}$ must be considered tentative. Nevertheless, negative $\mu^{182}\text{W}$ values of this magnitude are common in modern ocean island basalt systems (Rizo et al., 2019; Mundl-Petermeier et al., 2019; 2020) and have also been observed in the 3.6 Ga Schapenburg komatiites (Puchtel et al., 2016a), South Africa, and some spatially associated, Archean glacial diamictites (Mundl et al., 2018). By contrast, the majority of evolved Archean crustal rocks similar to those envisioned for the FTA contaminant, are characterized by positive ^{182}W anomalies (e.g., Touboul et al., 2014; Willbold et al., 2015; Rizo et al., 2016b).

6.6. Reconciling isotopic characteristics of the Jeiesiörova-Kevitsa komatiites

The combined chemical and isotopic data for the Jeiesiörova-Kevitsa komatiites are complex. The Nd and Hf isotopic data are consistent with derivation from a mantle domain that experienced at least one melting event prior to the one that produced the komatiitic magma. The Nd-Hf isotope characteristics of this mantle domain are similar to projections of the modern DMM to 2049 Ma. The ^{187}Os isotopic data indicate derivation from a broadly chondritic mantle source that was not strongly long-term melt depleted. Because the long-term effects of low degrees of partial melting on Re/Os, compared to Sm/Nd and Lu/Hf, are relatively small (Puchtel et al., 2004b; 2009b), the chondritic $^{187}\text{Os}/^{188}\text{Os}$ of the mantle source can be considered at least qualitatively consistent with the Nd and Hf isotopic data. By contrast, the fractionated Pt/Os of the mantle source, as revealed by the suprachondritic initial $^{186}\text{Os}/^{188}\text{Os}$, and the correspondingly fractionated HSE pattern for the source, requires an event that could have strongly modified HSE relative abundances early in Earth history. Finally, if the negative $\mu^{182}\text{W}$ value for the Jeiesiörova-Kevitsa komatiite system obtained after correction for crustal contamination is accurate, it requires invoking one of a limited number of possible processes to account for the projected anomalous composition for the Hf-W short-lived radiogenic isotope system.

To address these complexities, we consider two melting models. The parameters used in both models are as follows. (1) The mantle domain that ultimately became the source of the Jeiesiörova-Kevitsa komatiites initially had a composition consistent with that of the BSE. The chemical and isotopic parameters of this domain were compiled from Jacobsen and Wasserburg (1980), Hamilton et al. (1983), Bouvier et al. (2008), Hofmann (1988), Shirey and Walker (1998), Brandon et al. (2006), and Becker et al. (2006). (2) This domain had a

modal mineral composition typical of spinel lherzolite, with 65% olivine, 24% orthopyroxene, 10% clinopyroxene, and 1% spinel. (3) The melting of the domain was modeled using the batch melting equation of Shaw (1970). The bulk D^{Sm} , D^{Nd} , D^{Lu} , and D^{Hf} during melting were calculated to be 0.040, 0.024, 0.090, and 0.037, respectively, based on the above source modal composition and crystal-liquid D values for the respective mineral phases from Green (1994). The bulk D^{Re} and D^{Pt} under the redox conditions of $\Delta\text{FMQ} +0.90\pm0.33$ determined by Nicklas et al. (2019) were estimated to be 0.20 and 2.0, respectively (Rehkämper et al., 1999; Mallmann and O'Neill, 2007). The bulk D^{Os} for the low-degree partial melting, consistent with formation of a MORB-type magma, was adopted to be 50 (e.g., Gannoun et al., 2016 and references therein).

In Model 1, we consider an early, global differentiation event. This model requires a minimum degree of melting and a melt extraction that would lead to fractionation of Sm/Nd and Lu/Hf in the BSE domain from the chondritic value to that required to bring the $\varepsilon^{143}\text{Nd}$ and $\varepsilon^{176}\text{Hf}$ of the domain to the corrected values of +4.9 and +10.2, respectively, by the time of komatiite formation at 2049 Ma. The effect of this melt extraction event on the Re-Os and Pt-Os isotope systems is then considered. Model 2 considers the youngest-possible, maximum degree of melting event that would fractionate the Sm/Nd and Lu/Hf ratios in the BSE domain from the chondritic values to those calculated for the Jeesiörova-Kevitsa komatiite *source*, and again bring the $\varepsilon^{143}\text{Nd}$ and $\varepsilon^{176}\text{Hf}$ values of this domain to +4.9 and +10.2, respectively, by 2049 Ma. The effect of this melt extraction event on the Re-Os and Pt-Os isotope systems is then also considered.

For Model 1, we assume that the global differentiation event occurred 100 Ma after Solar System formation. This timing is consistent with the recent data on the early formation of the Moon at ~4.51 Ga (Barboni et al., 2017). The Jeesiörova-Kevitsa komatiite source would have evolved from chondritic ratios of $^{143}\text{Nd}/^{144}\text{Nd} = 0.506807$ and $^{176}\text{Hf}/^{177}\text{Hf} = 0.279863$ at 4.467 Ga to the initial ratios of $^{143}\text{Nd}/^{144}\text{Nd} = 0.510236\pm8$ and $^{176}\text{Hf}/^{177}\text{Hf} = 0.281762\pm18$ at 2049 Ma with long-term ratios of $^{147}\text{Sm}/^{144}\text{Nd} = 0.2123\pm5$ and $^{176}\text{Lu}/^{177}\text{Hf} = 0.0394\pm4$. Such modest change in $^{147}\text{Sm}/^{144}\text{Nd}$ and $^{176}\text{Lu}/^{177}\text{Hf}$ from the chondritic ratios of 0.1967 and 0.0336, respectively, would require only ~0.6% melting, followed by removal of the melt. At the same time, this melt extraction event would have decreased the $^{187}\text{Re}/^{188}\text{Os}$ in the melting residue from 0.4019 to 0.3912; the latter ratio is essentially identical to the calculated long-term $^{187}\text{Re}/^{188}\text{Os} = 0.397\pm8$ determined for the Jeesiörova-Kevitsa komatiite source. However,

such low-degree melting event would not have had any measurable effect on the Pt/Os ratio of the source. Therefore, Model 1 cannot explain the observed $^{186}\text{Os}/^{188}\text{Os}$ composition of the Jeesiörova-Kevitsa komatiite system.

For Model 2, using the lithophile trace element data, we first calculate the $^{147}\text{Sm}/^{144}\text{Nd}$ and $^{176}\text{Lu}/^{177}\text{Hf}$ ratios in the Jeesiörova-Kevitsa komatiite source to be 0.2912 and 0.0704, respectively. Modeling indicates that 9% melting would have generated a residual mantle domain with these Sm/Nd and Lu/Hf ratios. The latest time of the melting event required for the mantle domain with such Sm/Nd and Lu/Hf ratios to have evolved to the initial $\epsilon^{143}\text{Nd} = +4.9$ and $\epsilon^{176}\text{Hf} = +10.2$ by 2049 Ma is 2.45 Ga. Any event that occurred later than that would not have afforded sufficient time to develop such radiogenic $\epsilon^{143}\text{Nd}$ and $\epsilon^{176}\text{Hf}$ in the mantle domain. It is noteworthy that 2.45 Ga coincides with the ages of abundant mafic layered intrusions (Alapieti et al., 1990; Balashov et al., 1993; Amelin et al., 1995; Yang et al., 2016) and komatiitic lavas (Puchtel et al., 1997; 2016b) throughout the Fennoscandian Shield, indicating that large volumes of mafic magma were indeed produced in this part of the planet at that time.

Using this extent and time of melting, and applying a bulk $D^{\text{Re}} = 0.2$, model calculations indicate that the melting event that produced this light REE depletion would have reduced the $^{187}\text{Re}/^{188}\text{Os}$ ratio in the chondritic reservoir from 0.4019 to 0.2694 at 2.45 Ga. The resultant melt-depleted mantle domain, if allowed to evolve from 2.45 Ga until the time of the komatiite formation at 2049 Ma with this Re/Os ratio, would have developed a negative initial $\gamma^{187}\text{Os} = -0.8$, which is close to the observed value and well within the range of chondritic meteorites at 2049 Ma (**Fig. 10**). At the same time, this melting event would only have reduced the $^{190}\text{Pt}/^{188}\text{Os}$ ratio in the mantle domain by 0.5%. The 400 Ma of growth from 2.45 to 2.05 Ga with this slightly reduced Pt/Os ratio will decrease the $\mu^{186}\text{Os}$ value of the system by 0.4 ppm, not increase it by 29 ppm, as observed for the Jeesiörova-Kevitsa komatiite source. Hence, similarly to Model 1, Model 2 can account for the Nd, Hf, and ^{187}Os isotopic compositions of the komatiites, but it also fails for ^{186}Os .

One significant modification to Models 1 and 2, which would make it possible to account for all the observed isotopic and elemental data for the komatiites, is assuming early perturbation of the Pt/Os ratio in the komatiite source, without affecting Sm-Nd and Lu-Hf evolution, as well as not modifying Re-Os evolution outside of the chondritic range. The most parsimonious process that can achieve this perturbation is crystal-liquid fractionation of metal, given that fractional crystallization of metal can strongly fractionate Pt/Os, but only

modestly fractionates Re/Os, with solid metal-liquid metal bulk distribution coefficients in this order: $D_{Os} > D_{Re} \gg D_{Pt} > 1$ (e.g., Cook et al., 2004). Thus, addition to a mantle domain of evolved metal either through late accretion of a fragment of a differentiated planetesimal core, similar to the mechanism proposed by Fischer-Gödde et al. (2011), or addition of metal derived from the Earth's core, but partially crystallized at or near the core-mantle boundary, similar to the mechanism of core-mantle interaction proposed by Humayun (2011), can best achieve the Os isotopic requirements for the komatiites. Either process would have the added benefit of decreasing the $\mu^{182}W$ value of the mixed domain to a negative value, as suggested by the contamination-corrected $\mu^{182}W = -10$ for the komatiites.

Compilation of time-integrated $^{190}Pt/^{188}Os$ and $^{187}Re/^{188}Os$ ratios for IIAB, IIIAB, and IVA magmatic iron meteorites from Cook et al. (2004) and McCoy et al. (2011) reveals large, up to three orders of magnitude, variations in the $^{190}Pt/^{188}Os$ ratios that resulted from magmatic solid metal-liquid metal fractionation, accompanied by relatively small, about a factor of 2.5, variations in the $^{187}Re/^{188}Os$ ratios in the same magmatic systems. The $^{186,187}Os$ isotope systematics of the Jeziörova-Kevitsa komatiite source can be accounted for by mixing with a number of fractionated iron compositions. We performed mixing calculations for Os, Re, and Pt concentrations, as well as for W isotopic composition and W abundances, results of which are illustrated in **Fig. 12**. For the meteoritic endmember, we chose a low-Os group IIIAB iron meteorite, Tieraco Creek; similar results are also obtained for other low-Os irons from this group, e.g., Grant and Chupaderos. The mantle source endmember used in the mixing calculations was the BSE. The HSE and W abundance data and W isotopic composition for the endmembers were adopted from Cook et al. (2004), Becker et al. (2006), Arevalo and McDonough (2008), Wasson (1999), and Kruijer et al. (2017); these are presented in **Table 9**. Based on the results of calculations, addition of 0.2% of material with the composition of Tieraco Creek to a mantle domain of BSE composition can explain the HSE and W abundances and Os and W isotopic compositions of the mantle domain that gave rise to the Jeziörova-Kevitsa komatiite system. One caveat to this model is that the mixing event must have occurred very early, within the first 100 Ma of Solar System history, while late accretion of substantial mass was still possible.

Addition of fractionated metal to the Moon (and presumably Earth, too) in the form of a late accretionary component was advocated by Fischer-Gödde and Becker (2012) in their study of lunar impact melt breccias to explain suprachondritic Pt/Ir, Ru/Ir, and Pd/Ir ratios in some of the impact melts. It has also been argued that the impactors that created the largest impact basins on the Moon and Mars, and presumably Earth as well, were hundreds of km in

diameter (Ryder, 2002; Strom et al., 2005; Bottke et al., 2010) and represented by differentiated planetesimals and, thus, would be capable of generating large HSE and negative ^{182}W heterogeneities in portions of the mantle (Marchi et al., 2018, 2020). Cores of differentiated planetesimals have up to two orders of magnitude higher W abundances and up to 350 ppm less radiogenic $^{182}\text{W}/^{184}\text{W}$ than the modern BSE (Kleine et al., 2002, 2004; Schoenberg et al., 2002; Yin et al., 2002). Thus, any mantle domain to which an excess of this material was added would be ^{182}W -depleted. Such a mantle domain would also be expected to be enriched in HSE, compared to BSE, particularly in Pt and Pd.

In order to further evaluate the effect of addition of such material on $^{182}\text{W}/^{184}\text{W}$ and HSE abundances, we plot the $\mu^{182}\text{W}$ in the Jezirova-Kevitsa komatiite system versus the total calculated HSE abundances in its mantle source relative to those in the estimates for modern BSE (**Fig. 13**). This proportion corresponds to the fraction of the total HSE budget of the BSE added during late accretion assuming an HSE-free mantle following core formation. The $^{182}\text{W}/^{184}\text{W}$ of the BSE prior to late accretion is constrained by mass-balance calculations (e.g., Kleine and Walker, 2017) and is supported by the $^{182}\text{W}/^{184}\text{W}$ data for the lunar mantle of $+25\pm 5$ ppm (Kruijer et al., 2015; Touboul et al., 2015; Kruijer and Kleine, 2017). Calculations indicate that the estimated 10 ppm deficit in $\mu^{182}\text{W}$ is consistent with the calculated total HSE abundances in the source of the Jezirova-Kevitsa komatiite system and the results of the mixing calculations presented above.

A second type of processes to consider is interaction between Earth's core and mantle. This type of process was originally proposed by Walker et al. (1995, 1997) and Brandon et al. (1998, 1999) as a means of generating coupled $^{186,187}\text{Os}$ enrichments observed in some ocean island basalts. Later, Humayun (2011) proposed a mechanism of core-mantle interaction that involved outer core liquid becoming trapped in a cumulate pile of Fe-rich non-metallic precipitates (FeO, FeS, Fe₃Si) at the top of the core. This metal is presumed to undergo partial fractional crystallization. The resulting, chemically-evolved liquid is then incorporated into the base of the mantle, where mantle plumes subsequently entrain it. If this metal had a composition similar to that of evolved iron meteorites, such as Tieraco Creek, its entrainment into the plume that gave rise to the Jezirova-Kevitsa komatiites could explain the $^{186,187}\text{Os}$ isotope systematics of these komatiites. As with the grainy late accretion model, this mechanism has the added benefit of explaining the W isotope systematics of the komatiites. Based on the discovery of negative $\mu^{182}\text{W}$ values in modern ocean island basalts (Mundl et al., 2017), Rizo et al. (2019) and Mundl-Petermeier et al. (2020) have argued for some form

of core-mantle exchange/equilibration in order to transfer the inferred negative $\mu^{182}\text{W}$ value of the core ($\mu^{182}\text{W} = -220$) to plumes rising from the core-mantle boundary. Although ^{186}Os - ^{182}W data for the same rocks are still limited, all Hawaiian lavas with negative $\mu^{182}\text{W}$ (ranging between -7.8 and -20.2) are also characterized by positive $\mu^{186}\text{Os}$ (ranging between +18.6 and +62.8) and $\gamma^{187}\text{Os}$ (ranging between +2.0 and +7.2) values (**Fig. 14**). The projected negative $\mu^{182}\text{W}$ value for the Jeesiörova-Kevitsa komatiites is within the range of $\mu^{182}\text{W}$ values for Hawaii, implying that these komatiites could be a Proterozoic equivalent of Hawaiian picrites. This process would be expected to have decreased $^{182}\text{W}/^{184}\text{W}$ in the komatiite source relative to the ambient mantle, and also potentially have increased the HSE abundances over the ambient mantle levels, which is consistent with the observations.

We conclude that at present, both grainy late accretion and core-mantle interaction remain viable mechanisms for explaining the observed combined lithophile trace- and siderophile element isotope and abundance systematics of the Jeesiörova-Kevitsa komatiite system.

7. Conclusions

The Jeesiörova-Kevitsa komatiites of the Fennoscandian Shield are unique among the komatiite systems studied so far. They have a relatively young age of 2.05 Ga that has not been reported for the existing terrestrial komatiite record elsewhere, yet are as MgO-rich as some of their late Archean counterparts. These komatiites were derived from a contemporary DMM-like mantle source in terms of Nd-Hf isotope systematics; this source was characterized by a negative ^{182}W anomaly, total absolute HSE abundances ~20% higher than of the modern BSE, and chondritic initial $^{187}\text{Os}/^{188}\text{Os}$, but suprachondritic initial $^{186}\text{Os}/^{188}\text{Os}$ isotopic compositions. When considered together, these constraints are reconciled within a model that requires derivation of the parental Jeesiörova-Kevitsa komatiite magma from a mantle domain that experienced a melting and melt depletion event within the first 100 Ma of Solar System history. This mantle domain is argued here to have acquired its HSE and Os isotope systematics as a result of either (1) grainy late accretion of differentiated metallic impactors or (2) core-mantle interaction via addition of chemically fractionated outer core metal to the mantle source domain of the komatiites. The survival of the early-formed Jeesiörova-Kevitsa komatiite mantle source for ≥ 2.5 billion years implies that portions of the mantle remained poorly mixed with respect to HSE and W until at least the Paleoproterozoic.

Acknowledgements

This study was supported by NSF Petrology and Geochemistry Grants EAR 1447174 and EAR 1754186 to ISP, EAR-1624587 to RJW and AMP, and Academy of Finland Grant 281859 to EJH. AMP would like to acknowledge support from FWF grant V659-N29. We thank V.A. Puchtel and R.W. Nicklas for help with preparation of samples for trace element analysis and ICP-MS measurements. Thorough and constructive reviews by Klaus Mezger, David van Acken, and an anonymous reviewer helped improve the original version of the manuscript. We thank Catherine Chauvel for comments and editorial handling.

References

- Alapieti, T.T., Filen, B.A., Lahtinen, J.J., Lavrov, M.M., Smolkin, V.F., and Voitsekhovskiy, S.N., 1990. Early Proterozoic layered intrusions in the northeastern part of the Fennoscandian Shield. *Mineralogy and Petrology* **42**: 1-22.
- Alard, O., Griffin, W. L., Lorand, J.-P., Jackson, S. E., O'Reilly, S. Y., 2000. Non-chondritic distribution of the highly siderophile elements in mantle sulfides. *Nature* **407** (6806): 891-894.
- Albarède, F., Blichert-Toft, J., Vervoort, J. D., Gleason, J. D., Rosing, M., 2000. Hf-Nd isotope evidence for a transient dynamic regime in the early terrestrial mantle. *Nature* **404** (6777): 488-490.
- Amelin, Y.V., Heaman, L.M., and Semenov, V.S., 1995. U-Pb geochronology of layered mafic intrusions in the eastern Baltic Shield: implications for the timing and duration of Paleoproterozoic continental rifting. *Precambrian Research* **75**(1-2): 31-46.
- Archer, G. J., Mundl, A., Walker, R. J., Worsham, E. A., and Bermingham, K. R., 2017. High-precision analysis of $^{182}\text{W}/^{184}\text{W}$ and $^{183}\text{W}/^{184}\text{W}$ by negative thermal ionization mass spectrometry: Per-integration oxide corrections using measured $^{18}\text{O}/^{16}\text{O}$. *International Journal of mass-spectrometry* **414**: 80-86.
- Archer, G. J., Ash, R. D., Bullock, E. S., and Walker, R. J., 2014. Highly siderophile elements and ^{187}Re - ^{187}Os isotopic systematics of the Allende meteorite: Evidence for primary nebular processes and late-stage alteration. *Geochimica et Cosmochimica Acta* **131**: 402-414.
- Archer, G. J., Brennecka, G. A., Gleißner, P., Stracke, A., Becker, H., and Kleine, T., 2019. Lack of late-accreted material as the origin of ^{182}W excesses in the Archean mantle: Evidence from the Pilbara Craton, Western Australia. *Earth and Planetary Science Letters* **528**: 115841.
- Arevalo, R. and McDonough, W. F., 2008. Tungsten geochemistry and implications for understanding the Earth's interior. *Earth and Planetary Science Letters* **272** (3-4): 656-665.
- Balashov, Y. A., Bayanova, T. B., and Mitrofanov, F. P., 1993. Isotope data on the age and genesis of layered basic-ultrabasic intrusions in the Kola Peninsula and northern Karelia, northeastern Baltic Shield. *Precambrian Research* **64**(1-4): 197-205.
- Barboni, M., Boehnke, P., Keller, B., Kohl, I. E., Schoene, B., Young, E. D., and McKeegan, K. D., 2017. Early formation of the Moon 4.51 billion years ago. *Science Advances* **3** (1): e1602365.
- Barnes, S.-J., Naldrett, A. J., Gorton, M. P., 1985. The origin of the fractionation of platinum-group elements in terrestrial magmas. *Chemical Geology* **53** (3-4): 303-323.
- Barnes, S.-J., Often, M., 1990. Ti-rich komatiites from Northern Norway. *Contributions to Mineralogy and Petrology* **105** (1): 42-54.
- Barnes, S. J., Fiorentini, M. L., 2008. Iridium, ruthenium and rhodium in komatiites: Evidence for iridium alloy saturation. *Chemical Geology* **257** (1-2): 44-58.
- Becker, H., Horan, M. F., Walker, R. J., Gao, S., Lorand, J.-P., Rudnick, R. L., 2006. Highly siderophile element composition of the Earth's primitive upper mantle: Constraints from new data on peridotite massifs and xenoliths. *Geochimica et Cosmochimica Acta* **70** (17): 4528-4550.
- Begemann, F., Ludwig, K.R., Lugmair, G.W., Min., K., Nyquist, L.E., Patchett, P.J., Renne, P.R., Shih, C.-Y., Villa, I.M., and Walker, R.J., 2001. Call for an improved set of decay constants for geochronological use. *Geochimica et Cosmochimica Acta* **65**(1): 111-121.
- Blichert-Toft, J., Chauvel, C., and Albarède, F., 1997. Separation of Hf and Lu for high-precision isotope analysis of rock samples by magnetic sector multiple collector ICP-MS. *Contributions to Mineralogy and Petrology* **127**(3): 248-260.

- Blichert-Toft, J., 2001. On the Lu-Hf isotope geochemistry of silicate rocks. *Geostandards Newsletter* **25**, 41-56.
- Blichert-Toft, J., Puchtel, I. S., 2010. Depleted mantle sources through time: Evidence from Lu-Hf and Sm-Nd isotope systematics of Archean komatiites. *Earth and Planetary Science Letters* **297** (3-4): 598-606.
- Bottke, W. F., Walker, R. J., Day, J. M. D., Nesvorny, D., Elkins-Tanton, L., 2010. Stochastic Late Accretion to Earth, the Moon, and Mars. *Science* **330** (6010): 1527-1530.
- Boyet, M., Carlson, R. W., 2005. ^{142}Nd evidence for early (> 4.53 Ga) global differentiation of the silicate Earth. *Science* **309** (5734): 576-581.
- Brandon, A. D., Walker, R. J., Morgan, J. W., Norman, M. D., and Prichard, H. M., 1998. Coupled ^{186}Os and ^{187}Os evidence for core-mantle interaction. *Science* **208** (5369): 1570-1573.
- Brandon, A. D., Norman, M. D., Walker, R. J., and Morgan, J. W., 1999. ^{186}Os - ^{187}Os systematics of Hawaiian picrites. *Earth and Planetary Science Letters* **174** (1-2): 25-42.
- Brandon, A. D., Humayun, M., Puchtel, I. S., Zolensky, M., 2005. Re-Os isotopic systematics and platinum group element composition of the Tagish Lake carbonaceous chondrite. *Geochimica et Cosmochimica Acta* **69** (6): 1619-1631.
- Brandon, A.D., Walker, R.J., and Puchtel, I.S., 2006. Platinum-osmium isotope evolution of the Earth's mantle: Constraints from chondrites and Os-rich alloys. *Geochimica et Cosmochimica Acta* **70**(8): 2093-2103.
- Brenan, J. M., 2008. Re-Os fractionation by sulfide melt-silicate melt partitioning: A new spin. *Chemical Geology* **248** (3-4): 140-165.
- Brügmann G. E., Arndt N. T., Hofmann A. W., and Tobschall H. J., 1987. Noble metal abundances in komatiite suites from Alexo, Ontario, and Gorgona Island, Colombia. *Geochimica et Cosmochimica Acta* **51** (8): 2159-2169.
- Chou, C.-L., Shaw, D. M., Crocket, J. H., 1983. Siderophile trace elements in the Earth's oceanic crust and upper mantle. *Journal of Geophysical Research* **88** (S2): A507-A518.
- Cook D. L., Walker R. J., Horan M. F., Wasson J. T., and Morgan J. W., 2004. Pt-Re-Os systematics of Group IIAB and IIIAB iron meteorites. *Geochimica et Cosmochimica Acta* **68** (6): 1413-1431.
- Creaser, R. A., Papanastassiou, D. A., Wasserburg, G. J., 1991. Negative thermal ion mass-spectrometry of osmium, rhenium, and iridium. *Geochimica et Cosmochimica Acta* **55** (1): 397-401.
- Ding, S., Dasgupta, R., and Tsuno, K., 2014. Sulfur concentration of martian basalts at sulfide saturation at high pressures and temperatures - Implications for deep sulfur cycle on Mars. *Geochimica et Cosmochimica Acta* **131**: 227-246.
- Ding, S., Hough, T., Dasgupta, R., 2018. New high pressure experiments on sulfide saturation of high-FeO*— basalts with variable TiO_2 contents - Implications for the sulfur inventory of the lunar interior. *Geochimica et Cosmochimica Acta* **222**: 319-339.
- Fischer-Gödde, M., Becker, H., Wombacher, F., 2010. Rhodium, gold and other highly siderophile element abundances in chondritic meteorites. *Geochimica et Cosmochimica Acta* **74** (1): 356-379.
- Fischer-Gödde M., Becker H., and Wombacher F., 2011. Rhodium, gold and other highly siderophile elements in orogenic peridotites and peridotite xenoliths. *Chemical Geology* **280** (3-4): 365-383.
- Fischer-Gödde M. and Becker H., 2012. Osmium isotope and highly siderophile element constraints on ages and nature of meteoritic components in ancient lunar impact rocks. *Geochimica et Cosmochimica Acta* **77**: 135-156.

- Fonseca, R. O. C., Laurenz, V., Mallmann, G., Luguet, A., Hoehne, N., Jochum, K. P., 2012. New constraints on the genesis and long-term stability of Os-rich alloys in the Earth's mantle. *Geochimica et Cosmochimica Acta* **87**: 227-242.
- Fonseca, R. O. C., Mallmann, G., O'Neill, H. S. C., Campbell, I. H., Laurenz, V., 2011. Solubility of Os and Ir in sulfide melt: Implications for Re/Os fractionation during mantle melting. *Earth and Planetary Science Letters* **311** (3-4): 339-350.
- Fortin, M.-A., Riddle, J., Desjardins-Langlais, Y., Baker, D. R., 2015. The effect of water on the sulfur concentration at sulfide saturation (SCSS) in natural melts. *Geochimica et Cosmochimica Acta* **160**: 100-116.
- Foster, J. G., Lambert, D. D., Frick, L. R., Maas, R., 1996. Re-Os isotopic evidence for genesis of Archaean nickel ores from uncontaminated komatiites. *Nature* **382** (6593): 703-706.
- Galer, S. J. G., Goldstein, S. L., 1991. Early mantle differentiation and its thermal consequences. *Geochimica et Cosmochimica Acta* **55** (1-2): 227-239.
- Gangopadhyay, A. and Walker, R. J., 2003. Re-Os systematics of the ca. 2.7 Ga komatiites from Alexo, Ontario, Canada. *Chemical Geology* **196** (1-4): 147-162.
- Gangopadhyay, A., Walker, R. J., Hanski, E., Solheid, P. A., 2006. Origin of Paleoproterozoic Ti-enriched komatiitic rocks from Jeiesjörova, Kittilä Greenstone Complex, Finnish Lapland. *Journal of Petrology* **47**, 773-789.
- Gannoun A., Burton K. W., Day J. M. D., Harvey J., Schiano P., and Parkinson I. J., 2016. Highly Siderophile Element and Os Isotope Systematics of Volcanic Rocks at Divergent and Convergent Plate Boundaries and in Intraplate Settings. *Reviews in Mineralogy and Geochemistry* **81**: 651-724.
- Gast, P.W., Triton, G.R., Hedge, C.E., 1964. Isotopic composition of lead and strontium from Ascension and Gough islands. *Science* **145**, 1181-1185.
- Goldstein, S. L., Galer, S. J. G., 1992. On the trail of early mantle differentiation: $^{142}\text{Nd}/^{144}\text{Nd}$ ratios of early Archean rocks. *Eos* **73** (30): 323.
- Goldstein, S. J. and Jacobsen, S. B., 1988. Nd and Sr isotopic systematics of river water suspended material: implications for crustal evolution. *Earth and Planetary Science Letters* **87** (3): 249-265.
- Green T. H., 1994. Experimental studies of trace-element partitioning applicable to igneous petrogenesis - Sedona 16 years later. *Chemical Geology* **117** (1-4): 1-36.
- Hamilton, P. J., O'Nions, R. K., Bridgwater, D., Nutman, A. P., 1983. Sm-Nd studies of Archaean metasediments and metavolcanics from West Greenland and their implications for the Earth's early history. *Earth and Planetary Science Letters* **62** (2): 263-272.
- Hanski, E., Huhma, H. 2005. Central Lapland Greenstone Belt. In: M. Lehtinen, P. Nurmi, O.T. Rämö (Eds.), *Precambrian Bedrock of Finland – Key to the Evolution of the Fennoscandian Shield*, Elsevier, Amsterdam, p. 139-194.
- Hanski, E.J., Kamenetsky, V.S., 2013. Chrome spinel-hosted melt inclusions in primitive Paleoproterozoic volcanic rocks, northern Finland: evidence for coexistence and mixing of komatiitic and picritic magmas. *Chemical Geology* **343**, 25-37. DOI: 10.1016/j.chemgeo.2013.02.009.
- Hanski, E., Huhma, H., Rastas, P., Kamenetsky, V. S., 2001. The Paleoproterozoic komatiite-picrite association of Finnish Lapland. *Journal of Petrology* **42** (5): 855-876.
- Hart, S. R., Brooks, S., 1977. The geochemistry and evolution of Early Precambrian mantle. *Contributions to Mineralogy and Petrology* **61** (2): 109-128.
- Herzberg, C., Condie, K., and Korenaga, J., 2010. Thermal history of the Earth and its petrological expression. *Earth and Planetary Science Letters* **292** (1-2): 79-88.
- Hofmann, A. W., 1984. Geochemical mantle models. *Terra Cognita* **4**: 157-165.

- Hofmann, A. W., 1988. Chemical differentiation of the Earth: The relationship between mantle, continental crust and oceanic crust. *Earth and Planetary Science Letters* **90** (3): 297-314.
- Hofmann, A.W., 1997. Mantle geochemistry: the message from oceanic volcanism. *Nature* **385** (6613): 219-229.
- Holzheid, A. and Grove, T. L., 2002. Sulfur saturation limits in silicate melts and their implications for core formation scenarios for terrestrial planets. *American Mineralogist* **87** (2-3): 227-237.
- Horan, M. F., Walker, R. J., Morgan, J. W., Grossman, J. N., Rubin, A. E., 2003. Highly siderophile elements in chondrites. *Chemical Geology* **196** (1-4): 5-20.
- Huhma H., Mänttari I., Peltonen P., Kontinen A., Halkoaho T., Hanski E., Hokkanen T., Hölttä P., Juopperi H., Konnunaho J., Layahe Y., Luukkonen E., Pietikäinen K., Pulkkinen A., Sorjonen-Ward P., Vaasjoki M., and Whitehouse M., 2012. The age of the Archean greenstone belts in Finland. In: *The Archaean of the Karelia Province in Finland*. Hölttä P. (Ed). Geological Survey of Finland Special Paper 54: 74-105.
- Huhma, H., Hanski, E., Kontinen, A., Vuollo, J., Mänttari, I., Lahaye, Y. 2018. Sm-Nd and U-Pb isotope geochemistry of the Palaeoproterozoic mafic magmatism in eastern and northern Finland. *Geological Survey of Finland, Bulletin* 405, 150 p.
- Humayun M., 2011. A model for osmium isotopic evolution of metallic solids at the core-mantle boundary. *Geochemistry, Geophysics, Geosystems* **12** (3): 1-23.
- Ireland T. J., Walker R. J., and Brandon A. D., 2011. ^{186}Os - ^{187}Os systematics of Hawaiian picrites revisited: New insights into Os isotopic variations in ocean island basalts. *Geochimica et Cosmochimica Acta* **75** (16): 4456-4475.
- Jacobsen, S. B., 1988. Isotopic and chemical constraints on mantle-crust evolution. *Geochimica et Cosmochimica Acta* **52** (6): 1341-1350.
- Jacobsen, S. B., Wasserburg, G. J., 1980. Sm-Nd isotopic evolution of chondrites. *Earth and Planetary Science Letters* **50** (1): 139-155.
- Jacobsen, S. B., Yu, G., 2015. Extinct isotope heterogeneities in the mantles of Earth and Mars: Implications for mantle stirring rates. *Meteoritics & Planetary Science* **50** (4): 555-567.
- Jochum, K. P., Arndt, N. T., and Hofmann, A. W., 1991. Nb-Th-La in komatiites and basalts: constraints on komatiite petrogenesis and mantle evolution. *Earth and Planetary Science Letters* **107** (2): 272-289.
- Jugo, P. J., Luth, R. W., and Richards, J. P., 2005. An Experimental Study of the Sulfur Content in Basaltic Melts Saturated with Immiscible Sulfide or Sulfate Liquids at 1300°C and 1.0 GPa. *Journal of Petrology* **46** (4): 783-798.
- Keays, R. R., 1995. The role of komatiitic and picritic magmatism and S-saturation in the formation of ore deposits. *Lithos* **34** (1-3): 1-18.
- Kimura, K., Lewis, R. S., Anders, S., 1974. Distribution of gold and rhenium between nickel-iron and silicate melts; implications for abundance of siderophile elements on the Earth and Moon. *Geochimica et Cosmochimica Acta* **38** (5): 683-701.
- Kiseeva, E. S. and Wood, B. J., 2013. A simple model for chalcophile element partitioning between sulphide and silicate liquids with geochemical applications. *Earth and Planetary Science Letters* **383**: 68-81.
- Kiseeva, E. S. and Wood, B. J., 2015. The effects of composition and temperature on chalcophile and lithophile element partitioning into magmatic sulphides. *Earth and Planetary Science Letters* **424**: 280-294.
- Kiseeva, E. S., Fonseca, R. O. C., and Smythe, D. J., 2017. Chalcophile Elements and Sulfides in the Upper Mantle. *Elements* **113**: 111-116.

- Kleine, T., Münker, C., Mezger, K., Palme, H., 2002. Rapid accretion and early core formation on asteroids and the terrestrial planets from Hf-W chronometry. *Nature* **418** (6901): 952-955.
- Kleine, T., Mezger, K., Münker, C., Palme, H., Bischoff, A., 2004. ^{182}Hf - ^{182}W isotope systematics of chondrites, eucrites, and martian meteorites: Chronology of core formation and early mantle differentiation in Vesta and Mars. *Geochimica et Cosmochimica Acta* **68** (13): 2935-2946.
- Kleine, T., Walker, R. J., 2017. Tungsten isotopes in planets. *Annual Review of Earth and Planetary Sciences* **45** (1): 389-417.
- König, S., Münker, C., Hohl, S., Paulick, H., Barth, A. R., Lagos, M., Pfander, J., Büchl, A., 2011. The Earth's tungsten budget during mantle melting and crust formation. *Geochimica et Cosmochimica Acta* **75** (8): 2119-2136.
- Kruijer, T. S., Kleine, T., 2017. Tungsten isotopes and the origin of the Moon. *Earth and Planetary Science Letters* **475** (Supplement C): 15-24.
- Kruijer, T. S., Kleine, T., Fischer-Gödde, M., Sprung, P., 2015. Lunar tungsten isotopic evidence for the late veneer. *Nature* **520** (7548): 534-537.
- Kruijer, T. S., Kleine, T., Borg, L. E., Brennecka, G. A., Irving, A. J., Bischoff, A., Agee, C. B., 2017. The early differentiation of Mars inferred from Hf-W chronometry. *Earth and Planetary Science Letters* **474** (Supplement C): 345-354.
- Li, J. and Agee, C. B., 2001. The effect of pressure, temperature, oxygen fugacity and composition on partitioning of nickel and cobalt between liquid Fe-Ni-S alloy and liquid silicate: implications for the earth's core formation. *Geochimica et Cosmochimica Acta* **65** (11): 1821-1832.
- Liu, Y., Samaha, N.-T., and Baker, D. R., 2007. Sulfur concentration at sulfide saturation (SCSS) in magmatic silicate melts. *Geochimica et Cosmochimica Acta* **71** (7): 1783-1799.
- Lorand, J.-P., Alard, O., 2001. Platinum-group element abundances in the upper mantle: new constraints from in situ and whole-rock analyses of Massif Central xenoliths (France). *Geochimica et Cosmochimica Acta* **65** (16): 2789-2806.
- Luguet, A., Shirey, S. B., Lorand, J.-P., Horan, M. F., Carlson, R. W., 2007. Residual platinum-group minerals from highly depleted harzburgites of the Lherz massif (France) and their role in HSE fractionation of the mantle. *Geochimica et Cosmochimica Acta* **71** (12): 3082-3097.
- Maier, W. D., Barnes, S. J., Campbell, I. H., Fiorentini, M. L., Peltonen, P., Barnes, S. J., Smithies, R. H., 2009. Progressive mixing of meteoritic veneer into the early Earth's deep mantle. *Nature* **460** (7255): 620-623.
- Mallmann, G. and O'Neill, H. St C., 2007. The effect of oxygen fugacity on the partitioning of Re between crystals and silicate melt during mantle melting. *Geochimica et Cosmochimica Acta* **71** (11): 2837-2857.
- Marchi, S., Canup, R. M., and Walker, R. J., 2018. Heterogeneous delivery of silicate and metal to the Earth by large planetesimals. *Nature Geoscience* **11**: 77-81.
- Marchi S., Walker, R. J., and Canup, R. M., 2020. A compositionally heterogeneous martian mantle due to late accretion. *Science Advances* **6** (7): eaay2338.
- Mavrogenes, J. A. and O'Neill, H. S. C., 1999. The relative effects of pressure, temperature and oxygen fugacity on the solubility of sulfide in mafic magmas. *Geochimica et Cosmochimica Acta* **63** (7-8): 1173-1180.
- McCoy, T. J., Walker, R. J., Goldstein, J. I., Yang, J., McDonough, W. F., Rumble, D., Chabot, N. L., Ash, R. D., Corrigan, C. M., Michael, J. R., and Kotula, P. G., 2011. Group IVA irons: New constraints on the crystallization and cooling history of an asteroidal core with a complex history. *Geochimica et Cosmochimica Acta* **75** (22): 6821-6843.

- McDonough, W. F. and Sun, S. S., 1995. The composition of the Earth. *Chemical Geology* **120** (3-4): 223-253.
- McKenzie, D. and Bickle, M. J., 1988. The volume and composition of melt generated by extension of the lithosphere. *Journal of Petrology* **29** (3): 625-679.
- Morgan, J. W., 1985. Osmium isotope constraints on Earth's late accretionary history. *Nature* **317** (6039): 703-705.
- Morgan, J. W., 1986. Ultramafic xenoliths: Clues to Earth's late accretionary history. *Journal of Geophysical Research* **91** (B12): 12375-12387.
- Mundl, A., Touboul, M., Jackson, M. G., Day, J. M. D., Kurz, M. D., Lekic, V., Helz, R. T., Walker, R. J., 2017. Tungsten-182 heterogeneity in modern ocean island basalts. *Science* **356** (6333): 66-69.
- Mundl, A., Walker, R. J., Reimink, J. R., Rudnick, R. L., Gaschnig, R. M., 2018. Tungsten-182 in the upper continental crust: Evidence from glacial diamictites. *Chemical Geology* **494**: 144-152.
- Mundl-Petermeier, A., Walker, R.J., Jackson, M.G., Blichert-Toft, J., Kurz, M.D., and Haldórsson, S.A. (2019) Temporal evolution of primordial tungsten-182 and $^3\text{He}/^4\text{He}$ signatures in the Iceland mantle plume. *Chemical Geology* **525**, 245-259.
- Mundl-Petermeier, A., Walker, R.J., Fischer, R.A., Lekic, V., Jackson, M.G., and Kurz, M.D., 2020. Anomalous $\mu^{182}\text{W}$ signatures in high $^3\text{He}/^4\text{He}$ ocean island basalts – fingerprints of Earth's core? *Geochimica et Cosmochimica Acta* **271**: 194-211.
- Mungall, J. E. and Naldrett, A. J., 2008. Ore deposits of the platinum-group elements. *Elements* **4** (4): 253-258.
- Mutanen, T. 1997. Geology and ore petrology of the Akanvaara and Koitelainen mafic layered intrusions and the Keivitsa-Satovaara layered complex, northern Finland. Geological Survey of Finland, Bulletin 395, 233 p.
- Mutanen, T. 2005. The Akanvaara intrusion and the Keivitsa–Satovaara Complex, with stops at Kaikkivaara and Särkivaara intrusions. Geological Survey of Finland, Guide 51b, 124 p.
- Mutanen, T., Huhma, H. 2001. U-Pb geochronology of the Koitelainen, Akanvaara and Keivitsa layered intrusions and related rocks. Geological Survey of Finland, Special Paper **33**: 229–246.
- Naldrett, A. J., 2010. Secular Variation of Magmatic Sulfide Deposits and Their Source Magmas. *Economic Geology* **105** (3): 669-688.
- Nesbitt, R. W., Sun, S. S., Purvis, A. C., 1979. Komatiites: geochemistry and genesis. *Canadian Mineralogist* **17** (2): 165-186.
- Newsom, H. E., Sims, K. W. W., Noll, P. D., Jaeger, W. L., Maehr, S. A., Beserra, T. B., 1996. The depletion of tungsten in the bulk silicate Earth: Constraints on core formation. *Earth and Planetary Science Letters* **60** (5): 1155-1169.
- Nicklas, R. W., Puchtel, I. S., and Ash, R. D., 2016. High-precision determination of the oxidation state of komatiite lavas using vanadium liquid-mineral partitioning. *Chemical Geology* **433**: 36-45.
- Nicklas R.W., Puchtel I.S., Ash R.D., Piccoli P., Hanski E., Nisbet E.G., Waterton P., Pearson D.G., and Anbar A.D., 2019. Secular Mantle Oxidation across the Archean-Proterozoic Boundary: Evidence from V partitioning in Komatiites and Picrites. *Geochimica et Cosmochimica Acta* **250**: 49-75.
- O'Neill, H. St C. and Mavrogenes, J. A., 2002. The Sulfide Capacity and the Sulfur Content at Sulfide Saturation of Silicate Melts at 1400°C and 1 bar. *Journal of Petrology* **43** (6): 1049-1087.
- Peters B. J., Mundl-Petermeier A., Horan M. F., Carlson R. W. and Walker R. J., 2019. Chemical Separation of Tungsten and Other Trace Elements for TIMS Isotope Ratio

- Measurements Using Organic Acids. *Geostandards and Geoanalytical Research* 43 (2): 245-259.
- Puchtel, I.S., Haase, K.M., Hofmann, A.W., Chauvel, C., Kulikov, V.S., Garbe-Schönberg, C.D., and Nemchin, A.A., 1997. Petrology and geochemistry of crustally contaminated komatiitic basalts from the Vetreny Belt, southeastern Baltic Shield: Evidence for an early Proterozoic mantle plume beneath rifted Archean continental lithosphere. *Geochimica et Cosmochimica Acta* **61** (6): 1205-1222.
- Puchtel I. S., Hofmann A. W., Mezger K., Jochum K. P., Shchipansky A. A., and Samsonov A. V., 1998. Oceanic plateau model for continental crustal growth in the Archaean: A case study from the Kostomuksha greenstone belt, NW Baltic Shield. *Earth and Planetary Science Letters* 155 (1-2): 57-74.
- Puchtel, I. S. and Humayun, M., 2000. Platinum group elements in Kostomuksha komatiites and basalts: Implications for oceanic crust recycling and core-mantle interaction. *Geochimica et Cosmochimica Acta* 64 (24): 4227-4242.
- Puchtel, I. S., Humayun, M., 2005. Highly siderophile element geochemistry of ^{187}Os -enriched 2.8-Ga Kostomuksha komatiites, Baltic Shield. *Geochimica et Cosmochimica Acta* **69** (6): 1607-1618.
- Puchtel, I. S., Brandon, A. D., Humayun, M., 2004a. Precise Pt-Re-Os isotope systematics of the mantle from 2.7-Ga komatiites. *Earth and Planetary Science Letters* **224** (1-2): 157-174.
- Puchtel, I. S., Humayun, M., Campbell, A., Sproule, R., Leshner, C. M., 2004b. Platinum group element geochemistry of komatiites from the Alexo and Pyke Hill areas, Ontario, Canada. *Geochimica et Cosmochimica Acta* **68** (6): 1361-1383.
- Puchtel, I. S., Brandon, A. D., Humayun, M., Walker, R. J., 2005. Evidence for the early differentiation of the core from Pt-Re-Os isotope systematics of 2.8-Ga komatiites. *Earth and Planetary Science Letters* **237** (1-2): 118-134.
- Puchtel, I. S., Humayun, M., Walker, R. J., 2007. Os-Pb-Nd isotope and highly siderophile and lithophile trace element systematics of komatiitic rocks from the Volotsk suite, SE Baltic Shield. *Precambrian Research* **158** (1-2): 119-137.
- Puchtel, I. S., Walker, R. J., Anhaeusser, C. R., Gruau, G., 2009a. Re-Os isotope systematics and HSE abundances of the 3.5 Ga Schapenburg komatiites, South Africa: Hydrous melting or prolonged survival of primordial heterogeneities in the mantle? *Chemical Geology* **262** (3-4): 355-369.
- Puchtel, I. S., Walker, R. J., Brandon, A. D., Nisbet, E. G., 2009b. Pt-Re-Os and Sm-Nd isotope and HSE and REE systematics of the 2.7 Ga Belingwe and Abitibi komatiites. *Geochimica et Cosmochimica Acta* **73** (20): 6367-6389.
- Puchtel, I. S., Blichert-Toft, J., Touboul, M., Walker, R. J., Byerly, G., Nisbet, E. G., Anhaeusser, C. R., 2013. Insights into early Earth from Barberton komatiites: Evidence from lithophile isotope and trace element systematics. *Geochimica et Cosmochimica Acta* **108**: 63-90.
- Puchtel, I. S., Walker, R. J., Touboul, M., Nisbet, E. G., Byerly, G. R., 2014. Insights into Early Earth from the Pt-Re-Os isotope and Highly Siderophile Element abundance systematics of Barberton komatiites. *Geochimica et Cosmochimica Acta* **125**: 394-413.
- Puchtel, I. S., Blichert-Toft, J., Touboul, M., Horan, M. F., Walker, R. J., 2016a. The coupled ^{182}W - ^{142}Nd record of early terrestrial mantle differentiation. *Geochemistry, Geophysics, Geosystems* **17** (6): 2168-2193.
- Puchtel, I. S., Touboul, M., Blichert-Toft, J., Walker, R. J., Brandon, A. D., Nicklas, R. W., Kulikov, V. S., Samsonov, A. V., 2016b. Lithophile and siderophile element systematics of the mantle at the Archean-Proterozoic boundary: Evidence from 2.4 Ga komatiites. *Geochimica et Cosmochimica Acta* **180**: 227-255.

- Puchtel, I. S., Blichert-Toft, J., Touboul, M., and Walker, R. J., 2018. ^{182}W and HSE constraints from 2.7 Ga komatiites on the heterogeneous nature of the Archean mantle. *Geochimica et Cosmochimica Acta* **228**: 1-26.
- Rehkämper M., Halliday A. N., Fitton J. G., Lee D.-C., Wieneke M., and Arndt N. T., 1999. Ir, Ru, Pt and Pd in basalts and komatiites: New constraints for the geochemical behavior of the platinum group elements in the mantle. *Geochimica et Cosmochimica Acta* **63** (22): 3915-3934.
- Rizo H., Andrault D., Bennett N. R., Humayun M., Brandon A. D., Vlastelic I., Moine B., Poirier A., Bouhifd M. A., and Murphy D. T., 2019. ^{182}W evidence for core-mantle interaction in the source of mantle plumes. *Geochemical Perspectives Letters* **11**: 6-11.
- Rizo, H., Walker, R. J., Carlson, R. W., Horan, M. F., Mukhopadhyay, S., Manthos, V., Francis, D., Jackson, M. G., 2016a. Preservation of Earth-forming events in the tungsten isotopic composition of modern flood basalts. *Science* **352** (6287): 809-812.
- Rizo, H., Walker, R. J., Carlson, R. W., Touboul, M., Horan, M. F., Puchtel, I. S., Boyet, M., Rosing, M. T., 2016b. Early Earth differentiation investigated through ^{142}Nd , ^{182}W , and highly siderophile element abundances in samples from Isua, Greenland. *Geochimica et Cosmochimica Acta* **175**: 319-336.
- Rudnick, R.L., and Fountain, D.M., 1995. Nature and composition of the continental crust: a lower crustal perspective. *Reviews of Geophysics* **33** (3): 267-309.
- Rudnick, R.L., and Gao, S., 2014. Composition of the Continental Crust. *Treatise on Geochemistry*, Macmillan Publishers Limited: 1-51.
- Ryder, G., 2002. Mass flux in the ancient Earth-Moon system and benign implications for the origin of life on Earth. *Journal of Geophysical Research* **107** (E4): 6.1-6.13.
- Saverikko, M. 1985. The pyroclastic komatiite complex at Sattasvaara in northern Finland. *Bulletin of the Geological Society of Finland* **57**, 55-87.
- Schoenberg, R., Kamber, B. S., Collerson, K. D., Eugster, O., 2002. New W-isotope evidence for rapid terrestrial accretion and very early core formation. *Geochimica et Cosmochimica Acta* **66** (17): 3151-3160.
- Shirey, S. B., Walker, R. J., 1998. The Re-Os isotope system in cosmochemistry and high-temperature geochemistry. *Annual Reviews of Earth and Planetary Sciences* **26**: 423-500.
- Smoliar, M. I., Walker, R. J., Morgan, J. W., 1996. Re-Os ages of Group IIA, IIIA, IVA, and IVB iron meteorites. *Science* **271** (5762): 1099-1102.
- Smythe, D. J., Wood, B. J., and Kiseeva, E. S., 2017. The S content of silicate melts at sulfide saturation: New experiments and a model incorporating the effects of sulfide composition. *American Mineralogist* **102** (4): 795-803.
- Sossi, P.A., O'Neill, H.St.C., 2016. Liquidus temperatures of komatiites and the effect of cooling rate on element partitioning between olivine and komatiitic melt. *Contributions to Mineralogy and Petrology* **171**: 49.
- Steenstra, E.S., Berndt, J., Klemme, S., Rohrbach, A., Bullock, E.S., van Westrenen, W., 2020. An experimental assessment of the potential of sulfide saturation of the source regions of eucrites and angrites: Implications for asteroidal models of core formation, late accretion and volatile element depletions. *Geochimica et Cosmochimica Acta* **269**: 39-62.
- Steenstra, E. S., Seegers, A. X., Eising, J., Tomassen, B. G. J., Webers, F. P. F., Berndt, J., Klemme, S., Matveev, S., van Westrenen, W., 2018. Evidence for a sulfur-undersaturated lunar interior from the solubility of sulfur in lunar melts and sulfide-silicate partitioning of siderophile elements. *Geochimica et Cosmochimica Acta* **231**: 130-156.
- Strom, R. G., Malhotra, R., Ito, T., Yoshida, F., and Kring, D. A., 2005. The origin of planetary impactors in the inner Solar system. *Science* **309** (5742): 1847-1850.
- Sun, Z., Xiong, X., Wang, J., Liu, X., Li, L., Ruan, M., Zhang, L., and Takahashi, E., 2020. Sulfur abundance and heterogeneity in the MORB mantle estimated by copper

- partitioning and sulfur solubility modelling. *Earth and Planetary Science Letters* **538**: 116169.
- Thompson, P. M. E., Kempton, P. D., White, R. V., Kerr, A. C., Tarney, J., Saunders, A. D., Fitton, J. G., and McBirney, A., 2003. Hf-Nd isotope constraints on the origin of the Cretaceous Caribbean plateau and its relationship to the Galapagos plume. *Earth and Planetary Science Letters* **217** (1-2): 59-75.
- Törmänen, T., Konnunaho, J., Hanski, E., Moilanen, M., Heikura, P., 2016. The Paleoproterozoic komatiite-hosted PGE mineralization at Lomalampi, Central Lapland Greenstone Belt, northern Finland. *Mineralium Deposita* **51**: 411-430.
- Touboul, M., Puchtel, I. S., Walker, R. J., 2012. ^{182}W Evidence for Long-Term Preservation of Early Mantle Differentiation Products. *Science* **335**: 1065-1069.
- Touboul, M., Liu, J., O'Neil, J., Puchtel, I. S., Walker, R. J., 2014. New Insights into the Hadean Mantle Revealed by ^{182}W and Highly Siderophile Element Abundances of Supracrustal Rocks from the Nuvvuagittuq Greenstone Belt, Quebec, Canada. *Chemical Geology* **383**: 63-75.
- Touboul, M., Puchtel, I. S., Walker, R. J., 2015. Tungsten isotopic evidence for disproportional late accretion to the Earth and Moon. *Nature* **520** (7548): 530-533.
- Tusch, J., Sprung, P., van de Löcht, J., Hoffmann, J. E., Boyd, A. J., Rosing, M. T., and Münker, C., 2019. Uniform ^{182}W isotope compositions in Eoarchean rocks from the Isua region, SW Greenland: The role of early silicate differentiation and missing late veneer. *Geochimica et Cosmochimica Acta* **257**: 284-310.
- van Acken, D., Brandon, A. D., and Humayun, M., 2011. High-precision osmium isotopes in enstatite and Rumuruti chondrites. *Geochimica et Cosmochimica Acta* **75** (14): 4020-4036.
- Walker, R. J., 2016. Siderophile Elements in Tracing Planetary Formation and Evolution. *Geochemical Perspectives* **5** (1): 1-145.
- Walker, R. J., Morgan, J. W., and Horan, M. F., 1995. ^{187}Os enrichment in some plumes: evidence for core-mantle interaction? *Science* **269** (5225): 819-822.
- Walker, R. J., Morgan, J. W., Beary, E. S., Smoliar, M. I., Czamanske, G. K., and Horan, M. F., 1997. Applications of the ^{190}Pt - ^{186}Os isotope system to geochemistry and cosmochemistry. *Geochimica et Cosmochimica Acta* **61** (22): 4799-4807.
- Walker, R. J., Shirey, S. B., and Stecher, O., 1988. Comparative Re-Os, Sm-Nd and Rb-Sr isotope and trace element systematics for Archean komatiite flows from Munro Township, Abitibi belt, Ontario. *Earth and Planetary Science Letters* **87** (1-2): 1-12.
- Walker, R. J., Horan, M. F., Morgan, J. W., Becker, H., Grossman, J. N., Rubin, A. E., 2002. Comparative ^{187}Re - ^{187}Os systematics of chondrites: Implications regarding early solar system processes. *Geochimica et Cosmochimica Acta* **66** (23): 4187-4201.
- Walker, R. J., Bermingham, K., Liu, J., Puchtel, I. S., Touboul, M., and Worsham, E. A., 2015. In search of late-stage planetary building blocks. *Chemical Geology* **411**: 125-142.
- Wasson, J. T., 1999. Trapped melt in IIIAB irons; solid/liquid elemental partitioning during the fractionation of the IIIAB magma. *Geochimica et Cosmochimica Acta* **63** (18): 2875-2889.
- White, W. M. and Hofmann, A. W., 1982. Sr and Nd isotope geochemistry of oceanic basalts and mantle evolution. *Nature* **296** (5860): 821-825.
- White, W. M. and Patchett, J., 1984. Hf-Nd-Sr isotopes and incompatible element abundances in island arcs: implications for magma origins and crust-mantle evolution. *Earth and Planetary Science Letters* **67** (2): 167-185.
- Willbold, M., Elliott, T., Moor bath, S., 2011. The tungsten isotopic composition of the Earth's mantle before the terminal bombardment. *Nature* **477** (7363): 195-198.

- Willbold, M., Mojzsis, S. J., Chen, H. W., Elliott, T., 2015. Tungsten isotope composition of the Acasta Gneiss Complex. *Earth and Planetary Science Letters* **419**: 168-177.
- Wood, B. J. and Kiseeva, E. S., 2015. Trace element partitioning into sulfide: How lithophile elements become chalcophile and vice versa. *American Mineralogist* **100** (11-12): 2371-2379.
- Yang, S.-H., Hanski, E., Li, C., Maier, W. D., Huhma, H., Mokrushin, A. V., Latypov, R., Lahaye, Y., O'Brien, H., and Qu, W.-J., 2016. Mantle source of the 2.44-2.50-Ga mantle plume-related magmatism in the Fennoscandian Shield: evidence from Os, Nd, and Sr isotope compositions of the Monchepluton and Kemi intrusions. *Mineralium Deposita* **51**: 1055-1073.
- Yin, Q., Jacobsen, S. B., Yamashita, K., Blichert-Toft, J., Telouk, P., Albarede, F., 2002. A short timescale for terrestrial planet formation from Hf-W chronometry of meteorites. *Nature* **418** (6901): 949-952.
- Zindler, A., Jagoutz, E., Goldstein, S. L., 1982. Nd, Sr and Pb isotopic systematics in a three-component mantle: a new perspective. *Nature* **298** (5874): 519-523.

Figure captions

Fig. 1. (a) Geological map showing the distribution of the 2.05 Ga komatiites and related picrites in the Central Lapland Greenstone Belt, Finland (modified after Hanski et al., 2001), and the locations of the Jeesiörova komatiites and the Kevitsa mafic-ultramafic intrusion. (b) Geological map of the Kevitsa intrusion (modified after Mutanen and Huhma, 2001). (c) Location of drill hole DDH 814 and other drill holes penetrating Kevitsa komatiite dikes (red dots).

Fig. 2. Photomicrographs of olivine-phyric Kevitsa komatiite dikes. (a) Sample showing various olivine morphologies. (b) Olivine phenocrysts in a groundmass composed of clinopyroxene, plagioclase, olivine, brown amphibole, and oxide. (c) Close-up view of the groundmass. (d) Large olivine phenocryst with ragged grain boundaries. (e) Thin olivine-phyric vein injected into olivine-pyroxene cumulate of the Kevitsa intrusion. (f) Chilled margin with small olivine and clinopyroxene phenocrysts set in a fine-grained groundmass. Am = brown amphibole, Pl = plagioclase, Ol = olivine, Cp = clinopyroxene. Plane-polarized light in all panels except for (c).

Fig. 3. Thin section scanner micrographs in transmitted light showing olivine megacrysts and phenocrysts in the Kevitsa komatiite. Note the large chromite grains attached to the megacryst in (a).

Fig. 4. Abundances of selected major (wt.%) and lithophile trace (ppm) elements and W (ppb) plotted against MgO (wt.%) in the Jeesiörova and Kevitsa komatiites.

Fig. 5. BSE-normalized abundances of lithophile trace elements in the Jeesiörova (a) and Kevitsa (b) komatiites arranged in order of decreasing incompatibility during mantle melting. The normalizing value for W is from Arevalo and McDonough (2008) and, for the other elements, from Hofmann (1988).

Fig. 6. (a) Re-Os isochron diagram for whole-rock samples and olivine and chromite separates from the Jeesiörova and Kevitsa komatiites analyzed in this study. (b) Pt-Os isotopic data for chromite separates from the Jeesiörova komatiites.

Fig. 7. CI chondrite-normalized HSE abundances in whole-rock samples of the (a) Jeesiörova komatiites, (b) Kevitsa komatiites, and (c) olivine separates from the Kevitsa komatiites.

Fig. 8. Abundances of HSE in the Jeesiörova and Kevitsa komatiites and olivine separates (in ppb) plotted against MgO contents (wt.%).

Fig. 9. (a) ^{147}Sm - ^{143}Nd and (b) ^{176}Lu - ^{176}Hf isochron diagrams for the Jeesiörova-Kevitsa komatiites and clinopyroxene separates. See text for additional details. (c) Diagram illustrating average initial $\epsilon^{143}\text{Nd}$ and $\epsilon^{176}\text{Hf}$ values for the emplaced Jeesiörova-Kevitsa komatiites (light-green symbol) and their mantle source (dark-green symbol), as well as for the Chondritic Uniform Reservoir (CHUR) and Depleted MORB Mantle (DMM) calculated at the time of the emplacement (2049 Ma). The CHUR and DMM parameters

are from Jacobsen and Wasserburg (1980), Bouvier *et al.* (2008), and Blichert-Toft and Puchtel (2010). Note the coupled, or congruent, behavior between the two isotope systems in the source of the Jeesiörova-Kevitsa komatiites, after correction of the komatiite data for 1.0% crustal contamination.

Fig. 10. (a) Initial $^{187}\text{Os}/^{188}\text{Os}$ isotopic compositions, expressed in terms of $\gamma^{187}\text{Os}$, of Archean and Paleoproterozoic komatiite systems plotted as a function of age. The data are from Foster *et al.* (1996); Puchtel *et al.* (2004a; 2005; 2007; 2009a; 2009b; 2014; 2016a; 2016b; 2018), and this study. The data for chondritic meteorites are compiled from Walker *et al.* (2002), Brandon *et al.* (2005), and Fischer-Gödde *et al.* (2010). **(b).** Initial $^{186}\text{Os}/^{188}\text{Os}$ isotopic compositions, expressed in terms of $\mu^{186}\text{Os}$, of Archean and Paleoproterozoic komatiite systems studied to date, plotted as a function of age. The data are from Puchtel *et al.* (2004a, 2005, 2009b, 2014, 2016b; 2018), and this study. The data for chondritic meteorites are compiled from Horan *et al.* (2003), Brandon *et al.* (2005, 2006), and Fischer-Gödde *et al.* (2010). See text for more details.

Fig. 11. Calculated total Pt and Pd abundances in the sources of Archean and Paleoproterozoic komatiite systems plotted as per cent of the total Pt and Pd abundances in the estimates for the modern BSE of Becker *et al.* (2006). Data sources are as follows: 2.05 Ga Jeesiörova-Kevitsa - this study; 2.41 Ga Vetreny Belt - Puchtel *et al.* (2016b); 2.69 Ga Belingwe - Puchtel *et al.* (2009b); 2.72 Ga Pyke Hill and Alexo - Puchtel *et al.* (2004b); 2.72 Ga Boston Creek - Puchtel *et al.* (2018); 2.82 Ga Kostomuksha - Puchtel and Humayun (2005); 2.88 Ga Volotsk-Kamennozero - Puchtel *et al.* (2007); 3.26 Ga Weltevreden and 3.48 Ga Komati - Puchtel *et al.* (2014); 3.55 Ga Schapenburg - Puchtel *et al.* (2016a). Uncertainties are 2SD. See text for additional details.

Fig. 12. Diagram illustrating potential genetic relationships between long-term $^{187}\text{Re}/^{188}\text{Os}$ and $^{190}\text{Pt}/^{188}\text{Os}$ in IIIAB fractionated iron meteorites and in the Jeesiörova-Kevitsa (J-K) komatiite source. J-K source 1 and 2 denote the calculated Re/Os and Pt/Os in the source after and before the global melt depletion event, respectively, that is assumed to have occurred within the first 100 Ma of Solar System history and is necessary to account for the Nd-Hf isotope systematics of the Jeesiörova-Kevitsa komatiites. The data for the IIIAB iron meteorite Tieraco Creek used in the modeling are from Cook *et al.* (2004). Addition of 0.2% of Tieraco Creek metal to the mantle domain with BSE Re/Os and Pt/Os *prior* to the global melt depletion event can explain the $^{186,187}\text{Os}/^{188}\text{Os}$ isotope systematics of the Jeesiörova-Kevitsa komatiite source. See text for more details.

Fig. 13. $\mu^{182}\text{W}$ (ppm) versus total calculated HSE abundances in the sources of Archean and Paleoproterozoic komatiite systems studied to date relative to those in the estimates for the present-day BSE of Becker *et al.* (2006). This proportion corresponds to the fraction of the total HSE budget of the BSE added during late accretion assuming an HSE-free terrestrial mantle prior to late accretion. The $\mu^{182}\text{W}$ value of the BSE prior to late accretion is constrained via mass-balance calculations and is supported by the W isotopic data for the lunar mantle ($\mu^{182}\text{W} = +25 \pm 5$ ppm: Kruijer *et al.*, 2015; Touboul *et al.*, 2015).

The W isotopic data and estimates of the HSE contents for the komatiite systems are from Puchtel and Humayun (2005), Touboul et al. (2012), Puchtel et al. (2014; 2016a; 2016b; 2018), and this study.

Fig. 14. $\mu^{182}\text{W}$ versus $\mu^{186}\text{Os}$ data (ppm) for the Jeesiörova-Kevitsa komatiites and Hawaiian picrites. The Os data for Hawaii are compiled from Brandon et al. (1998) and Ireland et al. (2011), and the W data - from Mundl et al. (2017) and Mundl-Petermeier et al. (2020).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 1. Major and minor element data for Jeesiörova and Kevitsa komatiites.

Sample	12D-PPR	13-EJH	17.1-PPR	LP10	KD03	KD05	KD06	KD07	KD08	KD09	KD10	KD11	KD12	KD13	KD14
SiO ₂	44.8	43.8	43.0	43.6	46.6	45.0	44.9	44.6	44.1	43.4	43.5	44.1	45.1	45.0	46.9
TiO ₂	0.691	0.632	0.681	0.583	0.730	0.612	0.628	0.533	0.554	0.488	0.487	0.582	0.662	0.679	0.778
Al ₂ O ₃	7.70	7.47	8.01	6.94	10.1	8.55	8.74	7.34	7.94	6.50	6.35	8.20	9.09	9.57	10.6
Fe ₂ O _{3tot}	12.0	12.4	13.7	12.1	12.4	12.6	12.6	12.0	12.3	12.2	12.3	12.4	12.7	12.9	13.0
MnO	0.169	0.221	0.227	0.301	0.190	0.179	0.177	0.180	0.176	0.178	0.183	0.178	0.181	0.184	0.190
MgO	25.3	27.1	25.7	28.6	18.1	22.6	22.3	26.4	25.2	28.3	28.3	24.2	21.2	20.3	16.2
CaO	8.21	7.33	7.51	6.86	10.2	8.74	8.88	7.42	7.88	7.18	7.16	8.41	9.14	9.41	10.6
Na ₂ O	0.077	0.074	0.092	0.090	1.21	1.13	1.15	0.92	1.11	0.94	0.88	1.15	1.25	1.28	1.29
K ₂ O	0.004	0.018	0.056	0.019	0.068	0.040	0.047	0.135	0.086	0.043	0.119	0.062	0.054	0.069	0.089
P ₂ O ₅	0.053	0.033	0.034	0.027	0.063	0.039	0.040	0.039	0.036	0.031	0.028	0.039	0.045	0.047	0.056
LOI	11.3	7.20	6.39	7.49	0.13	0.63	0.55	1.72	1.40	1.47	2.15	1.25	1.01	1.22	0.09
Total	99.85	99.81	100.00	99.55	100.00	99.90	100.22	100.05	99.68	100.21	99.88	99.88	99.92	99.84	99.85
Cr	3834	3674	3267	3806	1847	2309	2241	2895	2843	3333	3280	2719	2317	2127	1557
Al ₂ O ₃ /TiO ₂	11.1	11.8	11.8	11.9	13.9	14.0	13.9	13.8	14.3	13.3	13.0	14.1	13.7	14.1	13.7

Data (in wt.% except for Cr in ppm) were recalculated on an anhydrous basis, but not re-normalized to 100% in order to preserve information on the quality of the analyses. Here and in Tables 2 and 3, samples from 12D-PPR to LP10 are Jeesiörova komatiites and those from KD03 to KD14 - Kevitsa komatiite dikes.

1
2
3
4
5
61574
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
381575
391576
40
411577
42
43
44
45
46
47
48
49

Table 2. Trace element data for Jeesiörova and Kevitsa komatiites.

Sample	12D-PPR	13-EJH	17.1-PPR	LP-10	KD-03	KD-05	KD-06	KD-07	KD-08	KD-09	KD-10	KD-11	KD-12	KD-13	KD-14	BSE
Sc	28.8	26.4	28.4	24.8	33.6	30.4	29.3	26.8	26.6	25.0	25.3	28.4	31.2	31.7	35.0	14.9
V	217	196	207	184	277	238	238	204	210	182	190	223	245	251	294	77.0
Ni	1547	1616	1535	1734	856	1173	1138	1479	1389	1611	1624	1322	1062	1014	748	2080
Cu	141	5.22	9.45	6.01	129	114	114	96	104	89.0	89.9	108	119	123	139	28.0
Ga	10.6	9.51	10.6	8.90	13.9	11.9	11.5	10.0	10.5	9.21	9.23	11.1	12.4	12.6	14.7	4.00
Th	0.0768	0.0722	0.0769	0.0677	0.0843	0.0752	0.0746	0.0643	0.0674	0.0584	0.0583	0.0696	0.0763	0.0794	0.0888	0.0813
W	n.d.	n.d.	n.d.	26.7	n.d.	9.93	8.76	n.d.	12.7	12.1	11.6	11.4	11.9	11.8	n.d.	13.0
U	0.0281	0.0269	0.0275	0.0240	0.0247	0.0208	0.0210	0.0190	0.0199	0.0173	0.0168	0.0212	0.0221	0.0225	0.0260	0.0203
Nb	0.359	0.331	0.345	0.294	0.379	0.335	0.320	0.267	0.296	0.263	0.266	0.310	0.355	0.347	0.393	0.618
La	0.687	0.658	0.589	0.561	0.652	0.569	0.557	0.489	0.501	0.442	0.439	0.522	0.596	0.610	0.687	0.614
Ce	2.23	2.13	1.98	1.79	2.26	1.93	1.90	1.70	1.75	1.51	1.49	1.83	2.03	2.11	2.40	1.60
Pr	0.414	0.410	0.391	0.345	0.502	0.429	0.416	0.361	0.387	0.328	0.333	0.399	0.448	0.460	0.526	0.242
Nd	2.71	2.65	2.61	2.25	3.43	2.91	2.87	2.50	2.62	2.25	2.30	2.77	3.06	3.13	3.62	1.19
Sm	1.19	1.15	1.18	1.01	1.57	1.34	1.35	1.15	1.19	1.04	1.05	1.25	1.41	1.45	1.66	0.387
Hf	0.803	0.775	0.782	0.706	1.036	0.924	0.913	0.797	0.822	0.720	0.723	0.873	0.944	0.967	1.093	0.268
Zr	28.4	25.2	28.2	24.3	36.9	31.9	32.0	26.5	28.3	25.3	25.0	29.3	33.2	34.5	39.1	9.71
Eu	0.566	0.268	0.311	0.343	0.646	0.553	0.525	0.462	0.468	0.381	0.372	0.501	0.593	0.593	0.677	0.146
Gd	1.81	1.69	1.63	1.54	2.20	1.91	1.90	1.60	1.70	1.49	1.48	1.79	1.99	2.04	2.32	0.513
Tb	0.311	0.289	0.307	0.264	0.381	0.326	0.325	0.282	0.293	0.257	0.259	0.312	0.348	0.355	0.404	0.0940
Dv	1.98	1.86	1.96	1.74	2.48	2.14	2.14	1.81	1.93	1.71	1.70	2.01	2.27	2.34	2.63	0.638
Y	9.79	9.20	9.51	8.34	12.3	10.5	10.7	9.07	9.76	8.50	8.45	10.0	11.0	11.4	12.9	3.94
Ho	0.398	0.373	0.387	0.353	0.504	0.433	0.432	0.362	0.386	0.340	0.339	0.399	0.445	0.463	0.533	0.142
Er	1.09	1.01	1.07	0.955	1.37	1.17	1.18	1.00	1.05	0.919	0.915	1.09	1.23	1.27	1.44	0.417
Tm	0.148	0.140	0.142	0.132	0.186	0.157	0.161	0.138	0.143	0.126	0.126	0.152	0.167	0.172	0.196	0.0643
Yb	0.911	0.880	0.929	0.817	1.21	1.02	1.02	0.87	0.923	0.815	0.816	0.964	1.08	1.11	1.26	0.414
Lu	0.129	0.127	0.129	0.117	0.170	0.145	0.147	0.125	0.132	0.115	0.116	0.137	0.154	0.158	0.179	0.0637
Ti/Zr	146	151	145	144	119	115	118	120	118	115	116	119	120	118	119	112
Nb/Nb*	0.566	0.550	0.586	0.546	0.585	0.587	0.568	0.545	0.582	0.593	0.601	0.588	0.602	0.570	0.576	1.00
W/W*	n.d.	n.d.	n.d.	2.07	n.d.	0.785	0.691	n.d.	1.09	1.19	1.16	0.931	0.902	0.874	n.d.	1.00
(La/Sm) _N	0.362	0.359	0.315	0.350	0.261	0.267	0.261	0.268	0.264	0.268	0.264	0.263	0.267	0.265	0.260	1.00
(Gd/Yb) _N	1.60	1.56	1.41	1.53	1.47	1.51	1.49	1.48	1.49	1.48	1.47	1.50	1.49	1.48	1.48	1.00

Abundances recalculated on an anhydrous basis in ppm, except for W (ppb). *N*- normalized to the BSE values of Hofmann (1988) for all elements except Ga (McDonough and Sun, 1995) and W (Arevalo and McDonough, 2008), as reported in the table.

Table 3. Highly siderophile element abundances (in ppb) and Re-Os isotopic data for Jeesiörova and Kevitsa komatiites and olivine and chromite separates.

Sample No.	Re	Os	Ir	Ru	Pt	Pd	$^{187}\text{Re}/^{188}\text{Os}$	$^{187}\text{Os}/^{188}\text{Os}$	$\gamma^{187}\text{Os}(2049)$
12D-PPR WR	0.9867	5.434	4.226	7.436	17.89	15.35	0.8766±0.0035	0.14236±7	-1.0
13-EJH WR	0.6650	4.269	3.279	6.235	18.26	14.87	0.7517±0.0030	0.13836±6	-0.7
17.1-PPR WR	0.8497	3.213	2.637	5.869	19.17	15.22	1.280±0.005	0.15817±7	+0.6
LP-10 WR	0.3756	4.435	3.631	6.657	16.75	13.93	0.4082±0.0016	0.12871±6	+1.3
KD-03 WR	1.188	1.528	1.472	2.573	22.31	19.01	3.805±0.015	0.24505±2	-0.1
KD-05 WR	1.007	2.149	1.845	3.549	19.77	16.92	2.277±0.009	0.19117±8	-0.9
KD-06 WR	1.010	5.528	3.006	3.824	19.57	16.49	0.8817±0.0035	0.14353±7	-0.1
KD-07 WR	0.8141	4.161	2.571	4.082	16.82	14.72	0.9452±0.0038	0.14662±1	+0.7
KD-08 WR	0.8693	3.684	2.824	4.300	17.77	15.32	1.141±0.005	0.15200±8	-0.6
KD-09 WR	0.7062	3.956	2.870	5.001	16.02	13.74	0.8620±0.0035	0.14307±6	+0.1
KD-10 WR	0.6813	3.190	2.434	5.027	16.22	13.60	1.032±0.004	0.14820±7	-0.6
KD-11 WR	0.9146	3.701	2.708	4.262	19.20	15.87	1.195±0.005	0.15471±7	+0.1
KD-12 WR	1.051	2.297	1.753	3.545	20.43	17.04	2.223±0.009	0.19087±10	+0.5
KD-13 WR	1.127	3.456	2.166	3.436	21.10	17.97	1.579±0.006	0.16626±10	-1.4
Replicate	1.109	2.063	1.933	3.384	21.04	17.76	2.617±0.010	0.20361±9	-0.3
KD-14 WR	1.284	2.035	1.593	2.168	23.68	20.01	3.076±0.012	0.21969±2	-0.2
12D-PPR Chr	1.195	122.7					0.0469±0.0004	0.11378±5	-0.8
17.1-PPR Chr	1.233	11.13					0.5341±0.0044	0.13226±8	+0.6
12D-PPR_1 Chr	1.688	156.7	102.2		27.85		0.0518±0.0010	0.11462±7	-0.2
12D-PPR_2 Chr	0.8898	141.0	94.4		33.03		0.0304±0.0010	0.11357±6	-0.5
12D-PPR_3 Chr	0.6325	149.8	102.5		36.05		0.0203±0.0009	0.11336±6	-0.4
12D-PPR_4 Chr	0.5442	153.3	103.6		38.92		0.0171±0.0009	0.11339±6	-0.2
12D-PPR_5 Chr	0.6303	150.2	101.7		47.32		0.0202±0.0009	0.11329±7	-0.4
12D-PPR_6 Chr	0.5624	157.5	108.0		52.97		0.0172±0.0009	0.11337±5	-0.3
13-EJH_1 Chr	0.2597	53.32	38.33		6.164		0.0234±0.0027	0.11394±6	+0.1
13-EJH_2 Chr	0.3449	53.59	38.45		7.314		0.0310±0.0026	0.11393±7	-0.2
KD-05 Ol	0.04468	3.952	2.262	2.296	1.525	1.426	0.0544±0.0009	0.11455±4	-0.4
KD-10 Ol	0.6178	3.126	2.285	4.158	1.936	1.698	0.9547±0.0050	0.14627±4	0.0
KD-05 Chr	0.4528	35.29					0.0617±0.0016	0.11497±6	-0.2
KD-08 Chr	0.5359	15.54					0.1660±0.0023	0.11837±8	-0.4
KD-10 Chr	1.197	78.14					0.0737±0.0011	0.11531±7	-0.3
KD-12 Chr	0.7288	7.693					0.4568±0.0062	0.12945±13	0.0
LP-10 Chr	1.014	55.28					0.0883±0.0019	0.11655±7	-0.2

HSE abundances recalculated on an anhydrous basis. Initial $\gamma^{187}\text{Os}$ values were calculated at $T = 2049$ Ma derived from the isochron using the parameters specified in the text. The HSE normalizing values are from Horan et al. (2003). *Replicate* – replicate digestion of the sample. Chromite separates 12D-PPR_1 through 12D-PPR_6 and 13-EJH_1 and 13-EJH_2 denote spiked subsamples of unspiked digestions for the Pt-Os study.

Table 4. High-precision Os isotopic data and Re/Os and Pt/Os ratios for chromite separates from the Jeesiörova komatiites.

Sample No.	$^{187}\text{Re}/^{188}\text{Os}$	$^{187}\text{Os}/^{188}\text{Os}$	$\gamma^{187}\text{Os}(T)$	$^{190}\text{Pt}/^{188}\text{Os}$	$^{184}\text{Os}/^{188}\text{Os}$	$^{186}\text{Os}/^{188}\text{Os}$	$\mu^{186}\text{Os}(T)$
12D-PPR_1 Chr	0.0518±0.0010	0.1146036±06	-0.23	0.0001697±34	0.0013031±2	0.1198369±6	+25±5
12D-PPR_2 Chr	0.0304±0.0010	0.1135064±05	-0.54	0.0002238±45	0.0013028±2	0.1198381±5	+34±4
12D-PPR_3 Chr	0.0203±0.0009	0.1134565±07	-0.27	0.0002299±46	0.0013023±3	0.1198377±7	+30±6
12D-PPR_4 Chr	0.0171±0.0009	0.1134060±05	-0.22	0.0002425±48	0.0013033±2	0.1198381±5	+33±4
12D-PPR_5 Chr	0.0202±0.0009	0.1134294±06	-0.29	0.0003009±60	0.0013033±2	0.1198378±6	+29±5
12D-PPR_6 Chr	0.0172±0.0009	0.1134048±06	-0.22	0.0003213±64	0.0013033±2	0.1198376±6	+27±5
13-EJH Chr	0.0272±0.0019	0.1138779±11	-0.11	0.0001204±24	0.0013024±4	0.1198370±10	+27±8

Initial $\gamma^{187}\text{Os}$ and $\mu^{186}\text{Os}$ values were calculated at $T = 2049$ Ma using the parameters specified in the text.

Table 5. Tungsten isotopic compositions of Kevitsa komatiites.

Sample	$\mu^{182}\text{W}$ ($\pm 2\text{SE}$)	$\mu^{183}\text{W}$ ($\pm 2\text{SE}$)
KD-08	-1.7 ± 3.6	-0.8 ± 4.3
<i>Replicate</i>	$+2.2 \pm 4.3$	-2.1 ± 4.8
KD-12	$+5.9 \pm 3.0$	$+2.1 \pm 3.6$
<i>Replicate</i>	-0.2 ± 3.3	-2.6 ± 4.1
Average ($\pm 2\text{SE}$)	$+1.5 \pm 3.3$	-0.8 ± 2.1

Uncertainties on W isotopic compositions for individual samples are the 2SE in-run uncertainties of the individual analyses.

Table 6. Sm-Nd isotope and concentration data for Jeesiörova and Kevitsa komatiites.

Sample	Sm, ppm	Nd, ppm	$^{147}\text{Sm}/^{144}\text{Nd}$	$\pm 2\text{SD}$	$^{143}\text{Nd}/^{144}\text{Nd}$	$\pm 2\text{SE}$	$\varepsilon^{143}\text{Nd}(\text{T})$
13-EJH WR	1.081	2.446	0.2673	0.0005	0.513768	0.000004	+3.5
17.1-PPR WR	1.123	2.445	0.2779	0.0006	0.513917	0.000003	+3.6
KD-05 WR	1.379	2.966	0.2811	0.0006	0.513959	0.000004	+3.6
KD-08 WR	1.176	2.546	0.2794	0.0006	0.513950	0.000003	+3.8
KD-10 WR	0.9099	1.950	0.2821	0.0006	0.513981	0.000003	+3.7
KD-12 WR	1.501	3.245	0.2798	0.0006	0.513946	0.000003	+3.7
13-EJH Cpx	0.9271	1.542	0.3636	0.0007	0.515069	0.000003	+3.5
17.1-PPR Cpx	0.9256	1.496	0.3743	0.0008	0.515217	0.000003	+3.6
KD-05 Cpx	0.6775	1.309	0.3131	0.0006	0.514395	0.000003	+3.7
KD-08 Cpx	0.6564	1.335	0.2975	0.0006	0.514177	0.000004	+3.5
KD-10 Cpx_1	0.6936	1.368	0.3066	0.0006	0.514317	0.000004	+3.9
KD-10 Cpx_2	0.7036	1.388	0.3066	0.0006	0.514316	0.000003	+3.8
KD-12 Cpx	0.8599	1.676	0.3104	0.0006	0.514366	0.000003	+3.8

Initial $\varepsilon^{143}\text{Nd}$ values were calculated at $T = 2049$ Ma using the parameters specified in the text. WR - whole-rock samples; Cpx - clinopyroxene separates.

Table 7. Lu-Hf isotope and concentration data for the Jeesiörova komatiites and USGS SRM BCR-1.

Sample	Lu (ppm)	Hf (ppm)	$^{176}\text{Lu}/^{177}\text{Hf}$	$\pm 2\text{SE}$	$^{176}\text{Hf}/^{177}\text{Hf}$	$\pm 2\text{SE}$	$\epsilon^{176}\text{Hf}(\text{T})$
13-EJH WR	0.1742	1.054	0.02343	0.00001	0.282630	0.000003	+8.6
13-EJH Cpx	0.1326	0.4958	0.03791	0.00001	0.283201	0.000004	+8.8
BCR-1	0.4980	4.949	0.01426	0.00001	0.282872	0.000004	
BCR-1*	0.4988 $\pm$ 0.0048	4.92 $\pm$ 0.06			0.282867	0.000015	

Initial $\epsilon^{176}\text{Hf}$ values were calculated at $T = 2049$ Ma, using the parameters specified in the text. *Denotes the *GeoRem* recommended values.

Table 8. Compositions of the emplaced Jeesiörova-Kevitsa komatiite magma, crustal contaminant endmember, and the calculated original komatiite magma.

	1. Emplaced komatiite magma	2. Crustal contaminant	3. Original komatiite magma
SiO₂	44.1	68.2	43.9
TiO₂	0.554	0.335	0.556
Al₂O₃	7.94	17.1	7.85
Fe₂O_{3tot}	12.3	3.28	12.4
MnO	0.176	0.048	0.177
MgO	25.2	1.35	25.4
CaO	7.88	4.01	7.92
Th	0.0674	3.51	0.0327
W	11.7	595	5.81
U	0.0199	0.980	0.0102
Nb	0.296	3.06	0.268
La	0.501	25.0	0.253
Ce	1.75	46.2	1.30
Pr	0.387	4.99	0.340
Nd	2.62	17.2	2.48
Sm	1.19	2.57	1.18
Hf	0.822	3.81	0.792
Gd	1.70	2.01	1.70
Tb	0.293	0.243	0.293
Dy	1.93	1.33	1.94
Ho	0.386	0.252	0.387
Er	1.05	0.704	1.06
Tm	0.143	0.0987	0.144
Yb	0.923	0.652	0.926
Lu	0.132	0.0970	0.133
Nb/Nb*	0.582	0.118	1.07
W/W*	1.00	1.00	1.00
(Nb/Th)_N	0.577	0.115	1.08
(La/Sm)_N	0.264	6.14	0.135
(Gd/Yb)_N	1.49	2.50	1.48
ε¹⁴³Nd(<i>T</i>)	+3.7±0.3	-14.5±1.0	+4.9±0.5
ε¹⁷⁶Hf(<i>T</i>)	+8.7±0.3	-21.8±1.9	+10.2±0.6
μ¹⁸²W	+1.5±3.3	+12.6±4.5	-10±5

1- emplaced magma composition for the Jeesiörova-Kevitsa komatiites; 2 - FTA composition compiled from the average Vodla Block tonalite composition of Puchtel et al. (2016b). 3 - original parental komatiite magma for the Jeesiörova-Kevitsa komatiites. The initial εNd and εHf values were calculated at $T = 2049$ Ma.

Table 9. Results of mixing calculations between an IIIAB iron meteorite endmember and the BSE mantle domain.

	Tieraco Creek	BSE	BSE+0.2% TC	Mantle source
Os, ppb	31.7	3.40±0.50	3.50±0.50	3.39±0.10
Re, ppb	3.10	0.289±0.007	0.295±0.010	0.285±0.010
Pt, ppb	2089	6.1±1.6	10.3±1.6	9.54±0.22
W, ppb	200	13.0±10	13.4±10	5.8±5
¹⁸⁷ Re/ ¹⁸⁸ Os	0.464	0.402	0.403	0.397±0.008
¹⁹⁰ Pt/ ¹⁸⁸ Os	0.0630	0.00171	0.00284	0.00269±0.008
$\mu^{182}\text{W}$	-335	0	-10	-10±5

The composition of the Jeziřova-Kevitsa komatiite mantle source is calculated to have resulted from mixing of 0.2% of material of the IIIAB iron meteorite Tieraco Creek into a mantle domain with the BSE composition. Uncertainties on the HSE abundances in the average estimate for the BSE are 1SE (Becker et al., 2006).

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4A.

Fig. 4B.

Fig. 5.

Fig. 6.

Fig. 7.

Fig. 8.

Fig. 9.

Fig. 10.

Fig. 11.

Fig. 12.

Fig. 13.

Fig. 14.

Supplementary Tables A1 and A2

[Click here to download Background dataset for online publication only: Supplementary Tables A1 and A2.docx](#)

Supplementary Table A3

[Click here to download Background dataset for online publication only: Supplementary Table A3.xlsx](#)

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.