Carbon/Nitrogen Metabolic Balance: Lessons from Cyanobacteria

Cheng-Cai Zhang, Cong-Zhao Zhou, Robert L Burnap, Ling Peng

► To cite this version:

Cheng-Cai Zhang, Cong-Zhao Zhou, Robert L Burnap, Ling Peng. Carbon/Nitrogen Metabolic Balance: Lessons from Cyanobacteria. Trends in Plant Science, 2018, 23 (12), pp.1116 - 1130. 10.1016/j.tplants.2018.09.008 . hal-02992919

HAL Id: hal-02992919

<https://hal.science/hal-02992919>

Submitted on 18 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Highlights

- Cyanobacteria are simple models among photosynthetic organisms for studying the metabolic regulation of carbon/nitrogen (C/N), the two most abundant nutrient elements for all living organisms.
- 2-oxoglutarate (2-OG), an intermediate from the Krebs cycle, serves as a carbon skeleton for nitrogen assimilation and as a signal of nitrogen limitation, whereas 2-phosphoglycolate (2-PG), an intermediate from photorespiration, acts as a signal of inorganic carbon limitation.
- The levels of 2-PG and 2-OG are inversely correlated, and their ratio reflects the C/N metabolic balance.
- The transcriptional activator NtcA is a sensor of 2-OG, and the transcriptional repressor NdhR is a sensor of both 2-OG and 2-PG. These regulators together balance C/N metabolic networks by switching on or off the expression of genes involved mainly in the uptake and assimilation of carbon and nitrogen sources.
- The signalling role of 2-OG and 2-PG in C/N balance is likely conserved in other photosynthetic organisms.

Carbon/nitrogen metabolic balance: lessons from cyanobacteria

Cheng-Cai Zhang^{1,5*}, Cong-Zhao Zhou², Robert L. Burnap³ and Ling Peng⁴

¹Key Laboratory of Algal Biology, Institute of Hydrobiology, the Chinese Academy of Sciences, Wuhan, Hubei 430072, People's Republic of China

²School of Life Sciences and Hefei National Laboratory for Physical Sciences at the Microscale, University of Science and Technology of China, Hefei, Anhui, 230027, People's Republic of China

³Department of Microbiology and Molecular Genetics, Henry Bellmon Research Center, Oklahoma State University, Stillwater, OK 74078, USA

⁴Aix-Marseille Université, CNRS, Centre Interdisciplinaire de Nanoscience de Marseille, CInaM UMR 7325, «Equipe Labellisée Ligue Contre le Cancer», 13288 Marseille, France
CInaM UMR 7325, 13288 Marseille, France

⁵Aix Marseille Univ, CNRS, LCB, Marseille, France

*Corresponding author: Zhang, C.-C. (cczhang@ihb.ac.cn).

Keywords: nitrogen metabolism, carbon fixation, photorespiration, signal transduction

31

32 **Abstract**

33 Carbon and nitrogen are the two most abundant nutrient elements for all living organisms,
34 and their metabolism is tightly coupled. What are the signalling mechanisms that cells use to
35 sense and control the carbon/nitrogen (C/N) metabolic balance following environmental
36 changes? Based on studies in cyanobacteria, it was found that 2-phosphoglycolate derived
37 from photorespiration and 2-oxoglutarate from the Krebs cycle act respectively as the carbon-
38 and nitrogen-starvation signals, and their concentration ratio likely reflects the status of the
39 C/N metabolic balance. We will present and discuss the regulatory principles underlying the
40 signalling mechanisms, which are likely to be conserved in other photosynthetic organisms.
41 These concepts may also contribute to developments in the field of biofuel engineering or
42 improvements in crop productivity.

43

44

45

Importance of carbon/nitrogen (C/N) balance

Carbon and nitrogen metabolism is essential for every biological system, since all major cellular components, including genetic materials, proteins, pigments, energy carrier molecules, etc., are derived from these activities. Carbon and nitrogen metabolism is tightly coupled in different living organisms. In prokaryotes and plants, such a coupling mechanism can be attributed to two major factors. First, the two elements are the most abundant in cells intensifying the requirement for coordination mechanisms to avoid metabolic inefficiencies; second, nitrogen assimilation depends on the availability of a carbon skeleton for biosynthesis, and consequently the limitation or oversupply of one element strongly affects the metabolism of the other. Therefore, the C/N stoichiometry in different organisms varies within a relatively narrow range; for example, the mass ratio is C/N=31/4 in phytoplankton [1].

Because properly balanced metabolism of carbon and nitrogen is necessary for optimal growth, different levels of regulation exist in cells in order to control the uptake and assimilation of various nitrogen and carbon sources whose supply may vary under different environmental conditions [2–4]. Such regulation may occur at various levels of control ranging from the allosteric modulation of the activity of nutrient assimilation proteins to a variety of mechanisms controlling the expression of genes encoding these structural proteins. The latter includes a rich variety of transcriptional and post-transcriptional mechanisms, some of which are still being experimentally clarified [5–7].

Whatever the type of regulation involved, two major questions arise: how do cells sense the metabolic status, and what signal transduction mechanisms do cells use to maintain a proper balance of nitrogen and carbon metabolism? Recently, tremendous progress has been made towards answering these questions by using cyanobacteria as a model for photosynthetic organisms, and this review serves to summarize the data and highlight the mechanisms underlying the signaling and transcriptional regulation involved in carbon and nitrogen metabolic control. We will also discuss the concept for controlling C/N metabolic balance. Cyanobacteria are the ancestors of plastids [8]; therefore, cyanobacteria and plants may share common features, especially in terms of the signaling mechanisms, for balancing C/N metabolism.

Coupling between carbon and nitrogen metabolic pathways

To facilitate understanding, we briefly summarize here the metabolic coupling of nitrogen and

carbon assimilation in cyanobacteria (Fig. 1). As for other autotrophic photosynthetic organisms, CO₂ is the primary carbon source for cyanobacteria, although some cyanobacterial strains can also use a sugar or other organic carbon as the carbon source [4,9,10]. However, as with eukaryotic algae, the availability of CO₂ is often growth-limiting because of specific physical factors such as the low diffusion rates in the aqueous environment. Moreover, dissolved CO₂ chemically speciates into several forms of inorganic carbon according to the general reaction describing the inorganic carbon pool:

Note that the equilibrium of the reaction is strongly pH dependent such that the fraction of the total inorganic pool existing as CO₂ is favored by acidic conditions, whereas bicarbonate HCO₃⁻ dominates under neutral and alkaline conditions. CO₃²⁻ can be ignored in general as it exists only in strong alkaline condition, which is seldom the case within normal cells. Cyanobacteria and algae are typically capable of acquiring either CO₂ or HCO₃⁻. Because both forms may be acquired and are rapidly interchanged within the cell, the symbol C_i is often utilized to refer to total inorganic carbon (i.e. C_i = CO₂ + HCO₃⁻) (Fig. 1). CO₂ fixation is catalyzed by Rubisco in the Calvin-Benson-Bassham (CBB) cycle. The reaction involves the carboxylation, using CO₂ and ribulose 1,5-biphosphate (RuBP), yielding two molecules of the three carbon 3-phosphoglycerate (3-PGA) [11–13]. However, Rubisco has a notoriously low affinity for CO₂ (K_m >150 μM) and, problematically, the enzyme also catalyzes the competing and wasteful oxygenase reaction between RuBP and O₂, which yields one molecule of the two-carbon 2-Phosphoglycolate (2-PG) and one 3-PGA rather the two 3-PGA molecules of the carboxylation reaction. The oxygenase reaction has resulted in the evolution of scavenging photorespiratory pathways, which recover and recycle the carbon that would otherwise be lost [11,13]. As a consequence of these factors, carbon fixation in cyanobacteria critically depends upon the so-called CO₂-concentrating mechanism (CCM) [4,14,15]. Cyanobacteria actively acquire C_i, either as HCO₃⁻ or as CO₂, using a complex set of transporters and uptake enzymes. Bicarbonate uptake involves active membrane transporters, such as the sodium-dependent bicarbonate transporter, SbtA or BicA. Energized CO₂ uptake occurs via the enzymatic hydration of CO₂ catalyzed by specialized forms of the NDH-1 complex located in thylakoid membrane. These complexes use redox energy to drive the conversion of CO₂ to HCO₃⁻ in the cytoplasm and simultaneously draw in CO₂ from the environment. Accordingly, these are not simple carbonic anhydrases, but enzymes capable of using metabolic energy to create a

disequilibrium in favor of HCO_3^- accumulation. Together, these systems produce and maintain high concentrations of HCO_3^- in the cytoplasm to supply the carboxysome, which encloses the entire cellular complement of Rubisco within a protein shell. The carboxysome is the first identified member of a large class of prokaryotic structures called *bacterial microcompartments*, which generally partition specific metabolic reactions within the cytoplasm of the bacterial species that possess them. In the case of the carboxysome, RuBP and HCO_3^- are imported through specific pores in the protein shell and 3-PGA is exported through pores in the shell to supply the CBB cycle, although the details of these transport process are only beginning to emerge. Similarly, the oxygenase reaction product, 2-PG, must also be able to diffuse out of the carboxysome, although again, the precise mechanisms for passage through the protein shell of the carboxysome remain to be elucidated [14]. Importantly, the carboxysome also contains a carbonic anhydrase with functions to convert imported HCO_3^- into CO_2 in the proximity of Rubisco. Therefore, given an abundant supply of HCO_3^- , the active sites of Rubisco within the carboxysome are saturated with CO_2 , which enables high rates of CO_2 fixation. This enormously diminishes the competing reaction with O_2 ; however under conditions of C_i limitation or in mutants that fail to accumulate sufficient HCO_3^- in the cytoplasm, significant rates of the oxygenase reaction have been observed to occur [11,13]. Thus, the CCM relies on the ability of the C_i -uptake systems to concentrate high concentrations of HCO_3^- within the cytoplasm in order to create a steep concentration gradient that promotes rapid net diffusion of HCO_3^- into the carboxysome. A substantial proportion of 3-PGA, produced by the CBB cycle, is converted to acetyl-CoA which feeds the essential central metabolic pathway, the Krebs cycle (also known as the citric acid cycle, tricarboxylic acid cycle or TCA cycle), thus leading to the production of intermediates for biosynthesis [4,13,16,17] (Fig. 1). As discussed below, the coordination of N- and C-metabolism depends, in part upon cytoplasmic concentrations of several metabolites affected by the efficiency of the CO_2 -fixation reactions.

Cyanobacteria can use different inorganic nitrogen sources, mostly in the form of nitrate or ammonium. Many cyanobacterial strains can synthesize the enzyme nitrogenase for N_2 fixation [3,9]. Whatever the form of inorganic nitrogen that is taken up into the cells, it is always converted into ammonium, the least costly nitrogen form for assimilation (Fig. 1). Nitrate requires two additional enzymes, nitrate reductase and nitrite reductase and reduced cofactors, in order to be reduced to ammonium, whereas N_2 fixation is even more

energetically expensive requiring both reduced cofactors and up to 16 ATP molecules per ammonium. Moreover, it depends on the nitrogenase complex which needs to operate under micro-oxic conditions, thereby requiring special adaptations. Some filamentous cyanobacteria, such as *Anabaena/Nostoc* PCC 7120 can form heterocysts, intercalated among vegetative cells, to provide a micro-oxic intracellular environment for N₂ fixation or, alternatively, N₂ fixation and photosynthetic O₂ evolution occur in the same cell, but the activities are diurnally regulated using a circadian clock so that N₂ fixation is restricted to the night [3,18–20]. Hierarchically, ammonium represses the use of alternative nitrogen sources, and heterocyst differentiation and N₂ fixation take place only when neither ammonium nor nitrate is available in the growth medium.

Two ammonium-assimilation pathways are known, the glutamine synthetase-glutamate synthase (GS-GOGAT) cycle, and direct assimilation through the reaction catalyzed by glutamate dehydrogenase (GDH). Both pathways require the carbon skeleton 2-oxoglutarate (2-OG, also known as α -ketoglutarate), one of the intermediates of the Krebs cycle [21,22] (Fig. 1). Therefore, ammonium assimilation using 2-OG provides the metabolic basis for coupling between nitrogen and carbon metabolism. In the GS-GOGAT cycle, GS assimilates ammonium using glutamate and ATP to synthesize glutamine, while GOGAT catalyzes glutamate formation using glutamine, 2-OG and NADPH. The second route for ammonium assimilation, catalyzed by GDH using 2-OG, does not play a major role in cyanobacteria, and the majority of cyanobacterial species whose genomes have been sequenced do not even contain the gene encoding GDH [21,23]. Thus, the GS-GOGAT cycle constitutes the major pathway for ammonium assimilation in cyanobacteria, and hence the central crossroad for carbon and nitrogen metabolism (Fig. 1). Glutamine and glutamate are also the major intracellular amino group donors for the synthesis of several other amino acids, and for purine and pyrimidine nucleobases, as well as other nitrogen-containing compounds [24,25].

Both carbon and nitrogen assimilation are under multiple control mechanisms depending on the availability or the nature of the nutrients in the environment. To understand how cyanobacteria regulate nitrogen and carbon metabolism, we will first focus on the nature of the signals that allow cells to sense the intracellular metabolic status when nitrogen and carbon metabolism becomes unbalanced, and then we will review the signaling mechanisms that allow cells to regain their metabolic balance in order to better adapt to environmental

changes. While the focus of this analysis is based upon the transcriptional regulation, it is important to note that researchers performing global transcriptome and proteome analyses have observed significant differences between fold-changes in transcript and the corresponding protein levels in cyanobacteria [26,27]. This indicates the existence of post-transcriptional factors in the control of gene expression in cyanobacteria. However, it is also important to note that there is generally a good qualitative correspondence between the levels of transcripts and proteins, with the latter typically exhibiting more muted fold changes in response to environmental changes.

The nitrogen-starvation signal 2-OG and its receptors

It has been postulated for a while that 2-OG acts as a signal in the control of nitrogen metabolism because of its role as a carbon skeleton in nitrogen assimilation and because its levels can vary according to nitrogen availability [28,29] (Fig. 1). Indeed, in both the filamentous heterocyst-forming diazotrophic cyanobacterium *Anabaena* PCC 7120 and the unicellular non-diazotrophic strain *Synechocystis* PCC 6803, 2-OG accumulates rapidly upon limitation of nitrogen supply in the growth medium [28,29]. A signaling function for 2-OG has also been proposed in a variety of biological systems, including bacteria, animals and humans [2,25,30–35]. In proteobacteria, for example, 2-OG is involved in the control of metabolism [25,36], although the signaling mechanism is in many ways different from that in cyanobacteria, as will be discussed later in this review.

Because 2-OG is a critical metabolite rapidly incorporated into a variety of N-containing compounds, one crucial question is how to experimentally discriminate its metabolic function from its signaling function, and how its direct effects are distinguished from the indirect effects of its metabolic derivatives. One way to tackle this problem is through the use of nonmetabolizable analogs, in which a stable bioisostere function replaces the keto group in 2-OG, the site of amidation in ammonium assimilation in the GS-GOGAT cycle. Specifically, several nonmetabolizable analogs of 2-OG were chemically synthesized and tested in *Anabaena* PCC 7120 [29,37–40]. This organism was chosen for these tests because it has, under nitrogen-limited conditions, a phenotype (formation of N₂-fixing heterocysts) which is easily observable under a light microscope [3,18]. Among the synthesized nonmetabolizable analogs of 2-OG, 2,2-difluoropentanedioic acid (DFPA) is particularly interesting (Fig. 2). DFPA was designed to mimic 2-OG by replacing the keto group with the stable and

nonmetabolizable fluorinated methylene entity [29]. Since fluorine is in general absent in biological systems and cells, fluorinated compounds can be readily tracked in intact cells using ¹⁹F-NMR, with little or no background interference [41,42]. Accumulation of DFPA in filaments of *Anabaena* PCC 7120 acts as a trigger for heterocyst differentiation, even when ammonium, which normally represses heterocyst differentiation, is present in the growth medium. DFPA can be recognized by both a 2-OG permease, KgtP, and a 2-OG receptor, NtcA [29,43]. These results provided the first in vivo evidence for a direct role of 2-OG as a nitrogen starvation signal in cyanobacteria.

Two 2-OG receptors have been known for a while in cyanobacteria, NtcA and PII [2,3]. The third, NdhR, was more recently established as a 2-OG receptor [44–46]. Here we will first focus on NtcA and PII, because NdhR is also a receptor of 2-PG [46–48], and will be discussed later in this review as a key element in C/N metabolic balance.

PII is a trimeric regulatory protein which is highly conserved in both bacteria and plants [2,35,49]. PII exerts its function mainly through protein-protein interactions mediated by the flexible and outward-extended T-loop present on each monomer. The binding of 2-OG determines the conformation of the T-loop, and hence the interaction with the partners in nitrogen metabolism [50]. Among the interacting partners of PII are PipX, which is involved in NtcA-mediated transcriptional control by acting as a co-activator of NtcA, and NAGK (N-acetyl-L-glutamate kinase), a key enzyme for arginine synthesis [51,52]. The interaction of PII with either PipX or NAGK occurs under nitrogen sufficiency, and thus when the intracellular 2-OG level is low [51–54]. PII relieves the feedback inhibition of Arg on the activity of NAGK, resulting in an increase of Arg for cyanophycin synthesis as a nitrogen reserve as a consequence of nitrogen sufficiency [52,53]; at the same time, PII sequesters PipX, preventing its role as a co-activator of NtcA [51,54]. Several reviews dealing with the detailed mechanism of PII signaling are available for readers interested in more details in this topic [2,35,49]. Importantly, additional proteins in the PII-family are recently reported as being involved in carbon assimilation [7,55]. The small PII-type protein StbB is physically linked and co-expressed with the bicarbonate uptake transporter, StbA. Structural and physiological analyses indicate that this protein is involved in cAMP sensing and responds to cyanobacterial inorganic carbon status. This indicates the existence of a direct regulatory connection between inorganic carbon availability and broader carbon metabolism. These findings resonate with the fact that C/N balance in other bacteria such as *E. coli* involves cAMP [25,33]; indeed, cAMP participates

in carbon catabolite repression and the cAMP levels are themselves regulated by 2-OG. While it remains to be seen whether this also applies to cyanobacteria, it points to the possibility that new modes of regulation affecting C/N balance may still be found.

NtcA is a dimeric global transcriptional factor belonging to the CRP/CAP family [3]. NtcA activates the expression of genes required for heterocyst development in *Anabaena* PCC 7120 and the use of alternative nitrogen sources under nitrogen deprivation. Recent structural studies of NtcA from both *Anabaena* PCC 7120 and *Synechocystis* PCC 6803 provide further insight into the molecular mechanism of 2-OG signaling mediated through NtcA [43,56]. Specifically, each NtcA monomer harbors a 2-OG binding pocket, and 2-OG binding generates two allosteric effects in NtcA: first, there is an increase in the contact area between the two monomers through the two long C-helices, and then this change propagates towards the DNA-binding domain by shortening the distance between the two helices responsible for DNA binding (Fig. 2). In the native or apo form of NtcA, the distance between the two DNA-binding helices is about 37 Å, whereas in the 2-OG-bound form, it becomes 34 Å, which corresponds to the distance between successive DNA major grooves [43]. Furthermore, transcriptional activity assays in vitro also reveal that interaction of 2-OG-free NtcA with its target promoters is not sufficient for activation of gene transcription, and 2-OG binding to NtcA is necessary to initiate the process of transcriptional control [57].

Based on all information available so far, a molecular model of the signaling mechanism in response to nitrogen starvation can be proposed. Upon nitrogen deprivation, 2-OG accumulates rapidly in cells; it binds to PII and regulates nitrogen metabolism by protein-protein interactions on the one hand, and it interacts with NtcA which activates genes involved in the uptake and assimilation of alternative nitrogen sources on the other (Fig. 2). For those cyanobacteria able to form heterocysts, such as *Anabaena* PCC 7120, 2-OG also serves as a trigger for the initiation of heterocysts, which fix atmospheric N₂ using nitrogenase [29,58].

Evidence for 2-PG as a signal of carbon starvation

CO₂ assimilation is catalyzed by Rubisco, the most abundant enzyme on earth. As mentioned above, CO₂ and O₂ are competitive substrates for the reaction with RuBP in the active site of Rubisco [11–13] (Fig. 3). Through the carboxylase activity of Rubisco, CO₂ is combined with RuBP to produce 3-PGA; The remainder of the CBB cycles serves to regenerate RuBP and consumes ATP as energy and NADPH as reducing power. Part of the 3-PGA pool is recycled

back to RuBP through the CBB pathway, while the rest of the 3-PGA is used to make sugar to feed carbon metabolism and the Krebs cycle (Fig. 3). Through the oxygenase activity of Rubisco, each RuBP molecule is converted to 1 molecule of 2-PG and 1 molecule of 3-PGA. 2-PG, being toxic, is then metabolized by the so-called photorespiration pathway [11–13,59]. Thus, the activity of Rubisco is dependent on the relative ratio of CO₂ and O₂.

Two transcriptional regulators have been extensively studied in cyanobacteria for the control of genes involved in carbon acquisition and/or metabolism: NdhR (CcmR) and CmpR [45,60]. A third one, an AbrB-type regulator, is also involved in the control of gene expression according to carbon availability in *Synechocystis* PCC 6803, but the underlying mechanism remains largely unknown [61]; the same is true for PacR, a LysR-type transcriptional regulator identified in *Anabaena* PCC 7120 [62]. Both NdhR and CmpR are LysR-type transcription factors, and their function in cyanobacteria has been well documented [4]. Briefly, CmpR is an activator of the *cmp* operon; this operon encodes the high-affinity bicarbonate transporter BCT1, which is induced under low-CO₂ conditions [60,63] (Fig. 3). NdhR is a repressor, whose target genes, among others, encode several CO₂ or bicarbonate uptake systems located on either the cytoplasmic (Na⁺-NDH-1, Sbt) or the thylakoid membranes (Ndh-1₃) [4,44–46,48,64,65] (Fig. 3). During carbon sufficiency, genes encoding CCM-related functions are therefore repressed by NdhR. LysR-type transcription factors are usually under allosteric control; thus, the existence of effectors that interact with either CmpR or NdhR was expected. Metabolites from photorespiration or CBB, whose levels could oscillate according to carbon availability, have been tested in various studies [45,47,48,63,64,66]. DNA gel-shift assays indicate that RuBP and 2-PG can both bind to CmpR to activate the *cmp* operon and enhance CO₂ uptake under carbon-limiting conditions [48,63]. However, the effect of 2-PG on the transcription of the *cmp* operon mediated by CmpR [47,66] will need to be confirmed in vivo, and the effect of both RuBP and 2-PG on CmpR has yet to be demonstrated from structural studies. In contrast, much progress has been made in identifying the effectors of NdhR, which is considered as the most critical regulator of cyanobacterial CCM [4]. Using surface plasmon resonance, it was shown that 2-OG and NADP⁺ could both interact with NdhR and enhance its DNA binding activity [48]. In a more recent study, 4 metabolites, 2-OG, 2-PG, RuBP, and NADP⁺, were investigated, among which 2-OG and 2-PG were found to interact with NdhR: 2-OG enhances the DNA-binding activity, and is therefore a co-repressor, while 2-PG prevents NdhR from DNA binding, thus acting as an anti-repressor or inducer [46]. These observations

were further confirmed in vivo by following the effects on gene expression after changes in the 2-OG or 2-PG levels. Indeed, increasing the cellular concentration of 2-OG by adding the membrane-permeable precursor, dimethyl 2-oxoglutarate (dmOG) (Fig. 2a), enhanced the repressive effect of NdhR [46], whereas decreasing the 2-PG pool in the cells by overexpressing a gene encoding phosphoglycolate phosphatase brought down the expression levels of the NdhR regulon [46,47].

Crystal structure analyses and structural simulation have provided a mechanism whereby 2-PG and 2-OG effectors control the activity of the NdhR repressor [46] (Fig. 3b). NdhR is a tetramer composed of two compact subunits and two extended subunits. Each subunit contains a DNA-binding domain (DBD) and a regulatory domain (RD), bridged by a long α -helical linker. The NdhR tetramer contains two 2-OG binding pockets formed at the interface between two RD domains. In the 2-OG/NdhR complex, the distance between the DNA-recognition helices in the two neighboring DBDs is 29 Å, which matches a continuous DNA major groove and is thus favorable for DNA binding. Therefore, 2-OG is a co-repressor that keeps NdhR bound to its DNA targets, providing a rationale for the effect of 2-OG on the repression of the NdhR regulon. 2-PG binds to NdhR at the intradomain cleft of each RD pair, and the distance between the two DNA-binding helices becomes 25 Å, which is too short to fit the DNA major groove. Thus, 2-PG acts as an anti-repressor or inducer of the NdhR regulon. Although 2-PG and 2-OG bind to different sites in NdhR with a similar affinity, their binding is mutually exclusive because of the structural incompatibility between the two-bound forms [46].

All the data summarized above, together with the fact that the 2-PG level transiently increases in cells upon carbon limitation [44], demonstrate that 2-PG acts as a carbon starvation signal, as already suggested by the effect of reduced 2-PG on the expression of genes related to CCM [47]. Furthermore, NdhR is another 2-OG receptor in addition to PII and NtcA in cyanobacteria.

C/N metabolic balance through the integration of 2-OG and 2-PG signals

Based on all the studies using different approaches, a model can be proposed to describe how cyanobacteria achieve C/N metabolic balance by transcriptional regulation (Fig. 4). Most of the data have been derived using three model cyanobacterial strains, namely *Synechococcus elongatus* PCC 7942, *Synechocystis* PCC 6803, and *Anabaena* PCC 7120.

Firstly, let us consider how changes in the levels of 2-OG and 2-PG are related to carbon and nitrogen metabolism. Cyanobacteria in general lack 2-OG dehydrogenase, thus have an incomplete Krebs cycle [16,17,67]; in such a case, 2-OG can be considered as an end-product of this metabolic pathway (Fig. 1). The intracellular pool of 2-OG, derived from the Krebs cycle, accumulates transiently upon, and serves as a signal for, nitrogen deprivation [28,29]. This is consistent with its role as a carbon skeleton for nitrogen assimilation through the GS-GOGAT cycle (Fig. 1). The level of 2-PG, a metabolite of photorespiration, increases transiently in cells following carbon limitation [44,47], and serves as a carbon starvation signal, which is directly linked to the activity of Rubisco (Fig. 3). Indeed, when the level of CO₂ is high, the higher CO₂/O₂ ratio favors the carboxylase activity of Rubisco, and thus the production of 3-PGA; in contrast, when CO₂ becomes limiting, the lower ratio of CO₂/O₂ favors the oxygenase activity of Rubisco, leading to the production of 2-PG. Because of the strong coupling between carbon and nitrogen metabolism, especially through the GS-GOGAT cycle (Fig. 1), nitrogen limitation can also be considered as carbon oversupply, and nitrogen oversupply also correlates with carbon limitation in cells. Altogether, our current knowledge suggests that the levels of 2-OG and 2-PG are somehow inversely correlated in cells, and their ratio may reflect the carbon and nitrogen metabolic status in cyanobacteria. The transient increase in 2-OG or 2-PG observed under a particular condition is characteristic of many signaling molecules, such as Ca²⁺, cAMP, etc. [68,69].

Secondly, combining with previously published proposal and hypothesis with recent data based on genetic, biochemical and structural studies [4,28,29,43,45–47,56,64,70], we suggest a model for the transcriptional control of C/N balance, which is illustrated in Fig. 4. When C/N metabolism is unbalanced because of nitrogen limitation (or carbon oversupply) (Fig. 4a), the 2-OG pool increases while the 2-PG pool decreases for the reasons explained above. Consequently, 2-OG binds to both NtcA and NdhR. The 2-OG/NtcA complex upregulates genes involved in the uptake of alternative nitrogen sources such as nitrate, or heterocyst differentiation and N₂ fixation for diazotrophic heterocyst-forming strains such as *Anabaena* PCC 7120 [3,18]; meanwhile, the 2-OG/NdhR complex downregulates CCM so that bicarbonate uptake goes down. PII also forms a complex with 2-OG and regulates nitrogen assimilation accordingly through protein-protein interaction [2]. Together, the increased uptake and assimilation of nitrogen sources and the decreased uptake of inorganic carbon help the cells to regain balanced C/N metabolism. Similarly, when cells face carbon limitation

(or nitrogen oversupply), Rubisco switches towards its oxygenation activity, leading to accumulation of a 2-PG pool while 2-OG levels decrease because less carbon will go to feed the Krebs cycle (Fig. 3, Fig. 4b). In such a case, NtcA exists in its apo form, which is unfavorable for DNA binding and activation of genes involved in the uptake and assimilation of nitrogen sources; at the same time, 2-PG forms a complex with NdhR, making it unable to bind to DNA, which relieves the repression of genes involved in carbon acquisition. Thus, decreased uptake and assimilation of nitrogen, alongside increased uptake of inorganic carbon, helps cells to rebalance their carbon and nitrogen metabolism.

Some aspects of this model still require further experimental studies. For example, the model assumes concomitant changes in the 2-OG and 2-PG pools upon changes in the availability of carbon or nitrogen sources, consistent with the fact that 2-OG and 2-PG cannot bind to the same molecule of NdhR at the same time [46]. Although transient changes in the pools of these two metabolites have been observed [28,29,44,47], it remains unknown to what extent and on what time-scale their changes are correlated, and the current data are sometimes even contradictory [4]. It will also be relevant to identify the input of other potential signals and regulators involved in C/N metabolism, as well as their interaction with the 2-OG and 2-PG signaling pathways. For example, could RuBP or 3-PGA act as effectors during carbon metabolism as suggested by some studies [4]?

The 2-OG and 2-PG signals control the uptake of nitrogen and carbon sources or the initiation step of heterocyst differentiation when N_2 fixation is needed, and 2-OG also extensively regulates later steps in nitrogen and carbon metabolism through PII and NtcA. Indeed, NtcA is not just required for nitrogen control, it is also a global regulator directly involved in the modulation of carbon metabolism, photosynthesis, and oxidative stress [3]. For example, *cmpR* is possibly a target of NtcA in *Anabaena* PCC 7120 [71], and *ccmK2* encoding a CCM-function protein was also identified by ChIP-seq as a direct target of NtcA in *Synechocystis* PCC 6803 following nitrogen starvation [72]. Although these results still require genetic confirmation, they do point out a much broader role of NtcA in the control of C/N metabolic balance than that summarized in Fig. 4. In a regulatory perspective, by switching the early steps in C/N metabolism on and off according to the availability of one nutrient relative to another, cyanobacterial cells ensure rapid responses to environmental changes, so that the C/N balance can be maintained while saving resources that would have to be spent to control later metabolic steps. In principle, 2-OG could regulate the C/N balance alone, as it

controls the early steps of both carbon and nitrogen acquisition through its action on NdhR and NtcA (Fig. 4). Why then do cyanobacterial cells require multiple signals to control the C/N metabolic balance? The answer may be linked to the response efficiency and the sensitivity of the control system. While the intracellular concentration of 2-OG, which acts as a carbon skeleton for nitrogen assimilation, may change rapidly following shifts in nitrogen availability, its variation may be slower and of a smaller amplitude following changes in carbon availability because the Krebs cycle is situated far downstream of inorganic-carbon assimilation steps (Fig. 1; Fig. 3). By also using 2-PG, whose levels are directly dependent on carbon availability through the activity of Rubisco, cyanobacterial cells can ensure a proper and timely balance of C/N metabolism.

In the proposed model, we concentrated on 2-PG and 2-OG as signals, and NtcA and NdhR as transcription factors, because these elements are the most thoroughly studied so far using biochemical, genetic and structural approaches. Other metabolites involved in carbon fixation or photorespiration are also potential signals for carbon metabolism, and other regulatory proteins may also exist [4]. For example, additional robustness in the regulatory systems may be afforded by linkage with cAMP levels as discussed above, although the evidence for such linkages are, for cyanobacteria, largely based upon circumstantial evidence and remain to be fully validated experimentally. In addition, the role of small, non-coding, regulatory RNA in controlling cyanobacterial gene expression is only beginning to be appreciated and, potentially, these could also influence the regulation of C/N balance. These include the recent discovery of a glutamine-responsive riboswitch that enables fast regulation of glutamine synthase, the key enzyme in the 2-OG utilizing GS-GOGAT cycle [5]. In the future, it will also be interesting to study how the 2-PG and 2-OG signaling pathways are modulated by, or respond to, environmental factors other than carbon and nitrogen availability, such as light/dark transition, and light quantity or quality. Other nutritional, developmental, or environmental factors may also shape C/N balance and the underlying signaling pathways. As an illustration to this idea, the C/N ratio differs in vegetative cells and heterocysts in *Anabaena* PCC 7120, with heterocysts accumulating a higher level of nitrogen reserves, consistent with their dedicated function of N₂ fixation for the whole filaments [73].

Cyanobacteria are a diverse group of prokaryotes. Marine cyanobacteria from oligotrophic oceans live in a much stable environment, with less nutritional variations than those from freshwater habitats. Consequently, many of these marine strains such as

Prochlorococcus species have evolved streamlined genomes, including the loss of many regulatory features [74]. Similarly, the interaction between NtcA and its target DNA in these strains become much less responsive to 2-OG, as a consequence of their long-term adaptation to the rather stable environment [75,76].

Tools used for and learned from the study of 2-OG and 2-PG signaling

The study of 2-OG and 2-PG signaling benefited and will continue to benefit from various tools established in the past. For example, a series of 2-OG analogs have been designed and synthesized for investigating the signaling role of 2-OG [29,37,39,40] (Fig. 2). In addition to the nonmetabolizable analog DFPA (described above), which gives a F-NMR signal with little background in living organisms, other 2-OG analogs include those that can mimic either the ketone form or the ketal form of 2-OG [37], which exist in equilibrium in living organisms. It was found that 2-MPA (Fig. 2a), a mimic of the ketone form of 2-OG, can play the same signaling role as 2-OG in the cyanobacterium *Anabaena* PCC 7120, while other analogs corresponding to mimics of the ketal form of 2-OG were unable to do so [37]. Attempts were also made to develop an affinity column to enrich or purify 2-OG receptors and binding proteins [77]. To that end, the structure-activity relationship of 2-OG signaling has been explored in order to determine to what extent 2-OG can be modified without affecting its signaling functions [37–40,77]. Two analogs, DFMPA and DMPA, obtained after modification of 2-OG at the C2 and C4 positions at the same time (Fig. 2a), still retained a signaling function in *Anabaena* PCC 7120, like 2-OG [38,39]. Based on this information, DMPA was conjugated to a solid resin by click chemistry, and the affinity resin obtained was used to purify the 2-OG receptor NtcA, which was overexpressed in *E. coli* [77]. Given the fact that 2-OG binding proteins, such as PII, NtcA abd NdhR, do not share a common motif for 2-OG recognition [43,46,50,56], different analogs of 2-OG may be conjugated to a solid resin for identification of new 2-OG binding proteins or receptors.

When 2-OG or its analogs are used, one problem is how to get them into the cells so that the intracellular pool of 2-OG can be manipulated. So far, two methods have proved successful. The first relies on the expression of the *kgtP* gene from *E. coli*, which encodes a permease for C5-compounds including 2-OG [29,58,78]. The second is the use of the membrane-permeable analog of 2-OG, dmOG (Fig. 2a), which is commercially available. Under the action of esterases, dmOG is converted to 2-OG in vivo, and hence the level of 2-OG can

be increased within the cells of either *E. coli* or *Synechocystis* PCC 6803 after treatment with dmOG [30,46].

Changes in the 2-OG pool in cells may be fast and may vary on an individual cell basis according to cell age or cell cycle progression. Thus, it is interesting to measure the 2-OG level and follow its changes in live single cells, and in real time. Several 2-OG biosensors based on the technique of FRET (fluorescence resonance energy transfer) have been developed [79–82]. These 2-OG biosensors rely on either 2-OG-dependent protein-protein interaction or protein conformational changes induced by 2-OG binding. At least in vitro, FRET efficiency of these 2-OG biosensors is quantitatively correlated to 2-OG concentrations.

Compared to the variety of tools available for studies on 2-OG signaling, far fewer have been developed for 2-PG. A 2-PG transporter system has been described in *Arabidopsis thaliana* [83], which may be used to increase the 2-PG pool in the cells. Overexpression of a gene encoding a phosphoglycolate phosphatase was proven to be successful for decreasing the internal pool of 2-PG in cyanobacteria [46,47].

A number of studies, including those reviewed here, revealed that 2-OG is not only an essential metabolite for biosynthesis, but also a signaling molecule in bacteria, plants, animals and humans [2,25,29–31,43,46]. Therefore, tools developed to study 2-OG signaling are of considerable interest to the scientific community in general.

Concluding remarks and future perspectives

Cyanobacteria are the ancestors of plastids, and they thus have much in common with algae and higher plants in terms of C/N metabolic coupling. Although the sensors/receptors involved may differ, the signalling nature of 2-OG and 2-PG is very likely a common feature in photosynthetic organisms. The signalling function of 2-OG is already well established in bacteria, plants, animals and humans, although the detailed mechanism in most cases remains poorly understood, except the conservation of the PII pathway in bacteria and plants, although distinct features have evolved for PII in algae and plants [2,35]. Several metabolites and amino acids affect nitrogen metabolism in plants, but it remains unclear whether they are directly involved in signalling in nitrogen control [84]. The signal for carbon limitation in plants is also not clear, but 2-PG and other metabolites from photorespiration and the CBB cycle are the best candidates in light of the current studies. Indeed, it has been shown in *Arabidopsis thaliana* that a *pglp1* mutant defective in the gene encoding 2-PG phosphatase accumulates

a high level of 2-PG but much less starch than the wild type [85]. These results are consistent with a possible role of 2-PG in signalling a carbon limitation. It is also reported recently that the Krebs cycle display a strikingly coordinated regulation upon nitrogen starvation in the diatom *Phaeodactylum tricornutum*, and this control is mediated through a conserved transcription factor bZIP14 [86]. These data clearly establish a certain parallel between eukaryotic algae and cyanobacteria in terms of nitrogen control. It would therefore be interesting to identify the receptors and the corresponding signalling pathways in plants. The concepts developed so far based on studies in cyanobacteria, and the various tools that are available, could help to stimulate research in this direction in plants.

As we commented in this review, increasing numbers of studies have provided evidence that photorespiration and the Krebs cycle are not only important metabolic pathways but also critical players in signalling in C/N metabolic control. Understanding the underlying mechanism of C/N balance is important not only for fundamental research but also for applied research. For example, bioenergy research largely consists of shifting the C/N balance towards the production of carbon-storage compounds for use as biofuel, and understanding different control points in C/N balance could therefore be instrumental in increasing the efficiency of biofuel production. Better understanding of the C/N balance could also be instructive for improving crop production, because properly engineering the control points of C/N balance could optimize the efficiency of the uptake, assimilation, and metabolism of carbon and nitrogen nutrients. Remarkably, a recent study demonstrates that two protein partners GRF4 and DELLA coordinate plant growth and C/N metabolism, and points to new breeding strategies for agriculture [87]. The broad scope of and the general interest in this field, combined with the essential function of C/N metabolic control, will promote further investigations and allow us to answer challenging questions in the future.

Acknowledgements

Work related to the topics in the authors' laboratories are supported by the Key Research Program of Frontier Sciences of the Chinese Academy of Sciences Grant QYZDJ-SSW-SMC016, the National Natural Science Foundation of China (www.nsfc.gov.cn) Grants 31630001, and the Strategic Priority Research Program of the Chinese Academy of Sciences Grant XDB08020300 and United States Department of Energy (DOE), Office of Science, Basic

Energy Sciences, grant no. DE-FG02-08ER15968. We thank Isabel Hanson for English corrections.

References

- 1 Giordano, M. (2013) Homeostasis: An underestimated focal point of ecology and evolution. *Plant Science* 211, 92–101
- 2 Chellamuthu, V.R. *et al.* (2013) From cyanobacteria to plants: Conservation of PII functions during plastid evolution. *Planta* 237, 451–462
- 3 Herrero, A. *et al.* (2004) Cellular differentiation and the NtcA transcription factor in filamentous cyanobacteria. *FEMS Microbiol. Rev.* 28, 469–487
- 4 Burnap, R. *et al.* (2015) Regulation of CO₂ Concentrating Mechanism in Cyanobacteria. *Life* 5, 348–371
- 5 Klähn, S. *et al.* (2018) A glutamine riboswitch is a key element for the regulation of glutamine synthetase in cyanobacteria. *Nucleic Acids Res.* DOI: 10.1093/nar/gky709
- 6 Du, J. *et al.* (2014) Characterisation of cyanobacterial bicarbonate transporters in *E. coli* shows that SbtA homologs are functional in this heterologous expression system. *PLoS One* 9, e115905
- 7 Selim, K.A. *et al.* (2018) P_{II}-like signaling protein SbtB links cAMP sensing with cyanobacterial inorganic carbon response. *Proc. Natl. Acad. Sci. U. S. A.* 115, 4861–4869
- 8 Ponce-Toledo, R.I. *et al.* (2017) An early-branching freshwater cyanobacterium at the origin of plastids. *Curr. Biol.* 27, 386–391
- 9 Rippka, R. *et al.* (1979) Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *J. Gen. Microbiol.* 111, 1–61
- 10 Zhang, C.C. *et al.* (1989) Molecular and genetical analysis of the fructose-glucose transport system in the cyanobacterium *Synechocystis* PCC6803. *Mol Microbiol* 3, 1221–1229
- 11 Bauwe, H. *et al.* (2010) Photorespiration: players, partners and origin. *Trends Plant Sci.* 15, 330–336
- 12 Moroney, J. V. *et al.* (2013) Photorespiration and carbon concentrating mechanisms: Two adaptations to high O₂, low CO₂ conditions. *Photosynth. Res.* 117, 121–131
- 13 Hagemann, M. and Bauwe, H. (2016) Photorespiration and the potential to improve

557 photosynthesis. *Curr. Opin. Chem. Biol.* 35, 109–116

558 14 Turmo, A. *et al.* (2017) Carboxysomes: metabolic modules for CO₂ fixation. *FEMS*
559 *Microbiol. Lett.* 364, fnx176

560 15 Price, G.D. *et al.* (2013) The cyanobacterial CCM as a source of genes for improving
561 photosynthetic CO₂ fixation in crop species. *J. Exp. Bot.* 64, 753–768

562 16 Young, J.D. *et al.* (2011) Mapping photoautotrophic metabolism with isotopically non
563 stationary ¹³C flux analysis. *Metab. Eng.* 13, 656–665

564 17 Xiong W. *et al.* (2015) The plasticity of cyanobacterial metabolism supports direct CO₂
565 conversion to ethylene. *Nat. Plants* 1, 15053

566 18 Zhang, C.C. *et al.* (2006) Heterocyst differentiation and pattern formation in
567 cyanobacteria: A chorus of signals. *Mol. Microbiol.* 59, 367–375

568 19 Herrero, A. *et al.* (2016) The multicellular nature of filamentous heterocyst-forming
569 cyanobacteria. *FEMS Microbiol Rev* 40, 831–854

570 20 Schneegurt, M.A. *et al.* (2000) Metabolic rhythms of a diazotrophic cyanobacterium,
571 *Cyanothece* sp. strain ATCC 51142, heterotrophically grown in continuous dark. *J.*
572 *Phycol.* 36, 107–117

573 21 Muro-Pastor, M.I. *et al.* (2005) Ammonium assimilation in cyanobacteria. *Photosynth.*
574 *Res.* 83, 135–150

575 22 Ohashi, Y. *et al.* (2011) Regulation of nitrate assimilation in cyanobacteria. *J. Exp. Bot.*
576 62, 1411–1424

577 23 Shih, P.M. *et al.* (2013) Improving the coverage of the cyanobacterial phylum using
578 diversity-driven genome sequencing. *Proc. Natl. Acad. Sci. U. S. A.* 110, 1053–1058

579 24 Reitzer, L. (2003) Nitrogen assimilation and global regulation in *Escherichia coli*. *Annu.*
580 *Rev. Microbiol.* 57, 155–76

581 25 Huergo, L.F. and Dixon, R. (2015) The emergence of 2-oxoglutarate as a master
582 regulator metabolite. *Microbiol. Mol. Biol. Rev.* 79, 419–435

583 26 Fulda, S. *et al.* (2006) Proteome analysis of salt stress response in the cyanobacterium
584 *Synechocystis* sp. strain PCC 6803. *Proteomics* 6, 2733–2745

585 27 Stöckel, J. *et al.* (2011) Diurnal rhythms result in significant changes in the cellular
586 protein complement in the cyanobacterium *Cyanothece* 51142. *PLoS One* 6, e16680

587 28 Muro-Pastor, M.I. *et al.* (2001) Cyanobacteria perceive nitrogen status by sensing
588 intracellular 2-oxoglutarate levels. *J. Biol. Chem.* 276, 38320–38328

589 29 Laurent, S. *et al.* (2005) Nonmetabolizable analogue of 2-oxoglutarate elicits
590 heterocyst differentiation under repressive conditions in *Anabaena* sp. PCC 7120.
591 *Proc. Natl. Acad. Sci. U. S. A.* 102, 9907–9912

592 30 Doucette, C.D. *et al.* (2011) α -ketoglutarate coordinates carbon and nitrogen
593 utilization via enzyme I inhibition. *Nat. Chem. Biol.* 7, 894–901

594 31 He, W. *et al.* (2004) Citric acid cycle intermediates as ligands for orphan G-protein-
595 coupled receptors. *Nature* 429, 188–193

596 32 Haas, R. *et al.* (2016) Intermediates of metabolism: From bystanders to signalling
597 molecules. *Trends Biochem. Sci.* 41, 460–471

598 33 You, C. *et al.* (2013) Coordination of bacterial proteome with metabolism by cyclic
599 AMP signalling. *Nature* 500, 301–306

600 34 Araujo, W.L. *et al.* (2014) 2-Oxoglutarate: linking TCA cycle function with amino acid,
601 glucosinolate, flavonoid, alkaloid, and gibberellin biosynthesis. *Front. Plant Sci.* 5, 552

602 35 Uhrig, R.G. *et al.* (2009) PII in higher plants: a modern role for an ancient protein. ,
603 *Trends Plant Sci.* 14, 505–511

604 36 Rabinowitz, J.D. and Silvavy, T.J. (2013) Systems biology: metabolite turns master
605 regulator. *Nature* 500, 283–4

606 37 Chen, H. *et al.* (2006) Studying the signaling role of 2-oxoglutaric acid using analogs
607 that mimic the ketone and ketal forms of 2-oxoglutaric acid. *Chem. Biol.* 13, 849–856

608 38 Liu, X. *et al.* (2013) Structural requirements of 2-oxoglutaric acid analogues to mimic
609 its signaling function. *Org. Lett.* 15, 4662–4665

610 39 Liu, X. *et al.* (2011) 2-Difluoromethylene-4-methylenepentanoic acid, a paradoxical
611 probe able to mimic the signaling role of 2-oxoglutaric acid in cyanobacteria. *Org. Lett.*
612 13, 2924–2927

613 40 Wang, Y. *et al.* (2014) Mimicking the 2-oxoglutaric acid signalling function using
614 molecular probes: insights from structural and functional investigations. *Org. Biomol.*
615 *Chem.* 12, 4723–9

616 41 Ziarelli, F. *et al.* (2012) High resolution magic angle spinning NMR to investigate
617 ligand-receptor binding events for mass-limited samples in liquids. *J. Pharm. Biomed.*
618 *Anal.* 59, 13–17

619 42 Chen, H. *et al.* (2013) ^{19}F NMR: a valuable tool for studying biological events. *Chem.*
620 *Soc. Rev.* 42, 7971

621 43 Zhao, M.-X. *et al.* (2010) Structural basis for the allosteric control of the global
622 transcription factor NtcA by the nitrogen starvation signal 2-oxoglutarate. *Proc. Natl.*
623 *Acad. Sci. U. S. A.* 107, 12487–92

624 44 Klähn, S. *et al.* (2015) Integrated transcriptomic and metabolomic characterization of
625 the low-carbon response using an *ndhR* mutant of *Synechocystis* sp. PCC 6803. *Plant*
626 *Physiol.* 169, 1787–1806

627 45 Wang, H.L. *et al.* (2004) Alterations in global patterns of gene expression in
628 *Synechocystis* sp. PCC 6803 in response to inorganic carbon limitation and the
629 inactivation of *ndhR*, a LysR Family Regulator. *J. Biol. Chem.* 279, 5739–5751

630 46 Jiang, Y.-L. *et al.* (2018) Coordinating carbon and nitrogen metabolic signaling through
631 the cyanobacterial global repressor NdhR. *Proc. Natl. Acad. Sci. U. S. A.* 115, 403–408

632 47 Haimovich-Dayana, M. *et al.* (2015) Does 2-phosphoglycolate serve as an internal signal
633 molecule of inorganic carbon deprivation in the cyanobacterium *Synechocystis* sp. PCC
634 6803? *Environ. Microbiol.* 17, 1794–1804

635 48 Daley, S.M.E. *et al.* (2012) Regulation of the cyanobacterial CO₂-concentrating
636 mechanism involves internal sensing of NADP⁺ and α -ketoglutarate levels by
637 transcription factor CcmR. *PLoS One* 7, e41286

638 49 Huergo, L.F. *et al.* PII signal transduction proteins: Nitrogen regulation and beyond. ,
639 *FEMS Microbiology Reviews*, 37. (2013) , 251–283

640 50 Fokina, O. *et al.* (2010) Mechanism of 2-oxoglutarate signaling by the *Synechococcus*
641 *elongatus* PII signal transduction protein. *Proc. Natl. Acad. Sci. U. S. A.* 107, 19760–
642 19765

643 51 Espinosa, J. *et al.* (2014) PipX, the coactivator of NtcA, is a global regulator in
644 cyanobacteria. *Proc. Natl. Acad. Sci. U. S. A.* 111, E2423–E2430

645 52 Llcer, J.L. *et al.* (2007) The crystal structure of the complex of PII and acetylglutamate
646 kinase reveals how PII controls the storage of nitrogen as arginine. *Proc. Natl. Acad.*
647 *Sci. U. S. A.* 104, 17644–17649

648 53 Maheswaran, M. *et al.* (2004) Complex formation and catalytic activation by the PII
649 signaling protein of N-acetyl-L-glutamate kinase from *Synechococcus elongatus* strain
650 PCC 7942. *J. Biol. Chem.* 279, 55202–55210

651 54 Espinosa, J. *et al.* (2006) Interaction network in cyanobacterial nitrogen regulation:
652 PipX, a protein that interacts in a 2-oxoglutarate dependent manner with PII and

653 NtcA. *Mol. Microbiol.* 61, 457–469

654 55 Wheatley, N.M. *et al.* (2016) A PII-Like Protein Regulated by Bicarbonate: Structural
655 and Biochemical Studies of the Carboxysome-Associated CPII Protein. *J. Mol. Biol.* 428,
656 4013–4030

657 56 Llacer, J.L. *et al.* (2010) Structural basis for the regulation of NtcA-dependent
658 transcription by proteins PipX and PII. *Proc. Natl. Acad. Sci. U. S. A.* 107, 15397–15402

659 57 Tanigawa, R. *et al.* (2002) Transcriptional activation of NtcA-dependent promoters of
660 *Synechococcus* sp. PCC 7942 by 2-oxoglutarate in vitro. *Proc. Natl. Acad. Sci. U. S. A.*
661 99, 4251–4255

662 58 Li, J.H. *et al.* (2003) An increase in the level of 2-oxoglutarate promotes heterocyst
663 development in the cyanobacterium *Anabaena* sp. strain PCC 7120. *Microbiology* 149,
664 3257–3263

665 59 Eisenhut, M. *et al.* (2008) The photorespiratory glycolate metabolism is essential for
666 cyanobacteria and might have been conveyed endosymbiontically to plants. *Proc.*
667 *Natl. Acad. Sci. U. S. A.* 105, 17199–204

668 60 Takahashi, Y. *et al.* (2004) Roles of CmpR, a LysR family transcriptional regulator, in
669 acclimation of the cyanobacterium *Synechococcus* sp. strain PCC 7942 to low-CO₂ and
670 high-light conditions. *Mol. Microbiol.* 52, 837–845

671 61 Lieman-Hurwitz, J. *et al.* (2009) A cyanobacterial AbrB-like protein affects the
672 apparent photosynthetic affinity for CO₂ by modulating low-CO₂-induced gene
673 expression. *Environ. Microbiol.* 11, 927–936

674 62 Picossi, S. *et al.* (2015) The LysR-type transcription factor PacR is a global regulator of
675 photosynthetic carbon assimilation in *Anabaena*. *Environ. Microbiol.* 17, 3341–3351

676 63 Nishimura, T. *et al.* (2008) Mechanism of low CO₂-induced activation of the *cmp*
677 bicarbonate transporter operon by a LysR family protein in the cyanobacterium
678 *Synechococcus elongatus* strain PCC 7942. *Mol. Microbiol.* 68, 98–109

679 64 Figge, R.M. *et al.* (2001) Characterization and analysis of an NAD(P)H dehydrogenase
680 transcriptional regulator critical for the survival of cyanobacteria facing inorganic
681 carbon starvation and osmotic stress. *Mol. Microbiol.* 39, 455–468

682 65 Woodger, F.J. *et al.* (2007) Transcriptional regulation of the CO₂-concentrating
683 mechanism in a euryhaline, coastal marine cyanobacterium, *Synechococcus* sp. strain
684 PCC 7002: Role of NdhR/CcmR. *J. Bacteriol.* 189, 3335–3347

685 66 Hackenberg, C. *et al.* (2012) Low-carbon acclimation in carboxysome-less and
686 photorespiratory mutants of the cyanobacterium *Synechocystis* sp. strain PCC 6803.
687 *Microbiology* 158, 398–413

688 67 Stanier, R.Y. and Cohen-Bazire, G. (1977) Phototrophic Prokaryotes: The
689 Cyanobacteria. *Annu. Rev. Microbiol.* 31, 225–274

690 68 Conrad, M. *et al.* (2014) Nutrient sensing and signaling in the yeast *Saccharomyces*
691 *cerevisiae*. *FEMS Microbiol. Rev.* 38, 254–299

692 69 Berridge, M.J. *et al.* (2003) Calcium signalling: Dynamics, homeostasis and
693 remodelling. *Nature Rev. Mol. Cell Biol.* 4, 517–529

694 70 Kaplan, A. and Reinhold, L. (1999) CO₂ concentrating mechanisms in microorganisms.
695 *Transport* 50, 539–570

696 71 Lopez-Igual, R. *et al.* (2012) N and C control of ABC-type bicarbonate transporter Cmp
697 and its LysR-type transcriptional regulator CmpR in a heterocyst-forming
698 cyanobacterium, *Anabaena* sp. *Environ. Microbiol.* 14, 1035–1048

699 72 Giner-Lamia, J. *et al.* (2017) Identification of the direct regulon of NtcA during early
700 acclimation to nitrogen starvation in the cyanobacterium *Synechocystis* sp. PCC 6803.
701 *Nucleic Acids Res.* 45, 11800–11820

702 73 Teramoto, T. *et al.* (2018) Soft X-ray imaging of cellular carbon and nitrogen
703 distributions in heterocystous cyanobacterium. *Plant Physiol.* DOI:
704 10.1104/pp.17.01767

705 74 Scanlan, D.J. *et al.* (2009) Ecological Genomics of Marine Picocyanobacteria. *Microbiol.*
706 *Mol. Biol. Rev.* 73, 249–299

707 75 Domínguez-Martín, M.A. *et al.* (2018) Differential NtcA responsiveness to 2-
708 oxoglutarate underlies the diversity of C/N balance regulation in *Prochlorococcus*.
709 *Front. Microbiol.* 8, 1–16

710 76 Domínguez-Martín, M.A. *et al.* (2018) Distinct features of C/N balance regulation in
711 *Prochlorococcus* sp. strain MIT9313. *FEMS Microbiol. Lett.* 365, fnx278

712 77 Wang, Y. *et al.* (2014) A “click” chemistry constructed affinity system for 2-oxoglutaric
713 acid receptors and binding proteins. *Org. Biomol. Chem.* 12, 6470–6475

714 78 Seol, W. and Shatkin, A.J. (1991) *Escherichia coli kgtP* encodes an α -ketoglutarate
715 transporter. *Proc. Natl. Acad. Sci. U. S. A.* 88, 3802–3806

716 79 Chen, H.L. *et al.* (2014) Fluorescence resonance energy transfer based on interaction

of PII and PipX proteins provides a robust and specific biosensor for 2-oxoglutarate, a central metabolite and a signalling molecule. *FEBS J.* 281, 1241–1255

80 Zhang, C. *et al.* (2013) Quantitative monitoring of 2-oxoglutarate in *Escherichia coli* cells by a fluorescence resonance energy transfer-based biosensor. *Appl. Microbiol. Biotechnol.* 97, 8307–8316

81 Lüddecke, J. *et al.* (2017) PII protein-derived FRET sensors for quantification and live-cell imaging of 2-oxoglutarate. *Sci. Rep.* 7, 1–13

82 Lüddecke, J. and Forchhammer, K. (2013) From PII signaling to metabolite sensing: A novel 2-oxoglutarate sensor that details PII - NAGK complex formation. *PLoS One* 8, 1–11

83 South, P.F. *et al.* (2017) Bile acid sodium symporter BASS6 can transport glycolate and is involved in photorespiratory metabolism in *Arabidopsis thaliana*. *Plant Cell* 29, 808–823

84 Gent, L. and Forde, B.G. (2017) How do plants sense their nitrogen status? *J. Exp. Bot.* 68, 2531–2540

85 Timm, S. *et al.* (2012) High-to-low CO₂ acclimation reveals plasticity of the photorespiratory pathway and indicates regulatory links to cellular metabolism of *Arabidopsis*. *PLoS One* 7, e42809

86 Matthijs, M. *et al.* (2017) The transcription factor bZIP14 regulates the TCA cycle in the diatom *Phaeodactylum tricornutum*. *EMBO J.* 61, 316–324

87 Li, S. *et al.* (2018) Modulating plant growth – metabolism coordination for sustainable agriculture. *Nature* 560, 595–560

Figure legends

Fig. 1. Coupling between carbon and nitrogen metabolism. Inorganic carbon, as CO₂ or HCO₃⁻, enters the cells and it is in the form of CO₂ that is assimilated into organic carbons through the CBB cycle. Note that the uptake of CO₂ is energetically driven by specialized forms of the NDH-1 complex in thylakoid membranes and results in the formation of HCO₃⁻ which is transported into Rubisco-containing carboxysomes before being reconverted to CO₂ (see text for details). Fixed carbon through the CBB cycle is further used for biosynthesis or to feed the Krebs cycle (incomplete in cyanobacteria in general due to the lack of the 2-OG

dehydrogenase), which produces intermediates for a variety of carbon compounds including lipids. Different forms of inorganic nitrogen can be used by cyanobacteria, and are assimilated in the form of NH_4^+ through the GS-GOGAT cycle using 2-OG as a carbon skeleton. Glu and Gln, the two amino acids produced from the GS-GOGAT cycle are also important nitrogen donors for the synthesis of a variety of nitrogen-containing compounds.

Fig. 2. (a) 2-OG and its analogs that can be used for the study of 2-OG regulation. 2-OG, 2-oxoglutarate; dmOG, dimethyl 2-oxoglutarate, a membrane permeable analog of 2-OG; DFPA, a fluorinated nonmetabolisable analog of 2-OG, which can be traced in vivo by ^{19}F -NMR; 2-MPA, a mimic of the ketone form of 2-OG; DFMPA, a hybrid of DFPA and 2-MPA; and DMPA, an analog of 2-OG that can be conjugated to resins for enrichment of 2-OG-binding proteins. (b) Schematic representation on the signalling mechanism of NtcA based on structural studies. NtcA is a homodimer and a transcriptional activator, each subunit containing an effector-binding domain (EBD) and a DNA-binding domain (DBD). The two helices in the DBDs (in green) interact with target DNA. When 2-OG binds to NtcA at the EBDs, structural conformation changes are induced, shortening the distance between the two DNA-binding helices from 37 to 34 Å, optimal for interaction with DNA and thus transcriptional control.

Fig. 3. Mechanism of carbon metabolic control.

(a) Outline of the transcriptional control. The activator CmpR and the repressor NdhR are the two major transcriptional factors for the regulation of CCM (CO_2 concentrating mechanism). CmpR controls the *cmp* operon, encoding a high affinity bicarbonate transport system, whereas NdhR represses several inorganic carbon uptake systems, including Ndh-1₃, a high-affinity CO_2 uptake system; SbtA, a high-affinity bicarbonate transport system; and BicA, a low affinity bicarbonate transport system. The signal that regulates CmpR is not yet fully understood. NdhR is under the dual control of 2-PG and 2-OG, acting as an anti-repressor and co-repressor, respectively. When CO_2 is limiting, 2-PG is produced by the oxygenase activity of Rubisco and binds to NdhR. The 2-PG/NdhR complex is unable to fix to DNA, thus the CCM-related operon is expressed, leading to enhanced CO_2 /bicarbonate uptake. If CO_2 level is sufficient, the carboxylase activity of Rubisco is favored, leading to CO_2 fixation. Under such a condition, the 2-PG level is low, while that of 2-OG is high, leading to the formation of the 2-OG/NdhR complex and a stronger repression of the CCM-related gene expression.

(b) Mechanism of action of NdhR. NdhR is a tetramer and a repressor, with two compact subunits (in cyan) and two extended subunits (in orange). Under nitrogen starvation, 2-OG binds at the inter-domain cleft between the RDs (regulatory domain). In the 2-OG bound form, the distance between the two helices at the corresponding DBDs (DNA-binding domains) is 29 Å, favorable for DNA binding, thus transcriptional repression. Under a carbon starvation, 2-PG binds at the intra-domain cleft of the RDs, leading to shortening of the distance between the two DNA-binding helices to 25 Å, unfavorable for DNA binding, thus relieving transcriptional repression.

Fig. 4. A model on the mechanism of C/N metabolic balance in cyanobacteria.

(a) Under nitrogen limitation or carbon oversupply, the 2-OG level increases while that of 2-PG decreases. The 2-OG/NtcA complex activates genes involved in nitrogen uptake and assimilation, while the 2-OG/NdhR complex represses the CCM (CO₂ concentrating mechanism)-related gene expression so that carbon uptake decreases. The joint action of the two transcriptional complexes help the cells to regain C/N balance.

(b) Under carbon limitation or nitrogen oversupply, the 2-OG level drops while that of 2-PG increases. The apo form of NtcA has a much lower DNA binding activity, thus leading to a decrease in nitrogen uptake and assimilation. Meanwhile, the 2-PG/NdhR complex cannot act as a repressor, so that genes related to CCM is expressed. Consequently, the C/N balance is achieved.

Outstanding questions for future research

- The key enzyme for autotrophic growth, Rubisco (ribulose 1,5-bisphosphate carboxylase/oxygenase), is a bifunctional enzyme. What are the regulatory factors that allow cells to switch between CO₂ fixation, catalyzed by the carboxylase activity, or photorespiration, catalyzed by the oxygenase activity?
- How do the 2-phosphoglycolate (2-PG) and 2-oxoglutarate (2-OG) signaling pathways, and hence the C/N balance, respond to changes other than C/N availability, such as light intensity?
- Are there other signals involved in C/N balance in cyanobacteria? If there are, how are different signals integrated to coordinate C/N metabolic balance?
- To what extent are changes in 2-PG and 2-OG levels correlated in cells? How do 2-PG and 2-OG signaling affect other regulations required for C/N balance such as redirection of metabolic fluxes or enzymatic activities?
- To what extent are the signaling mechanisms identified from cyanobacteria conserved in eukaryotic algae and plants?

Figure 2

a