Complete genome sequence of *Methanofervidicoccus* sp. A16, a thermophilic methanogen isolated from Mid Cayman Rise hydrothermal vent

Liping Wang, Qiliang Lai, Mohamed Jebbar, Zongze Shao, Karine Alain

► To cite this version:

Liping Wang, Qiliang Lai, Mohamed Jebbar, Zongze Shao, Karine Alain. Complete genome sequence of *Methanofervidicoccus* sp. A16, a thermophilic methanogen isolated from Mid Cayman Rise hydrothermal vent. *Marine Genomics*, 2020, 53, pp.100768. 10.1016/j.margen.2020.100768 . hal-02995883

HAL Id: hal-02995883

<https://hal.science/hal-02995883>

Submitted on 16 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Complete genome sequence of *Methanofervidicoccus* sp. A16, a thermophilic methanogen isolated from Mid Cayman Rise hydrothermal vent

Liping Wang^{a,c}, Qiliang Lai^{a,c}, Mohamed Jebbar^{b,c}, Zongze Shao^{a,c,*}, Karine Alain^{b,c,*}

^a Key Laboratory of Marine Genetic Resources, Third Institute of Oceanography, Ministry of Natural Resources of PR China, Xiamen 361005, China

^b Univ Brest, CNRS, IFREMER, Laboratoire de Microbiologie des Environnements Extrêmes LM2E, F-29280 Plouzané, France

^c LIA1211 MICROBSEA, Sino-French International Laboratory of Deep-Sea Microbiology, Xiamen-Plouzané, France

ARTICLE INFO

Keywords:

Thermophilic
Methanogenesis
Methanofervidicoccus sp.
Hydrothermal vent

ABSTRACT

Methanofervidicoccus sp. A16 is a novel thermophilic and obligate hydrogenotrophic methanogen isolated from a deep-sea hydrothermal vent chimney sample at the Mid Cayman spreading center, Caribbean Sea. Here we report the complete genome of strain A16, which has one circular chromosome of 1,485,358 bp with a mean G + C content of 35.01 mol%. The complete genome harbors 1442 predicted protein-encoding genes. Genes involved in hydrogenotrophic methane production and N₂ fixation were identified in this genome. This study expands our knowledge of methanogenesis at high temperatures and the involvement of these microorganisms in the carbon and nitrogen cycles of deep-sea hydrothermal environments.

1. Introduction

Methanogenic archaea are strictly anaerobic prokaryotes that are widely distributed in anoxic environments such as wetlands, paddy fields, sediments and digestive tracts (Thauer et al., 2008). They produce methane as the terminal step of organic matter fermentation. Of the 500–600 million tons of methane released into the atmosphere each year, 69% originate from microbial metabolism (Conrad, 2009). Methanogens can use carbon dioxide and H₂, simple carbon compounds (acetate, formate, etc) or methylated compounds (methylamine, dimethylsulfide, choline, etc) as energy and carbon sources, and produce methane as a metabolic waste product. Nowadays, a number of hyperthermophilic or thermophilic methanogenic archaea belonging to the order *Methanococcales* have been isolated from a variety of marine hydrothermal environments, including *Methanococcus igneus* Kol 5^T, *Methanoterris formicicus* Mc-S-70^T, *Methanothermococcus okinawensis* IH1^T, *Methanococcus jannaschii* DSM 2661^T, *Methanocaldococcus infernus* ME^T, *Methanocaldococcus indicus* SL43^T, *Methanocaldococcus villosus* KIN24-T80^T, *Methanocaldococcus bathoardescens* JH146^T and further species (Jones et al., 1983; Burggraf et al., 1990; Jeanthon et al., 1998; Takai et al., 2002; L'Haridon et al., 2003; Takai et al., 2004; Bellack et al., 2011; Stewart et al., 2015).

The A16 genome sequence and its annotation is reported in the study. Strain A16 was isolated from a deep-sea hydrothermal vent

chimney sample collected in a depth of 2295 m from Von Damm vent field (Hole to Hell site) (18°22.6007' N, 81°47.8909' W) at the Mid Cayman spreading center, Caribbean Sea. It was enriched and purified in DSM 141 medium, at 70 °C, under an atmosphere of H₂/CO₂ (80/20, 0.2 MPa). Strain A16 was identified as a potential novel species of the genus *Methanofervidicoccus*, with highest 16S rRNA sequence similarity of 98.6% to *Methanofervidicoccus abyssi* HHB^T, which was reported recently as a novel methanogen belonging to the order *Methanococcales* (Sakai et al., 2019). Strain A16 is an obligate anaerobic archaeon that can use CO₂ and H₂ as the sole carbon and energy sources to produce CH₄.

2. Data description

Genomic DNA of strain A16 was extracted with the QIAGEN Genomic-tip 20/G (QIAGEN, Düsseldorf, Germany) kit following the manufacturer's standard protocol. The complete genome sequence of strain A16 was determined by Shanghai Majorbio Bio-pharm Technology Co., Ltd. (Shanghai, China), using the PacBio RSII platform (Table 1). For whole genome sequencing, a 10 kb DNA fragment library was constructed and sequenced on the PacBio RSII sequencing platform, with one SMRT cell (MajorBio Co., Shanghai, China). After quality control, reads were assembled by HGAP 3.0 (Chin et al., 2013). Gene prediction was performed using Glimmer version 3.02 (<http://>

Abbreviations: ANI, average nucleotide identity; CDS, coding DNA sequences; COG, clusters of orthologous groups; GO, gene ontology; KEGG, Kyoto encyclopedia of genes and genomes; ORFs, open reading frames

* Corresponding authors at: LIA1211 MICROBSEA, Sino-French International Laboratory of Deep-Sea Microbiology, Xiamen-Plouzané, France.

E-mail addresses: shaozz@163.com (Z. Shao), Karine.Alain@univ-brest.fr (K. Alain).

<https://doi.org/10.1016/j.margen.2020.100768>

Received 11 September 2019; Received in revised form 21 January 2020; Accepted 17 March 2020

Available online 26 March 2020

1874-7787/ © 2020 Elsevier B.V. All rights reserved.

Table 1
General features of *Methanofervidicoccus* sp. A16.

Items	Description
General features	
Superkingdom	Archaea
Strain	<i>Methanofervidicoccus</i> sp. A16
Geographic location	Atlantic Ocean: Caribbean Sea, Von Damm vent field (Hole to Hell site) at the Mid Cayman spreading center deep-sea hydrothermal vent chimney sample
Sample type	deep-sea hydrothermal vent chimney sample
Latitude and longitude	18°22.6007' N, 81°47.8909' W
Depth	~2295 m
Collection date	June 2013
Biotic relationship	Free-living
Genome characteristics	
Sequencing platform	PacBio RSII
Assembly method	HGAP 3.0
Genome coverage	650×
Finishing quality	Complete genome
NCBI accession number	CP022242
BioProject	PRJNA391986
BioSample	SAMN07281457
Size (bp)	1,485,358
DNA G + C content (%)	35.01
CDSs	1422
tRNAs	34
16S-23S-5S rRNAs	2-2-2

cbcb.umd.edu/software/glimmer/). In addition, rRNA and tRNA genes were identified using Barnap 0.4.2 and tRNA scan-SE v1.3.1 softwares. Functional annotation was performed using NCBI prokaryotic genome annotation pipeline (Tatusova et al., 2016), the Rapid Annotation using Subsystem Technology (RAST) pipeline (<http://rast.nmpdr.org/>) (Overbeek et al., 2014), and the the MicroScope Microbial Genome

Annotation and Analysis Platform (MaGe) (Vallet et al., 2009). Deduced gene products were blasted against KEGG, COG, String and GO databases.

The complete genome of the strain A16 consists of a single circular chromosome with a total length of 1,485,358 bp and an average G + C content of 35.01 mol% (Fig. 1). A total of 1442 protein-coding sequences (CDSs) were predicted, with a summed up gene length of 1,178,139 bp, giving a coding intensity of 78.71%. Majority of the CDSs (1206/1422, 84.81%) could be assigned to a putative function according to COG categories, while the rest were annotated as hypothetical proteins. 34 tRNA genes for 21 amino acids, two 5S rRNAs, two 16S rRNA, two 23S rRNA, 2 miscellaneous RNAs (misc RNA) and 2 pseudogenes were also identified in the genome.

Strain A16 can use H₂ as electron donor and CO₂ as carbon source for growth and produce CH₄. Genes coding for the full pathway of hydrogenotrophic methanogenesis were identified in the genome, which included formylmethanofuran dehydrogenase (Fwd, CFE53_04850, CFE53_05015, CFE53_06085, CFE53_06090, CFE53_06095, CFE53_06100, CFE53_06105, and CFE53_06110), formylmethanofuran-tetrahydromethanopterin N-formyltransferase (Ftr, CFE53_06700), N5, N10-methenyltetrahydromethanopterin cyclohydrolase (Mch, CFE53_06735), F₄₂₀-dependent methylenetetrahydromethanopterin dehydrogenase (Mtd, CFE53_05635), F₄₂₀-dependent N5, N10-methylenetetrahydromethanopterin reductase (Mer, CFE53_02485), N5-methyltetrahydromethanopterin: coenzyme M methyltransferase (Mtr, CFE53_01150, CFE53_01155, CFE53_01160, CFE53_01170, CFE53_01175, CFE53_01180, CFE53_01185), Methyl coenzyme M reductase (Mcr, CFE53_01190, CFE53_01195, CFE53_01200, CFE53_01205 and CFE53_01210), and heterodisulfide reductase (Hdr, CFE53_02215, CFE53_06410, CFE53_06415, CFE53_06755 and CFE53_06760) (Table S1).

Fig. 1. Circular map of the chromosome in the genome of *Methanofervidicoccus* sp. A16. From the outside to the center: label of genome size, CDSs on forward strand (colored by COG categories), CDSs on reverse strand (colored by COG categories), RNA genes (tRNAs and rRNA operons), DNA G + C content and GC skew. CDSs are depicted in different colors according to COG categories.

At the time of writing, twenty-three genomes belonging to the order *Methanococcales* were available in the NCBI and JGI databases. Based on core genome analysis, strain A16 featured the closest phylogenetic relationship to *Methanofervidicoccus abyssi* HHB^T (Fig. S1). The average nucleotide identity (ANI) value between strain A16 and HHB^T were 85.2%, which was below standard ANI criteria for species identity (95–96%) (Richter and Rosselló-Mora, 2009). Further genome sequence analysis of strain A16 revealed 116 unique genes mainly involved in Energy production and conversion, Amino acid transport and metabolism, Replication, recombination and repair, Cell wall, membrane and envelope biogenesis and Inorganic ion transport and metabolism, when compared with HHB^T (Table S2). A minimum set of six genes, *nifH* (CFE53_01675), *nifD* (CFE53_01660), *nifK* (CFE53_01655), *nifE* (CFE53_01645), *nifN* (CFE53_01640) and *nifB* (CFE53_05660), coding for structural and biosynthetic components of nitrogenase (Santos et al., 2012) are present in the A16 genome, while these are absent in strain HHB^T. In addition, two genes (CFE53_01665 and CFE53_01670) coding for nitrogen-fixation associated regulatory protein P-II are also located in this gene cluster. Comparative genome analyses further showed that the gene cluster for nitrogen fixation also exists in other sixteen strains of the order *Methanococcales*, although they show some differences in gene synteny, as shown in Fig. 2 and Fig. S1. Among these strains, *Methanothermococcus thermolithotrophicus* DSM 2095^T (Belay et al., 1984; Leigh, 2000), *Methanococcus maripaludis* DSM 2067^T (Blank et al., 1995), *M. maripaludis* S2 (Kessler et al., 1998; Santos et al., 2012), *Methanococcus aeolicus* Nankai-3^T (Kendall et al., 2006), *Methanotorricus formicicus* Mc-S-70^T (Takai et al., 2004) and *Methanocaldococcus* sp. FS406-22 (Mehta and Baross, 2006), have been reported as diazotrophic methanogens. In this study, the genome analysis of strain A16 will help to understand the ecological role of this species in carbon and nitrogen cycle in submarine hydrothermal environments.

3. Nucleotide sequence accession number

Belay, N., Sparling, R., Daniels, L., 1984. Dinitrogen fixation by a thermophilic methanogenic bacterium. *Nature* 312, 286–288.

Bellack, A., Huber, H., Rachel, R., Wanner, G., Wirth, R., 2011. *Methanocaldococcus villosus* sp. nov., a heavily flagellated archaeon that adheres to surfaces and forms cell–cell contacts. *Int. J. Syst. Evol. Microbiol.* 61, 1239–1245.

Blank, C., Kessler, P.S., Leigh, J.A., 1995. Genetics in methanogens: transposon insertion mutagenesis of a *Methanococcus maripaludis* nifH gene. *J. Bacteriol.* 177, 5773–5777.

Burggraf, S., Fricke, H., Neuner, A., Kristjansson, J.K., Rouvier, P., Mandelco, L., et al., 1990. *Methanococcus igneus* sp. nov., a novel hyperthermophilic methanogen from a shallow submarine hydrothermal system. *Syst. Appl. Microbiol.* 13, 263–269.

Chin, C., Alexander, D.H., Marks, P., Klammer, A., Drake, J.P., Heiner, C., et al., 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat. Methods* 10, 563–569.

Conrad, R., 2009. The global methane cycle: recent advances in understanding the microbial processes involved. *Environ. Microbiol. Rep.* 1, 285–292.

Jeanthon, C., L'Haridon, S., Reysenbach, A.L., Vernet, M., Messner, P., Sleytr, U.B., Prieur, D., 1998. *Methanococcus infernus* sp. nov., a novel hyperthermophilic lithotrophic methanogen isolated from a deep-sea hydrothermal vent. *Int. J. Syst. Bacteriol.* 48,

- 913–919.
- Jones, W.J., Leigh, J.A., Mayer, F., Woese, C.R., Wolfe, R.S., 1983. *Methanococcus jannaschii* sp. nov., an extremely thermophilic methanogen from a submarine hydrothermal vent. Arch. Microbiol. 136, 254–261.
- Kendall, M.M., Liu, Y., Sieprawskalupa, M., Stetter, K.O., Whitman, W.B., Boone, D.R., 2006. *Methanococcus aeolicus* sp. nov., a mesophilic, methanogenic archaeon from shallow and deep marine sediments. Int. J. Syst. Evol. Microbiol. 56, 1525–1529.
- Kessler, P.S., Blank, C., Leigh, J.A., 1998. The *nif* gene operon of the methanogenic archaeon *Methanococcus maripaludis*. J. Bacteriol. 180, 1504–1511.
- Leigh, J.A., 2000. Nitrogen fixation in methanogens: the archaeal perspective. Curr. Iss. Mol. Biol. 2, 125–131.
- L'Haridon, S., Reysenbach, A.L., Banta, A., Messner, P., Schumann, P., Stackebrandt, E., Jeanthon, C., 2003. *Methanocaldococcus indicus* sp. nov., a novel hyperthermophilic methanogen isolated from the Central Indian Ridge. Int. J. Syst. Evol. Microbiol. 53, 1931–1935.
- Mehta, M.P., Baross, J.A., 2006. Nitrogen fixation at 92 degrees C by a hydrothermal vent archaeon. Science 314, 1783–1786.
- Overbeek, R.A., Olson, R., Pusch, G.D., Olsen, G.J., Davis, J.J., Disz, T., et al., 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). Nucleic Acids Res. 42, 206–214.
- Richter, M., Rosselló-Mora, R., 2009. Shifting the genomic gold standard for the prokaryotic species definition. Proc. Natl. Acad. Sci. U. S. A. 106, 19126–19131.
- Sakai, S., Takaki, Y., Miyazaki, M., Ogawara, M., Yanagawa, K., Miyazaki, J., Takai, K., 2019. *Methanofervidicoccus abyssi* gen. nov., sp. nov., a hydrogenotrophic methanogen, isolated from a hydrothermal vent chimney in the Mid Cayman spreading center, the Caribbean Sea. Int. J. Syst. Evol. Microbiol. 69, 1225–1230.
- Santos, P.C.D., Fang, Z., Mason, S.W., Setubal, J.C., Dixon, R., 2012. Distribution of nitrogen fixation and nitrogenase-like sequences amongst microbial genomes. BMC Genomics 13, 162.
- Stewart, L.C., Jung, J.H., Kim, Y.T., Kwon, S.W., Park, C.S., Holden, J.F., 2015. *Methanocaldococcus bathoardescens* sp. nov., a hyperthermophilic methanogen isolated from a volcanically active deep-sea hydrothermal vent. Int. J. Syst. Evol. Microbiol. 65, 1280–1283.
- Takai, K., Inoue, A., Horikoshi, K., 2002. *Methanothermococcus okinawensis* sp. nov., a thermophilic, methane-producing archaeon isolated from a Western Pacific deep-sea hydrothermal vent system. Int. J. Syst. Evol. Microbiol. 52, 1089–1095.
- Takai, K., Nealson, K.H., Horikoshi, K., 2004. *Methanoterris formicicus* sp. nov., a novel extremely thermophilic, methane-producing archaeon isolated from a black smoker chimney in the Central Indian Ridge. Int. J. Syst. Evol. Microbiol. 54, 1095–1100.
- Tatusova, T., Dicuccio, M., Badretdin, A., Chetvernin, V., Nawrocki, E.P., Zaslavsky, L., et al., 2016. NCBI prokaryotic genome annotation pipeline. Nucleic Acids Res. 44, 6614–6624.
- Thauer, R.K., Kaster, A., Seedorf, H., Buckel, W., Hedderich, R., 2008. Methanogenic archaea: ecologically relevant differences in energy conservation. Nat. Rev. Microbiol. 6, 579–591.
- Vallenet, D., Engelen, S., Mornico, D., Cruveiller, S., Fleury, L., Lajus, A., et al., 2009. MicroScope: a platform for microbial genome annotation and comparative genomics. Database 2009.