Quantitative Magnetic Resonance Imaging Assessment of the Quadriceps Changes during an Extreme Mountain Ultramarathon

Hoai-Thu Nguyen, Thomas Grenier, Benjamin Leporq, Caroline Le Goff, Benjamin Gilles, Sylvain Grange, Rémi Grange, Grégoire Millet, Olivier Beuf, Pierre Croisille, et al.

► To cite this version:

Hoai-Thu Nguyen, Thomas Grenier, Benjamin Leporq, Caroline Le Goff, Benjamin Gilles, et al.. Quantitative Magnetic Resonance Imaging Assessment of the Quadriceps Changes during an Extreme Mountain Ultramarathon. *Medicine and Science in Sports and Exercise*, 2021, 53 (4), pp.869-881. 10.1249/mss.0000000000002535 . hal-02998029

HAL Id: hal-02998029

<https://hal.science/hal-02998029>

Submitted on 3 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quantitative Magnetic Resonance Imaging assessment of the Quadriceps changes during an extreme mountain ultramarathon

Hoai-Thu Nguyen¹, MS, Thomas Grenier², PhD, Benjamin Leporq², PhD, Caroline Le Goff³, MD, Benjamin Gilles⁴, PhD, Sylvain Grange^{1,5}, MD, Rémi Grange^{1,5}, MD, Grégoire P. Millet⁶, Pr, PhD, Olivier Beuf², PhD, Pierre Croisille^{1,5}, Pr, MD, PhD, Magalie Viallon^{1,5}, PhD

¹ Univ-Lyon, UJM-Saint-Etienne, INSA-Lyon, Université Claude Bernard Lyon 1, CNRS, Inserm, CREATIS UMR 5220, U1206, F-42023, Saint-Etienne, France.

² Univ Lyon, INSA-Lyon, Université Claude Bernard Lyon 1, UJM-Saint Etienne, CNRS, Inserm, CREATIS UMR 5220, U1206, F-69621, Villeurbanne, France.

³ Department of Clinical Chemistry, University of Liège, CHU Sart-Tilman, Liège, Belgium.

⁴ LIRMM, Université Montpellier, CNRS, F-34095 Montpellier, France.

⁵ Department of Radiology, Centre Hospitalier Universitaire de Saint Etienne, Avenue Albert Raimond, Université de Saint Etienne, F-42055 Saint Etienne, France.

⁶ Institute of Sport Sciences, University of Lausanne, Lausanne, Switzerland.

Email and postal address of corresponding author:

magalie.viallon@creatis.insa-lyon.fr

Service de radiologie

CHU de Saint Etienne

Avenue Albert Raimond

42055 Saint Etienne Cedex 2.

25 **ABSTRACT**

26 **Introduction/Purpose**

27 Extreme ultra-endurance races are growing in popularity but their effects on skeletal
28 muscles remain mostly unexplored. This longitudinal study explores physiological
29 changes in mountain ultramarathon (MUM) athletes' quadriceps using quantitative
30 magnetic resonance imaging (qMRI) coupled with serological biomarkers.

31 The study aimed to monitor the longitudinal effect of the race and recovery, and to
32 identify local inflammatory and metabolic muscle responses by codetection of biological
33 markers.

34 **Methods**

35 An automatic image processing framework was designed to extract imaging-based
36 biomarkers from qMRI acquisitions of the upper legs of 20 finishers at three time points.
37 The longitudinal impact of the race was demonstrated by analyzing the image markers
38 with dedicated biostatistical analysis.

39 **Results**

40 Our framework allows a reliable calculation of statistical data not only inside the whole
41 quadriceps volume but also within each individual muscle heads. Local changes in MRI
42 parameters extracted from quantitative maps were described and found significantly
43 correlated with principal serological biomarkers of interest. A decrease in the PDFF after
44 the race and a stable paramagnetic susceptibility value were found. Pairwise post hoc
45 tests suggested that the recovery process differs among the muscle heads.

46 **Conclusion**

47 This longitudinal study conducted during a prolonged and extreme mechanical stress,
48 showed that quantitative MRI-based markers of inflammation and metabolic response
49 can detect local changes related to the prolonged exercise, with differentiated
50 involvement of each head of the quadriceps muscle as expected in such eccentric load.
51 Consistent and efficient extraction of the local biomarkers enables to highlight
52 interplay/interactions between blood and MRI biomarkers. This work indeed proposes
53 an automatized analytic framework to tackle the time consuming and mentally
54 exhausting segmentation task of muscle heads in large multi-time-points cohorts.

55 **KEYWORDS**

56 Musculoskeletal; statistical analysis; image processing; functional variation

INTRODUCTION

Mountain ultra-endurance running consists of running/walking outdoors over a distance longer than the traditional marathon on mountain trails with significant upward and downward slopes. This sport has become increasingly popular over the last decade.

Because of the inherent challenge for the body, the mountain ultramarathon (MUM) is considered an outstanding model to investigate the adaptive responses to extreme load and mechanical stress(1) in skeletal muscles. Studies have been carried out to explore the impact of these prolonged cardiovascular and muscle efforts, demonstrating, for instance, multiple visceral changes with inflammatory effects(2–4) or an impairment of neuromuscular function(5). Among all muscles, the quadriceps are particularly targeted because extensive downhill running causes repeated eccentric contractions and high mechanical stress, ultimately leading to muscle damage(6,7).

Magnetic Resonance Imaging (MRI) is a reliable, non-irradiating and non-invasive imaging method for tissue characterization and quantitative assessment of tissue integrity through its magnetic properties. Indeed T2 MR relaxation times, and its corresponding rates R_2 ($=1/T_2$) denote the characteristic time constants of the recovery back toward equilibrium of the xy (transverse) components of the magnetization. After being disturbed, the MR signal source, i.e. nuclear magnetization, does not spontaneously recover very fast, but relies almost entirely on interactions of hydrogen nuclei with the surrounding material to re-equilibrate(8). Because T2 is sensitive to low-frequency components, in tenth of ms, water molecules diffuse and sample many different environments on the cellular level within the timescale of relaxation. In many (although by no means all) situations, very rapid exchange may occur between bulk water and bound and interfacial water in biological systems. This is believed to be the origin for many T2 changes in various pathologies, such as edematous changes following insults to tissue. Changes in tissue water and protein content have been shown to affect relaxation. The measure of

average water proton relaxation rate will depend on how effectively, and at what rate, these effects are spread through the rest of the water population. Moreover, the existence of water in separate compartments that are only slowly exchanging gives rise to more complex behavior(9). Many studies witness that edematous areas in the muscle appear bright in T2-weighted images because the T2 relaxation time becomes longer, and T2 maps show elevated quantitative values in the matching area(10,11). T2 maps have, therefore, been used to monitor inflammation in muscle diseases, before and after medical interventions or treatment(12–14), or to predict return to play(15). T2* is the "effective" T2 resulting from inhomogeneities in the main magnetic field or susceptibility-induced field distortions produced in the tissue, related to the presence of chemical or paramagnetic substances such as fat or hemorrhage(16). However, the first step in decoding the various effects of prolonged exercise stress on the body using qMRI markers is to develop systematic and comprehensive noninvasive exploration strategies to consistently extract the biomarkers reflecting the physiological consequences of such challenges.

From that perspective, magnetic resonance imaging (MRI) appears to be a unique imaging modality for extracting anatomical and structural features of muscle tissue(17,18). Recently, quantitative imaging methods such as chemical shift-encoded MRI(19,20) and MR relaxometry mapping(21,22) have allowed users to understand chemical alterations noninvasively at the imaged pixel size.

Several postprocessing methods need to be sequentially applied to extract a quantitative index from MRI data. These methods include the segmentation of muscle heads on multiple large 3D images, the extraction of image features in each area of interest, and statistical analysis. Each of these steps is an area of research in its own right. Although various alternatives have been proposed to target each of these challenges(17,18), they are rarely applied to cohorts or longitudinal studies.

The aim of the study was to quantify the response of the quadriceps muscle to a MUM's extreme loading conditions with quantitative MRI (qMRI) and to monitor the longitudinal effect of race and recovery. To do so, we propose an automatized and integrated analysis pipeline that enables the extraction of local quantitative imaging markers. Hence, the relation of these new markers with standard blood biomarkers was investigated to provide further insight into the local inflammatory and metabolic muscle responses.

MATERIALS AND METHODS

Subjects and experimental study design

We performed a longitudinal study on the athletes who participated in the "Tor des Géants" MUM in 2014, including three MRI sessions carried out just before the start and, for those who finished the race, immediately upon arrival and after 48 hours of recovery. Biological samples were also drawn at four time points with an additional sample obtained at half-race. The flowchart of the study is shown in Figure 1.

This study was approved by the local ethical committee (Aosta Valley, Azienda USL 101/946), and the experimental plan was conducted in accordance with the Helsinki Declaration (2001). Subjects were recruited through mailing and public announcements to registered runners by race organizers. Exclusion criteria were smoking, substance abuse, regular intake of medications, medical or psychiatric illness, and any contraindication to MRI (e.g., claustrophobia, nonremovable metal devices) or abnormalities detected upon laboratory screening. There were 50 experienced runners who volunteered and provided informed written consent to participate in this study.

The Tor des Géants is a 330 km long ultradistance trail running, with considerable positive/negative elevation changes (+24,000 m) in the Valley of Aosta (Italy). It is considered

one of the most difficult mountain marathon races in the world since the ultra-endurance activity is associated with high altitude exposure and sleep deprivation. The altitude along the course ranges between 322 and 3,300 m, with 25 mountain passes over 2,000 m. The maximum time allowed to complete the race is 150 h, and in 2014, the best performance was 71 h 49 min among 740 starters and 446 (60%) finishers (<http://www.tordesgeants.it/>).

The present experimental design was thus a longitudinal study with repeated assessments at four checkpoints:

- The first point (prerace: Pre) was at the start location. The data collection was performed within 4 days before the race and consisted of MRI acquisition and biological sampling.
- The second point was located halfway through the race (middle: Mid). Only biological sampling was performed.
- The third one was at the arrival of the race (arrival: Post): athletes who finished the race were transported by car to the laboratory and were evaluated (MRI and biological sampling) within 1 h after finishing the race.
- The last point (recovery: Post+3) was 48-72 h after arrival time. Both MRI and biological sampling were acquired.

MRI acquisitions

At Pre, Post and Post+3 time points, MRI acquisitions were performed on site using a mobile 1.5 T MR scanner system (MAGNETOM Avanto, Siemens Healthcare, Erlangen, Germany, located within a truck from Alliance Medical, England). A standard coil configuration was used: a flexible 4-channel array surface coil, combined with 4 elements of the spine coil embedded in the bed, resulting in an 8-channel coil in total.

Three MRI acquisitions of the legs were sequentially performed:

- A three-dimensional (3D) isotropic gradient dual-echo sequence was used. The coronal acquisition included the entire upper leg (from the tibial tuberosity to the anterior superior iliac spine). The reconstruction of the water and fat images from the acquired dual-echo data sets was performed inline (Syngo software, Siemens healthcare, Erlangen, Germany) using a Dixon approach enabling four 3D isotropic in-phase, out-phase, fat and water coronal images to be calculated inline on the MR scanner, hereafter denoted water (W), fat (F), in-phase (IN) and out-phase (OUT) images, respectively.
- A 3D spoiled gradient echo sequence (3D GRE) was used for simultaneous estimation of proton density fat fraction (PDFF), T2*, and internal magnetic susceptibility (χ) quantification. Eight echoes were acquired in the transverse plane with a flyback readout gradient (first echo: 1.58 ms and echo spacing: 2.52 ms). TR and flip angle were adjusted to minimize the T1-related bias. Phase and magnitude images were systematically reconstructed. Prescription of localization was performed using anatomic images from an isotropic 3D gradient echo acquisition. For standardization purposes, the central partition in the z-direction was planned at a 15 cm distance from the upper part of the patella using the sagittal multiplanar reconstruction of the first acquired 3D gradient isotropic sequence.
- A 2D multiecho T2 weighted spin-echo sequence (meSE) with 16 echo times (TEs) ranging from 10 to 178 ms, with its central slice was also planned at the same location as previously described.

Each individual sequence was individually optimized. Anatomical regions explored by each technique are summarized in Figure 2a, and the main MR parameters are listed in Supplemental Digital Content 1, MR imaging acquisition parameters.

MR Image postprocessing

PDFF, χ , and T2* were calculated from 3D-GRE acquisition using the reconstruction method described in (19–22) which encompassed two main steps: fat-water separation (provided parametric T2* and PDFF maps) and magnetic susceptibility quantification. Meanwhile, T2 parametric maps were calculated from the meSE sequences.

An automatic segmentation based on external-force-driven deformable registration was performed on the T1-W images (23) to obtain the quadriceps heads of interest.

Briefly, dedicated in-house postprocessing pipelines were required 1) to simultaneously compute the main parametric maps (fat fraction – PDFF, internal magnetic susceptibility (χ) and T2* from the 3D GRE sequence, T2 parametric map from the meSE sequence) and 2) to correct the results of the automatic muscle segmentation and to adjust their resolutions to different image sequences (Figure 2). These steps allowed us to extract metrics from the four parametric maps: PDFF, χ , T2* and T2. All details about parametric map reconstruction and muscle segmentation are provided in Supplemental Digital Contents 2, quantitative maps computation, and 3, muscle segmentation procedure.

Biological sampling and analysis

Blood and urinary samples were collected at each of the four sessions (Pre, Mid, Post, Post+3) within 10 min after arrival at each key point. Blood samples were drawn from an antecubital vein into a dry, heparinized, or EDTA tube according to the analysis to be performed and immediately centrifuged. Since it was not possible to carry out all the investigations on the same day by point-of-care technologies, plasma and serum were frozen at -80°C within 20 min after blood collection for later analysis of muscle injury markers and biochemical variables.

The hematology parameters (hemoglobin, red blood cells, white blood cells) were directly analyzed by a pocH-100i™ automated hematology analyzer (Sysmex, Villepinte, France).

Cobas 8000 (RocheDiagnostics, Mannheim, Germany) was used to perform serial

determinations for C-reactive protein (CRP), urinary creatinine, creatinine, calcium, chlorine, potassium, sodium, and cholesterol. The osmolality and urinary osmolality were measured on an Arkray Osmo Station OM-6050 (Menarini, Florence, Italy).

All blood biomarkers analyzed in this study are listed in Supplemental Digital Content 4, complete list of 58 biological markers analyzed in the study.

Statistical analysis

Missing values in biological datasets were estimated (see Supplemental Digital Contents 5, biological data preprocessing for missing data) to employ paired statistical tests and for subsequent correlation analysis between MRI and biological markers. The significance level was set at $P < .05$ for all statistical tests.

Longitudinal statistical analysis

For the qMRI longitudinal data, we used a statistical analysis of repeated measures with adaptation to the data normality, as the normal distribution could not be assumed. For each qMRI calculated index of each of the 9 muscle volumes (4 quadriceps muscle heads per leg and the total quadriceps volume), we tested the normality of the data at 3 time points with the Shapiro-Wilk test. For the global effect test, one-way ANOVA designed for repeated measures was conducted for all data normally distributed at all the time points; otherwise, the Friedman test was employed.

While performing ANOVA, the sphericity of the data was verified by using Mauchly's test. If the sphericity assumption was violated, the Greenhouse-Geisser correction method was used on the P-value of ANOVA. After the global effect test, a post hoc test was performed to compare each pair of time points. The type of post hoc test depended on the normality of differences between two time points: dependent t-test for normally distributed differences and

229 Wilcoxon signed rank test otherwise. The obtained P-values were adjusted with the Bonferroni
230 adjustment method for multiple comparisons.

231 A similar strategy was applied to the biological marker data with 58 variables at 4 time points.
232 Relative variations of a biomarker X at each time point ($RV(X_n), n \in \{Mid, Post, Post + 3\}$)
233 compared to time point Pre were also computed with:

$$234 \quad RV(X_n) = \frac{\overline{X_n} - \overline{X_{Pre}}}{\overline{X_{Pre}}} \times 100\%$$

235 where $\overline{X_n}$ is the average value of biomarker X at time point n .

236 *Associations between image markers and biological markers*

237 The potential associations between image markers and biological markers were evaluated by
238 calculating the repeated measure correlation coefficient(24) between each image marker and
239 each biological marker.

240

RESULTS

MR markers: parametric mapping analysis

Of the 50 athletes that were included in the study, 31 finishers completed the race (62%) in an average time of 124 ± 13 h. Among them, only those with quadriceps muscle MRI datasets available at all the time points ($n=24$) and with quantitative maps free of artifacts at all three time points were retained for final analysis ($n=20$). Artifacts leading to data exclusions were mostly motion-related. They were specifically due to uncontrolled motion of the legs typically observed immediately after long races potentially combined with those related to sudden sleep attacks during MR acquisitions. Demographic data and global qMRI metrics collected at each time point are reported in Supplemental Digital Content 6, demographic data and qMRI metrics of the subject population.

The first step was to explore differences among all quadriceps muscle heads. Figure 3 shows a matrix of the t-test results when comparing qMRI metrics (χ , PDFF, T2, T2*) between all muscle heads at the three MR acquisition time points (Pre, Post, Post+3). When focusing only on right/left differences of the corresponding muscle heads, χ , T2* and PDFF showed similar tendency. At the same time, there was no significant difference between right and left muscle heads for χ and T2* at any time point, and PDFF exhibited a significant difference between vastus lateralis (VL) heads only at a single time point (Post+3). On the other hand, the T2 metric showed a different pattern with a significant difference between the rectus femoris (RF) heads of the two legs at the three time points (Pre, Post, Post+3) and a significant difference between the left and right vastus medialis (VM) at time point Pre. Different muscle heads also had different qMRI values, especially the RF, which showed significant differences compared to all the other muscle heads most of the time. These results highlight the need to consider

individual muscle behavior separately and that pooling quadriceps muscle heads may result in a loss of information.

When analyzing temporal changes on the repeated measures of qMRI metrics, we observed a significant time effect (i.e., race effect) on T2* and T2 mean values. This is shown in Figures 4 and 5 with typical source images (see also Supplemental Digital Content 6 - Demographic data and qMRI metrics of the subject population, for summary values and Supplemental Digital Content 7 - Specific longitudinal T2* analysis, for a local analysis of right VI in T2* for an example subject). Both T2* and T2 were significantly longer at arrival for most of the muscle heads. They significantly decreased after recovery for the VM and the vastus intermedius (VI) while not returning to baseline values at that time of measurement.

PDFF and χ showed only a small time effect (Figure 6-7). Pooling all muscles, PDFF had a tendency to decrease after the race (2.88 ± 0.53) compared to baseline (3.16 ± 0.45) while increased to higher values than baseline (3.17 ± 0.46) after 48 h of recovery. When considering each muscle, the time effect reached significance for both VI, right VM and left VL muscle heads (Figure 6). Despite a similar trend as the PDFF, time-effect changes in χ did not reach significance for most muscle heads, except for left VL and left VM (L).

When focusing on T2* and T2 findings, the vastus group exhibited stronger variations than the RF muscle group. The VM and VI had larger changes than the VL or the average of all muscles. T2* values appeared less sensitive to muscle changes than T2, as illustrated in Figure 3, but most of the statistical tests were significant except for some muscle heads on both legs between time points Post and Post+3.

Biological marker analysis

The longitudinal variations in biological markers throughout the race are shown in Supplemental Digital Content 8, Values of urinary and blood biomarkers at 4 time points. As

main blood biomarkers of muscle damage, serum creatine kinase (CK) and myoglobin levels peaked at +6598% and +4159% at Mid, while serum lactate dehydrogenase (LDH) levels peaked at +240% at Post and were elevated at Mid at +197% (all P-values of post hoc test against Pre are less than 0.01, see Figure 8). For most of the biomarkers, the biomarker values at Pre were significantly different from those at the other time points (Mid, Post, Post+3), while the difference between Mid and Post was nonsignificant.

Correlation between qMRI features and main blood biomarkers

We presented our results with a heatmap produced within R software (Supplemental Digital Content 9, Correlation between biological markers and image features extracted from quadriceps muscles in each qMRI quantitative map). An interactive version of this heatmap can be found online (http://amp.pharm.mssm.edu/clustergrammer/viz_sim_mats/5bed953490fbf7585f5680c2/Correlation_Mean.txt) thanks to Clustergrammer(25). Significant correlations between qMRI features and biological markers were found:

- The T2 mean extracted from the whole quadriceps showed a significant time effect at all time points of the race and a relatively strong correlation with blood/urine biomarkers such as myoglobin (0.69), CK (0.78) and LDH (0.72). At the level of individual muscles, the correlation coefficient is higher for VI and VM muscles.

- T2* exhibited similar results but with lower correlation coefficients: 0.54 for myoglobin, 0.57 for CK and 0.61 for LDH.

- Meanwhile, neither the mean global PDFF nor χ showed any significant correlation coefficients with biological markers in this study.

DISCUSSION

While ultraloading events such as MUMs are recognized to endanger the musculoskeletal system, little is known about the exact consequences of the acute biomechanical burden on musculoskeletal tissues. Although (26) recommends the use of MRI as the method of choice for studying endurance-related damage, to date, there has been only one longitudinal observational field cohort study, focusing primarily on changes in overall body composition (27) or morphological tissue damage(28).

The challenge and novelty of this work are two-fold: 1) first, to provide a technological approach specifically developed enabling the extraction of qMRI parameters for longitudinal skeletal muscle studies; 2) then, to demonstrate that qMRI brings unique insights into the response of quadriceps muscle exposed to prolonged mechanical stress of a 330 km ultra-endurance event.

To do so, we built a standardizing and automatic analysis pipeline that segments, calculates, and extracts all qMRI markers in each quadriceps muscle's heads that can be robustly propagated to the different follow-up time points, thus enabling longitudinal monitoring.

Beyond the scope of this work, our first contribution is to empower a comprehensive and multiparametric quantification of MRI biomarkers in thigh skeletal muscle, including T2 and T2* relaxation times, together with fat content (PDFF) and internal magnetic susceptibility χ . Secondly, the analysis framework enabled analysis of each quadriceps muscle head over different time points (see Supplemental Digital Content 6, Demographic data and qMRI metrics of the subject population, and Figures 4-7).

Our main findings are that the cumulative biomechanical stress of a 330 km MUM onto quadriceps muscles translated in our subjects to a clear inflammatory burden with an increase in T2 and T2* relaxation times. On the other hand, muscular metabolic and energy storage

status probed fat content (PDFF) remained relatively stable with a small decrease in PDFF, contrasting with the high energy expenditure that mainly relies on lipids. Below is a detailed discussion of each qMRI finding.

T2 and T2 changes: MR markers of the muscular inflammatory burden*

T2 and T2* changes (Supplemental Digital Content 6, Demographic data and qMRI metrics of the subject population, and Figures 4, 5) are an expression of the well-known sensitivity of qMRI T2 and T2* relaxation time parameters to overall water content variations and water redistribution between intracellular and extracellular space(18,22,29–32). Our findings confirm previous studies that reported with other approaches a significant inflammatory response after MUM(5,33), mainly related to repetitive eccentric muscle contractions resulting in microscopic muscle damage and subsequent edema (5,6,34). In our study, T2 and T2* remained elevated even after a few days of recovery, suggesting that 48-72 h is not sufficient for a full recovery in most subjects and confirming previous findings by Robach *et al*(35).

Relaxometry changes demonstrated at the muscular level were corroborated by a large increase in skeletal myocyte injury-related biomarkers (CK, myoglobin and LDH) that was also measured in our subjects. These blood biomarkers are known to witness microlesions and early signs of cellular inflammatory processes preceding water redistribution from intracellular to extracellular compartments and have also been shown to reflect the extent of cellular damage(33,36). Among the other findings in the biological profiles, WBC activation, particularly neutrophil activation, is known to trigger cellular and humoral inflammatory responses(37) characterized by elevated CRP levels and the presence of leucocytes in the extracellular and extravascular spaces in response to long-term exercise(33,38). The intrinsic strength of MR imaging relies on its capacity to precisely reveal the local changes, while the evident correlation of quantified T2 and T2* parameters with biological markers most likely

could reflect a difference in the temporal response of the compared biomarkers. If the tendency are similar, the amplitude of the correlation could only be due to a difference in the temporal occurrence of the inflammation signature at the blood and tissue level, hence also during the tissue recovery process.

The development of peripheral and muscular edema together with a total body water increase have been reported in the context of ultramarathon running(27), with a 6% increase in total body water after a 1,200 km run over 17 consecutive days and an increase in total body water associated with tissue edema(39). Our measures of the total quadriceps muscle volume (excluding superficial tissues) illustrated the same trend but with a small increase in quadriceps muscle volume (+3.3% at the end of the race and +3.4% after 48-72 h recovery relative to baseline; see Supplemental Digital Content 6, Demographic data and qMRI metrics of the subject population): number of voxels from the 3D GRE sequence, expressing muscle volume). This observation suggests that among the potential mechanisms involved, the T2 and T2* changes are more likely related to water redistribution than to a net muscular tissue water content increase.

An interesting finding is the observable difference in T2 and T2* response among the different quadriceps muscle heads. Indeed, while the VI and VL exhibited both higher T2 and T2* increases, the RF showed almost no variations. These findings confirm previous studies exploring MUM and long-distance running adaptation strategies. Sloniger *et al.*(40) For instance, uphill running requires considerably greater activation of the vastus group and less activation of the RF than horizontal running(40). Ultrarunners are also known to modify their running pattern by increasing their stride frequency and by reducing the vertical oscillations to minimize the load and the associated pain occurring in lower limbs mainly during the eccentric phases(41). Impact forces when running downhill can be moderated by increasing knee flexion at initial contact and reducing stride length(42). This shock attenuation strategy is also

associated with a less pronounced heel strike and a forward leaning trunk. Altogether, these protective mechanical adaptations are likely to be specific to MUM and to differentially impact the quadriceps muscles, inducing less stimulation of the RF compared to the vastus muscles, which is clearly illustrated here.

From a mechanistic point of view, T2 and T2* MR relaxation measures are well established and sensitive markers of exercise response. For instance, Le Rumeur *et al.*(32) have shown that T2 relaxation times can detect different effects of dynamic exercise in trained and untrained subjects and that there was a correlation between T2 increase and work intensity in healthy volunteers. Beyond the observed mean variations already discussed, individual variations observed in T2 and T2* values (Figure 4 for T2* and Figure 5 for T2) along such long-distance running are also probably determined by a complex system of regulatory mechanisms including individual response to inflammatory processes at each compartment level, and the intrinsic differences between T2 and T2* mechanisms can probably provide further insights. Indeed, T2* mapping differentiates from T2 mapping by its ability to characterize the relaxation of the transverse magnetization that is influenced by macroscopic (inhomogeneities of the magnetic field) and mesoscopic (structure of the tissue) magnetic field inhomogeneities(43–47). For example, reduced T2* values have already shown the potential to describe structural alterations suggestive of ischemic alterations, collagen areas or hemorrhage, for instance, in myocardium and extracardiac tissues(48,49). According to this, histopathologic substrates of T2* and a correlation of T2* to flow analyses have been demonstrated(44–47,50). Recently, reduced T2* values have been described in a group of hypertrophic cardiomyopathy patients potentially triggered through relative ischemia(51). In our case, explanation for the observed increased T2* values could be related to increased perfusion and/or local variation of oxymyoglobin and oxyhemoglobin as oxygen suppliers are increased while deoxymyoglobin is decreased. Note

also that increased $T2^*$ contrast will be associated with oxygenated proteins, while deoxygenated proteins that are paramagnetic would reduce local $T2^*$ values.

Contrary to $T2^*$, $T2$ in muscle is known to have three components: ~18% of the signal had a very short $T2$ (10 ms), 67% of the signal had an intermediate $T2$ (~40 ms), and the remaining 15% of the signal had a long $T2$ (~170 ms)(52). The short $T2$ component is the water bound to macromolecules, and the intermediate and long components represent intracellular and interstitial water, respectively. The literature also clearly describes a unidirectional flow of water from the “innermost” to “outermost” of three compartments connected in series (intracellularly bound water -> free intracellular water -> interstitial water -> lymphatic vessels). The latter three are subject to considerable variations as a result of normal physiological changes in the muscle. In our study and due to MR acquisition technical constraints, the relaxation in the tissue could be modeled only as a unique compartment. Using a single exponential to describe $T2$ decay in muscle limits our understanding of the exact contributions of different physiological and biochemical responses to exercise.

PDFF and muscular lipid storage

Our study analyzed noninvasively the local variations in fat content in skeletal muscles during an ultra-endurance running event. With PDFF mapping, MRI is indeed the only noninvasive method that can enable simultaneous, reliable, and repeatable fat fraction estimates within each muscle's heads not achievable by alternative methods such as MR spectroscopy(53).

Overall, muscular fat content showed a small decrease at the end of the race, but the changes were not statistically significant except in the most solicited muscle heads (vastus group) (see Figure 3).

Exploring muscular fat content variations during an MUM was of special interest since lipids are the main source of energy in low- to moderate-intensity exercises, such as long-distance events. Additionally, the highest rates of fat oxidation are observed in endurance athletes below 65%(54,55) which is in accordance with what has been measured on the Tor des Géants MUM by Maufrais et al., who reported an average exercise intensity below 50% of VO_{2max} (56).

Muscle fat content is mainly represented by small quantities of triglycerides (TG) that are present as lipid droplets located between or inside muscle fibers (intramyocellular TG, or IMTG), while most fat storage is mainly located in subcutaneous and deep visceral adipose tissue(57). There is now clear evidence that the IMTG pool acts as a buffer regulating fatty acid (FA) flux in skeletal muscle both during exercise to maintain intracellular FA (metabolite) concentrations and in conditions of increased plasma FA provision(58,59). Our findings are in line with other studies that also reported no or minimal reduction of IMTG using biopsies before and after exercise(60–62). Using 1H -MR spectroscopy, other authors reported larger changes with an ~25% IMTG decrease after 2-3 h of exhaustive running at 50-65% VO_{2max} (63,64). Both approaches have been challenged as suffering from intrinsic bias: the direct biopsy approach is subject to between-biopsy variability (reported up to 23%), while 1H -MR spectroscopy modeling accuracy may be biased by water content changes and assumptions of FA compositions in IMTG(65). All existing studies have explored fat content changes in short-duration (up to 3 h) and high-intensity (up to 80% VO_{2max}) exercises that profoundly differ from MUM, which intrinsically limits comparison.

While the IMTG pool appeared to behave as a buffer to facilitate mitochondrial oxidation and skeletal muscle energy supply, our blood lipid profiles showed a marked reduction in circulating TG and low-density lipoprotein (LDL), since the lipoprotein lipase released by endothelial cells during exercise has been shown to increase FA availability(66). At the same time, we observed an increase in high-density lipoprotein (HDL) cholesterol that coincides

with significant increases in lipoprotein lipase activity(67) as already reported in long-distance exercise(68).

Paramagnetic susceptibility χ

Magnetic susceptibility χ is a physical property that describes the response of a medium being placed within a magnetic field(69). A positive susceptibility value characterizes a paramagnetic medium (O₂, salts) with a magnetic field increase, while most biological tissues, including water and fat, are diamagnetic with negative values inducing a local decrease of the initial magnetic field.

All magnetic susceptibility measured in our subjects and all time points were paramagnetic coherently with the metabolic consumption of oxygen in tissues: the weakly diamagnetic oxyhemoglobin releases strongly paramagnetic O₂ and deoxyhemoglobin molecules(70). The magnetic susceptibility χ remained very stable with only a minimal increase at arrival of 1.75±0.43 ppm vs. 1.49±0.34 ppm at baseline (see Supplemental Digital Content 8, Values of urinary and blood biomarkers at 4 time points), as expected in low-intensity aerobic exercise where the average intensity remains below 50% of VO₂max(56), explaining the lack of significant variations over time. Indeed, measurable changes in magnetic susceptibility-related measures in skeletal muscles reported in the literature occurred only in critical conditions such as cuff compression tests or ischemia(71). Our differences among muscle heads and groups are, however, in line with findings by others(71). This finding is explained by the differences among muscles in slow-twitch oxidative muscle fibers, different capillary density, and myoglobin content.

From a broader perspective, the proposed approach could also serve in diseases for longitudinal monitoring of muscular disorder progression or improvement due to various treatments.

Indeed, the motivation for both athletes and physicians during this study was also to serve and encourage technological developments that could, beyond a deeper understanding of muscle physiology during extreme exercise, also help, in turn, be a crucial adjunct to improve patient diagnosis, follow-up, and care, evaluate the effects of personalized-designed therapies, a highly critical objective not in clinical routine. Indeed, there are several clinical scenarios in which pathologies lead to inflammation or muscle wasting. Moreover, such an approach could be coupled to the other strengths of MRI, including in vivo dynamic imaging of muscle function and kinematics(72–75), muscle microstructure using diffusion MRI or measures of chemical changes associated with metabolic states and glycogen consumption using MR spectroscopy(76).

Limitations

While we have studied the average response for all the subjects, there is an interindividual response heterogeneity which is likely influenced by factors such as age(77), training status(78), performance level, experience, and many others that likely modulated the inflammatory response but could not be taken into account due to our limited sample size.

In this paper, we used simple statistical descriptors to analyze the quantitative indexes extracted at each time point and within each muscle head, while radiomic features could have been explored for each quantitative index.

While considering the results on PDFF, it should be kept in mind that the PDFF values in our study are especially low since our subjects are professional athletes and there are estimation errors in PDFF map computation.

Recently, more specific acquisition methods and modeling, combining ultrashort-TE acquisition(79) and muscle-specific models of diffusion, might enable short-term distinguishing the T2 of each muscle compartment (T2 intracellular, T2 in the membrane, T2 extracellular) and thus elucidating the different causes of changes in the parameters.

Conclusion

Using a model of prolonged and extreme mechanical stress, we showed that a dedicated analysis pipeline enables the extraction of local quantitative imaging markers of inflammation and metabolic response and the detection of local changes related to exercise. Our study confirms that prolonged MUMs induce singular muscle damages compared to other sporting events, while the extreme eccentric load (due to the 24000 m of downhill running) is likely the main trigger. When applied with proposed segmentation and analysis pipeline, qMRI can monitor these changes while differentiating the involvement of the heads of the quadriceps muscle.

There are many potential applications; for example, in sports science, our approach could be used to explore the impact of other exercise challenges or to quantify the effects of training strategies or training programs on specific individuals. Considering that MRI can also provide high-spatial-resolution anatomical images of muscle injuries and trauma, as well as information on muscle microstructure and metabolic variables such as glycogen consumption, qMRI would be a highly relevant noninvasive modality for investigating the relationship among force loss, muscle microstructure, performance and recovery in athletes as well as muscle wasting and disability in patients.

523 **ACKNOWLEDGEMENTS**

524 This work has been performed within the framework of LABEX PRIMES (ANR-11-LABX-
525 0063) of Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-
526 0007) and France Life Imaging (ANR-11-INBS-0006) operated by the French National
527 Research Agency (ANR). H-T.Nguyen is working under doctoral contract with 488 SIS
528 doctoral school of Université de Lyon. The authors also thank warmly all ultra-trailers that
529 volunteered for the MUST project, Dr Guido Girardini from the Valle d'Aosta Regional for his
530 unfailing support and help, Hospital the Aosta Valley Regional Government, the VDA trailers
531 and Courmayeur city representatives for their technical assistance.

532 **CONFLICT OF INTEREST**

533 The authors declare that there is no conflict of interest. The results of the present study do not
534 constitute endorsement by ACSM and are presented clearly, honestly, and without fabrication,
535 falsification, or inappropriate data manipulation.

REFERENCES

1. Millet GP, Millet GY. Ultramarathon is an outstanding model for the study of adaptive responses to extreme load and stress. *BMC Med* [Internet]. 2012;10(1):1–3. Available from: <http://dx.doi.org/10.1186/1741-7015-10-77>
2. Andonian P, Viallon M, Le Goff C, et al. Shear-wave elastography assessments of quadriceps stiffness changes prior to, during and after prolonged exercise: A longitudinal study during an extreme mountain ultra-marathon. *PLoS One*. 2016;11(11):1–21.
3. Zanchi D, Viallon M, Goff C Le, et al. Extreme mountain ultra-marathon leads to acute but transient increase in cerebral water diffusivity and plasma biomarkers levels changes. *Front Physiol*. 2017;7(JAN).
4. Belli T, Macedo DV, De Araújo GG, et al. Mountain ultramarathon induces early increases of muscle damage, inflammation, and risk for acute renal injury. *Front Physiol*. 2018;9:1368.
5. Saugy J, Place N, Millet GY, Degache F, Schena F, Millet GP. Alterations of Neuromuscular Function after the World's Most Challenging Mountain Ultra-Marathon. *PLoS One* [Internet]. 2013;8(6):1–11. Available from: <http://dx.doi.org/10.1371/journal.pone.0065596>
6. Clarkson PM, Hubal MJ. Exercise-induced muscle damage in humans. *Am J Phys Med Rehabil*. 2002;81(11 Supp):S52-69.
7. Guilhem G, Cornu C, Maffiuletti NA, Guével A. Neuromuscular adaptations to isoload versus isokinetic eccentric resistance training. *Med Sci Sports Exerc*. 2013;45(2):326–35.
8. Brown RW, Cheng YCN, Haacke EM, Thompson MR, Venkatesan R. Magnetic

560 Resonance Imaging: Physical Principles and Sequence Design: Second Edition. 2014.

561 9. Saab G, Thompson RT, Marsh GD. Multicomponent T2 relaxation of in vivo skeletal
562 muscle. *Magn Reson Med*. 1999;42:150–7.

563 10. Ababneh ZQ, Ababneh R, Maier SE, et al. On the correlation between T2 and tissue
564 diffusion coefficients in exercised muscle: Quantitative measurements at 3T within the
565 tibialis anterior. *Magn Reson Mater Physics, Biol Med*. 2008;21:273–8.

566 11. Ploutz-Snyder LL, Nyren S, Cooper TG, Potchen EJ, Meyer RA. Different effects of
567 exercise and edema on T2 relaxation in skeletal muscle. *Magn Reson Med*.
568 1997;37:676–82.

569 12. Ran J, Ji S, Morelli JN, Wu G, Li X. T2 mapping in dermatomyositis/polymyositis and
570 correlation with clinical parameters. *Clin Radiol*. 2018;73(12):1057.e13-1057.e8.

571 13. Maillard SM, Jones R, Owens C, et al. Quantitative assessment of MRI T2 relaxation
572 time of thigh muscles in juvenile dermatomyositis. *Rheumatology*. 2004;43(5):603–8.

573 14. Thuny F, Lairez O, Roubille F, et al. Post-conditioning reduces infarct size and edema
574 in patients with ST-segment elevation myocardial infarction. *J Am Coll Cardiol*.
575 2012;59(24):2175–81.

576 15. Biglands JD, Grainger AJ, Robinson P, et al. MRI in acute muscle tears in athletes: can
577 quantitative T2 and DTI predict return to play better than visual assessment? *Eur*
578 *Radiol*. 2020;

579 16. Welsch GH, Hennig FF, Krinner S, Trattnig S. T2 and T2* Mapping. *Current*
580 *Radiology Reports*. 2014. p. 60.

581 17. Froeling M, Oudeman J, Strijkers GJ, et al. Muscle Changes Detected with Diffusion-
582 Tensor Imaging after Long-Distance Running. *Radiology [Internet]*. 2015;274(2):548–

583 62. Available from: <http://dx.doi.org/10.1148/radiol.14140702>

584 18. Maeo S, Ando Y, Kanehisa H, Kawakami Y. Localization of damage in the human leg
585 muscles induced by downhill running. *Sci Rep.* 2017;7(1):5769.

586 19. Leporq B, Le Troter A, Le Fur Y, et al. Combined quantification of fatty infiltration, T
587 1-relaxation times and T 2*-relaxation times in normal-appearing skeletal muscle of
588 controls and dystrophic patients. *Magn Reson Mater Physics, Biol Med [Internet].*
589 2017 Aug;30(4):407–15. Available from: <https://doi.org/10.1007/s10334-017-0616-1>

590 20. Leporq B, Ratiney H, Pilleul F, Beuf O. Liver fat volume fraction quantification with
591 fat and water T1 and T2*estimation and accounting for NMR multiple components in
592 patients with chronic liver disease at 1.5 and 3.0 T. *Eur Radiol.* 2013;23(8):2175–86.

593 21. Tawara N, Nitta O, Kuruma H, Niitsu M, Itoh A. T2 mapping of muscle activity using
594 ultrafast imaging. *Magn Reson Med Sci.* 2011;10(2):85–91.

595 22. Patten C, Meyer RA, Fleckenstein JL. T2 mapping of muscle. *Semin Musculoskelet*
596 *Radiol.* 2003;7(4):297–305.

597 23. Gilles B, De Bourguignon C, Croisille P, Millet G, Beuf O, Viallon M. Automatic
598 segmentation for volume quantification of quadriceps muscle head: a longitudinal
599 study in athletes enrolled in extreme mountain ultra-marathon. In: *ISMRM2016:*
600 *International Society for Magnetic Resonance in Medicine.* Singapore;

601 24. Bakdash JZ, Marusich LR. Repeated measures correlation. *Front Psychol.*
602 2017;42(2):261–7.

603 25. Fernandez NF, Gundersen GW, Rahman A, et al. Clustergrammer, a web-based
604 heatmap visualization and analysis tool for high-dimensional biological data. *Sci Data.*
605 2017;4:170151.

- 606 26. Lucas SJE, Helge JW, Schütz UHW, Goldman RF, Cotter JD. Moving in extreme
607 environments: Extreme loading; carriage versus distance. *Extrem Physiol Med*.
608 2016;5:6.
- 609 27. Schütz UHW, Billich C, König K, et al. Characteristics, changes and influence of body
610 composition during a 4486 km transcontinental ultramarathon: Results from the
611 Transeurope Footrace mobile whole body MRI-project. *BMC Med*. 2013;11(1):122.
- 612 28. Theysohn JM, Kraff O, Maderwald S, et al. MRI of the ankle joint in healthy non-
613 athletes and in marathon runners: Image quality issues at 7.0 T compared to 1.5 T.
614 *Skeletal Radiol*. 2013;42(2):261–7.
- 615 29. Fulford J, Eston RG, Rowlands A V, Davies RC. Assessment of magnetic resonance
616 techniques to measure muscle damage 24 h after eccentric exercise. *Scand J Med Sci*
617 *Sport*. 2014;25(1):e28-39.
- 618 30. Fleckenstein JL. Muscle water shifts, volume changes, and proton T2 relaxation times
619 after exercise. *J Appl Physiol*. 1993;74(4):2047–8.
- 620 31. Fleckenstein JL, Watumull D, McIntire DD, Bertocci LA, Chason DP, Peshock RM.
621 Muscle proton T2 relaxation times and work during repetitive maximal voluntary
622 exercise. *J Appl Physiol*. 1993;74(6):2855–9.
- 623 32. Le Rumeur E, Carre F, Bernard AM, Bansard JY, Rochcongar P, De Certaines JD.
624 Multiparametric classification of muscle T1 and T2 relaxation times determined by
625 magnetic resonance imaging. The effects of dynamic exercise in trained and untrained
626 subjects. *Br J Radiol*. 1994;67(794):150–6.
- 627 33. Skenderi KP, Kavouras SA, Anastasiou CA, Yiannakouris N, Matalas AL. Exertional
628 rhabdomyolysis during a 246-km continuous running race. *Med Sci Sports Exerc*.
629 2006;38(6):1054–7.

- 630 34. Fridén J, Lieber RL. Eccentric exercise-induced injuries to contractile and cytoskeletal
631 muscle fibre components. *Acta Physiol Scand*. 2001;171(2):321–6.
- 632 35. Robach P, Boisson RC, Vincent L, et al. Hemolysis induced by an extreme mountain
633 ultra-marathon is not associated with a decrease in total red blood cell volume. *Scand J*
634 *Med Sci Sport*. 2014;24(1):18–27.
- 635 36. Overgaard K, Lindstrom T, Ingemann-Hansen T, Clausen T. Membrane leakage and
636 increased content of Na⁺-K⁺ pumps and Ca²⁺ in human muscle after a 100-km run. *J*
637 *Appl Physiol*. 2002;92(5):1891–8.
- 638 37. McCarthy DA, Dale MM. The Leucocytosis of Exercise: A Review and Model. *Sport*
639 *Med*. 1988;6(6):333–63.
- 640 38. Hikida RS, Staron RS, Hagerman FC, Sherman WM, Costill DL. Muscle fiber necrosis
641 associated with human marathon runners. *J Neurol Sci*. 1983;59(2):185–203.
- 642 39. Knechtle B, Duff B, Schulze I, Kohler G. A multi-stage ultra-endurance run over 1,200
643 km leads to a continuous accumulation of total body water. *J Sport Sci Med*.
644 2008;7(3):357–64.
- 645 40. Sloniger MA, Cureton KJ, Prior BM, Evans EM. Lower extremity muscle activation
646 during horizontal and uphill running. *J Appl Physiol* [Internet]. 1997;83(6):2073–9.
647 Available from: <http://jap.physiology.org/content/83/6/2073>
- 648 41. Degache F, Morin JB, Oehen L, et al. Running mechanics during the world's most
649 challenging mountain ultramarathon. *Int J Sports Physiol Perform*. 2016;11(5):608–14.
- 650 42. Gottschall JS, Kram R. Ground reaction forces during downhill and uphill running. *J*
651 *Biomech*. 2005;38(3):445–52.
- 652 43. Van Oorschot JWM, Gho JMIH, Van Hout GPJ, et al. Endogenous contrast MRI of

cardiac fibrosis: Beyond late gadolinium enhancement. *J Magn Reson Imaging*. 2015;41(5):1181–9.

44. Van Nierop BJ, Bax NAM, Nelissen JL, et al. Assessment of myocardial fibrosis in mice using a T2*-weighted 3D radial magnetic resonance imaging sequence. *PLoS One*. 2015;10(6):e0129899.

45. Huang S yun, Li X hua, Huang L, et al. T2* Mapping to characterize intestinal fibrosis in crohn’s disease. *J Magn Reson Imaging*. 2018;48(3).

46. Zia MI, Ghugre NR, Connelly KA, et al. Characterizing myocardial edema and hemorrhage using quantitative t2 and t2* mapping at multiple time intervals post ST-segment elevation myocardial infarction. *Circ Cardiovasc Imaging*. 2012;5(5):566–72.

47. Messroghli DR, Moon JC, Ferreira VM, et al. Clinical recommendations for cardiovascular magnetic resonance mapping of T1, T2, T2 and extracellular volume: A consensus statement by the Society for Cardiovascular Magnetic Resonance (SCMR) endorsed by the European Association for Cardiovascular Imagin. *J Cardiovasc Magn Reson*. 2017;19(1):75.

48. Manka R, Paetsch I, Schnackenburg B, Gebker R, Fleck E, Jahnke C. BOLD cardiovascular magnetic resonance at 3.0 tesla in myocardial ischemia. *J Cardiovasc Magn Reson*. 2010;12(1):54.

49. Jahnke C, Gebker R, Manka R, Schnackenburg B, Fleck E, Paetsch I. Navigator-gated 3D blood oxygen leveldependent CMR at 3.0-T for detection of stress-induced myocardial ischemic reactions. *JACC Cardiovasc Imaging*. 2010;3(4):375–84.

50. Aguor ENE, Arslan F, Van De Kolk CWA, et al. Quantitative T2 * assessment of acute and chronic myocardial ischemia/reperfusion injury in mice. *Magn Reson Mater Physics, Biol Med*. 2012;25(5):369–79.

51. Lira FS, Yamashita AS, Uchida MC, et al. Low and moderate, rather than high intensity strength exercise induces benefit regarding plasma lipid profile. *Diabetol Metab Syndr*. 2010;3:31.
52. Belton PS, Jackson RR, Packer KJ. Pulsed NMR studies of water in striated muscle. I. Transverse nuclear spin relaxation times and freezing effects. *BBA - Gen Subj*. 1972;286(1):16–25.
53. Nemeth A, Segrestin B, Leporq B, et al. Comparison of MRI-derived vs. traditional estimations of fatty acid composition from MR spectroscopy signals. *NMR Biomed*. 2018;31(9):e3991.
54. Achten J, Gleeson M, Jeukendrup AE. Determination of the exercise intensity that elicits maximal fat oxidation. *Med Sci Sports Exerc*. 2002;42(4):405–12.
55. Jeukendrup AE, Saris WHM, Wagenmakers AJM. Fat metabolism during exercise: A review - Part II: Regulation of metabolism and the effects of training. *Int J Sports Med*. 1998;19(5):203–302.
56. Maufrais C, Millet GP, Schuster I, Rupp T, Nottin S. Progressive and biphasic cardiac responses during extreme mountain ultramarathon. *Am J Physiol - Hear Circ Physiol* [Internet]. 2016;310(10):H1340–8. Available from: <http://ajpheart.physiology.org/content/310/10/H1340>
57. Spriet LL. Regulation of skeletal muscle fat oxidation during exercise in humans. *Med Sci Sports Exerc*. 2002;34(9):1477–84.
58. Watt MJ, Cheng Y. Triglyceride metabolism in exercising muscle. *Biochim Biophys Acta - Mol Cell Biol Lipids*. 2017;1862(10 Pt B):1250–9.
59. Van Loon LJC. Use of intramuscular triacylglycerol as a substrate source during exercise in humans. *J Appl Physiol*. 2004;97(4):1170–87.

- 701 60. Stankiewicz-Choroszuca B, Górski J. Effect of decreased availability of substrates on
702 intramuscular triglyceride utilization during exercise. *Eur J Appl Physiol Occup*
703 *Physiol.* 1978;40(1):27–35.
- 704 61. Kiens B, Essen-Gustavsson B, Christensen NJ, Saltin B. Skeletal muscle substrate
705 utilization during submaximal exercise in man: effect of endurance training. *J Physiol.*
706 1993;469(459–78).
- 707 62. Guo Z, Burguera B, Jensen MD. Kinetics of intramuscular triglyceride fatty acids in
708 exercising humans. *J Appl Physiol.* 2000;89(5):2057–64.
- 709 63. Krssak M, Petersen KF, Bergeron R, et al. Intramuscular glycogen and
710 intramyocellular lipid utilization during prolonged exercise and recovery in man: A
711 ¹³C and ¹H nuclear magnetic resonance spectroscopy study. *J Clin Endocrinol Metab.*
712 2000;85(2):748–54.
- 713 64. Décombaz J, Schmitt B, Ith M, et al. Postexercise fat intake repletes intramyocellular
714 lipids but no faster in trained than in sedentary subjects. *Am J Physiol - Regul Integr*
715 *Comp Physiol.* 2001;281(3):R760-9.
- 716 65. Watt MJ, Heigenhauser GJF, Spriet LL. Intramuscular triacylglycerol utilization in
717 human skeletal muscle during exercise: Is there a controversy? *J Appl Physiol.*
718 2002;93(4):1185–95.
- 719 66. Kiens B, Roepstorff C, Glatz JFC, et al. Lipid-binding proteins and lipoprotein lipase
720 activity in human skeletal muscle: Influence of physical activity and gender. *J Appl*
721 *Physiol.* 2004;97(4):1209–18.
- 722 67. Ferguson M a, Alderson NL, Trost SG, Essig D a, Burke JR, Durstine JL. Effects of
723 four different single exercise sessions on lipids, lipoproteins, and lipoprotein lipase. *J*
724 *Appl Physiol.* 1998;85(3):1169–74.

68. Frias MA, Virzi J, Golaz O, Gencer B, Mach F, Vuilleumier N. Impact of long distance rowing on biological health: A pilot study. *Clin Biochem.* 2018;52(142–147).
69. Schenck JF. The role of magnetic susceptibility in magnetic resonance imaging: MRI magnetic compatibility of the first and second kinds. *Med Phys.* 1996;23(6):815–50.
70. Wang Y, Liu T. Quantitative susceptibility mapping (QSM): Decoding MRI data for a tissue magnetic biomarker. *Magn Reson Med.* 2015;73(1):82–101.
71. Wang C, Zhang R, Zhang X, et al. Noninvasive measurement of lower extremity muscle oxygen extraction fraction under cuff compression paradigm. *J Magn Reson Imaging.* 2016;43(5):1148–58.
72. Drace JE, Pelc NJ. Measurement of skeletal muscle motion in vivo with phase-contrast MR imaging. *J Magn Reson Imaging.* 1994;4(2):157–63.
73. Pappas GP, Asakawa DS, Delp SL, Zajac FE, Drace JE. Nonuniform shortening in the biceps brachii during elbow flexion. *J Appl Physiol.* 2002;92(6):2381–9.
74. Asakawa DS, Blemker SS, Gold GE, Delp SL. In vivo motion of the rectus femoris muscle after tendon transfer surgery. *J Biomech.* 2002;35(8):1029–37.
75. Zhong X, Epstein FH, Spottiswoode BS, Helm PA, Blemker SS. Imaging two-dimensional displacements and strains in skeletal muscle during joint motion by cine DENSE MR. *J Biomech.* 2008;41(3):532–40.
76. Kent-Braun J a, McCully KK, Chance B. Metabolic effects of training in humans: a ³¹P-MRS study. *J Appl Physiol.* 1990;69(3):1165–70.
77. Manfredi TG, Fielding RA, O'Reilly KP, Meredith CN, Lee HY, Evans WJ. Plasma creatine kinase activity and exercise-induced muscle damage in older men. *Med Sci Sports Exerc.* 1991;23(9):1028–34.

- 748 78. Newton MJ, Morgan GT, Sacco P, Chapman DW, Nosaka K. Comparison of
749 Responses to Strenuous Eccentric Exercise of the Elbow Flexors Between Resistance-
750 Trained and Untrained Men. *J Strength Cond Res.* 2008;22(2):597–607.
- 751 79. Saab G, Thompson RT, Marsh GD. Multicomponent T2 relaxation of in vivo skeletal
752 muscle. *Magn Reson Med.* 1999;42(1):150–7.
- 753 80. Kikinis R, Pieper SD, Vosburgh KG. 3D Slicer: A Platform for Subject-Specific Image
754 Analysis, Visualization, and Clinical Support. In: Jolesz FA, editor. *Intraoperative*
755 *Imaging and Image-Guided Therapy* [Internet]. New York, NY: Springer New York;
756 2014. p. 277–89. Available from: https://doi.org/10.1007/978-1-4614-7657-3_19
- 757
- 758

FIGURE LEGENDS

Figure 1: Flowchart summarizing the main steps of our study. MRI was performed at three time points: before (Pre), and participants who finished the race, immediately after (Post) and ~48-72 h after the race (Post+3). Meanwhile, biological sampling was performed at four time points (an additional sample was performed at half-race (Mid)).

Figure 2: MRI input sequences superposed on the isotropic water image (a), the coronal view of the automatic segmentation displayed with the isotropic water image as a background (b) and its 3D volume (c) rendered with 3DSlicer(80).

Figure 3: t-test matrix with color-coded P-values for multiple comparisons of qMRI metrics (χ , PDFF, T2, T2*) between muscle heads at all three acquisition time points (Pre, Post, Post+3). Abbreviations: VL – Vastus Lateralis, RF – Rectus Femoris, VM – Vastus Medialis, VI – Vastus Intermedius, r – right, l – left. For easy viewing, dotted diagonals highlight inter-leg (right/left) comparisons of the same muscle heads at each time point, while all other boxes are intra-leg and/or inter-head comparisons.

Figure 4: Variation of T2* mean in the individual muscle heads of all finishers with an example of T2* maps at the three MR acquisition time points relative to the race of the same subject. A P-value less than .05 indicates a significant change between two time points. Abbreviations: VL – Vastus Lateralis, RF – Rectus Femoris, VM – Vastus Medialis, VI – Vastus Intermedius.

Figure 5: Variation of T2 mean in the individual muscle heads of all finishers with an example of T2 maps at the three MR acquisition time points relative to the race of the same subject. A P-value less than .05 indicates a significant change between two time points. Abbreviations: VL – Vastus Lateralis, RF – Rectus Femoris, VM – Vastus Medialis, VI – Vastus Intermedius.

Figure 6: Variation of PDFF mean in the individual muscle heads of all finishers with an example of PDFF maps at the three acquisition time points relative to the race of the same

subject. A P-value less than .05 indicates significant changes between two time points.

Abbreviations: VL – Vastus Lateralis, RF – Rectus Femoris, VM – Vastus Medialis, VI –

Vastus Intermedius.

Figure 7: Variation of χ mean in the individual muscle heads of all finishers with an example

of χ maps at the three acquisition time points relative to the race of the same subject. A P-value

less than .05 indicates a significant change between two time points. Abbreviations: VL –

Vastus Lateralis, RF – Rectus Femoris, VM – Vastus Medialis, VI – Vastus Intermedius.

Figure 8: P-values of statistical tests in our longitudinal analysis on the blood and urinary

biomarker datasets. A P-value less than .05 indicates a significant change between two time

points.

793 **Supplemental Digital Contents**

794 SDC1.docx MR imaging acquisition parameters

795 SDC2.docx Quantitative maps computation

796 SDC3.docx Muscle segmentation

797 SDC4.docx List of 58 biological markers

798 SDC5.docx Biological data preprocessing for missing data

799 SDC6.docx Demographic data and qMRI metrics of the subject population

800 SDC7.docx Specific longitudinal T2* analysis

801 SDC8.docx Values of urinary and blood biomarkers at 4 time points

802 SDC9.docx Correlation between biological markers and image features extracted from
803 quadriceps muscles in each qMRI quantitative map