Environmental determinants of leaf litter ant community composition along an elevational gradient

Mélanie Fichaux, Jason Vleminckx, Elodie A Courtois, Jacques Delabie, Jordan Galli, Shengli Tao, Nicolas Labrière, Jérôme Chave, Christopher Baraloto, Jérôme Orivel

► To cite this version:

Mélanie Fichaux, Jason Vleminckx, Elodie A Courtois, Jacques Delabie, Jordan Galli, et al.. Environmental determinants of leaf litter ant community composition along an elevational gradient. Biotropica, 2020, 10.1111/btp.12849 . hal-03001673

HAL Id: hal-03001673

<https://hal.science/hal-03001673>

Submitted on 12 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Environmental determinants of leaf-litter ant community composition along an elevational gradient

Journal:	<i>Biotropica</i>
Manuscript ID	BITR-19-276.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	20-May-2020
Complete List of Authors:	<p>Fichaux, Mélanie; CNRS, UMR Ecologie des Forêts de Guyane (EcoFoG), AgroParisTech, CIRAD, INRA, Université de Guyane, Université des Antilles</p> <p>Vleminckx, Jason; CNRS, UMR Ecologie des Forêts de Guyane (EcoFoG), AgroParisTech, CIRAD, INRA, Université de Guyane, Université des Antilles; Florida International University Department of Biological Sciences, Biological Sciences</p> <p>Courtois, Elodie; Laboratoire Ecologie, Evolution, Interactions des Systèmes Amazoniens (LEEISA), Université de Guyane, CNRS, IFREMER; Centre of Excellence PLECO (Plant and Vegetation Ecology), Department of Biology, University of Antwerp</p> <p>Delabie, Jacques; Centro de Pesquisa do Cacau, Laboratório de Mirmecologia; Universidad Estadual de Santa Cruz, DCAA</p> <p>Galli, Jordan; CNRS, UMR Ecologie des Forêts de Guyane (EcoFoG), AgroParisTech, CIRAD, INRA, Université de Guyane, Université des Antilles; Naturalia Environnement, Site Agroparc, 20 rue Lawrence Durell, BP 31285</p> <p>Tao, Shengli; CNRS, Laboratoire Evolution et Diversité Biologique, Université Paul Sabatier</p> <p>Labrière, Nicolas; CNRS, Laboratoire Evolution et Diversité Biologique, Université Paul Sabatier</p> <p>Chave, Jérôme; CNRS, Laboratoire Evolution et Diversité Biologique, Université Paul Sabatier</p> <p>Baraloto, Christopher; Florida International University Department of Biological Sciences, Biological Sciences</p> <p>Orivel, Jerome; CNRS, UMR Ecologie des Forêts de Guyane (EcoFoG), AgroParisTech, CIRAD, INRA, Université de Guyane, Université des Antilles</p>
Keywords:	ants, climate, elevation, environmental filtering, French Guiana, functional traits, soil composition

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fichaux *et al.*

Environmental Determinants of Ant Community Composition

1 Environmental determinants of leaf-litter ant community composition along an elevational 2 gradient

3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Mélanie Fichaux^{1*}, Jason Vleminckx^{1,2*}, Elodie A. Courtois^{3,4}, Jacques H. C Delabie^{5,6}, Jordan
2 Galli^{1,7}, Shengli Tao⁸, Nicolas Labrière⁸, Jérôme Chave⁸, Christopher Baraloto², Jérôme Orivel¹

¹ CNRS, UMR Ecologie des Forêts de Guyane (EcoFoG), AgroParisTech, CIRAD, INRA,
Université de Guyane, Université des Antilles, Campus agronomique, BP 316, 97379, Kourou
cedex, France

² Department of Biological Sciences, Florida International University 11200 S.W. 8th Street
Miami, FL 33199, USA

³ Laboratoire Ecologie, Evolution, Interactions des Systèmes Amazoniens (LEEISA), Université
de Guyane, CNRS, IFREMER, Cayenne, France

⁴ Department of Biology, Centre of Excellence PLECO (Plant and Vegetation Ecology), University
of Antwerp, Wilrijk, Belgium

⁵ Laboratório de Mirmecologia, CEPEC, CEPLAC, Caixa Postal 7, Itabuna, BA 45600-970, Brazil

⁶ Departamento de Ciências Agrárias e Ambientais, Universidade Estadual de Santa Cruz, Rodovia
Jorge Amado Km 16, Ilheus, BA 45662-900, Brazil

⁷ Naturalia Environnement, Site Agroparc, 20 rue Lawrence Durell, BP 31285, 84911 Avignon
Cedex 9, France

⁸ Laboratoire Evolution et Diversité Biologique UMR 5174, CNRS, Université Paul Sabatier, IRD,
118 route de Narbonne, 31062 Toulouse, France

* Co-first authors

Corresponding author: Mélanie Fichaux, fichaux.mel@gmail.com

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

25 Associate Editor: Jennifer Powers

26 Handling Editor: Stephen Yanoviak

27 Received 24 September 2019; revision accepted 1 June 2020.

For Peer Review Only

Abstract

Ant communities are extremely diverse and provide a wide variety of ecological functions in tropical forests. Here we investigated the abiotic factors driving ant composition turnover across an elevational gradient at Mont Itoupé, French Guiana. Mont Itoupé is an isolated mountain whose top is covered by cloud forests, a biogeographical rarity that is likely to be threatened according to climate change scenarios in the region. We examined the influence of six soil, climatic and LiDAR-derived vegetation structure variables on leaf-litter ant assembly (267 species) across nine 0.12-ha plots disposed at three elevations (ca. 400, 600 and 800m asl). We tested (a) whether species co-occurring within a same plot or a same elevation were more similar in terms of taxonomic, functional and phylogenetic composition, than species from different plots/elevations, and (b) which environmental variables significantly explained compositional turnover among plots. We found that the distribution of species and traits of ant communities along the elevational gradient was significantly explained by a turnover of environmental conditions, particularly in soil phosphorus and sand content, canopy height and mean annual relative humidity of soil. Our results shed light on the role exerted by environmental filtering in shaping ant community assembly in tropical forests. Identifying the environmental determinants of ant species distribution along tropical elevational gradients could help predicting the future impacts of global warming on biodiversity organization in vulnerable environments such as cloud forests.

Key words: ants; climate; elevation; environmental filtering; French Guiana; functional traits; soil composition.

1
2
3 51 **1. INTRODUCTION**
4
5
6 52

7
8 53 Determining how environmental factors drive biodiversity patterns is one of the fundamental
9
10 54 goals in ecology. Predictable changes in diversity and composition of plant and animal
11
12 55 communities are observed along broad elevational or latitudinal gradients (Willig *et al.* 2003,
13
14 56 Hillebrand 2004, McCain & Grytnes 2010). Elevational gradients are particularly valuable to
15
16
17 57 study biodiversity patterns, given that they often span sharp gradients in abiotic conditions even
18
19 58 over small spatial scales (e.g. Hodkinson 2005, Kraft *et al.* 2011, Hoiss *et al.* 2012). For instance,
20
21 59 recurrent patterns observed with increasing elevation involve a continuous decrease of
22
23
24 60 temperature (McCain & Grytnes 2010), or an increase in soil moisture and organic carbon
25
26 61 content (He *et al.* 2016).

27
28 62 Niche models predict that environmental conditions select species with particular traits to
29
30 63 establish and persist within an area (Keddy 1992). The effect of environmental filtering may be
31
32
33 64 studied by comparing observed functional or phylogenetic diversity of communities with those
34
35 65 expected under nulls models generated by drawing species at random from a regional species
36
37
38 66 pool (e.g. Gotelli 2000). At local to landscape scales, homogeneous environmental conditions
39
40 67 should generate assemblages of species that are functionally and phylogenetically (if traits are
41
42 68 conserved) more similar than expected by chance (i.e. functional/phylogenetic clustering,
43
44
45 69 respectively). Several studies examining biodiversity patterns along elevational gradients
46
47 70 reported a functional or phylogenetic overdispersion of species in lowland assemblages, shifting
48
49 71 to a functional or phylogenetic clustering of species in highland assemblages (Graham *et al.*
50
51 72 2009, Machac *et al.* 2011, Dehling *et al.* 2014). These results can be interpreted as the effect of
52
53
54 73 increasing strength of environmental filtering on shaping communities (Purschke *et al.* 2013),
55
56 74 which may be caused by the reduction of temperatures with increasing elevation.
57
58
59
60

Here we aim to measure the variations in ant community composition and to determine the environmental factors that shape these variations along a Neotropical altitudinal gradient. Ants (Hymenoptera: Formicidae) represent an ideal model for studying the determinants of species distribution and coexistence because they are abundant and ecologically dominant in terrestrial ecosystems (Hölldobler & Wilson 1990) and they perform a wide variety of ecological functions such as predation, scavenging and seed dispersal (Folgarait 1998, Del Toro *et al.* 2012). Furthermore, previous studies have shown that ant community composition can change markedly along environmental gradients (e.g. Bihn *et al.* 2010, Yates *et al.* 2011, Arnan *et al.* 2014, Groc *et al.* 2014, Silva & Brandão 2014, Smith *et al.* 2014, Bishop *et al.* 2015, Fontanilla *et al.* 2019). Nevertheless, few studies have investigated changes in ant community composition along elevational gradients in tropical regions (Brühl *et al.* 1999, Dunn *et al.* 2009, Smith *et al.* 2014, Nowrouzi *et al.* 2016). Elevational gradients deserve a special attention because they are characterized by a sharp change of various abiotic conditions, especially temperature and humidity, which are predicted to shape ant species distribution. For instance, Nowrouzi *et al.* (2016) have emphasized marked species turnovers between 600 and 800m, which in their study site corresponded to a transition between lowland and cloud forests. Considering the rise of the cloud layers that is predicted by climate change scenarios and associated increases in temperature and decreases in relative humidity (Helmer *et al.* 2019, Los *et al.* 2019), it is urgent to characterize species distributions in these environments to assess threats to their persistence.

To address this gap, we assess how taxonomic, functional and phylogenetic composition of Neotropical ant assemblages are shaped by different environmental parameters along an elevational gradient. We focused on leaf-litter ants because they are easily sampled with a standardized and generalizable collection protocol and their taxonomy has been well described in the region (Groc *et al.* 2009, Fichaux *et al.* 2019). We collected leaf-litter ants at Mont Itoupé

(French Guiana), a mountain which represents a particular biogeographic interest, because of its isolation and its relatively high elevation (ca. 800m) in the eastern Guiana Shield. The top of the mountain is covered by cloud forests, a biogeographic rarity that is likely to be threatened according to scenarios of climate change in this region. Climatic parameters, such as temperature and humidity, may represent major drivers of ant species distributions (e.g. Sanders *et al.* 2007, Dunn *et al.* 2009, Silva & Brandão 2014, Arnan *et al.* 2015). Variation in habitat characteristics such as nutrient availability, vegetation cover as well as soil texture may also play an important role (Vasconcelos *et al.* 2003, Chen *et al.* 2015, Blatrix *et al.* 2016, Schmidt *et al.* 2016).

We sampled along these gradients to address two main objectives. First, we assessed if ant community structure differed from a random distribution of species along the elevational gradient. A stronger effect of environmental filtering is generally observed at highest elevations, leading to a clustered assemblage structure (Graham *et al.* 2009, Machac *et al.* 2011, Dehling *et al.* 2014, Smith *et al.* 2014). Thus, we expected to observe clustered patterns among the leaf-litter assemblages at Mont Itoupé, as a result of the effects of the environmental filtering particularly at high altitude. Second, we identified the main environmental determinants (soil, climate and vegetation structure as measured by LiDAR) of the taxonomic, functional and phylogenetic composition of ant assemblages. A decrease in ant diversity along elevational gradients is a common pattern (Brühl *et al.* 1999, Machac *et al.* 2011, Longino *et al.* 2014, Fontanilla *et al.* 2019) which may be explained by the variation in climatic or soil parameters. We therefore expected both climate and soil variables to be strongly correlated with variation in leaf-litter ant assemblages' composition. Moreover, ant species turnover has been shown to be highly coordinated with tree species turnover in the region, independently from environmental gradients (Vleminckx *et al.* 2019). We may therefore expect a change of ant species composition with

elevation considering previous evidence that forest structure and composition change along elevational gradients (Swenson *et al.* 2011).

2. METHODS

2.1. Experimental design

We collected ants during the dry season in November 2014, at Mont Itoupé (3°01'10.32"N, 53°04'45.90"W), French Guiana. Mont Itoupé, located in the heart of the National Park of the Amazon in French Guiana, peaks at an altitude of 830m asl, representing one of the highest peaks within 250km. A total of nine 0.12-ha plots were established at approximately 400m, 600m and 800m asl, with three plots per elevation range, spanning a total area of ca. 10km² (Figure 1). The plots were chosen to span the range of variation in both climatic and soil conditions across the site. Mean annual rainfall in the area reached 2584 mm (< 140 mm of difference across elevations) while mean annual temperature ranged from 23.0 at 400m to 21.6°C at 800m (worldclim.com).

Each plot represented an area of 30m x 40m, within which we established a grid system of 20 sampling points separated by at least 10m, according to the Ants of Leaf Litter Protocol described in Agosti & Alonso (2000). At each sampling point, we collected leaf-litter ants using pitfall traps and the mini-Winkler method (for more details, see Bestelmeyer *et al.* 2000). Pitfall traps were left in the ground for 72 hours while the mini-Winkler extractors were installed for 48 hours. Because pitfall traps and mini-Winkler extractors were used as complementary traps, we pooled data from both methods in our analyses. Thus, only a single occurrence was reported for a species collected in both traps at the same sampling point.

2.2. Ant identification

Ants were identified at the species level whenever possible, or assigned to a morpho-species code. Species identification was mainly based on online identification keys published on Antwiki (http://www.antwiki.org/wiki/Category:Identification_key) and keys developed by John T. Longino (<http://ants.biology.utah.edu/AntsofCostaRica.html>). We also used the reference collection of the Laboratory Ecofog (Kourou, French Guiana). For the morphospecies with problematic morphological identification (i.e. species from the genera *Hypoponera*, *Nylanderia*, *Pheidole* and *Solenopsis*), we also sequenced the 16S rRNA barcode (at least three specimens per morphospecies) using the protocol developed by Kocher *et al.* (2016). DNA sequences were compared to a local reference barcode library for the ant species of French Guiana (under development; unpublished data). Samples are housed in the Laboratory Ecofog, with voucher specimens deposited in the Laboratorio de Mirmecologia, Cocoa Research Centre CEPEC/CEPLAC (Itabuna, BA, Brazil) under the references #5761 (mini-Winkler traps) and #5762 (pitfall traps).

2.3. Morphological data

We measured nine morphological attributes (Table 1) using an ocular micrometer accurate to 0.01mm mounted on a Leica M80 dissecting microscope (Leica Microsystems, Heerbrugg, Switzerland). Traits were selected based on their expected link with ecological strategies related to resource use (Table 1). Measures were performed for all the species collected (i.e. 267 species), using at least six randomly selected (minor-caste) workers per species whenever possible (Table S1).

2.4. Phylogenetic data

1
2
3 170 We produced a genus-level phylogenetic tree and calculated phylogenetic distances among all
4
5 171 inventoried genera ($n = 56$) using the phylogenetic tree produced in a recent publication
6
7 172 (Blanchard & Moreau 2016). Because two genera sampled in this study (*Gigantiops* and
8
9 173 *Rasopone*) were missing in the tree of Blanchard & Moreau (2016), our resulting tree contained
10
11 174 54 genera (Figure S1).
12
13
14
15 175

176 2.5. Environmental data

177 We measured environmental parameters from soil, climatic and LiDAR data.

178 2.5.1. Soil data

179 In each plot, soil samples were collected at ten locations from three different soil layers (0-10,
180 10-20, 20-30cm), following the procedure described in Baraloto *et al.* (2011). The 10 samples for
181 each depth were then bulked into a single composite sample of ca. 500g of soil. Composite
182 sample were then transported to the Laboratory Ecofog and dried to reach constant weight at
183 25°C, sieved to 2mm and sent to the INRA Arras soil analysis laboratory (Arras, France) for
184 physical and chemical analyses. A set of nine physico-chemical properties were measured (Table
185 S2): the percentage of sand, silt and clay, soil pH, the percentage of organic matter (OM), the
186 carbon-to-nitrogen ratio (C/N), and the soil P, Na and K contents. Particle size analysis of sand,
187 silt and clay were quantitatively performed by their settling rates in an aqueous solution using a
188 hydrometer. Soil pH was measured in 1 M potassium chloride solution. The organic matter
189 content was determined based on the loss of gases after ignition for 2h at 360°C. The total
190 amount of nitrogen (N) and carbon (C) in all forms in soil were quantitatively analyzed using a
191 dynamic flash combustion system coupled with a gas chromatographic separation system and
192 thermal conductivity detection system. Extractable Phosphorus (P) was assayed using the Olsen
193 method (Olsen *et al.* 1954) based on the extraction of phosphate from the soil by 0.5 N sodium

bicarbonate solution adjusted to pH 8.5. A semi-quantitative method was used to determine the amount of soil exchangeable sodium (Na) and potassium (K) residing on the soil colloid exchange sites by displacement with ammonium acetate solution buffered to pH 7.0.

2.5.2. Climatic data

Air temperature (in °C) and relative humidity (in %) were measured in all plots using micro-environmental sensors (HOBO U23-001) as described in Tymen *et al.* (2017). The weather sensors were placed within the area of each plot except for Plot 6, for which the sensor was placed halfway between Plot 5 and 6 because of a logistical issue. Since this sensor was not far away for the sampled area and at the same elevation, the climatic data were used as a proxy for environmental data of this plot (see Figure 1).

The two parameters (air temperature and relative humidity) were monitored for one year, from 1 January 2015 to 31 December 2015 with one measurement every 30 minutes. Using the local micro-environmental data, we defined four climatic variables for each plot (Table S2): T_{MEAN} – mean annual Temperature, RH_{MEAN} – mean annual Relative Humidity, T_{MIN} – minimum temperature of the coldest month, RH_{MIN} – minimum Relative Humidity of the driest month.

2.5.3. LiDAR data

LiDAR data were acquired in early August 2014 using a LMS-Q560 RIEGL laser range finder (wave length 1550nm) on board an aircraft flying at ca. 600m above the ground. A total of 64 km² were covered with a scan angle ranging from $\pm 25^\circ$. Point density slightly exceeded 20 pts/m² (pulse density ca. 13 pulses/m²), with a ground point density of 0.31 pts/m². Digital Elevation Model (DEM) at 1m resolution was interpolated from ground LiDAR points using Lastools (Insenburg n.d.). Digital Surface Model (DSM) was built using Quick Terrain Modeler

(free trial version, <http://appliedimagery.com>) at the same resolution. Canopy Height Model (CHM) was then calculated as the difference between DSM and DEM. When we ground-surveyed our plots, multiple GPS point positions were acquired for each plot using a hand-held GPS (accuracy 5–10m). To further improve plot geo-referencing, we adjusted plot locations against LiDAR CHM as in Réjou-Méchain *et al.* (2015): the location of all surveyed trees (DBH ≥ 30 cm) inferred from ground positioning was first compared to that deduced from CHM. Tree GPS coordinates were then shifted until best match with the CHM, resulting in horizontal shifts typically of less than 15m. With the improved plot locations, we then derived three variables for the plots from LiDAR data (Table S2): canopy height (C_H), canopy cover (C_{COV}) and the gap fraction (gap). Canopy height was calculated as the mean value of CHM pixels inside each plot. Canopy cover was calculated using LAStools (*lascanopy -cov*), with the height threshold set at 5m for separating canopy and non-canopy points. Gap fraction was also calculated following the approach of Morsdorf *et al.* (2006), which was defined as the percent of vegetation echoes to all (ground and vegetation) echoes.

2.6. Data analysis

All analyses were conducted using R 3.5.1 statistical software (R Core Team 2018). The taxonomic, functional and environmental datasets are available in supplementary material (Tables S1, S2 and S3).

2.6.1. Functional heterogeneity among species

The mean value of traits ($n = 9$) per species were normalized by using a Box-Cox transformation, then all traits (except Weber's length) were standardized by dividing their values by Weber's length to correct for individual body size (see Table 1). A Principal Component Analysis (PCA)

on the species $\times$ traits matrix (hereafter, PCA_{TRAITS}) was then performed to eliminate trait redundancy (see Figure S2), using the R package “FactoMineR” (Lê *et al.* 2008). Subsequent analyses were based on the first three principal components (except for the fourth-corner analysis for which we also used individual traits; see below), which jointly explained 77% of the overall trait inertia (Table S4).

2.6.2. Environmental heterogeneity among plots

Environmental variables were normalized (using Box-Cox transformation) and standardized (z-score transformation) prior to analyses, using the R package “forecast” (Hyndman & Khandakar 2008, Hyndman *et al.* 2019). We then performed a PCA on all environmental variables (n = 17), hereafter the PCA_{ENV}, to examine associations among variables and characterize environmental differences among plots. The difference among the three elevations for each environmental variable was tested using a Kruskal-Wallis test.

2.6.3. Community-wide taxonomic, functional and phylogenetic structure

A community that is taxonomically, functionally or phylogenetically clustered within plot corresponds to a situation where species co-occurring within these plots share more similar taxa, traits, or are phylogenetically closer-related, respectively, than species from different plots. This clustering was quantified using the I_{ST}, τ _{ST} and Π _{ST} statistics which calculate the taxonomic, functional and phylogenetic turnover among plots, respectively (Hardy & Senterre 2007). The three indices are calculated as followed:

$I_{ST} = 1 - TD_w/TD_a$ (Taxonomic turnover) eq. 1

$\tau_{ST} = 1 - FD_w/FD_a$ (Functional turnover) eq. 2

$$\Pi_{ST} = 1 - PD_w/PD_a \text{ (Phylogenetic turnover)} \quad \text{eq. 3}$$

where TD corresponds to the Taxonomic Diversity calculated using Simpson index with species occurrence data; FD and PD correspond, respectively, to the mean Functional Dissimilarity (here, Euclidean) between distinct species and the mean Phylogenetic Distance (mean divergence time based on the phylogenetic tree) between distinct genera (thus FD and PD are based on species/genus presence-absence), within (w) and among (a) plots. τ_{ST} and Π_{ST} thus quantify the relative increase of the mean functional divergence and phylogenetic distance between species sampled among plots versus within plots, respectively. Species, trait and phylogenetic clustering is observed if I_{ST} , τ_{ST} or $\Pi_{ST} > 0$, respectively, while negative values indicate overdispersion (Hardy & Senterre 2007). The overall clustering within plot was calculated using the mean I_{ST} , τ_{ST} and Π_{ST} values among all pairs of plots. These mean values were tested by comparing their observed value with 999 null values obtained under a null model where species occurrences are randomized across plots (Hardy 2008). We then used the pairwise values of these statistics among all pairs ($n = 36$) of different plots in regression models to test the effect of environmental dissimilarity on taxonomic, functional and phylogenetic turnover (see next section). Turnover values were also quantified and tested among pairs of altitudes (three comparisons, 400-600m, 400-800m and 600-800m). These indices were calculated using the R package “spacodiR” (Eastman *et al.* 2013).

2.6.4. Environmental determinants of taxonomic, functional and phylogenetic composition

We used Multiple Regression on distance Matrices (MRM; Lichstein 2007) to quantify and test the association between environmental dissimilarity and taxonomic (I_{ST}), phylogenetic (Π_{ST}) and functional (τ_{ST}) turnover of ant assemblages among plots, using the R package “ecodist” (Goslee

1
2
3 288 & Urban 2007). Statistical significance of each regression coefficient was assessed through
4
5 289 residuals permutation tests ($n = 9,999$ permutations). Pairwise τ_{ST} values corresponded more
6
7 290 exactly to the functional turnover among plots calculated using the three first axes of the
8
9 291 PCA_{TRAITS} together, as well as each of these axes individually. Analyzing the functional turnover
10
11 292 based on individual PCA axes can help identifying significant effects on traits that may be
12
13 293 obscured when all traits are taken into account in the analysis. The examination of trait loadings
14
15 294 on the PCA axes can help identifying those traits (Figure 2). It is worth noting that all of our tests
16
17 295 may suffer from a limited statistical power due to the number of pairwise turnover values ($n = 36$)
18
19 296 investigated in the MRM. Thus, while the significant signals observed may reflect some
20
21 297 ecological reality, the absence of signals should also be taken with caution as they could mean
22
23 298 that our sampling design did not allow capturing a significant association.
24
25
26
27

28 299 Three types of MRM models were used for each of the six response variable (I_{ST} , τ_{ST}
29
30 300 calculated over the three first axes of the PCA_{TRAITS} , τ_{ST} calculated for each of these three axes,
31
32 301 and Π_{ST}). The first model (Model 1) corresponded to a simple regression testing the effect of the
33
34 302 overall environmental (Euclidean) dissimilarity (calculated over the 17 environmental variables)
35
36 303 on each response variable. Model 2 corresponded to a multiple regression model where we tested
37
38 304 the relative effects (regression coefficients) of the dissimilarity of several environmental
39
40 305 variables, using a reduced set of variables with limited collinearity to limit the number of tests.
41
42 306 To do so, we retained ecologically relevant variables with r -Spearman correlation ≤ 0.6 for
43
44 307 climate and soil variables separately (Figures S3-4). In doing so, four soil variables (the
45
46 308 percentage of sand, OM, soil pH and P) and one climatic variable (RH_{MEAN}) were retained for
47
48 309 analyses. For LiDAR data, because the variation in the variables “canopy cover” and “gap
49
50 310 fraction” remained virtually unchanged across plots (see Table S2), we only retained the variable
51
52 311 “canopy height” (C_H) for analyses. This latter variable was positively correlated to the variables
53
54
55
56
57
58
59
60

“canopy cover” and “gap fraction” (Figure S5). In Model 3, we tested the relative effects of the dissimilarity in the scores of the two first axes of the PCA_{ENV}. In each of the three models, the spatial dependence among plots was taken into account by including spatial distance (Euclidean, untransformed) among the explanatory variables.

Finally, to have more accurate insights regarding the associated pairs of environmental variables and traits, we performed a fourth-corner analysis (Legendre *et al.* 1997), using the R packages “ade4” (Dray & Dufour 2007) and “adespatial” (Dray *et al.* 2019). The fourth-corner analysis combines three matrices – plots × environmental variables (R), plots × species occurrences (L) and species × traits (Q) – into a matrix of trait-environment associations. The latter is then used to quantify and test pairwise associations between traits and environmental variables, using 9,999 randomizations with the so-called model “6” which combines models “2” and “4” as described in Dray *et al.* (2014), in order to avoid type-I error inflation.

3. RESULTS

We collected a total of 31,865 ant individuals belonging to 267 species, 56 genera and 11 subfamilies along the elevational gradient at Mont Itoupé (Table S3). The total number of species slightly decreased with increasing elevation (resp. 198, 176 and 161 species at 400m, 600m and 800m asl.). Across plots, the number of observed species ranged from 85 (P6; 600m asl.) to 129 (P9; 400m asl.). The average number of species per sample was higher at the lowest elevations (400m: mean ± sd = 25.45 ± 2.69; 600m: mean ± sd = 20.38 ± 4.48; 800m: mean ± sd = 20.63 ± 1.60). We also calculated the Simpson’s index and found very similar values along the elevational gradient (mean ± sd = 0.98 ± <0.01 for each elevation).

3.1. Functional heterogeneity

The PCA performed on species traits ($\text{PCA}_{\text{TRAITS}}$) showed that the first three axes jointly explained 77% of the overall trait inertia (38%, 26% and 13%, respectively; Figures 2 and S6; Table S4). Plot scores on the first axis were highly positively correlated to HL, HW, PW and negatively to WL. The second axis was mostly associated with FL, SL and EL which were highly positively correlated to each other; the third axis was negatively associated to ML (Figures 2 and S6, Table S4).

3.2. Environmental heterogeneity

The PCA performed on the matrix of environmental data (hereafter, PCA_{ENV}) revealed that axes 1 and 2 explained 60.5% and 20.7% of the overall environmental inertia across the nine plots, respectively (Figure 3; Table S5). Among the variables retained for analyses, PCA scores indicated that relative humidity (RH_{MEAN}) and canopy height (C_{H}) were strongly correlated to axis 1 (resp. $r = 0.93$ and $r = -0.90$), while pH was strongly correlated to axis 2 ($r = 0.85$). We found that the following variables significantly (or almost) varied across elevations (Kruskal-Wallis rank tests): OM ($X^2 = 7.20$, $p = 0.03$), C_{H} ($X^2 = 6.00$, $p = 0.05$), P ($X^2 = 2.40$, $p = 0.06$) and RH_{MEAN} ($X^2 = 5.42$, $p = 0.07$). In contrast, no significant variation was found in sand ($X^2 = 2.40$, $p = 0.30$) and soil pH ($X^2 = 0.61$, $p = 0.74$) across elevations.

3.3. Taxonomic, functional and phylogenetic structure of ant assemblages

The leaf-litter ant assemblages displayed significant taxonomic and trait clustering at the plot level (Table 2). Trait clustering was mostly explained by a significant clustering observed for the scores of the second axis of the $\text{PCA}_{\text{TRAITS}}$ (Table 2), which accounted for 26% of the overall functional inertia. The latter axis (i.e. axis 2 of the $\text{PCA}_{\text{TRAITS}}$) was mostly associated to relative

femur and scape length (Figure 2; Table S4). In contrast, we did not find any significant phylogenetic clustering (Table 2). We further found that the taxonomic clustering was mostly due to differences in species composition between 800 and 600m-elevated plots, and that the functional clustering was mostly due to differences in trait composition between the most elevated plots (800m) and the other plots (Table 2). No phylogenetic clustering signal was found at the plot or the elevation level.

3.4. Effect of environmental variables on taxonomic, functional and phylogenetic composition

To facilitate the reading hereafter, we will sometimes omit to use the term “dissimilarity” when describing that the “dissimilarity of A significantly explains the dissimilarity of B”, and will simply write that “A significantly explains B” but we must keep in mind that we only deal with dissimilarity values among plots in our regression models. The overall environmental dissimilarity significantly explained the taxonomic (I_{ST}) and the overall functional (τ_{ST}) turnover among plots (Table 3) but not the phylogenetic (Π_{ST}) turnover. Moreover, the scores of the first axis (strongly associated to the Weber’s length, the relative head length, the relative head width and the pronotum width) and second axis (strongly associated to the relative scape and the relative femur lengths) of the PCA_{TRAITS} were significantly explained by the overall environmental dissimilarity.

Meanwhile, the plot scores of the first axis of the PCA_{ENV} (which was strongly associated with temperature, relative humidity and vegetation structure; Table S5) significantly explained the variation in the overall taxonomic and functional composition, as well as the variation in the plot scores of axes 1 and 2 of the PCA_{TRAITS} (Table 3). When examining the relative effects of each of the six selected environmental variable (sand, pH, OM, P, RH_{MEAN} and C_H), we found

1
2
3 384 that soil phosphorus content and canopy height significantly explained the taxonomic
4
5 385 composition (I_{ST}) but not the overall functional composition (τ_{ST}), the latter being only explained
6
7 386 by the percentage of sand (with a marginal significance; $p = 0.081$). Although the second axis of
8
9
10 387 the PCA_{TRAITS} displayed the strongest association with environment and with the first axis of the
11
12 388 PCA_{ENV} , it was not significantly explained by any individual environmental variable, despite a
13
14 389 relatively high coefficient value obtained with C_H , which may result from a lack of power. Axis 1
15
16
17 390 and 3 of the PCA_{TRAITS} , however, were both significantly explained by soil phosphorus, while
18
19 391 axis 1 was also significantly explained by pH and axis 3 by the percentage of sand. The
20
21 392 phylogenetic (Π_{ST}) turnover was (marginally) significantly explained by soil pH only.

22
23
24 393 The fourth-corner analysis indicated that the percentage of sand was negatively related to
25
26 394 the relative scape, eye and femur lengths, and to the second and third axes of the PCA_{TRAITS}
27
28 395 (Table 4) which were highly associated to the variation of these three traits, while the third axis
29
30 396 was also strongly correlated to the relative mandible length (Table S4). Phosphorus concentration
31
32 397 was positively related to relative head length, whereas it was negatively related to Weber's length
33
34
35 398 and the first axis of the PCA_{TRAITS} . Mean annual relative humidity was negatively related to
36
37 399 relative scape, eye and femur lengths of ants (Table 4).

40
41
42 401 **4. DISCUSSION**
43
44
45 402

46
47 403 This study highlights the existence of associations between key abiotic environmental variables
48
49 404 and the distribution of species and functional traits among Neotropical ant assemblages at a local
50
51 405 (i.e. 10km²) scale. Ant species and functional composition were mostly explained by canopy
52
53 406 height and the first axis of the PCA_{ENV} which was strongly associated to the relative humidity
54
55
56 407 and temperature (and thus elevation), and to a lesser extent by soil phosphorus concentration,
57
58
59
60

sand and pH (Table 3). The observed within-plot and within-elevation functional clustering (Table 2) likely arose from the filtering of these environmental conditions.

Climate represents a major environmental filter and restricts the number and identity of species that can survive and establish at different locations. Among climatic filters, temperature has been shown to be a key parameter influencing patterns of ant species distributions (e.g. Sanders *et al.* 2007, Dunn *et al.* 2009, Silva & Brandão 2014). Other studies have also reported a relationship between temperature and ant functional composition (Stuble *et al.* 2013, Arnan *et al.* 2014). Relative humidity may be even more constraining for leaf-litter ants than temperature (Kaspari & Weiser 2000, Menke & Holway 2006). These studies reported results that were consistent with our findings. Indeed, we found that the variation in the dissimilarity of plot scores along the first axis of the PCA_{ENV}, which was highly explained by variation in soil relative humidity, temperature and elevation (Figure 3; Table S5), significantly explained the taxonomic and functional turnover. As highlighted by the PCA generated on environmental data (PCA_{ENV}; Figure 3), the top of the mountain (ca. 800m asl), which is covered by cloud forests, is characterized by slightly higher levels of humidity (mean_{RH} = 99.3%) compared to lower elevations (between ca. 400 and 600m asl; mean_{RH} = 96.5%). The top forests may therefore provide a refuge area for certain species preferring relatively colder and more humid conditions. Changes in species and functional composition may thus reflect the presence of climate-specialists at higher elevations. More specifically, our results suggest that smaller ant species, as well as species with relative smaller eyes or smaller legs, were found in sites with higher levels of humidity (thus at higher elevation). These traits are characteristics of hypogaeic ant species, i.e. ant species that nest and forage within the litter or into the soil (Weiser & Kaspari 2006).

Along Mont Itoupé, turnovers in species and functional composition were also associated to variation in phosphorus concentration (Table 3). Phosphorus limitation may explain variation in

plant diversity (Wright *et al.* 2011, Vleminckx *et al.* 2017), which in turn is likely to influence the diversity and density of leaf-litter associated invertebrates (McGlynn & Salinas 2007, Vleminckx *et al.* 2019). In their study, McGlynn and Salinas (2007) found that environments richer in phosphorus had greater litter invertebrate densities (mostly detritivores). Thus, the association between phosphorus availability and the species and functional composition of ant communities may be indirectly explained by the variation in the density of prey for predatory ant species. The fourth-corner analysis performed in this study reveal that phosphorus concentration was positively related to the relative head length and negatively related to body length (i.e. Weber's length), features that are characteristics of small hypogaecic ant species belonging to genera such as *Carebara*, *Solenopsis* and *Strumigenys*. Those predatory ant genera may thus benefit from the higher phosphorus concentrations that may favor the abundance of small prey. Such results contrast with those obtained in a study conducted by Jacquemin *et al.* (2012) in which they found a decrease of predatory ant species in plots enriched in Carbon, Nitrogen and Phosphorus. In their study, predatory ants seemed to be limited by habitat loss due to the increased litter decomposition. The positive effect of higher levels of phosphorus for small predatory ant species found in our study may have been counterbalanced by habitat loss in their study, since the amount of leaf litter at the highest elevation, i.e. at high levels of phosphorus, was much more important than at the lowest ones.

In addition to climatic factors, nutrient availability and soil texture can also influence invertebrate composition by imposing constraints on species living in ground habitats (Vasconcelos *et al.* 2003, Schmidt *et al.* 2016, Costa-Milanez *et al.* 2017), but also indirectly via an effect of these variables on tree community assembly (Vleminckx *et al.* 2019). In our study, a significant association was found between the functional composition and the soil texture (percentage of sand) turnover across plots (Table 3), especially for the third axis of the PCA_{TRAITS}

which was mostly associated with the relative mandible length (Table S4). We lack hypotheses to explain that result. In addition, the percentage of sand was negatively related to individual functional traits of ants (i.e. the relative scape and femur lengths). Particular functional traits of ant species rather than the global functional space of ant assemblages may thus be linked to soil texture. Short scape and eye, relative to body size, are typical of hypogaeic ant species (Brandão *et al.* 2012), foraging and nesting inside the leaf-litter. Sandy soils may thus favor movements and nest establishment of hypogaeic species within the litter, compared to clay-rich soils.

CONCLUSIONS

Elevational gradients stretching from lowland forests to lower mountain cloud forests represent a major source of environmental variation in tropical regions which, in our study, has been shown to influence the distribution of leaf-litter ants. In particular, the taxonomic and functional turnover of ant communities were mostly explained by soil phosphorus content, climatic (temperature, relative humidity) variables and vegetation structure, while trait variation seemed to be also influenced by soil texture and pH. Our results shed light on the unique biodiversity value of cloud forests, a particularly rare ecosystem in the eastern Guiana Shield, likely to be threatened by climate change, with scenarios predicting an intensification of drought events and an increase of temperature across the region (Esquivel-Muelbert *et al.* 2019). Indeed, the congruence between species and functional turnover along the humidity gradient highlights that a loss of species and functional diversity represent real threats in this regional biodiversity hotspot.

ACKNOWLEDGMENTS

We thank Sandrine Etienne for her work on the process of molecular data, and Aurélie Dourdain for producing the map of the site. Thanks are also due to the national park managers for allowing our research program in the core area of the *Parc Amazonien de Guyane*. Financial support for this study was provided by an *Investissement d’Avenir* grant of the *Agence Nationale de la Recherche* (CEBA, ANR-10-LABX-25-01) through a PhD fellowship to MF and the funding of the DIADEMA (Dissecting Amazonian Diversity by Enhancing a Multiple taxonomic-groups Approach) and DIAMOND (Dissecting And MONitoring amazonian Diversity) projects, by the *Programme Convergence* 2007-2013, *Région Guyane* from the European community (BREGA, 757/2014/SGAR/DE/BSF) and by the PO-FEDER 2014-2020, *Région Guyane* (BiNG, GY0007194).

DATA AVAILABILITY

Data available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.00000> (Fichaux *et al.* 2020).

REFERENCES

AGOSTI, D., and L. ALONSO. 2000. The ALL Protocol. *In* D. Agosti, J. Majer, L. Alonso, and T. Schultz (Eds.) *Ants: standard methods for measuring and monitoring biodiversity*. pp. 204–206, Washington D.C. USA.

ARNAN, X., X. CERDÁ, and J. RETANA. 2014. Ant functional responses along environmental gradients. *Journal of Animal Ecology* 83: 1398–1408.

- ARNAN, X., X. CERDÁ, and J. RETANA. 2015. Partitioning the impact of environment and spatial structure on alpha and beta components of taxonomic, functional, and phylogenetic diversity in European ants. *PeerJ* 3: e1241.
- BARALOTO, C., S. RABAUD, Q. MOLTO, L. BLANC, C. FORTUNEL, B. HÉRAULT, N. DÁVILA, I. MESONES, M. RIOS, E. VALDERRAMA, and P. V. A. FINE. 2011. Disentangling stand and environmental correlates of aboveground biomass in Amazonian forests. *Global Change Biology* 17: 2677–2688.
- BESTELMEYER, B. T., D. AGOSTI, L. E. ALONSO, C. R. F. BRANDÃO, W. L. BROWN JR., J. H. C. DELABIE, and R. SILVESTRE. 2000. Field techniques for the study of ground-dwelling ants: an overview, description, and evaluation. *In* D. Agosti, J. Majer, L. Alonso, and T. Schultz (Eds.) *Ants: standard methods for measuring and monitoring biodiversity*. pp. 122–144, Washington D.C. USA.
- BIHN, J. H., G. GEBAUER, and R. BRANDL. 2010. Loss of functional diversity of ant assemblages in secondary tropical forests. *Ecology* 91: 782–792.
- BISHOP, T. R., M. P. ROBERTSON, B. J. VAN RENSBURG, and C. L. PARR. 2015. Contrasting species and functional beta diversity in montane ant assemblages. *Journal of Biogeography* 42: 1776–1786.
- BLANCHARD, B. D., and C. S. MOREAU. 2016. Defensive traits exhibit an evolutionary trade-off and drive diversification in ants. *Evolution* 71: 315–328.
- BLATRIX, R., C. LEBAS, C. GALKOWSKI, P. WEGNEZ, R. PIMENTA, and D. MORICHON. 2016. Vegetation cover and elevation drive diversity and composition of ant communities (Hymenoptera: Formicidae) in a Mediterranean ecosystem. *Myrmecological News* 22: 119–127.
- BRANDÃO, C. R. F., R. R. SILVA, and J. H. C. DELABIE. 2012. Neotropical ants (Hymenoptera) functional groups: nutritional and applied implications. *In* J. R. P. Parra (Ed.) *Insect bioecology and nutrition for integrated pest management*. pp. 213–236, CRC, Boca Raton.

1
2
3 528 BRÜHL, C. A., M. MOHAMED, and K. E. LINSSENMAIR. 1999. Altitudinal distribution of leaf litter
4
5 529 ants along a transect in primary forests on Mount Kinabalu, Sabah, Malaysia. *Journal of Tropical*
6
7 530 *Ecology* 15: 265–277.
8
9
10 531 CHEN, X., B. ADAMS, C. BERGERON, A. SABO, and L. HOOPER-BUI. 2015. Ant community structure
11
12 532 and response to disturbances on coastal dunes of Gulf of Mexico. *Journal of Insect Conservation*
13
14 533 19: 1–13.
15
16
17 534 COSTA-MILANEZ, C. B. DA, J. D. MAJER, P. DE T. A. CASTRO, and S. P. RIBEIRO. 2017. Influence of
18
19 535 soil granulometry on average body size in soil ant assemblages: implications for bioindication.
20
21 536 *Perspectives in Ecology and Conservation* 15: 102–108.
22
23
24 537 DAVIDSON, D. W., S. C. COOK, and R. R. SNELLING. 2004. Liquid-feeding performances of ants
25
26 538 (Formicidae): ecological and evolutionary implications. *Oecologia* 139: 255–266.
27
28
29 539 DEHLING, D. M., S. A. FRITZ, T. TÖPFER, M. PÄCKERT, P. ESTLER, K. BÖHNING-GAESE, and M.
30
31 540 SCHLEUNING. 2014. Functional and phylogenetic diversity and assemblage structure of frugivorous
32
33 541 birds along an elevational gradient in the tropical Andes. *Ecography* 37: 1047–1055.
34
35
36 542 DEL TORO, I., R. R. RIBBONS, and S. L. PELINI. 2012. The little things that run the world revisited:
37
38 543 a review of ant-mediated ecosystem services and disservices (Hymenoptera: Formicidae).
39
40 544 *Myrmecological News* 17: 133–146.
41
42
43 545 DRAY, S., and A.-B. DUFOUR. 2007. The ade4 package: implementing the duality diagram for
44
45 546 ecologists. *Journal of Statistical Software* 22: 1–20.
46
47
48 547 DRAY, S., P. CHOLER, S. DOLÉDEC, P. R. PERES-NETO, W. THUILLER, S. PAVOINE, and C. J. F. TER
49
50 548 BRAAK. 2014. Combining the fourth-corner and the RLQ methods for assessing trait responses to
51
52 549 environmental variation. *Ecology* 95: 14–21.
53
54
55 550 DRAY, S., D. BAUMAN, G. BLANCHET, D. BORCARD, S. CLAPPE, G. GUENARD, T. JOMBART, G.
56
57 551 LAROCQUE, P. LEGENDRE, N. MADI, and H. H. WAGNER. 2019. adespatial: multivariate multiscale

- spatial analysis. R package version 0.3-4, <https://CRAN.R-project.org/package=adespatial>.
- DUNN, R. R. ET AL. 2009. Climatic drivers of hemispheric asymmetry in global patterns of ant species richness. *Ecology Letters* 12: 324–333.
- EASTMAN, J., T. PAINE, and O. HARDY. 2013. *spacodiR*: spatial and phylogenetic analysis of community diversity. R package version 0.13-0115, <https://CRAN.R-project.org/package=spacodiR>.
- ESQUIVEL-MUELBERT, A. ET AL. 2019. Compositional response of Amazon forests to climate change. *Global Change Biology* 25: 39–56.
- FEENER JR., D. H., J. R. B. LIGHTON, and G. A. BARTHOLOMEW. 1988. Curvilinear allometry, energetics and foraging ecology: a comparison of leaf-cutting ants and army ants. *Functional Ecology* 2: 509–520.
- FICHAUX, M., B. BÉCHADE, J. DONALD, A. WEYNA, J. H. C. DELABIE, J. MURIENNE, C. BARALOTO, and J. ORIVEL. 2019. Habitats shape taxonomic and functional composition of Neotropical ant assemblages. *Oecologia* 189: 201–513.
- FICHAUX, M., J. VLEMINCKX, E. A. COURTOIS, J. H. C. DELABIE, J. GALLI, S. TAO, N. LABRIÈRE, J. CHAVE, C. BARALOTO, J. ORIVEL. 2020. Data from: Environmental determinants of leaf-litter ant community composition along an elevational gradient. Dryad Digital Repository. doi:10.5061/dryad.00000
- FOLGARAIT, P. J. 1998. Ant biodiversity and its relationship to ecosystem functioning: a review. *Biodiversity and Conservation* 7: 1221–1244.
- FONTANILLA, A. M., A. NAKAMURA, Z. XU, M. CAO, R. L. KITCHING, Y. TANG, and C. J. BURWELL. 2019. Taxonomic and functional ant diversity along tropical, subtropical, and subalpine elevational transects in Southwest China. *Insects* 128.

1
2
3 575 FOWLER, H. G., L. C. FORTI, C. R. F. BRANDÃO, J. H. C. DELABIE, and H. L. VASCONCELOS. 1991.
4
5 576 Ecologia nutricional de formigas. *In* A. R. Panizzi and J. R. P. Parra (Eds.) Ecologia nutricional de
6
7 577 insetos e suas implicações no manejo de pragas. pp. 131–223, São Paulo, Brazil: Editora Manole.
8
9
10 578 GOSLEE, S. C., and D. L. URBAN. 2007. The ecodist package for dissimilarity-based analysis of
11
12 579 ecological data. *Journal of Statistical Software* 22: 1–19.
13
14
15 580 GOTELLI, N. J. 2000. Null model analysis of species co-occurrence patterns. *Ecology* 81: 2606–
16
17 581 2621.
18
19 582 GRAHAM, C. H., J. L. PARRA, C. RAHBK, and J. A. MCGUIRE. 2009. Phylogenetic structure in
20
21 583 tropical hummingbird communities. *Proceedings of the National Academy of Sciences* 106:
22
23 584 19673–19678.
24
25
26 585 GROC, S., J. ORIVEL, A. DEJEAN, J.-M. MARTIN, M.-P. ETIENNE, B. CORBARA , and J. H. C. DELABIE.
27
28 586 2009. Baseline study of the leaf-litter ant fauna in a French Guianese forest. *Insect Conservation*
29
30 587 and *Diversity* 2: 183–193.
31
32
33 588 GROC, S., J. H. C. DELABIE, F. FERNÁNDEZ, M. LEPONCE, J. ORIVEL, R. SILVESTRE, H. L.
34
35 589 VASCONCELOS, and A. DEJEAN. 2014. Leaf-litter ant communities (Hymenoptera: Formicidae) in
36
37 590 a pristine Guianese rain-forest: Stable functional structure versus high species turnover.
38
39 591 *Myrmecological News* 19: 43–51.
40
41
42 592 GRONENBERG, W., J. TAUTZ, and B. HÖLLDOBLER. 1993. Fast trap jaws and giant neurons in the
43
44 593 ant *Odontomachus*. *Science* 262: 561–563.
45
46
47 594 HARDY, O. J. 2008. Testing the spatial phylogenetic structure of local communities: Statistical
48
49 595 performances of different null models and test statistics on a locally neutral community. *Journal of*
50
51 596 *Ecology* 96: 914–926.
52
53
54 597 HARDY, O. J., and B. SENTERRE. 2007. Characterizing the phylogenetic structure of communities
55
56 598 by an additive partitioning of phylogenetic diversity. *Journal of Ecology* 95: 493–506.
57
58
59
60

- HE, X., E. HOU, Y. LIU, and D. WEN. 2016. Altitudinal patterns and controls of plant and soil nutrient concentrations and stoichiometry in subtropical China. *Scientific Reports* 6: 24261.
- HELMER, E. H., E. A. GERSON, L. S. BAGGETT, B. J. BIRD, T. S. RUZYCKI, and S. M. VOGGESSER. 2019. Neotropical cloud forests and páramo to contract and dry from declines in cloud immersion and frost S. Lötters (Ed.). *PLoS ONE* 14: e0213155.
- HILLEBRAND, H. 2004. On the generality of the latitudinal diversity gradient. *The American Naturalist* 163: 192–211.
- HODKINSON, I. D. 2005. Terrestrial insects along elevation gradients: species and community responses to altitude. *Biological Reviews* 80: 489–513.
- HOISS, B., J. KRAUSS, S. G. POTTS, S. ROBERTS, and I. STEFFAN-DEWENTER. 2012. Altitude acts as an environmental filter on phylogenetic composition, traits and diversity in bee communities. *Proceedings of the Royal Society B: Biological Sciences* 279: 4447–4456.
- HÖLLDOBLER, B., and E. O. WILSON. 1990. *The Ants*. Harvard University Press.
- HYNDMAN, R. J., and Y. KHANDAKAR. 2008. Automatic time series forecasting: the forecast package for R. *Journal of Statistical Software* 27: 1–23.
- HYNDMAN, R., G. ATHANASOPOULOS, C. BERGMEIR, G. CACERES, L. CHHAY, M. O'HARA-WILD, F. PETROPOULOS, S. RAZBASH, E. WANG, and F. YASMEEN. 2019. forecast: forecasting functions for time series and linear models. R package version 8.7, <URL: <http://pkg.robjhyndman.com/forecast>>.
- INSENBURG, M. LAStools—efficient LiDAR processing software (version 160921, academic) obtained from <http://rapidlasso.com/LAStools>.
- JACQUEMIN, J., M. MARAUN, Y. ROISIN, and M. LEPONCE. 2012. Differential response of ants to nutrient addition in a tropical Brown Food Web. *Soil Biology and Biochemistry* 46: 10–17.
- KASPARI, M. 1993. Body size and microclimate use in Neotropical granivorous ants. *Oecologia* 96:

1
2
3 623 500–507.
4
5 624 KASPARI, M., and M. D. WEISER. 1999. The size–grain hypothesis and interspecific scaling in ants.
6
7 625 Functional Ecology 13: 530–538.
8
9
10 626 KASPARI, M., and M. D. WEISER. 2000. Ant activity along moisture gradients in a Neotropical
11
12 627 forest. Biotropica 32: 703–711.
13
14
15 628 KEDDY, P. A. 1992. Assembly and response rules: two goals for predictive community ecology.
16
17 629 Journal of Vegetation Science 3: 157–164.
18
19 630 KOCHER, A., J.-C. GANTIER, P. GABORIT, L. ZINGER, H. HOLOTA, S. VALIERE, I. DUSFOUR, R.
20
21 631 GIROD, A. L. BAÑULS, and J. MURIENNE. 2016. Vector soup: high-throughput identification of
22
23 632 Neotropical phlebotomine sand flies using metabarcoding. Molecular Ecology Resources 17: 172–
24
25 633 182.
26
27
28 634 KRAFT, N. J. B., L. S. COMITA, J. M. CHASE, N. J. SANDERS, N. G. SWENSON, T. O. CRIST, J. C.
29
30 635 STEGEN, M. VELLEND, B. BOYLE, M. J. ANDERSON, H. V. CORNELL, K. F. DAVIES, A. L.
31
32 636 FREESTONE, B. D. INOUE, S. P. HARRISON, and MYERS. 2011. Disentangling the drivers of β
33
34 637 diversity along latitudinal and elevational gradients. Science 1755–1758.
35
36
37 638 LÊ, S., J. JOSSE, and F. HUSSON. 2008. FactoMineR: an R package for multivariate analysis. Journal
38
39 639 of Statistical Software 25: 1–18.
40
41
42 640 LEGENDRE, P., R. GALZIN, and M. L. HARMELIN-VIVIEN. 1997. Relating behavior to habitat:
43
44 641 solutions to the fourth-corner problem. Ecology 78: 547–562.
45
46
47 642 LICHSTEIN, J. W. 2007. Multiple regression on distance matrices: a multivariate spatial analysis
48
49 643 tool. Plant Ecology 188: 117–131.
50
51
52 644 LONGINO, J. T., M. G. BRANSTETTER, and R. K. COLWELL. 2014. How ants drop out: ant abundance
53
54 645 on tropical mountains. PloS one 9: e104030.
55
56 646 LOS, S. O., F. A. STREET-PERROTT, N. J. LOADER, C. A. FROYD, A. CUNÍ-SANCHEZ, and R. A.

- MARCHANT. 2019. Sensitivity of a tropical montane cloud forest to climate change, present, past and future: Mt. Marsabit, N. Kenya. *Quaternary Science Reviews* 218: 34–48.
- MACHAC, A., M. JANDA, R. R. DUNN, and N. J. SANDERS. 2011. Elevational gradients in phylogenetic structure of ant communities reveal the interplay of biotic and abiotic constraints on diversity. *Ecography* 34: 364–371.
- MCCAIN, C. M., and J.-A. GRYTNES. 2010. Elevational gradients in species richness. *In* *Encyclopedia of Life Sciences (ELS)*. pp. 1–10, John Wiley & Sons, Ltd, Chichester, UK.
- MCGLYNN, T. P., and D. J. SALINAS. 2007. Phosphorus limits tropical rain forest litter fauna. *Biotropica* 39: 50–53.
- MENKE, S. B., and D. A. HOLWAY. 2006. Abiotic factors control invasion by Argentine ants at the community scale. *Journal of Animal Ecology* 75: 368–376.
- MORS DORF, F., B. KÖTZ, E. MEIER, K. I. ITTEN, and B. ALLGÖWER. 2006. Estimation of LAI and fractional cover from small footprint airborne laser scanning data based on gap fraction. *Remote Sensing of Environment* 104: 50–61.
- NOWROUZI, S., A. A. ANDERSEN, S. MACFADYEN, K. M. STAUNTON, J. VANDERWAL, and S. K. A. ROBSON. 2016. Ant diversity and distribution along elevational gradients in the Australian wet tropics: the importance of seasonal moisture stability. *PloS ONE* 11: e0153420.
- OLSEN S., C. COLE, F. WATANABE, and L. DEAN. 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. USDA Circular Nr 939, US Gov. Print. Office, Washington, D.C.
- PURSCHKE, O., B. C. SCHMID, M. T. SYKES, P. POSCHLOD, S. G. MICHALSKI, W. DURKA, I. KÜHN, M. WINTER, and H. C. PRENTICE. 2013. Contrasting changes in taxonomic, phylogenetic and functional diversity during a long-term succession: insights into assembly processes. *Journal of Ecology* 101: 857–866.

1
2
3 671 R CORE TEAM. 2018. R: A Language and Environment for Statistical Computing.
4
5 672 RÉJOU-MÉCHAIN, M., B. TYMEN, L. BLANC, S. FAUSET, T. R. FELDPAUSCH, A. MONTEAGUDO, O.
6
7 673 L. PHILLIPS, H. RICHARD, and J. CHAVE. 2015. Using repeated small-footprint LiDAR acquisitions
8
9 674 to infer spatial and temporal variations of a high-biomass Neotropical forest. *Remote Sensing of*
10
11 675 *Environment* 169: 93–101.
12
13
14 676 SANDERS, N. J., J.-P. LESSARD, M. C. FITZPATRICK, and R. R. DUNN. 2007. Temperature, but not
15
16 677 productivity or geometry, predicts elevational diversity gradients in ants across spatial grains.
17
18 678 *Global Ecology and Biogeography* 16: 640–649.
19
20
21 679 SCHMIDT, F. A., J. H. SCHOEREDER, and M. D. N. CAETANO. 2016. Ant assemblage and
22
23 680 morphological traits differ in response to soil compaction. *Insectes Sociaux* 64: 219–225.
24
25
26 681 SILVA, R. R., and C. R. F. BRANDÃO. 2014. Ecosystem-wide morphological structure of leaf-litter
27
28 682 ant communities along a tropical latitudinal gradient. *PLoS ONE* 9: e93049.
29
30
31 683 SMITH, M. A., W. HALLWACHS, and D. H. JANZEN. 2014. Diversity and phylogenetic community
32
33 684 structure of ants along a Costa Rican elevational gradient. *Ecography* 37: 720–731.
34
35
36 685 STUBLE, K. L., S. L. PELINI, S. E. DIAMOND, D. A. FOWLER, R. R. DUNN, and N. J. SANDERS. 2013.
37
38 686 Foraging by forest ants under experimental climatic warming: a test at two sites. *Ecology and*
39
40 687 *Evolution* 3: 482–491.
41
42
43 688 SWENSON, N. G., P. ANGLADA-CORDERO, and J. A. BARONE. 2011. Deterministic tropical tree
44
45 689 community turnover: evidence from patterns of functional beta diversity along an elevational
46
47 690 gradient. *Proc. R. Soc. B.* 278: 1707.
48
49 691 TYMEN, B., G. VINCENT, E. A. COURTOIS, J. HEURTEBIZE, J. DAUZAT, I. MARECHAUX, and J. CHAVE.
50
51 692 2017. Quantifying micro-environmental variation in tropical rainforest understory at landscape
52
53 693 scale by combining airborne LiDAR scanning and a sensor network. *Annals of Forest Science* 74:
54
55 694 32.
56
57
58
59
60

- 695 VASCONCELOS, H. L., A. C. C. MACEDO, and J. M. S. VILHENA. 2003. Influence of topography on
696 the distribution of ground-dwelling ants in an Amazonian forest. *Studies on Neotropical Fauna and*
697 *Environment* 38: 115–124.
- 698 VLEMINCKX, J., J.-L. DOUCET, J. MORIN-RIVAT, A. B. BIWOLÉ, D. BAUMAN, O. J. HARDY, A.
699 FAYOLLE, J.-F. GILLET, K. DAÏNOU, A. GOREL and T. DROUET. 2017. The influence of spatially
700 structured soil properties on tree community assemblages at a landscape scale in the tropical forests
701 of southern Cameroon. *Journal of Ecology* 105: 354–366.
- 702 VLEMINCKX, J., H. SCHIMANN, T. DECAËNS, M. FICHAUX, V. VEDEL, G. JAOUEN, M. ROY, E.
703 LAPIED, J. ENGEL, A. DOURDAIN, P. PETRONELLI, J. ORIVEL and C. BARALOTO. 2019. Coordinated
704 community structure among trees, fungi and invertebrate groups in Amazonian rainforests.
705 *Scientific reports* 9: 11337.
- 706 WEBER, N. A. 1938. The biology of the fungus-growing ants. Part 4. Additional new forms. Part 5.
707 The Attini of Bolivia. *Revista de Entomologia* 7: 154–206.
- 708 WEISER, M. D., and M. KASPARI. 2006. Ecological morphospace of New World ants. *Ecological*
709 *Entomology* 31: 131–142.
- 710 WILLIG, M. R., D. M. KAUFMAN, and R. D. STEVENS. 2003. Latitudinal gradients of biodiversity:
711 pattern, process, scale, and synthesis. *Annual Review of Ecology, Evolution, and Systematics* 34:
712 273–309.
- 713 WRIGHT, S. K., J. B. YAVITT, N. WURZBURGER, B. L. TURNER, E. V. J. TANNER, E. J. SAYER, L. S.
714 SANTIAGO, M. KASPARI, L. O. HEDIN, K. E. HARMS, M. N. GARCIA, and M. D. CORRE. 2011.
715 Potassium, phosphorus, or nitrogen limit root allocation, tree growth, or litter production in a
716 lowland tropical forest. *Ecology* 92: 1616–1625.
- 717 YATES, M., H. GIBB, and N. R. ANDREW. 2011. Habitat characteristics may override climatic
718 influences on ant assemblage composition: A study using a 300-km climatic gradient climatic

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

719 gradient. Australian Journal of Entomology 59: 332–338.

720 YATES, M. L., N. R. ANDREW, M. BINNS, and H. GIBB. 2014. Morphological traits: predictable

721 responses to macrohabitats across a 300 km scale. PeerJ 2: e271.

722

For Peer Review Only

TABLE 1. List of morphological traits measured and their hypothesized ecological functions. WL = Weber's length; HW = head width; HL = head length; PW = pronotum width; FL = femur length; SL = scape length; EL = eye length; CL = clypeus length; ML = mandible length.

Traits	Hypothesized ecological functions
WL	Proxy for body size, related to several physiological and ecological traits, such as habitat use (Weber 1938, Kaspari & Weiser 1999).
HW	Indicator of mandibular musculature, related to trophic position (Kaspari 1993, Weiser & Kaspari 2006).
HL	Indicator of body size (Kaspari & Weiser 1999) and body mass which determines the quantity of resources consumed.
PW	Predictor of body mass (Kaspari & Weiser 1999).
FL	Indicator of foraging speed, related to habitat complexity (Feener Jr. <i>et al.</i> 1988).
SL	Related to sensory abilities (Weiser & Kaspari 2006), indicative of an ant's ability to navigate and move through its surroundings (Yates <i>et al.</i> 2014).
EL	Indicator of feeding behavior (predatory ants have smaller eyes); hypogaeic ant species have smaller eyes compared to epigaeic ant species (Weiser & Kaspari 2006, Yates <i>et al.</i> 2014).
CL	Related to liquid absorption abilities (Davidson <i>et al.</i> 2004); clypeus length is here used as a surrogate for liquid feeding habits in ants.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ML Longer mandibles are associated with larger prey consumption (Fowler *et al.* 1991). In some cases (i.e. trap-jaw ants), sublinear to linear and long mandibles are used as extremely rapid trap for hunting and killing fast preys (Gronenberg *et al.* 1993).

For Peer Review Only

TABLE 2. Taxonomic, functional and phylogenetic turnover of ant assemblages, averaged over all pairs of plots (Global turnover) or for each pair of elevations (in m: 800-600, 800-400, 600-400). I_{ST} = taxonomic turnover (species level); τ_{ST} = functional turnover (species level); Π_{ST} = phylogenetic turnover (genus level). The functional turnover was calculated using the three first axes of PCA_{TRAITS} (τ_{ST}) and each of these axes separately ($\tau_{ST}PC1$; $\tau_{ST}PC2$; $\tau_{ST}PC3$). Turnover values were multiplied by 10^4 to facilitate the reading. Significant values are emphasized in bold and with asterisks (*** $p \leq 0.001$; ** $p \leq 0.01$; * $p \leq 0.05$).

	Global turnover	800-600	800-400	600-400
I_{ST}	40.98***	36.31*	31.83	17.26
τ_{ST}	19.31**	14.51**	30.88*	-5.46
$\tau_{ST}PC1$	10.18	-10.44	-9.28	-7.79
$\tau_{ST}PC2$	36.24**	39.62**	92.41*	0.76
$\tau_{ST}PC3$	3.20	-9.20	-2.01	-9.44
Π_{ST}	1.05	9.10	7.52	-4.68

TABLE 3. Regression coefficients quantifying the effect on the turnover in community composition (I_{ST} = taxonomic turnover; τ_{ST} = functional turnover; Π_{ST} = phylogenetic turnover), of the overall environmental dissimilarity (Model 1), the dissimilarity of individual environmental variables (Model 2) and the dissimilarity in the scores of the two first axes of the PCA_{ENV} (PC1, PC2; Model 3). For the functional turnover, the effect of environmental variables was evaluated for the three first axes of PCA_{TRAITS} (τ_{ST}) together and for each of these axes separately ($\tau_{ST}PC1$; $\tau_{ST}PC2$; $\tau_{ST}PC3$). Sand = percentage of sand; pH = soil pH; OM = percentage of organic matter; P = Olsen-phosphorus concentration; RH_{MEAN} = mean annual relative humidity; C_H = canopy height). Regression coefficients are multiplied by 10^4 to facilitate the reading as they sometimes reach very low yet significant values. Significant coefficients are emphasized in bold ($***p \leq 0.01$; $**p \leq 0.01$; $*p \leq 0.05$; $\dagger p \leq 0.1$). The regression coefficients were obtained by including the spatial distance in each of the three models in order to remove spatial effects (see methods).

	I_{ST}	τ_{ST}	$\tau_{ST}PC1$	$\tau_{ST}PC2$	$\tau_{ST}PC3$	Π_{ST}
Model 1						
Global env. effect	0.08*	0.20***	0.17*	0.47***	-0.06	-0.02
Model 2						
sand	1.55	10.17†	-0.54	17.08	14.55*	-2.83
pH	-6.95	0.22	14.59†	-14.40	-1.03	10.41†
OM	0.91	-2.08	-0.32	4.93	-7.64	-5.90

P	7.90†	9.42	35.61**	3.22	-13.59†	2.14
RH _{MEAN}	-7.62	7.75	21.61	11.26	-11.51	13.15
C _H	21.99*	15.34	-8.14	43.75	3.40	-13.49
<i>Model 3</i>						
PC1	34.18†	96.57**	67.75†	233.89***	-24.70	-8.90
PC2	9.82	-7.09	28.34	-53.53	-2.23	-0.37

750

751

TABLE 4. Results from the fourth-corner analysis quantifying the strength of pairwise trait-environment associations. Significant relationships are highlighted in bold (** $p \leq 0.01$; * $p \leq 0.05$; † $p \leq 0.07$). HL = relative head length; HW = relative head width; ML = relative mandible length; CL = relative clypeus length; SL = relative scape length; EL = relative eye length; WL = Weber's length; FL = relative femur length; PW = relative pronotum width; PC1/PC2/PC3: axes 1, 2 and 3 of the PCA_{TRAITS}; sand = percentage of sand; pH = soil pH; OM = percentage of organic matter; P = Olsen-phosphorus concentration; RH_{MEAN} = mean annual relative humidity; C_H = canopy height.

	sand	pH	OM	P	RH _{MEAN}	C _H
HL	0.03	0.05	-0.02	0.07*	0.04	-0.00
HW	0.01	0.04	-0.02	0.05	0.01	-0.02
ML	0.03	0.04	-0.01	0.02	0.01	-0.01
CL	-0.03	0.02	-0.03	0.01	-0.02	0.03
SL	-0.07*	0.02	-0.05	-0.05	-0.07*	0.07
EL	-0.10**	0.00	-0.06	-0.08	-0.10*	0.09
WL	-0.03	-0.04	0.01	-0.08**	-0.05	0.00
FL	-0.08†	0.01	-0.05	-0.10	-0.10**	0.09
PW	0.01	0.04	-0.02	0.05	0.01	-0.01
PC1	-0.04	-0.04	0.01	-0.08*	-0.05	0.01

PC2	-0.08*	0.03	-0.06	-0.06	-0.08	0.09
PC3	-0.05*	-0.02	-0.02	-0.02	-0.02	0.02

760

For Peer Review Only

1
2
3 761 FIGURE 1. Location of the study area and the nine sampling plots and environmental sensors at
4
5 762 Mont Itoupé. The weather sensor of Plot 6 was located halfway between Plot 5 and Plot 6 (for more
6
7 763 details, see section 2.5.2).
8
9
10 764
11
12 765 FIGURE 2. Principal Component Analysis of trait data in each plot. HL = relative head length;
13
14 766 HW = relative head width; ML = relative mandible length; CL = relative clypeus length; SL =
15
16 767 relative scape length; EL = relative eye length; WL = Weber's length; FL = relative femur length;
17
18 768 PW = relative pronotum width.
19
20
21 769
22
23 770 FIGURE 3. Principal Component Analysis of environmental data in each plot. Plots of the same
24
25 771 elevational band are represented by green circles (800m), red squares (600m) and orange triangles
26
27 772 (400m). elev = elevation (m); sand = percentage of sand; silt = percentage of silt; clay = percentage
28
29 773 of clay; pH = soil pH; OM = percentage of organic matter; C/N = carbon-to-nitrogen ratio; P =
30
31 774 Olsen-phosphorus concentration; Na = sodium concentration; K = potassium concentration; T_min
32
33 775 = minimum temperature of the coldest month; T_mean = mean annual temperature; RH_min =
34
35 776 minimum relative humidity of the driest month; RH_mean = mean annual relative humidity; C_cov
36
37 777 = canopy cover; C_h = canopy height; gap = gap fraction. Variables retained for analyses are
38
39 778 highlighted in blue.
40
41
42
43
44 779
45
46
47 780
48
49
50
51
52
53
54
55
56
57
58
59
60

781

For Peer Review Only

SUPPORTING INFORMATION

Environmental determinants of leaf-litter ant community composition along an elevational gradient

Mélanie Fichaux^{1*}, Jason Vleminckx^{1,2*}, Elodie A. Courtois^{3,4}, Jacques H. C Delabie^{5,6}, Jordan Galli^{1,7}, Shengli Tao⁸, Nicolas Labrière⁸, Jérôme Chave⁸, Christopher Baraloto², Jérôme Orivel¹

¹ CNRS, UMR Ecologie des Forêts de Guyane (EcoFoG), AgroParisTech, CIRAD, INRA, Université de Guyane, Université des Antilles, Campus agronomique, BP 316, 97379, Kourou cedex, France

² Department of Biological Sciences, Florida International University 11200 S.W. 8th Street Miami, FL 33199, USA

³ Laboratoire Ecologie, Evolution, Interactions des Systèmes Amazoniens (LEEISA), Université de Guyane, CNRS, IFREMER, Cayenne, France

⁴ Department of Biology, Centre of Excellence PLECO (Plant and Vegetation Ecology), University of Antwerp, Wilrijk, Belgium

⁵ Laboratório de Mirmecologia, CEPEC, CEPLAC, Caixa Postal 7, Itabuna, BA 45600-970, Brazil

⁶ Departamento de Ciências Agrárias e Ambientais, Universidade Estadual de Santa Cruz, Rodovia Jorge Amado Km 16, Ilheus, BA 45662-900, Brazil

⁷ Naturalia Environnement, Site Agroparc, 20 rue Lawrence Durell, BP 31285, 84911 Avignon Cedex 9, France

⁸ Laboratoire Evolution et Diversité Biologique UMR 5174, CNRS, Université Paul Sabatier, IRD, 118 route de Narbonne, 31062 Toulouse, France

* Co-first authors

Corresponding author: Mélanie Fichaux, fichaux.mel@gmail.com

FIGURE S1. Phylogenetic tree of ant genera collected along Mont Itoupé.

FIGURE S2. Correlation plots showing correlations between the nine morphological traits. ML = relative mandible length; CL = relative clypeus length; HL = relative head length; HW = relative head width; PW = relative pronotum width; WL = Weber’s length; EL = relative eye length; SL = relative scape length; FL = relative femur length.

FIGURE S3. Correlation plots showing correlations between the nine soil variables. Clay = percentage of clay; sand = percentage of sand; silt = percentage of silt; P = Olsen-phosphorus concentration; K = potassium concentration; Na = sodium concentration; C/N = carbon-to-nitrogen ratio; OM = percentage of organic matter; pH = soil pH.

FIGURE S4. Correlation plots showing correlations between the four climatic variables.
 T_{MIN} = minimum temperature of the coldest month; T_{MEAN} = mean annual temperature; RH_{MIN} = minimum relative humidity of the driest month; RH_{MEAN} = mean annual relative humidity.

FIGURE S5. Correlation plots showing correlations between the three LiDAR-derived variables. C_{COV} = Canopy cover; C_H = Canopy height; Gap = Gap fraction.

FIGURE S6. Third and fourth axis of a principal component analysis of trait data. HL = relative head length; HW = relative head width; ML = relative mandible length; CL = relative clypeus length; SL = relative scape length; EL = relative eye length; WL = Weber’s length; FL = relative femur length; PW = relative pronotum width.

TABLE S1. Mean value of traits per species (HL = head length; HW = head width; ML = mandible length; CL = clypeus length; SL = scape length; EL = eye length; WL = Weber's length; FL = femur length; PW = pronotum width).

Species	HL	HW	ML	CL	SL	EL	WL	FL	PW
<i>Acanthognathus brevicornis</i>	0.83	0.64	0.94	0.26	0.63	0.15	0.79	0.69	0.41
<i>Acanthognathus ocellatus</i>	0.77	0.59	0.76	0.24	0.74	0.13	0.76	0.69	0.39
<i>Acanthognathus</i> sp.IT14.01	0.80	0.65	0.67	0.24	0.68	0.15	0.81	0.75	0.42
<i>Acromyrmex hystrix</i>	1.91	2.30	1.46	0.57	2.64	0.38	3.20	4.26	1.40
<i>Acromyrmex subterraneus brunneus</i>	1.44	1.64	1.11	0.50	1.86	0.27	2.30	2.71	1.02
<i>Acropyga decedens</i>	0.56	0.53	0.31	0.11	0.45	0.03	0.59	0.46	0.35
<i>Acropyga exsanguis</i>	0.48	0.51	0.29	0.10	0.38	0.03	0.51	0.42	0.33
<i>Acropyga fuhrmanni</i>	0.48	0.50	0.28	0.10	0.36	0.04	0.47	0.38	0.31
<i>Acropyga guianensis</i>	0.55	0.50	0.30	0.12	0.45	0.03	0.54	0.45	0.34
<i>Acropyga smithii</i>	0.41	0.40	0.22	0.07	0.26	0.03	0.41	0.25	0.26
<i>Anochetus horridus</i>	1.18	1.02	1.16	0.24	1.41	0.22	1.89	1.49	0.62
<i>Anochetus inermis</i>	1.00	0.95	0.73	0.26	1.03	0.16	1.52	1.12	0.58
<i>Anochetus mayri</i>	0.75	0.76	0.49	0.20	0.66	0.09	1.05	0.72	0.46
<i>Anochetus targionii</i>	0.98	1.02	0.71	0.15	0.95	0.16	1.32	0.96	0.57
<i>Apterostigma acre</i>	0.81	0.65	0.46	0.16	0.71	0.13	1.13	0.91	0.53
<i>Apterostigma auriculatum</i>	1.19	0.85	0.70	0.31	1.16	0.25	1.96	2.05	0.74
<i>Apterostigma avium</i>	1.06	0.72	0.57	0.28	1.40	0.21	1.91	2.14	0.64
<i>Apterostigma chocoense</i>	1.00	0.71	0.53	0.27	1.18	0.19	1.63	1.54	0.59
<i>Apterostigma ierense</i>	0.97	0.69	0.51	0.27	0.92	0.18	1.28	1.32	0.73
<i>Apterostigma pilosum</i>	0.91	0.66	0.48	0.20	0.82	0.15	1.29	1.18	0.55
<i>Apterostigma robustum</i>	1.18	0.86	0.63	0.35	1.12	0.19	1.72	1.62	0.80
<i>Azteca</i> sp.IT14.01	0.65	0.68	0.33	0.19	0.57	0.17	0.77	0.79	0.44
<i>Basiceros manni</i>	1.50	1.18	0.85	0.47	1.34	0.23	2.38	1.98	1.11
<i>Basiceros militaris</i>	1.48	1.39	0.52	0.44	0.81	0.11	1.58	1.13	0.83
<i>Brachymyrmex cavernicola</i>	0.56	0.48	0.28	0.19	0.59	0.09	0.62	0.58	0.34
<i>Brachymyrmex pictus</i>	0.43	0.39	0.21	0.13	0.44	0.10	0.47	0.52	0.27
<i>Brachymyrmex</i> sp.JTL07	0.34	0.29	0.15	0.08	0.32	0.08	0.34	0.33	0.20
<i>Camponotus atriceps</i>	1.97	1.62	0.86	0.62	2.38	0.53	2.86	2.68	1.30
<i>Camponotus excisus</i>	0.98	0.96	0.42	0.30	1.05	0.33	1.25	1.09	0.82
<i>Camponotus fastigatus</i>	1.39	1.17	0.60	0.43	1.78	0.36	2.01	1.84	1.03
<i>Camponotus latangulus</i>	1.08	1.13	0.44	0.32	1.15	0.31	1.46	1.31	0.92
<i>Camponotus linnaei</i>	1.34	1.17	0.54	0.46	1.73	0.36	2.04	1.74	1.22
<i>Camponotus novogranadensis</i>	1.22	1.08	0.58	0.39	1.37	0.32	1.54	1.43	0.84
<i>Camponotus rapax</i>	2.36	1.92	1.09	0.78	3.04	0.46	3.61	3.72	1.55
<i>Camponotus salvini</i>	1.31	1.12	0.64	0.42	1.94	0.34	1.90	2.14	1.04
<i>Camponotus simillimus</i>	2.11	1.56	0.89	0.61	3.07	0.57	3.36	3.66	1.29
<i>Camponotus</i> gp. <i>Tanaemyrmex</i> sp.3	1.52	0.99	0.57	0.50	2.00	0.46	2.50	2.17	0.98
<i>Carebara brevipilosa</i>	0.46	0.42	0.26	0.05	0.29	0.02	0.48	0.37	0.26

<i>Carebara reina</i>	0.34	0.29	0.14	0.06	0.17	0.02	0.31	0.18	0.20
<i>Carebara</i> sp.KW14.01	0.29	0.25	0.14	0.03	0.16	0.01	0.29	0.18	0.15
<i>Carebara urichi</i>	0.42	0.41	0.26	0.09	0.29	0.02	0.49	0.33	0.25
<i>Crematogaster brasiliensis</i>	0.63	0.66	0.33	0.20	0.66	0.15	0.79	0.79	0.39
<i>Crematogaster carinata</i>	0.56	0.60	0.29	0.17	0.57	0.14	0.68	0.71	0.37
<i>Crematogaster curvispinosa</i>	0.58	0.56	0.29	0.20	0.51	0.14	0.63	0.57	0.37
<i>Crematogaster flavosensitiva</i>	0.49	0.47	0.26	0.17	0.48	0.14	0.52	0.49	0.30
<i>Crematogaster levior</i>	0.55	0.54	0.28	0.18	0.56	0.13	0.66	0.64	0.34
<i>Crematogaster limata</i>	0.70	0.72	0.36	0.22	0.73	0.17	0.83	0.85	0.43
<i>Crematogaster sotobosque</i>	0.49	0.47	0.27	0.17	0.52	0.14	0.58	0.59	0.33
<i>Crematogaster tenuicula</i>	0.55	0.52	0.29	0.17	0.60	0.13	0.68	0.73	0.35
<i>Crematogaster torosa</i>	0.74	0.81	0.36	0.22	0.53	0.18	0.85	0.70	0.47
<i>Cryptopone holmgreni</i>	0.94	0.95	0.64	0.19	0.58	0.00	1.34	0.70	0.64
<i>Cyphomyrmex peltatus</i>	0.51	0.56	0.30	0.18	0.51	0.12	0.74	0.57	0.37
<i>Cyphomyrmex rimosus</i>	0.56	0.57	0.33	0.20	0.57	0.13	0.80	0.69	0.37
<i>Discothyrea denticulata</i>	0.43	0.37	0.18	0.10	0.22	0.01	0.43	0.25	0.26
<i>Discothyrea sexarticulata</i>	0.39	0.35	0.17	0.05	0.20	0.02	0.38	0.23	0.24
<i>Dolichoderus bidens</i>	1.62	1.42	0.75	0.45	1.81	0.38	2.40	2.62	1.06
<i>Dolichoderus bispinosus</i>	1.31	1.47	0.57	0.37	1.35	0.30	1.72	1.75	0.93
<i>Dolichoderus gagates</i>	0.85	0.94	0.43	0.27	0.85	0.21	1.12	1.08	0.62
<i>Dolichoderus imitator</i>	1.37	0.91	0.74	0.48	2.05	0.40	2.59	2.32	0.88
<i>Dolichoderus lutosus</i>	1.16	1.08	0.48	0.38	1.04	0.28	1.54	1.20	0.62
<i>Dolichoderus smithi</i>	1.07	1.03	0.46	0.39	0.87	0.29	1.24	1.20	0.86
<i>Ectatomma edentatum</i>	1.82	1.48	1.21	0.73	1.78	0.35	2.94	2.51	1.20
<i>Ectatomma lugens</i>	2.21	1.77	1.52	0.72	2.27	0.58	3.93	3.40	1.49
<i>Ectatomma tuberculatum</i>	2.40	1.94	1.75	0.97	2.43	0.57	3.75	3.13	1.59
<i>Eurhopalothrix bolau</i>	0.61	0.66	0.24	0.21	0.36	0.08	0.72	0.55	0.41
<i>Fulakora orizabana</i>	0.57	0.52	0.46	0.09	0.35	0.00	0.72	0.36	0.33
<i>Gigantiops destructor</i>	2.18	2.16	1.21	0.90	2.52	1.56	3.25	4.46	1.12
<i>Gnamptogenys acuminata</i>	1.02	0.94	0.66	0.26	0.85	0.29	1.61	1.21	0.83
<i>Gnamptogenys annulata</i>	1.26	1.14	0.86	0.34	0.88	0.27	2.00	1.29	1.03
<i>Gnamptogenys haenschii</i>	1.80	1.76	1.30	0.57	1.45	0.16	2.68	1.98	1.46
<i>Gnamptogenys horni</i>	0.73	0.61	0.42	0.19	0.48	0.14	0.98	0.62	0.54
<i>Gnamptogenys interrupta</i>	0.97	0.83	0.63	0.25	0.60	0.17	1.33	0.79	0.64
<i>Gnamptogenys mecotyle</i>	1.19	1.12	0.88	0.32	1.17	0.29	1.93	1.36	1.05
<i>Gnamptogenys moelleri</i>	1.19	1.00	0.65	0.36	1.29	0.21	1.72	1.63	0.75
<i>Gnamptogenys regularis</i>	0.74	0.69	0.45	0.20	0.52	0.19	1.04	0.70	0.59
<i>Gnamptogenys relict</i>	0.67	0.55	0.35	0.18	0.49	0.08	0.85	0.57	0.44
<i>Gnamptogenys sulcata</i>	1.14	1.12	0.76	0.31	1.01	0.39	1.90	1.56	0.86
<i>Gnamptogenys tortuolosa</i>	1.68	1.51	1.08	0.46	1.52	0.43	2.58	2.02	1.25
<i>Heteroponera flava</i>	0.86	0.74	0.44	0.28	0.52	0.19	1.15	0.68	0.58
<i>Hylomyrma blandiens</i>	0.83	0.80	0.54	0.25	0.61	0.22	1.12	0.78	0.58
<i>Hylomyrma immanis</i>	1.18	1.15	0.79	0.41	0.90	0.28	1.61	1.25	0.83

1										
2										
3										
4	<i>Hylomyrma reginae</i>	1.04	1.02	0.67	0.29	0.76	0.23	1.43	1.07	0.74
5	<i>Hypoponera foreli</i>	0.58	0.52	0.32	0.10	0.43	0.03	0.77	0.47	0.37
6	<i>Hypoponera</i> sp.02.FGSGt	0.73	0.65	0.39	0.19	0.65	0.04	1.03	0.70	0.50
7	<i>Hypoponera</i> sp.12.FGSGt	0.46	0.36	0.24	0.08	0.30	0.02	0.56	0.31	0.26
8	<i>Hypoponera</i> sp.13.FGSGt	0.51	0.44	0.27	0.09	0.33	0.04	0.64	0.40	0.32
9	<i>Hypoponera</i> sp.B	0.89	0.78	0.51	0.22	0.77	0.04	1.30	0.84	0.61
10	<i>Hypoponera</i> sp.IT14.01	0.64	0.54	0.35	0.18	0.58	0.05	0.88	0.65	0.40
11	<i>Hypoponera</i> sp.IT14.02	0.74	0.69	0.41	0.18	0.70	0.04	1.06	0.72	0.50
12	<i>Hypoponera</i> sp.IT14.04	0.46	0.39	0.21	0.08	0.33	0.04	0.58	0.34	0.30
13	<i>Hypoponera</i> sp.IT14.05	0.59	0.51	0.32	0.14	0.47	0.03	0.82	0.47	0.40
14	<i>Hypoponera</i> sp.MI15.02	0.62	0.49	0.32	0.11	0.45	0.03	0.83	0.59	0.38
15	<i>Hypoponera</i> sp.MI15.05	0.82	0.75	0.46	0.12	0.70	0.05	1.17	0.85	0.55
16	<i>Labidus coecus</i>	1.04	0.96	0.71	0.18	0.77	0.04	1.42	1.41	0.63
17	<i>Leptogenys gaigei</i>	1.44	0.97	0.65	0.52	1.58	0.33	2.39	1.85	0.75
18	<i>Leptogenys langi</i>	0.88	0.60	0.46	0.27	0.79	0.16	1.30	0.88	0.49
19	<i>Leptogenys</i> sp.05.FGSGt	0.96	0.57	0.41	0.36	0.90	0.22	1.47	1.03	0.52
20	<i>Leptogenys</i> sp.IT14.04	1.14	0.94	0.74	0.36	1.17	0.26	1.83	1.22	0.73
21	<i>Mayaponera constricta</i>	1.48	1.23	1.03	0.42	1.56	0.26	2.47	1.88	0.95
22	<i>Megalomyrmex balzani</i>	1.97	1.61	0.99	0.52	2.66	0.43	2.81	3.31	1.18
23	<i>Megalomyrmex drifti</i>	0.54	0.44	0.28	0.13	0.44	0.14	0.69	0.48	0.34
24	<i>Megalomyrmex incisus</i>	0.64	0.54	0.36	0.17	0.59	0.20	0.89	0.78	0.43
25	<i>Megalomyrmex silvestrii</i>	0.56	0.46	0.30	0.16	0.73	0.18	0.77	0.71	0.36
26	<i>Megalomyrmex</i> sp.MI15.01	0.78	0.62	0.42	0.25	0.79	0.22	1.18	0.98	0.52
27	<i>Mycetophylax faunulus</i>	0.74	0.78	0.46	0.19	0.64	0.13	1.01	1.08	0.49
28	<i>Mycetophylax morschi</i>	0.50	0.50	0.32	0.13	0.40	0.09	0.66	0.53	0.37
29	<i>Mycetophylax strigatus</i>	0.66	0.73	0.41	0.20	0.65	0.11	1.01	0.89	0.45
30	<i>Mycocepurus smithii</i>	0.60	0.62	0.40	0.15	0.52	0.12	0.84	0.63	0.40
31	<i>Myrmelachista guyanensis</i>	0.47	0.47	0.17	0.11	0.26	0.10	0.54	0.32	0.30
32	<i>Myrmicocrypta</i> sp.IT14.01	0.68	0.64	0.48	0.22	0.63	0.07	0.92	0.68	0.44
33	<i>Myrmicocrypta</i> sp.IT14.02	0.62	0.53	0.37	0.12	0.55	0.06	0.77	0.62	0.34
34	<i>Myrmicocrypta</i> sp.IT14.03	0.64	0.53	0.39	0.19	0.64	0.07	0.78	0.66	0.36
35	<i>Neocerapachys splendens</i>	0.77	0.58	0.31	0.15	0.40	0.04	0.99	0.56	0.43
36	<i>Neoponera apicalis</i>	2.74	2.32	2.16	0.77	3.08	0.93	4.47	4.19	1.62
37	<i>Neoponera commutata</i>	3.75	3.24	2.96	1.07	3.81	0.98	5.64	5.02	2.46
38	<i>Neoponera crenata</i>	2.34	1.96	1.29	0.85	2.29	0.57	3.58	2.00	1.51
39	<i>Neoponera foetida</i>	2.55	2.28	1.98	1.04	2.36	0.69	4.22	3.46	0.69
40	<i>Neoponera procidua</i>	2.86	2.80	2.28	0.65	2.41	0.33	4.12	3.02	1.90
41	<i>Neoponera verena</i>	2.27	1.79	1.58	0.88	2.63	0.82	3.75	3.03	1.47
42	<i>Nylanderia fulva</i>	0.69	0.57	0.35	0.16	0.89	0.15	0.88	0.96	0.43
43	<i>Nylanderia guatemalensis</i>	0.64	0.56	0.33	0.20	0.78	0.15	0.81	0.77	0.41
44	<i>Nylanderia</i> sp.IT14.01	0.63	0.52	0.33	0.15	0.78	0.10	0.78	0.80	0.39
45	<i>Nylanderia</i> sp.SL13.03	0.69	0.58	0.37	0.17	0.85	0.16	0.87	0.89	0.42
46	<i>Nylanderia steinheili</i>	0.54	0.49	0.26	0.15	0.69	0.14	0.69	0.65	0.35
47										
48										
49										
50										
51										
52										
53										
54										
55										
56										
57										
58										
59										
60										

<i>Ochetomyrmex neopolitus</i>	0.54	0.51	0.29	0.17	0.42	0.13	0.65	0.49	0.32
<i>Ochetomyrmex semipolitus</i>	0.51	0.47	0.29	0.16	0.35	0.11	0.55	0.39	0.30
<i>Octostruma balzani</i>	0.48	0.54	0.18	0.17	0.27	0.04	0.54	0.35	0.33
<i>Octostruma betschi</i>	0.46	0.53	0.20	0.16	0.28	0.04	0.54	0.37	0.33
<i>Octostruma iheringi</i>	0.60	0.73	0.23	0.23	0.36	0.04	0.74	0.50	0.42
<i>Octostruma obtusidens</i>	0.58	0.70	0.23	0.18	0.35	0.05	0.78	0.53	0.47
<i>Octostruma</i> sp.JTL035	0.52	0.64	0.21	0.18	0.31	0.04	0.69	0.43	0.37
<i>Odontomachus caelatus</i>	3.22	2.72	1.95	0.61	3.09	0.52	3.81	3.55	1.56
<i>Odontomachus chelifer</i>	3.84	2.63	2.32	0.60	3.96	0.66	4.71	4.71	1.51
<i>Odontomachus haematodus</i>	2.42	1.93	1.47	0.41	2.28	0.42	2.87	2.44	1.09
<i>Odontomachus laticeps</i>	2.39	1.86	1.45	0.42	2.11	0.39	3.01	2.61	1.01
<i>Odontomachus meinerti</i>	1.94	1.53	1.16	0.32	1.82	0.30	2.14	1.87	0.91
<i>Odontomachus scalptus</i>	2.66	2.14	1.59	0.45	2.64	0.47	3.24	2.91	1.14
<i>Pachycondyla crassinoda</i>	3.79	3.70	2.64	0.87	3.42	0.74	5.67	4.21	2.68
<i>Pachycondyla harpax</i>	1.70	1.73	1.08	0.38	1.43	0.31	2.54	1.71	1.24
<i>Pachycondyla impressa</i>	2.91	2.80	1.94	0.61	2.46	0.52	4.15	2.95	2.08
<i>Pachycondyla</i> nr. <i>lenkoi</i>	2.13	2.00	1.33	0.50	1.73	0.31	2.79	1.92	1.41
<i>Pheidole aenescens</i>	0.65	0.55	0.38	0.17	0.92	0.16	0.87	1.01	0.38
<i>Pheidole alienata</i>	0.63	0.59	0.40	0.18	0.89	0.14	0.79	1.07	0.38
<i>Pheidole allarmata</i>	0.54	0.57	0.45	0.13	0.61	0.08	0.74	0.61	0.38
<i>Pheidole araneoides</i>	0.89	0.68	0.53	0.25	1.35	0.18	1.36	1.59	0.53
<i>Pheidole biconstricta</i>	0.82	0.73	0.52	0.24	1.00	0.16	1.09	1.18	0.48
<i>Pheidole bruesi</i>	0.56	0.51	0.33	0.18	0.66	0.12	0.70	0.68	0.34
<i>Pheidole bulliceps</i>	0.72	0.62	0.46	0.20	0.94	0.16	1.00	1.08	0.43
<i>Pheidole cataractae</i>	0.51	0.48	0.31	0.16	0.69	0.11	0.67	0.74	0.29
<i>Pheidole coffeicola</i>	0.60	0.49	0.34	0.16	0.86	0.14	0.83	0.96	0.35
<i>Pheidole cramptoni</i>	0.54	0.51	0.32	0.16	0.48	0.11	0.65	0.65	0.34
<i>Pheidole cursor</i>	0.60	0.48	0.36	0.17	0.84	0.14	0.82	0.90	0.35
<i>Pheidole deima</i>	0.72	0.80	0.51	0.22	0.70	0.15	0.98	1.00	0.53
<i>Pheidole embolopyx</i>	0.57	0.49	0.33	0.16	0.71	0.13	0.75	0.77	0.34
<i>Pheidole fissiceps</i>	0.61	0.49	0.35	0.16	0.89	0.15	0.85	0.95	0.35
<i>Pheidole flavens</i>	0.40	0.38	0.24	0.12	0.35	0.08	0.44	0.35	0.25
<i>Pheidole fracticeps</i>	0.41	0.41	0.23	0.13	0.33	0.09	0.49	0.40	0.23
<i>Pheidole gemmula</i>	0.42	0.41	0.24	0.14	0.33	0.08	0.42	0.36	0.25
<i>Pheidole</i> gp. <i>fallax</i> sp.D	0.59	0.56	0.44	0.16	0.80	0.12	0.84	0.94	0.38
<i>Pheidole</i> gp. <i>fallax</i> sp.F	0.64	0.58	0.37	0.18	0.88	0.16	0.88	0.98	0.39
<i>Pheidole</i> gp. <i>tristis</i> sp.B3	0.53	0.52	0.34	0.13	0.49	0.09	0.62	0.60	0.32
<i>Pheidole</i> gp. <i>tristis</i> sp.C	0.45	0.41	0.28	0.13	0.46	0.09	0.56	0.47	0.27
<i>Pheidole</i> gp. <i>tristis</i> sp.D	0.61	0.56	0.43	0.17	0.81	0.11	0.85	0.88	0.38
<i>Pheidole</i> gp. <i>tristis</i> sp.H1	0.52	0.51	0.34	0.14	0.52	0.10	0.68	0.58	0.33
<i>Pheidole guajirana</i>	0.39	0.37	0.22	0.13	0.33	0.08	0.46	0.36	0.23
<i>Pheidole jeannei</i>	0.57	0.56	0.37	0.17	0.64	0.12	0.68	0.73	0.34
<i>Pheidole latti</i>	0.59	0.48	0.34	0.14	0.91	0.14	0.85	1.02	0.34

1									
2									
3									
4	<i>Pheidole longiseta</i>	0.63	0.57	0.37	0.18	0.81	0.15	0.81	0.39
5	<i>Pheidole lovejoyi</i>	0.62	0.49	0.36	0.17	0.99	0.15	0.94	1.10
6	<i>Pheidole meinertopsis</i>	0.44	0.45	0.27	0.13	0.42	0.09	0.53	0.46
7	<i>Pheidole microps</i>	0.52	0.51	0.35	0.13	0.46	0.05	0.64	0.51
8	<i>Pheidole midas</i>	0.61	0.56	0.35	0.19	0.71	0.12	0.73	0.72
9	<i>Pheidole nana</i>	0.52	0.50	0.33	0.13	0.49	0.07	0.61	0.59
10	<i>Pheidole neoschultzi</i>	0.34	0.33	0.18	0.09	0.29	0.08	0.36	0.30
11	<i>Pheidole nitella</i>	0.36	0.34	0.21	0.11	0.31	0.08	0.41	0.31
12	<i>Pheidole</i> nr. <i>araneoides</i>	0.74	0.59	0.45	0.20	1.51	0.17	1.18	1.67
13	<i>Pheidole</i> nr. <i>cramptoni</i>	0.53	0.50	0.32	0.14	0.46	0.11	0.64	0.53
14	<i>Pheidole</i> nr. <i>fimbriata</i> sp.A	0.68	0.67	0.53	0.18	0.74	0.14	0.98	0.85
15	<i>Pheidole</i> nr. <i>fimbriata</i> sp.C	0.58	0.59	0.42	0.16	0.58	0.13	0.73	0.71
16	<i>Pheidole</i> nr. <i>fimbriata</i> sp.D	0.42	0.41	0.26	0.12	0.40	0.08	0.49	0.46
17	<i>Pheidole</i> nr. <i>sagax</i>	0.61	0.59	0.34	0.17	0.71	0.14	0.79	0.78
18	<i>Pheidole</i> nr. <i>seeldrayersii</i>	0.60	0.57	0.35	0.17	0.80	0.15	0.81	0.87
19	<i>Pheidole</i> nr. <i>subarmata</i>	0.38	0.36	0.23	0.11	0.32	0.09	0.44	0.34
20	<i>Pheidole</i> nr. <i>tristis</i>	0.59	0.59	0.44	0.17	0.62	0.12	0.81	0.79
21	<i>Pheidole pedana</i>	0.37	0.37	0.21	0.12	0.27	0.08	0.41	0.30
22	<i>Pheidole rugiceps</i>	0.52	0.46	0.32	0.14	0.70	0.11	0.66	0.78
23	<i>Pheidole sagax</i>	0.56	0.52	0.33	0.17	0.68	0.13	0.73	0.72
24	<i>Pheidole sculptor</i>	0.43	0.40	0.27	0.12	0.41	0.09	0.47	0.45
25	<i>Pheidole</i> sp.IT14.01	0.75	0.75	0.52	0.23	0.75	0.13	1.00	0.98
26	<i>Pheidole</i> sp.IT14.08	0.42	0.43	0.23	0.12	0.34	0.08	0.43	0.34
27	<i>Pheidole</i> sp.MI15.06	0.44	0.41	0.25	0.14	0.41	0.09	0.49	0.49
28	<i>Pheidole subarmata</i>	0.43	0.41	0.28	0.13	0.43	0.09	0.55	0.47
29	<i>Pheidole terribilis</i>	0.74	0.82	0.64	0.19	0.87	0.16	1.15	0.95
30	<i>Pheidole tristis</i>	0.73	0.68	0.47	0.23	0.73	0.13	0.96	0.91
31	<i>Pheidole zelata</i>	0.66	0.63	0.39	0.19	0.86	0.15	0.86	1.04
32	<i>Pheidole zeteki</i>	0.42	0.42	0.20	0.12	0.29	0.08	0.40	0.29
33	<i>Platythyrea pilosula</i>	1.51	1.23	0.76	0.41	1.11	0.42	2.31	1.61
34	<i>Prionopelta amabilis</i>	0.53	0.46	0.26	0.09	0.29	0.03	0.67	0.33
35	<i>Procryptocerus scabriusculus</i>	1.30	1.40	0.49	0.35	0.86	0.33	1.68	1.04
36	<i>Procryptocerus schmitti</i>	0.87	0.88	0.27	0.27	0.57	0.27	1.11	0.61
37	<i>Procryptocerus</i> sp.IT14.01	1.50	1.52	0.30	0.43	0.87	0.40	1.89	1.11
38	<i>Pseudomyrmex tenuis</i>	1.27	1.13	0.52	0.20	0.60	0.83	1.78	1.18
39	<i>Pseudoponera stigma</i>	1.01	0.99	0.65	0.24	0.75	0.09	1.48	0.85
40	<i>Pseudoponera succedanea</i>	0.86	0.81	0.53	0.24	0.60	0.10	1.23	0.69
41	<i>Rasopone arhuaca</i>	1.28	1.13	0.86	0.23	1.03	0.16	1.85	1.27
42	<i>Rasopone</i> sp.JTL047	1.29	1.23	0.82	0.21	0.95	0.14	1.82	1.06
43	<i>Rhopalothrix</i> sp.JTL014	0.34	0.41	0.16	0.12	0.20	0.01	0.46	0.29
44	<i>Rogeria besucheti</i>	0.69	0.64	0.36	0.21	0.47	0.08	0.75	0.49
45	<i>Rogeria blanda</i>	0.81	0.74	0.42	0.25	0.55	0.14	1.05	0.70
46	<i>Rogeria germaini</i>	0.76	0.67	0.35	0.16	0.53	0.08	0.84	0.63
47									
48									
49									
50									
51									
52									
53									
54									
55									
56									
57									
58									
59									
60									

<i>Rogeria innotabilis</i>	0.60	0.52	0.28	0.08	0.40	0.08	0.62	0.52	0.35
<i>Rogeria scobinata</i>	0.55	0.48	0.27	0.16	0.35	0.08	0.59	0.41	0.35
<i>Rogeria</i> sp.IT14.01	0.45	0.38	0.22	0.10	0.30	0.03	0.47	0.33	0.29
<i>Rogeria</i> sp.IT14.06	0.50	0.44	0.25	0.10	0.33	0.03	0.51	0.35	0.31
<i>Rogeria</i> sp.SL13.01	0.54	0.49	0.26	0.09	0.36	0.07	0.56	0.43	0.34
<i>Rogeria subarmata</i>	0.77	0.67	0.34	0.16	0.50	0.15	0.88	0.66	0.50
<i>Sericomyrmex bondari</i>	0.96	1.27	0.69	0.32	0.86	0.18	1.53	1.28	0.80
<i>Sericomyrmex mayri</i>	1.11	1.48	0.79	0.33	0.90	0.20	1.65	1.55	0.98
<i>Sericomyrmex parvulus</i>	0.78	0.97	0.56	0.24	0.62	0.15	1.15	0.93	0.63
<i>Solenopsis bicolor</i>	0.52	0.45	0.23	0.10	0.31	0.05	0.56	0.41	0.30
<i>Solenopsis</i> sp.01	0.49	0.42	0.26	0.09	0.40	0.05	0.57	0.46	0.29
<i>Solenopsis</i> sp.02	0.47	0.40	0.23	0.07	0.34	0.05	0.57	0.44	0.26
<i>Solenopsis</i> sp.03	0.46	0.39	0.24	0.07	0.28	0.04	0.50	0.31	0.26
<i>Solenopsis</i> sp.09	0.59	0.54	0.25	0.11	0.36	0.05	0.67	0.48	0.36
<i>Solenopsis</i> sp.11	0.34	0.30	0.15	0.06	0.18	0.04	0.35	0.18	0.20
<i>Solenopsis</i> sp.12	0.40	0.34	0.21	0.07	0.26	0.04	0.44	0.27	0.23
<i>Solenopsis</i> sp.13	0.39	0.33	0.17	0.06	0.22	0.04	0.42	0.25	0.22
<i>Solenopsis</i> sp.14	0.35	0.30	0.18	0.04	0.24	0.04	0.40	0.27	0.20
<i>Solenopsis</i> sp.15	0.38	0.31	0.19	0.07	0.26	0.04	0.42	0.28	0.21
<i>Solenopsis</i> sp.17	0.42	0.35	0.19	0.08	0.27	0.04	0.45	0.30	0.23
<i>Solenopsis</i> sp.21	0.34	0.28	0.16	0.05	0.22	0.03	0.35	0.23	0.19
<i>Solenopsis</i> sp.23	0.41	0.34	0.21	0.06	0.29	0.03	0.42	0.34	0.22
<i>Solenopsis</i> sp.24	0.34	0.27	0.15	0.06	0.17	0.04	0.34	0.19	0.18
<i>Solenopsis</i> sp.25	0.30	0.24	0.14	0.04	0.16	0.03	0.30	0.18	0.15
<i>Solenopsis</i> sp.28	0.48	0.41	0.22	0.09	0.34	0.05	0.52	0.38	0.26
<i>Solenopsis virulens</i>	0.83	0.78	0.44	0.24	0.74	0.12	1.08	0.99	0.50
<i>Strumigenys auctidens</i>	0.47	0.36	0.19	0.15	0.17	0.05	0.48	0.27	0.25
<i>Strumigenys beebei</i>	0.55	0.39	0.14	0.16	0.26	0.08	0.57	0.33	0.26
<i>Strumigenys borgmeieri</i>	0.69	0.48	0.48	0.21	0.57	0.09	0.77	0.72	0.34
<i>Strumigenys cincinnata</i>	0.45	0.33	0.10	0.15	0.22	0.04	0.45	0.31	0.21
<i>Strumigenys cosmostela</i>	0.48	0.44	0.32	0.11	0.32	0.05	0.57	0.39	0.28
<i>Strumigenys crassicornis</i>	0.48	0.40	0.20	0.15	0.24	0.05	0.49	0.36	0.27
<i>Strumigenys denticulata</i>	0.41	0.35	0.36	0.12	0.27	0.04	0.46	0.30	0.24
<i>Strumigenys diabola</i>	0.73	0.70	1.11	0.23	0.75	0.09	0.89	0.83	0.42
<i>Strumigenys dyseides</i>	0.38	0.32	0.24	0.09	0.25	0.02	0.38	0.29	0.21
<i>Strumigenys elongata</i>	0.54	0.48	0.41	0.13	0.47	0.05	0.61	0.46	0.30
<i>Strumigenys perparva</i>	0.36	0.33	0.22	0.10	0.24	0.04	0.39	0.26	0.21
<i>Strumigenys schulzi</i>	0.46	0.38	0.12	0.17	0.24	0.06	0.50	0.32	0.25
<i>Strumigenys smithii</i>	0.59	0.58	0.43	0.16	0.49	0.08	0.75	0.62	0.37
<i>Strumigenys spathula</i>	0.52	0.49	0.36	0.12	0.42	0.05	0.59	0.44	0.27
<i>Strumigenys subedentata</i>	0.46	0.44	0.32	0.11	0.21	0.06	0.53	0.39	0.27
<i>Strumigenys trinidadensis</i>	0.67	0.65	0.55	0.18	0.59	0.11	0.85	0.73	0.38
<i>Strumigenys trudifera</i>	0.65	0.51	0.77	0.18	0.57	0.04	0.70	0.65	0.32

<i>Strumigenys villiersi</i>	0.48	0.35	0.16	0.15	0.25	0.05	0.57	0.37	0.22
<i>Strumigenys waiwai</i>	0.31	0.27	0.20	0.07	0.22	0.03	0.36	0.22	0.18
<i>Tatuidris tatusia</i>	0.67	0.86	0.37	0.18	0.33	0.05	0.72	0.41	0.55
<i>Thaumatomyrmex soesilae</i>	0.65	0.80	0.79	0.17	0.59	0.19	1.04	0.76	0.53
<i>Trachymyrmex cornetzi</i>	0.73	0.79	0.50	0.21	0.75	0.14	1.11	1.00	0.52
<i>Trachymyrmex farinosus</i>	1.01	1.23	0.77	0.32	0.92	0.21	1.63	1.48	0.81
<i>Trachymyrmex mandibularis</i>	1.02	1.34	0.78	0.30	1.21	0.22	1.57	1.61	0.78
<i>Trachymyrmex nr. farinosus</i>	1.22	1.34	0.87	0.40	1.11	0.25	1.48	1.64	0.88
<i>Trachymyrmex</i> sp.A	1.27	1.50	0.92	0.38	1.14	0.26	2.06	1.90	1.02
<i>Trachymyrmex</i> sp.B	0.94	1.01	0.71	0.27	0.89	0.19	1.51	1.39	0.68
<i>Tranopelta gilva</i>	0.64	0.63	0.38	0.19	0.42	0.06	0.76	0.52	0.42
<i>Typhlomyrmex pusillus</i>	0.43	0.36	0.22	0.11	0.27	0.01	0.57	0.27	0.26
<i>Wasmannia auropunctata</i>	0.45	0.40	0.20	0.14	0.37	0.11	0.46	0.44	0.27
<i>Wasmannia rochai</i>	0.49	0.49	0.24	0.17	0.40	0.11	0.55	0.44	0.33

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

TABLE S2. Leaf-litter ant sampling plots, coordinates, elevation and environmental parameters measured in the study. Lat. = latitude; long. = longitude; elev. = elevation (m); sand = percentage of sand; silt = percentage of silt; clay = percentage of clay; pH = soil pH; OM = percentage of organic matter; C/N = carbon-to-nitrogen ratio; P = Olsen-phosphorus concentration; Na = sodium concentration; K = potassium concentration; T_{MIN} = minimum temperature of the coldest month; T_{MEAN} = mean annual temperature; RH_{MIN} = minimum relative humidity of the driest month; RH_{MEAN} = mean annual relative humidity; C_{COV} = canopy cover; C_H = canopy height; gap = gap fraction. Variables retained for analyses are highlighted in bold.

Plot	lat.	long.	elev.	sand	silt	clay	pH	OM	C/N	P	Na	K	T _{MIN}	T _{MEAN}	RH _{MIN}	RH_{MEAN}	C _{COV}	C_H	gap
IT-01	-53.08	3.03	818.85	22.47	32.77	44.76	4.54	8.16	12.96	2.27	3.66	0.12	17.44	21.08	69.34	99.54	0.98	21.05	0.85
IT-02	-53.08	3.02	822.86	29.42	29.77	40.81	4.60	9.35	13.07	2.53	4.15	0.13	17.65	20.99	64.77	99.18	1.00	22.69	0.91
IT-03	-53.09	3.02	801.74	28.57	9.00	62.44	4.74	6.13	12.87	1.73	2.77	0.09	17.68	21.18	71.13	99.27	0.99	20.36	0.88
IT-04	-53.10	3.03	581.82	13.39	8.36	78.25	5.54	5.59	12.24	1.27	2.66	0.07	19.37	22.35	59.74	97.53	1.00	30.82	0.95
IT-05	-53.10	3.02	578.16	15.78	9.31	74.91	4.73	5.68	12.13	1.20	2.72	0.08	19.72	22.67	54.38	95.41	1.00	32.53	0.96
IT-06	-53.10	3.01	623.70	8.98	11.83	79.92	4.61	5.83	12.61	1.53	2.69	0.09	19.32	22.27	58.23	96.56	1.00	34.20	0.95
IT-07	-53.11	3.03	464.55	6.70	6.44	86.85	4.47	4.66	12.11	1.80	2.24	0.08	20.34	23.21	57.53	96.86	0.99	32.15	0.94
IT-08	-53.11	3.02	434.11	36.61	48.44	14.96	6.22	3.83	9.18	2.00	2.43	0.09	19.77	23.14	58.57	96.78	1.00	39.11	0.95
IT-09	-53.11	3.02	422.30	8.16	8.64	83.21	5.66	5.14	9.77	1.80	3.06	0.10	20.44	23.36	58.08	95.83	1.00	37.09	0.97

TABLE S3. Number of occurrences of species collected at each elevation (between 0 and 60 per elevation), according to their classification. A dash indicates the absence of a species at an elevation.

Species occurrences at each elevation	400m	600m	800m
Subfamily AGROECOMYRMECINAE Carpenter, 1930			
Tribe Agroecomyrmecini Carpenter, 1930			
<i>Tatuidris tatusia</i> Brown & Kempf, 1968	1	2	-
Subfamily AMBLYOPONINAE Forel, 1893			
Tribe Amblyoponini Forel, 1893			
<i>Fulakora orizabana</i> (Brown, 1960)	1	-	-
<i>Prionopelta amabilis</i> Borgmeier, 1949	8	5	26
Subfamily DOLICHODERINAE Forel, 1878			
Tribe Dolichoderini Forel, 1878			
<i>Azteca</i> sp.IT14.01	1	-	-
<i>Dolichoderus bidens</i> (Linnaeus, 1758)	1	-	-
<i>Dolichoderus bispinosus</i> (Olivier, 1792)	1	1	-
<i>Dolichoderus gagates</i> Emery, 1890	-	2	-
<i>Dolichoderus imitator</i> Emery, 1894	1	1	-
<i>Dolichoderus lutosus</i> (Smith, 1858)	1	-	-
<i>Dolichoderus smithi</i> MacKay, 1993	-	1	-
Subfamily DORYLINAE Leach, 1815			
Tribe Cerapachyini Forel, 1893			
<i>Neocerapachys splendens</i> (Borgmeier, 1957)	-	2	1
Tribe Ecitonini Forel, 1893			
<i>Labidus coecus</i> (Latreille, 1802)	5	5	3
Subfamily ECTATOMMINAE Emery, 1895			
Tribe Ectatommini Emery, 1895			
<i>Ectatomma edentatum</i> Roger, 1863	12	5	3
<i>Ectatomma lugens</i> Emery, 1894	21	20	-
<i>Ectatomma tuberculatum</i> (Olivier, 1792)	1	1	-
<i>Gnamptogenys acuminata</i> (Emery, 1896)	-	1	2
<i>Gnamptogenys annulata</i> (Mayr, 1887)	-	-	1
<i>Gnamptogenys haenschi</i> (Emery, 1902)	1	2	3
<i>Gnamptogenys horni</i> (Santschi, 1929)	13	12	11
<i>Gnamptogenys interrupta</i> (Mayr, 1887)	1	-	-
<i>Gnamptogenys mecotyle</i> Brown, 1958	1	-	1
<i>Gnamptogenys moelleri</i> (Forel, 1912)	1	5	9
<i>Gnamptogenys regularis</i> Mayr, 1870	-	1	-
<i>Gnamptogenys relictata</i> (Mann, 1916)	-	-	1
<i>Gnamptogenys sulcata</i> (Smith, 1858)	1	-	-
<i>Gnamptogenys tortuolosa</i> (Smith, 1858)	-	1	-
Tribe Typhlomymecini Emery, 1911			

<i>Typhlomyrmex pusillus</i> Emery, 1894	-	-	2
<hr/>			
Subfamily FORMICINAE Latreille, 1809	<hr/>		
Tribe Camponotini Forel, 1878			
<i>Camponotus atriceps</i> (Smith, 1858)	15	23	1
<i>Camponotus excisus</i> Mayr, 1870	1	-	-
<i>Camponotus fastigatus</i> Roger, 1863	1	2	2
<i>Camponotus latangulus</i> Roger, 1863	-	1	-
<i>Camponotus linnaei</i> Forel, 1886	1	1	-
<i>Camponotus novogranadensis</i> Mayr, 1870	24	12	-
<i>Camponotus rapax</i> (Fabricius, 1804)	1	4	-
<i>Camponotus salvini</i> Forel, 1899	1	-	-
<i>Camponotus simillimus</i> (Smith, 1862)	-	-	2
<i>Camponotus</i> gp. <i>Tanaemyrmex</i> sp.3	1	-	-
Tribe Gigantiopini Ashmead, 1905			
<i>Gigantiops destructor</i> (Fabricius, 1804)	1	-	-
Tribe Lasiini Ashmead, 1905			
<i>Acropyga decedens</i> (Mayr, 1887)	-	1	2
<i>Acropyga exsanguis</i> (Wheeler, 1909)	1	-	-
<i>Acropyga fuhrmanni</i> (Forel, 1914)	-	1	1
<i>Acropyga guianensis</i> Weber, 1944	-	1	-
<i>Acropyga smithii</i> Forel, 1893	8	2	-
Tribe Myrmelachistini Forel, 1912			
<i>Myrmelachista guyanensis</i> Wheeler, 1934	-	1	-
Tribe Plagiolepidini Forel, 1886			
<i>Brachymyrmex cavernicola</i> Wheeler, 1938	8	7	4
<i>Brachymyrmex pictus</i> Mayr, 1887	1	-	-
<i>Brachymyrmex</i> sp.JTL07	4	2	2
<i>Nylanderia fulva</i> (Mayr, 1862)	1	-	-
<i>Nylanderia guatemalensis</i> (Forel, 1885)	10	2	15
<i>Nylanderia</i> sp.IT14.01	3	1	-
<i>Nylanderia</i> sp.SL13.03	1	1	2
<i>Nylanderia steinheili</i> (Forel, 1893)	27	24	8
<hr/>			
Subfamily HETEROPONERINAE Bolton, 2003	<hr/>		
Tribe Heteroponerini Bolton, 2003			
<i>Heteroponera flava</i> Kempf, 1962	-	-	2
<hr/>			
Subfamily MYRMICINAE Lepeletier de Saint-Fargeau, 1835	<hr/>		
Tribe Attini Smith, 1858			
<i>Acanthognathus brevicornis</i> Smith, 1944	-	-	1
<i>Acanthognathus ocellatus</i> Mayr, 1887	2	-	2
<i>Acanthognathus</i> sp.IT14.01	-	-	1
<i>Acromyrmex hystrix</i> (Latreille, 1802)	-	-	1
<i>Acromyrmex subterraneus brunneus</i> (Forel, 1912)	2	2	1
<i>Apterostigma acre</i> Latke, 1997	5	5	4

<i>Apterostigma auriculatum</i> Wheeler, 1925	2	-	1
<i>Apterostigma avium</i> Lattke, 1997	-	1	-
<i>Apterostigma chocoense</i> Lattke, 1997	1	3	1
<i>Apterostigma ierense</i> Weber, 1937	1	-	-
<i>Apterostigma pilosum</i> Mayr, 1865	21	6	8
<i>Apterostigma robustum</i> Emery, 1896	2	2	1
<i>Basiceros manni</i> Brown & Kempf, 1960	-	-	1
<i>Basiceros militaris</i> (Weber, 1950)	-	1	1
<i>Cyphomyrmex peltatus</i> Kempf, 1966	10	13	2
<i>Cyphomyrmex rimosus</i> (Spinola, 1851)	26	11	14
<i>Eurhopalothrix bolaii</i> (Mayr, 1870)	-	-	1
<i>Mycetophylax faunulus</i> (Wheeler, 1925)	1	-	-
<i>Mycetophylax morschi</i> (Emery, 1888)	1	-	-
<i>Mycetophylax strigatus</i> (Mayr, 1887)	2	-	-
<i>Mycocepurus smithii</i> (Forel, 1893)	-	2	-
<i>Myrmicocrypta</i> sp.IT14.01	-	-	1
<i>Myrmicocrypta</i> sp.IT14.02	1	1	-
<i>Myrmicocrypta</i> sp.IT14.03	11	2	1
<i>Octostruma balzani</i> (Emery, 1894)	16	5	19
<i>Octostruma bettschi</i> Perrault, 1988	13	15	21
<i>Octostruma iheringi</i> (Emery, 1888)	1	-	-
<i>Octostruma obtusidens</i> Longino, 2013	2	1	1
<i>Octostruma</i> sp.JTL035	-	1	2
<i>Procryptocerus scabriusculus</i> Forel, 1899	1	-	-
<i>Procryptocerus schmitti</i> Forel, 1901	-	-	1
<i>Procryptocerus</i> sp.IT14.01	1	-	-
<i>Rhopalothrix</i> sp.JTL014	1	-	-
<i>Sericomyrmex bondari</i> Borgmeier, 1937	1	2	5
<i>Sericomyrmex mayri</i> Forel, 1912	2	-	-
<i>Sericomyrmex parvulus</i> Forel, 1912	-	-	1
<i>Trachymyrmex cornetzi</i> (Forel, 1912)	2	-	-
<i>Trachymyrmex farinosus</i> (Emery, 1894)	5	3	-
<i>Trachymyrmex mandibularis</i> Weber, 1938	1	-	1
<i>Trachymyrmex</i> nr. <i>farinosus</i>	-	1	-
<i>Trachymyrmex</i> sp.A	3	3	-
<i>Trachymyrmex</i> sp.B	1	1	-
<i>Tranopelta gilva</i> Mayr, 1866	1	-	3
<i>Wasmannia auropunctata</i> (Roger, 1863)	13	8	3
<i>Wasmannia rochai</i> Forel, 1912	2	1	-
Tribe Crematogastrini Forel, 1893			
<i>Crematogaster brasiliensis</i> Mayr, 1878	17	12	3
<i>Crematogaster carinata</i> Mayr, 1862	-	1	-
<i>Crematogaster curvispinosa</i> Mayr, 1862	1	1	-

<i>Crematogaster flavosensitiva</i> Longino, 2003	16	5	-
<i>Crematogaster levior</i> Longino, 2003	-	4	-
<i>Crematogaster limata</i> Smith, 1858	15	-	-
<i>Crematogaster sotobosque</i> Longino, 2003	20	2	13
<i>Crematogaster tenuicula</i> Forel, 1904	-	15	-
<i>Crematogaster torosa</i> Mayr, 1870	1	-	-
Tribe Dacetini Forel, 1892			
<i>Strumigenys auctidens</i> (Bolton, 2000)	17	8	3
<i>Strumigenys beebeyi</i> (Wheeler, 1915)	5	-	1
<i>Strumigenys borgmeieri</i> Brown, 1954	1	-	-
<i>Strumigenys cincinnata</i> (Kempf, 1975)	-	5	-
<i>Strumigenys cosmostela</i> Kempf, 1975	4	2	-
<i>Strumigenys crassicornis</i> Mayr, 1887	1	-	3
<i>Strumigenys denticulata</i> Mayr, 1887	54	47	55
<i>Strumigenys diabola</i> Bolton, 2000	1	-	-
<i>Strumigenys dyseides</i> Bolton, 2000	2	-	4
<i>Strumigenys elongata</i> Roger, 1863	21	4	17
<i>Strumigenys perparva</i> Brown, 1958	3	4	17
<i>Strumigenys schulzi</i> Emery, 1894	3	5	14
<i>Strumigenys smithii</i> Forel, 1886	1	4	7
<i>Strumigenys spathula</i> Lattke & Goitía, 1997	1	-	-
<i>Strumigenys subedentata</i> Mayr, 1887	2	1	4
<i>Strumigenys trinidadensis</i> Wheeler, 1922	1	-	6
<i>Strumigenys trudifera</i> Kempf & Brown, 1969	10	2	-
<i>Strumigenys villiersi</i> (Perrault, 1986)	1	-	2
<i>Strumigenys waiwai</i> (Sosa-Calvo <i>et al.</i> , 2010)	-	1	-
Tribe Formicoxenini Forel, 1893			
<i>Ochetomyrmex neopolitus</i> Fernández, 2003	16	21	11
<i>Ochetomyrmex semipolitus</i> Mayr, 1878	7	5	2
Tribe Pheidolini Emery, 1877			
<i>Pheidole aenescens</i> Wilson, 2003	1	2	3
<i>Pheidole alienata</i> Borgmeier, 1929	-	1	-
<i>Pheidole allarmata</i> Wilson, 2003	20	16	25
<i>Pheidole araneoides</i> Wilson, 2003	13	11	1
<i>Pheidole biconstricta</i> Mayr, 1870	2	2	29
<i>Pheidole bruesi</i> Wheeler, 1911	-	1	4
<i>Pheidole bulliceps</i> Wilson, 2003	3	1	2
<i>Pheidole cataractae</i> Wheeler, 1916	1	-	-
<i>Pheidole coffeicola</i> Borgmeier, 1934	1	-	-
<i>Pheidole cramptoni</i> Wheeler, 1916	-	2	-
<i>Pheidole cursor</i> Wilson, 2003	9	5	1
<i>Pheidole deima</i> Wilson, 2003	-	14	1
<i>Pheidole embolopyx</i> Brown, 1968	4	1	3

1				
2				
3	<i>Pheidole fissiceps</i> Wilson, 2003	1	-	-
4	<i>Pheidole flavens</i> Roger, 1863	37	20	47
5	<i>Pheidole fracticeps</i> Wilson, 2003	1	1	-
6	<i>Pheidole gemmula</i> Wilson, 2003	-	-	4
7	<i>Pheidole</i> gp. <i>fallax</i> sp.D	10	13	-
8	<i>Pheidole</i> gp. <i>fallax</i> sp.F	1	15	5
9	<i>Pheidole</i> gp. <i>tristis</i> sp.B3	-	-	3
10	<i>Pheidole</i> gp. <i>tristis</i> sp.C	9	20	5
11	<i>Pheidole</i> gp. <i>tristis</i> sp.D	1	3	-
12	<i>Pheidole</i> gp. <i>tristis</i> sp.H1	-	1	1
13	<i>Pheidole guajirana</i> Wilson, 2003	40	23	12
14	<i>Pheidole jeannei</i> Wilson, 2003	2	4	3
15	<i>Pheidole lattkei</i> Wilson, 2003	1	1	-
16	<i>Pheidole longiseta</i> Wilson, 2003	4	3	3
17	<i>Pheidole lovejoyi</i> Wilson, 2003	6	6	2
18	<i>Pheidole meinertopsis</i> Wilson, 2003	3	-	-
19	<i>Pheidole microps</i> Wilson, 2003	3	1	9
20	<i>Pheidole midas</i> Wilson, 2003	4	-	-
21	<i>Pheidole nana</i> Emery, 1894	1	1	-
22	<i>Pheidole neoschultzi</i> LaPolla, 2006	18	11	1
23	<i>Pheidole nitella</i> Wilson, 2003	51	48	43
24	<i>Pheidole</i> nr. <i>araneoides</i>	-	-	3
25	<i>Pheidole</i> nr. <i>cramptoni</i>	-	5	5
26	<i>Pheidole</i> nr. <i>fimbriata</i> sp.A	1	-	-
27	<i>Pheidole</i> nr. <i>fimbriata</i> sp.C	3	2	3
28	<i>Pheidole</i> nr. <i>fimbriata</i> sp.D	5	-	-
29	<i>Pheidole</i> nr. <i>sagax</i>	1	3	-
30	<i>Pheidole</i> nr. <i>seeldrayersii</i>	2	-	-
31	<i>Pheidole</i> nr. <i>subarmata</i>	-	-	1
32	<i>Pheidole</i> nr. <i>tristis</i>	-	1	-
33	<i>Pheidole pedana</i> Wilson, 2003	27	15	9
34	<i>Pheidole rugiceps</i> Wilson, 2003	5	4	18
35	<i>Pheidole sagax</i> Wilson, 2003	2	6	2
36	<i>Pheidole sculptior</i> Forel, 1893	1	1	1
37	<i>Pheidole</i> sp.IT14.01	1	-	-
38	<i>Pheidole</i> sp.IT14.08	1	-	-
39	<i>Pheidole</i> sp.MI15.06	-	1	-
40	<i>Pheidole subarmata</i> Mayr, 1884	3	32	11
41	<i>Pheidole terribilis</i> Wilson, 2003	3	6	-
42	<i>Pheidole tristis</i> (Smith, 1858)	-	-	1
43	<i>Pheidole zelata</i> Wilson, 2003	3	3	-
44	<i>Pheidole zeteki</i> Smith, 1947	1	-	-
45	Tribe Pogonomyrmecini Ward <i>et al.</i> , 2015			

1				
2				
3	<i>Hylomyrma blandiens</i> Kempf, 1961	11	6	1
4	<i>Hylomyrma immanis</i> Kempf, 1973	2	8	20
5	<i>Hylomyrma reginae</i> Kutter, 1977	-	3	21
6				
7	Tribe Solenopsidini Forel, 1893			
8	<i>Carebara brevipilosa</i> Fernández, 2004	2	2	-
9	<i>Carebara reina</i> Fernández, 2004	8	5	9
10	<i>Carebara</i> sp.KW14.01	7	6	17
11	<i>Carebara urichi</i> (Wheeler, 1922)	8	6	4
12	<i>Megalomyrmex balzani</i> Emery, 1894	3	-	2
13	<i>Megalomyrmex drifti</i> Kempf, 1961	2	-	1
14	<i>Megalomyrmex incisus</i> Smith, 1947	-	1	1
15	<i>Megalomyrmex silvestrii</i> Wheeler, 1909	2	3	2
16	<i>Megalomyrmex</i> sp.MI15.01	1	-	-
17	<i>Rogeria besucheti</i> Kugler, 1994	-	-	1
18	<i>Rogeria blanda</i> (Smith, 1858)	1	-	1
19	<i>Rogeria germaini</i> Emery, 1894	-	-	2
20	<i>Rogeria innotabilis</i> Kugler, 1994	1	-	2
21	<i>Rogeria scobinata</i> Kugler, 1994	5	-	-
22	<i>Rogeria</i> sp.IT14.01	3	-	-
23	<i>Rogeria</i> sp.IT14.06	1	-	-
24	<i>Rogeria</i> sp.SL13.01	10	-	-
25	<i>Rogeria subarmata</i> (Kempf, 1961)	-	1	-
26	<i>Solenopsis bicolor</i> (Emery, 1906)	7	2	-
27	<i>Solenopsis</i> sp.01	7	21	23
28	<i>Solenopsis</i> sp.02	1	4	2
29	<i>Solenopsis</i> sp.03	15	26	7
30	<i>Solenopsis</i> sp.09	4	2	1
31	<i>Solenopsis</i> sp.11	32	2	23
32	<i>Solenopsis</i> sp.12	23	3	5
33	<i>Solenopsis</i> sp.13	37	38	20
34	<i>Solenopsis</i> sp.14	7	3	17
35	<i>Solenopsis</i> sp.15	46	50	54
36	<i>Solenopsis</i> sp.17	13	5	1
37	<i>Solenopsis</i> sp.21	21	12	12
38	<i>Solenopsis</i> sp.23	4	3	4
39	<i>Solenopsis</i> sp.24	7	5	2
40	<i>Solenopsis</i> sp.25	5	-	-
41	<i>Solenopsis</i> sp.28	1	2	-
42	<i>Solenopsis virulens</i> (Smith, 1858)	7	2	9
43				
44	Subfamily PONERINAE Lepeletier de Saint-Fargeau, 1835			
45				
46	Tribe Ponerini Lepeletier de Saint-Fargeau, 1835			
47	<i>Anochetus horridus</i> Kempf, 1964	11	7	6
48	<i>Anochetus inermis</i> André, 1889	2	2	15
49				
50				
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				

<i>Anochetus mayri</i> Emery, 1884	-	-	1
<i>Anochetus targionii</i> Emery, 1894	2	2	1
<i>Cryptopone holmgreni</i> (Wheeler, 1925)	-	1	2
<i>Hypoponera foreli</i> (Mayr, 1887)	12	8	24
<i>Hypoponera</i> sp.02.FGSGt	10	-	5
<i>Hypoponera</i> sp.12.FGSGt	14	6	19
<i>Hypoponera</i> sp.13.FGSGt	5	5	12
<i>Hypoponera</i> sp.B	14	9	3
<i>Hypoponera</i> sp.IT14.01	3	4	8
<i>Hypoponera</i> sp.IT14.02	8	8	24
<i>Hypoponera</i> sp.IT14.04	-	-	2
<i>Hypoponera</i> sp.IT14.05	9	2	-
<i>Hypoponera</i> sp.MI15.02	-	1	-
<i>Hypoponera</i> sp.MI15.05	-	-	1
<i>Leptogenys gaigei</i> Wheeler, 1923	-	3	1
<i>Leptogenys langi</i> Wheeler, 1923	1	1	2
<i>Leptogenys</i> sp.05.FGSGt	2	1	7
<i>Leptogenys</i> sp.IT14.04	-	-	1
<i>Mayaponera constricta</i> (Mayr, 1884)	8	4	6
<i>Neoponera apicalis</i> (Latreille, 1802)	1	5	2
<i>Neoponera commutata</i> (Roger, 1860)	-	3	1
<i>Neoponera crenata</i> (Roger, 1861)	-	-	1
<i>Neoponera foetida</i> (Linnaeus, 1758)	-	1	-
<i>Neoponera procidua</i> (Emery, 1890)	-	1	-
<i>Neoponera verena</i> Forel, 1922	3	5	-
<i>Odontomachus caelatus</i> Brown, 1976	2	2	-
<i>Odontomachus chelifer</i> (Latreille, 1802)	5	9	10
<i>Odontomachus haematodus</i> (Linnaeus, 1758)	1	5	-
<i>Odontomachus laticeps</i> Roger, 1861	4	2	1
<i>Odontomachus meinerti</i> Forel, 1905	8	9	11
<i>Odontomachus scalptus</i> Brown, 1978	4	-	4
<i>Pachycondyla crassinoda</i> (Latreille, 1802)	5	9	-
<i>Pachycondyla harpax</i> (Fabricius, 1804)	21	22	15
<i>Pachycondyla impressa</i> (Roger, 1861)	1	-	2
<i>Pachycondyla</i> nr. <i>lenkoi</i>	1	-	-
<i>Platythyrea pilosula</i> (Smith, 1858)	-	1	-
<i>Pseudoponera stigma</i> (Fabricius, 1804)	-	4	2
<i>Pseudoponera succedanea</i> (Roger, 1863)	-	-	6
<i>Rasopone arhuaca</i> (Forel, 1901)	-	-	1
<i>Rasopone</i> sp.JTL047	-	-	3
<i>Thaumatomyrmex soesilae</i> Makhan, 2007	-	1	-

Subfamily PROCERATIINAE Emery, 1895

Tribe Proceratiini Emery, 1895

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

<i>Discothyrea denticulata</i> Weber, 1939	1	1	1
<i>Discothyrea sexarticulata</i> Borgmeier, 1954	11	1	7
<hr/>			
Subfamily PSEUDOMYRMECINAE Smith, 1952			
<hr/>			
Tribe Pseudomyrmecini Smith, 1952			
<i>Pseudomyrmex tenuis</i> (Fabricius, 1804)	3	3	-
<hr/>			

For Peer Review Only

TABLE S4. Eigenvalues and trait loadings of the three first axes of a PCA performed on the matrix of mean trait values for each species (PCA_{TRAITS}).

	Axis 1	Axis 2	Axis 3
Eigenvalue	3.46	2.33	1.21
Percentage of variance	38.38	25.58	13.48
Cumulative percentage of variance	38.38	63.96	77.44
Trait loadings			
- Weber's length	-0.83	-0.08	0.15
- Relative head width	0.89	0.21	0.02
- Relative head length	0.88	0.17	-0.09
- Relative pronotum width	0.80	0.10	0.24
- Relative femur length	-0.44	0.81	-0.17
- Relative scape length	-0.27	0.83	-0.29
- Relative eye length	-0.27	0.70	0.42
- Relative clypeus length	0.35	0.55	0.52
- Relative mandible length	0.30	0.28	-0.76

TABLE S5. Eigenvalues and trait loadings of the two first axes of a PCA performed on the matrix of the 17 environmental variables measured at each plot (PCA_{ENV}). T_{MIN} = minimum temperature of the coldest month; T_{MEAN} = mean annual temperature; Gap = Gap fraction; C_H = Canopy height; C_{COV} = Canopy cover; Clay = percentage of clay; pH = soil pH; Silt = percentage of silt; Phosphorus = Olsen-phosphorus concentration; sand = percentage of sand; K = potassium concentration; C/N = carbon-to-nitrogen ratio; Na = sodium concentration; OM = percentage of organic matter; RH_{MIN} = minimum relative humidity of the driest month; RH_{MEAN} = mean annual relative humidity. The variables retained for analyses are highlighted in bold.

	Axis 1	Axis 2
Eigenvalue	10.28	3.52
Percentage of variance	60.45	20.70
Cumulative percentage of variance	60.45	81.15
Trait loadings		
- T _{MIN}	-0.98	0.01
- T _{MEAN}	-0.97	0.15
- Gap	-0.92	0.01
- C_H	-0.90	0.38
- C _{COV}	-0.86	-0.04
- Clay	-0.58	-0.73
- pH	-0.42	0.85
- Silt	0.54	0.70
- Phosphorus	0.59	0.54
- Sand	0.60	0.57
- K	0.63	0.48
- C/N	0.68	-0.69
- Na	0.72	0.12
- OM	0.77	-0.43
- RH _{MIN}	0.89	0.07
- Elevation	0.91	-0.30
- RH_{MEAN}	0.93	0.00