Microbial community structure reveals instability of nutritional symbiosis during the evolutionary radiation of Amblyomma ticks

Florian Binetruy, Marie Buysse, Quentin Lejarre, Roxanne Barosi, Manon Villa, Nil Rahola, Christophe Paupy, Diego Ayala, Olivier Duron

► To cite this version:

Florian Binetruy, Marie Buysse, Quentin Lejarre, Roxanne Barosi, Manon Villa, et al.. Microbial community structure reveals instability of nutritional symbiosis during the evolutionary radiation of Amblyomma ticks. *Molecular Ecology*, 2020, 29 (5), pp.1016-1029. 10.1111/mec.15373 . hal-03001756

HAL Id: hal-03001756

<https://hal.science/hal-03001756>

Submitted on 18 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Microbial community structure reveals instability of nutritional symbiosis
during evolutionary radiation of *Amblyomma* ticks**

Florian Binetruy^{a*}, Marie Buysse^a, Quentin Lejarre^{a,b}, Roxanne Barosi^a, Manon Villa^a, Nil
Rahola^{a,b}, Christophe Paupy^a, Diego Ayala^{a,b} and Olivier Duron^a

^a MIVEGEC (Maladies Infectieuses et Vecteurs : Ecologie, Génétique, Evolution et
Contrôle), Centre National de la Recherche Scientifique (CNRS) - Institut pour la Recherche
et le Développement (IRD) - Université de Montpellier (UM), Montpellier, France

^b CIRMF (Centre International de Recherches Médicales de Franceville), Franceville, Gabon

* Correspondance: florian.binetruy@ird.fr

Abstract

Mutualistic interactions with microbes have facilitated the adaptation of major eukaryotic lineages to restricted diet niches. Hence, ticks with their strictly blood-feeding lifestyle are associated with intracellular bacterial symbionts through an essential B vitamin supplementation. In this study, examination of the whole bacterial diversity in 25 tick species of the *Amblyomma* genus showed that three intracellular bacteria, *Coxiella*-like endosymbionts (LE), *Francisella*-LE and *Rickettsia*, are remarkably common. No other bacterium is so uniformly present in *Amblyomma* ticks. Almost all *Amblyomma* species were found to harbour a nutritive obligate symbiont, *Coxiella*-LE or *Francisella*-LE, able to synthesize B vitamins. However, despite the co-evolved and obligate nature of these mutualistic interactions, the structure of microbiomes does not mirror the *Amblyomma* phylogeny with a clear exclusion pattern between *Coxiella*-LE and *Francisella*-LE across tick species. *Coxiella*-LE, but not *Francisella*-LE, form evolutionarily stable associations with ticks commonly leading to co-cladogenesis. We further evidenced symbiont replacements during radiation of *Amblyomma*, with recent, and likely ongoing, invasions by *Francisella*-LE and subsequent replacements of ancestral *Coxiella*-LE through transient co-infections. Nutritional symbiosis in *Amblyomma* ticks is thus not a stable evolutionary state, but instead arises from conflicting origins between unrelated but competing symbionts with similar metabolic capabilities.

Introduction

Macro-organisms harbour complex microbial communities living inside and on their body (Margulis, 1993; Theis et al., 2016). These microbial communities, known as the microbiomes, can determine pivotal phenotypic traits of their hosts, driving a variety of ecological and evolutionary processes including major nutritive, reproductive and immune functions (Gould et al., 2018; Groussin et al., 2017; Hanning & Diaz-Sanchez, 2015; Ley et al., 2008; Turner, James, & Poole, 2013). As hosts vary in the microbiomes they harbour, an associated functionally important phenotypic variation exists within host populations (Falony et al., 2016; Ferrari & Vavre, 2011; Jaenike, 2012; Oliver, Russell, Moran, & Hunter, 2003; Scarborough, Ferrari, & Godfray, 2005). In arthropods, these microbiomes notably include highly specialized intracellular bacteria depending almost exclusively on maternal (transovarial) transmission to ensure their persistence in host populations (Moran, McCutcheon, & Nakabachi, 2008; Wernegreen, 2012). Some of these maternally inherited symbionts are essential for the life cycle of their arthropod hosts: They are obligate mutualists able to synthesize biochemical products favouring the specialization of arthropods to novel habitats or to particular feeding niches such as strict haematophagy or phloemophagy (Moran et al., 2008; Wernegreen, 2012). Overall, these mutualistic interactions have facilitated the radiation of major arthropod lineages, leading to remarkable host–symbiont phylogenetic congruence with a strict co-cladogenesis pattern in many cases (Chen, Li, & Aksoy, 1999; Duron et al., 2017; Jousset, Desclaux, & Coeur d’acier, 2009; Moran, Tran, & Gerardo, 2005; Takiya, Tran, Dietrich, & Moran, 2006).

Arthropods and beneficial maternally inherited symbionts can form evolutionary stable associations lasting for millions of years, but that are not necessarily permanent (Bennett & Moran, 2015; McCutcheon, Boyd, & Dale, 2019; Moran et al., 2008; Wernegreen, 2012).

Recent phylogenetic reconstructions suggest that beneficial symbiotic relationships can break down: Recently acquired symbionts can replace ancestral beneficial symbionts and provide similar benefits to the host (McCutcheon et al., 2019; Sudakaran, Kost, & Kaltenpoth, 2017). An alternative scenario is that recently acquired symbionts may cooperate with ancestral beneficial symbionts (Moran et al. 2008; Vautrin & Vavre, 2009). Vertical transmission actually locks the different symbionts together as coinfection and creates then privileged situations for symbiont–symbiont interactions, especially cooperation and dependence between symbionts. Functions of ancestral beneficial symbionts may be complemented by recently acquired cosymbionts and their coexistence can be ultimately stable over millions of years (Meseguer et al 2017). New beneficial symbionts often originate from microbes abundant in the host environment, potentially including entomopathogens, parasites vectored by arthropods or other maternally inherited symbionts, primarily facultative (i.e. not essential) for host survival (Koga & Moran, 2014; Matsuura et al., 2018; McCutcheon et al., 2019; Sachs, Skophammer, & Regus, 2011). These facultative symbionts, however, determine important traits in arthropods: protection against natural enemies, adaptation to changing environments or reproductive traits (Engelstädter & Hurst, 2009; Ferrari & Vavre, 2011; Moran et al., 2008; Oliver et al., 2003). Contrary to beneficial obligate symbionts, facultative symbionts undergo occasional horizontal transfers (HT) across arthropod species, resulting in limited phylogenetic congruence between hosts and symbionts (Duron, Wilkes, & Hurst, 2010; Jousset, Cœur d’Acier, Vanlerberghe-Masutti, & Duron, 2013; Russell et al., 2009). Overall, the diverse range of microbial lifestyle strategies creates a complex web of interactions mediating the dynamics of beneficial symbioses in arthropods (McCutcheon et al., 2019).

79 Co-existence of symbionts within microbial communities is expected to involve interactions
80 ranging from cooperation to competition and that can, in turn, determine aggregation and
81 exclusion patterns (Ferrari & Vavre, 2011; Moran et al., 2008; Vautrin & Vavre, 2009).
82 Exclusion patterns have been recently detected between maternally inherited symbionts of
83 ticks, suggesting that replacements of beneficial symbionts occur in this system (Duron et al.,
84 2017). Among arthropods, ticks (Arachnida: Ixodidae) are well known to engage in symbiotic
85 associations with at least 10 different genera of maternally inherited bacteria (Ahanitig,
86 Trinachartvanit, Baimai, & Grubhoffer, 2013; Duron et al., 2017). Ticks are specialized for an
87 exclusive diet of vertebrate blood, and have evolved intimate interactions with beneficial
88 symbionts that provide essential B vitamins and co-factors deficient in the blood diet (Bonnet,
89 Binetruy, Hernández-Jarguín, & Duron, 2017; Duron et al., 2017, 2018; Gerhart, Moses, &
90 Raghavan, 2016; Gottlieb, Lalzar, & Klasson, 2015; Guizzo et al., 2017; Hunter et al., 2015;
91 Olivieri et al., 2019; Smith, Driscoll, Gillespie, & Raghavan, 2015). Approximately two thirds
92 of tick species harbour *Coxiella*-like endosymbionts (*Coxiella*-LE hereafter), which are
93 required for tick survival and reproduction (Gottlieb et al., 2015; Guizzo et al., 2017; Smith et
94 al., 2015; Zhong, Jasinskas, & Barbour, 2007). *Coxiella*-LE genomes encode pathways for the
95 biosynthesis of major B vitamins and co-factors that fit closely with the expected nutritional
96 complements required for strict haematophagy (Gottlieb et al., 2015; Guizzo et al., 2017;
97 Smith et al., 2015). *Coxiella*-LE are abundant in two organs of ticks: ovaries, that is
98 consistent with vertical transmission into developing oocytes, and Malpighian tubules, where
99 B vitamins are possibly synthesized (Buysse, Plantard, McCoy, Duron, & Menard, 2019;
100 Wang et al., 2018)(Lalzar et al 2012). Owing to their maternal inheritance and beneficial
101 nature, *Coxiella*-LE are present in most individuals within infected host species [16, 41]. In
102 the *Rhipicephalus* tick genus, the acquisition of *Coxiella*-LE was followed by co-

diversification resulting in deeply congruent *Rhipicephalus*–*Coxiella*-LE phylogenies (Duron et al., 2017).

In a few tick species, however, *Coxiella*-LE are present at much lower frequencies than expected for obligate nutritional symbionts, suggesting that they are instead facultative symbionts in these hosts (Duron et al., 2015). Phylogenetic evidence corroborates this hypothesis, since closely related *Coxiella*-LE may infect distantly related tick species suggesting recurrent HT of some *Coxiella*-LE (Duron et al., 2015). In other tick species, no *Coxiella*-LE were detected at all, but alternative obligate beneficial symbionts have been identified or hypothesized (Duron et al., 2017, 2018; Gerhart et al., 2016; Kurtti et al., 2015; Olivieri et al., 2019). Indeed, *Francisella*-like endosymbionts (*Francisella*-LE) are commonly found in tick species lacking *Coxiella*-LE. A recent analysis of endosymbiotic communities in 81 tick species also showed that there is a significant exclusion pattern between *Francisella*-LE and *Coxiella*-LE (Duron et al., 2017). Like *Coxiella*-LE, *Francisella*-LE are essential for tick nutrition: Ticks deprived of their *Francisella*-LE completely cease development but resume normal growth upon supplementation with B vitamins (Duron et al., 2018). Genomes of *Francisella*-LE contain roughly the same biosynthesis pathways of B vitamins and co-factors as observed in *Coxiella*-LE genomes (Duron et al., 2018; Gerhart et al., 2016). *Francisella*-LE also presents the same tropism than *Coxiella*-LE: *Francisella*-LE are abundant in ovaries and Malpighian tubules of the ticks they infect (Duron et al 2018, current biology). Although *Francisella*-LE and *Coxiella*-LE are distantly related, they have converged towards an analogous nutritional mutualism with ticks [32, 33].

While *Coxiella*-LE symbioses are likely ancestral in ticks, replacements by *Francisella*-LE having recently transitioned to an endosymbiotic lifestyle (Duron et al., 2018; Gerhart et al.,

2016) appear across the tick phylogeny (Duron et al., 2017). Yet, the factors favouring *Francisella*-LE over *Coxiella*-LE in this evolutionary dynamic are not well understood. In this study, we examined the evolutionary dynamic of tick microbiomes, with a focus on beneficial nutritional symbioses, in an ecologically diverse genus of hard ticks, *Amblyomma*. This genus is the third largest in the family Ixodidae, with its species primarily occupying the tropical zones. The centre of species diversity is on the American continent, where half of all the 130 *Amblyomma* species are found (Guglielmone, Estrada-Peña, Keirans, & Robbins, 2003). The *Amblyomma* genus includes major vectors of tick-borne disease agents, including the lone star tick, *A. americanum*, which is the primary vector of *Ehrlichia* spp. (Childs & Paddock, 2003). Only a few studies have examined the microbial diversity in *Amblyomma* species, showing that they are infected either by *Coxiella*-LE or by *Francisella*-LE in addition to other maternally inherited bacteria (Binetrui, Dupraz, Buysse, & Duron, 2019; Budachetri et al., 2014; Clay et al., 2008; Duron et al., 2017; Gerhart et al., 2016). Here, we investigated the variation in tick microbial communities at different geographic and phylogenetic scales using a representative collection of specimens covering ca. 20% of *Amblyomma* species diversity. First, we reconstructed the *Amblyomma* phylogeny through the sequencing of large nuclear rDNA sequences (18S, ITS1, 5.8S, ITS2, and 28S rDNA). Second, we extensively characterized bacterial communities in *Amblyomma* species through a DNA barcoding approach targeting the 16S rDNA. Third, we traced the evolutionary histories of *Coxiella*-LE and *Francisella*-LE using multilocus sequence typing (MLST) systems. While the *Coxiella*-LE MLST already exists (Duron et al., 2015), in this study we developed a specific *Francisella*-LE MLST. Finally, we compared *Amblyomma* phylogeny with microbiome structure and further used co-phylogenetics, by comparing *Amblyomma*, *Coxiella*-LE and *Francisella*-LE phylogenies, to reveal the global dynamics of symbiotic interactions.

Materials and methods

Tick collection and processing

We examined a total of 144 tick specimens belonging to 25 *Amblyomma* species (1–11 specimens per species) collected from field sites in America and Africa or from laboratory colonies (Supplementary Table S1). Samples were preserved in 70% ethanol until use. To eliminate external (i.e. cuticular) microbes, tick specimens were surface cleaned with bleach prior to DNA extraction (Binetruy, Dupraz, et al., 2019). A few specimens (*A. loculosum*, $n=4$; *A. sculptum*, $n=1$; *Amblyomma* sp., $n=1$, obtained from a previous study (Duron et al., 2017)) were, however, not bleach-treated prior to DNA extraction (Supplementary Table S1). All tick DNA was individually extracted using the DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany).

Molecular typing of ticks

To reconstruct *Amblyomma* phylogeny, we sequenced almost complete rDNA sequences, including the 5' end of 18S, the entire ITS1, 5.8S and ITS2 sequences, and the 3' end of 28S rDNA (Supplementary Table S2) from 44 specimens (indicated in yellow in supplementary Table S1). Depending on the tick species, the size of the rDNA amplicon obtained varies from 4270 bp to 5054 bp. PCR products were purified using the kit CleanPCR (CleanNA, Waddinxveen, The Netherlands), fragmented to a size of 300 bp and further used to construct libraries with the kit Nextera XT (Illumina, San Diego, California, USA). These libraries were then pooled together, indexed and pair-end sequenced on a MiSeq (Illumina) sequencer using a flow cell equipped with a V3, 600-cycle reagent cartridge.

We obtained >24 million reads after quality filtering and removing adaptors using the Cutadapt tool on a Galaxy workbench (Goecks, Nekrutenko, Taylor, & The Galaxy Team,

2010; M. Martin, 2011). Paired-end reads were *de novo* assembled using metaSPAdes v3.11 with k-mer sizes of 21, 33, 55, and 73 bp (Nurk, Meleshko, Korobeynikov, & Pevzner, 2017). The resulting metaSPAdes contigs were binned into tick nuclear DNA and non-tick nuclear DNA groups through megablast in the GenBank nucleotide collection (Boratyn et al., 2013). These contigs were scaffolded, when applicable, by aligning to the complete sequences of the rRNA genes operon of *A. americanum* (GenBank AF291874), *A. hebraeum* (GenBank KY457489) and the partial sequence of *A. marmoreum* (GenBank KY457492) using MEGA (Kumar, Stecher, & Tamura, 2016).

Bacterial metabarcoding

A 251-bp portion of the V4 variable region of the bacterial 16S rRNA gene was amplified individually for each DNA sample using a Multiplex PCR Kit (Qiagen) and universal primers (16SV4F: 5'-GTGCCAGCMGCCGCGGTAA-3' and 16SV4R: 5'-GGACTACHVGGGTWTCTAATCC-3') (Galan et al., 2016). Amplified bacterial 16S rDNA products were purified and sequenced using an Illumina MiSeq platform (GenSeq, Montpellier University) and 251-bp end sequence reads were obtained. All bioinformatic analyses were conducted using the pipeline Frogs (<https://github.com/geraldinepascal/FROGS>) (Escudié et al., 2018) as previously described (Binetruy et al., 2019). One step of post-process operational taxonomic unit (OTU) affiliation was additionally performed through the Frogs pipeline on a Galaxy workbench (Escudié et al., 2018; Goecks et al., 2010). This step consists in aggregating OTUs that share 97% of identity within 99% of the amplicon length, and it reduced the probability of keeping artefactual OTUs by resolving multi-hit ambiguities. To control the contamination during these procedure negative controls were performed (three negative extraction controls were included in all extraction series, two negative PCR controls were included in all PCR series).

Moreover, OTUs having a maximal abundance in negative controls were discarded and false-positive OTUs were removed by filtering OTU representing less than 0.005% of the OTU total abundance (Bokulich et al., 2013). Following this procedure, the microbiome was determined individually for almost all *Amblyomma* specimens ($n=142/143$); however, the amount of DNA for the single specimen of *A. sculptum* was not sufficient.

A phylogenetic tree using OTU sequences and beta-diversity matrices based on this tree was assessed with FastTree and GUnifrac packages in R (J. Chen et al., 2012; Price, Dehal, & Arkin, 2009), using the computational procedure described in (Binetruy, Dupraz, et al., 2019). Multidimensional scaling (MDS) plots were then generated using the package ggplot2 in R (Wickham, 2016). Permutational multivariate analysis of variance (PERMANOVA) implemented in the vegan package in R or pairwise PERMANOVA (Arbizu, 2017/2018) was further performed on the generalized UniFrac ($\alpha = 0.5$) dissimilarity matrix to evaluate the potential impact of the presence of symbionts and co-infection on bacterial diversity. The P -values of the pairwise PERMANOVA were corrected for multiple comparisons using Holm's method (Holm, 1979). A Mantel test was used to examine the association between microbial diversity (GUnifrac distance) and *Amblyomma* phylogeny using the ecodist package in R ($n=9999$ permutations) (Goslee & Urban, 2007).

Multilocus typing of Coxiella-LE and Francisella-LE

Coxiella-LE were genotyped through nested or semi-nested PCR amplification and sequencing of three housekeeping genes (16S rRNA, *rpoB* and *groEL*) previously developed for the *Coxiella* MLST methodology (Duron et al., 2015) (Supplementary Table S2). No *Francisella*-LE MLST was previously developed and we thus used the published genome of *Francisella*-LE F-Om strain (isolated from the soft tick *Ornithodoros moubata* [32]; GenBank accession number: QAPC000000000) as reference to design specific PCR primers

for five genes (16S rRNA, *rpoB*, *groEL*, *ftsZ* and *gyrB*; Supplementary Table S2). Positive PCR products were purified and sequenced in both directions by Eurofins (Ebersberg, Germany). Sequence chromatograms were manually cleaned with Chromas Lite (http://www.technelysium.com.au/chromas_lite.html) and aligned with CLUSTALW implemented in MEGA 7 (Kumar et al., 2016; Thompson, Gibson, & Higgins, 2002).

Phylogenetic analyses

Phylogenetic relationships were assessed using *Amblyomma*, *Coxiella*-LE and *Francisella*-LE sequences produced in this study and additional sequences available in GenBank (including *Amblyomma*, *Coxiella*-LE and *Francisella*-LE relatives and outgroups). GBLOCKS (Castresana, 2000) was used to remove poorly aligned positions and to obtain unambiguous sequence alignments. All sequence alignments were also checked for putative recombinant regions using the RDP3 analysis package (D. P. Martin et al., 2010). The best fitting evolutionary models were determined using the Akaike information criterion and Bayesian information criterion with MEGA 7 (Kumar et al., 2016). Phylogenetic analyses were based on Bayesian inferences (BI) with MrBayes v3.2.7 (Ronquist et al., 2012). Two replicate analyses were run for 1 million generations. For each replicate, we ran one cold chain and three hot chains of the Markov chain Monte Carlo method, using a random starting tree and sampling trees every 100 generations and discarding 25% as burn-in. The remaining trees were used to calculate 50% majority-rule consensus trees.

The BI phylogenies of *Amblyomma*, *Coxiella*-LE and *Francisella*-LE were then used to conduct co-phylogenetic analyses using the Procrustean Approach to Cophylogeny (PACo) package in R (Balbuena, Míguez-Lozano, & Blasco-Costa, 2013; Hutchinson, Cagua, Balbuena, Stouffer, & Poisot, 2017). The significance of the co-phylogenetic tests was

established by 100,000 random permutations of the two-association matrix. To test an effect of *Rickettsia* co-infection on the evolution of *Francisella*-LE and *Coxiella*-LE, linear regressions were computed in R using the matrix of genetic distances of these symbionts and the presence/absence of *Rickettsia* as an explicative variable.

Ethics statement

The use of the genetic resources was declared to the French Ministry of the Environment (reference TREL19028117S/156) and to Gabon government (entry authorization #AE16008/PR/ANPN/SE/CS/AEPN and #research authorization #AR0013/16/MESRS/CENAREST/CG/CST/CSAR).

Results

Phylogeny of Amblyomma ticks

We first reconstructed the phylogenetic relationships among the 25 *Amblyomma* species using BI analyses based on large fragments of nuclear rDNA sequences obtained from one-to-two specimens per species (Figure 1). The findings support a monophyletic origin of the five African *Amblyomma* species (AF group): They cluster together in a robust clade nested among the 20 New World *Amblyomma* species (NW group), suggesting an American origin of these African species (Figure 1). Moreover, the phylogeny of *Amblyomma* also parallels the tick host-range at least in the NW group: Closely related *Amblyomma* species often share the same host species, such as *A. dissimile* and *A. rotundatum* feeding on poikilotherms (reptiles and amphibians) or *A. latepunctatum*, *A. scalpturatum* and *A. naponense* feeding mainly on tapirs and Suidae (Figure 1). However, in other cases, closely related *Amblyomma* species use different host species, such as *A. ovale*, a generalist species feeding on a diversity

of domestic and wild animals, and *A. varium*, a specialized species feeding on arboreal vertebrates.

Microbial diversity and symbiont prevalence

We further examined the whole bacterial diversity of 24 out of the 25 *Amblyomma* species ($n=142$ specimens, one-to-ten specimens were examined per species) via high-throughput 16S rDNA sequencing. The amount of DNA for the 25th *Amblyomma* species, *A. sculptum* ($n=1$), was not sufficient to perform bacterial barcoding (this specimen was already PCR-typed for *Coxiella*-LE and *Francisella*-LE, as further detailed). After filtration of false-positive OTUs and contaminants (Binetruy, Dupraz, et al., 2019; Birer, Tysklind, Zinger, & Duplais, 2017), 4,364,360 reads distributed in 195 OTUs were obtained (Table S3). *Coxiella*-LE and *Francisella*-LE were the most abundant bacterial genera representing 33.6% and 33.1% of the total number of reads, respectively (Figure 2 and Supplementary Table S1). Other bacterial genera were found: In most cases, each represented a negligible part of the 16S rDNA reads when present. A remarkable exception to this pattern was the presence of abundant intracellular bacteria belonging to the *Rickettsia* genus in 16 *Amblyomma* species (Figure 2 and Supplementary Table S1). Other exceptions included the *A. romitii* samples for which no *Coxiella*-LE and *Francisella*-LE reads were detected: Most reads were assigned to a tick-borne pathogen, *Ehrlichia* sp. In *A. loculosum* and *Amblyomma* sp. samples, *Coxiella*-LE reads were detected but reads of other bacteria were more abundant. Since the *A. loculosum* and *Amblyomma* sp. samples were not bleach-treated prior to DNA extraction (as done for the other samples), the abundant presence of these bacteria may be due to the cuticular bacteriome, as recently observed in another *Amblyomma* species (Binetruy, Dupraz, et al., 2019).

The structure of microbial diversity was not globally impacted by the phylogenetic proximity among *Amblyomma* species: The dendrogram of microbial diversity did not parallel the *Amblyomma* phylogeny (Mantel two-tailed test, $R=0.16$, $P=0.19$) (Supplementary Figure S1). However, the MDS plot suggested an effect of tick species on bacterial diversity (Supplementary Figure S2), as confirmed by the PERMANOVA analysis ($R^2=0.51$, $P=0.001$). Further testing showed a clear separation between *Coxiella*-LE-infected tick species/specimens and *Francisella*-LE-infected ones (pairwise PERMANOVA analysis, $R^2=0.31$, adjusted P for multiple comparisons=0.0003; Figure 3A). *Rickettsia* also structures the bacterial diversity but to a lesser extent than *Coxiella*-LE and *Francisella*-LE (PERMANOVA, $R^2=0.11$, $P=0.001$; Figure 3B). The *Coxiella*-LE and *Francisella*-LE clusters are actually structured into two sub-clusters each, fitting with the presence of *Rickettsia* (Figure 3c) as corroborated by PERMANOVA analyses: (1) *Coxiella*-LE without *Rickettsia* vs. *Coxiella*-LE with *Rickettsia* ($R^2=0.2$, adjusted P for multiple comparisons=0.0015), and (2) *Francisella*-LE without *Rickettsia* vs. *Francisella*-LE with *Rickettsia* ($R^2=0.22$, adjusted P for multiple comparisons=0.0015).

Of the 25 *Amblyomma* species (including *A. sculptum*) examined for the presence of *Coxiella*-LE and *Francisella*-LE, 24 were infected by one or both of these symbionts: 11 *Amblyomma* species harbour only *Coxiella*-LE, 13 species only *Francisella*-LE and three species both (Figure 2, Supplementary Table S1). Only *A. romitii* was not infected by *Coxiella*-LE and *Francisella*-LE but this may be explained by the presence of the tick-borne pathogen *Ehrlichia* sp., which may mask the presence of other bacteria. In the 24 infected *Amblyomma* species, *Coxiella*-LE and *Francisella*-LE were not randomly associated (Fisher's exact test, $P=0.001$): These two symbionts co-occurred in the same tick species less frequently than expected by chance (exclusion pattern), meaning that their distribution across tick species was

strongly dependent on each other. Conversely, neither *Coxiella*-LE nor *Francisella*-LE showed a non-random association with *Rickettsia* (Fisher's exact tests, $P=0.99$ and 0.68 , respectively). However, *Coxiella*-LE and *Francisella*-LE had a patchy and quite uniform distribution along the *Amblyomma* phylogeny: While some closely related *Amblyomma* species were infected by the same symbiont genus (e.g. *A. dissimile* and *A. rotundatum* by *Francisella*-LE, or *A. latepunctatum*, *A. sculpturatum* and *A. naponense* by *Coxiella*-LE), others were not (e.g. *A. americanum*, infected by *Coxiella*-LE, and *A. oblongoguttatum* by *Francisella*-LE) (Figure 2). This distribution pattern paralleled partly the tick host-range: Indeed, *A. dissimile* and *A. rotundatum* that are related and specialized for poikilotherms were both infected by *Francisella*-LE (Figure 1, 2).

In the tick species they infect, *Coxiella*-LE and *Francisella*-LE were present in most specimens examined (Figure 2, Supplementary Table S1). However, a more contrasted pattern was apparent in the three *Amblyomma* species that were co-infected by *Coxiella*-LE and *Francisella*-LE: (i) in *A. geayi*, of the 10 examined specimens, seven were infected by *Coxiella*-LE in one locality, but in other localities, two specimens were infected by *Francisella*-LE and one was co-infected by *Coxiella*-LE and *Francisella*-LE; (ii) in *A. latepunctatum*, of the four examined specimens, three specimens of the same locality were co-infected by *Coxiella*-LE and *Francisella*-LE but one specimen from another locality was only infected by *Coxiella*-LE; (iii) in *A. sculptum*, the single examined specimen was co-infected by *Coxiella*-LE and *Francisella*-LE (Figure 2, Supplementary Table S1). In contrast to *Coxiella*-LE and *Francisella*-LE, the prevalence of *Rickettsia* was heterogeneous, with infection frequencies ranging from 14% to 100% depending on tick species (Figure 2, Supplementary Table S1). In all cases, *Rickettsia* was found with either *Coxiella*-LE (seven

Amblyomma species) or *Francisella*-LE (seven species) or both (two species). Neither
Rickettsia nor *Coxiella*-LE nor *Francisella*-LE shows infection-biased sex ratio: within each
Amblyomma species, the prevalence of infection did not differ between males and females
(Fisher's exact test, all $P > 0.1$).

Evolutionary history of Coxiella-LE and Francisella-LE symbioses

Sequencing of three *Coxiella* MLST genes (16S rRNA, *rpoB* and *groEL*) led to the
identification of 14 genetically different *Coxiella*-LE in a subset of 32 specimens representing
the 14 infected *Amblyomma* species (one to four specimens per species were examined). Each
Amblyomma species was infected by a genetically distinct *Coxiella*-LE and no variation of
Coxiella-LE was observed among specimens belonging to the same *Amblyomma* species. We
observed no sign of recombination in the *Coxiella*-LE data set (all $P > 0.05$ for the
GENECONV and RDP recombination-detection tests) and we thus used the 16S rRNA, *rpoB*
and *groEL* concatenated sequences for BI analyses. Comparisons with other sequences
available on GenBank showed that the *Coxiella*-LE of *Amblyomma* are polyphyletic: They
were scattered into different well-supported clusters among *Coxiella*-LE of other tick species
(Supplementary Figure S3). Indeed, the *Coxiella*-LE of *A. variegatum*, *A. tholloni* and *A.*
splendidum form a monophyletic clade that is more closely related to the *Coxiella*-LE of
Ixodes tick species than to the *Coxiella*-LE of other *Amblyomma* species. Similarly, the
Coxiella-LE clade of *A. cajennense*, *A. sculptum* and *A. americanum* is more closely related to
the *Coxiella*-LE of *Dermacentor* tick species. This pattern is suggestive of recurrent HT
events of *Coxiella*-LE among tick species. However, the *Coxiella*-LE clusters of *Amblyomma*
species can be gathered into two main groups, one with all the *Coxiella*-LE of NW
Amblyomma species and the other with all the *Coxiella*-LE of AF *Amblyomma* species
(Supplementary Figure S3). This pattern suggests an effect of phylogeographic drivers in

structuring the evolution of *Coxiella*-LE. This is strongly supported by the co-phylogeny analysis between *Coxiella*-LE and *Amblyomma* phylogenies: There is a significant topological congruence between their phylogeny (PACo analysis, $P=0.0001$; Figure 4A). This shows that co-cladogenesis with *Coxiella*-LE occurred during the radiation of *Amblyomma*.

Sequencing of five *Francisella* MLST genes (16S rRNA, *rpoB*, *groEL*, *ftsZ* and *gyrB*) led to the identification of 15 genetically different *Francisella*-LE in a subset of 28 tick specimens representing the 13 infected *Amblyomma* species (one to four specimens per species were used). There was only one *Francisella*-LE in each *Amblyomma* species, except *A. pacae* and *A. rotundatum* in which two and three genetically distinct *Francisella*-LE, respectively, were present in specimens from different localities. Each *Amblyomma* species harbours genetically distinct *Francisella*-LE, except for *A. geayi* and *A. latepunctatum* that harbour identical *Francisella*-LE on the basis of their MSLT sequences (Supplementary Figure S4). Owing to the lack of *Francisella*-LE *rpoB*, *groEL*, *ftsZ* and *gyrB* gene sequences available in GenBank before this study (with the exception of two published *Francisella*-LE genomes), the BI phylogenetic analysis between the *Francisella*-LE of *Amblyomma* and those of other tick species (with sequences available in GenBank) was made using only their 16S rRNA nucleotidic sequences (Supplementary Figure S5). No *Francisella*-LE subclade specific to *Amblyomma* exists along the 16S rRNA phylogenetic tree: The *Francisella*-LE of *Amblyomma* are instead scattered among *Francisella*-LE of other tick genera suggesting recurrent HT events among unrelated tick species (Supplementary Figure S5). However, the inner topology of the *Francisella*-LE clade based on 16S rRNA gene sequences remained too poorly resolved in many cases (as shown by low support values of inner branches) to infer the exact relatedness among all *Francisella*-LE. We thus further reconstructed the phylogenetic relationships between *Francisella*-LE using BI analyses based on their 16S rRNA, *rpoB*,

groEL, *ftsZ* and *gyrB* nucleotidic sequences. We observed no sign of recombination in the *Francisella*-LE data set (all $P > 0.05$ for the GENECONV and RDP recombination-detection tests) and we thus used the 16S rRNA, *rpoB*, *groEL*, *ftsZ* and *gyrB* concatenated sequences for analyses. Conversely to *Coxiella*-LE, there was no apparent co-cladogenesis or phylogeographic pattern along the *Francisella*-LE phylogeny (Supplementary Figure S4). Indeed, the *Francisella*-LE of *A. sculptum* (NW group) is closely related to the *Francisella*-LE of an unrelated *Amblyomma* species, *A. paulopunctatum* (AF group). In addition, the *Francisella*-LE of *A. sculptum* and *A. paulopunctatum* are more closely related to the *Francisella*-LE of the soft tick *O. moubata* than to the *Francisella*-LE of other *Amblyomma* species. The co-phylogeny analysis also showed no significant signal of congruence between the *Francisella*-LE and *Amblyomma* phylogenies (PACo analysis, $P = 0.06$): Only HT events seem to have impacted the distribution *Francisella*-LE across *Amblyomma* phylogeny (Figure 4B). Interestingly, while co-infection with *Rickettsia* along the *Coxiella*-LE phylogeny is random (linear model, adjusted $R^2 = -0.01$, F-stat=0.60, $P = 0.56$; Supplementary Figure S6A), it is not so along the *Francisella*-LE phylogeny (adjusted $R^2 = 0.15$, F-stat=6.90, $P = 0.002$; Figure S6B): Co-infections with *Rickettsia* are more common with certain *Francisella*-LE subclades than with others.

Discussion

Three intracellular bacterial genera, *Coxiella*-LE, *Francisella*-LE and *Rickettsia*, are widespread across the 25 species of *Amblyomma* ticks we examined in this study. Only a few other bacteria have been detected and none is so uniformly present in *Amblyomma*. However, the structure of the microbiomes does not mirror the *Amblyomma* phylogeny and closely related *Amblyomma* commonly harbour divergent microbiomes. As expected, almost all *Amblyomma* species were found to harbour a nutritive obligate symbiont, *Coxiella*-LE or

Francisella-LE, both able to synthesize B vitamins but with a clear exclusion pattern between them: *Coxiella*-LE was found as a single infection in 11 *Amblyomma* species, *Francisella*-LE in 13 species and co-infection was seen in only three species. Despite the co-evolved and obligate interactions of ticks with their mutualistic partners, we detected evidence of symbiont replacements during radiation of *Amblyomma*, raising questions regarding the ecological and evolutionary factors underlying replacements.

The comparison of symbiont and tick phylogenies revealed that the *Coxiella*-LE symbiosis is ancient and arose in the early evolution of the *Amblyomma* genus. Hence, some *Coxiella*-LE are specialized for their *Amblyomma* hosts, with an ancient acquisition followed by co-diversification, meaning that the persistence of *Coxiella*-LE through vertical transmission is stable over the duration of *Amblyomma* species diversification. A very similar co-diversification pattern has also been reported for *Coxiella*-LE symbiosis in the *Rhipicephalus* tick genus (Duron et al., 2017). However, the spread of *Coxiella*-LE was more complex in *Amblyomma*: The infections found in some *Amblyomma* species are distantly related and do not form an *Amblyomma*-specific clade. Rather, phylogenetics shows that *Coxiella*-LE of *Amblyomma* are actually scattered among *Coxiella*-LE of other tick genera such as *Ixodes* and *Dermacentor*. Only extensive HT of *Coxiella*-LE among tick genera may explain these phylogenetic incongruences. Since facultative, but not obligate, symbionts can undergo HT between host species (Bennett & Moran, 2015; McCutcheon et al., 2019; Nancy A. Moran et al., 2008; Wernegreen, 2012), this suggests that some *Coxiella*-LE are facultative symbionts of ticks. Interestingly, *Coxiella*-LE is a facultative symbiont in some *Ixodes* species, such as *I. ricinus* and *I. uriae* (Duron et al., 2017; Duron, Jourdain, & McCoy, 2014; Duron et al., 2015). The phylogenetic proximity of *Coxiella*-LE of *Ixodes* spp. with the *Coxiella*-LE of *A. variegatum*, *A. splendidum* and *A. tholloni* thus suggests that a facultative *Coxiella*-LE of

451 *Ixodes* spp. had an early jump to the *Amblyomma* ancestor of these species before replacing
 452 the ancestral obligate symbiont and evolving obligate nutritional symbioses with current
 453 species.
 454

455 The infection dynamics of *Francisella*-LE is different to that of *Coxiella*-LE. While the
 456 *Francisella*-LE of ticks form a monophyletic clade within the *Francisella* genus (Duron et al.,
 457 2017, 2018), we observed frequent HT events between unrelated tick species. No co-
 458 cladogenesis signal with *Amblyomma* is apparent along the phylogenies, meaning that current
 459 *Francisella*-LE arose only recently in this genus. The presence of unrelated *Francisella*-LE in
 460 *Amblyomma* further indicates that several independent acquisitions of *Francisella*-LE have
 461 occurred during the radiation of this tick genus. These acquisitions have likely come at the
 462 expense of the *Coxiella*-LE with their ultimate replacement by *Francisella*-LE. The AF
 463 *Amblyomma* group is illustrative of this process: Most species (*A. splendidum*, *A. variegatum*,
 464 *A. tholloni* and *A. loculosum*) are infected by *Coxiella*-LE with a strong co-cladogenesis
 465 pattern, but one species (*A. paulopunctatum*) is infected by *Francisella*-LE. This pattern
 466 suggests that *Francisella*-LE has replaced the *Coxiella*-LE primarily present in the *A.*
 467 *paulopunctatum* ancestor. Other examples include the monophyletic group formed by *A.*
 468 *americanum*, *A. oblongoguttatum*, *A. cajennense* and *A. sculptum*: *Coxiella*-LE has co-
 469 diverged with all species but one, since here *Francisella*-LE eliminated the *Coxiella*-LE
 470 primarily present in the *A. oblongoguttatum* ancestor. In addition, *A. sculptum* is co-infected
 471 at the individual level by an ancestral *Coxiella*-LE (i.e. showing a co-cladogenesis pattern
 472 with *Coxiella*-LE of the *A. sculptum* relatives) and a recently acquired *Francisella*-LE. This
 473 pattern suggests that HT of *Francisella*-LE within *Amblyomma* communities is recent, likely
 474 ongoing, and that co-infections with ancestral *Coxiella*-LE are only transitory. The
 475 preferential association of some *Francisella*-LE with *Rickettsia* further implies that these co-

infections may be important drivers since some *Rickettsia* are also able to synthesize folate (B9 vitamin) (Hunter et al., 2015) and thus to participate in nutritional symbiosis along with *Francisella*-LE. Under this hypothesis, *Francisella*-LE and *Rickettsia* are cooperating together, and each may fulfil essential metabolic functions not ensured by the others. They may also act together to replace ancestral *Coxiella*-LE.

Some biological traits of these symbioses are indicative of how and why *Francisella*-LE and some *Coxiella*-LE are both eliminating ancestral *Coxiella*-LE in *Amblyomma* ticks. *Francisella*-LE or *Coxiella*-LE were occasionally detected in the salivary glands of several tick species (Budachetri et al., 2014; Buysse et al., 2019; Klyachko, Stein, Grindle, Clay, & Fuqua, 2007) suggesting that ticks may inject part of their symbionts during feeding. Ticks, unlike other arthropod vectors, often attach and aggregate on the host for several days to obtain a meal, a process termed ‘co-feeding’. The spatiotemporal proximity of ticks during co-feeding may favour the HT of *Francisella*-LE and *Coxiella*-LE between conspecifics but also between different tick species, as commonly observed for tick-borne pathogens (Voordouw, 2015; Wright, Sonenshine, Gaff, & Hynes, 2015). This process may lead to local or systemic infections in vertebrates since a few cases of opportunistic *Coxiella*-LE infections have been reported after tick feeding (Shivaprasad et al., 2008; Vapniarsky, Barr, & Murphy, 2012; Woc-Colburn et al., 2008, p.). This mode of transmission may be particularly significant for *Francisella*-LE and *Coxiella*-LE by leading to co-infections with ancestral *Coxiella*-LE in ticks. Interestingly, we found two *Amblyomma* species, *A. geayi* and *A. latepunctatum*, that are co-infected by their respective ancestral *Coxiella*-LE but that also share the same *Francisella*-LE, which is also closely related to the *Francisella*-LE of *A. dissimile*. The genetic proximity of *Francisella*-LE in these three unrelated *Amblyomma* species is suggestive of recent HT events through co-feeding: While these three *Amblyomma*

species feed on very different vertebrate hosts, hampering the possibility of co-feeding, the immature stages of *A. dissimile* are commonly found in diverse mammals or birds (Binetruy, Chevillon, de Thoisy, Garnier, & Duron, 2019; Guglielmone & Nava, 2010; Scott & Durden, 2015), suggesting that they may be ecological bridges driving HT of *Francisella*-LE across tick species.

By replacing the ancestral *Coxiella*-LE, the novel symbiont colonizes a pre-adapted tick physiological environment that requires the provision of B vitamin: Thus, it must be able to synthesize these compounds as the ancestral *Coxiella*-LE did. The presence of several B vitamin biosynthesis pathways is ancestral in the *Coxiella* and *Francisella* genera, and all members of these genera, including pathogenic species that are also all intracellular, have conserved these abilities through their radiation (Duron et al., 2018; Gerhart et al., 2016; Meibom & Charbit, 2010; Rowe & Huntley, 2015; Smith et al., 2015; van Schaik, Chen, Mertens, Weber, & Samuel, 2013): All *Francisella*-LE and *Coxiella*-LE are already pre-adapted to nutritional symbioses with ticks. Beyond B vitamins, the replacement of ancestral *Coxiella*-LE suggests that the new symbiont could supply an additional benefit that the ancestral *Coxiella*-LE was unable to supply to ticks, thereby out-competing them. The genomes of ancient beneficial endosymbionts have lost most of their gene contents from their ancestor, being usually small in size and dense in gene content but also suffering Muller's ratchet, with fixation of deleterious mutations through genetic drift (McCutcheon et al., 2019; McCutcheon & Moran, 2012; N. A. Moran, 1996; Rispe & Moran, 2000). This evolution towards massive genomic reduction is obvious for the *Coxiella*-LE of *A. americanum*, which have a genome of only 0.66 Mb (Smith et al., 2015), but not for the *Francisella*-LE of *A. maculatum*: Its genome is 1.56 Mb, and although half is pseudogenized, it may have a higher biosynthetic capability (Gerhart et al., 2016). Similar variation is also reported between

Coxiella-LE with some genomes reaching ca. 1.5 Mb (Gottlieb et al., 2015; Ramaiah & Dasch, 2018), suggesting that some *Coxiella*-LE have greater biosynthetic capabilities than others. However, other processes may act on the *Coxiella*-LE replacement. Indeed, ancestral *Coxiella*-LE may have evolved too reduced (degraded) genomes and become maladapted, opening the road to replacement by a new symbiont. This degeneration–replacement model has been proposed for other arthropods such as cicadas (Campbell et al., 2015; Łukasik et al., 2018; Matsuura et al., 2018), but replacements are expected to be transient making them difficult to observe (McCutcheon et al., 2019). In *Amblyomma*, the observations of three species with co-infections by ancestral *Coxiella*-LE and recently acquired *Francisella*-LE may correspond to this transient state before extinction of *Coxiella*-LE.

That ecological specialization to strict haematophagy is driven by nutritional symbiotic interactions is beyond doubt for ticks. The present study, nevertheless, shows that nutritional symbiosis in the *Amblyomma* genus is not stable state, being impacted by competition between *Coxiella*-LE and *Francisella*-LE or between *Coxiella*-LE themselves. We potentially underestimate the amplitude of this dynamics: our intraspecific sampling was low for some tick species and this may have led to an underestimation of the *Francisella*-LE frequency. Indeed, in the cases of low prevalence, most samples will be found not infected by *Francisella*-LE while other members of the species, not sampled and tested, are in fact infected. Precisely, this pattern is found in this study: we found variation *Francisella*-LE infection pattern between sampling localities in three tick species in which just few few specimens were infected, which clearly indicates this potential for false negatives arising from insufficient sampling. Anyway, ticks now march on with their recently acquired *Francisella*-LE, at the expense of *Coxiella*-LE, but the precise mechanisms providing the advantage to *Francisella*-LE remain to be determined, including endosymbiosis intrinsic factors, such as

competition between symbionts with similar metabolic capabilities and the differential degree of genome reduction.

References

- Ahantarig, A., Trinachartvanit, W., Baimai, V., & Grubhoffer, L. (2013). Hard ticks and their bacterial endosymbionts (or would be pathogens). *Folia Microbiologica*, 58(5), 419–428. doi: 10.1007/s12223-013-0222-1
- Arbizu, P. M. (2018). *Pairwise multilevel comparison using adonis. Contribute to pmartinezarbizu/pairwiseAdonis development by creating an account on GitHub* [R]. Retrieved from <https://github.com/pmartinezarbizu/pairwiseAdonis> (Original work published 2017)
- Balbuena, J. A., Míguez-Lozano, R., & Blasco-Costa, I. (2013). PACo: A Novel Procrustes Application to Cophylogenetic Analysis. *PLOS ONE*, 8(4), e61048. doi: 10.1371/journal.pone.0061048
- Bennett, G. M., & Moran, N. A. (2015). Heritable symbiosis: The advantages and perils of an evolutionary rabbit hole. *Proceedings of the National Academy of Sciences of the United States of America*, 112(33), 10169–10176. doi: 10.1073/pnas.1421388112
- Binetruy, F., Chevillon, C., de Thoisy, B., Garnier, S., & Duron, O. (2019). Survey of ticks in French Guiana. *Ticks and Tick-Borne Diseases*, 10(1), 77–85. doi: 10.1016/j.ttbdis.2018.09.003
- Binetruy, F., Dupraz, M., Buysse, M., & Duron, O. (2019). Surface sterilization methods impact measures of internal microbial diversity in ticks. *Parasites & Vectors*, 12(1), 268. doi: 10.1186/s13071-019-3517-5
- Birer, C., Tysklind, N., Zinger, L., & Duplais, C. (2017). Comparative analysis of DNA extraction methods to study the body surface microbiota of insects: A case study with ant cuticular bacteria. *Molecular Ecology Resources*, 17(6), e34–e45. doi: 10.1111/1755-0998.12688

575 Bokulich, N. A., Subramanian, S., Faith, J. J., Gevers, D., Gordon, J. I., Knight, R., ... Caporaso, J. G.
 576 (2013). Quality-filtering vastly improves diversity estimates from Illumina amplicon
 577 sequencing. *Nature Methods*, 10(1), 57–59. doi: 10.1038/nmeth.2276

578 Bonnet, S. I., Binetruy, F., Hernández-Jarguín, A. M., & Duron, O. (2017). The Tick Microbiome: Why
 579 Non-pathogenic Microorganisms Matter in Tick Biology and Pathogen Transmission. *Frontiers*
 580 *in Cellular and Infection Microbiology*, 7. doi: 10.3389/fcimb.2017.00236

581 Boratyn, G. M., Camacho, C., Cooper, P. S., Coulouris, G., Fong, A., Ma, N., ... Zaretskaya, I. (2013).
 582 BLAST: a more efficient report with usability improvements. *Nucleic Acids Research*, 41(Web
 583 Server issue), W29–W33. doi: 10.1093/nar/gkt282

584 Budachetri, K., Browning, R. E., Adamson, S. W., Dowd, S. E., Chao, C.-C., Ching, W.-M., & Karim, S.
 585 (2014). An insight into the microbiome of the *Amblyomma maculatum* (Acari: Ixodidae).
 586 *Journal of Medical Entomology*, 51(1), 119–129. doi: 10.1603/me12223

587 Buysse, M., Plantard, O., McCoy, K. D., Duron, O., & Menard, C. (2019). Tissue localization of *Coxiella*-
 588 like endosymbionts in three European tick species through fluorescence in situ hybridization.
 589 *Ticks and Tick-Borne Diseases*, 10(4), 798–804. doi: 10.1016/j.ttbdis.2019.03.014

590 Campbell, M. A., Leuven, J. T. V., Meister, R. C., Carey, K. M., Simon, C., & McCutcheon, J. P. (2015).
 591 Genome expansion via lineage splitting and genome reduction in the cicada endosymbiont
 592 *Hodgkinia*. *Proceedings of the National Academy of Sciences*, 112(33), 10192–10199. doi:
 593 10.1073/pnas.1421386112

594 Castresana, J. (2000). Selection of Conserved Blocks from Multiple Alignments for Their Use in
 595 Phylogenetic Analysis. *Molecular Biology and Evolution*, 17(4), 540–552. doi:
 596 10.1093/oxfordjournals.molbev.a026334

597 Chen, J., Bittinger, K., Charlson, E. S., Hoffmann, C., Lewis, J., Wu, G. D., ... Li, H. (2012). Associating
 598 microbiome composition with environmental covariates using generalized UniFrac distances.
 599 *Bioinformatics (Oxford, England)*, 28(16), 2106–2113. doi: 10.1093/bioinformatics/bts342

- Chen, X., Li, S., & Aksoy, S. (1999). Concordant evolution of a symbiont with its host insect species: molecular phylogeny of genus *Glossina* and its bacteriome-associated endosymbiont, *Wigglesworthia glossinidia*. *Journal of Molecular Evolution*, 48(1), 49–58.
- Childs, J. E., & Paddock, C. D. (2003). The ascendancy of *Amblyomma americanum* as a vector of pathogens affecting humans in the United States. *Annual Review of Entomology*, 48, 307–337. doi: 10.1146/annurev.ento.48.091801.112728
- Clay, K., Klyachko, O., Grindle, N., Civitello, D., Oleske, D., & Fuqua, C. (2008). Microbial communities and interactions in the lone star tick, *Amblyomma americanum*. *Molecular Ecology*, 17(19), 4371–4381.
- Duron, O., Binetruy, F., Noël, V., Cremaschi, J., McCoy, K. D., Arnathau, C., ... Chevillon, C. (2017). Evolutionary changes in symbiont community structure in ticks. *Molecular Ecology*, 26(11), 2905–2921. doi: 10.1111/mec.14094
- Duron, O., Jourdain, E., & McCoy, K. D. (2014). Diversity and global distribution of the *Coxiella* intracellular bacterium in seabird ticks. *Ticks and Tick-Borne Diseases*, 5(5), 557–563. doi: 10.1016/j.ttbdis.2014.04.003
- Duron, O., Morel, O., Noël, V., Buysse, M., Binetruy, F., Lancelot, R., ... Vial, L. (2018). Tick-Bacteria Mutualism Depends on B Vitamin Synthesis Pathways. *Current Biology*, 28(12), 1896–1902.e5. doi: 10.1016/j.cub.2018.04.038
- Duron, O., Noël, V., McCoy, K. D., Bonazzi, M., Sidi-Boumedine, K., Morel, O., ... Chevillon, C. (2015). The Recent Evolution of a Maternally-Inherited Endosymbiont of Ticks Led to the Emergence of the Q Fever Pathogen, *Coxiella burnetii*. *PLoS Pathogens*, 11(5), e1004892. doi: 10.1371/journal.ppat.1004892
- Duron, O., Wilkes, T. E., & Hurst, G. D. D. (2010). Interspecific transmission of a male-killing bacterium on an ecological timescale. *Ecology Letters*, 13(9), 1139–1148. doi: 10.1111/j.1461-0248.2010.01502.x

625 Engelstädter, J., & Hurst, G. D. D. (2009). The Ecology and Evolution of Microbes that Manipulate
 626 Host Reproduction. *Annual Review of Ecology, Evolution, and Systematics*, 40(1), 127–149.
 627 doi: 10.1146/annurev.ecolsys.110308.120206

628 Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., ... Pascal, G. (2018). FROGS:
 629 Find, Rapidly, OTUs with Galaxy Solution. *Bioinformatics (Oxford, England)*, 34(8), 1287–
 630 1294. doi: 10.1093/bioinformatics/btx791

631 Falony, G., Joossens, M., Vieira-Silva, S., Wang, J., Darzi, Y., Faust, K., ... Raes, J. (2016). Population-
 632 level analysis of gut microbiome variation. *Science (New York, N.Y.)*, 352(6285), 560–564. doi:
 633 10.1126/science.aad3503

634 Ferrari, J., & Vavre, F. (2011). Bacterial symbionts in insects or the story of communities affecting
 635 communities. *Philosophical Transactions of the Royal Society of London. Series B, Biological*
 636 *Sciences*, 366(1569), 1389–1400. doi: 10.1098/rstb.2010.0226

637 Galan, M., Razzauti, M., Bard, E., Bernard, M., Brouat, C., Charbonnel, N., ... Cosson, J.-F. (2016). 16S
 638 rRNA Amplicon Sequencing for Epidemiological Surveys of Bacteria in Wildlife. *MSystems*,
 639 1(4). doi: 10.1128/mSystems.00032-16

640 Gerhart, J. G., Moses, A. S., & Raghavan, R. (2016). A Francisella-like endosymbiont in the Gulf Coast
 641 tick evolved from a mammalian pathogen. *Scientific Reports*, 6, 33670. doi:
 642 10.1038/srep33670

643 Goecks, J., Nekrutenko, A., Taylor, J., & The Galaxy Team. (2010). Galaxy: a comprehensive approach
 644 for supporting accessible, reproducible, and transparent computational research in the life
 645 sciences. *Genome Biology*, 11(8), R86. doi: 10.1186/gb-2010-11-8-r86

646 Goslee, S. C., & Urban, D. L. (2007). The ecodist Package for Dissimilarity-based Analysis of Ecological
 647 Data. *Journal of Statistical Software*, 22(1), 1–19. doi: 10.18637/jss.v022.i07

648 Gottlieb, Y., Ghanim, M., Gueguen, G., Kontsedalov, S., Vavre, F., Fleury, F., & Zchori-Fein, E. (2008).
 649 Inherited intracellular ecosystem: symbiotic bacteria share bacteriocytes in whiteflies. *FASEB*

650 *Journal: Official Publication of the Federation of American Societies for Experimental Biology*,
651 22(7), 2591–2599. doi: 10.1096/fj.07-101162

652 Gottlieb, Y., Lalar, I., & Klasson, L. (2015). Distinctive Genome Reduction Rates Revealed by Genomic
653 Analyses of Two Coxiella-Like Endosymbionts in Ticks. *Genome Biology and Evolution*, 7(6),
654 1779–1796. doi: 10.1093/gbe/evv108

655 Gould, A. L., Zhang, V., Lamberti, L., Jones, E. W., Obadia, B., Korasidis, N., ... Ludington, W. B. (2018).
656 Microbiome interactions shape host fitness. *Proceedings of the National Academy of*
657 *Sciences*, 115(51), E11951–E11960. doi: 10.1073/pnas.1809349115

658 Groussin, M., Mazel, F., Sanders, J. G., Smillie, C. S., Lavergne, S., Thuiller, W., & Alm, E. J. (2017).
659 Unraveling the processes shaping mammalian gut microbiomes over evolutionary time.
660 *Nature Communications*, 8, 14319. doi: 10.1038/ncomms14319

661 Guglielmone, A. A., Estrada-Peña, A., Keirans, J. E., & Robbins, R. G. (2003). *Ticks (Acari: Ixodida) of*
662 *the Neotropical Zoogeographic Region*. Universiteit Utrecht, J. Bovy-Verbeek (public relations
663 ICTTD).

664 Guglielmone, A. A., & Nava, S. (2010). Hosts of Amblyomma dissimile Koch, 1844 and Amblyomma
665 rotundatum Koch, 1844 (Acari: Ixodidae). *Zootaxa*, 2541, 27–49.

666 Guizzo, M. G., Parizi, L. F., Nunes, R. D., Schama, R., Albano, R. M., Tirloni, L., ... Oliveira, P. L. (2017). A
667 Coxiella mutualist symbiont is essential to the development of Rhipicephalus microplus.
668 *Scientific Reports*, 7(1), 17554. doi: 10.1038/s41598-017-17309-x

669 Hanning, I., & Diaz-Sanchez, S. (2015). The functionality of the gastrointestinal microbiome in non-
670 human animals. *Microbiome*, 3. doi: 10.1186/s40168-015-0113-6

671 Holm, S. (1979). A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian Journal of*
672 *Statistics*, 6(2), 65–70. Retrieved from JSTOR.

673 Hosokawa, T., Koga, R., Kikuchi, Y., Meng, X.-Y., & Fukatsu, T. (2010). Wolbachia as a bacteriocyte-
674 associated nutritional mutualist. *Proceedings of the National Academy of Sciences of the*
675 *United States of America*, 107(2), 769–774. doi: 10.1073/pnas.0911476107

676 Hunter, D. J., Torkelson, J. L., Bodnar, J., Mortazavi, B., Laurent, T., Deason, J., ... Zhong, J. (2015). The
 677 Rickettsia Endosymbiont of Ixodes pacificus Contains All the Genes of De Novo Folate
 678 Biosynthesis. *PLoS ONE*, 10(12). doi: 10.1371/journal.pone.0144552
 679 Hutchinson, M. C., Cagua, E. F., Balbuena, J. A., Stouffer, D. B., & Poisot, T. (2017). paco:
 680 implementing Procrustean Approach to Cophylogeny in R. *Methods in Ecology and Evolution*,
 681 8(8), 932–940. doi: 10.1111/2041-210X.12736
 682 Jaenike, J. (2012). Population genetics of beneficial heritable symbionts. *Trends in Ecology &*
 683 *Evolution*, 27(4), 226–232. doi: 10.1016/j.tree.2011.10.005
 684 Jousselin, E., Cœur d’Acier, A., Vanlerberghe-Masutti, F., & Duron, O. (2013). Evolution and diversity
 685 of Arsenophonus endosymbionts in aphids. *Molecular Ecology*, 22(1), 260–270. doi:
 686 10.1111/mec.12092
 687 Jousselin, E., Desdevises, Y., & Coeur d’acier, A. (2009). Fine-scale cospeciation between
 688 Brachycaudus and Buchnera aphidicola: bacterial genome helps define species and
 689 evolutionary relationships in aphids. *Proceedings. Biological Sciences*, 276(1654), 187–196.
 690 doi: 10.1098/rspb.2008.0679
 691 Klyachko, O., Stein, B. D., Grindle, N., Clay, K., & Fuqua, C. (2007). Localization and visualization of a
 692 coxiella-type symbiont within the lone star tick, Amblyomma americanum. *Applied and*
 693 *Environmental Microbiology*, 73(20), 6584–6594. doi: 10.1128/AEM.00537-07
 694 Koga, R., & Moran, N. A. (2014). Swapping symbionts in spittlebugs: evolutionary replacement of a
 695 reduced genome symbiont. *The ISME Journal*, 8(6), 1237–1246. doi: 10.1038/ismej.2013.235
 696 Koga, R., Nikoh, N., Matsuura, Y., Meng, X.-Y., & Fukatsu, T. (2013). Mealybugs with distinct
 697 endosymbiotic systems living on the same host plant. *FEMS Microbiology Ecology*, 83(1), 93–
 698 100. doi: 10.1111/j.1574-6941.2012.01450.x
 699 Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis
 700 Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*, 33(7), 1870–1874. doi:
 701 10.1093/molbev/msw054

702 Kurtti, T. J., Felsheim, R. F., Burkhardt, N. Y., Oliver, J. D., Heu, C. C., & Munderloh, U. G. (2015).
703 *Rickettsia buchneri* sp. nov., a rickettsial endosymbiont of the blacklegged tick *Ixodes*
704 *scapularis*. *International Journal of Systematic and Evolutionary Microbiology*, 65(Pt 3), 965–
705 970. doi: 10.1099/ij.s.0.000047

706 Ley, R. E., Hamady, M., Lozupone, C., Turnbaugh, P., Ramey, R. R., Bircher, J. S., ... Gordon, J. I. (2008).
707 Evolution of mammals and their gut microbes. *Science (New York, N.Y.)*, 320(5883), 1647–
708 1651. doi: 10.1126/science.1155725

709 Łukasik, P., Nazario, K., Leuven, J. T. V., Campbell, M. A., Meyer, M., Michalik, A., ... McCutcheon, J. P.
710 (2018). Multiple origins of interdependent endosymbiotic complexes in a genus of cicadas.
711 *Proceedings of the National Academy of Sciences*, 115(2), E226–E235. doi:
712 10.1073/pnas.1712321115

713 Margulis, L. (1993). Origins of species: acquired genomes and individuality. *Biosystems*, 31(2), 121–
714 125. doi: 10.1016/0303-2647(93)90039-F

715 Martin, D. P., Lemey, P., Lott, M., Moulton, V., Posada, D., & Lefevre, P. (2010). RDP3: a flexible and
716 fast computer program for analyzing recombination. *Bioinformatics (Oxford, England)*,
717 26(19), 2462–2463. doi: 10.1093/bioinformatics/btq467

718 Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
719 *EMBnet.Journal*, 17(1), 10–12. doi: 10.14806/ej.17.1.200

720 Matsuura, Y., Kikuchi, Y., Hosokawa, T., Koga, R., Meng, X.-Y., Kamagata, Y., ... Fukatsu, T. (2012).
721 Evolution of symbiotic organs and endosymbionts in lygaeid stinkbugs. *The ISME Journal*,
722 6(2), 397–409. doi: 10.1038/ismej.2011.103

723 Matsuura, Y., Moriyama, M., Łukasik, P., Vanderpool, D., Tanahashi, M., Meng, X.-Y., ... Fukatsu, T.
724 (2018). Recurrent symbiont recruitment from fungal parasites in cicadas. *Proceedings of the*
725 *National Academy of Sciences*, 115(26), E5970–E5979. doi: 10.1073/pnas.1803245115

726 McCutcheon, J. P., Boyd, B. M., & Dale, C. (2019). The Life of an Insect Endosymbiont from the Cradle
727 to the Grave. *Current Biology: CB*, 29(11), R485–R495. doi: 10.1016/j.cub.2019.03.032

728 McCutcheon, J. P., & Moran, N. A. (2012). Extreme genome reduction in symbiotic bacteria. *Nature*
 729 *Reviews Microbiology*, 10(1), 13–26. doi: 10.1038/nrmicro2670
 730 Meibom, K. L., & Charbit, A. (2010). The unraveling panoply of *Francisella tularensis* virulence
 731 attributes. *Current Opinion in Microbiology*, 13(1), 11–17. doi: 10.1016/j.mib.2009.11.007
 732 Moran, N. A. (1996). Accelerated evolution and Muller's ratchet in endosymbiotic bacteria.
 733 *Proceedings of the National Academy of Sciences of the United States of America*, 93(7),
 734 2873–2878. doi: 10.1073/pnas.93.7.2873
 735 Moran, Nancy A., McCutcheon, J. P., & Nakabachi, A. (2008). Genomics and Evolution of Heritable
 736 Bacterial Symbionts. *Annual Review of Genetics*, 42(1), 165–190. doi:
 737 10.1146/annurev.genet.41.110306.130119
 738 Moran, Nancy A., Tran, P., & Gerardo, N. M. (2005). Symbiosis and insect diversification: an ancient
 739 symbiont of sap-feeding insects from the bacterial phylum Bacteroidetes. *Applied and*
 740 *Environmental Microbiology*, 71(12), 8802–8810. doi: 10.1128/AEM.71.12.8802-8810.2005
 741 Nakabachi, A., Yamashita, A., Toh, H., Ishikawa, H., Dunbar, H. E., Moran, N. A., & Hattori, M. (2006).
 742 The 160-kilobase genome of the bacterial endosymbiont *Carsonella*. *Science (New York, N.Y.)*,
 743 314(5797), 267. doi: 10.1126/science.1134196
 744 Nurk, S., Meleshko, D., Korobeynikov, A., & Pevzner, P. A. (2017). metaSPAdes: a new versatile
 745 metagenomic assembler. *Genome Research*, 27(5), 824–834. doi: 10.1101/gr.213959.116
 746 Oliver, K. M., Russell, J. A., Moran, N. A., & Hunter, M. S. (2003). Facultative bacterial symbionts in
 747 aphids confer resistance to parasitic wasps. *Proceedings of the National Academy of Sciences*
 748 *of the United States of America*, 100(4), 1803–1807. doi: 10.1073/pnas.0335320100
 749 Olivieri, E., Epis, S., Castelli, M., Varotto Boccazzi, I., Romeo, C., Desirò, A., ... Sassera, D. (2019).
 750 Tissue tropism and metabolic pathways of *Midichloria mitochondrii* suggest tissue-specific
 751 functions in the symbiosis with *Ixodes ricinus*. *Ticks and Tick-Borne Diseases*. doi:
 752 10.1016/j.ttbdis.2019.05.019

753 Price, M. N., Dehal, P. S., & Arkin, A. P. (2009). FastTree: computing large minimum evolution trees
 754 with profiles instead of a distance matrix. *Molecular Biology and Evolution*, 26(7), 1641–
 755 1650. doi: 10.1093/molbev/msp077

756 Ramaiah, A., & Dasch, G. A. (2018). Genome Sequence of Coxiella-Like Endosymbiont Strain CLE-
 757 RmD, a Bacterial Agent in the Cattle Tick, Rhipicephalus microplus Deutsch Strain. *Genome*
 758 *Announcements*, 6(13). doi: 10.1128/genomeA.00003-18

759 Rispe, C., & Moran, N. A. (2000). Accumulation of Deleterious Mutations in Endosymbionts: Muller's
 760 Ratchet with Two Levels of Selection. *The American Naturalist*, 156(4), 425–441. doi:
 761 10.1086/303396

762 Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D. L., Darling, A., Höhna, S., ... Huelsenbeck, J. P.
 763 (2012). MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and Model Choice Across a
 764 Large Model Space. *Systematic Biology*, 61(3), 539–542. doi: 10.1093/sysbio/sys029

765 Rowe, H. M., & Huntley, J. F. (2015). From the Outside-In: The Francisella tularensis Envelope and
 766 Virulence. *Frontiers in Cellular and Infection Microbiology*, 5, 94. doi:
 767 10.3389/fcimb.2015.00094

768 Russell, J. A., Goldman-Huertas, B., Moreau, C. S., Baldo, L., Stahlhut, J. K., Werren, J. H., & Pierce, N.
 769 E. (2009). Specialization and geographic isolation among Wolbachia symbionts from ants and
 770 lycaenid butterflies. *Evolution; International Journal of Organic Evolution*, 63(3), 624–640.
 771 doi: 10.1111/j.1558-5646.2008.00579.x

772 Sachs, J. L., Skophammer, R. G., & Regus, J. U. (2011). Evolutionary transitions in bacterial symbiosis.
 773 *Proceedings of the National Academy of Sciences*, 108(Supplement 2), 10800–10807. doi:
 774 10.1073/pnas.1100304108

775 Scarborough, C. L., Ferrari, J., & Godfray, H. C. J. (2005). Aphid protected from pathogen by
 776 endosymbiont. *Science (New York, N.Y.)*, 310(5755), 1781. doi: 10.1126/science.1120180

777 Scott, J. D., & Durden, L. A. (2015). *Amblyomma dissimile* Koch (Acari: Ixodidae) parasitizes bird
778 captured in Canada. *Systematic and Applied Acarology*, 20(8), 854–860. doi:
779 10.11158/saa.20.8.2

780 Shivaprasad, H. L., Cadenas, M. B., Diab, S. S., Nordhausen, R., Bradway, D., Crespo, R., &
781 Breitschwerdt, E. B. (2008). Coxiella-like infection in psittacines and a toucan. *Avian Diseases*,
782 52(3), 426–432. doi: 10.1637/8192-120707-Reg

783 Smith, T. A., Driscoll, T., Gillespie, J. J., & Raghavan, R. (2015). A Coxiella-Like Endosymbiont Is a
784 Potential Vitamin Source for the Lone Star Tick. *Genome Biology and Evolution*, 7(3), 831–
785 838. doi: 10.1093/gbe/evv016

786 Sudakaran, S., Kost, C., & Kaltenpoth, M. (2017). Symbiont Acquisition and Replacement as a Source
787 of Ecological Innovation. *Trends in Microbiology*, 25(5), 375–390. doi:
788 10.1016/j.tim.2017.02.014

789 Takiya, D. M., Tran, P. L., Dietrich, C. H., & Moran, N. A. (2006). Co-cladogenesis spanning three
790 phyla: leafhoppers (Insecta: Hemiptera: Cicadellidae) and their dual bacterial symbionts.
791 *Molecular Ecology*, 15(13), 4175–4191. doi: 10.1111/j.1365-294X.2006.03071.x

792 Theis, K. R., Dheilly, N. M., Klassen, J. L., Brucker, R. M., Baines, J. F., Bosch, T. C. G., ... Bordenstein, S.
793 R. (2016). Getting the Hologenome Concept Right: an Eco-Evolutionary Framework for Hosts
794 and Their Microbiomes. *MSystems*, 1(2), e00028-16. doi: 10.1128/mSystems.00028-16

795 Thompson, J. D., Gibson, T. J., & Higgins, D. G. (2002). Multiple sequence alignment using ClustalW
796 and ClustalX. *Current Protocols in Bioinformatics*, Chapter 2, Unit 2.3. doi:
797 10.1002/0471250953.bi0203s00

798 Turner, T. R., James, E. K., & Poole, P. S. (2013). The plant microbiome. *Genome Biology*, 14, 209. doi:
799 10.1186/gb-2013-14-6-209

800 van Schaik, E. J., Chen, C., Mertens, K., Weber, M. M., & Samuel, J. E. (2013). Molecular pathogenesis
801 of the obligate intracellular bacterium *Coxiella burnetii*. *Nature Reviews. Microbiology*, 11(8),
802 561–573. doi: 10.1038/nrmicro3049

803 Vapniarsky, N., Barr, B. C., & Murphy, B. (2012). Systemic Coxiella-like infection with myocarditis and
804 hepatitis in an eclectus parrot (*Eclectus roratus*). *Veterinary Pathology*, 49(4), 717–722. doi:
805 10.1177/0300985811409251

806 Vautrin, E., & Vavre, F. (2009). Interactions between vertically transmitted symbionts: cooperation or
807 conflict? *Trends in Microbiology*, 17(3), 95–99. doi: 10.1016/j.tim.2008.12.002

808 Voordouw, M. J. (2015). Co-feeding transmission in Lyme disease pathogens. *Parasitology*, 142(2),
809 290–302. doi: 10.1017/S0031182014001486

810 Wang, M., Zhu, D., Dai, J., Zhong, Z., Zhang, Y., & Wang, J. (2018). Tissue Localization and Variation of
811 Major Symbionts in *Haemaphysalis longicornis*, *Rhipicephalus haemaphysaloides*, and
812 *Dermacentor silvarum* in China. *Applied and Environmental Microbiology*, 84(10). doi:
813 10.1128/AEM.00029-18

814 Wernegreen, J. J. (2012). Endosymbiosis. *Current Biology: CB*, 22(14), R555–561. doi:
815 10.1016/j.cub.2012.06.010

816 Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis* (2nd ed.). Retrieved from
817 <https://www.springer.com/gp/book/9783319242750>

818 Woc-Colburn, A. M., Garner, M. M., Bradway, D., West, G., D’Agostino, J., Trupkiewicz, J., ...
819 Nordhausen, R. W. (2008). Fatal coxiellosis in Swainson’s Blue Mountain Rainbow Lorikeets
820 (*Trichoglossus haematodus moluccanus*). *Veterinary Pathology*, 45(2), 247–254. doi:
821 10.1354/vp.45-2-247

822 Wright, C. L., Sonenshine, D. E., Gaff, H. D., & Hynes, W. L. (2015). *Rickettsia parkeri* Transmission to
823 *Amblyomma americanum* by Cofeeding with *Amblyomma maculatum* (Acari: Ixodidae) and
824 Potential for Spillover. *Journal of Medical Entomology*, 52(5), 1090–1095. doi:
825 10.1093/jme/tjv086

826 Zhong, J., Jasinskas, A., & Barbour, A. G. (2007). Antibiotic treatment of the tick vector *Amblyomma*
827 *americanum* reduced reproductive fitness. *PloS One*, 2(5), e405. doi:
828 10.1371/journal.pone.0000405

829

830 **Acknowledgements**

831 Financial support was provided by the French government's Programmes Investissement
832 d'Avenir (Laboratoire d'Excellence CEBA, the MicroBIOMES Strategic Project 2016–2018
833 and the MiTick Annual Project 2016) and recurring funding from CNRS and IRD (OD, MB,
834 DA). FB benefits from a PhD fellowship financed by the CEBA Laboratoire d'Excellence and
835 Montpellier University. We wish to thank the platforms MGX (Montpellier Genomics and
836 Bioinformatics Facility) and GenSeq (Montpellier University), for the technical help and the
837 high-throughput sequencing. We are grateful to the members of Institut Pasteur de Guyane,
838 particularly Benoît de Thoisy and Agathe Chavy, for technical support. Special
839 acknowledgement to the “Agence Nationale de la Preservation de la Nature” (ANPN) and the
840 “Centre National de la Recherche Scientifique et Technologique of Gabon” (CENAREST)
841 that authorized and facilitated the access to the national parks in Gabon, and Marc F. Ngangué
842 (ANPN) that participated to the field collections in Gabon. We also thank Frédéric Delsuc,
843 Xavier Bailly, Laurence Mouton and Christine Chevillon for helpful discussions. The authors
844 acknowledge the IRD itrop HPC (South Green Platform) at IRD Montpellier for providing
845 HPC resources that have contributed to the research results reported in this paper. We also
846 acknowledge useful discussions with members of the Tiques et Maladies à Tiques (TMT)
847 working group from the Réseau Ecologique des Interactions Durables (REID).

848

849 **Conflict of interest**

850 The authors declare that they have no conflict of interest.

Figure legends

Figure 1 Phylogenetic relationship of the 25 *Amblyomma* species examined in this study. The phylogenetic tree was inferred using Bayesian inferences (BI) from a large fragment of rDNA sequences (18S, ITS1, 5.8S, ITS2, and 28S rDNA; 3312 bp unambiguously aligned; best-fit approximation for the evolutionary model: GTR+G+I), the numbers on nodes indicate the BI probability. One to two specimens per *Amblyomma* species (45 specimens in total) were analysed. The vertebrate species on which tick species mainly feed (host-range) is shown. White circles, African group (AF); black circles, New World group (NW).

Figure 2 Diversity of the microbiome and bacterial prevalence in *Amblyomma* species. The left part of the Figure shows the *Amblyomma* phylogenetic tree adapted from Figure 1. One to two specimens per *Amblyomma* species (45 specimens in total) were included in the phylogenetic analysis. In the middle part, the heatmap shows the abundance and diversity of the nine most abundant OTUs characterized in the 16S rRNA bacterial data set for each tick species. The number on the right of the heatmap indicates the number of tick specimens per species used to construct the heatmap (one to ten specimens per *Amblyomma* species (142 specimens in total) were included in the microbiome analysis). On the right of the figure, the prevalence of the three most common symbionts is illustrated by the coloured squares (*Coxiella*-LE, blue square; *Francisella*-LE, green square and *Rickettsia*, red square): full square indicate a prevalence of 100%, while empty or partly colored squares indicate prevalences from 0 to <100% (the numbers besides these squares indicate the prevalences). The amount of DNA for the single specimen of *A. sculptum* was not sufficient to perform the 16S rRNA barcoding, therefore the heatmap for this species is not applicable (N/A). In this latter species, the presence of *Coxiella*-LE and *Francisella*-LE was asserted through specific

PCR assays as detailed in the text. White circles, *Amblyomma* African group (AF); black circles, *Amblyomma* New World group (NW).

Figure 3 Structuration of the *Amblyomma* microbiome. **A** Multidimensional scaling plot (MDS) of the microbial diversity of each *Amblyomma* species depending on the presence of both *Coxiella*-LE (red dots) and *Francisella*-LE (green dots) or the absence of obligate symbiont (blue dots). **B** MDS plot of the microbial diversity depending on the presence (red dots) / absence (blue dots) of *Rickettsia*. **C** MDS plot of the microbial diversity depending on the association / no association between the three symbionts (*Coxiella*-LE, *Francisella*-LE and *Rickettsia*): C_R (red dots) correspond to *Amblyomma* specimens co-infected by *Coxiella*-LE and *Rickettsia*; C (brownish dots) to specimens only infected by *Coxiella*-LE; F (green dots) to specimens only infected by *Francisella*-LE; F_R to specimens co-infected by *Francisella*-LE and *Rickettsia*; C_F_R (light blue dots) to specimens multi-infected by *Coxiella*-LE, *Francisella*-LE and *Rickettsia*; dark blue dots to specimens not infected by these symbionts. Each MDS plot is based on the generalized Unifrac ($\alpha = 0.5$) distance matrix and all confidence ellipses are been drawn with the confidence level of 0.95.

Figure 4 Evolutionary histories between *Amblyomma* and their nutritional symbionts. **A** Co-phylogeny between the phylogenies of *Amblyomma* (on the left) and *Coxiella*-LE (on the right): The tick phylogeny is based on 3312-bp nuclear rDNA sequences (18S, ITS1, 5.8S, ITS2, and 28S rDNA; best-fit approximation for the evolutionary model: GTR+G+I); the *Coxiella*-LE phylogeny was reconstructed using a concatenated gene sequence (16S rRNA, *rpoB* and *groEL* concatenated sequences, 2045 unambiguously aligned base pairs; best-fit approximation for the evolutionary model: GTR+G+I). **B** Cophylogeny between the phylogenies of *Amblyomma* (on the left) and *Francisella*-LE (on the right): The tick

900 phylogeny is based on 3312-bp nuclear rDNA sequences (18S, ITS1, 5.8S, ITS2, and 28S
901 rDNA; best-fit approximation for the evolutionary model: GTR+G+I); the *Francisella*-LE
902 phylogeny was reconstructed using a concatenated gene sequence (16S rRNA, *rpoB*, *groEL*,
903 *ftsZ* and *gyrB* concatenated sequences, 3506 unambiguously aligned base pairs; best-fit
904 approximation for the evolutionary model: GTR+G+I). Each phylogeny was reconstructed
905 using BI and node numbers are the posterior probabilities (only values >70 are shown). White
906 circles, *Amblyomma* African group (AF); black circles, New World group (NW).