**Protein scaffold involving MSMEG_1285 maintain cell wall organization
and mediates penicillin sensitivity in mycobacteria**

Romain Veyron-Churlet^{1*}, Jean-Michel Saliou¹ and Camille Locht¹

¹ Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 8204 - CIIL - Center
for Infection and Immunity of Lille, F-59000 Lille, France

* Corresponding author: Romain Veyron-Churlet, CNRS-UMR8204, Institut Pasteur de Lille, 1 rue du
Pr Calmette, F-59021 Lille, France. Tel: +33 (0)3 20 87 11 55. Email: [romain.veyron-
churlet@ibl.cnrs.fr](mailto:romain.veyron-churlet@ibl.cnrs.fr)

Running title: PPI and penicillin sensitivity in mycobacteria

Keywords: *Mycobacterium*, tuberculosis, protein-protein interactions, antibiotic resistance, penicillin

Abbreviations: BACTH, Bacterial Adenylate Cyclase Two-Hybrid; CW, Cell Wall; HbhA, Heparin
Binding HemAgglutinin; LB, Luria Bertani; MB, Membrane; MBP, Maltose Binding Protein; *Msmeg*,
Mycobacterium smegmatis; *Mtb*, *Mycobacterium tuberculosis*; PBP, Penicillin-Binding Protein; PG,
peptidoglycan; PPI, protein-protein interactions; TB, tuberculosis; TBS, Tris-Buffered Saline.

Abstract

Protein-protein interactions are key in mycobacterial physiology, notably during the biosynthesis of the very peculiar mycobacterial cell wall. In this paper, we demonstrate that MSMEG_1285 interacts with PonA1, a bifunctional penicillin-binding protein involved in peptidoglycan biosynthesis. Deletion of *MSMEG_1285* enhances *Mycobacterium smegmatis* resistance to penicillin antibiotics, a phenotype that is exacerbated by the additional deletion of *hbhA*. This also led to a substantial decrease in the amounts of porins in the cell wall, which are necessary for the import of small and hydrophilic β -lactams. Deletion of both *MSMEG_1285* and *hbhA* provoked an over-representation of several enzymes involved in peptidoglycan degradation. Thus, we propose that MSMEG_1285 is part of a protein scaffold, which also involves PonA1 and HbhA, and that it is responsible for the tight regulation of peptidoglycan hydrolysis. This study provides a better understanding of the mycobacterial physiology, which is an essential step for strengthening the action of drugs that specifically target peptidoglycan biosynthesis.

Introduction

Mycobacterium tuberculosis (*Mtb*) is the main causative agent of tuberculosis (TB) in humans, which is the leading cause of death globally due to a single infectious agent. In 2017, TB killed an estimated 1.6 million people, according to the World Health Organization [1]. In addition, there is an alarming increase in multi-drug resistant TB cases (0.6 million cases in 2017). Hence, understanding the mechanisms leading to increased antimicrobial resistance is urgently needed to develop potent novel strategies against TB.

Deciphering the Protein-Protein Interactions (PPI) in pathogenic bacteria may help to understand the physiology of the cell and to elucidate host-pathogen interactions, in which proteins play crucial roles [2]. In addition, the study of PPI may facilitate the discovery of protein functions by the ‘guilty by association’ principle [3]. PPI are key factors in *Mtb* physiology and virulence [4]. For instance, interactions between the mycobacterial virulence factors ESAT-6 and CFP-10 are necessary for their

stability and secretion [5-7]. In addition, ESAT-6 alone or in complex with CFP-10 is able to interact with host proteins favoring dissemination of *Mtb* in the lungs [8], to traffic into the endoplasmic reticulum and to reduce antigen presentation by host cells [9]. PPI also play a central role in the biosynthesis of mycobacterial cell wall (CW) components. As the *Mtb* CW is very peculiar, they represent attractive targets for new, specific antitubercular drugs. For example, mycolic acid biosynthesis, which is targeted by the first-line drug isoniazid, relies on specialized and interconnected protein complexes [10-12]. Thus, deciphering the PPI network of *Mtb* should lead to a better understanding of TB pathogenesis and to the identification of new and critical steps required for infection.

We have shown that the *Mtb* Heparin-binding haemAgglutinin (HbhA_*Mtb*) interacts with Rv0613c and that MSMEG_1285, the Rv0613c orthologue in *Mycobacterium smegmatis* (*Msmeg*), is necessary for the proper cell-surface exposure of HbhA [13]. We wondered whether Rv0613c binds to other proteins in addition to HbhA and whether Rv0613c/MSMEG_1285 is involved in the proper localization of other mycobacterial proteins. To address this issue, we used here the Bacterial Adenylate Cyclase Two-Hybrid (BACTH) system to identify additional proteins that can interact with Rv0613c and discovered interactions between Rv0613c and the mycobacterial transglycosylase/D,D-transpeptidase PonA1. As *ponA1* was previously shown to play a role in *Msmeg* resistance to β -lactams antibiotics [14], we further demonstrate that deletion of *MSMEG_1285* increases the resistance of *Msmeg* to ampicillin. This phenotype was augmented by the additional deletion of *hbhA_Msmeg*. This increased resistance was directly linked to lower amounts of porins in the CW of both Δ *MSMEG_1285* and Δ *MSMEG_1285*+ Δ *hbhA_Msmeg* and to an over-representation of several enzymes involved in peptidoglycan (PG) hydrolysis. These data indicate that MSMEG_1285 plays an important role in the organization of the mycobacterial CW, which has a significant impact on the sensitivity to antibiotics, such as β -lactams.

Results and Discussion

Full-length Rv0613c interacts with PonA1

To identify additional interactants of Rv0613c, we performed a screening using the BACTH system [15] with pKT25_*rv0613c* as a bait and the *Mtb* Erdman library contained in pUT18C (Table S1). A positive clone was selected accordingly to its red phenotype on MacConkey agar plates and was further streaked onto fresh plates to confirm the phenotype. DNA was then extracted and the pUT18C insert was sequenced. This led to the identification of a portion of *ponA1* (*rv0050*), coding for amino acids 321 to 515. pUT18C_*ponA1*₃₂₁₋₅₁₅ was then introduced into *E. coli* DHM1 containing pKT25_*rv0613c* to confirm the phenotype seen in the BACTH system (Figure 1A). In parallel, the negative controls with pKT25 or pUT18C did not yield any red colony (Figure 1A). This phenotype was further confirmed using Luria Bertani (LB) agar plates supplemented with IPTG and X-gal by the blue colony phenotype in the strain harboring pUT18C_*ponA1*₃₂₁₋₅₁₅ together with pKT25_*rv0613c* (Figure 1B).

We have previously shown that HbhA interacts with a truncated version of Rv0613c, spanning residues 314 to 550 (Rv0613c₃₁₄₋₅₅₀) [13]. However, truncated Rv0613c₃₁₄₋₅₅₀ was not able to interact with PonA1₃₂₁₋₅₁₅ (data not shown), inferring that the N-terminal and/or C-terminal moieties of Rv0613c are/is necessary to maintain the interaction between Rv0613c and PonA1₃₂₁₋₅₁₅.

PonA1, also named Penicillin-Binding Protein 1 (PBP1), is a mycobacterial transglycosylase/D,D-transpeptidase, which is involved in PG biosynthesis [16]. The N-terminal region of PonA1 contains a transglycosylase domain (residues 37-218), whereas the C-terminal domain region contains a penicillin-binding transpeptidase domain (residues 310-600) [17], introducing a 4-3 crosslink in two parallel strands of the PG [18, 19]. The portion of PonA1 (PonA1₃₂₁₋₅₁₅) that interacts with Rv0613c is fully contained within the transpeptidase domain of the protein. The *Mtb* H37Rv genome codes for two bifunctional PBP (PonA1 and PonA2), whereas the *Msmeg* genome codes for three PBP (PonA1,

PonA2 and PonA3) [20]. PonA1 from *Mtb* displays 80% of sequence identity with PonA1 from *Msmeg*.

Rv0613c-⁶His is able to co-precipitate Maltose Binding Protein (MBP)-PonA1₃₂₁₋₅₁₅

E. coli BL21(DE3) was co-transformed with pET24a or pET24a_rv0613c, and pMAL-c2X or pMAL-c2X::ponA1₃₂₁₋₅₁₅. Total lysates were purified using Ni-NTA chromatography. As expected, MBP-PonA1₃₂₁₋₅₁₅ was not retained by the Ni-NTA column, whereas Rv0613c-⁶His (93 kDa) could be purified by Ni-NTA chromatography, in the presence of either co-expressed MBP or MBP-PonA1₃₂₁₋₅₁₅ (Figure 2, upper panel). Whereas for a similar amount of purified protein, Rv0613c-⁶His did not co-precipitate MBP (51 kDa), it did co-precipitate MBP-PonA1₃₂₁₋₅₁₅ (63 kDa) (Figure 2, upper panel), which was confirmed by mass spectrometry analysis of the corresponding band (data not shown).

The various fractions were further analyzed by western-blotting using an anti-MBP antibody. Confirming the data obtained Coomassie-blue staining, MBP-PonA1₃₂₁₋₅₁₅ did not bind to the Ni-NTA column, but co-expression with Rv0613c-⁶His resulted in binding to the Ni-NTA column (Figure 2, lower panel). These results thus indicate that MBP-PonA1₃₂₁₋₅₁₅ binds to Rv0613c, thereby confirming Rv0613c-PonA1₃₂₁₋₅₁₅ interaction identified in the BACTH system.

Deletion of *MSMEG_1285* and *hbhA_Msmeg* affects the resistance of *Msmeg* to ampicillin

It has been shown that a *Msmeg* ponA1 transposon mutant is more sensitive to β -lactam antibiotics than its parental strain, inferring that PonA1 is involved in intrinsic *Msmeg* resistance to β -lactams [14]. Considering the Rv0613c/PonA1 interaction, we wondered whether the deletion of the rv0613c orthologue in *Msmeg* (*MSMEG_1285*) would also influence the intrinsic resistance of *Msmeg* to β -lactams, such as ampicillin. Overnight cultures (i.e. at the end of exponential growth phase) of *Msmeg* mc²155 and the deletion mutant Δ *MSMEG_1285* (Δ *MS1285*) were therefore spread onto 7H11 agar plates containing ampicillin (ranging from 10 to 500 μ g/ml). Surprisingly, the absence of *MSMEG_1285* increased the resistance of *Msmeg* to ampicillin, as Δ *MS1285* was able to grow on

concentrations above 250 µg/ml (Figure 3A), in contrast to *ponA1* transposon mutant [14]. As a control, we used rifampicin (ranging from 0.1 to 10 µg/ml), which does not target PG crosslinking, and found that the sensitivity to rifampicin was not altered by the deletion of *MSMEG_1285* (Figure 3A).

As Rv0613c interacts with both PonA1 and HbhA [13], we wondered whether the deletion of *hbhA_Msmeg* (*MSMEG_0919*) would also affect the sensitivity to ampicillin. We thus generated a *hbhA_Msmeg* deletion mutant ($\Delta MS0919$), as well as a double deletion mutant $\Delta MS1285 + \Delta MS0919$ (Figure 4A and 4B). The single deletion of *MS0919* did not affect the sensitivity to ampicillin, nor to rifampicin (Figure 3A). However, the double deletion further increased ampicillin resistance as $\Delta MS1285 + \Delta MS0919$ was able to grow at concentrations above 500 µg/ml, whereas sensitivity for rifampicin was not influenced (Figure 3A). Plating of serial dilutions at an intermediate ampicillin concentration (50 µg/ml) confirmed that $\Delta MS1285 + \Delta MS0919$ is more resistant to ampicillin than $\Delta MS1285$, which, in turn, is more resistant than *Msmeg* mc²155 or $\Delta MS0919$, whereas the sensitivity to rifampicin was not affected (Figure 3B).

Proteomics analysis of the *Msmeg* deletion mutants

Four *Msmeg* porins (MspA to MspD) have been shown to be involved in antibiotic resistance by directly transporting small β-lactam antibiotics, such as ampicillin [21]. To determine whether the deletion of *MS1285* and/or *MS0919* impacts on the presence of these porins in the bacterial membrane (MB) or CW, we performed proteomic analyses on MB and CW preparations from *Msmeg* mc²155, $\Delta MS0919$, $\Delta MS1285$ and $\Delta MS1285 + \Delta MS0919$. We used a spectral counting approach [22] and defined a ratio for each protein as the total number of spectra for each mutant divided by the total number of spectra for *Msmeg* mc²155 (for convenience, the absence of any spectrum was replaced by 0.5, when needed). Only ratios ≥ 5 were considered as significant. Controls confirmed that neither *HbhA_Msmeg* nor *MSMEG_1285* were detected in the corresponding deletion mutants (Table 1). In addition, the total amounts (CW plus MB fractions) of PonA1 and PonA2 were barely affected,

inferring that any difference in the ampicillin sensitivity was not linked to the total amount of D,D-transpeptidases (Table 1), whereas PonA3 (MSMEG_4388) was not detected in any sample tested (data not shown). Although the difference in the distribution of PonA1 in the CW and MB fractions between deletion mutants and wild-type strain seems to be rather modest, an impact of this difference on CW properties can nevertheless not be excluded (Table 1).

As expected, peptides corresponding to MspA, B, C, and D (LIASTGDSVTTYGEPWNmN and TLTVQQWDTFLNGVFPLDR) were only detected in the CW fraction, confirming the efficacy of the cell fractionation (Table 1). As the first peptide is common to the four porins and the second one is common to MspA, MspB and MspC, it was not possible to distinguish between the four Msp-like porins. The absence of HbhA_ *Msmeg* slightly decreased the abundance of the porins by two fold, whereas the absence of MSMEG_1285 (with or without HbhA_ *Msmeg*) reduced their abundance by at least eight fold (Table 1). Thus, the higher ampicillin resistance of $\Delta MS1285$ and $\Delta MS1285 + \Delta MS0919$ mutants compared to *Msmeg* mc²155 is paralleled by a decreased level of porins in the mycobacterial CW, thereby most likely limiting the import of small β -lactams like ampicillin.

The deletion of *MS0919* together with that of *MS1285* did not further decrease the amount of porins in the CW, despite the fact that the double mutant is more resistant to ampicillin than either strain containing a single deletion (Figure 3). Therefore, we further analyzed the proteomic content of CW plus MB fractions from $\Delta MS1285 + \Delta MS0919$ and compared them to those of *Msmeg* mc²155. Among 31 over-represented proteins (ratio ≥ 5) in the double mutant (Table S3), the most strongly over-represented one was MSMEG_3145 (Table 1), the orthologue of the PG endopeptidase RipA in *Mtb* [23]. RipA_ *Msmeg* was also slightly over-represented in the single deletion mutant $\Delta MS0919$ but not in $\Delta MS1285$. At least two other proteins involved in PG hydrolysis were detected with a ratio ≥ 5 : MSMEG_6815, which displays similarities with phage lysins and with the *Mtb* PG hydrolase Rv2525c [24], and MSMEG_6281, known as the PG amidase Ami1_ *Msmeg* [25] (Table 1). Interestingly, a transcriptomics study showed that *ripA_Mtb* (*rv1477*), *rv2525c* and *ami1_Mtb* (*rv3717*) are co-

regulated, suggesting a controlled and complementary role in the PG hydrolysis for the three corresponding enzymes in *Mtb* [26].

Overexpression of *ripA_Mtb* or *ami1* increases resistance of *Msmeg* to ampicillin

As deletion of *ripA* increases susceptibility to β -lactams like carbenicillin [27], we investigated whether over-representation of RipA in $\Delta MS1285+\Delta MS0919$ could account for the increased resistance to ampicillin. Whereas constitutive overexpression of *ripA_Msmeg* using the expression vector pVV16 was detrimental to the growth of *Msmeg* and yielded no colony (data not shown), as previously demonstrated with an inducible system [28], overexpression of *ripA_Mtb* was well tolerated in *Msmeg*. Using this recombinant strain of *Msmeg*, we found that overexpression of *ripA_Mtb* in *Msmeg* mc²155 increased resistance to ampicillin by one order of magnitude at 500 μ g/ml (Figure 5).

The deletion of *ami1_Msmeg* was also reported to increase *Msmeg* susceptibility to ampicillin by 2 fold [25]. Thus, we wondered whether over-representation of either *ami1_Msmeg* or *ami1_Mtb* could account for the increased ampicillin resistance observed in $\Delta MS1285+\Delta MS0919$. We found that overexpression of *ami1_Msmeg* and *ami1_Mtb* also increased the resistance to ampicillin by two and three orders of magnitude, respectively, at 500 μ g/ml (Figure 5). As MSMEG_6815 was overrepresented in both $\Delta MS0919$ and $\Delta MS1285+\Delta MS0919$ (Table 1), this protein is most likely not responsible for the increased resistance to ampicillin seen only with the double mutant.

The basal resistance observed in both $\Delta MS1285$ and $\Delta MS1285+\Delta MS0919$ is due to the reduced presence of porins in the CW, whereas further increased resistance to ampicillin in the case of $\Delta MS1285+\Delta MS0919$ is linked to the over-representation of both *RipA_Msmeg* and *Ami1_Msmeg* (Figure 3 and Table 1).

Effect of $\Delta MS1285+\Delta MS0919$ complementation on penicillin and antitubercular drug resistance

The double mutant was transformed with pMV361, pMV361_MS0919, pMV361_MS1285 and pMV361_MS1285_MS0919. As expected, pMV361 and pMV361_MS0919 did not modify the increased resistance to ampicillin (50 µg/ml), in contrast to pMV361_MS1285, which restored the sensitivity to a level similar to that of *Msmeg* mc²155 or Δ MS0919 (Figure 6). Surprisingly, simultaneous expression of MS1285 and MS0919 did not restore sensitivity to ampicillin. This was not due to a lack of expression of MS0919 (located downstream of MS1285 in the pMV361 construct), as HbhA_*Msmeg* was detected by western-blot analysis (Figure 4C). Instead, it is likely that the low amount of HbhA_*Msmeg* in Δ MS1285+ Δ MS0919 containing pMV361_MS1285_MS0919 was not enough to fully restore the phenotype, in comparison with the level of endogenous HbhA_*Msmeg* in *Msmeg* mc²155 (Figure 4C).

We also tested carbenicillin (1,000 µg/ml) as another antibiotic belonging to the penicillin family. Similar to ampicillin, Δ MS1285+ Δ MS0919 was one log more resistant to carbenicillin than *Msmeg* mc²155 or Δ MS0919 (Figure 6). Complementation by pMV361_MS1285 was enough to restore sensitivity to carbenicillin (Figure 6). Thus, deletion of both MS1285 and MS0919 increases resistance to several members of the penicillin family like ampicillin and carbenicillin.

To examine whether this phenotype was limited to antibiotics targeting the PG biosynthesis, we also tested the antitubercular drugs isoniazid (5 µg/ml) and ethionamide (5 µg/ml). However, Δ MS1285+ Δ MS0919 showed no difference in sensitivity to these drugs at the tested concentrations, as compared to *Msmeg* mc²155 (Figure 6).

Conclusion

In the present study, we show that Rv0613c interacts with a region of PonA1 fully contained within its transpeptidase domain. We have previously uncovered interactions between HbhA and Rv0613c, as well as between HbhA and a domain within the second extracellular loop of MmpL14 [13]. Another study demonstrated that a C-terminal portion of PonA1, different from the portion interacting with

Rv0613c, interacts with RipA [29]. Based on this PPI network, we hypothesize that Rv0613c/MSMEG_1285 may be a scaffold protein responsible for the formation of protein complexes involved in CW organization and PG biosynthesis (Figure 7). Thus, deletion of *rv0613c* or *MSMEG_1285* result in (i) an improper localization of both PonA1 and HbhA, as previously shown for HbhA by atomic force microscopy [13], (ii) an accumulation of RipA, MSMEG_6815 and Ami1, when *hbhA* is additionally deleted, and (iii) a decrease in porin levels and the subsequent increased resistance to β -lactams in *Msmeg*. This hypothesis is consistent with the CW localization of MSMEG_1285 (and porins) as evidenced by our proteomic analyses of the various *Msmeg* strains (Table 1) and by another independent proteomics study [30]. Interestingly, it was recently shown that PPI between HtrA and LppZ are responsible for the control of Ami3_*Msmeg* toxicity, inferring a tight spatiotemporal coordination of PG expansion and division [31]. Moreover, deciphering the MmpL3 interactome led to the identification of functional links between the MmpL protein family and the assembling and export of PG [32]. Here, we hypothesize that Rv0613c/MSMEG_1285, interacting with PonA1 and HbhA, might be part of a protein scaffold that is necessary for the proper localization of PG hydrolyzing enzymes (RipA, MSMEG_6815 and Ami1) (Figure 7), as PG degradation is a process that is tightly regulated in mycobacteria [16]. The characterization of such protein complexes at the CW level will facilitate the understanding of *Mtb* physiology and may pave the way for strengthening the action of drugs targeting mycobacterial PG biosynthesis, like carbapenems [33, 34].

Methods

Bacterial strains and media

E. coli TOP10 (Invitrogen, Carlsbad, CA, USA) was used for cloning, *E. coli* DHM1 was used for BACTH assays and *E. coli* BL21(DE3) (Agilent, Santa Clara, CA, USA) was used for purification of recombinant proteins (Table S1). The *E. coli* strains were grown in LB medium (MP Biomedicals, Illkirch, France) at 37 °C supplemented with ampicillin (100 μ g/ml), apramycin (25 μ g/ml) and/or kanamycin (25 μ g/ml), when required. *Msmeg* mc²155, the deletion mutants Δ MS0919, Δ MS1285, Δ MS1285+ Δ MS0919 and the corresponding complemented strains (Table S1) were grown in

262 Middlebrook 7H9 broth or on 7H11 agar plates supplemented with OADC enrichment (Becton
263 Dickinson, Le Pont-de-Claix, France), apramycin (25 µg/ml), hygromycin (50 µg/mL) and/or
264 kanamycin (25 µg/mL), when required. Growth curves in Middlebrook 7H9 supplemented with
265 OADC did not manifest any growth difference between wild-type, simple and/or double mutants (data
266 not shown).

267

268 **Plasmid constructions**

269 DNA encoding full-length Rv0613c and truncated Rv0613c₃₁₄₋₅₅₀ were amplified by PCR using *Mtb*
270 H37Rv genomic DNA and the pairs of primers [pKT25_rv0613c_dir + pKT25_rv0613c_rev] and
271 [pKT25_rv0613c₃₁₄₋₅₅₀_dir + pKT25_rv0613c₃₁₄₋₅₅₀_rev] (Table S2). The PCR fragments were
272 digested with PstI and KpnI and ligated into pKT25 digested with the same enzymes, generating
273 pKT25_rv0613c and pKT25_rv0613c₃₁₄₋₅₅₀ (Table S1). The *rv0613c* gene was also amplified by PCR
274 using the pair of primers [pET24a_rv0613c_dir + pET24a_rv0613c_rev] (Table S2). The PCR
275 fragment was digested with NdeI and HindIII and ligated into pET24a digested with the same
276 enzymes, generating pET24a_rv0613c (Table S1). DNA encoding the portion 321-515 of PonA1 was
277 amplified by PCR using *Mtb* H37Rv genomic DNA and the pair of primers [pMAL-c2X_ponA1₃₂₁₋
278 ₅₁₅_dir + pMAL-c2X_ponA1₃₂₁₋₅₁₅_rev] (Table S2). The PCR fragment was digested with EcoRI and
279 HindIII and inserted into pMAL-c2X digested with the same enzymes, generating pMAL-c2X
280 _ponA1₃₂₁₋₅₁₅ (Table S1). The genes *ripA* and *amiI* were amplified by PCR using *Msmeg* mc²155 or
281 *Mtb* H37Rv genomic DNA and the pairs of primers [pVV16_ripA_Msmeg_dir +
282 pVV16_ripA_Msmeg_rev], [pVV16_ripA_Mtb_dir + pVV16_ripA_Mtb_rev],
283 [pVV16_amiI_Msmeg_dir + pVV16_amiI_Msmeg_rev] and [pVV16_amiI_Mtb_dir +
284 pVV16_amiI_Mtb_rev] (Table S2). The PCR fragments were digested with NdeI and HindIII and
285 ligated into pVV16 digested with the same enzymes, generating pVV16_ripA_Msmeg,
286 pVV16_ripA_Mtb, pVV16_amiI_Msmeg and pVV16_amiI_Mtb, respectively (Table S1). The
287 *hbhA_Msmeg* (*MSMEG_0919*) gene was amplified by PCR using *Msmeg* mc²155 genomic DNA and
288 the pair of primers [pMV361_MS0919_dir + pMV361_MS0919_rev] (Table S2). The PCR fragment
289 was digested with ClaI and HpaI and ligated into pMV361 digested with the same enzymes,

generating pMV361_MS0919 (Table S1). The *MSMEG_1285* gene was amplified by PCR using *Msmeg* mc²155 genomic DNA and the pair of primers [pMV361_MS1285_dir + pMV361_MS1285_rev] (Table S2). The PCR fragment was digested with MunI and HindI and ligated into pMV361 or pMV361_MS0919 digested with the same enzymes, generating pMV361_MS1285 or pMV361_MS1285_MS0919, respectively (Table S1). All plasmids were checked by sequencing.

BACTH assays

One hundred ng of the *Mtb* Erdman library contained in pUT18C (Table S1) was electroporated into *E. coli* DHM1 containing pKT25_rv0613c. Transformants were plated onto MacConkey agar (Becton Dickinson) supplemented with ampicillin (100 µg/ml), kanamycin (25 µg/ml), IPTG (0.5 mM) and maltose 1%. After 6 days at 30 °C, a clone was selected based on its red color and transferred onto fresh MacConkey agar plates to confirm the red phenotype. DNA was extracted from the selected colony and pUT18C insert was sequenced using the Seq_pUT18C_dir primer (Table S2). To confirm the phenotype, pUT18C containing *ponA1*₃₂₁₋₅₁₅ was re-introduced into *E. coli* DHM1 containing either pKT25_rv0613c or pKT25_rv0613c₃₁₄₋₅₅₀. Transformants were plated onto both MacConkey agar or LB agar supplemented with ampicillin (100 µg/ml), kanamycin (25 µg/ml), IPTG (0.5 mM) and X-gal (50 µg/ml) and incubated for 3 days at 30 °C.

Co-precipitation assays

E. coli BL21(DE3) was co-transformed with pET24a_rv0613c and pMAL-c2X_ponA1₃₂₁₋₅₁₅. As a control, *E. coli* BL21(DE3) was co-transformed with pMAL-c2X_ponA1₃₂₁₋₅₁₅ together with pET24a, or with pET24a_rv0613c together with pMAL-c2X. *E. coli* cells were grown until OD₆₀₀ = 0.5 and induced with 1 mM IPTG at 37°C for 3 hrs. Cells were disrupted using a French pressure cell in purification buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 10% glycerol, 1 mM DTT, 1 mM EDTA, 10 mM imidazole). Each lysate was subjected to HisPur Ni-NTA chromatography (ThermoFisher, Waltham, MA, USA) following manufacturer's protocol. Recombinant (and co-precipitating) proteins were eluted in 1 ml of purification buffer containing 200 mM imidazole. Eluted proteins were then analyzed on SDS-PAGE and Coomassie-blue staining.

318

319 **Immunoblot analysis**

320 Total proteins were resolved by SDS-PAGE using 12% acrylamide gels and then electro-transferred
321 onto a nitrocellulose membrane. The membrane was saturated with Tris-Buffered Saline (TBS) pH 7.5
322 containing 0.05% Tween 80 and 5% milk and probed overnight at 4 °C with anti-HbhA monoclonal
323 antibody VF2 [35], anti-Hsp65/GroL1 monoclonal antibody (Abcam, Paris, France, dilution 1:1,000),
324 or anti-MBP (NEB Biolabs, Evry, France, dilution 1:10,000) in TBS-Tween-3% milk. Finally, the
325 membrane was incubated for 1 hr at room temperature with goat anti-mouse horseradish peroxidase-
326 conjugated secondary antibodies (Abcam, diluted 1:5,000) in TBS-Tween-3% milk. The blots were
327 developed using the Amersham ECL Prime Western-Blotting Detection Reagent (GE Healthcare,
328 Velizy-Villacoublay, France) and chemiluminescence was detected using the Amersham Imager 600
329 (GE Healthcare).

330

331 **Generation of deletion mutants in *Msmeg***

332 To delete *hbhA_Msmeg* (*MSMEG_0919*) in *Msmeg* mc²155 and in the Δ *MSI285* mutant, we followed
333 the protocol previously described by van Kessel *et al.* [36]. Briefly, upstream and downstream regions
334 of *hbhA_Msmeg* were amplified from *Msmeg* mc²155 genomic DNA using the pairs of primers
335 [Up_*hbhA_Msmeg_dir* + Up_*hbhA_Msmeg_rev*] and [Down_*hbhA_Msmeg_dir* +
336 Down_*hbhA_Msmeg_rev*] (Table S2). These upstream and downstream fragments were inserted at
337 either side of an apramycin resistance cassette in pCR-BluntII-TOPO (ThermoFisher) (Table S1).
338 After digestion by XbaI and HindIII, the allelic exchange substrate was electroporated into competent
339 *Msmeg* mc²155 and the Δ *MSI285* mutant containing pJV53 (Table S1), following induction by 0.2%
340 acetamide for 3 hrs. Several apramycin-resistant clones were selected and tested for the insertion of
341 the apramycin resistance cassette by PCR using the pairs of primers [Up_*hbhA_Msmeg_dir* +
342 Down_*hbhA_Msmeg_rev*], [Up_*hbhA_Msmeg_dir* + Seq_*apramycin_rev*] and [Seq_*apramycin_dir* +
343 Down_*hbhA_Msmeg_rev*] (Table S2).

344

345 **Drug susceptibility testing**

Susceptibility of *Msmeg*, the deletion mutants and the complemented strains to several antibiotics (rifampicin, ampicillin, carbenicillin, isoniazid, ethionamide) was determined on Middlebrook 7H11 solid medium containing OADC enrichment with various drug concentrations. Overnight cultures of *Msmeg* were all diluted at $OD_{600} = 0.5$ and 5 μ L of this dilution or 5 μ L of ten-fold serial dilutions (10^{-1} to 10^{-5}) of each culture were plated and incubated at 37°C for 3 days.

Fractionation protocol

Msmeg and mutant strains were disrupted in a French pressure cell and the lysates were centrifuged at $3,000 \times g$ for 10 min at 4 °C to eliminate cellular debris and non-lysed cells. The supernatants were pelleted at $27,000 \times g$ for 30 min at 4 °C in order to collect CW fractions. The remaining supernatants were centrifuged at $100,000 \times g$ for 2 hrs at 4 °C to separate the membrane MB fractions from the cytosolic fractions. Each fraction was resuspended in an appropriate volume of denaturing buffer (0.1% β -mercaptoethanol, 0.0005% bromophenol blue, 10% glycerol, 2% SDS, 63 mM Tris-HCl, pH 6.8).

Mass spectrometry proteomic analysis

Mass spectrometry proteomic analysis was performed as described [13].

Proteomic Data Analysis

Proteomic data analysis was performed as described [13], using a *Msmeg* UniProt database (strain ATCC 700084/mc (2)155, taxo 246196, October 2014, 8805 entries) fused with recombinant trypsin and a list of classical contaminants (119 entries).

Author Contributions

RVC conceived and designed the research. RVC and JMS performed the experiments. RVC and JMS analyzed the data. RVC and CL wrote the manuscript. All authors have read and approved the final version of the manuscript. The authors declare they have no competing interests.

373

374 **Acknowledgments**

375 We thank Dr Alain Baulard (Institut Pasteur de Lille, France) for kindly providing the *Mtb* Erdman
376 library in pUT18C vector.

377

378 **Conflict of interest**

379 The authors declare no conflict of interest.

380

381 **References**

- 382 1. World Health Organization (2018) Global Tuberculosis Report.
- 383 2. Velasco-Garcia, R. & Vargas-Martinez, R. (2012) The study of protein-protein interactions in
384 bacteria, *Can J Microbiol.* **58**, 1241-57.
- 385 3. Schauer, K. & Stingl, K. (2009) 'Guilty by Association' - Protein-Protein Interactions (PPIs) in
386 Bacterial Pathogens, *Genome Dyn.* **6**, 48-61.
- 387 4. Veyron-Churlet, R. & Loch, C. (2019) In Vivo Methods to Study Protein-Protein Interactions as
388 Key Players in *Mycobacterium Tuberculosis* Virulence, *Pathogens.* **8**.
- 389 5. Renshaw, P. S., Panagiotidou, P., Whelan, A., Gordon, S. V., Hewinson, R. G., Williamson, R. A.
390 & Carr, M. D. (2002) Conclusive evidence that the major T-cell antigens of the *Mycobacterium*
391 *tuberculosis* complex ESAT-6 and CFP-10 form a tight, 1:1 complex and characterization of the
392 structural properties of ESAT-6, CFP-10, and the ESAT-6*CFP-10 complex. Implications for
393 pathogenesis and virulence, *J Biol Chem.* **277**, 21598-603.
- 394 6. Renshaw, P. S., Lightbody, K. L., Veverka, V., Muskett, F. W., Kelly, G., Frenkiel, T. A., Gordon,
395 S. V., Hewinson, R. G., Burke, B., Norman, J., Williamson, R. A. & Carr, M. D. (2005) Structure and
396 function of the complex formed by the tuberculosis virulence factors CFP-10 and ESAT-6, *EMBO J.*
397 **24**, 2491-8.
- 398 7. Brodin, P., de Jonge, M. I., Majlessi, L., Leclerc, C., Nilges, M., Cole, S. T. & Brosch, R. (2005)
399 Functional analysis of early secreted antigenic target-6, the dominant T-cell antigen of *Mycobacterium*

400 *tuberculosis*, reveals key residues involved in secretion, complex formation, virulence, and
 401 immunogenicity, *J Biol Chem.* **280**, 33953-9.

402 8. Kinhikar, A. G., Verma, I., Chandra, D., Singh, K. K., Weldingh, K., Andersen, P., Hsu, T., Jacobs,
 403 W. R., Jr. & Laal, S. (2010) Potential role for ESAT6 in dissemination of *M. tuberculosis* via human
 404 lung epithelial cells, *Mol Microbiol.* **75**, 92-106.

405 9. Sreejit, G., Ahmed, A., Parveen, N., Jha, V., Valluri, V. L., Ghosh, S. & Mukhopadhyay, S. (2014)
 406 The ESAT-6 protein of *Mycobacterium tuberculosis* interacts with beta-2-microglobulin (beta2M)
 407 affecting antigen presentation function of macrophage, *PLoS Pathog.* **10**, e1004446.

408 10. Veyron-Churlet, R., Guerrini, O., Mourey, L., Daffe, M. & Zerbib, D. (2004) Protein-protein
 409 interactions within the Fatty Acid Synthase-II system of *Mycobacterium tuberculosis* are essential for
 410 mycobacterial viability, *Mol Microbiol.* **54**, 1161-72.

411 11. Veyron-Churlet, R., Bigot, S., Guerrini, O., Verdoux, S., Malaga, W., Daffe, M. & Zerbib, D.
 412 (2005) The biosynthesis of mycolic acids in *Mycobacterium tuberculosis* relies on multiple specialized
 413 elongation complexes interconnected by specific protein-protein interactions, *J Mol Biol.* **353**, 847-58.

414 12. Cantaloube, S., Veyron-Churlet, R., Haddache, N., Daffe, M. & Zerbib, D. (2011) The
 415 *Mycobacterium tuberculosis* FAS-II dehydratases and methyltransferases define the specificity of the
 416 mycolic acid elongation complexes, *PLoS One.* **6**, e29564.

417 13. Veyron-Churlet, R., Dupres, V., Saliou, J. M., Lafont, F., Raze, D. & Locht, C. (2018)
 418 Rv0613c/MSMEG_1285 Interacts with HBHA and Mediates Its Proper Cell-Surface Exposure in
 419 Mycobacteria, *Int J Mol Sci.* **19**.

420 14. Billman-Jacobe, H., Haite, R. E. & Coppel, R. L. (1999) Characterization of a *Mycobacterium*
 421 *smegmatis* mutant lacking penicillin binding protein 1, *Antimicrob Agents Chemother.* **43**, 3011-3.

422 15. Karimova, G., Pidoux, J., Ullmann, A. & Ladant, D. (1998) A bacterial two-hybrid system based
 423 on a reconstituted signal transduction pathway, *Proc Natl Acad Sci U S A.* **95**, 5752-6.

424 16. Kieser, K. J. & Rubin, E. J. (2014) How sisters grow apart: mycobacterial growth and division,
 425 *Nat Rev Microbiol.* **12**, 550-62.

426 17. Filippova, E. V., Kieser, K. J., Luan, C. H., Wawrzak, Z., Kiryukhina, O., Rubin, E. J. &
 427 Anderson, W. F. (2016) Crystal structures of the transpeptidase domain of the *Mycobacterium*

428 *tuberculosis* penicillin-binding protein PonA1 reveal potential mechanisms of antibiotic resistance,
 429 *FEBS J.* **283**, 2206-18.

430 18. Sauvage, E., Kerff, F., Terrak, M., Ayala, J. A. & Charlier, P. (2008) The penicillin-binding
 431 proteins: structure and role in peptidoglycan biosynthesis, *FEMS Microbiol Rev.* **32**, 234-58.

432 19. Maitra, A., Munshi, T., Healy, J., Martin, L. T., Vollmer, W., Keep, N. H. & Bhakta, S. (2019)
 433 Cell wall peptidoglycan in *Mycobacterium tuberculosis*: An Achilles' heel for the TB-causing
 434 pathogen, *FEMS Microbiol Rev.* **43**, 548-575.

435 20. Patru, M. M. & Pavelka, M. S., Jr. (2010) A role for the class A penicillin-binding protein PonA2
 436 in the survival of *Mycobacterium smegmatis* under conditions of nonreplication, *J Bacteriol.* **192**,
 437 3043-54.

438 21. Danilchanka, O., Pavlenok, M. & Niederweis, M. (2008) Role of porins for uptake of antibiotics
 439 by *Mycobacterium smegmatis*, *Antimicrob Agents Chemother.* **52**, 3127-34.

440 22. Nahnsen, S., Bielow, C., Reinert, K. & Kohlbacher, O. (2013) Tools for label-free peptide
 441 quantification, *Mol Cell Proteomics.* **12**, 549-56.

442 23. Hett, E. C., Chao, M. C., Steyn, A. J., Fortune, S. M., Deng, L. L. & Rubin, E. J. (2007) A partner
 443 for the resuscitation-promoting factors of *Mycobacterium tuberculosis*, *Mol Microbiol.* **66**, 658-68.

444 24. Bellinzoni, M., Haouz, A., Miras, I., Magnet, S., Andre-Leroux, G., Mukherjee, R., Shepard, W.,
 445 Cole, S. T. & Alzari, P. M. (2014) Structural studies suggest a peptidoglycan hydrolase function for
 446 the *Mycobacterium tuberculosis* Tat-secreted protein Rv2525c, *J Struct Biol.* **188**, 156-64.

447 25. Senzani, S., Li, D., Bhaskar, A., Ealand, C., Chang, J., Rimal, B., Liu, C., Joon Kim, S., Dhar, N.
 448 & Kana, B. (2017) An Amidase_3 domain-containing *N*-acetylmuramyl-L-alanine amidase is required
 449 for mycobacterial cell division, *Sci Rep.* **7**, 1140.

450 26. Saint-Joanis, B., Demangel, C., Jackson, M., Brodin, P., Marsollier, L., Boshoff, H. & Cole, S. T.
 451 (2006) Inactivation of Rv2525c, a substrate of the twin arginine translocation (Tat) system of
 452 *Mycobacterium tuberculosis*, increases beta-lactam susceptibility and virulence, *J Bacteriol.* **188**,
 453 6669-79.

454 27. Hett, E. C., Chao, M. C., Deng, L. L. & Rubin, E. J. (2008) A mycobacterial enzyme essential for
 455 cell division synergizes with resuscitation-promoting factor, *PLoS Pathog.* **4**, e1000001.

28. Chao, M. C., Kieser, K. J., Minami, S., Mavrici, D., Aldridge, B. B., Fortune, S. M., Alber, T. & Rubin, E. J. (2013) Protein complexes and proteolytic activation of the cell wall hydrolase RipA regulate septal resolution in mycobacteria, *PLoS Pathog.* **9**, e1003197.
29. Hett, E. C., Chao, M. C. & Rubin, E. J. (2010) Interaction and modulation of two antagonistic cell wall enzymes of mycobacteria, *PLoS Pathog.* **6**, e1001020.
30. Chiaradia, L., Lefebvre, C., Parra, J., Marcoux, J., Burlet-Schiltz, O., Etienne, G., Tropis, M. & Daffe, M. (2017) Dissecting the mycobacterial cell envelope and defining the composition of the native mycomembrane, *Sci Rep.* **7**, 12807.
31. Wu, K. J., Boutte, C. C., Ioerger, T. R. & Rubin, E. J. (2019) *Mycobacterium smegmatis* HtrA Blocks the Toxic Activity of a Putative Cell Wall Amidase, *Cell Rep.* **27**, 2468-2479 e3.
32. Belardinelli, J. M., Stevens, C. M., Li, W., Tan, Y. Z., Jones, V., Mancina, F., Zgurskaya, H. I. & Jackson, M. (2019) The MmpL3 interactome reveals a complex crosstalk between cell envelope biosynthesis and cell elongation and division in mycobacteria, *Sci Rep.* **9**, 10728.
33. Dubee, V., Triboulet, S., Mainardi, J. L., Etheve-Quelquejeu, M., Gutmann, L., Marie, A., Dubost, L., Hugonnet, J. E. & Arthur, M. (2012) Inactivation of *Mycobacterium tuberculosis* l,d-transpeptidase LdtMt(1) by carbapenems and cephalosporins, *Antimicrob Agents Chemother.* **56**, 4189-95.
34. Diacon, A. H., van der Merwe, L., Barnard, M., von Groote-Bidlingmaier, F., Lange, C., Garcia-Basteiro, A. L., Sevene, E., Ballell, L. & Barros-Aguirre, D. (2016) beta-Lactams against Tuberculosis--New Trick for an Old Dog?, *N Engl J Med.* **375**, 393-4.
35. Raze, D., Verwaerde, C., Deloison, G., Werkmeister, E., Coupin, B., Loyens, M., Brodin, P., Rouanet, C. & Loch, C. (2018) Heparin-Binding Hemagglutinin Adhesin (HBHA) Is Involved in Intracytosolic Lipid Inclusions Formation in Mycobacteria, *Front Microbiol.* **9**, 2258.
36. van Kessel, J. C., Marinelli, L. J. & Hatfull, G. F. (2008) Recombineering mycobacteria and their phages, *Nat Rev Microbiol.* **6**, 851-7.

Table

Table 1. Total number of spectra for each protein in Membrane (MB) and Cell Wall (CW) fractions.

Protein	Fraction	mc ² 155	$\Delta MS0919$	$\Delta MS1285$	$\Delta MS1285 + \Delta MS0919$
MSMEG_1285	CW	2	1	0	0
	MB	0	0	0	0
	Total	2	1	0	0
MSMEG_0919 (HbhA_ <i>Msmeg</i>)	CW	7	0	10	0
	MB	10	0	9	0
	Total	17	0	19	0
MSMEG_6900 (PonA1)	CW	21	7	16	13
	MB	12	17	14	10
	Total	33	24	30	23
MSMEG_6201 (PonA2)	CW	16	9	16	12
	MB	8	11	6	5
	Total	24	20	22	17
MspA/B/C/D	CW	31	15	4	3
	MB	0	0	0	0
	Total	31	15	4	3
MSMEG_3145 (RipA_ <i>Msmeg</i>)	CW	0	3	0	6
	MB	0	1	0	5
	Total	0	4	0	11
MSMEG_6815	CW	1	5	0	3
	MB	0	1	1	3
	Total	1	6	1	6
MSMEG_6281 (Ami1_ <i>Msmeg</i>)	CW	2	2	2	7
	MB	0	0	1	4
	Total	2	2	3	11

Figure Legends

Figure 1. Rv0613c interacts with PonA1₃₂₁₋₅₁₅ in the BACTH system. (A) Red colony phenotype observed on MacConkey agar plates and (B) Blue colony phenotype on LB agar plates supplemented with IPTG and X-gal for *E. coli* transformed with pKT25_*rv0613c* and pUT18C_*ponA1*₃₂₁₋₅₁₅ (middle) in contrast to *E. coli* transformed with pKT25 and pUT18C_*ponA1*₃₂₁₋₅₁₅ (left) or with pKT25_*rv0613c* and pUT18C (right).

Figure 2. Rv0613c-⁶His co-precipitates MBP-PonA1₃₂₁₋₅₁₅. Coomassie-blue staining (upper panel) and immunoblot analysis using anti-MBP antibodies (lower panel) of total lysates of *E. coli* BL21(DE3) co-transformed with pET24a and pMAL-c2X_*ponA1*₃₂₁₋₅₁₅ (lane 1), with

pET24a_*rv0613c* and pMAL-c2X (lane 2), or with pET24a_*rv0613c* and pMAL-c2X_*ponA1*₃₂₁₋₅₁₅ (lane 3) and the corresponding elution fraction after purification by Ni-NTA chromatography. Protein marker (M). The arrows indicate the protein MBP-PonA1₃₂₁₋₅₁₅ (63 kDa).

Figure 3. Deletion of *MSMEG_1285* increases ampicillin resistance and the phenotype is exacerbated by additional deletion of *hbhA_Msmeg* (*MSMEG_0919*). (A) Plating of *Msmeg* mc²155, Δ *MS0919*, Δ *MS1285* and Δ *MS1285*+ Δ *MS0919* onto 7H11+OADC containing either Ampicillin (10-500 μ g/ml) or Rifampicin (0.1-10 μ g/ml) and (B) Serial dilutions (from 10⁰ to 10⁻⁵, anticlockwise) of *Msmeg* mc²155, Δ *MS0919*, Δ *MS1285* and Δ *MS1285*+ Δ *MS0919* onto 7H11+OADC containing either Ampicillin (50 μ g/ml) or Rifampicin (1 μ g/ml).

Figure 4. Generation of Δ *MS0919* and Δ *MS1285*+ Δ *MS0919* in *Msmeg* mc²155. (A) Genomic organization of the *hbhA_Msmeg* (*MSMEG_0919*) locus in *Msmeg* mc²155 and the Δ *MS1285* mutant and schematic representation for the insertion of the apramycin resistance cassette. The large black arrows depict *hbhA_Msmeg* (complete, upper line; interrupted, lower line). The grey arrows show the *hbhA_Msmeg* flanking genes and the white arrow indicates the apramycin resistance cassette. (B) Western-blot analysis using anti-Hsp65/GroL1 (upper panel) and anti-HbhA VF2 (lower panel) monoclonal antibodies on total lysates of *Msmeg* mc²155, Δ *MS0919*, Δ *MS1285* and Δ *MS1285*+ Δ *MS0919*. (C) Western-blot analysis using anti-Hsp65/GroL1 (upper panel) and anti-HbhA VF2 (lower panel) monoclonal antibodies on total lysates of *Msmeg* mc²155, the simple mutant Δ *MS0919* and the double mutant Δ *MS1285*+ Δ *MS0919* containing pMV361, pMV361_*MS0919* (+*MS0919*), pMV361_*MS1285* (+*MS1285*) or pMV361_*MS1285*_MS0919 (+*MS1285*_MS0919).

Figure 5. Overexpression of *ripA_Mtb* or *ami1* (*Msmeg* or *Mtb*) in *Msmeg* increases ampicillin resistance. Serial dilutions (from 10⁰ to 10⁻⁵) of *Msmeg* mc²155 containing pVV16 (empty vector) or overexpressing *ripA_Mtb*, *ami1_Msmeg* or *ami1_Mtb* were plated onto 7H11+OADC without (left panel) and with Ampicillin (right panel, 500 μ g/ml).

522

523 **Figure 6. Complementation of the double mutant $\Delta MS1285+\Delta MS0919$ with single *MS1285***
524 **restores sensitivity to the penicillin antibiotics, ampicillin and carbenicillin.** Serial dilutions (from
525 10^0 to 10^{-5} , anticlockwise) of *Msmeg* mc²155, $\Delta MS0919$ and $\Delta MS1285+\Delta MS0919$ containing
526 pMV361, pMV361_*MS0919* (+*MS0919*), pMV361_*MS1285* (+*MS1285*) or
527 pMV361_*MS1285_MS0919* (+*MS1285_MS0919*) were plated onto 7H11+OADC containing
528 Ampicillin (50 µg/ml), Carbenicillin (1,000 µg/ml), Isoniazid (5 µg/ml) or Ethionamide (5 µg/ml).

529

530 **Figure 7. Proposed global scheme of interactions for Rv0613c/MSMEG_1285.** PPI network (A) In
531 *Mtb* or *Msmeg* for Rv0613c or MS1285 and (B) In $\Delta rv0613c$ or $\Delta MS1285$ deletion mutant. The
532 representation of Msp-like porins only depict *Msmeg* cell wall, as no such porins were identified and
533 characterized in *Mtb*.

534

535 **Supplemental Information**

536 **Supplemental Table S1. List of strains and plasmids used in this study.**

537

538 **Supplemental Table S2. List of primers used in this study.**

539

540 **Supplemental Table S3. Total number of spectra for the 31 over-represented proteins (ratio ≥ 5)**
541 **in the $\Delta MS1285+\Delta MS0919$ double mutant.**